

Mean-Field Critical Behaviour for Percolation in High Dimensions

Takashi Hara ^{1,*} and Gordon Slade ^{2,**}

¹ Courant Institute of Mathematical Sciences, N. Y. U., 251 Mercer Street, New York, NY 10012, USA

² Department of Mathematics and Statistics, McMaster University, Hamilton, Ontario, Canada L8S4K1

Abstract. The triangle condition for percolation states that $\sum_{x,y} \tau(0,x) \tau(x,y) \cdot \tau(y,0)$ is finite at the critical point, where $\tau(x,y)$ is the probability that the sites x and y are connected. We use an expansion related to the lace expansion for a self-avoiding walk to prove that the triangle condition is satisfied in two situations: (i) for nearest-neighbour independent bond percolation on the d -dimensional hypercubic lattice, if d is sufficiently large, and (ii) in more than six dimensions for a class of “spread-out” models of independent bond percolation which are believed to be in the same universality class as the nearest-neighbour model. The class of models in (ii) includes the case where the bond occupation probability is constant for bonds of length less than some large number, and is zero otherwise. In the course of the proof an infrared bound is obtained. The triangle condition is known to imply that various critical exponents take their mean-field (Bethe lattice) values ($\gamma = \beta = 1, \delta = \Delta_t = 2, t \geq 2$) and that the percolation density is continuous at the critical point. We also prove that $\nu_2 = 1/2$ in (i) and (ii), where ν_2 is the critical exponent for the correlation length.

Table of Contents

1. Introduction	334
1.1. The Models	335
1.2. Main Results	337
1.3. Overview of the Proof	339
1.4. Organization, Notation, and Definitions	346
2. The Expansion	348
2.1. Derivation of the Expansion	349

* Supported by the Nishina Memorial Foundation and NSF grant PHY-8896163. Address after September 1, 1989: Department of Mathematics, University of Texas at Austin, Austin, TX 78712, USA. Electronic address: hara @ emx.utexas.edu

** Supported by NSERC grant A9351. Electronic address: slade @ sscvax.mcmaster.ca

2.2. Bounds in x -Space for Each Term of the Expansion	354
2.3. Bounds in k -Space for Each Term of the Expansion	358
3. Diagrammatic Estimates	360
3.1. Bounds on $\sum_x h_n(0, x)$	361
3.2. Bounds on $\sum_x x ^2 h_n(0, x)$	364
3.3. Proof of Lemma 3.3	367
4. Proof of the Main Results (Theorem 1.1) for the Nearest-Neighbour Model.	368
4.1. General Structure	368
4.2. Proof of Lemmas 4.1 and 4.2	370
4.3. Proof of Proposition 4.3	371
4.4. Proof that $\nu_2 = 1/2$	378
5. Proof of the Main Results (Theorem 1.2) for the Spread-Out Models	379
5.1. Proof of Theorem 1.2 Assuming Lemma 5.1	380
5.2. Proof of Lemma 5.1	383

1. Introduction

In the last decade significant progress has been made in the rigorous understanding of critical phenomena, particularly concerning its mean-field behaviour in high dimensions. For Ising and ϕ^4 spin systems one of the important ideas has been the combination of the infrared bound [16] with correlation inequalities. This method, together with other ideas, has led to considerable understanding of these models in four or more dimensions, including a proof of the triviality of ϕ^4 field theories in more than four dimensions and a proof that many critical exponents take their mean-field values above four dimensions [35, 1, 15, 4, 3].

For the self-avoiding walk no general proof of an infrared bound is known, and the methods which were successful for the spin systems cannot be applied. An alternate approach was introduced by Brydges and Spencer [12], who used the lace expansion to prove mean-field critical behaviour for the weakly self-avoiding walk in more than four dimensions. This method was further developed in [31–33], where mean-field critical behaviour was proved for the strictly self-avoiding walk above some undetermined dimension $d_0 > 4$, and in [25], where the infinite self-avoiding walk was constructed in high dimensions.

For percolation there is also no general proof of an infrared bound, and in fact there are indications that the infrared bound is violated in less than six dimensions [14, 41]. However based on an analogy with the bubble diagram, which played an important role in the analysis of the Ising and ϕ^4 models, Aizenman and Newman [6] introduced an unverified condition, the so-called triangle condition, which was shown by them to imply mean-field behaviour for the susceptibility in percolation models. The triangle condition is expected to hold above six dimensions. Subsequently further implications of the triangle condition were obtained in [8, 28]. In this paper we prove that the triangle condition is satisfied in two situations: (i) for independent nearest-neighbour Bernoulli bond percolation in sufficiently high dimensions, and (ii) in more than six dimensions for a class of “spread-out” models of independent bond percolation, which includes certain

finite range models as well as certain models in which the bond occupation probability decays exponentially as a function of the length of the bond. In the course of the proof we obtain a uniform infrared bound below the critical point, which is valid in high dimensions for model (i), and above six dimensions for the models in (ii). The method of proof can also be applied to site percolation, and yields the same results (with a suitable interpretation of the spread-out models). Consequences of the triangle condition are that the critical exponents γ, β, δ , and Δ_t (defined below) exist and take their mean-field values, and that the percolation density is continuous at the critical point.

The models in class (ii) are believed to be in the same universality class as the nearest neighbour-model. It is known that for the nearest neighbour model the upper critical dimension is at least six [13, 38], and the proof of this fact can be extended to the models in (ii) [39]. Thus our result strongly supports the conjecture that for these models the upper critical dimension is equal to six.

The proof of these results is based on an expansion for the two-point function which is related to the lace expansion for self-avoiding walk. The expansion is used to treat a percolation model as a perturbation of the random walk model whose transition probabilities are proportional to the percolation bond occupation probabilities. Similar methods can also be applied to branched polymers [21].

1.1. The Models

We consider independent Bernoulli bond percolation on the infinite d -dimensional hypercubic lattice \mathbb{Z}^d . To each unordered *bond* (pair of distinct sites) $b = \{x, y\}$ ($x, y \in \mathbb{Z}^d$) a random variable n_b is associated, which takes the values 0 and 1. The set of random variables $\{n_b\}$ is independent, and the distribution of n_b is given by

$$\text{Prob}(n_b = 1) = p_b, \quad \text{Prob}(n_b = 0) = 1 - p_b.$$

We require \mathbb{Z}^d -invariance (translation, reflection and rotation by $\pi/2$) for the $P_{\{x,y\}} = P_{\{0,y-x\}}$.

We consider the following possibilities for p_b :

(i) the nearest-neighbour model:

$$P_{\{0,x\}} = \begin{cases} p & \text{if } x \text{ is a nearest neighbour of } 0 \\ 0 & \text{otherwise.} \end{cases}$$

(ii) The spread-out models:

$$P_{\{0,x\}} = p \cdot L^{-d} g(x/L),$$

where $g: \mathbb{R}^d \rightarrow [0, \infty)$ is a given function which is normalized so that $\int g(x) d^d x = 1$, and is invariant under rotations by $\pi/2$ and reflections in the coordinate hyperplanes. The parameter L will be taken to be large. (This type of limit to study mean-field behaviour is related to the so-called *Kac limit* [23, 26].) A basic example is

$$g(x) = \begin{cases} 1 & \text{if } \|x\|_\infty \equiv \max_{1 \leq i \leq d} |x_i| \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

We require that g decay exponentially at infinity (i.e., there exist $C, \varepsilon > 0$ such that $g(x) \leq C \exp[-\varepsilon \|x\|_\infty]$). Then models (i) and (ii) are expected to be in the same universality class. The bond density p is the only parameter in these models (apart from an additional parameter h we shall mention briefly to define the critical exponent δ). For the models in (ii) we will show that the triangle condition is satisfied for $d > 6$, if L is sufficiently large and g is piecewise differentiable.

If $n_b = 1$ we say that b is *occupied*, while if $n_b = 0$ we say that b is *vacant*. We use $\text{Prob}_p(E)$ to denote the probability of an event E with respect to the joint distribution of the $\{n_b\}$, and denote expectation with respect to this distribution by $\langle \cdot \rangle_p$.

Given a bond configuration $\{n_b\}$, two sites x and y in the lattice are said to be *connected* if there exists a path from x to y which consists of occupied bonds. The *connected cluster* $C(x)$ of x is the random set of sites defined by

$$C(x) = \{y \in \mathbb{Z}^d: y \text{ is connected to } x\}.$$

The number of sites in $C(x)$ is denoted by $|C(x)|$.

We define the two-point function

$$\tau_p(x, y) = \text{Prob}_p(y \text{ is connected to } x), \tag{1.1}$$

the susceptibility

$$\chi(p) = \sum_x \tau_p(0, x) = \langle |C(0)| \rangle_p, \tag{1.2}$$

the percolation density

$$P_\infty(p) = \text{Prob}_p(|C(0)| = \infty), \tag{1.3}$$

and the magnetization

$$M(p, h) = 1 - \sum_{1 \leq n < \infty} e^{-hn} \text{Prob}_p(|C(0)| = n). \tag{1.4}$$

We also define two correlation lengths

$$\xi(p) = - \left[\lim_{n \rightarrow \infty} \frac{1}{n} \ln \tau_p(0, (n, 0, \dots, 0)) \right]^{-1} \tag{1.5}$$

and

$$\xi_2(p) = \left(\frac{\sum_x |x|^2 \tau_p(0, x)}{\sum_x \tau_p(0, x)} \right)^{1/2}. \tag{1.6}$$

To simplify the notation we will often omit the subscript p .

For the nearest-neighbour model it has been known for thirty years that (except for the trivial case $d = 1$) there is a critical value $p_c \in (0, 1)$ (depending on the dimension) such that the percolation density vanishes for $p < p_c$ and is nonzero for $p > p_c$ [10, 19]. Such a critical value of p also exists for the models in (ii), for the g 's we will consider. Recently it has been proved that p_c can also be characterized as $\sup\{p: \chi(p) < \infty\}$ [27, 2]. In this paper we are concerned with the critical

behaviour of the model, i.e., the behaviour of functions such as those defined above in the vicinity of $(p, h) = (p_c, 0)$. By analogy with other statistical mechanical models, and in agreement with numerical calculations, this behaviour is expected to be in the form of power laws, and we introduce the critical exponents $\gamma, \beta, \delta, \Delta_{t+1}, \nu$ and ν_2 as follows:

$$\chi(p) \sim (p_c - p)^{-\gamma} \quad \text{as } p \uparrow p_c, \quad (1.7)$$

$$P_\infty(p) \sim (p - p_c)^\beta \quad \text{as } p \downarrow p_c, \quad (1.8)$$

$$M(p_c, h) \sim h^{1/\delta} \quad \text{as } h \downarrow 0, \quad (1.9)$$

$$\langle |C(0)|^{t+1} \rangle / \langle |C(0)|^t \rangle \sim (p_c - p)^{-\Delta_{t+1}} \quad \text{as } p \uparrow p_c, \quad (1.10)$$

$$\xi(p) \sim (p_c - p)^{-\nu} \quad \text{as } p \uparrow p_c, \quad (1.11)$$

$$\xi_2(p) \sim (p_c - p)^{-\nu_2} \quad \text{as } p \uparrow p_c. \quad (1.12)$$

Here $f(p) \sim |p - p_c|^{-\lambda}$ is defined to mean that there exist positive constants C_1 and C_2 such that

$$C_1 |p - p_c|^{-\lambda} \leq f(p) \leq C_2 |p - p_c|^{-\lambda}.$$

There are other critical exponents that can be defined (see [17, 24, 37]), but these are the ones for which we can conclude mean-field values. The exponent δ is also often defined by the (formally) equivalent relation

$$\text{Prob}_{p_c} (|C(0)| = n) \sim n^{-1-1/\delta}.$$

So far very little has been rigorously proved about the existence of the above critical exponents. See [17, 24] for a review.

On the Bethe lattice (Cayley tree), it can be shown that γ, β, δ , and Δ_{t+1} exist and have the values $\gamma = \beta = 1$, $\delta = \Delta_{t+1} = 2$ for $t + 1 = 2, 3, 4, \dots$ [17]. The Bethe lattice critical exponents are known as the *mean-field values*, and it is expected that for the models (i) and (ii) in more than six dimensions all critical exponents take their mean-field values. The definition of ν and ν_2 is problematic on the Bethe lattice since these exponents are defined using the Euclidean structure, but with a proper definition it can be shown that $\nu_2 = 1/2$ [17]. On the hypercubic lattice it is expected that ν and ν_2 are equal, so the mean field values of these exponents is $1/2$. On the other hand, it has been rigorously shown by Chayes and Chayes [13] and by Tasaki [38] that if all the critical exponents exist, then they cannot simultaneously take their mean-field values in any dimension less than six. Thus the upper critical dimension of the system is expected to be six.

1.2. Main Results

Aizenman and Newman [6] introduced an unverified condition, the triangle condition, and showed that it implies that (1.7) holds with $\gamma = 1$. The triangle condition states that

$$\nabla(p_c) < \infty, \quad (1.13)$$

where $\mathcal{V}(p)$ is the triangle diagram

$$\mathcal{V}(p) = \sum_{x,y} \tau_p(0, x) \tau_p(x, y) \tau_p(y, 0). \tag{1.14}$$

Since the susceptibility $\chi(p) = \sum_x \tau(0, x)$ diverges as p approaches p_c from below for the models (i) and (ii) we are considering [6], the statement that the triangle diagram is finite at the critical point is not without content. Denoting by $\hat{\tau}_p(k)$ the Fourier transform of the two-point function, defined by

$$\hat{\tau}_p(k) = \sum_x \tau_p(0, x) e^{ik \cdot x},$$

the triangle diagram can be written

$$\mathcal{V}(p) = (2\pi)^{-d} \int_{[-\pi, \pi]^d} d^d k \hat{\tau}_p(k)^3.$$

If an infrared bound

$$\hat{\tau}_p(k) \leq C(p) k^{-2}, \text{ as } k \rightarrow 0,$$

were known, it would then follow that $\mathcal{V}(p) < \infty$ for $d > 6$. A uniform bound on $C(p)$ for $p < p_c$ would thus be tantamount to (1.13). Such a uniform bound, and hence the triangle condition, is expected to hold in more than six dimensions.

In addition to the fact that the triangle condition implies $\gamma = 1$, Barsky and Aizenman [8] have shown that the triangle condition implies that (1.8) and (1.9) hold with $\beta = 1$ and $\delta = 2$, and Nguyen [28] has shown that it implies that (1.10) holds with $\Delta_t = 2$ for $t = 2, 3, 4, \dots$. It follows from (1.8) that the percolation density P_∞ is continuous at the critical point, a fact which has been proved until now only for $d = 2$ [30]. (Continuity of P_∞ has been shown for all other values of p in [5].) Our main result is that the triangle condition is satisfied (i) for the nearest-neighbour model if the dimension d is sufficiently large, and (ii) for the spread-out models if L is sufficiently large, for $d > 6$. Hence all the above consequences of this condition also hold in these situations.

In this paper we prove the following theorems.

Theorem 1.1. *For nearest-neighbour independent Bernoulli bond percolation on \mathbb{Z}^d there is a $d_0 > 6$ such that for $d \geq d_0$ the infrared bound*

$$\hat{\tau}_p(k) \leq \text{const } k^{-2}, \text{ uniformly in } p < p_c$$

holds. In addition the triangle condition is satisfied, i.e.,

$$\mathcal{V}(p_c) = \sum_{x,y} \tau_{p_c}(0, x) \tau_{p_c}(x, y) \tau_{p_c}(y, 0) < \infty.$$

Theorem 1.2. *The infrared bound*

$$\hat{\tau}_p(k) \leq \text{const } k^{-2}, \text{ uniformly in } p < p_c,$$

and the triangle condition both hold for $d > 6$, for the spread-out models (ii), if L is sufficiently large (depending on d and g) and if $\frac{\partial^d g}{\partial x_1 \dots \partial x_d}$ is piecewise continuous

and g satisfies the following conditions:

$$g e^{\delta \|x\|_1} \in L_\infty(\mathbb{R}^d) \text{ for some } \delta > 0, \\ \int g(x) d^d x = 1,$$

$\int |\partial^I g(x)| d^d x < \infty$, where the derivative is interpreted as a distribution, and

$$\partial^I = \prod_{\mu \in I} \frac{\partial}{\partial x_\mu} \text{ and } I \subset \{1, 2, \dots, d\},$$

g is invariant under rotations by $\pi/2$ and reflections in the coordinate hyperplanes.

As a consequence of the proofs of these theorems, we will also show that the exponent ν_2 for the correlation length ξ_2 exists and takes its mean-field value

$$\nu_2 = 1/2 \tag{1.15}$$

for the nearest neighbour model if $d \geq d_0$ and for the spread-out models if $d > 6$ and $L \geq L_0$. However a more complicated analysis is required to control the correlation length ξ . It is proved in [20] that $\nu = 1/2$, under the same hypotheses as (1.15).

Also, Eq. (4.10) below gives an upper bound for the critical probability for the nearest-neighbour model in high dimensions, which together with a well-known lower bound states that there is a constant $C > 0$ (independent of d) such that

$$\frac{1}{2d-1} \leq p_c \leq \frac{1}{2d} + \frac{C}{d^2}.$$

We have done little to obtain the best possible value of d_0 in Theorem 1.1. Our current best estimate is $d_0 = 48$, obtained by a slightly more complicated analysis than that presented in this paper. This value can doubtless be improved, but a new idea will be needed to obtain the triangle condition for the nearest-neighbour model right down to the expected upper critical dimension of six. The fact that we are unable to do much better than $d_0 = 48$ suggests that we still do not have a very efficient expansion for percolation. (For the self-avoiding walk the situation is better: the lace expansion can be used to show that the bubble diagram is finite and $\gamma = 1, \nu_2 = 1/2$ for the self-avoiding walk if $d \geq 7$ [22]. Here γ is the exponent which measures the rate of divergence of the generating function at the critical point, and ν_2 is defined as for percolation. The upper critical dimension for self-avoiding walk is expected to be 4, so $d \geq 5$ should be optimal.) However in view of the fact that all of the models we are considering are believed to be in the same universality class, the conjecture of universality, together with Theorem 1.2, strongly supports mean-field behaviour for the nearest-neighbour model in more than six dimensions.

The method of proof involves an expansion whose convergence is assured by taking $1/d$ to be small in Theorem 1.1 and $1/L$ to be small in Theorem 1.2.

1.3. Overview of the Proof

In this section we describe the general structure of the proof of Theorem 1.1. The same ideas are used to prove Theorem 1.2. The basic structure of the proof is the

same as that used in [31], where in particular it was shown that the bubble diagram for the self-avoiding walk is finite at the critical point, in sufficiently high dimensions. (Proofs with this type of structure, in different contexts, can be found in [11, 36].) In order to focus on the main ideas here we make some simplifications and omit some details, deferring the complete proof to the remainder of the paper.

We define the following quantities:

$$T(p) = \nabla(p) - 1 = \sum_{x,y} \tau(0, x) \tau(x, y) \tau(y, 0) - \tau(0, 0)^3, \tag{1.16}$$

$$W(p) = \sum_x |x|^2 \tau(0, x)^2. \tag{1.17}$$

The massless gaussian propagator is given by

$$C(x, y) = (2\pi)^{-d} \int d^d k \frac{e^{ik \cdot (y-x)}}{1 - D(k)}, \text{ where } D(k) = d^{-1} \sum_{\mu=1}^d \cos k_\mu.$$

We also introduce the gaussian quantities T_G and W_G corresponding to $T(p)$ and $W(p)$, defined by replacing τ by C in (1.16) and (1.17). It is not difficult to show that for $d \geq 7$ there are constants K_1 and K_2 such that $T_G \leq K_1 d^{-1}$ and $W_G \leq K_2 d^{-1}$ (e.g., using Appendix B of [20] or Lemma 3.1 of [31]).

1.3.1. General Framework. The proof that the triangle condition is satisfied in high dimensions is accomplished by showing that the following three statements hold:

- (i) For $p < p_c$, $T(p)$ and $W(p)$ are continuous functions of p .
- (ii) For $p \leq 1/2d$, $T(p) \leq T_G \leq K_1 d^{-1}$ and $W(p) \leq W_G \leq K_2 d^{-1}$.
- (iii) Let d be sufficiently large and fix any $p \in [1/2d, p_c)$. If

$$T(p) \leq 4 K_1 d^{-1}, \quad W(p) \leq 4 K_2 d^{-1}, \quad \text{and} \quad 2dp \leq 4, \tag{1.18}$$

then in fact

$$T(p) \leq 3 K_1 d^{-1}, \quad W(p) \leq 3 K_2 d^{-1}, \quad \text{and} \quad 2dp \leq 3. \tag{1.19}$$

(In reality the precise statement of (iii) is more involved and can be found in Sect. 4.) In the course of the proof of (iii) the infrared bound is obtained.

Together (i), (ii) and (iii) imply that there is a forbidden region in the graph of $T(p)$ or $W(p)$, as depicted in Fig. 1. Therefore (1.19) holds and hence by definition of $T(p)$ it follows that

$$\nabla(p) \leq 1 + 3 K_1 d^{-1}, \text{ for all } p < p_c. \tag{1.20}$$

But since $\tau_p(x, y)$ is an increasing and continuous function of p [5], it follows from (1.20) and the monotone convergence theorem that

$$\nabla(p_c) = \lim_{p \uparrow p_c} \nabla(p) \leq 1 + 3 K_1 d^{-1},$$

and hence the triangle condition holds.

The proofs of (i) and (ii) are simple and are given in Sect. 4.2. The difficult part of the proof is to obtain (iii). This is done by obtaining an expression for the Fourier transform $\hat{\tau}_p(k)$, which under the assumption (1.18) can be shown to be a




Fig. 1. Forbidden region (shaded) in the graph of T or W versus p , for $p < p_c$.

small perturbation of the massless gaussian propagator $\hat{C}(k) = [1 - D(k)]^{-1}$. Hence $T(p)$ and $W(p)$ are close to their gaussian counterparts T_G and W_G , and thus the improved bounds on $T(p)$ and $W(p)$ in (1.19) follow.

For the self-avoiding walk the analogue of step (iii) was obtained using the lace expansion. Here we derive and use an analogous expansion for percolation, which yields an expression for $\hat{\tau}_p(k)$. The lace expansion was derived in [12] using an expansion followed by resummation, but it can also be derived using the inclusion-exclusion relation [34]. It is the latter approach that we use for percolation. (In a similar spirit, Park [29] has used the inclusion-exclusion relation to study intersection probabilities for simple random walk.)

1.3.2. The Expansion. We describe the expansion in detail in Sect. 2, but let us here give the basic idea of the expansion and explain how it can be used to obtain (iii). We begin with some definitions. Two sites x and y are said to be *doubly connected* (in a given configuration) if there exist two self-avoiding walks from x to y , consisting of occupied bonds, which are distinct in the sense that they do not share a common bond (although they may share common sites). If x and y are connected, but not doubly connected, then there must be at least one bond in the connected cluster of x whose removal would disconnect x and y . Such a bond is referred to as a *pivotal* bond for the connection of x and y . There is a natural order for the set of pivotal bonds, namely the first pivotal bond is the pivotal bond $b = \{u, v\}$ such that one endpoint, say u , of b is doubly connected to x . Either the other endpoint v of b is doubly connected to y , in which case there are no further pivotal bonds for the connection of x and y , or it is not. In the latter case, the second pivotal bond is the one for which one endpoint is doubly connected to v , and so on. This leads to the picture of a cluster joining x and y shown in Fig. 2. This picture was also used in [27]. It is convenient to always regard a site as being doubly connected to itself. In Fig. 2(b), the pivotal bonds divide the cluster into parts which are mutually avoiding in the sense that no two can share a common




Fig. 2. (a) A configuration in which x and y are connected. (b) Schematic representation of the configuration in (a)

site. This represents a kind of repulsive interaction similar to that of self-avoiding walk.

The event that 0 is connected to x is then the disjoint union of the event that 0 is doubly connected to x and the event that there is a pivotal bond for the connection (and hence a first pivotal bond for the connection). We denote the former event symbolically as

$$\{0 \text{ is doubly connected to } x\} = 0 \text{ } \text{---} \text{ } \text{---} \text{ } x,$$

and the latter as

$$\{0 \text{ is connected to } x \text{ but not doubly}\} = \bigcup_{(u,v)} 0 \text{ } \text{---} \text{ } \text{---} \text{ } u \text{ } \text{---} \text{ } v \text{ } \text{---} \text{ } x. \quad (1.21)$$

Here the *ordered* bond (u, v) is the first pivotal bond for the connection, so 0 is doubly connected to u . The union in (1.21) is disjoint. Using this notation, and the convention that $\tau(x, x) = 1$, we have

$$\tau(0, x) = \text{Prob}(0 \text{ } \text{---} \text{ } \text{---} \text{ } x) + \sum_{(u,v)} \text{Prob}(0 \text{ } \text{---} \text{ } \text{---} \text{ } u \text{ } \text{---} \text{ } v \text{ } \text{---} \text{ } x). \quad (1.22)$$

At this point we need to make the following definitions.

Definition 1.3. Given a bond configuration $\{n_b\}$ and a set A of sites, we define

(a) the connected cluster of x in $\mathbb{Z}^d \setminus A$:

$$C^A(x) = \{y \in \mathbb{Z}^d: y \text{ and } x \text{ are connected using bonds having no endpoint in } A\}.$$

If $y \in C^A(x)$ we say that y and x are connected in $\mathbb{Z}^d \setminus A$,

(b) the reduced two-point function:

$$\tau_p^A(x, y) = \text{Prob}_p(y \in C^A(x)),$$

(c) the connected cluster of x after the bond $\{u, v\}$ has been made vacant:

$$\tilde{C}^{\{u,v\}}(x) = \{y \in C(x): y \text{ remains connected to } x \text{ after } n_{\{u,v\}} \text{ is set equal to zero}\}.$$

Now we use a lemma from [6] (which is stated and proved in Sect. 2.1) to write the summand in the second term on the right side of Eq. (1.22) as follows:

$$\text{Prob} (0 \text{ --- } \overset{u}{\curvearrowright} \text{ --- } v \text{ --- } x) = p \langle I[0 \text{ --- } u] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle. \quad (1.23)$$

The factor of p on the right side is due to the fact that $\{u, v\}$ is occupied, while the restricted two-point function and the double connection of 0 and u are due to the fact that (u, v) is the first pivotal bond. Next we replace the restricted two-point function in (1.23) by

$$\tau^{\tilde{C}^{(u,v)}(0)}(v, x) = \tau(v, x) - [\tau(v, x) - \tau^{\tilde{C}^{(u,v)}(0)}(v, x)]. \quad (1.24)$$

This yields the equation

$$\begin{aligned} \tau(0, x) = & \text{Prob} (0 \text{ --- } x) + p \sum_{(u,v)} \text{Prob}(0 \text{ --- } u) \tau(v, x) \\ & - p \sum_{(u,v)} \langle I[0 \text{ --- } u] (\tau(v, x) - \tau^{\tilde{C}^{(u,v)}(0)}(v, x)) \rangle. \end{aligned} \quad (1.25)$$

Our goal is to manipulate the right-hand side of (1.25) so as to obtain terms which either involve a convolution with τ evaluated at x , as in the second term on the right side, or involve a multiple connection from x and no explicit two-point function, as in the first term on the right side. (The reason for this will become clear when we take Fourier transforms.) To this end we first observe that

$$\tau(v, x) - \tau^{\tilde{C}^{(u,v)}(0)}(v, x) = \langle I[v \text{ is connected to } x \text{ in } \mathbb{Z}^d \text{ but not in } \mathbb{Z}^d \setminus \tilde{C}^{(u,v)}(0)] \rangle. \quad (1.26)$$

The event on the right side of (1.26) is such that **every** occupied self-avoiding walk from v to x must pass through $\tilde{C}^{(u,v)}(0)$ (which within the inner expectation (1.26) on the right side of (1.25) represents a given **deterministic** fixed set of sites). As an example of a situation that can occur in this event, we consider the configuration in Fig. 3. There (u_1, v_1) is the first pivotal bond such that every occupied self-avoiding walk from v to u_1 passes through $\tilde{C}^{(u,v)}(0)$, and z' is the




Fig. 3. An example of a configuration contributing to the right side of Eq. (1.26). Thick lines represent the bonds connecting the sites in $\tilde{C}^{(u,v)}(0)$. Thin lines represent the bonds connecting v and x . The point t is a point in $\tilde{C}^{(u,v)}(0)$ on an occupied self-avoiding walk from z' to u_1

latter endpoint of the previous pivotal bond $\{z, z'\}$. Other possibilities and special cases exist, for example v could be doubly connected to x , but we ignore such cases in this overview. By another application of the lemma quoted above (1.23), the contribution to (1.26) due to the configuration in Fig. 3 is equal to

$$\sum_{(u_1, v_1)} \sum_{z'} \langle I[E(v, z', (u_1, v_1))] \tau^{\tilde{C}^{(u_1, v_1)}(v)}(v_1, x) \rangle, \tag{1.27}$$

where $E(v, z', (u_1, v_1))$ is the event that (a) v is connected to z' via a pivotal bond $\{z, z'\}$, (b) z' is doubly connected to u_1 , with one of the occupied paths passing through a site in $\tilde{C}^{(u, v)}(0)$, and (c) $\{u_1, v_1\}$ is occupied and is the first pivotal bond such that every occupied self-avoiding walk from v to u_1 passes through $\tilde{C}^{(u, v)}(0)$. As before we will replace the restricted two-point function in (1.27) using the analogue of (1.24).

This allows us to write (1.25) as

$$\begin{aligned} \tau(0, x) = & \text{Prob}(0 \text{ --- } x) + p \sum_{(u, v)} \text{Prob}(0 \text{ --- } u) \tau(v, x) \\ & - p \sum_{\substack{(u, v) \\ (u_1, v_1)}} \sum_{z'} \langle I[0 \text{ --- } u] \langle I[E(v, z', (u_1, v_1))] \rangle \rangle \tau(v_1, x) \\ & + \text{other cases} + \text{remainder}. \end{aligned} \tag{1.28}$$

The remainder comes from the use of (1.24) in (1.27). Let us now for simplicity ignore the other cases and the remainder in (1.28). We take the Fourier transform of (1.28), and use the fact that the Fourier transform of a convolution is the product of the Fourier transforms. Solving the resulting equation for $\hat{\tau}_p(k)$ yields

$$\hat{\tau}(k) = \frac{1 + \hat{g}_0(k)}{1 - 2dp D(k) - \hat{\Pi}_0(k) + \hat{\Pi}_1(k)} + \text{corrections}, \tag{1.29}$$

where

$$\hat{g}_0(k) = \sum_{x \neq 0} \text{Prob}(0 \text{ --- } x) e^{ik \cdot x}, \tag{1.30}$$

$$2dp D(k) + \hat{\Pi}_0(k) = p \sum_{(u, v)} \text{Prob}(0 \text{ --- } u) e^{ik \cdot v}, \tag{1.31}$$

(the first term on the left side of (1.31) is the $u = 0$ term on the right side) and

$$\hat{\Pi}_1(k) = p \sum_{\substack{(u, v) \\ (u_1, v_1)}} \sum_{z'} \langle I[0 \text{ --- } u] \langle I[E(v, z', (u_1, v_1))] \rangle \rangle e^{ik \cdot v_1}. \tag{1.32}$$

1.3.3. Bounding Terms in the Expansion. We can estimate \hat{g}_0 , $\hat{\Pi}_0$, and $\hat{\Pi}_1$ using the van den Berg-Kesten inequality [9]. In the form that we need it the van den Berg-Kesten inequality states the following. Let V_1, \dots, V_n be sets of paths in the lattice, and let E_i , $i = 1, \dots, n$, be the event that at least one of the paths in V_i is occupied. Then

$$\begin{aligned} \text{Prob}[\text{there exist pairwise distinct occupied paths } \omega_1 \in V_1, \dots, \omega_n \in V_n] \\ \leq \text{Prob}[E_1] \dots \text{Prob}[E_n]. \end{aligned} \tag{1.33}$$

The inequality (1.33) with $n = 2$ can be used to estimate (1.30) and (1.31) by taking $V_1 = V_2$ to be the set of all paths from 0 to x :

$$|\hat{g}_0(k)| \leq \sum_{x \neq 0} \tau(0, x)^2 \leq T(p), \tag{1.34}$$

$$|\hat{\Pi}_0(k)| \leq 2dp \sum_{u \neq 0} \text{Prob}(0 \text{ --- } u) \leq 2dp \sum_{x \neq 0} \tau(0, x)^2 \leq 2dp T(p). \tag{1.35}$$

The estimation of (1.32) is more involved. We will use the inequality

$$I[E(v, z', (u_1, v_1))] \leq \sum_t I[t \in \tilde{C}^{(u, v)}(0)] I[\bar{E}(v, z', t, (u_1, v_1))], \tag{1.36}$$

where $\bar{E}(v, z', t, (u_1, v_1))$ is the event that (a) v is connected to z' , (b) z' is doubly connected to u_1 , with one of the occupied paths passing through t , (c) (u_1, v_1) is occupied, and (d) all of these connections are given by *distinct* paths. With (1.33), this gives

$$\begin{aligned} |\hat{\Pi}_1(k)| &\leq p \sum_{\substack{(u, v) \\ (u_1, v_1)}} \sum_{t, z'} \langle I[0 \text{ --- } u] I[t \in \tilde{C}^{(u, v)}(0)] \langle I[\bar{E}(v, z', t, (u_1, v_1))] \rangle \rangle \\ &\leq 2dp^2 \sum_{(u, v), u_1, t, z'} \langle I[0 \text{ --- } u] I[t \in \tilde{C}^{(u, v)}(0)] \rangle \tau(v, z') \tau(z', t) \tau(t, u_1) \tau(z', u_1) \\ &\leq 2dp^2 \sum_{(u, v), u_1, t, z', w} \tau(0, u) \tau(0, w) \tau(w, u) \tau(w, t) \tau(v, z') \tau(z', t) \tau(t, u_1) \tau(z', u_1). \end{aligned}$$

To better visualize this inequality we will introduce a diagrammatic notation in which it is represented by

$$|\hat{\Pi}_1| \leq 2dp \text{  }. \tag{1.37}$$

(One factor of p is inherent in the diagram.) As we shall show in Sects. 2.3 and 3.1, the right side of (1.37) can be bounded in terms of $T(p)$ to give

$$|\hat{\Pi}_1| \leq (2dp)^2 (1 + T(p))^2 [T(p)/3 + (2T(p)/3d)^{1/2} + 1/d]. \tag{1.38}$$

The right side of (1.38) is $O(d^{-1})$ for large d , under the assumption (1.18). Analogous upper bounds can be obtained for first and second k -derivatives of \hat{g}_0 , $\hat{\Pi}_0$, and $\hat{\Pi}_1$ involving $W(p)$ as well as $T(p)$.

1.3.4. Bounds on $\hat{\tau}(k)$. Proof Completed. In this section we describe how (1.19) can be obtained, assuming (1.18). Incorporating the corrections in Eq. (1.29) into corrections to \hat{g}_0 and $\hat{\Pi}_0 - \hat{\Pi}_1$, $\hat{\tau}(k)$ can be written in the form

$$\hat{\tau}(k) = \frac{1 + \hat{g}(k)}{1 - 2dp D(k) - \hat{\Pi}(k, p)}, \tag{1.39}$$

where

$$\hat{g}(k) = \hat{g}_0(k) + \text{corrections},$$

and

$$\hat{\Pi}(k, p) = \hat{\Pi}_0(k) - \hat{\Pi}_1(k) + \text{corrections}.$$

Now for any $p < p_c$,

$$1 - 2dp D(k) - \hat{\Pi}(k, p) = 1 - 2dp D(k) - \hat{\Pi}(k, p) - [1 - 2dp - \hat{\Pi}(0, p)] + [1 - 2dp - \hat{\Pi}(0, p)].$$

Using (1.18), it follows from (1.34) that $1 + \hat{g}(0) \approx 1$ for large d , and hence (since $\hat{\tau}(0) = \chi(p)$) the last term on the right side is

$$1 - 2dp - \hat{\Pi}(0, p) = [1 + \hat{g}(0)] \chi(p)^{-1} \approx \chi(p)^{-1} \geq 0. \tag{1.40}$$

Therefore

$$\hat{\tau}(k) \leq \frac{1 + |\hat{g}(k)|}{2dp(1 - D(k)) + \hat{\Pi}(0, p) - \hat{\Pi}(k, p)}.$$

The difference of $\hat{\Pi}$'s in the denominator can be controlled in terms of the second k -derivative of $\hat{\Pi}$ using (1.18), along the lines we have outlined above for bounding $\hat{\Pi}_1$. The assumption on W in (1.18) is used at this point. This allows us to absorb the difference of $\hat{\Pi}$'s into the gaussian term $2dp(1 - D(k))$, which with (1.34) yields an infrared bound. The fact that $2dp > 1$ is used at this point. We then write $T(p)$ and $W(p)$ in terms of $\hat{\tau}(k)$, and using this infrared bound obtain the improved estimate (1.19) for $T(p)$ and $W(p)$. The inequality $2dp \leq 3$ follows from (1.40) and the fact that $\hat{\Pi}$ is $O(d^{-1})$. Although the proof of the infrared bound initially uses the assumption (1.18), once we have derived the stronger statement (1.19) the assumption holds automatically.

This completes the outline of the proof. The remainder of the paper is devoted to giving a rigorous proof of Theorems 1.1 and 1.2, following this outline. In the next section we describe the organization of the remainder of the paper and summarize some notation and definitions.

1.4. Organization, Notation and Definitions

In this section we describe the organization of the remainder of the paper and summarize some definitions. We also introduce a convenient diagrammatic notation.

In Sect. 2 a detailed derivation of the expansion for the two-point function is given, and estimates are given for the terms of the expansion in both x -space and k -space. These estimates are given by diagrams as in (1.37). In Sect. 3 it is shown how the diagrams can be bounded by products of T , W and a small number of related quantities which play a role in the precise version of the assumption (1.18). In Sect. 4 the statements (i), (ii) and the precise version of (iii) (from Sect. 1.3.1) are proved, completing the proof of Theorem 1.1. In Sect. 4.4 the proof that $\nu_2 = 1/2$ for the nearest-neighbour model when $d \geq d_0$ is given. Finally in Sect. 5 the modifications to the proof of Theorem 1.1 which are necessary to prove Theorem 1.2 are described.

We now list several definitions, some of which were made in the last section and some of which are new.

Definition 1.4. (a) A *bond* is an unordered pair of distinct sites $\{x, y\}$. An *ordered bond* is denoted (x, y) . A *path* from x to y is a self-avoiding walk (not necessarily

nearest-neighbour for the spread-out models) from x to y , considered as a set of bonds. Given a bond configuration $\{n_b\}$, an *occupied path* is a path consisting of occupied bonds. Two paths are *distinct* if they have no bonds in common. (Distinct paths may have common sites.)

(b) Given a bond configuration, two sites x and y are *connected* if there is an occupied path from x to y . They are *doubly connected* if there are at least two distinct occupied paths from x to y . A site x is always considered to be doubly-connected to itself.

(c) Given a set of sites $A \subset \mathbb{Z}^d$ and a bond configuration, two sites x and y are *connected in A* if there is an occupied path from x to y consisting of bonds whose endpoints both lie in A . The sites x and y are *doubly-connected in A* if there are at least two distinct occupied paths from x to y consisting of bonds whose endpoints both lie in A . Two sites x and y are *connected through A* if there is at least one occupied path from x to y and if in addition every occupied path from x to y has at least one bond with an endpoint in A . Similarly we define x and y to be *doubly-connected through A* if x and y are doubly-connected and connected through A .

(d) Given a bond configuration, the connected cluster $C(x)$ of x is given by

$$C(x) = \{y \in \mathbb{Z}^d : y \text{ is connected to } x\}.$$

Given a set A of sites, we define

$$C^A(x) = \{y \in \mathbb{Z}^d : y \text{ and } x \text{ are connected in } \mathbb{Z}^d \setminus A\},$$

and the restricted two-point function

$$\tau_p^A(x, y) = \text{Prob}_p(y \in C^A(x)).$$


Given a bond $\{u, v\}$, we define

$$\tilde{C}^{(u,v)}(x) = \{y \in \mathbb{Z}^d : y \text{ is connected to } x \text{ in the new configuration obtained by setting } n_{\{u,v\}} = 0\}.$$

(e) Given a bond configuration, a bond $\{u, v\}$ (occupied or not) is called *pivotal* for the connection from x to y if either $x \in C(u)$ and $y \in C(v)$, or $x \in C(v)$ and $y \in C(u)$, but $y \notin \tilde{C}^{(u,v)}(x)$. Similarly an ordered bond (u, v) is pivotal for the connection from x to y if $x \in \tilde{C}^{(u,v)}(u)$, $y \in \tilde{C}^{(u,v)}(v)$, $y \notin \tilde{C}^{(u,v)}(x)$. If x and y are connected then there is a natural order to the set of occupied pivotal bonds for the connection from x to y (providing there exists one or more occupied pivotal bond), and each of these pivotal bonds can be ordered in a natural way, as follows. The *first pivotal bond from x to y* is the **ordered** occupied pivotal bond (u, v) such that u is doubly connected to x . If (u, v) is the first pivotal bond for the connection from x to y , then the *second pivotal bond* is the first pivotal bond for the connection from v to y , and so on.


Finally we describe the diagrammatic notation that will be used for the remainder of the paper. This notation has the virtue of making cumbersome expressions more transparent. We will denote the event that there is an occupied path between x and y by a solid line joining x and y : x ——— y . In a diagram

consisting of several such solid lines it is always to be understood that there are *distinct* paths making the required connections. For example, the diagram



denotes the event that y is connected to z , z is connected to u , z is connected to x , and x is connected to u , all by distinct paths.


We will also use a Feynman diagram notation in which full propagators are represented by wavy lines, unlabelled vertices are summed over the lattice, and labelled vertices are fixed. In addition we will shade in any loop for which all vertices on the loop may coincide. In unshaded loops the summation is constrained such that at least two of the vertices must be distinct. For example the diagram



represents

$$\tau(y, z) \sum_{u, x} \tau(z, u) \tau(u, x) \tau(x, z) = \tau(y, z) V(p),$$

while the diagram



represents

$$\tau(y, z) \left[\sum_{u, x} \tau(z, u) \tau(u, x) \tau(x, z) - \tau(z, z)^3 \right] = \tau(y, z) [V(p) - 1].$$

A pair of wavy lines terminating together in bars denotes two propagators ending at two sites y and y' and carries a factor of $p_{\{y, y'\}}$ and a sum over all ordered bonds (y, y') . For example,

$$0 \text{ (loop) } = \sum_{(y, y')} p_{\{y, y'\}} \tau(0, y) \tau(y', 0).$$

2. The Expansion

In this section we derive the expansion for $\tau_p(0, x)$ which is the main tool used in the proof of Theorems 1.1 and 1.2. The results of this section are valid for general \mathbb{Z}^d -invariant bond percolation models, long or short range. (In fact, Propositions 2.3 and 2.4 hold without the assumption of \mathbb{Z}^d -invariance.) To simplify the

connected bond cluster of the origin which remains after setting $n_{(u,v)} = 0$. The event that E occurs on B is defined to be the set consisting of those bond configurations for which the new configuration obtained by setting $n_b = 0$ for all $b \notin B$ is a configuration in the event E . The lemma involves events satisfying:

$$E \text{ occurs and the ordered bond } (u, v) \text{ is pivotal for the connection from } 0 \text{ to } x \\ \Rightarrow E \text{ occurs on } \tilde{C}^{(u,v)}(0)_b. \quad (2.5)$$

An example of an E satisfying (2.5) is the event that 0 is doubly connected to u .

For simplicity we consider only the case $p < p_c$, which is sufficient for our needs.

Lemma 2.1. *Let E be an event satisfying (2.5), and fix $p < p_c$. Then*

$$\langle I[E \text{ occurs and } (u, v) \text{ is occupied and pivotal for the connection from } 0 \text{ to } x] \rangle \\ = p_{uv} \langle I[E \text{ occurs and } u \in C(0)] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle.$$

Proof. The proof is by conditioning on $\tilde{C}^{(u,v)}(0)$. Since $p < p_c$, this cluster is finite with probability one. Let

$$\mathcal{E} = \{\text{finite connected bond clusters } B: 0, u \in B_s \text{ and } E \text{ occurs on } B\}.$$

Since E satisfies (2.5), apart from a set of measure zero

$$\{E \text{ occurs and } (u, v) \text{ is pivotal}\} = \bigcup_{B \in \mathcal{E}} \{\tilde{C}^{(u,v)}(0)_b = B \text{ and } (u, v) \text{ is pivotal}\}.$$

Therefore

$$\langle I[E \text{ occurs and } (u, v) \text{ is occupied and pivotal}] \rangle \\ = p_{uv} \langle I[E \text{ occurs and } (u, v) \text{ is pivotal}] \rangle \\ = p_{uv} \sum_{B \in \mathcal{E}} \text{Prob}((u, v) \text{ is pivotal} | \tilde{C}^{(u,v)}(0)_b = B) \text{Prob}(\tilde{C}^{(u,v)}(0)_b = B).$$

Since

$$\{\tilde{C}^{(u,v)}(0)_b = B \text{ and } (u, v) \text{ is pivotal}\} \\ = \{\tilde{C}^{(u,v)}(0)_b = B \text{ and } v \text{ is connected to } x \text{ in } \mathbb{Z}^d \setminus B_s\},$$

and since the events $\{\tilde{C}^{(u,v)}(0)_b = B\}$ and $\{v \text{ is connected to } x \text{ in } \mathbb{Z}^d \setminus B_s\}$ are independent, the conditional probability is equal to

$$\langle I[v \text{ is connected to } x \text{ in } \mathbb{Z}^d \setminus B_s] \rangle = \tau^{B_s}(v, x).$$

Therefore the above sum can be written

$$p_{uv} \sum_{B \in \mathcal{E}} \tau^{B_s}(v, x) \langle I[\tilde{C}^{(u,v)}(0)_b = B] \rangle = p_{uv} \sum_{B \in \mathcal{E}} \langle \tau^{\tilde{C}^{(u,v)}(0)}(v, x) I[\tilde{C}^{(u,v)}(0)_b = B] \rangle \\ = p_{uv} \langle I[u \in \tilde{C}^{(u,v)}(0) \text{ and } E \text{ occurs on } \tilde{C}^{(u,v)}(0)] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle \\ = p_{uv} \langle I[u \in \tilde{C}^{(u,v)}(0) \text{ and } E \text{ occurs}] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle. \quad (2.6)$$

The last equality is due to the fact that if E occurs, but not on $\tilde{C}^{(u,v)}(0)_b$, then by (2.5) (u, v) cannot be pivotal, in which case either $v \in \tilde{C}^{(u,v)}(0)$ or v is not connected to x , and hence $\tau^{\tilde{C}^{(u,v)}(0)}(v, x) = 0$.




Fig. 4. Examples of configurations contributing to $I_D(v, x; A)$

But now if $u \in C(0) \setminus \tilde{C}^{(u,v)}(0)$ then 0 is connected to u through $\{v\}$, and hence $\tau^{\tilde{C}^{(u,v)}(0)}(v, x) = 0$. Thus the right side of (2.6) is equal to

$$p_{uv} \langle I[E \text{ occurs and } u \in C(0)] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle,$$

and the lemma is proved. \square

We now use this lemma, with E the event that 0 is doubly connected to u , to write the second term on the right side of (2.3) as

$$\sum_{(u,v)} p_{uv} \langle I[0 \text{ } \langle \! \! \rangle \! \! \rangle \! \! \rangle u] \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \rangle.$$

To implement the inclusion-exclusion principle, we replace the restricted two point function by

$$\tau^{\tilde{C}^{(u,v)}(0)}(v, x) = \tau(v, x) - [\tau(v, x) - \tau^{\tilde{C}^{(u,v)}(0)}(v, x)]. \tag{2.7}$$

This gives (2.1) with $N = 0$, if we define

$$H_0(0, y') = \sum_{y \neq 0} p_{yy'} \langle I[0 \text{ } \langle \! \! \rangle \! \! \rangle \! \! \rangle y] \rangle = \sum_y p_{yy'} g_0(0, y) \tag{2.8a}$$

and

$$R_0(0, x) = \sum_{(u,v)} p_{uv} \langle I[0 \text{ } \langle \! \! \rangle \! \! \rangle \! \! \rangle u] \{ \tau(v, x) - \tau^{\tilde{C}^{(u,v)}(0)}(v, x) \} \rangle. \tag{2.8b}$$

To proceed further we rewrite the difference of two-point functions occurring in the remainder R_0 using the following lemma. In preparation for the statement of the lemma we make the following definition, which is illustrated in Fig. 4. Given a set A of sites, define

$$\begin{aligned} I_D(v, x; A) = & I[v \text{ and } x \text{ are doubly connected through } A] \\ & + I[\exists z' \neq v \text{ such that } z' \text{ is connected to } v \text{ in } \mathbb{Z}^d \setminus A, \\ & z' \text{ is an endpoint of a pivotal bond for the connection of } v \text{ and } x, \\ & \text{and } z' \text{ is doubly connected to } x \text{ through } A]. \end{aligned} \tag{2.9}$$

Lemma 2.2. *Given a set A of sites and two fixed sites v and x ,*

$$\tau(v, x) - \tau^A(v, x) = \langle I_D(v, x; A) \rangle + \sum_{(y,y')} p_{yy'} \langle I_D(v, y; A) \tau^{\tilde{C}^{(y,y')}(v)}(y', x) \rangle.$$

Proof. By definition of τ^A ,

$$\tau(v, x) - \tau^A(v, x) = \langle I[v \text{ and } x \text{ are connected through } A] \rangle.$$

Each configuration contributing to the right side belongs to exactly one of the following two cases.

Case 1. The sites v and x are doubly connected through A . This contributes to I_D , giving the first term on the right side of (2.9).

Case 2. The sites v and x are connected through A but are not doubly connected through A . In this case there is at least one pivotal bond for the connection from v to x , and we subdivide this case into the following cases.

Case 2a. There is no pivotal bond (w, w') (for the connection from v to x) such that w is connected to v through A . This gives the other contribution to I_D in (2.9). There z' is the latter endpoint of the last pivotal bond for the connection of v to x .

Case 2b. There is a pivotal bond whose first endpoint is connected to v through A . In this case there is a first such pivotal bond (y, y') . The contribution due to this case to $\tau(v, x) - \tau^A(v, x)$ is

$$\sum_{(y, y')} \langle I[(y, y') \text{ is the first pivotal bond (for the connection from } v \text{ to } x) \text{ whose first endpoint is connected to } v \text{ through } A] \rangle.$$

Let E be the event that v and y are connected through A , and there is no pivotal bond (w, w') for the connection from v to y such that w is connected to v through A . Then E satisfies (2.5) (with (u, v) replaced by (y, y') and 0 by v) and $I[E] = I_D(v, y; A)$, and hence by Lemma 2.1 the above expression is equal to the last term on the right side of the equation in the statement of the lemma. This completes the proof of Lemma 2.2. \square

We now return to the derivation of the expansion. Beginning with the equation

$$\tau(0, x) = \delta_{x,0} + g_0(0, x) + \sum_{v \neq 0} p_{0v} \tau(v, x) + \sum_{y'} \Pi_0(0, y') \tau(y', x) - R_0(0, x),$$

in which g_0 , Π_0 and R_0 were defined in Eqs. (2.4), (2.8a), and (2.8b), we use Lemma 2.2, with $A = \tilde{C}^{\{u,v\}}(0)$, to replace the difference of two-point functions occurring in R_0 . This leads to a nested expectation in R_0 . In order to specify unambiguously to which expectation the set $\tilde{C}^{\{u,v\}}(0)$ (which is defined in terms of a given bond configuration) corresponds, we use subscripts in nested expectations, i.e., $\tilde{C}_n^{\{u,v\}}(0)$ denotes $\tilde{C}^{\{u,v\}}(0)$ defined by the bond configuration corresponding to the configuration $\langle \cdot \rangle_{(n)}$. In applying Lemma 2.2 to (2.8b), $\tilde{C}^{\{u,v\}}(0)$ is random in the expectation of (2.8b), but deterministic with respect to the expectation produced by application of Lemma 2.2. Using subscripts we have

$$\begin{aligned} R_0(0, x) &= \sum_{(u,v)} p_{uv} \langle I[0 \text{ --- } u] \langle I_D(v, x; \tilde{C}_0^{\{u,v\}}(0)) \rangle_{(1)} \rangle_{(0)} \\ &\quad + \sum_{(u,v)} p_{uv} \langle I[0 \text{ --- } u] \sum_{(y,y')} p_{yy'} \langle I_D(v, y; \tilde{C}_0^{\{u,v\}}(0)) \tau_1^{(y,y')(v)}(y', x) \rangle_{(1)} \rangle_{(0)}. \end{aligned}$$

The first term gives g_1 . In the second term we replace the restricted two-point function $\tau_1^{\tilde{C}_0^{\{u,v\}}(v)}$ by an unrestricted two-point function plus a correction as in (2.7). The term involving the unrestricted two-point function gives rise to a term $\sum_{y'} \Pi_1(0, y') \tau(y', x)$, where Π_1 (like g_1) contains two nested expectation values.

The correction term, containing a difference of restricted and unrestricted two-point functions, is the remainder $R_1(0, x)$. Now the same procedure can be iterated, beginning with the use of Lemma 2.2 to rewrite the difference of two-point functions, and so on. The details are straightforward but tedious, and lead to the following result. To abbreviate the notation, we write

$$\tilde{C}_{j-1} = \tilde{C}_{j-1}^{(y_j, y_j)}(y'_{j-1}),$$

and

$$I_D^j = I_D(y'_j, y_{j+1}; \tilde{C}_{j-1}).$$

Proposition 2.3. For $N \geq 0$ and for $p < p_c$,

$$\begin{aligned} \tau(0, x) = & \delta_{x,0} + \sum_{n=0}^N (-1)^n g_n(0, x) + \sum_{v \neq 0} p_{0v} \tau(v, x) \\ & + \sum_{n=0}^N (-1)^n \sum_{y'} \Pi_n(0, y') \tau(y', x) + (-1)^{N+1} R_N(0, x), \end{aligned}$$

where

$$g_0(0, x) = \text{Prob}(0 \text{ --- } x) - \delta_{0,x},$$

and for $n \geq 1$,

$$\begin{aligned} g_n(0, x) = & \sum_{(y_1, y'_1)} p_{y_1 y'_1} \cdots \sum_{(y_n, y'_n)} p_{y_n, y'_n} \langle I[0 \text{ --- } y_1] \langle I_D^1 \langle I_D^2 \langle I_D^3 \cdots \langle I_D^{n-1} \\ & \cdot \langle I_D(y'_n, x; \tilde{C}_{n-1}) \rangle_{(n)} \rangle_{(n-1)} \cdots \rangle_{(3)} \rangle_{(2)} \rangle_{(1)} \rangle_{(0)}, \end{aligned}$$

where $y'_0 = 0$. Also, for $n \geq 0$,

$$\Pi_n(0, y') = \sum_y p_{yy'} g_n(0, y).$$

Finally

$$\begin{aligned} R_N(0, x) = & \sum_{(y_1, y'_1)} p_{y_1 y'_1} \cdots \sum_{(y_{N+1}, y'_{N+1})} p_{y_{N+1}, y'_{N+1}} \langle I[0 \text{ --- } y_1] \\ & \cdot \langle I_D^1 \langle I_D^2 \cdots \langle I_D^N \cdot \{\tau(y'_{N+1}, x) - \tau^{\tilde{C}_N}(y'_{N+1}, x)\} \rangle_{(N)} \cdots \rangle_{(2)} \rangle_{(1)} \rangle_{(0)}. \end{aligned}$$

Remark. In the nested expectation

$$\begin{aligned} & \langle I[0 \text{ --- } y_1] \langle I_D^1 \langle I_D(y'_2, x; \tilde{C}_1) \rangle_{(2)} \rangle_{(1)} \rangle_{(0)} \\ = & \langle I[0 \text{ --- } y_1] \langle I_D(y'_1, y_2; \tilde{C}_0^{(y_1, y'_1)}(0)) \langle I_D(y'_2, x; \tilde{C}_1^{(y_2, y'_2)}(y'_1)) \rangle_{(2)} \rangle_{(1)} \rangle_{(0)}, \end{aligned}$$

which occurs in $g_2(0, x)$, it is worth emphasizing again that within the innermost expectation, $\tilde{C}_1^{(y_2, y'_2)}(y'_1)$ represents a deterministic set of sites, which is random in the middle expectation. Similarly $\tilde{C}_0^{(y_1, y'_1)}(0)$ is deterministic in the middle expectation, but random in the outer expectation. The situation is entirely analogous for g_n and R_N .




Fig. 5. Diagrammatic representation for h_0 , h_1 , and h_2 , using the notation introduced in Sect. 1.4

Proposition 2.4. *The quantities g_n , Π_n and R_N in the expansion (given in Proposition 2.3) for the two-point function $\tau(0, x)$ satisfy the following bounds, for $n, N \geq 0$:*

$$\begin{aligned}
 0 &\leq g_n(0, x) \leq h_n(0, x), \\
 0 &\leq \Pi_n(0, y') \leq \sum_y p_{yy'} h_n(0, y), \\
 0 &\leq R_N(0, x) \leq \sum_{(y, y')} p_{yy'} h_N(0, y) \tau(y', x).
 \end{aligned}
 \tag{2.15}$$

Proof. We begin with g_0 and Π_0 . By definition,

$$g_0(0, x) = \text{Prob}(0 \text{ --- } \text{blob} \text{ --- } x) - \delta_{0,x},$$

and by the van den Berg-Kesten inequality the right side is less than $\tau(0, x)^2 - \delta_{0,x} = h_0(0, x)$. This is the desired bound on g_0 . Hence, by definition of Π_0 (2.8 a),

$$\Pi_0(0, y') \leq \sum_y p_{yy'} h_0(0, y).$$

This is the desired bound on Π_0 . The bound on R_0 is similar.

To bound g_n, Π_n and R_N for $n, N \geq 1$, we use the following lemma:

Lemma 2.5.

(a) $\langle I_D(y', x; A) \rangle \leq \sum_u I[u \in A] \text{ --- } \text{blob} \text{ --- } x,$ (2.16)

(b) $\langle I_D(y'_i, y_{i+1}; A) I[v \in \tilde{C}^{(y_{i+1}, y_{i+1})}(y'_i)] \rangle$
 $\leq \sum_u I[u \in A] \left(\text{blob} + \text{blob} \right).$ (2.17)

The proof of Lemma 2.5 is deferred to the end of this section. Now we show how the lemma can be used to complete the proof of Proposition 2.4. The basic

idea is simple: the nested expectations given for $g_n, \Pi_n,$ and R_N in Proposition 2.3 are estimated using Lemma 2.5, working from the inside out.

We begin with $g_n,$ and consider the formula for g_n given in Proposition 2.3. We first estimate the innermost expectation $\langle I_D \rangle_{(n)}$ using Lemma 2.5 (a). (Within this expectation, $\tilde{C}_{n-1}^{(y_n, y'_n)}(y'_{n-1})$ is deterministic; it is random within the next expectation.) This gives

$$\langle I_D(y'_n, x; \tilde{C}_{n-1}^{(y_n, y'_n)}(y'_{n-1})) \rangle_{(n)} \leq \sum_{u_n} I[u_n \in \tilde{C}_{n-1}^{(y_n, y'_n)}(y'_{n-1})] \cdot \left(\begin{array}{c} u_n \\ \text{---} \\ y'_n \text{---} \end{array} \rightarrow x \right).$$

Then we estimate the next expectation (which now contains a factor $I[u_n \in \tilde{C}_{n-1}^{(y_n, y'_n)}(y'_{n-1})]$ due to the above bound on $\langle I_D \rangle_{(n)}$ using Lemma 2.5 (b). This gives

$$\begin{aligned} & \langle I_D(y'_{n-1}, y_n; \tilde{C}_{n-2}^{(y_{n-1}, y'_{n-1})}(y'_{n-2})) I[u_n \in \tilde{C}_{n-1}^{(y_n, y'_n)}(y'_{n-1})] \rangle_{(n-1)} \\ & \leq \sum_{u_{n-1}} I[u_{n-1} \in \tilde{C}_{n-2}^{(y_{n-1}, y'_{n-1})}(y'_{n-2})] \cdot \left[\begin{array}{c} y'_{n-1} \text{---} u_n \\ \text{---} \\ u_{n-1} \text{---} y_n \end{array} + \begin{array}{c} y'_{n-1} \text{---} u_n \\ \text{---} \\ u_{n-1} \text{---} y_n \end{array} \right] \end{aligned}$$

Repeating this process gives the estimate for g_n given in Proposition 2.4.

The bound on Π_n then follows from the fact (Proposition 2.3) that

$$\Pi_n(0, y') = \sum_y p_{yy'} g_n(0, y).$$

Finally, we bound R_N using the formula given in Proposition 2.3 and the inequality

$$0 \leq \tau(y', x) - \tau^A(y', x) \leq \tau(y', x) \tag{2.18}$$

to obtain

$$R_N(0, x) \leq \sum_{y_{N+1}} \Pi_N(0, y'_{N+1}) \tau(y'_{N+1}, x). \quad \square$$

This completes the proof of Proposition 2.4, given Lemma 2.5. Now we give the proof of Lemma 2.5.

Proof of Lemma 2.5.

(a) By definition (Eq. 2.9, see also Fig. 4),

$$\begin{aligned} \langle I_D(y', x; A) \rangle &= \langle I[y' \text{ and } x \text{ are doubly connected through } A] \rangle \\ &+ \langle I[\exists z \neq y' \text{ such that } z \text{ is connected to } y' \text{ in } \mathbb{Z}^d \setminus A, \\ &z \text{ is an endpoint of a pivotal bond for the connection of } y' \text{ and } x, \\ &\text{and } z \text{ is doubly connected to } x \text{ through } A] \rangle. \end{aligned}$$

For any bond configuration contributing to the first term on the right side, there exists a site $u \in A$ and three *distinct* occupied paths connecting y' and x, y' and $u,$ and x respectively. Therefore by the van den Berg-Kesten inequality the first term on the right side is bounded above by

$$\sum_{u \in A} \tau(y', u) \tau(u, x) \tau(x, y').$$

Furthermore, for any bond configuration contributing to the second term on the right side of (2.19), there is a site $u \in A$ and four *distinct* paths connecting y' and z , z and x , z and u , u and x . Note that $z \neq y'$. Thus by the van den Berg-Kesten inequality the second term is bounded above by

$$\sum_{u \in A} \sum_{\substack{z \\ z \neq y'}} \tau(y', z) \tau(z, x) \tau(z, u) \tau(u, x).$$

Combining these two gives (a).

(b) By definition of I_D ,

$$\begin{aligned} &\langle I_D(y'_i, y_{i+1}; A) I[v \in \tilde{C}^{(y_{i+1}, y'_{i+1})}(y'_i)] \rangle \\ &\leq \langle I[y_{i+1} \text{ and } y'_i \text{ are doubly connected through } A, \text{ and } v \text{ is connected to } y'_i] \rangle \\ &\quad + \langle I[\exists \text{ a last pivotal bond } (z, z') \text{ for the connection from } y'_i \text{ to } y_{i+1}, z' \text{ and } \\ &\quad y_{i+1} \text{ are doubly connected through } A, z \text{ and } y'_i \text{ are connected in } \mathbb{Z}^d \setminus A, \\ &\quad \text{and } v \text{ is connected to } y'_i] \rangle. \end{aligned} \tag{2.20}$$

We first consider the case $y'_i \neq y_{i+1}$. For a configuration contributing to the first term on the right side of (2.20), there are sites $u \in A$ and $w \in \mathbb{Z}^d$ and five distinct occupied paths joining y'_i to u , u to y_{i+1} , y'_i to w , w to y_{i+1} , and w to v . (Since y_{i+1} and y'_i are doubly connected through A , each path connecting y'_i to y_{i+1} goes through A , and hence $u \in A$ can be chosen such that w and u lie on distinct paths from y'_i to y_{i+1} .) Thus by the van den Berg-Kesten inequality the first term on the right side of (2.20) is bounded above by

$$\sum_{u \in A} \left(\begin{array}{c} y'_i \quad v \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ u \quad y_{i+1} \end{array} \right). \tag{2.21 a}$$

The second term can be bounded similarly by

$$\begin{aligned} &\text{Prob} \left[\exists z' \in \mathbb{Z}^d, u \in A \text{ such that } \begin{array}{c} y'_i \quad z' \quad v \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ u \quad y_{i+1} \end{array} \text{ or } \begin{array}{c} y'_i \quad v \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ u \quad y_{i+1} \end{array} \right] \\ &\leq \sum_{\substack{u \in A \\ z' \in \mathbb{Z}^d}} \left[\begin{array}{c} y'_i \quad z' \quad v \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ u \quad y_{i+1} \end{array} + \begin{array}{c} y'_i \quad v \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ u \quad y_{i+1} \end{array} \right], \end{aligned} \tag{2.21 b}$$

using the van den Berg-Kesten inequality in the last step. In (2.21 b), $y'_i \neq z'$, because z' is the latter endpoint of a pivotal bond. Combining these two estimates gives (2.17) for the case $y'_i \neq y_{i+1}$, apart from checking that the loop in the first term on the right side of (2.21 b) cannot shrink to a point. However, the case where that loop skrink to a point is already accounted for in the second term of (2.21 b). (The loop in (2.21 a) cannot shrink to a point when $y'_i \neq y_{i+1}$.) This proves (2.17) for the case when $y'_i \neq y_{i+1}$.

When $y'_i = y_{i+1}$, it follows from the definition of I_D that

$$\begin{aligned} \langle I_D(y'_i, y_{i+1}; A) I[v \in \tilde{C}^{(y_{i+1}, y'_{i+1})}(y'_i)] \rangle &\leq I[y'_i \in A] \langle I[v \in \tilde{C}^{(y_{i+1}, y'_{i+1})}(y'_i)] \rangle \\ &\leq I[y'_i \in A] \text{Prob}(y'_i \text{---} v) \leq I[y'_i \in A] \tau(y'_i, v). \end{aligned} \tag{2.22}$$

However the upper bound (2.22) is bounded above by the second term in (2.21 b) (which includes it as the case $y'_i = z' = u = y_{i+1}$). Thus (2.17) holds also in the case $y'_i = y_{i+1}$. \square

2.3. Bounds in k -Space for Each Term of the Expansion

In this section we obtain bounds on $\hat{g}_n(k)$, $\hat{\Pi}_n(k)$ and $\hat{R}_N(k)$ which will be used in Sects. 4 and 5 in deriving the infrared bound. These bounds are given in Proposition 2.6, and follow in a straightforward way from the x -space bounds obtained in Sect. 2.2. In the statement of the bounds, derivatives with respect to k_μ are denoted by ∂_μ , and $|x|$ ($\equiv \|x\|_2$) is the Euclidean distance from x to the origin.

Proposition 2.6. *The following bounds are satisfied:*

$$\begin{aligned} |\hat{g}_n(k)| &\leq \sum_x h_n(0, x), \\ |\partial_\mu^s \hat{g}_n(k)| &\leq d^{-1} \sum_x |x|^2 h_n(0, x), \quad s = 1, 2, \\ |\hat{\Pi}_n(k)| &\leq \sum_v p_{0v} \sum_x h_n(0, x), \\ |\partial_\mu^s \hat{\Pi}_n(k)| &\leq d^{-1} \left[\sum_v p_{0v} \sum_x |x|^2 h_n(0, x) + \sum_v p_{0v} |v|^2 \sum_x h_n(0, x) \right], \quad s = 1, 2, \\ 0 \leq \hat{\Pi}_n(0) - \hat{\Pi}_n(k) &\leq \frac{|k|^2}{2d} \left[\sum_v p_{0v} \sum_x |x|^2 h_n(0, x) + \sum_v p_{0v} |v|^2 \sum_x h_n(0, x) \right], \\ |\hat{R}_N(k)| &\leq \sum_v p_{0v} \chi(p) \sum_x h_N(0, x), \\ |\partial_\mu^s \hat{R}_N(k)| &\leq d^{-1} \chi(p) \left[\sum_v p_{0v} \sum_x |x|^2 h_N(0, x) + \sum_v p_{0v} |v|^2 \sum_x h_N(0, x) \right] \\ &\quad + d^{-1} \sum_v p_{0v} \sum_x h_n(0, x) \sum_y |y|^2 \tau(0, y), \quad s = 1, 2. \end{aligned}$$

Proof. These bounds all follow in a straightforward way from the x -space bounds given in Proposition 2.4. Beginning first with the bounds which do not involve derivatives, it follows from Proposition 2.4 that

$$\begin{aligned} |\hat{g}_n(k)| &= \left| \sum_x e^{ik \cdot x} g_n(0, x) \right| \leq \sum_x g_n(0, x) \leq \sum_x h_n(0, x), \\ |\hat{\Pi}_n(k)| &= \left| \sum_{y'} e^{ik \cdot y'} \Pi_n(0, y') \right| \leq \sum_{y'} \Pi_n(0, y') \leq \sum_v p_{0v} \sum_y h_n(0, y), \end{aligned}$$

and

$$|\hat{R}_N(k)| = \left| \sum_x e^{ik \cdot x} R_N(0, x) \right| \leq \sum_x R_N(0, x) \leq \sum_v p_{0v} \chi(p) \sum_x h_N(0, x).$$

To obtain the bound on $\partial_\mu^s \hat{g}_n(k)$, $s = 1, 2$, we use symmetry and Proposition 2.4 as follows:

$$\begin{aligned} |\partial_\mu^s \hat{g}_n(k)| &= \left| \sum_x (ix_\mu)^s e^{ik \cdot x} g_n(0, x) \right| \\ &\leq \sum_x x_\mu^2 g_n(0, x) = d^{-1} \sum_x |x|^2 g_n(0, x) \leq d^{-1} \sum_x |x|^2 h_n(0, x). \end{aligned}$$

The bound on $\partial_\mu^s \hat{I}_n(k)$ is obtained in much the same way. We first note that

$$|\partial_\mu^s \hat{I}_n(k)| \leq d^{-1} \sum_{y'} |y'|^2 I_n(0, y') \leq d^{-1} \sum_{y,v} p_{0v} |y+v|^2 h_n(0, y).$$

By symmetry the cross term $2y \cdot v$ gives no contribution to the sum on the right side, and hence

$$|\partial_\mu^s \hat{I}_n(k)| \leq d^{-1} \sum_{y,v} p_{0v} (|y|^2 + |v|^2) h_n(0, y).$$

The bounds on $|\partial_\mu^s \hat{R}_N(k)|$ can be obtained in a similar fashion.

Finally, again using symmetry, we find

$$\begin{aligned} \hat{I}_n(0) - \hat{I}_n(k) &= \sum_{y'} [1 - \cos k \cdot y'] I_n(0, y') \leq \frac{1}{2} \sum_{y'} \left(\sum_{\mu=1}^d k_\mu y'_\mu \right)^2 \sum_y p_{yy'} h_n(0, y) \\ &= \frac{1}{2} \sum_{\mu,v} k_\mu k_v \sum_{y,v} p_{0v} (y_\mu + v_\mu) (y_v + v_v) h_n(0, y) \\ &= \frac{1}{2} \sum_{\mu=1}^d k_\mu^2 \sum_{y,v} p_{0v} (|y_\mu|^2 + |v_\mu|^2) h_n(0, y) \\ &= \frac{|k|^2}{2d} \sum_{y,v} p_{0v} (|y|^2 + |v|^2) h_n(0, y). \end{aligned} \tag{2.23}$$

Note that for the nearest neighbour model this expression gives the upper bound

$$\hat{I}_n(0) - \hat{I}_n(k) \leq \frac{\pi^2}{4} (1 - D(k)) 2dp \sum_y (|y|^2 + 1) h_n(0, y), \tag{2.24}$$

since

$$1 - D(k) \equiv \frac{1}{d} \sum_{\mu=1}^d (1 - \cos k_\mu) \geq 2\pi^{-2} \frac{|k|^2}{d}.$$

The corresponding expression for the spread-out models will be given in Sect. 5. It follows immediately from the first equality in (2.23) that the left side is nonnegative.

This completes the proof of Proposition 2.6. \square

3. Diagrammatic Estimates

In the last section bounds were obtained for $\partial_\mu^s \hat{g}_n(k)$, $\partial_\mu^s \hat{\Pi}_n(k)$ and $\partial_\mu^s \hat{R}_N(k)$, $s = 0, 1, 2$, in terms of $\sum_x h_n(0, x)$ and $\sum_x |x|^2 h_n(0, x)$, for any \mathbb{Z}^d -invariant model.


The definition of $h_n(0, x)$ is given in Eqs. (2.10) and (2.11).

In this section we derive upper bounds (also valid in any \mathbb{Z}^d -invariant model) on $\sum_x h_n(0, x)$ and $\sum_x |x|^2 h_n(0, x)$, in terms of quantities \bar{T} , \bar{W} and \bar{H} which are introduced in the next definition. These quantities will figure in the precise version of (1.18) and for the nearest neighbour model should be thought of as being $O(d^{-1})$ uniformly in $p < p_c$. Indeed we will prove that such uniform bounds hold in Sect. 4. (Related estimates which show that \bar{T} , \bar{W} and \bar{H} go to zero as L approaches infinity will be obtained in Sect. 5 for the spread-out models.)

Definition 3.1. For a, a_1 , and a_2 in \mathbb{Z}^d we define

$$T_a = \sum_{x,y} \tau(0, x) \tau(x, y) \tau(y, a) - \delta_{a,0}, \tag{3.1}$$


$$W_a = \sum_x |x|^2 \tau(0, x) \tau(x, a), \tag{3.2}$$

$$H_{a_1, a_2} = \sum_{x,y,z} |x|^2 \tau(0, x) \tau(x, y) \tau(x, u) \tau(0, u) \tau(0, z) \tau(u, v) \tau(v, y + a_2) \tau(v, z + a_1). \tag{3.3}$$


We write T_0 and W_0 simply as T and W . We also define

$$\bar{T} = \sup_{a \in \mathbb{Z}^d} T_a, \quad \bar{W} = \sup_{a \in \mathbb{Z}^d} W_a, \quad \bar{H} = \sup_{a_1, a_2 \in \mathbb{Z}^d} H_{a_1, a_2}. \tag{3.4}$$

Finally we define

$$W'_a = \sum_x |x|^2 \tau(0, x+a) \tau(y, x) \equiv \sum_{x,y} p_{0y} |x|^2 \tau(0, x+a) \tau(y, x)$$


and

$$\bar{W}' = \sup_a W'_a.$$

We simply write W' for W'_0 .

In Sects. 3.1 and 3.2 the following lemma is proved.

Lemma 3.2.

(a)
$$0 \leq \sum_x h_0(0, x) \leq \frac{T}{3},$$

and for $n \geq 1$,

$$0 \leq \sum_x h_n(0, x) \leq (1 + T)^2 \left[\sum_v p_{0v} \frac{T}{3} + 2 \left(\sum_v p_{0v}^2 T/3 \right)^{1/2} + 2\bar{p} \right] r^{n-1},$$

where

$$\bar{p} = \sup_v p_{0v}$$

and

$$r = (1 + T + \bar{T}) \left[\bar{p} + \sum_v p_{0v} \bar{T} \right].$$

$$(b) \quad 0 \leq \sum_x |x|^2 h_0(0, x) \leq W,$$

$$0 \leq \sum_x |x|^2 h_1(0, x) \leq W' + 6T(1 + T) \bar{W}' + 6\bar{W}(1 + T) \left[\bar{p} + \sum_v p_{0v} \bar{T} \right],$$

and for $n \leq 1$,

$$0 \leq \sum_x |x|^2 h_n(0, x) \leq (2n + 1) \left(\frac{n}{2} \sum_v p_{0v} + (n + 1)r \right) (1 + T)^2 r^{n-1} \bar{W} \\ + (2n + 1) \left[\frac{n}{2} \right] (1 + T)^2 r^{n-1} \bar{W}' + (2n + 1) \left[\frac{n-1}{2} \right] (1 + T)^2 r^{n-2} \bar{H} \left(\sum_v p_{0v} \right)^2,$$

where $\lceil x \rceil$ denotes the largest integer which does not exceed x .

Remark. The exact form of the upper bounds in the lemma is not important unless a good value of d_0 is sought. Otherwise the relevant feature, for the nearest-neighbour model, of these bounds is that under the assumption that \bar{T} , \bar{W} , \bar{W}' , \bar{H} are all $O(d^{-1})$ and $2dp$ is $O(1)$, the bounds imply

$$(a) \quad 0 \leq \sum_x h_n(0, x) \leq \begin{cases} O(d^{-1}) & n = 0 \\ O(d^{-1})^n & n = 1, 2, \dots \end{cases} \text{ and}$$

$$(b) \quad 0 \leq \sum_x |x|^2 h_n(0, x) \leq \begin{cases} O(d^{-1}) & n = 0, 1 \\ n^2 O(d^{-1})^{n-1} & n = 2, 3, 4, \dots \end{cases}$$

(The $n \geq 2$ bound in (b) is believed not to be optimal, but it does suffice for our purposes.) An analogous remark applies for the spread-out models, as described in Sect. 5.

The proof of Lemma 3.2 (a) is given in Sect. 3.1 and the proof of part (b) in Sect. 3.2. Finally in Sect. 3.3 we prove the following bounds on T_a and W'_a :

Lemma 3.3.

$$T_a \leq T + (3T/2d)^{1/2}, \\ W' \equiv W'_0 \leq \sum_v p_{0v} W + \sum_v p_{0v} |v| (WT/3)^{1/2} + \left(\sum_v p_{0v}^2 |v|^2 W \right)^{1/2}, \\ W'_a \leq 2 \sum_v p_{0v} \bar{W} + 2 \sum_v p_{0v} |v|^2 \frac{T}{3} + 4 \left(\sum_v p_{0v}^2 |v|^4 \frac{T}{3} \right)^{1/2} + 4p_{0a} |a|^2.$$

3.1. Bounds on $\sum_x h_n(0, x)$

In this and the next section we make extensive use of the graphical notation introduced in Sect. 1.4 and the graphical representation for h_n given in Eqs.

(2.11–14) and Fig. 5. This section is devoted to the proof of Lemma 3.2(a). The proof uses translation invariance and repeated application of the simple inequality

$$\left| \sum_x f(x) g(x) \right| \leq \sup_x |f(x)| \sum_x |g(x)|. \tag{3.5}$$

This inequality is used to reduce $\sum_x h_n(0, x)$ to a product of basic units.

The case $n = 0$ is the simplest. We just observe that

$$T = \sum_{x,y} \tau(0, x) \tau(x, y) \tau(y, 0) - 1 \geq 3 \sum_{x \neq 0} \tau(0, x) \tau(x, 0),$$

and hence

$$\sum_x h_0(0, x) = \sum_{x \neq 0} \tau(0, x)^2 \leq \frac{T}{3}.$$

For $n \geq 1$ we use translation invariance to rewrite (2.11) as

$$\sum_x h_n(0, x) = \sum_{a_1, \dots, a_n} \tilde{A}_1(a_1) \prod_{i=2}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n), \tag{3.6}$$

where

$$\begin{aligned} \tilde{A}_1(a) &= \sum_x A_1(x, 0, a) = \text{[Diagram: A jagged path from 0 to a]}, \\ \tilde{A}_2(a, a') &= \sum_x A_2(0, a, x, x + a') = \sum_x \left[\text{[Diagram: Two jagged paths from 0 to x and x+a']} \right] + \left[\text{[Diagram: A square loop from 0 to x+a']} \right], \\ \tilde{A}_3(a) &= \sum_x A_3(0, a, x) = \text{[Diagram: A jagged path from 0 to a]}. \end{aligned} \tag{3.7}$$


Then we apply (3.5) repeatedly to obtain


$$\sum_x h_n(0, x) \leq \left(\sup_a \tilde{A}_1(a) \right) \left(\sup_{a'} \left\{ \sum_a \tilde{A}_2(a, a') \right\} \right)^{n-1} \left(\sum_a \tilde{A}_3(a) \right). \tag{3.8}$$

The last factor on the right side of (3.8) is $1 + T$. To complete the proof of Lemma 3.2(a) we use (3.8) and the following lemma.

Lemma 3.4. *The following inequalities hold for any $a \in \mathbb{Z}^d$:*

- (a) $\text{[Diagram: Jagged path from 0 to a]} \leq \sum_v p_{0v} T/3 + 2 \left(\sum_v p_{0v}^2 T/3 \right)^{1/2} + 2\bar{p},$
- (b) $\text{[Diagram: Square loop from 0 to a]} \leq \sum_v p_{0v} \bar{T} + \bar{p},$

(c)  $\leq (1 + T) \left[\sum_v p_{0v} T/3 + 2 \left(\sum_v p_{0v}^2 T/3 \right)^{1/2} + 2\bar{p} \right],$

(d)  $\leq (1 + T) \left[\sum_v p_{0v} \bar{T} + \bar{p} \right],$

(e)  $\leq \bar{T} \left[\sum_v p_{0v} \bar{T} + \bar{p} \right].$

Proof. (a) Using the Schwarz inequality and the inequality used to bound $\sum_x h_0(0, x)$ we obtain

$$\begin{aligned} \text{Diagram} &\equiv \sum_{x,v} p_{0v} \tau(v, x) \tau(x, a) = \sum_v p_{0v} \left[\sum_{x \neq v, a} \tau(v, x) \tau(x, a) + 2\tau(v, a) \right] \\ &\leq \sum_v p_{0v} T/3 + 2 \sum_{v \neq a} p_{0v} \tau(v, a) + 2p_{0a} \\ &\leq \sum_v p_{0v} T/3 + 2 \left(\sum_v p_{0v}^2 T/3 \right)^{1/2} + 2\bar{p}. \end{aligned}$$

(b) Using translation invariance the left side is equal to

$$\sum_v p_{0v} \sum_{x,y} \tau(-v, x) \tau(x, y) \tau(y, a).$$

By considering separately in the sum over v the term $v = -a$, we obtain the upper bound

$$\sum_{v \neq a} p_{0v} \bar{T} + p_{0a}(1 + T) \leq \sum_v p_{0v} \bar{T} + \bar{p}.$$

(c) By (3,5),

$$\text{Diagram} \leq \sup_a \left[\sum_x \begin{matrix} x+a' & a \\ \text{Diagram} \\ x & 0 \end{matrix} \right].$$

The first factor on the right side is equal to $1 + T$, and the second can be bounded using (a) and translation invariance.

(d) and (e) are proved similarly. \square

Now the desired bound on $\sum_x h_n(0, x)$ follows immediately from (3.8) and (c–e), and the proof of Lemma 3.2 (a) is complete. \square

3.2. Bounds on $\sum_x |x|^2 h_n(0, x)$

In this section we prove Lemma 3.2(b). The proof is similar to the proof of Lemma 3.2(a) given in the last section, although the situation here is more complicated due to the presence of the factor $|x|^2$.

The case $n = 0$ is the simplest, since by definition of W

$$\sum_x |x|^2 h_0(0, x) = \sum_x |x|^2 \tau(0, x)^2 = W.$$

For higher order terms the basic idea is to use (3.5) together with the triangle inequality, as we now illustrate for the case $n = 1$. By definition of h_1 ,

$$\sum_x |x|^2 h_1(0, x) = \sum_{z, u, x} |x|^2 \text{0} \begin{array}{c} z \quad u \\ \text{shaded triangles} \end{array} x \tag{3.9}$$

and by the triangle inequality

$$|x|^2 \leq (|x - u| + |u - z| + |z|)^2 \leq 3(|x - u|^2 + |u - z|^2 + |z|^2). \tag{3.10}$$

We insert (3.10) into (3.9) and consider separately the case where both of the shaded triangles in (3.9) are points. In this special case (3.9) contributes W' , and hence

$$\sum_x |x|^2 h_1(0, x) \leq W' + 3 \sum_{z, u, x} (|x - u|^2 + |u - z|^2 + |z|^2) \text{0} \begin{array}{c} z \quad u \\ \text{shaded triangles} \end{array} x$$

where at least one of the shaded triangles on the right side is not a point. Each of the three terms in the summation on the right side is now treated separately, using (3.5), translation invariance and Lemma 3.4. For example,

$$\begin{aligned} \sum_{z, u, x} |z|^2 \text{0} \begin{array}{c} z \quad u \\ \text{shaded triangles} \end{array} x &\leq \sup_a \left(\sum_v |v|^2 v \begin{array}{c} a \\ \text{shaded triangle} \end{array} \right) \left(\text{0} \begin{array}{c} \text{shaded triangle} \end{array} \right) \\ &\leq \bar{W} \sup_a \left(\begin{array}{c} a \\ \text{shaded triangle} \end{array} \right) (1 + T) \leq \bar{W} \left[\sum_v p_{0v} \bar{T} + \bar{p} \right] (1 + T). \end{aligned}$$

The term involving $|x - u|^2$ is handled similarly, and satisfies the same upper bound. The remaining term is

$$\begin{aligned} \sum_x |x|^2 \text{0} \begin{array}{c} x \\ \text{shaded triangle} \end{array} &\leq 2 \left(\begin{array}{c} \text{shaded triangle} \\ 0 \end{array} \right) \left(\begin{array}{c} \text{shaded triangle} \\ 0 \end{array} \right) \sup_{a_1, a_2} \left(\sum_x |x|^2 \begin{array}{c} 0 \text{---} x \\ a_1 \text{---} x + a_2 \end{array} \right) \\ &\leq 2(1 + T) T \bar{W}', \end{aligned}$$

where we have used the fact that one of the two triangles is not a point. As a result,

$$\sum_x |x|^2 h_1(0, x) \leq W' + 6T(1 + T) \bar{W}' + 6\bar{W}(1 + T) \left[\sum_v p_{0v} \bar{T} + \bar{p} \right],$$

which is the desired bound.

The case $n \geq 2$ can be handled in a similar fashion, although the additional factors of \tilde{A}_2 make the analysis more involved. It is at this stage that it becomes necessary to work with the quantity \bar{H} , as we are unable to bound the diagrams encountered solely in terms of \bar{T} and \bar{W} .

It is convenient to use the following expression for $h_n(0, x)$, which is equivalent to (2.11). To write the expression we recall the definition of $A_3(u, v, x)$ in (2.14) and define

$$B_1(s, t, u, v) = \begin{array}{c} t \text{ --- } \text{~~~~} v \\ | \\ s \text{ --- } \text{~~~~} u \end{array} \tag{3.11}$$

and

$$B_2(u, v, s, t) = \begin{array}{c} s \\ | \\ \text{triangle} \\ | \\ u \end{array} \delta_{v,s} + \begin{array}{c} v \text{ --- } \text{~~~~} s \\ | \\ u \text{ --- } \text{~~~~} t \end{array} \tag{3.12}$$

Then (2.11) can be rewritten

$$\begin{aligned} h_n(0, x) &= \sum_{\substack{s_k, t_k \\ u_k, v_k}} A_3(0, s_1, t_1) \prod_{i=1}^n B_1(s_i, t_i, u_i, v_i) \prod_{i=2}^n B_2(u_{i-1}, v_{i-1}, s_i, t_i) A_3(u_n, v_n, x) \\ &\equiv \sum h_n(0, \vec{s}, \vec{t}, \vec{u}, \vec{v}, x). \end{aligned} \tag{3.13}$$

To bound $\sum_x |x|^2 h_n(0, x)$ we use the triangle inequality as for the case $n = 1$:

$$\begin{aligned} |x|^2 &\leq (2n + 1) (|t_1|^2 + |v_1 - t_1|^2 + |s_2 - v_1|^2 + \dots \\ &\quad + \{[|s_n - u_n|^2 + |u_n - x|^2] \text{ or } [|t_n - v_n|^2 + |v_n - x|^2]\}), \end{aligned} \tag{3.14}$$

where in the term in brace brackets in (3.14) the first expression is used if n is even and the second if n is odd.

The effect of each term on the right side of (3.14) is treated separately, and we proceed term by term. Beginning with the $|t_1|^2$ term, and using an unlabelled sum to denote summation over all vertices, a minor change in the organization gives

$$\begin{aligned} &(2n + 1)^{-1} \sum |t_1|^2 h_n(0, \vec{s}, \vec{t}, \vec{u}, \vec{v}, x) \\ &\leq \sum \left(|t_1|^2 \begin{array}{c} 0 \\ | \\ \text{triangle} \\ | \\ t_1 + a_0 \end{array} \right) \left(\begin{array}{c} 0 \text{ --- } \text{~~~~} y \\ | \\ a_0 \text{ --- } \text{~~~~} y + a_1 \end{array} \right) \prod_{i=2}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n) \\ &\leq \bar{W} \sup_{a_1} \left(\begin{array}{c} 0 \\ | \\ \text{triangle} \\ | \\ a_1 \end{array} \right) \left(\sup_{a'} \sum_a \tilde{A}_2(a, a') \right)^{n-1} \sum_a \tilde{A}_3(a). \end{aligned} \tag{3.15}$$

Now Lemma 3.4 applies as before.

The $|v_1 - t_1|^2$ term is treated as follows:

$$\begin{aligned}
 & (2n+1)^{-1} \sum |v_1 - t_1|^2 h_n(0, \vec{s}, \vec{t}, \vec{u}, \vec{v}, x) \\
 & \leq \sum \left(\begin{array}{c} 0 \\ \text{[Diagram: square with shaded bottom-left corner]} \\ a_0 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ a_0 \text{ [Diagram: wavy line]} x+a_1 \\ |x|^2 \end{array} \right) \prod_{i=2}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n) \\
 & \leq \sum_{a_0} \tilde{A}_3(a_0) \bar{W}' \left(\sup_{a'} \sum_a \tilde{A}_2(a, a') \right)^{n-1} \sum_a \tilde{A}_3(a), \tag{3.16}
 \end{aligned}$$

and again Lemma 3.4 can be applied.

The term involving $|s_2 - v_1|^2$ can be organized as follows:

$$\begin{aligned}
 & (2n+1)^{-1} \sum |s_2 - v_1|^2 h_n(0, \vec{s}, \vec{t}, \vec{u}, \vec{v}, x) \\
 & = \sum \left(\begin{array}{c} 0 \\ \text{[Diagram: square with shaded bottom-left corner]} \\ a_0 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} y \\ a_0 \text{ [Diagram: wavy line]} y+a_1 \\ |z|^2 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} z \\ a_1 \text{ [Diagram: wavy line]} z+b \\ |z|^2 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ b \text{ [Diagram: wavy line]} x+a_2 \end{array} \right) \\
 & \cdot \prod_{i=3}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n) \\
 & \leq \left(\sum_a \tilde{A}_3(a) \right)^2 \left(\sup_{a'} \sum_{a'} \tilde{A}_2(a, a') \right)^{n-2} \left(\sup_{a_0} \begin{array}{c} 0 \text{ [Diagram: wavy line]} \\ a_0 \text{ [Diagram: wavy line]} \end{array} \right) \bar{W} \left(\sup_{a_2} \begin{array}{c} 0 \text{ [Diagram: wavy line]} \\ a_2 \text{ [Diagram: wavy line]} \end{array} \right)
 \end{aligned}$$

Now Lemma 3.4 can be applied.

To estimate the term involving $|u_2 - s_2|^2$ we define

$$\tilde{A}'_2(a, a') = \sum_x \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ a \text{ [Diagram: wavy line]} x+a' \\ \text{[Diagram: square with shaded bottom-left corner]} \end{array} + \begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ a \text{ [Diagram: wavy line]} x+a' \\ \text{[Diagram: square with shaded top-right corner]} \end{array} \right),$$

and write

$$\begin{aligned}
 & (2n+1)^{-1} \sum |u_2 - s_2|^2 h_n(0, \vec{s}, \vec{t}, \vec{u}, \vec{v}, x) \\
 & \leq \sum \tilde{A}_3(a_0) \tilde{A}'_2(a_0, a_1) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ a_1 \text{ [Diagram: wavy line]} a_2+x \\ |x|^2 \end{array} \right) \prod_{i=3}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n) \\
 & \leq \bar{W} \sum_v p_{0v} \left(\sum_a \tilde{A}_3(a) \right)^2 \left(\sup_{a'} \sum_{a'} \tilde{A}'_2(a, a') \right) \left(\sup_{a'} \sum_a \tilde{A}_2(a, a') \right)^{n-2}.
 \end{aligned}$$

Now Lemma 3.4 can be applied.

Finally we treat the term $|u_2 - t_3|^2$, which is the most complicated one, and in which we use \bar{H} . The contribution due to the term involving the second term on the right side of (3.12) is equal to

$$\begin{aligned}
 & \sum \tilde{A}_3(a_0) \tilde{A}'_2(a_0, a_1) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} x \\ a_1 \text{ [Diagram: wavy line]} x+a_2 \\ |y|^2 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} y \\ a_2 \text{ [Diagram: wavy line]} y+b \\ |z|^2 \end{array} \right) \left(\begin{array}{c} 0 \text{ [Diagram: wavy line]} z \\ b \text{ [Diagram: wavy line]} z+a_3 \end{array} \right) \\
 & \cdot \prod_{i=4}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n),
 \end{aligned}$$

which can be bounded above as was done for the $|s_2 - v_1|^2$ term. The contribution due to the first term on the right side of (3.12) is equal to

$$\begin{aligned} & \sum \tilde{A}_3(a_0) \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right) \\ & \cdot \left[\begin{array}{c} \text{Diagram 3} \\ |u_2 - t_3|^2 \end{array} \right] \cdot \prod_{i=4}^n \tilde{A}_2(a_{i-1}, a_i) \tilde{A}_3(a_n) \\ & \leq \bar{H} \left(\sum_v p_{0v} \right)^2 \left(\sum_a \tilde{A}_3(a) \right)^2 \left(\sup_a \sum_{a'} \tilde{A}'_2(a, a') \right) \left(\sup_{a'} \sum_a \tilde{A}_2(a, a') \right)^{n-3}, \end{aligned}$$

and again Lemma 3.4 can be applied.

This illustrates the method for bounding $\sum_x |x|^2 h_n(0, x)$. By applying Lemma 3.4 and carefully counting the number of terms of each type which occurs, the upper bound in the statement of the lemma can be obtained. We omit the straightforward but tedious details. \square

3.3. Proof of Lemma 3.3

In this section we prove Lemma 3.3. We begin with the inequality for T_a . Fix $a \neq 0$. Using the fact that $\hat{\tau}(k) \geq 0$ [6],

$$\begin{aligned} T_a &= (2\pi)^{-d} \int d^d k [\hat{\tau}(k)^3 - 1] e^{ik \cdot a} \\ &= (2\pi)^{-d} \int d^d k [\hat{\tau}(k) - 1]^2 [\hat{\tau}(k) + 2] e^{ik \cdot a} + 3\tau(0, a) \\ &\leq (2\pi)^{-d} \int d^d k [\hat{\tau}(k) - 1]^2 [\hat{\tau}(k) + 2] + 3\tau(0, a) = T + 3\tau(0, a). \end{aligned}$$

Now for $a \neq 0$ it follows by symmetry that there are at least $2d$ terms which give the same contributions as $\tau(0, a)^2$ in

$$B \equiv \sum_{x \neq 0} \tau(0, x)^2 \leq T/3.$$

This leads to the desired inequality for T_a .

To obtain the inequality for W' we argue using the Schwarz inequality as follows:

$$\begin{aligned} W' &\equiv \sum_{x,y} p_{0y} |x|^2 \tau(0, x) \tau(x, y) \leq \sum_y p_{0y} \sum_x \tau(0, x) \tau(x, y) [|x| |x - y| + |x| |y|] \\ &\leq \sum_y p_{0y} W + \sum_y p_{0y} \left(\sum_{x \neq y} \tau(0, x) \tau(y, x) |x| |y| + \tau(0, y) |y|^2 \right) \\ &= \sum_y p_{0y} W + \sum_y p_{0y} |y| \sum_{x \neq y} \tau(0, x) |x| \tau(x, y) + \sum_y p_{0y} |y| |y| \tau(0, y) \\ &\leq \sum_y p_{0y} W + \sum_y p_{0y} |y| (WB)^{1/2} + \left(\sum_y p_{0y}^2 |y|^2 W \right)^{1/2}. \end{aligned}$$

Finally, the bound on W'_a is obtained similarly:

$$\begin{aligned}
 W'_a &\equiv \sum_{x,y} p_{0y} |x|^2 \tau(y, x) \tau(0, x+a) \\
 &\leq 2 \sum_{x,y} p_{0y} \tau(y, x) \tau(0, x+a) [|x-y|^2 + |y|^2] \\
 &= 2 \sum_{x,y} p_{0y} \tau(0, x) \tau(0, x+a+y) [|x|^2 + |y|^2] \\
 &= 2 \sum_y p_{0y} \sum_x \tau(0, x) \tau(0, x+a+y) |x|^2 \\
 &\quad + 2 \sum_y p_{0y} |y|^2 \left[\sum_{x \neq 0, -y-a} \tau(0, x) \tau(0, x+a+y) + 2\tau(a, y) \right] \\
 &\leq 2 \sum_y p_{0y} \bar{W} + 2 \sum_y p_{0y} |y|^2 B + 4 \sum_{y \neq a} p_{0y} |y|^2 \tau(a, y) + 4p_{0a} |a|^2 \\
 &\leq 2 \sum_y p_{0y} \bar{W} + 2B \sum_y p_{0y} |y|^2 + 4 \left(B \sum_{y \neq a} p_{0y}^2 |y|^4 \right)^{1/2} + 4p_{0a} |a|^2.
 \end{aligned}$$

The desired bound now follows from the inequality $B \leq T/3$. \square

4. Proof of the Main Results (Theorem 1.1) for the Nearest-Neighbour Model

In this section we use the results of Sect. 2 and 3 to prove Theorem 1.1. The general structure of the proof follows the outline given in Sect. 1.3 and is described in Sect. 4.1. Throughout this section we consider primarily the nearest-neighbour model, for which $p_{0v} = p$ if $|v| = 1$ and $p_{0v} = 0$ otherwise. In particular, $\sum_v p_{0v} = 2dp$. We use this and similar facts in the bounds given in Lemma 3.2, without further mention. However since Lemmas 4.1 and 4.2 hold quite generally, we state and prove these lemmas for both the nearest-neighbour and the spread-out models introduced in Sect. 1.1.

We use the notation

$$D_L(k) = \sum_x p_{0x} e^{ik \cdot x}$$

for the spread-out models at the gaussian critical value p_G , i.e., the value of p for which $D_L(0) = 1$. For the nearest neighbour model at its critical value $p_G = 1/2d$ we write simply

$$D(k) = d^{-1} \sum_\mu \cos k_\mu$$

in place of $D_L(k)$, and we use D_G to represent either D or D_L .

4.1. General Structure

The fact that the triangle diagram $\mathcal{V}(p)$ is bounded uniformly in $p < p_c$ is proved using the following three results. Lemmas 4.1 and 4.2 are valid for both the nearest neighbour model and the spread-out models introduced in Sect. 1.1. Proposi-

tion 4.3 is valid as stated only for the nearest-neighbour model; a variation valid for the spread-out models is proved in Sect. 5.1.

Lemma 4.1. *For both the models (i) and (ii) of Sect. 1.1, if $0 \leq p \leq p_G$ then $\tau_p(x, y) \leq C_G(x, y)$, where $C_G(x, y)$ denotes the gaussian propagator at $p = p_G$:*

$$C_G(x, y) = (2\pi)^{-d} \int d^d k \frac{e^{ik \cdot (y-x)}}{1 - D_G(k)}.$$

Lemma 4.2. *For both models (i) and (ii) of Sect. 1.1, T , W_a , and H_{a_1, a_2} are continuous in p , for $p < p_c$ and for all $a, a_1, a_2 \in \mathbb{Z}^d$.*

Proposition 4.3. *For the nearest-neighbour model (i), there is a universal constant $d_0 > 6$ such that for $d \geq d_0$ and for any fixed $p \in [1/2 d, p_c)$, $P_4 \Rightarrow P_3$, where P_α is the statement that the following set of inequalities holds:*

$$\begin{aligned} T &\leq \alpha K_T d^{-1}, & W &\leq \alpha K_W d^{-1}, & 2dp &\leq \alpha, \\ W_a &\leq \alpha K'_W d^{-1} & \text{for } \|a\|_1 &\leq 2\chi(p) \{(d+2) \ln[5\chi(p)] + 2 \ln d\}, \\ H_{a_1, a_2} &\leq 10 \alpha K_W d^{-1} & \text{for } \max_{i=1, 2} \|a_i\|_1 &\leq 2\chi(p) \{(5d+2) \ln[5\chi(p)] + 2 \ln d\}. \end{aligned}$$

In Proposition 4.3 K_T and K_W are universal constants such that for $d \geq 7$,

$$T_G \equiv \sum_{x, y} C(0, x) C(x, y) C(y, 0) - C(0, 0)^3 \leq \frac{1}{25} K_T d^{-1},$$

and

$$W_G \equiv \sum_x |x|^2 C(0, x)^2 \leq \frac{1}{30} K_W d^{-1}.$$

The existence of such constants follows from Appendix B of [20] or Lemma 3.1 of [31]. (In the notation of [31], $C(0, x) = \sum_{T=0}^\infty N_0(x, T)$.) The constant K'_W is a (large) universal constant which depends only on K_T and K_W and is determined in the proof of the proposition. The statement P_α , for fixed $p < p_c$, involves only *finitely* many inequalities, since $\chi(p) < \infty$. For $\|a\|_1$ or $\|a_i\|_1$ violating the inequalities in P_α , it follows from the exponential decay of $\tau_p(x, y)$ that $W_a \leq d^{-1}$ and $H_{a_1, a_2} \leq d^{-1}$; see the proof of Lemma 4.4.

It follows immediately from Lemma 4.1 that for $p \in [0, 1/2 d]$ the first three inequalities in the definition of P_α hold with $\alpha = 1$. The other inequalities in the definition of P_1 can be proved for $p \in [0, 1/2 d]$ by using Lemma 4.1 to bound all propagators by the gaussian propagator, and then analyzing the resulting gaussian quantities. This can be done using the method of Sect. 4.3.3(d–e), by putting $\hat{G} = 1$ and $\hat{\Gamma} = 0$ there. It follows from the fact that P_1 holds for $p \leq 1/2 d$, together with Lemma 4.2 and Proposition 4.3 that there are forbidden regions in the graphs of T , W , W_a and H_{a_1, a_2} as a function of p , for $p < p_c$, as illustrated in Fig. 1. Therefore P_3 holds and so in particular $\mathcal{V}(p) = T(p) + 1$ is bounded

uniformly in $p < p_c$. Since $\tau_p(x, y)$ is an increasing and continuous function of p [5], it then follows from the monotone convergence theorem that

$$\nabla(p_c) = \lim_{p \uparrow p_c} \nabla(p) \leq 3 K_T d^{-1} + 1,$$

and hence the triangle condition is satisfied.

In the course of the proof of Proposition 4.3 it is shown that if P_4 holds, then the infrared bound stated in Theorem 1.1 follows. But as we have noted it follows from Lemmas 4.1 and 4.2 and Proposition 4.3 that in fact P_3 holds, and hence the weaker statement P_4 holds, and the infrared bound follows. Also, it follows almost automatically from the proof that the relation $\xi_2(p)^2 \sim \chi(p)$ holds, and hence $\nu_2 = 1/2$.

The proof of Theorem 1.1 has now been reduced to proving Lemmas 4.1 and 4.2 and Proposition 4.3, and showing that the infrared bound follows from P_4 . This is done in the remaining subsections. The two lemmas are elementary and do not use the expansion. The proof of Proposition 4.3 is the significant part of the analysis.

4.2. Proof of Lemmas 4.1 and 4.2

In this section we prove Lemmas 4.1 and 4.2.

Proof of Lemma 4.1. Fix any $p \in [0, p_G]$. Any bond configuration in which x and y are connected contains an occupied path ω from x to y . We denote the sum over all paths from x to y by $\sum'_{\omega: x \rightarrow y}$, and the sum over all simple (not necessarily self-avoiding) walks from x to y , consisting of bonds for which $p_b \neq 0$, by $\sum_{\omega: x \rightarrow y}$. Then

$$\begin{aligned} \tau_p(x, y) &\leq \sum'_{\omega: x \rightarrow y} \text{Prob}_p(\omega \text{ is occupied}) = \sum'_{\omega: x \rightarrow y} \prod_{b \in \omega} p_b \\ &\leq \sum_{\omega: x \rightarrow y} \prod_{b \in \omega} p_b \leq C_G(x, y), \end{aligned}$$

where in the last inequality we used $p \leq p_G$. \square

Proof of Lemma 4.2. In [18] it was proved that for quite general finite range models, if $\chi(p) < \infty$ then $\tau_p(x, y)$ decays exponentially. Moreover in [6] it was shown that

$$\tau_p(0, x) \leq \exp[-\|x\|_1/\chi(p)] \quad \text{for model (i).}$$

For model (ii) a similar exponential decay holds if g has compact support, and for general exponentially decaying g exponential decay of $\tau_p(x, y)$ (for $p < p_c$) can be shown using the same argument as that used for the Ising model in [7]. The exponential decay of the two-point function implies that T , W_a , and H_{a_1, a_2} are finite for $p < p_c$.

Continuity of these quantities for $p < p_c$ then follows from the fact that $\tau_p(x, y)$ is monotonic and continuous in $p \in [0, 1]$ [5], together with the monotone convergence theorem. \square

4.3. *Proof of Proposition 4.3.* In this section we restrict our attention to the nearest-neighbour model. We prove Proposition 4.3 and show that the infrared bound follows from P_4 , thereby completing the proof of Theorem 1.1. We begin with a preliminary lemma in which bounds on \bar{T} , \bar{W} and \bar{H} are obtained from the assumption P_4 .

4.3.1. *Bounds on \bar{T} , \bar{W} , and \bar{H} .* In this section we prove the following lemma.

Lemma 4.4. *If $p < p_c$ and we assume P_4 , then*

$$\begin{aligned} \bar{T} &\leq 4d^{-1} [K_T + (3K_T/8)^{1/2}] \equiv c_1 d^{-1}, \\ W_1 &\leq d^{-1} [4K_W + 4(K_W K_T/3)^{1/2} + (2K_T/3)^{1/2}] \equiv c_2 d^{-1}, \\ \bar{W} &\leq 4K'_W d^{-1}, \quad \bar{H} \leq 40K_W d^{-1}, \\ W' &\leq [16K_W + 16(K_W K_T/3)^{1/2} + 4(2K_W)^{1/2}] d^{-1} \equiv c_3 d^{-1}, \\ \bar{W}' &\leq 8d^{-1} [4K'_W + 4K_T/3 + 2(2K_T/3)^{1/2} + 1] \leq c_4 [K'_W + 1] d^{-1}. \end{aligned}$$

Here W_1 denotes W_a where a is any nearest neighbour of the origin. Note that the constants c_1, c_2, c_3 and c_4 depend only on K_T and K_W , and not on K'_W .

Proof. By Lemma 3.3, for any $a \in \mathbb{Z}^d$ we have

$$T_a \leq T + [3T/2d]^{1/2},$$

and the bound on \bar{T} then follows if we assume P_4 .

To obtain the inequality for W_1 we proceed as follows: Denoting a nearest neighbour of the origin by e ,

$$\begin{aligned} W_1 &= \sum_x |x|^2 \tau(0, x) \tau(x, e) \\ &\leq \sum_x |x| \tau(0, x) |x - e| \tau(e, x) + \sum_{x \neq e} |x| \tau(0, x) |e| \tau(e, x) \\ &\quad + |e| \tau(0, e) |e| \tau(e, e). \end{aligned}$$

Applying the Schwarz inequality to the first two terms on the right side, and using the fact that

$$\tau(0, e)^2 \leq (2d)^{-1} \sum_{x \neq 0} \tau(0, x)^2 \leq (2d)^{-1} T/3,$$

gives

$$W_1 \leq W + \left[W \cdot \sum_{x \neq e} \tau(e, x)^2 \right]^{1/2} + (T/6d)^{1/2} \leq W + [WT/3]^{1/2} + (T/6d)^{1/2}.$$

The assumption P_4 then gives the bound $W_1 \leq c_2 d^{-1}$.

We next give the proof that P_4 implies the bound $\bar{W} \leq 4K'_W d^{-1}$. Although P_4 only involves W_a for $\|a\|_1 \leq 2\chi(p) \{(d+2) \ln[5\chi(p)] + 2 \ln d\}$, it is possible to estimate $\bar{W} = \sup_a W_a$ by using the exponential decay of the two-point function to bound W_a for large values of $\|a\|_1$. In fact, we shall show that

$$W_a \leq d^{-1} \quad \text{for} \quad \|a\|_1 \geq 2\chi(p) \{(d+2) \ln[5\chi(p)] + 2 \ln d\}. \quad (4.2)$$

Given (4.2), it follows that

$$\bar{W} \leq \max \{4K_W d^{-1}, 4K'_W d^{-1}, d^{-1}\} = 4K'_W d^{-1}.$$

To prove (4.2) we use the fact [6] that $\tau(0, x) \leq \exp[-\|x\|_1/\chi(p)]$ to obtain

$$\begin{aligned} W_a &\leq \sum_x |x|^2 \exp[-(\|x\|_1 + \|x - a\|_1)/\chi(p)] \\ &\leq \sum_x |x|^2 \exp[-\|x\|_1/2\chi(p)] \exp[-\|a\|_1/2\chi(p)] \\ &\leq d[5\chi(p)]^{d+2} \exp[-\|a\|_1/2\chi(p)]. \end{aligned}$$

The inequality for \bar{H} is proved similarly, and we omit the proof. The bounds on W' and \bar{W}' follow from P_4 and Lemma 3.3. \square

4.3.2. *Bounds on $\hat{\tau}(k)$.* In Proposition 2.3 an expansion was given for the two-point function $\tau(0, x)$. Since $\tau(0, x)$ decays exponentially for $p < p_c$, the Fourier transform $\hat{\tau}(k)$ exists. Taking the Fourier transform of both sides of the expansion and solving for $\hat{\tau}(k)$ yields

$$\hat{\tau}(k) = \frac{\hat{G}^{(N)}(k)}{1 - 2dpD(k) - \hat{\Pi}^{(N)}(k)}, \tag{4.3}$$

for any $p \in [0, p_c)$ and for any nonnegative integer N , where

$$\hat{G}^{(N)}(k) = 1 + \sum_{n=0}^N (-1)^n \hat{g}_n(k) + (-1)^{N+1} \hat{R}_N(k),$$

and

$$\hat{\Pi}^{(N)}(k) = \sum_{n=0}^N (-1)^n \hat{\Pi}_n(k).$$

In this section we prove the following lemma.

Lemma 4.5. *Fix $p \in [1/2d, p_c)$ and $N \geq 0$, and assume P_4 . There exists a $d_0 > 6$, independent of p , such that for $d \geq d_0$,*

$$|\hat{\Pi}^{(N)}(k)| \leq cd^{-1}, \quad |\partial_\mu^s \hat{\Pi}^{(N)}(k)| \leq c'd^{-2}, \quad s = 1, 2. \tag{4.4}$$

If in addition N is sufficiently large (depending on d and p), then

$$\hat{F}(k) \equiv 1 - 2dpD(k) - \hat{\Pi}^{(N)}(k) \geq (1 - c''d^{-1})(1 - D(k)), \tag{4.5}$$

$$|\hat{G}^{(N)}(k) - 1| \leq cd^{-1}, \quad |\partial_\mu^s \hat{G}^{(N)}(k)| \leq c'd^{-2}, \quad s = 1, 2, \tag{4.6}$$

and

$$0 \leq \hat{\tau}(k) \leq (1 + c'''d^{-1})(1 - D(k))^{-1}. \tag{4.7}$$

The constants c, c', c'' and c''' depend only on K_T and K_W (and not on K'_W, p , or d).

Remark. For $p \in [0, 1/2d]$, $\hat{\tau}(k) \leq \hat{\tau}(0) = \chi(p) \leq \chi(1/2d) < \infty$. (It is known that $\hat{\tau}(k)$ is nonnegative [6].) Together with (4.7) this shows that for large d the infrared bound stated in Theorem 1.1 is a consequence of P_4 .

Proof of Lemma 4.5. The bounds (4.4)–(4.6) are simply a combination of the results in Proposition 2.6, Lemma 3.2, and Lemma 4.4 with the assumption P_4 . In the course of the proof, we use c and c' to denote universal constants which may depend on K_T and K_W , but not on K'_W , p , or d . These constants may take on different values in different occurrences. First, by Lemma 3.2, Lemma 4.4 and P_4 ,

$$0 \leq \sum_x h_n(0, x) \leq \begin{cases} cd^{-1} & n = 0, 1 \\ (c'd^{-1})^n & n \geq 2 \end{cases} \tag{4.8}$$

and

$$0 \leq \sum_x |x|^2 h_n(0, x) \leq \begin{cases} cd^{-1} & n = 0, 1 \\ n^2 (c'd^{-1})^{n-1} & n \geq 2 \end{cases}. \tag{4.9}$$

Now using Proposition 2.6 and (4.8–9) we immediately obtain

$$|\hat{H}^{(N)}(k)| \leq 2dp \sum_{n=0}^N \sum_x h_n(0, x) \leq cd^{-1}.$$

The bound on $\partial_\mu^s \hat{H}^{(N)}(k)$ is obtained similarly.

To bound $|\hat{G}^{(N)}(k) - 1|$ we use Proposition 2.6 to obtain

$$|\hat{G}^{(N)}(k) - 1| \leq \sum_{n=0}^N \sum_x h_n(0, x) + 2dp \chi(p) \sum_x h_N(0, x),$$

and then note that the first term on the right side is bounded above by cd^{-1} , while the second is bounded above by $2dp \chi(p) (c'd^{-1})^N$. For N sufficiently large, this last quantity is bounded above by cd^{-1} , and the desired bound on $|\hat{G}^{(N)}(k) - 1|$ follows.

The bounds on $|\partial_\mu^s \hat{G}^{(N)}(k)|$ are obtained in a similar fashion. To estimate $|\partial_\mu^s \hat{R}_N(k)|$ we use the bound from Proposition 2.6 and the inequality

$$\sum_x |x|^2 \tau(0, x) \leq \sum_x |x|^2 \exp[-\|x\|_1 / \chi(p)] \leq d(3\chi(p))^{d+2}.$$

To obtain (4.5) we first observe that by (4.6),

$$1 - 2dp - \hat{H}^{(N)}(0) = \hat{G}^{(N)}(0) \chi(p)^{-1} \geq 0.$$

Therefore $2dp \leq 1 + cd^{-1}$, and also

$$\begin{aligned} 1 - 2dp D(k) - \hat{H}^{(N)}(k) &= 1 - 2dp D(k) - \hat{H}^{(N)}(k) - [1 - 2dp - \hat{H}^{(N)}(0)] \\ &\quad + [1 - 2dp - \hat{H}^{(N)}(0)] \\ &\geq 2dp [1 - D(k)] + \hat{H}^{(N)}(0) - \hat{H}^{(N)}(k). \end{aligned}$$

By Proposition 2.6 and (2.24),

$$\begin{aligned} \hat{H}^{(N)}(0) - \hat{H}^{(N)}(k) &\geq - \sum_{\substack{n=1 \\ n:\text{odd}}}^N [\hat{H}_n(0) - \hat{H}_n(k)] \\ &\geq \frac{\pi^2}{4} (1 - D(k)) \cdot 2dp \left[- \sum_{\substack{n=1 \\ n:\text{odd}}}^N \sum_x (|x|^2 + 1) h_n(0, x) \right]. \end{aligned}$$

Now, using (4.8–9) gives (4.5).

Finally the infrared bound (4.7) follows from (4.3) and (4.5–6). \square

4.3.3. *Proof that $P_4 \Rightarrow P_3$.* In this section we use the consequences of P_4 given in Lemma 4.5 to derive P_3 . This completes the proof of Proposition 4.3 and thus of Theorem 1.1. We fix an N large enough (depending on p) that (4.5–7) hold, and omit the superscript (N) to simplify the notation. As usual we use c, c' and c'' to denote universal constants which depend only on K_T and K_W , but not on K'_W, p , or d . These constants may represent different values in different occurrences. Throughout this section we fix $p \in [1/2d, p_c)$.

The statement P_3 consists of five inequalities (given in Proposition 4.3). We obtain these one by one, beginning with $2dp \leq 3$

(a) $2dp \leq 3$: In the proof of Lemma 4.5 it was shown that for $p \in [1/2d, p_c)$,

$$1 - 2dp - \hat{\Pi}(0) \geq 0,$$

and hence by (4.4)

$$2dp \leq 1 - \hat{\Pi}(0) \leq 1 + cd^{-1}. \tag{4.10}$$

The right side of (4.10) is less than 3 for $d \geq c/2$.

(b) The bound on T : By definition

$$\begin{aligned} T &= (2\pi)^{-d} \int d^d k [\hat{\tau}(k)^3 - \tau(0, 0)^3] \\ &= (2\pi)^{-d} \int d^d k [(2 + \hat{\tau}(k)) (\hat{\tau}(k) - 1)^2 - (2 + \tau(0, 0)) (\tau(0, 0) - 1)^2], \end{aligned} \tag{4.11}$$

and T_G is obtained by replacing $\hat{\tau}(k)$ by $[1 - D(k)]^{-1}$. (In fact, for percolation $\tau(0, 0) = 1$ and the second term on the right side of (4.11) is zero.) By (4.7), for $d \geq 4c'''$,

$$0 \leq 2 + \hat{\tau}(k) \leq 2 + (1 + c'''d^{-1}) [1 - D(k)]^{-1} \leq 2 + \frac{5}{4} [1 - D(k)]^{-1}. \tag{4.12}$$

Also,

$$\hat{\tau}(k) - 1 = \frac{2dpD(k) + \hat{G}(k) - 1 + \hat{\Pi}(k)}{1 - 2dpD(k) - \hat{\Pi}(k)}.$$

We define

$$\hat{F}(k) = 1 - 2dpD(k) - \hat{\Pi}(k). \tag{4.13}$$

Then by (4.4–6),

$$\begin{aligned} [\hat{\tau}(k) - 1]^2 &\leq \hat{F}(k)^{-2} 2 \{ (2dpD(k))^2 + (|\hat{G}(k) - 1| + |\hat{\Pi}(k)|)^2 \} \\ &\leq 20 [1 - D(k)]^{-2} (D(k)^2 + c'd^{-2}), \end{aligned} \tag{4.14}$$

for d sufficiently large. Using the fact that the integral of $[1 - D(k)]^{-m}$ is bounded uniformly in $d \geq 7$ for $m = 1, 2, 3$ (see Appendix B of [20] or Eq. (3.12) of [31] for a proof), it follows from (4.11), (4.12) and (4.14) that

$$\begin{aligned} T &\leq 20 (2\pi)^{-d} \int d^d k (2 + \frac{5}{4} [1 - D(k)]^{-1}) D(k)^2 [1 - D(k)]^{-2} + c''d^{-2} \\ &\leq 25 T_G + c''d^{-2} \leq K_T d^{-1} + c''d^{-2} \leq 3 K_T d^{-1}, \end{aligned}$$

for d sufficiently large, where we used the fact that $C(0, 0) - 1 \leq cd^{-1}$ (which follows from Appendix B of [20] or Lemma 3.1 of [31]) together with the form of T_G described below (4.11).

(c) The bound on W : By definition and the Parseval relation,

$$W = \sum_x [|x| \tau(0, x)]^2 = \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k [\partial_\mu \hat{\tau}(k)]^2. \tag{4.15}$$

Differentiation of (4.3) gives

$$\begin{aligned} \partial_\mu \hat{\tau}(k) &= \hat{F}(k)^{-2} \hat{G}(k) 2dp \partial_\mu D(k) + \hat{F}(k)^{-1} \partial_\mu \hat{G}(k) \\ &\quad + \hat{F}(k)^{-2} \hat{G}(k) \partial_\mu \hat{\Pi}(k). \end{aligned} \tag{4.16}$$

By (4.15–16)

$$\begin{aligned} W \leq \sum_{\mu=1}^d 3(2\pi)^{-d} \int d^d k \{ &\hat{F}(k)^{-4} 9[\hat{G}(k) \partial_\mu D(k)]^2 + \hat{F}(k)^{-2} (\partial_\mu \hat{G}(k))^2 \\ &+ \hat{F}(k)^{-4} (\hat{G}(k) \partial_\mu \hat{\Pi}(k))^2 \}. \end{aligned} \tag{4.17}$$

We use (4.5) to estimate the powers of $\hat{F}(k)$ occurring in the three terms on the right-hand side. By (4.6), the first term (including the sum over μ) is bounded by $30 W_G$ for d sufficiently large, and by (4.6) the second is bounded above by cd^{-3} . To bound the third term, we first note that by symmetry $\partial_\mu \hat{\Pi}(k)$ is equal to zero for any k for which $k_\mu = 0$. Denoting by \tilde{k} the result of replacing the μ^{th} component of k by zero, it follows from Taylor’s theorem that

$$\partial_\mu \hat{\Pi}(k) = \partial_\mu \hat{\Pi}(k) - \partial_\mu \hat{\Pi}(\tilde{k}) = k_\mu \partial_\mu^2 \hat{\Pi}(k^*),$$

where k^* is a point on the line segment joining k and \tilde{k} . By (4.4) and (4.6), the third term in (4.17) is bounded above by

$$cd^{-4} (2\pi)^{-d} \int d^d k [1 - D(k)]^{-4} k^2.$$

Since $k^2 d^{-1} \leq \frac{\pi^2}{2} [1 - D(k)]$, this is bounded above by $c'd^{-3}$. This gives the desired bound on W , for d sufficiently large.

(d) The bound on W_a : For $|a| = 1$, it was shown in Lemma 4.4 that $W_a \leq c_2 d^{-1}$ and hence $W_a \leq 3K'_W d^{-1}$ if we take $K'_W \geq c_2/3$.

For the case $|a| > 1$ we again use Fourier transforms and write

$$W_a = \sum_x |x|^2 \tau(0, x) \tau(x, a) = -(2\pi)^{-d} \int d^d k \hat{\tau}(k) \sum_{\mu=1}^d \hat{\tau}_{\mu\mu}(k) e^{ik \cdot a}, \tag{4.18}$$

where we have used subscripts to denote partial derivatives with respect to k_μ . Now

$$\hat{\tau}_{\mu\mu}(k) = \hat{F}^{-1} \hat{G}_{\mu\mu} - 2\hat{F}^{-2} \hat{G}_\mu \hat{F}_\mu - \hat{F}^{-2} \hat{G} \hat{F}_{\mu\mu} + 2\hat{F}^{-3} \hat{G} \hat{F}_\mu^2, \tag{4.19}$$

with

$$\hat{F}_\mu = -2dp D_\mu - \hat{\Pi}_\mu = 2p \sin k_\mu - \hat{\Pi}_\mu$$

and

$$\hat{F}_{\mu\mu} = -2dpD_{\mu\mu} - \hat{\Pi}_{\mu\mu} = 2p \cos k_\mu - \hat{\Pi}_{\mu\mu}.$$

Therefore

$$W_a = - \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k e^{ik \cdot a} \cdot [\hat{\tau}^2 \hat{G}^{-1} \hat{G}_{\mu\mu} - 2\hat{\tau}^3 \hat{G}^{-2} \hat{G}_\mu \hat{F}_\mu - \hat{\tau}^3 \hat{G}^{-1} \hat{F}_{\mu\mu} + 2\hat{\tau}^4 \hat{G}^{-2} \hat{F}_\mu^2]. \quad (4.20)$$

We estimate the terms on the right side using (4.4), (4.6) and (4.7), together with the fact that the integral of $[1 - D(k)]^{-m}$ is uniformly bounded for $m \leq 3$ and $d \geq 7$. The first term (including the sum over μ) is less than cd^{-1} , and since $\hat{F}_\mu \leq c'd^{-1}$ the second term is less than cd^{-2} . The estimate on the third term is more involved.

The contribution to W_a arising from the gaussian part $2p \cos k_\mu$ of $\hat{F}_{\mu\mu}$ in the third term on the right side of (4.20) is

$$A_a \equiv 2dp (2\pi)^{-d} \int d^d k \hat{\tau}(k)^3 [1 + \hat{G}(k)^{-1} - 1] d^{-1} \sum_{\mu=1}^d \cos k_\mu e^{ik \cdot a}.$$

To estimate the contribution A'_a to A_a due to the 1 in the square brackets we write

$$d^{-1} \sum_{\mu=1}^d \cos k_\mu = (2d)^{-1} \sum_{v, |v|=1} e^{ik \cdot v}$$

to get

$$\begin{aligned} |A'_a| &\leq 2dp (2d)^{-1} \sum_{v, |v|=1} \left| (2\pi)^{-d} \int d^d k \hat{\tau}(k)^3 e^{ik \cdot (a+v)} \right| \\ &= 2dp (2d)^{-1} \sum_{v, |v|=1} T_{a+v} \leq 3\bar{T} \leq 3c_1 d^{-1}. \end{aligned}$$

(Note that $a + v \neq 0$ here since $|a| \neq 1$.) Then by (4.6–7) and the fact that $2dp \leq 3$ we have

$$|A_a| \leq |A'_a| + |A_a - A'_a| \leq 3c_1 d^{-1} + cd^{-1} \leq c'd^{-1}.$$

Using (4.4) and (4.6–7) it is not difficult to see that the contribution to W_a due to the $\hat{\Pi}_{\mu\mu}$ term in $\hat{F}_{\mu\mu}$ in the third term on the right side of (4.20) is also bounded by cd^{-1} .

Finally the last term on the right hand side of (4.20) can be bounded above by cd^{-1} using an argument involving Taylor’s Theorem which is similar to that used in the last paragraph of (c). (Note that it is necessary to exploit the fact that $F_\mu(0) = 0$ if we are to avoid quantities which diverge in more than six dimensions.) All the constants c and c' encountered here depend only on K_T and K_W , so it is possible to choose K'_W (depending on K_T and K_W) sufficiently large that the desired bound $W_a \leq 3K'_W d^{-1}$ holds, and that the corresponding gaussian quantity is less than $K'_W d^{-1}$ (as claimed in Sect. 4.1).

(e) The bound on H : Using the Fourier transform, and then integrating by parts, we can rewrite Eq. (3.3) as

$$\begin{aligned}
 H_{a_1, a_2} &= \int \prod_{j=1}^3 \frac{d^d k_j}{(2\pi)^d} e^{i(k_1 \cdot a_1 - k_2 \cdot a_2)} \hat{\tau}(k_1)^2 \hat{\tau}(k_2)^2 \hat{\tau}(k_1 - k_2) \hat{\tau}(k_1 - k_3) \\
 &\quad \cdot \hat{\tau}(k_3 - k_2) \left[- \sum_{\mu=1}^d \partial_\mu^2 \hat{\tau}(k_3) \right] \\
 &= \sum_{\mu=1}^d \int \prod_{j=1}^3 \frac{d^d k_j}{(2\pi)^d} e^{i(k_1 \cdot a_1 - k_2 \cdot a_2)} \hat{\tau}(k_1)^2 \hat{\tau}(k_2)^2 \hat{\tau}(k_1 - k_2) \\
 &\quad \cdot \{ \hat{\tau}(k_1 - k_3) (\partial_\mu \hat{\tau}(k_3)) (\partial_\mu \hat{\tau}(k_3 - k_2)) - \hat{\tau}(k_3 - k_2) (\partial_\mu \hat{\tau}(k_3)) (\partial_\mu \hat{\tau}(k_1 - k_3)) \}.
 \end{aligned}$$

Taking absolute values of the integrand on the right side and using the symmetry between k_1 and k_2 leads to

$$\begin{aligned}
 H_{a_1, a_2} &\leq \sum_{\mu=1}^d \int \prod_{j=1}^3 \frac{d^d k_j}{(2\pi)^d} \hat{\tau}(k_1)^2 \hat{\tau}(k_2)^2 \hat{\tau}(k_1 - k_2) \\
 &\quad \cdot \hat{\tau}(k_1 - k_3) \{ (\partial_\mu \hat{\tau}(k_3))^2 + (\partial_\mu \hat{\tau}(k_3 - k_2))^2 \}.
 \end{aligned}$$

Rewriting this upper bound in x -space gives

$$H_{a_1, a_2} \leq \sum_{\mu=1}^d \sum_{x, a} |x_\mu| |x_\mu - a_\mu| \left(\begin{array}{c} \text{Diagram 1} \\ + \\ \text{Diagram 2} \end{array} \right). \tag{4.21}$$

Now we use the basic inequality (3.5) to bound the right side. First,

$$\begin{aligned}
 &\sum_{\mu=1}^d \sum_{x, a} |x_\mu| |x_\mu - a_\mu| \begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \\
 &\leq \left[\sup_a \sum_{\mu=1}^d \sum_x |x_\mu| |x_\mu - a_\mu| \begin{array}{c} \text{Diagram 2} \\ 0 \end{array} \right] \begin{array}{c} \text{Diagram 3} \\ 0 \end{array} \\
 &\leq W \left[\sup_a \begin{array}{c} 0 \\ \text{Diagram 4} \\ a \end{array} \right] (1 + T) \leq W(1 + T)(1 + \bar{T}), \tag{4.22}
 \end{aligned}$$

where in the second inequality the sum over x was estimated using the Schwarz inequality. The second term on the right side of (4.21) can be bounded similarly:

$$\sum_{\mu=1}^d \sum_{x,a} |x_\mu| |x_\mu - a_\mu| \begin{array}{c} \text{diagram} \\ 0 \end{array} \leq \sum_{\mu=1}^d \begin{array}{c} \text{diagram} \\ \text{diagram} \end{array} \sup_a \sum_x |x_\mu| |x_\mu - a_\mu| \begin{array}{c} \text{diagram} \\ 0 \end{array} \leq \begin{array}{c} \text{diagram} \\ \text{diagram} \end{array} W. \quad (4.23)$$

Now we use (4.7) to bound the diagram on the right side of (4.23) in terms of its gaussian value:

$$\begin{array}{c} \text{diagram} \\ \text{diagram} \end{array} \leq (1 + c''' d^{-1})^6 \begin{array}{c} \text{diagram} \\ \text{diagram} \end{array} \Big|_{\text{gaussian}}. \quad (4.24)$$

The gaussian diagram on the right side is finite for $d > 6$ and we have verified numerically that it is bounded by 3 uniformly in $d \geq 9$.

For d sufficiently large, depending on K_T and K_W , it follows from (4.21–24) that

$$H_{a_1, a_2} \leq \frac{3}{2} \left(1 + \begin{array}{c} \text{diagram} \\ \text{diagram} \end{array} \Big|_{\text{gaussian}} \right) W \leq 30 K_W d^{-1}. \quad (4.25)$$

This is the desired bound. Also, this calculation shows that the gaussian counterpart of H_{a_1, a_2} is bounded above by $10 K_W d^{-1}$ for d sufficiently large, as claimed in Sect. 4.1. \square

4.4. Proof that $v_2 = 1/2$

By definition [Eq. (1.6)],

$$\xi_2(p)^2 = - \sum_{\mu=1}^d \frac{\hat{t}_{\mu\mu}(0)}{\hat{t}(0)}. \quad (4.26)$$

Since $\hat{F}^{-1} = \hat{G}^{-1} \hat{t}$, $\hat{F}_\mu(0) = 0$, and $\hat{t}(0) = \chi(p)$, it follows from (4.19) and (4.26) that

$$\begin{aligned} \xi_2(p)^2 &= - \sum_{\mu=1}^d [\hat{G}(0)^{-1} \hat{G}_{\mu\mu}(0) - \chi(p) \hat{G}(0)^{-1} \hat{F}_{\mu\mu}(0)] \\ &= - \sum_{\mu=1}^d \hat{G}(0)^{-1} [\hat{G}_{\mu\mu}(0) - \chi(p) (2p - \hat{\Pi}_{\mu\mu}(0))]. \end{aligned}$$

By Lemma 4.5 (since we have proved P_3, P_4 does hold), it follows that for $1/2d \leq p < p_c$,

$$\xi_2(p)^2 = O(d^{-1}) + [1 + O(d^{-1})]\chi(p),$$

and hence for some constant $C > 0$,

$$C^{-1}\chi(p) \leq \xi_2(p)^2 \leq C\chi(p) \quad \text{for } p \in [1/2d, p_c]. \tag{4.27}$$

Since $\chi(p) \sim (p_c - p)^{-1}$ follows from the triangle condition, $\xi_2(p) \sim (p_c - p)^{-1/2}$, i.e., $\nu_2 = 1/2$.

5. Proof of the Main Results (Theorem 1.2) for the Spread-out Models

In this section we prove Theorem 1.2 by adapting the proof of Theorem 1.1 to the spread-out models introduced in Sect. 1.1. These models are defined by

$$p_{0x} = p \cdot L^{-d}g(x/L),$$

where g is a nonnegative function on \mathbb{R}^d such that $\partial^I g$ ($\partial^I g \equiv \prod_{\mu \in I} \partial_\mu g, I \subset \{1, 2, \dots, d\}$) is piecewise continuous, which satisfies:

$$\begin{aligned} \int g(x) dx &= 1, \quad g \cdot e^{\delta \|x\|_1} \in L_\infty(\mathbb{R}^d) \text{ for some } \delta > 0, \\ \int |\partial^I g(x)| dx &< \infty, \text{ where the derivative is interpreted as a distribution,} \\ g &\text{ is invariant under rotations by } \pi/2 \text{ and reflections} \\ &\text{ in the coordinate hyperplanes.} \end{aligned} \tag{5.1}$$

To simplify the technicalities we have required that g be exponentially decaying, although some weaker decay can also be handled. The fact that Theorem 1.2 holds independently of the exact form of g is an illustration of universality.

Just as for the nearest-neighbour model, $p_c = \sup \{p \mid \chi(p) < \infty\} \in (0, 1)$. The only significant difference between the analysis of the model (ii) and the nearest-neighbour model is due to the change in gaussian propagator. (The situation is similar for the long-range weakly self-avoiding walk studied in [40], although in that work the $1/r^2$ decay considered was sufficiently slow to change the upper critical dimension, unlike the models under study here.)

The hypotheses on g are used indirectly, via certain of their consequences, to prove Theorem 1.2. To state these consequences we introduce the following notation. We consider simple random walks with transition probabilities

$$p_{\partial x}^{(L)} = p_L \cdot L^{-d}g(x/L). \tag{5.2}$$

Here p_L is defined so that $D_L(0) = 1$, where

$$D_L(k) \equiv p_L \sum_x L^{-d}g(x/L) e^{ik \cdot x}. \tag{5.3}$$

The gaussian propagator (in k -space) for the model is then

$$\hat{C}_L(k) = [1 - D_L(k)]^{-1}, \tag{5.4}$$

and we define T_L and W_L the same way that T_G and W_G were defined (under Proposition 4.3), using this propagator. We also define

$$S = \sum_x |x|^2 p_{0x}^{(L)}. \tag{5.5}$$

(As will be seen in Sect. 5.2, $S = O(L^2)$; clearly $S \geq 1$.)

To prove Theorem 1.2 we will use the following lemma.

Lemma 5.1. *The hypotheses (5.1) on g imply that given $d > 6$ and $\varepsilon > 0$, there is an L_0 , depending on ε and g , such that for $L \geq L_0$ the following conditions are satisfied:*

$$\sup_x p_{0x}^{(L)}, \quad \sum_x (p_{0x}^{(L)})^2 \leq \varepsilon/S, \tag{5.6}$$

$$\sup_x p_{0x}^{(L)} |x|^2 \leq \varepsilon, \tag{5.7}$$

$$\sum_x (p_{0x}^{(L)})^2 |x|^4 \leq S\varepsilon, \tag{5.8}$$


$$1 - D_L(k) \geq |k|^2/3\pi^2 d, \tag{5.9}$$

$$C_L(0, 0) - 1 \leq \varepsilon/S, \tag{5.10}$$

$$T_L \leq \varepsilon/S, \tag{5.11}$$

$$W_L \leq \varepsilon, \tag{5.12}$$

$$(2\pi)^{-d} \int d^d k \sum_{\mu=1}^d |\partial_\mu^2 D_L(k)| [1 - D_L(k)]^{-3} \leq 40\varepsilon, \tag{5.13}$$



$$\leq 10. \tag{5.14}$$

gaussian

The values of the constants appearing in these conditions are not sharp and can be adjusted.

This section is organized as follows. In Sect. 5.1 Theorem 1.2 is proved, assuming Lemma 5.1. In Sect. 5.2 the proof of Lemma 5.1 is given.

5.1. Proof of Theorem 1.2 Assuming Lemma 5.1

In this section we prove that the triangle condition and the infrared bound follow if conditions (5.6–14) are satisfied, for ε sufficiently small. Theorem 1.2 then follows from Lemma 5.1. The basic structure of the proof is exactly as in Theorem 1.1: the result follows from Lemmas 4.1 and 4.2 and the following analogue of Proposition 4.3, together with a proof that the infrared bound follows from the analogue of P_4 . The proof that $v_2 = 1/2$ proceeds the same way as for the nearest-neighbour model in Sect. 4.4.

The analogue of Proposition 4.3 is the following.

Proposition 5.2. *There is an $\varepsilon_0 > 0$ such that if an independent bond percolation model on \mathbb{Z}^d ($d > 6$) satisfies conditions (5.6–14) for some $\varepsilon \leq \varepsilon_0$, then for any fixed*

$p \in [p_L, p_c)$, P_4 implies P_3 , where P_α is the statement that the following set of inequalities holds:

$$\begin{aligned} T &\leq \alpha \cdot 25\varepsilon/S, & W &\leq \alpha \cdot 30\varepsilon, & \frac{p}{p_L} &\leq \alpha, \\ W_a &\leq \alpha K' \varepsilon \text{ for } \|a\|_1 \leq M_1(p), \\ H_{a_1, a_2} &\leq \alpha \cdot 500\varepsilon \text{ for } \max_{i=1,2} \|a_i\|_1 \leq M_2(p). \end{aligned}$$

The universal constant K' is determined in the proof of the proposition. The quantities $M_i(p)$ are finite for $p < p_c$ and are defined such that

$$W_a \leq \varepsilon \text{ for } \|a\|_1 \geq M_1(p), \quad H_{a_1, a_2} \leq \varepsilon \text{ for } \max_{i=1,2} \|a_i\|_1 \geq M_2(p).$$

The existence of such constants follows from the exponential decay of $\tau_p(x, y)$ (discussed in the proof of Lemma 4.2), as for the nearest-neighbour model.

Proof of Proposition 5.2. The remainder of this section is devoted to the proof of Proposition 5.2. In the course of the proof it will be shown that the infrared bound is a consequence of P_4 . The first step is the following lemma.

Lemma 5.3. *If $p < p_c$ and we assume P_4 , with ε sufficiently small, then there are constants c_i , independent of p and K' , such that*

$$\bar{T} \leq c_1 \varepsilon^{1/2}, \quad \bar{W} \leq 4K' \varepsilon, \quad \bar{H} \leq 2000\varepsilon, \quad W' \leq c_2 \varepsilon, \quad W'_a \leq c_3 K' \varepsilon.$$

Proof. The bounds on \bar{W} and \bar{H} follow immediately from P_4 and the definition of $M_i(p)$. The bound on \bar{T} follows from P_4 and the fact that by Lemma 3.3,

$$T_a \leq T + (3T/2d)^{1/2}.$$

The bounds on W' and W'_a also follow directly from Lemma 3.3, conditions (5.6–14) and P_4 . To bound $\sum_v p_{0v} |v|$ we use the Schwarz inequality:

$$\sum_v p_{0v} |v| \leq \left[\sum_x p_{0x} \cdot \sum_y p_{0y} |y|^2 \right]^{1/2} = \frac{p}{p_L} S^{1/2}.$$

In conditions (5.6–14) the powers of S are chosen in such a way as to cancel the powers of S which arise from the bounds on W'_1 and W'_a of Lemma 3.3. \square

Lemma 5.3 combined with Lemma 3.2 yields the following bounds, in which c and c' are constants independent of p and K' , and ε is taken sufficiently small depending on K' ,

$$0 \leq \sum_x h_n(0, x) \leq \begin{cases} c\varepsilon/S & \text{for } n = 0, 1 \\ \frac{\varepsilon}{S} (c'\varepsilon)^{(n-1)/2} & \text{for } n \geq 2 \end{cases}, \quad (5.15)$$

$$0 \leq \sum_x |x|^2 h_n(0, x) \leq \begin{cases} c\varepsilon & \text{for } n = 0, 1 \\ c\varepsilon n^2 (c'\varepsilon)^{(n-2)/2} & \text{for } n \geq 2 \end{cases}. \quad (5.16)$$

These bounds can be used in conjunction with Proposition 2.6 to control $\hat{\tau}(k)$, just as in Sect. 4.3.2.

Indeed, using the notation under (4.3), (2.2) becomes

$$\hat{\tau}(k) = \hat{G}^{(N)}(k) / \left[1 - \frac{p}{p_L} D_L(k) - \hat{\Pi}^{(N)}(k) \right].$$

Proceeding exactly as in the proof of Lemma 4.5 the following inequalities, in which the constants $c, c',$ and c'' are independent of d and K' , are obtained.

$$|\hat{\Pi}^{(N)}(k)| \leq c\epsilon/S, \quad |\partial_\mu^s \hat{\Pi}^{(N)}(k)| \leq c'\epsilon/d, \quad s = 1, 2. \tag{5.17}$$

If in addition N is sufficiently large, depending on p , then for $p \in [p_L, p_c)$,

$$\hat{F}(k) \equiv 1 - \frac{p}{p_L} D_L(k) - \hat{\Pi}^{(N)}(k) \geq (1 - c''\epsilon)(1 - D_L(k)), \tag{5.18}$$

$$|\hat{G}^{(N)}(k) - 1| \leq c\epsilon/S, \quad |\partial_\mu^s \hat{G}^{(N)}(k)| \leq c'\epsilon/d, \quad s = 1, 2, \tag{5.19}$$

and

$$\hat{\tau}(k) \leq (1 + c''\epsilon)(1 - D_L(k))^{-1}. \tag{5.20}$$

The proof of (5.18) uses (5.9), and the fact that (5.19) holds for sufficiently large N uses the exponential decay of $\tau_p(x, y)$.

We now turn to the proof that P_4 implies P_3 , and proceed step by step through the five inequalities in the statement of P_4 . Most of the analysis is identical to that of Sect. 4.3.3, using (5.17–20), and we refer the reader to that section. Now ϵ plays the role of $1/d$. To simplify the notation we omit the superscript (N) .

(a) $\frac{p}{p_L} \leq 3$: This is identical to the nearest neighbour case in Sect. 4.4.3(a).

(b) The bound on T : This bound can also be obtained by the same argument as in the nearest neighbour case, by faithfully following Sect. 4.3.3(b). Note that the integral of $(1 - D_L)^{-m}$ ($m = 1, 2, 3$) is bounded, in fact close to one, by conditions (5.10–11) and the Hölder inequality. In the last step (5.10) is used.

(c) The bound on W : This case follows 4.3.3(c).

(d) The bound on W_a : The treatment of this case is the only one which differs significantly from the nearest neighbour case. Direct calculation as in (4.20) gives

$$W_a = - \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k e^{ik \cdot a} \cdot [\hat{\tau}^2 \hat{G}^{-1} \hat{G}_{\mu\mu} - 2\hat{\tau}^3 \hat{G}^{-2} \hat{G}_\mu \hat{F}_\mu - \hat{\tau}^3 \hat{G}^{-1} \hat{F}_{\mu\mu} + 2\hat{\tau}^4 \hat{G}^{-2} \hat{F}_\mu^2], \tag{5.21}$$

with

$$\hat{F}_\mu(k) = - \frac{p}{p_L} \partial_\mu D_L(k) - \partial_\mu \hat{\Pi}(k), \tag{5.22}$$

and

$$\hat{F}_{\mu\mu}(k) = - \frac{p}{p_L} \partial_\mu^2 D_L(k) - \partial_\mu^2 \hat{\Pi}(k). \tag{5.23}$$

In (5.21) the first term and the contributions to the second and third terms from the derivatives of $\hat{\Pi}$ can be bounded just as in Sect. 4.3.3(d), using (5.17–20). Their sum is bounded by $O(\epsilon)$.

The fourth term is treated as follows. First we use

$$\hat{F}_\mu^2 \leq 2 \left(\frac{p}{p_L} \partial_\mu D_L \right)^2 + 2 (\partial_\mu \hat{\Pi})^2,$$

and for the second term use the mean value theorem and (5.17) to bound $|\partial_\mu \hat{\Pi}|$ by $O(\varepsilon)|k_\mu|$, and then argue as in Sect. 4.3.3(d) to bound this contribution to (5.21) by $O(\varepsilon)^2$. The contribution from the first term is bounded by

$$\text{const} \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k (\partial_\mu D_L)^2 (1 - D_L)^{-4} = \text{const} W_L \leq \text{const} \varepsilon,$$

using (5.12).

Now we are left with two terms: the contributions from derivatives of D_L in the second and third terms of (5.21). We first consider the second term and use (5.19–20) and the Schwarz inequality to obtain

$$\begin{aligned} \left| \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k 2 \hat{\tau}^3 \hat{G}^{-2} \hat{G}_\mu \partial_\mu D_L \right| &\leq \text{const} \sum_{\mu=1}^d \int d^d k |\hat{G}_\mu \partial_\mu D_L| (1 - D_L)^{-3} \\ &\leq \text{const} \left[\sum_{\mu} \int d^d k (\partial_\mu D_L)^2 (1 - D_L)^{-4} \sum_{\nu} \int d^d l \hat{G}_\nu^2 (1 - D_L)^{-2} \right]^{1/2} \\ &\leq O(\varepsilon) W_L^{1/2} \leq O(\varepsilon^{3/2}). \end{aligned}$$

Finally we consider the term

$$\left| \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k \hat{\tau}^3 \hat{G}^{-1} \partial_\mu^2 D_L e^{ik \cdot a} \right| \leq \text{const} \int d^d k \sum_{\mu=1}^d |\partial_\mu^2 D_L| (1 - D_L)^{-3}. \quad (5.24)$$

The integral on the right side of (5.24) is bounded above by $O(\varepsilon)$, by condition (5.13).

As a result of the above estimates, we have

$$W_a \leq c\varepsilon,$$

where c is independent of ε and K' . Choosing $K' = c/3$ gives the desired result.

(e) The bound on H : The argument of Sect. 4.3.3(e) can be used here. The condition (5.14) is used. \square

5.2. Proof of Lemma 5.1

In this section we prove Lemma 5.1, which together with Sect. 5.1 completes the proof of Theorem 1.2.

5.2.1. *Verification of Conditions (5.6–8)*. The gaussian critical value p_L is given by

$$p_L = \left[\sum_x L^{-d} g(x/L) \right]^{-1}.$$

The sum over x in this expression is a Riemann sum approximation to $\|g\|_1 = 1$, and hence

$$\lim_{L \rightarrow \infty} p_L = 1. \tag{5.25}$$

By definition,

$$S = p_L L^2 \sum_x L^{-d} (x/L)^2 g(x/L) \sim L^2 \|x^2 g\|_1, \tag{5.26}$$

and hence $S = O(L^2)$. Since $g \in L_\infty$, it follows from (5.2) that $\sup_x p_{0x}^{(L)} = O(L^{-d})$. Since $g \in L_2$,

$$\sum_x (p_{0x}^{(L)})^2 = p_L^2 L^{-d} \sum_x L^{-d} g(x/L)^2 \sim L^{-d} \|g\|_2^2 = O(L^{-d}).$$

This gives the condition (5.6), if L is taken sufficiently large.

Similarly, it follows from the fact that $x^2 g \in L_\infty$ that

$$\sup_x p_{0x}^{(L)} |x|^2 = O(L^{-d+2}),$$

which yields (5.7). Finally, the fact that $x^2 g \in L_2$ implies

$$\sum_x (p_{0x}^{(L)})^2 |x|^4 = O(L^{-d+4}),$$

yielding (5.8). The big O notation is used to represent an upper bound involving constants which are independent of L , but may depend on d and g .

5.2.2. Basic Properties of the Gaussian Propagator and (5.9). In this section we show that the condition (5.9) is satisfied, and obtain some further results which will be used to estimate T_L , etc., in the following sections.

Recall the definitions:

$$D_L(k) = p_L \sum_x L^{-d} g(x/L) e^{ik \cdot x}$$

and

$$\hat{C}_L(k) = [1 - D_L(k)]^{-1}.$$

For a function h defined on \mathbb{R}^d , we define a transformation

$$\tilde{h}_L(k) = p_L \sum_x L^{-d} h(x/L) e^{ik \cdot x}.$$

In this notation, $D_L(k) = \tilde{g}_L(k)$. We also use the notation $|I|$ to represent the cardinality of a set $I \subset \{1, \dots, d\}$, and write $\partial^I = \prod_{v \in I} \frac{\partial}{\partial x_v}$. The expressions $\partial^I h$ and $\|\partial^I h\|_1$ are to be interpreted in terms of distributions.

Lemma 5.4. *Suppose that h vanishes at infinity, that $\frac{\partial^d h}{\partial x_1 \partial x_2 \cdots \partial x_d}$ is piecewise continuous, and that \tilde{h}_L exists for all L . Then for L sufficiently large,*

$$|\tilde{h}_L(k)| \leq 2 \|h\|_1. \tag{5.27}$$

Also, for any $I \subset \{1, \dots, d\}$ and for any k ,

$$|\tilde{h}_L(k)| \leq 2 \|\partial^I h\|_1 \left| \prod_{v \in I} 2L \sin(k_v/2) \right|^{-1}. \tag{5.28}$$

Proof. The bound (5.27) follows from (5.25) and the fact that $\sum_x L^{-d} |h(x/L)| \rightarrow \|h\|_1$ as $L \rightarrow \infty$. The bound (5.28) is proved using summation by parts, as follows. To simplify the notation, we assume that $1 \in I$. Then

$$\begin{aligned} \tilde{h}_L(k) &= p_L \sum_x L^{-d} h(x/L) e^{-ik_1/2} [e^{ik_1(x_1+1)} - e^{ik_1x_1}] [2i \sin(k_1/2)]^{-1} \exp \left[\sum_{v=2}^d ik_v x_v \right] \\ &= -p_L e^{-ik_1/2} [2i \sin(k_1/2)]^{-1} \sum_x L^{-d} [h(x/L) - h(x/L - \hat{e}_1/L)] e^{ik \cdot x}, \end{aligned} \tag{5.29}$$

where $\hat{e}_1 = (1, 0, \dots, 0)$. To avoid nonilluminating complications, we consider henceforth only the case of continuously differentiable h ; the general case can be treated similarly. By the mean value theorem (5.29) is equal to

$$-p_L e^{-ik_1/2} [2i \sin(k_1/2)]^{-1} \sum_x L^{-d} \frac{1}{L} \partial_1 h(x^*/L) e^{ik \cdot x},$$

where x^* is a point on the line joining x/L to $(x - \hat{e}_1)/L$. Iteration of this procedure gives

$$\tilde{h}_L(k) = (-1)^{|I|} p_L \prod_{v \in I} [e^{-ik_v/2} / 2L i \sin(k_v/2)] \sum_x L^{-d} \partial^I h(x^{**}/L) e^{ik \cdot x},$$

where $\|(x^{**} - x)/L\|_\infty \leq L^{-1}$. Therefore

$$|\tilde{h}_L(k)| \leq p_L \prod_{v \in I} |2L \sin(k_v/2)|^{-1} \sum_x L^{-d} |\partial^I h(x^{**}/L)| \sim \prod_{v \in I} |2L \sin(k_v/2)|^{-1} \|\partial^I h\|_1.$$

This gives the desired bound. \square

The following lemma is proved using Lemma 5.4. Fix $M > 0$ such that the integral of g over the region $\|x\|_1 \leq M$ is strictly positive. We write $\delta = 3\pi \|\partial_1 g\|_1 / 2L$ and $\delta_1 = \pi / LM$, and use c to denote a constant which is independent of L and whose value may change from one occurrence to another. We also fix a small $\varepsilon > 0$ (e.g., $\varepsilon = 1/5$). It seems likely that the bounds in which ε appears, in the remainder of this section, can be strengthened to the corresponding $\varepsilon = 0$ bounds; however the bounds we obtain are sufficient for our needs.

Lemma 5.5.

$$\lim_{L \rightarrow \infty} D_L(k) = 0, \quad \text{for all } k \neq 0. \tag{5.30}$$

For L sufficiently large, the following inequalities hold:

$$|D_L(k)| \leq \frac{2}{3}, \quad \text{for } \|k\|_\infty \geq \delta, \tag{5.31}$$

$$1 - D_L(k) \geq ck^2 L^2, \quad \text{for } \|k\|_\infty \leq \delta_1, \tag{5.32}$$

$$1 - D_L(k) \geq ck^2 L^{2-\varepsilon}, \quad \text{for } \delta_1 \leq \|k\|_\infty \leq \delta \text{ (if } \delta_1 < \delta), \tag{5.33}$$

$$|\partial_\mu^s D_L(k)| \leq 2L^s \|\partial^I(x_\mu^s g)\|_1 \prod_{v \in I} |2L \sin(k_v/2)|^{-1}, \quad \text{for } s = 0, 1, 2 \text{ and all } k, I, \tag{5.34}$$

$$|\partial_\mu D_L(k)| \leq 2L^2 |k_\mu| \|x_\mu^2 g\|_1, \quad \text{for all } k. \tag{5.35}$$

Proof. Equation (5.30) follows from (5.28) with $h = g$ and $I = \{\mu\}$, $\mu = 1, \dots, d$. To prove (5.31), suppose that $|k_v| \geq \delta$, and let $I = \{v\}$. Then by (5.28) and the definition of δ ,

$$|D_L(k)| \leq \pi L^{-1} |k_v|^{-1} \|\partial_v g\|_1 \leq 2/3.$$

To prove (5.32) we suppose $\|k\|_\infty \leq \delta_1$, and use symmetry to write

$$\begin{aligned} 1 - D_L(k) &= p_L \sum_x L^{-d} g(x/L) (1 - \cos k \cdot x) \geq p_L 2\pi^{-2} \sum_{x: |k \cdot x| \leq \pi} L^{-d} g(x/L) (k \cdot x)^2 \\ &\geq p_L 2\pi^{-2} \sum_{x: \|x\|_1 \leq LM} L^{-d} g(x/L) (k \cdot x)^2 \\ &= p_L 2\pi^{-2} k^2 L^2 \sum_{x: \|x\|_1 \leq LM} L^{-d} g(x/L) (x_1/L)^2 \\ &\sim 2\pi^{-2} k^2 L^2 \int_{\|x\|_1 \leq M} g(x) x_1^2 d^d x > ck^2 L^2, \end{aligned}$$

for L sufficiently large.

The inequality (5.33) is proved as follows. Fix k such that $\delta_1 \leq \|k\|_\infty \leq \delta$, and let $X_k = \{x: 1 - \cos k \cdot x > L^{-\epsilon}\}$. Then

$$\begin{aligned} 1 - D_L(k) &\geq p_L L^{-\epsilon} \sum_{x \in X_k} L^{-d} g(x/L) \geq k^2 d^{-1} \delta^{-2} L^{-\epsilon} p_L \sum_{x \in X_k} L^{-d} g(x/L) \\ &\sim ck^2 L^{2-\epsilon} \left[1 - \sum_{x \notin X_k} L^{-d} g(x/L) \right]. \end{aligned}$$

It suffices to show that the sum on the right side goes to zero as L goes to infinity. The domain of summation consists of those x for which $1 - \cos k \cdot x \leq L^{-\epsilon}$. Since for any x we can find an integer n (depending on $k \cdot x$) such that $1 - \cos k \cdot x \geq 2\pi^{-2} (k \cdot x - 2n\pi)^2$, this domain is contained in the set of all x such that $x/L \in Y_k$, where

$$Y_k = \left\{ y \in \mathbb{Z}^d: |\hat{k} \cdot y - 2n\pi|^2 \leq \frac{\pi^2}{2} L^{-\epsilon}, \text{ for some } n \in \mathbb{Z} \right\}.$$

Here we have written $\hat{k} \equiv kL$, with $\|\hat{k}\|_\infty \in [\pi/M, 3\pi \|\partial_1 g\|_1/2]$. For $R > 0$, let $B_R = \{y: \|y\|_\infty \leq R\}$. Since g decays exponentially, given $\varrho > 0$ we can choose an R such that

$$\sum_{\frac{x}{L} \notin B_R} L^{-d} g(x/L) < \varrho.$$

But then we can choose L sufficiently large that

$$\sum_{\frac{x}{L} \in Y_k \cap B_R} L^{-d} g(x/L) < \varrho,$$

since the sum on the left side converges to zero as L goes to infinity, because g is bounded and the measure of $Y_k \cap B_R$ goes to zero as L goes to infinity. This proves (5.33).

The inequality (5.34) follows from (5.27) and (5.28). To prove (5.35) we use symmetry to write

$$\begin{aligned} |\partial_\mu D_L(k)| &= \left| p_L \sum_x L^{-d} g(x/L) x_\mu \sin(k_\mu x_\mu) \prod_{v \neq \mu} e^{ik_v x_v} \right| \\ &\leq p_L \sum_x L^{-d} g(x/L) |x_\mu| |k_\mu x_\mu| \sim L^2 |k_\mu| \|x_\mu^2 g\|_1. \quad \square \end{aligned}$$

We now use the bounds given in Lemma 5.5 to estimate $\hat{C}_L(k)$. These estimates will in turn be used to estimate T_G and W_G . The condition (5.9) is an immediate consequence of (5.37) below, if we take L sufficiently large.

Lemma 5.6. *For*

$$k \neq 0, \quad \hat{C}_L(k) \rightarrow 1 \text{ as } L \rightarrow \infty. \tag{5.36}$$

In addition

$$0 \leq \hat{C}_L(k) \leq \begin{cases} 3 & \text{if } \|k\|_\infty \geq \delta \\ O(L^{-2+\varepsilon})|k|^{-2} & \text{if } \|k\|_\infty \leq \delta \end{cases}, \tag{5.37}$$

$$|\hat{C}_L(k) - 1| \leq \begin{cases} 3|D_L(k)| & \text{if } \|k\|_\infty \geq \delta \\ O(L^{-2+\varepsilon})|k|^{-2} & \text{if } \|k\|_\infty \leq \delta \end{cases}, \tag{5.38}$$

$$|\partial_\mu \hat{C}_L(k)| \leq \begin{cases} 9|\partial_\mu D_L(k)| & \text{if } \|k\|_\infty \geq \delta \\ O(L^{-2+2\varepsilon})|k_\mu| |k|^{-4} & \text{if } \|k\|_\infty \leq \delta \end{cases}. \tag{5.39}$$

Here the big O denotes upper bounds involving constants which are independent of L , but may depend on d or g . Also $|k|$ denotes the euclidean length of k .

Proof. The limit (5.36) follows immediately from (5.4) and (5.30). For (5.37), we use (5.31–33). Since $\hat{C}_L(k) - 1 = D_L(k)/[1 - D_L(k)]$, the first bound in (5.38) follows from (5.31), and the second from (5.32–33) and the fact that $|D_L(k)| \leq 1$. Similarly (5.39) follows from calculation of the derivative, (5.31–33) and (5.35). \square

5.2.3. *Conditions (5.10), (5.11), (5.14).* The following lemma will be used to estimate T_G for this model.

Lemma 5.7. *For $d > 2$,*

$$(2\pi)^{-d} \int d^d k |\hat{C}_L(k) - 1| \leq O(L^{-d+\varepsilon}) \left[(\ln L)^d + \frac{c_d}{d-2} \right].$$

For $d > 2N, N \geq 2$,

$$(2\pi)^{-d} \int d^d k |\hat{C}_L(k) - 1|^N \leq O(L^{-d+\varepsilon N}) \left[1 + \frac{c'_d}{d-2N} \right].$$

The constants c_d and c'_d depend only on d and remain bounded as $d \downarrow 2$ in the first bound and $d \downarrow 2N$ in the second.

Proof. We divide the domain of integration into two parts: $\|k\|_\infty \leq \delta$ and $\|k\|_\infty \geq \delta$, and use (5.38) to bound the integrand. This gives

$$\begin{aligned} (2\pi)^{-d} \int d^d k |\hat{C}_L(k) - 1|^N &\leq 3^N (2\pi)^{-d} \int_{\|k\|_\infty \geq \delta} d^d k |D_L(k)|^N \\ &\quad + O(L^{-(2-\varepsilon)N}) (2\pi)^{-d} \int_{\|k\|_\infty \leq \delta} d^d k |k|^{-2N} \\ &\leq 3^N (2\pi)^{-d} \int_{\|k\|_\infty \geq \delta} d^d k |D_L(k)|^N \\ &\quad + O(L^{-(2-\varepsilon)N}) \Omega_d (2\pi)^{-d} \frac{(\sqrt{d}\delta)^{d-2N}}{d-2N}, \end{aligned}$$

where Ω_d is the volume of the unit sphere in \mathbb{R}^d . The first term on the right side is estimated using (5.34), as follows.

The region $\|k\|_\infty \geq \delta$ is the disjoint union, over $I \subset \{1, 2, \dots, d\}$, $I \neq \emptyset$, of

$$R_I \equiv \{k \in \mathbb{R}^d \mid \delta < |k_\nu| < \pi \text{ for } \nu \in I, |k_\mu| \leq \delta \text{ for } \mu \notin I\}.$$

By (5.34),

$$\begin{aligned} (2\pi)^{-d} \int_{R_I} d^d k |D_L(k)|^N &\leq 2^N \|\partial^I g\|_1^N (2\pi)^{-d} \int_{R_I} d^d k \prod_{\nu \in I} [\pi(2L|k_\nu|)^{-1}]^N \\ &\leq c^N L^{-N|I|} \delta^{d-|I|} \begin{cases} |\ln \delta|^{|I|} & \text{if } N = 1 \\ \delta^{|I|(1-N)} & \text{if } N > 1 \end{cases}. \end{aligned}$$

This gives the bounds in the statement of the lemma. \square

Corollary 5.8. For $d > 2$,

$$|C_L(0, 0) - 1| \leq O(L^{-d+\varepsilon}) \left[(\ln L)^d + \frac{c_d}{d-2} \right]. \tag{5.40}$$

For $d > 6$,

$$T_L \leq O(L^{-d+3\varepsilon}) \left(1 + \frac{c'_d}{d-6} \right). \tag{5.41}$$

Proof. The first inequality is an immediate consequence of the first statement in Lemma 5.7. The second follows from the second statement in the lemma and the fact that by definition

$$\begin{aligned} T_L &= \sum_{x,y} C_L(0, x) C_L(x, y) C_L(y, 0) - C_L(0, 0)^3 \\ &= (2\pi)^{-d} \int d^d k (\hat{C}_L(k) + 2)(\hat{C}_L(k) - 1)^2 - (C_L(0, 0) + 2)(C_L(0, 0) - 1)^2. \quad \square \end{aligned}$$

In view of (5.26), Corollary 5.8 ensures that the conditions (5.10) and (5.11) are satisfied for sufficiently large L . The condition (5.14) follows immediately from the following lemma.

Lemma 5.9. For $d > 6$,

$$\lim_{L \rightarrow \infty} \left(\text{diagram} \right) = 1, \tag{5.42}$$

where the lines in the diagram denote the gaussian propagator $\hat{C}_L(k)$.

Proof. Fix $d > 6$. By (5.36) and (5.37), $|\hat{C}_L(k)|$ is bounded above by $|k|^{-2}$ for k near zero, and $\hat{C}_L(k)$ approaches one pointwise as L goes to infinity. Since the replacement of $\hat{C}_L(k)$ by $|k|^{-2}$ in the diagram in (5.42) yields a finite result for $d > 6$, the conclusion follows by the dominated convergence theorem. \square

A similar argument can be employed to prove that as $L \rightarrow \infty$, T_L converges to zero and $C_L(0, 0)$ converges to one. However this is not sufficient for the conditions (5.10–11), and thus we presented the more detailed bounds of Corollary 5.8.

5.2.4. Bounds on Quantities with Derivatives: Conditions (5.12) and (5.13). The condition (5.12) on W_L is an immediate consequence of the following lemma.

Lemma 5.10. For $d > 6$,

$$W_L \leq O(L^{2+4\epsilon-d}) \left(1 + \frac{c_d}{d-6} \right). \tag{5.43}$$

Proof. By definition,

$$W_L = \sum_{\mu=1}^d (2\pi)^{-d} \int d^d k (\partial_\mu \hat{C}_L(k))^2.$$

We divide the domain of integration as in Lemma 5.7 and use (5.39) to obtain

$$W_L \leq 81 (2\pi)^{-d} \int_{\|k\|_\infty \geq \delta} d^d k |\partial_\mu D_L(k)|^2 + O(L^{-4+4\epsilon}) \int_{\|k\|_\infty \leq \delta} d^d k |k|^{-6}.$$

The second term on the right side gives the second term on the right side of (5.43). The integral in the first term on the right side can be estimated by writing the integral as a sum over integrals over R_I , exactly as was done in the proof of Lemma 5.7, and using (5.34) with $s = 1$. The only difference here is the extra factor of L^2 which arises from $s = 1$ in (5.34). \square

Finally we prove the following bound, which gives the condition (5.13).

Lemma 5.11. For $d > 6$,

$$(2\pi)^{-d} \int d^d k |\partial_\mu^2 D_L| (1 - D_L)^{-3} \leq O(L^{2+3\epsilon-d}) \left[(\ln L)^d + \frac{c_d}{d-6} \right]. \tag{5.44}$$

Proof. Using (5.37) and (5.33) gives

$$(2\pi)^{-d} \int d^d k |\partial_\mu^2 D_L| (1 - D_L)^{-3} \leq 27 (2\pi)^{-d} \int_{\|k\|_\infty \geq \delta} d^d k |\partial_\mu^2 D_L| \\ + O(L^{-6+3\epsilon+2}) \int_{\|k\|_\infty \leq \delta} d^d k |k|^{-6}.$$

The second term on the right side gives the second term on the right side of (5.44). For the integral over $\|k\|_\infty \geq \delta$ we use (5.34) with $s = 2$, as in Lemmas 5.7 and 5.10, to obtain the desired result. \square

Acknowledgements. It is a pleasure to thank Michael Aizenman for many stimulating and valuable conversations concerning the subject of this paper. G.S. also wishes to thank him for making possible a visit to the Courant Institute in May of 1988, during which this work was begun. We thank David Brydges, Chuck Newman, Alan Sokal, and Hal Tasaki for interesting and useful conversations.

References

1. Aizenman, M.: Geometric analysis of φ^4 fields and Ising models, Parts I and II. *Commun. Math. Phys.* **86**, 1–48 (1982)
2. Aizenman, M., Barsky, D.J.: Sharpness of the phase transition in percolation models. *Commun. Math. Phys.* **108**, 489–526 (1987)
3. Aizenman, M., Fernández, R.: On the critical behaviour of the magnetization in high dimensional Ising models. *J. Stat. Phys.* **44**, 393–454 (1986)
4. Aizenman, M., Graham, R.: On the renormalized coupling constant and the susceptibility in $\lambda\varphi^4$ field theory and the Ising model in four dimensions. *Nucl. Phys.* **B225** [FS9], 261–288 (1983)
5. Aizenman, M., Kesten, H., Newman, C.M.: Uniqueness of the infinite cluster and continuity of connectivity functions for short and long range percolation. *Commun. Math. Phys.* **111**, 505–531 (1987)
6. Aizenman, M., Newman, C.M.: Tree graph inequalities and critical behaviour in percolation models. *J. Stat. Phys.* **36**, 107–143 (1984)
7. Aizenman, M., Simon, B.: Local Ward identities and the decay of correlations in ferromagnets. *Commun. Math. Phys.* **77**, 137–143 (1980)
8. Barsky, D.J., Aizenman, M.: Percolation critical exponents under the triangle condition. Preprint (1988)
9. van den Berg, J., Kesten, H.: Inequalities with applications to percolation and reliability, *J. Appl. Prob.* **22**, 556–569 (1985)
10. Broadbent, S.R., Hammersley, J.M.: Percolation processes. I. Crystals and mazes. *Proc. Camb. Phil. Soc.* **53**, 629–641 (1957)
11. Brydges, D.C., Fröhlich, J., Sokal, A.D.: A new proof of the existence and nontriviality of the continuum φ_2^4 and φ_3^4 quantum field theories. *Commun. Math. Phys.* **91**, 141–186 (1983)
12. Brydges, D.C., Spencer, T.: Self-avoiding walk in 5 or more dimensions. *Commun. Math. Phys.* **97**, 125–148 (1985)
13. Chayes, J. T., Chayes, L.: On the upper critical dimension of Bernoulli percolation. *Commun. Math. Phys.* **113**, 27–48 (1987)
14. Essam, J.W.: Percolation Theory. *Rep. Prog. Phys.* **43**, 833–912 (1980)
15. Fröhlich, J.: On the triviality of φ_d^4 theories and the approach to the critical point in $d \geq 4$ dimensions. *Nucl. Phys.* **B200** [FS4], 281–296 (1982)
16. Fröhlich, J., Simon, B., Spencer, T.: Infrared bounds, phase transitions, and continuous symmetry breaking. *Commun. Math. Phys.* **50**, 79–95 (1976)

17. Grimmett, G.: Percolation, Berlin Heidelberg New York: Springer 1989
18. Hammersley, J.M.: Percolation processes. Lower bounds for the critical probability. *Ann. Math. Statist.* **28**, 790–795 (1957)
19. Hammersley, J.M.: Bornes supérieures de la probabilité critique dans un processus de filtration. In: *Le Calcul des Probabilités et ses Applications 17–37* CNRS Paris (1959)
20. Hara, T.: Mean field critical behaviour of correlation length for percolation in high dimensions. Preprint (1989)
21. Hara, T., Slade, G.: On the upper critical dimension of lattice trees and lattice animals. Submitted to *J. Stat. Phys.*
22. Hara, T., Slade, G.: Unpublished
23. Kac, M., Uhlenbeck, G.E., Hemmer, P.C.: On the van der Waals theory of the vapor-liquid equilibrium. I. Discussion of a one-dimensional model. *J. Math. Phys.* **4**, 216–288 (1963)
24. Kesten, H.: Percolation theory and first passage percolation. *Ann. Probab.* **15**, 1231–1271 (1987)
25. Lawler, G.: The infinite self-avoiding walk in high dimensions. To appear in *Ann. Probab.* (1989)
26. Lebowitz, J.L., Penrose, O.: Rigorous treatment of the van der Waals-Maxwell theory of the liquid-vapor transition. *J. Math. Phys.* **7**, 98–113 (1966)
27. Menshikov, M.V., Molchanov, S.A., Sidorenko, A.F.: Percolation theory and some applications. *Itogi Nauki i Tekhniki* (Series of Probability Theory, Mathematical Statistics, Theoretical Cybernetics) **24**, 53–110 (1986). English translation. *J. Soviet Math.* **42**, 1766–1810 (1988)
28. Nguyen, B.G.: Gap exponents for percolation processes with triangle condition. *J. Stat. Phys.* **49**, 235–243 (1987)
29. Park, Y.M.: Direct estimates on intersection probabilities of random walks. To appear in *J. Stat. Phys.*
30. Russo, L.: On the critical percolation probabilities. *Z. Wahrsch. Verw. Gebiete.* **56**, 229–237 (1981)
31. Slade, G.: The diffusion of self-avoiding random walk in high dimensions. *Commun. Math. Phys.* **110**, 661–683 (1987)
32. Slade, G.: The scaling limit of self-avoiding random walk in high dimensions. *Ann. Probab.* **17**, 91–107 (1989)
33. Slade, G.: Convergence of self-avoiding random walk to Brownian motion in high dimensions. *J. Phys. A: Math. Gen.* **21**, L417–L420 (1988)
34. Slade, G.: The lace expansion and the upper critical dimension for percolation, Lectures notes from the A.M.S. Summer Seminar, Blacksburg, June 1989
35. Sokal, A.D.: A rigorous inequality for the specific heat of an Ising or φ^4 ferromagnet. *Phys. Lett.* **71A**, 451–453 (1979)
36. Sokal, A.D., Thomas, L.E.: Exponential convergence to equilibrium for a class of random walk models. *J. Stat. Phys.* **54**, 797–828 (1989)
37. Stauffer, D.: Introduction to percolation theory. Taylor and Francis, London Philadelphia (1985)
38. Tasaki, H.: Hyperscaling inequalities for percolation. *Commun. Math. Phys.* **113**, 49–65 (1987)
39. Tasaki, H.: Private communication
40. Yang, W., Klein, D.: A note on the critical dimension for weakly self avoiding walks. *Prob. Th. Rel. Fields* **79**, 99–114 (1988)
41. Ziff, R.M., Stell, G.: Critical behaviour in three-dimensional percolation: Is the percolation threshold a Lifshitz point? Preprint (1988)

Communicated by M. Aizenman

Received February 8, 1989; in revised form July 6, 1989

