

UC Berkeley

UC Berkeley Previously Published Works

Title

Measurement of CP observables for the decays $B \pm \rightarrow D\bar{C}P0K\pm$

Permalink

<https://escholarship.org/uc/item/7d38p8t3>

Journal

Physical Review D - Particles, Fields, Gravitation and Cosmology, 73(5)

ISSN

1550-7998

Authors

Aubert, B
Barate, R
Boutigny, D
et al.

Publication Date

2006

DOI

10.1103/PhysRevD.73.051105

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Measurement of CP observables for the decays $B^\pm \rightarrow D_{CP}^0 K^\pm$

- B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. S. Best,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,^{6,*} Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulson,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu Todyshev,¹² M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroeseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretskii,¹⁹ D. G. Hitlin,¹⁹ J. S. Minamora,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winkelmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ B. Spaan,²³ T. Brandt,²⁴ M. Dickopp,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneau,²⁵ P. Grenier,^{25,†} E. Latour,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,‡} M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² J. A. Nash,³² M. B. Nikolic,³² W. Panduro Vazquez,³² X. Chai,³³ M. J. Charles,³³ W. F. Mader,³³ U. Mallik,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ G. Schott,³⁵ N. Arnaud,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ F. Le Diberder,³⁶ V. Lepeltier,³⁶ A. M. Lutz,³⁶ A. Oyanguren,³⁶ T. C. Petersen,³⁶ S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ A. Stocchi,³⁶ W. F. Wang,³⁶ G. Wormser,³⁶ C. H. Cheng,³⁷ D. J. Lange,³⁷ D. M. Wright,³⁷ A. J. Bevan,³⁸ C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ K. A. George,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ C. L. Brown,⁴⁰ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ M. G. Green,⁴⁰ D. A. Hopkins,⁴⁰ P. S. Jackson,⁴⁰ T. R. McMahon,⁴⁰ S. Ricciardi,⁴⁰ F. Salvatore,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴² M. P. Kelly,⁴² G. D. Lafferty,⁴² M. T. Naisbit,⁴² J. C. Williams,⁴² J. I. Yi,⁴² C. Chen,⁴³ W. D. Hulsbergen,⁴³ A. Jawahery,⁴³ D. Kovalskyi,⁴³ C. K. Lae,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ R. Kofler,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ S. Saremi,⁴⁴ H. Staengle,⁴⁴ S. Y. Willocq,⁴⁴ R. Cowan,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ H. Kim,⁴⁶ P. M. Patel,⁴⁶ C. T. Potter,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ V. Lombardo,⁴⁷ F. F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ J. Reidy,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ P. Taras,⁴⁹ F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ N. Cavallo,^{51,§} G. De Nardo,⁵¹ F. Fabozzi,^{51,§} C. Gatto,⁵¹ L. Lista,⁵¹ D. Monorchio,⁵¹ P. Paolucci,⁵¹ D. Piccolo,⁵¹ C. Sciacca,⁵¹ M. Baak,⁵² H. Bulten,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ T. Allmendinger,⁵⁴ G. Benelli,⁵⁴ K. K. Gan,⁵⁴ K. Honscheid,⁵⁴ D. Hufnagel,⁵⁴ P. D. Jackson,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ T. Pulliam,⁵⁴ A. M. Rahimi,⁵⁴ R. Ter-Antonyan,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵

B. AUBERT *et al.*

PHYSICAL REVIEW D 73, 051105 (2006)

- F. Galeazzi,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ M. Benayoun,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ B. L. Hartfiel,⁵⁷ M. J. J. John,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ L. Roos,⁵⁷ G. Therin,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ J. Panetta,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ D. E. Wagoner,⁶¹ J. Biesiada,⁶² N. Danielson,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,⁶³ A. D'Orazio,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ E. O. Olaiya,⁶⁵ F. F. Wilson,⁶⁵ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Legendre,⁶⁶ B. Mayer,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ W. Park,⁶⁷ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ J. R. Wilson,⁶⁷ T. Abe,⁶⁸ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ J. C. Dingfelder,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hrynowska,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ D. B. MacFarlane,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va'vra,⁶⁸ N. van Bakel,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ L. Wilden,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ V. Jain,⁷⁰ B. Pan,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappeler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶ F. Martinez-Vidal,⁷⁶ R. S. Panvini,^{77,||} Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ M. T. Graham,⁸⁰ J. J. Hollar,⁸⁰ J. R. Johnson,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰ A. Mihalyi,⁸⁰ A. K. Mohapatra,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy⁴Institute of High Energy Physics, Beijing 100039, China⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany⁹University of Bristol, Bristol BS8 1TL, United Kingdom¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹³University of California at Irvine, Irvine, California 92697, USA¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA¹⁵University of California at Riverside, Riverside, California 92521, USA¹⁶University of California at San Diego, La Jolla, California 92093, USA¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA¹⁹California Institute of Technology, Pasadena, California 91125, USA²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA²¹University of Colorado, Boulder, Colorado 80309, USA

MEASUREMENT OF CP OBSERVABLES FOR THE DECAYS $B^\pm \rightarrow D_{CP}^0 K^\pm$ PHYSICAL REVIEW D 73, 051105 (2006)²²*Colorado State University, Fort Collins, Colorado 80523, USA*²³*Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany*²⁴*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*²⁵*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*²⁶*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*²⁷*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*²⁹*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*³¹*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*³²*Imperial College London, London, SW7 2AZ, United Kingdom*³³*University of Iowa, Iowa City, Iowa 52242, USA*³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*³⁵*Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*³⁶*Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*³⁷*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*³⁸*University of Liverpool, Liverpool L69 7ZE, United Kingdom*³⁹*Queen Mary, University of London, E1 4NS, United Kingdom*⁴⁰*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*⁴¹*University of Louisville, Louisville, Kentucky 40292, USA*⁴²*University of Manchester, Manchester M13 9PL, United Kingdom*⁴³*University of Maryland, College Park, Maryland 20742, USA*⁴⁴*University of Massachusetts, Amherst, Massachusetts 01003, USA*⁴⁵*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*⁴⁶*McGill University, Montréal, Québec, Canada H3A 2T8*⁴⁷*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*⁴⁸*University of Mississippi, University, Mississippi 38677, USA*⁴⁹*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*⁵⁰*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*⁵¹*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*⁵²*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*⁵³*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁴*Ohio State University, Columbus, Ohio 43210, USA*⁵⁵*University of Oregon, Eugene, Oregon 97403, USA*⁵⁶*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*⁵⁷*Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France*⁵⁸*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*⁵⁹*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*⁶⁰*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*⁶¹*Prairie View A&M University, Prairie View, Texas 77446, USA*⁶²*Princeton University, Princeton, New Jersey 08544, USA*⁶³*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*⁶⁴*Universität Rostock, D-18051 Rostock, Germany*⁶⁵*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*⁶⁶*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*⁶⁷*University of South Carolina, Columbia, South Carolina 29208, USA*⁶⁸*Stanford Linear Accelerator Center, Stanford, California 94309, USA*⁶⁹*Stanford University, Stanford, California 94305-4060, USA*⁷⁰*State University of New York, Albany, New York 12222, USA*⁷¹*University of Tennessee, Knoxville, Tennessee 37996, USA*⁷²*University of Texas at Austin, Austin, Texas 78712, USA*⁷³*University of Texas at Dallas, Richardson, Texas 75083, USA*⁷⁴*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*⁷⁵*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*⁷⁶*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*⁷⁷*Vanderbilt University, Nashville, Tennessee 37235, USA*⁷⁸*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*⁷⁹*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*⁸⁰*University of Wisconsin, Madison, Wisconsin 53706, USA*⁸¹*Yale University, New Haven, Connecticut 06511, USA*

(Received 27 December 2005; published 15 March 2006)

We present a study of the decay $B^- \rightarrow D_{(CP)}^0 K^-$ and its charge conjugate, where $D_{(CP)}^0$ is reconstructed in CP -even, CP -odd, and non- CP flavor eigenstates, based on a sample of $232 \times 10^6 Y(4S) \rightarrow B\bar{B}$ decays collected with the *BABAR* detector at the PEP-II e^+e^- storage ring. We measure the partial-rate charge asymmetries $A_{CP\pm}$ and the ratios $R_{CP\pm}$ of the $B \rightarrow D^0 K$ decay branching fractions as measured in $CP\pm$ and non- CP D^0 decays: $A_{CP+} = 0.35 \pm 0.13(\text{stat}) \pm 0.04(\text{syst})$, $A_{CP-} = -0.06 \pm 0.13(\text{stat}) \pm 0.04(\text{syst})$, $R_{CP+} = 0.90 \pm 0.12(\text{stat}) \pm 0.04(\text{syst})$, and $R_{CP-} = 0.86 \pm 0.10(\text{stat}) \pm 0.05(\text{syst})$.

DOI: 10.1103/PhysRevD.73.051105

PACS numbers: 11.30.Er, 13.25.Hw, 14.40.Nd

A theoretically clean measurement of the angle $\gamma = \arg(-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*)$ of the Cabibbo-Kobayashi-Maskawa matrix V can be obtained from the study of $B^- \rightarrow D^{(*)0} K^{(*)-}$ decays [1] by exploiting the interference between the $b \rightarrow c\bar{s}$ and $b \rightarrow u\bar{s}$ decay amplitudes [2,3]. Among the proposed methods, the one originally suggested by Gronau, London, and Wyler (GLW) exploits the interference between $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow \bar{D}^0 K^-$ when the D^0 and \bar{D}^0 mesons decay to the same CP eigenstate.

The results of the GLW analyses are usually expressed in terms of the ratios $R_{CP\pm}$ of charge-averaged partial rates and of the partial-rate charge asymmetries $A_{CP\pm}$,

$$R_{CP\pm} = \frac{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) + \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)}{[\Gamma(B^- \rightarrow D^0 K^-) + \Gamma(B^+ \rightarrow \bar{D}^0 K^+)]/2}, \quad (1)$$

$$A_{CP\pm} = \frac{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) - \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)}{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) + \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)}. \quad (2)$$

Here, $D_{CP\pm}^0 = (D^0 \pm \bar{D}^0)/\sqrt{2}$ are the CP eigenstates of the neutral D meson system, and we have followed the notation used in [4]. Neglecting $D^0 - \bar{D}^0$ mixing [5], the observables $R_{CP\pm}$ and $A_{CP\pm}$ are related to the angle γ , the magnitude r of the ratio of the amplitudes for the processes $B^- \rightarrow \bar{D}^0 K^-$ and $B^- \rightarrow D^0 K^-$, and the relative strong phase δ between these two amplitudes, through the relations $R_{CP\pm} = 1 + r^2 \pm 2r \cos\delta \cos\gamma$ and $A_{CP\pm} = \pm 2r \sin\delta \sin\gamma / R_{CP\pm}$ [2]. Theoretical expectations for r are in the range ≈ 0.1 – 0.2 [2,6], in agreement with the 90% C.L. upper limits on r set by *BABAR* ($r < 0.23$) and *Belle* ($r < 0.18$) through the study of $B^- \rightarrow DK^-$, $D \rightarrow K^+ \pi^-$ decays [7].

In this paper we present the measurements of $R_{CP\pm}$ and $A_{CP\pm}$. The ratios $R_{CP\pm}$ are computed using the relations $R_{CP\pm} \approx R_\pm / R$, where the quantities $R_{(\pm)}$ are defined as

^{*}Also with the Johns Hopkins University, Baltimore, MD 21218, USA.

[†]Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France.

[‡]Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

[§]Also with Università della Basilicata, Potenza, Italy.

^{||}Deceased.

$$R_{(\pm)} = \frac{\mathcal{B}(B^- \rightarrow D_{(CP\pm)}^0 K^-) + \mathcal{B}(B^+ \rightarrow \bar{D}_{(CP\pm)}^0 K^+)}{\mathcal{B}(B^- \rightarrow D_{(CP\pm)}^0 \pi^-) + \mathcal{B}(B^+ \rightarrow \bar{D}_{(CP\pm)}^0 \pi^+)}. \quad (3)$$

Several systematic uncertainties cancel out in the measurement of these double ratios. We also express the CP -sensitive observables in terms of three independent quantities:

$$x_\pm = \frac{R_{CP+}(1 \mp A_{CP+}) - R_{CP-}(1 \mp A_{CP-})}{4}, \quad (4)$$

$$r^2 = x_\pm^2 + y_\pm^2 = \frac{R_{CP+} + R_{CP-} - 2}{2}, \quad (5)$$

where $x_\pm = r \cos(\delta \pm \gamma)$ and $y_\pm = r \sin(\delta \pm \gamma)$ are the same CP parameters as were measured by the *BABAR* Collaboration with $B^- \rightarrow DK^-$, $D \rightarrow K_S^0 \pi^- \pi^+$ decays [8]. This choice allows the results of the two measurements to be expressed in a consistent manner.

The measurements use a sample of 232 million $Y(4S)$ decays into $B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II asymmetric-energy B factory. Since the *BABAR* detector is described in detail elsewhere [9], only the components that are crucial to this analysis are summarized here. Charged-particle tracking is provided by a five-layer silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH). For charged-particle identification, ionization energy loss in the DCH and SVT, and Cherenkov radiation detected in a ring-imaging device (DIRC) are used. Photons are identified by the electromagnetic calorimeter (EMC), which comprises 6580 thallium-doped CsI crystals. These systems are mounted inside a 1.5-T solenoidal superconducting magnet. We use the GEANT [10] software to simulate interactions of particles traversing the detector, taking into account the varying accelerator and detector conditions.

We reconstruct $B^- \rightarrow D^0 h^-$ decays, where the prompt track h^- is a kaon or a pion. D^0 candidates are reconstructed in the CP -even eigenstates $\pi^-\pi^+$ and K^-K^+ (D_{CP+}^0), in the CP -odd eigenstates $K_S^0\pi^0$, $K_S^0\phi$ and $K_S^0\omega$ (D_{CP-}^0), and in the non- CP , flavor eigenstate $K^-\pi^+$. ϕ candidates are reconstructed in the K^-K^+ channel and ω candidates in the $\pi^-\pi^+\pi^0$ channel. We optimize our event selection to minimize the statistical error on the $B^- \rightarrow D_{(CP)}^0 K^-$ signal yield, determined for each D^0 decay channel using simulated signal and background events.

The prompt particle h is required to have a momentum greater than $1.4 \text{ GeV}/c$ and the number of photons associated to its Cherenkov ring is required to be greater than four to improve the quality of the reconstruction. We reject a candidate track if its Cherenkov angle does not agree within 4σ standard deviations (σ) with either the pion or kaon hypothesis, or if it is identified as an electron by the DCH and the EMC. Particle identification (PID) information from the drift chamber and, when available, from the DIRC, must be consistent with the kaon hypothesis for the K meson candidate in $D^0 \rightarrow K^- \pi^+$, $D^0 \rightarrow K^- K^+$, and $\phi \rightarrow K^- K^+$ decays and with the pion hypothesis for the π^\pm meson candidates in $D^0 \rightarrow \pi^- \pi^+$ and $\omega \rightarrow \pi^+ \pi^- \pi^0$ decays.

Neutral pions are reconstructed by combining pairs of photon candidates with energy deposits larger than 70 MeV that are not matched to charged tracks. The $\gamma\gamma$ invariant mass is required to be in the range $115\text{--}150 \text{ MeV}/c^2$ and the total π^0 energy must be greater than 200 MeV . To improve momentum resolution, the invariant mass of the two photons from candidate π^0 's used in the B meson reconstruction is constrained to the nominal π^0 mass [11].

Neutral kaons are reconstructed from pairs of oppositely charged tracks with invariant mass within $7.8 \text{ MeV}/c^2$ ($\sim 3\sigma$) of the nominal K^0 mass. We also require that the ratio between the flight length in the plane transverse to the beam direction and its error be greater than 2. The ϕ mesons are reconstructed from two oppositely charged kaons with invariant mass in the range $1.008 < M(K^+ K^-) < 1.032 \text{ GeV}/c^2$. We also require $|\cos\theta_{\text{hel}}(\phi)| > 0.4$, where $\theta_{\text{hel}}(\phi)$ is the angle between the flight direction of one of the ϕ daughters and the D^0 flight direction, in the ϕ rest frame. The ω mesons are reconstructed from $\pi^+ \pi^- \pi^0$ combinations with invariant mass in the range $0.763 < M(\pi^+ \pi^- \pi^0) < 0.799 \text{ GeV}/c^2$. We define θ_N as the angle between the normal to the ω decay plane and the D^0 momentum in the ω rest frame, and $\theta_{\pi\pi}$ as the angle between the flight direction of one of the three pions in the ω rest frame and the flight direction of one of the other two pions in their center-of-mass (CM) frame. The quantities $\cos\theta_N$ and $\cos\theta_{\pi\pi}$ follow $\cos^2\theta_N$ and $\sin^2\theta_{\pi\pi}$ distributions for the signal and are almost flat for wrongly reconstructed or false ω candidates. We require the product $\cos^2\theta_N \sin^2\theta_{\pi\pi} > 0.08$. The invariant mass of a D^0 candidate, $M(D^0)$, must be within 2.5σ of the mean fitted mass, with resolution σ ranging from 4 to $20 \text{ MeV}/c^2$ depending on the D^0 decay mode. For $D^0 \rightarrow \pi^- \pi^+$, the invariant mass of the $(h^- \pi^+)$ system, where π^+ is the pion from D^0 , and h^- is the prompt track from B^- taken with the kaon mass hypothesis, must be greater than $1.9 \text{ GeV}/c^2$ to reject background from $B^- \rightarrow D^0 \pi^-$, $D^0 \rightarrow K^- \pi^+$ and $B^- \rightarrow K^{*0} \pi^-$, $K^{*0} \rightarrow K^- \pi^+$ decays. To improve the D^0 momentum resolution, for all the D^0 decay channels the candidate invariant mass is constrained to the nominal D^0 mass [11].

We reconstruct B meson candidates by combining a D^0 candidate with a track h . For the $D^0 \rightarrow K^- \pi^+$ mode, the charge of the track h must match that of the kaon from the D^0 meson decay. We select B meson candidates using the beam-energy-substituted mass $m_{\text{ES}} = \sqrt{(E_i^{*2}/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - p_B^2}$ and the energy difference $\Delta E = E_B^* - E_i^*/2$, where the subscripts i and B refer to the initial $e^+ e^-$ system and the B candidate, respectively, and the asterisk denotes the CM [$Y(4S)$] frame. The m_{ES} distributions for $B^- \rightarrow D^0 h^-$ signals are Gaussian functions centered at the B mass with a resolution of $2.6 \text{ MeV}/c^2$, which do not depend on the decay mode or on the nature of the prompt track. In contrast, the ΔE distributions depend on the mass assigned to the prompt track and on the D^0 momentum resolution. We evaluate ΔE with the kaon mass hypothesis so that the distributions are Gaussian and centered near zero for $B^- \rightarrow D^0 K^-$ events and shifted by approximately 50 MeV for $B^- \rightarrow D^0 \pi^-$ events. The $B^- \rightarrow D^0 K^-$ ΔE resolution is about 17 MeV for all the D^0 decay modes. All B candidates are selected with m_{ES} within 3σ of the mean value and with ΔE in the range $-0.16 < \Delta E < 0.23 \text{ GeV}$.

To reduce background from continuum production of light quarks, we construct a linear Fisher discriminant [12] based on the following quantities: (i) $L_0 = \sum_i p_i$ and $L_2 = \sum_i p_i \cos^2\theta_i$, evaluated in the CM frame, where p_i is the momentum, and θ_i is the angle with respect to the thrust axis of the B candidate of charged tracks and neutral clusters not used to reconstruct the B ; (ii) $|\cos\theta_T|$, where θ_T is the angle between the thrust axes of the B candidate and of the remaining tracks and clusters, evaluated in the CM frame; (iii) $|\cos\theta_B|$, where θ_B is the polar angle of the B candidate in the CM frame.

For events with multiple $B^- \rightarrow D^0 h^-$ candidates (1%–7% of the selected events, depending on the D^0 decay mode), we choose that with the smallest χ^2 formed from the differences of the measured and true masses of the candidate B , D^0 , π^0 (only for $D^0 \rightarrow K_S^0 \pi^0$, $K_S^0 \omega$), $\phi(D^0 \rightarrow K_S^0 \phi)$, $\omega(D^0 \rightarrow K_S^0 \omega)$, scaled by the mass spread. The total reconstruction efficiencies, based on simulated signal events, are 39% ($K^- \pi^+$), 31% ($K^- K^+$), 30% ($\pi^- \pi^+$), 17% ($K_S^0 \pi^0$), 20% ($K_S^0 \phi$), and 7% ($K_S^0 \omega$).

The main contributions to the background from $B\bar{B}$ events come from the processes $B \rightarrow D^* h$ ($h = \pi, K$), $B^- \rightarrow D^0 \rho^-$, misreconstructed $B^- \rightarrow D^0 h^-$, and from charmless B decays to the same final state as the signal: for instance, the process $B^- \rightarrow K^- K^+ K^-$ is a background for $B^- \rightarrow D^0 K^-$, $D^0 \rightarrow K^- K^+$. These charmless backgrounds have similar ΔE and m_{ES} distribution as the $D^0 K^-$ signal and we call them “peaking $B\bar{B}$ backgrounds.”

For each D^0 decay mode an extended unbinned maximum likelihood fit to the selected data events determines yields for two signal channels, $B^- \rightarrow D^0 \pi^-$ and $B^- \rightarrow D^0 K^-$, and four kinds of backgrounds: candidates selected

B. AUBERT *et al.*

either from continuum or from $B\bar{B}$ events, in which the prompt track is either a pion or a kaon.

The fit uses as input ΔE and a particle identification probability for the prompt track based on the Cherenkov angle θ_C , the momentum p , and the polar angle θ of the track.

The extended likelihood function \mathcal{L} is defined as

$$\mathcal{L} = \exp\left(-\sum_{i=1}^6 n_i\right) \prod_{j=1}^N \left[\sum_{i=1}^6 n_i \mathcal{P}_i(\vec{x}_j; \vec{\alpha}_i) \right], \quad (6)$$

where N is the total number of observed events and n_i is the yield of the i th event category. The six functions $\mathcal{P}_i(\vec{x}_j; \vec{\alpha}_i)$ are the probability density functions (PDFs) for the variables \vec{x}_j , given the set of parameters $\vec{\alpha}_i$. They are evaluated as a product $\mathcal{P}_i = \mathcal{P}_{1i}(\Delta E) \times \mathcal{P}_{2i}(\theta_C)$.




FIG. 1. Distributions of ΔE for events enhanced in the $B \rightarrow D^0 K$ signal. Top: $B^- \rightarrow D^0 K^-$, $D^0 \rightarrow K^- \pi^+$; middle: $B^- \rightarrow D_{CP+}^0 K^-$; bottom: $B^- \rightarrow D_{CP-}^0 K^-$. Solid curves represent projections of the maximum likelihood fit; dashed, dash-dotted and dotted curves represent the $B \rightarrow D^0 K$, $B \rightarrow D^0 \pi$ and background contributions.

TABLE I. Yields from the maximum likelihood fit. The quoted uncertainties are statistical.

D^0 mode	$N(D\pi^+)$	$N(D\pi^-)$	$N(DK^+)$	$N(DK^-)$
$K^- \pi^+$	8151 ± 95	7899 ± 93	649 ± 29	611 ± 28
$K^- K^+$	705 ± 28	690 ± 28	26 ± 9	70 ± 10
$\pi^- \pi^+$	256 ± 18	219 ± 17	18 ± 7	17 ± 7
$K_S^0 \pi^0$	707 ± 29	677 ± 29	39 ± 9	42 ± 9
$K_S^0 \phi$	176 ± 14	157 ± 13	15 ± 5	13 ± 4
$K_S^0 \omega$	235 ± 17	230 ± 17	25 ± 7	14 ± 6

The ΔE distribution for $B^- \rightarrow D^0 K^-$ signal events is parametrized with a Gaussian function. The ΔE distribution for $B^- \rightarrow D^0 \pi^-$ is parametrized with the same Gaussian function used for $B^- \rightarrow D^0 K^-$ with an additional shift, computed event by event as a function of the prompt track momentum, arising from the wrong mass assignment to the prompt track. The offset and width of the Gaussian functions are determined from data together with the yields.

The ΔE distribution for the continuum background is parametrized with a linear function whose slope is determined from off-resonance data. The ΔE distribution for the nonpeaking $B\bar{B}$ background is empirically parametrized with the sum of a Gaussian function and an exponential function when the prompt track is a pion, and with an exponential function when the prompt track is a kaon. The parameters are determined from simulated events. The ΔE distribution for the peaking charmless $B\bar{B}$ background is parametrized with the same Gaussian function used for the $B^- \rightarrow D^0 K^-$ signal. The yield of the $B\bar{B}$ peaking background is estimated from the sidebands of the D^0 invariant mass distribution and fixed in the fit.

The parametrization of the particle identification PDF is performed by fitting with two Gaussian functions the background-subtracted distribution of the difference between the reconstructed and expected Cherenkov angles of kaon and pion samples. The parametrization is performed as a function of the momentum and polar angle of the track. Pions and kaons are selected from a pure $D^{*+} \rightarrow D^0 \pi^+$, $D^0 \rightarrow K^- \pi^+$ control sample.

The results of the fit are summarized in Table I. Figure 1 shows the distributions of ΔE for the $K^- \pi^+$, $CP+$ and $CP-$ modes after enhancing the $B \rightarrow D^0 K$ purity by requiring that the prompt track be consistent with the kaon hypothesis. The total PDF, normalized by the fitted signal

TABLE II. Measured ratios $R_{CP\pm}$ and $A_{CP\pm}$ for CP -even and CP -odd D decay modes. The first error is statistical, the second is systematic. R_{CP-} and A_{CP-} are corrected for the CP -even dilution described in the text.

D^0 mode	R_{CP}	A_{CP}
$CP+$	$0.90 \pm 0.12 \pm 0.04$	$0.35 \pm 0.13 \pm 0.04$
$CP-$	$0.86 \pm 0.10 \pm 0.05$	$-0.06 \pm 0.13 \pm 0.04$

MEASUREMENT OF CP OBSERVABLES FOR THE DECAYS $B^\pm \rightarrow D_{CP}^0 K^\pm$

PHYSICAL REVIEW D 73, 051105 (2006)

and background yields, integrated over the Cherenkov angle variable and modified to take into account the tighter selection criteria, is overlaid in the figure.

The ratios $R_{CP\pm}$ are computed for the five CP modes using the relations in Eq. (3). A number of systematic uncertainties, as the uncertainty associated to the tracking efficiency and the uncertainty on the D^0 decay branching fractions, cancel out in the measurement of the double ratio. The relations $R_{CP\pm} = R_\pm/R$ hold neglecting the magnitude r_π of the ratio of the amplitudes of the $B^- \rightarrow \bar{D}^0 \pi^-$ and $B^- \rightarrow D^0 \pi^-$ processes [6] ($r_\pi \sim r \frac{\lambda^2}{1-\lambda^2} \lesssim 0.012$, where $\lambda \approx 0.22$ [11] is the sine of the Cabibbo angle). This assumption is considered further when we discuss the systematic uncertainties. The quantities R_\pm/R are computed from the ratios of the $B \rightarrow DK$ and $B \rightarrow D\pi$ yields in Table I, scaled by correction factors taking into account small differences in the selection efficiency between $B \rightarrow DK$ and $B \rightarrow D\pi$. These correction factors are evaluated from simulated events and range between 0.982 ± 0.018 and 1.020 ± 0.031 depending on the D^0 decay mode. The results for the CP -even and CP -odd combinations are listed in Table II.

The partial-rate charge asymmetries $A_{CP\pm}$ are calculated from the measured yields of positive and negative $B \rightarrow DK$ decays in Table I. The results for the CP -even and CP -odd combinations are reported in Table II.

In the case of $D^0 \rightarrow K_S^0 \phi$, $\phi \rightarrow K^+ K^-$, and $D^0 \rightarrow K_S^0 \omega$, $\omega \rightarrow \pi^+ \pi^- \pi^0$, the values of R_{CP-} and A_{CP-} quoted in Table II are obtained after correcting the measured values to take into account the dilution from a CP -even background arising from $B^- \rightarrow D^0 h^-$, $D^0 \rightarrow K_S^0(K^- K^+)_{\text{non-}\phi}$ and $D^0 \rightarrow K_S^0(\pi^- \pi^+ \pi^0)_{\text{non-}\omega}$ decays. For the $K_S^0 \phi$ channel we exploit the investigation performed by *BABAR* of the $D^0 \rightarrow K_S^0 K^+ K^-$ Dalitz plot [13] to estimate the level of the CP -even background (0.160 ± 0.006 relative to the $K_S^0 \phi$ signal) and the corresponding R_{CP-} and A_{CP-} dilution. For the $K_S^0 \omega$ channel there is little information on this background. We estimate the amount of $D^0 \rightarrow K_S^0(\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$ background (0.25 ± 0.05 relative to the $K_S^0 \omega$ signal) from the $\cos\theta_N$ distribution of $B^- \rightarrow$

$D^0 \pi^-$, $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ candidates, and assume the CP -even content of this background to be $(50 \pm 29)\%$.

Systematic uncertainties in the ratios $R_{CP\pm}$ and in the CP asymmetries $A_{CP\pm}$ are listed in Table III. They arise both from the uncertainties on the signal yields, extracted through the unbinned maximum likelihood fit, and from the assumptions used to compute $R_{CP\pm}$ and $A_{CP\pm}$. The correlations between the different sources of systematic errors, when non-negligible, are considered when combining the two CP -even or the three CP -odd modes.

The uncertainties on the fitted signal yields are due to the imperfect knowledge of the ΔE and PID PDFs and of the peaking background yields, and are evaluated by varying the parameters of the PDFs and the peaking background yields by $\pm 1\sigma$ and taking the difference in the signal yields. The uncertainties in the branching fractions used in the simulation of the B decays that contribute to the $B\bar{B}$ background are also taken into account. The yields of the $B\bar{B}$ and continuum backgrounds found in data are consistent with what is expected from the simulation. In the $K_S^0 \phi$ and $K_S^0 \omega$ channels we also take into account the uncertainties in the dilution factors due to the imperfect knowledge of the levels of the CP -even backgrounds from $B^- \rightarrow D^0 K^-$, $D^0 \rightarrow K_S^0(K^- K^+)_{\text{non-}\phi}$ and $D^0 \rightarrow K_S^0(\pi^- \pi^+ \pi^0)_{\text{non-}\omega}$ decays.

A possible bias in the measured $A_{CP\pm}$ may come from an intrinsic detector charge asymmetry due to asymmetries in acceptance or tracking and particle identification efficiencies. An upper limit on this bias has been obtained from the measured asymmetries in the processes $B^- \rightarrow D^0 h^-$, $D^0 \rightarrow K^- \pi^+$ and $B^- \rightarrow D_{CP\pm}^0 \pi^-$, where CP violation is expected to be negligible. From the average asymmetry, $(-1.8 \pm 0.9)\%$, we obtain the limit $\pm 2.7\%$ for the bias. This has been added in quadrature to the total systematic uncertainty on the CP asymmetry.

For the branching fraction ratios $R_{CP\pm}$ two additional sources of uncertainty are the correction factors used to scale the yield ratios, and the assumption that $R_{CP\pm} = R_\pm/R$. The scaling factor, estimated from simulated events, is a double ratio of efficiencies, $\varepsilon_\pm^{K/\pi}/\varepsilon^{K/\pi}$, where

TABLE III. Systematic uncertainties on the observables $R_{CP\pm}$ and $A_{CP\pm}$ after combination of the two CP -even and the three CP -odd D^0 decay modes.

Source	ΔR_{CP+} (%)	ΔR_{CP-} (%)	ΔA_{CP+} (%)	ΔA_{CP-} (%)
Background ΔE PDF	1.3	1.1	1.1	0.4
PID PDF	0.1	0.1	0.2	0.2
Peaking background yields	3.0	4.2	2.6	2.2
Opposite- CP background	...	1.3	...	1.0
Detector charge asymmetry	2.7	2.7
$\varepsilon_\pm^{K/\pi}/\varepsilon^{K/\pi}$	1.0	1.1
r_π	2.2	2.1
Total	4.1	5.1	3.9	3.7

B. AUBERT *et al.*

$\varepsilon_{(\pm)}^{K/\pi}$ denotes the ratio between the selection efficiencies of $B \rightarrow D_{(CP\pm)}^0 K$ and $B \rightarrow D_{(CP\pm)}^0 \pi$. In the double ratio the systematic uncertainties arising from possible discrepancies between data and simulation are negligible, and only the contribution from the limited statistics of the simulated samples remains. The assumption $R_{CP\pm} = R_{\pm}/R$ introduces a relative uncertainty $\pm 2r_{\pi} \cos\delta_{\pi} \cos\gamma$ on $R_{CP\pm}$, where δ_{π} is the relative strong phase between the amplitudes $A(B^- \rightarrow \bar{D}^0 \pi^-)$ and $A(B^- \rightarrow D^0 \pi^-)$. Since $|\cos\delta_{\pi} \cos\gamma| \leq 1$ and $r_{\pi} \lesssim 0.012$, we assign a relative uncertainty $\pm 2.4\%$ to $R_{CP\pm}$, which is completely anticorrelated between R_{CP+} and R_{CP-} .

We quote the measurements in terms of x_{\pm} and r^2 ,

$$x_+ = -0.082 \pm 0.053(\text{stat}) \pm 0.018(\text{syst}), \quad (7)$$

$$x_- = +0.102 \pm 0.062(\text{stat}) \pm 0.022(\text{syst}), \quad (8)$$

$$r^2 = -0.12 \pm 0.08(\text{stat}) \pm 0.03(\text{syst}). \quad (9)$$

The measured values of x_{\pm} are consistent with those found, on a slightly smaller data sample, with the $B^- \rightarrow DK^-$, $D \rightarrow K_S^0 \pi^- \pi^+$ decays, and the precision is comparable [8].

PHYSICAL REVIEW D 73, 051105 (2006)

In conclusion, we have reconstructed $B^- \rightarrow D^0 K^-$ decays with D^0 mesons decaying to non- CP , CP -even and CP -odd eigenstates. We have improved the measurements of $R_{CP\pm}$ and $A_{CP\pm}$ [14,15], and we have also expressed the results in terms of the same x_{\pm} parameters as were measured with $B^- \rightarrow DK^-$, $D \rightarrow K_S^0 \pi^- \pi^+$ through a Dalitz plot analysis of the D final state [8], with a comparable precision. These measurements, combined with the existing measurements of the $B \rightarrow DK$ decays, will improve the knowledge of the angle γ and the parameter r .

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (the Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A.P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

-
- [1] Reference to the charge-conjugate state is implied here and throughout the text unless otherwise stated.
 - [2] M. Gronau and D. Wyler, Phys. Lett. B **265**, 172 (1991); M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991).
 - [3] D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997); A. Giri, Y. Grossman, A. Soffer, and J. Zupan, Phys. Rev. D **68**, 054018 (2003).
 - [4] M. Gronau, Phys. Rev. D **58**, 037301 (1998).
 - [5] Y. Grossman, A. Soffer, and J. Zupan, Phys. Rev. D **72**, 031501(R) (2005).
 - [6] M. Gronau, Phys. Lett. B **557**, 198 (2003).
 - [7] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **72**, 032004 (2005); M. Saigo *et al.* (*Belle* Collaboration), Phys. Rev. Lett. **94**, 091601 (2005), updated in hep-ex/0508048.
 - [8] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **95**, 121802 (2005).
 - [9] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
 - [10] S. Agostinelli *et al.* (*GEANT4* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
 - [11] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
 - [12] R. A. Fisher, Ann. Eugen. **7**, 179 (1936).
 - [13] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **72**, 052008 (2005).
 - [14] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 202002 (2004). Note that R_{CP} in this reference is defined as R_+ in this paper.
 - [15] K. Abe *et al.* (*Belle* Collaboration), hep-ex/0601032.