

*Physics*

*Physics Research Publications*

---

*Purdue University*

*Year 2006*

---

Measurement of the  
B-s(0)-(B)over-bar(s)(0) oscillation  
frequency

A. Abulencia, D. Acosta, J. Adelman, T. Affolder, T. Akimoto, M. G. Albrow, D. Ambrose, S. Amerio, D. Amidei, A. Anastassov, K. Anikeev, A. Annovi, J. Antos, M. Aoki, G. Apollinari, J. F. Arguin, T. Arisawa, A. Artikov, W. Ashmanskas, A. Attal, F. Azfar, P. Azzi-Bacchetta, P. Azzurri, N. Bacchetta, H. Bachacou, W. Badgett, A. Barbaro-Galtieri, V. E. Barnes, B. A. Barnett, S. Baroiant, V. Bartsch, G. Bauer, F. Bedeschi, S. Behari, S. Belforte, G. Bellettini, J. Bellinger, A. Belloni, E. Ben Haim, D. Benjamin, A. Beretvas, J. Beringer, T. Berry, A. Bhatti, M. Binkley, D. Bisello, R. E. Blair, C. Blocker, B. Blumenfeld, A. Bocci, A. Bodek, V. Boisvert, G. Bolla, A. Bolshov, D. Bortolotto, J. Boudreau, A. Boveia, B. Brau, C. Bromberg, E. Brubaker, J. Budagov, H. S. Budd, S. Budd, K. Burkett, G. Busetto, P. Bussey, K. L. Byrum, S. Cabrera, M. Campanelli, M. Campbell, F. Canelli, A. Canepa, D. Carlsmith, R. Carosi, S. Carron, B. Casal, M. Casarsa, A. Castro, P. Catastini, D. Cauz, M. Cavalli-Sforza, A. Cerri, L. Cerrito, S. H. Chang, J. Chapman, Y. C. Chen, M. Chertok, G. Chiarelli, G. Chlachidze, F. Chlebana, I. Cho, K. Cho, D. Chokheli, J. P. Chou, P. H. Chu, S. H. Chuang, K. Chung, W. H. Chung, Y. S. Chung, M. Ciljak, C. I. Ciobanu, M. A. Ciocci, A. Clark, D. Clark, M. Coca, G. Compostella, M. E. Convery, J. Conway, B. Cooper, K. Copic, M. Cordelli, G. Cortiana, F. Crescioli, A. Cruz, C. C. Almenar, J. Cuevas, R. Culbertson, D. Cyr, S. DaRonco, S. D'Auria, M. D'Onofrio, D. Dagenhart, P. De Barbaro, S. De Cecco, A. Deisher, G. De Lentdecker, M. Dell'Orso, F. D. Paoli, S. Demers, L. Demortier, J. Deng, M. Deninno, D. De Pedis, P. F. Derwent, G. P. Di Giovanni, B. Di Ruzza, C. Dionisi, J. R. Dittmann, P. DiTuro, C. Dorr, S. Donati, M. Donega, P. Dong, J. Donini, T. Dorigo, S. Dube, K. Ebina, J. Efron, J. Ehlers, R. Erbacher, D. Errede, S. Errede, R. Eusebi, H. C. Fang, S. Farrington, I. Fedorko, W. T. Fedorko, R. G. Feild, M. Feindt, J. P. Fernandez, R. Field, G. Flanagan, L. R. Flores-Castillo, A. Foland, S. Forrester, G. W. Foster, M. Franklin, J. C. Freeman, H. J. Frisch, I. Furic, M. Gallinaro, J. Galyardt, J. E. Garcia, M. G. Sciveres, A. F. Garfinkel, C. Gay, H. Gerberich, D. Gerdes, S.

Giagu, P. Giannetti, A. Gibson, K. Gibson, C. Ginsburg, N. Giokaris, K. Giolo, M. Giordani, P. Giromini, M. Giunta, G. Giurgiu, V. Glagolev, D. Glenzinski, M. Gold, N. Goldschmidt, J. Goldstein, G. Gomez, G. Gomez-Ceballos, M. Goncharov, O. Gonzalez, I. Gorelov, A. T. Goshaw, Y. Gotra, K. Goulianos, A. Gresele, M. Griffiths, S. Grinstein, C. Grosso-Pilcher, R. C. Group, U. Grundler, J. G. Da Costa, Z. Gunay-Unalan, C. Haber, S. R. Hahn, K. Hahn, E. Halkiadakis, A. Hamilton, B. Y. Han, J. Y. Han, R. Handler, F. Happacher, K. Hara, M. Hare, S. Harper, R. F. Harr, R. M. Harris, K. Hatakeyama, J. Hauser, C. Hays, A. Heijboer, B. Heinemann, J. Heinrich, M. Herndon, D. Hidas, C. S. Hill, D. Hirschbuehl, A. Hocker, A. Holloway, S. Hou, M. Houlden, S. C. Hsu, B. T. Huffman, R. E. Hughes, J. Huston, J. Incandela, G. Introzzi, M. Iori, Y. Ishizawa, A. Ivanov, B. Iyutin, E. James, D. Jang, B. Jayatilaka, D. Jeans, H. Jensen, E. J. Jeon, S. Jindariani, M. Jones, K. K. Joo, S. Y. Jun, T. R. Junk, T. Kamon, J. Kang, P. E. Karchin, Y. Kato, Y. Kemp, R. Kephart, U. Kerzel, V. Khotilovich, B. Kilminster, D. H. Kim, H. S. Kim, J. E. Kim, M. J. Kim, S. B. Kim, S. H. Kim, Y. K. Kim, L. Kirsch, S. Klimentenko, M. Klute, B. Knuteson, B. R. Ko, H. Kobayashi, K. Kondo, D. J. Kong, J. Konigsberg, A. Korytov, A. V. Kotwal, A. Kovalev, A. Kraan, J. Kraus, I. Kravchenko, M. Kreps, J. Kroll, N. Krumnack, M. Kruse, V. Krutelyov, S. E. Kuhlmann, Y. Kusakabe, S. Kwang, A. T. Laasanen, S. Lai, S. Lami, S. Lammel, M. Lancaster, R. L. Lander, K. Lannon, A. Lath, G. Latino, I. Lazzizzera, T. LeCompte, J. Lee, J. Lee, Y. J. Lee, S. W. Lee, R. Lefevre, N. Leonardo, S. Leone, S. Levy, J. D. Lewis, C. Lin, C. S. Lin, M. Lindgren, E. Lipeles, T. M. Liss, A. Lister, D. O. Litvintsev, T. Liu, N. S. Lockyer, A. Loginov, M. Loreti, P. Loverre, R. S. Lu, D. Lucchesi, P. Lujan, P. Lukens, G. Lungu, L. Lyons, J. Lys, R. Lysak, E. Lytken, P. Mack, D. MacQueen, R. Madrak, K. Maeshima, T. Maki, P. Maksimovic, S. Malde, G. Manca, F. Margaroli, R. Marginean, C. Marino, A. Martin, V. Martin, M. Martinez, T. Maruyama, P. Mastrandrea, H. Matsunaga, M. E. Mattson, R. Mazini, P. Mazzanti, K. S. McFarland, P. McIntyre, R. McNulty, A. Mehta, S. Menzemer, A. Menzione, P. Merkel, C. Mesropian, A. Messina, M. von der Mey, T. Miao, N. Miladinovic, J. Miles, R. Miller, J. S. Miller, C. Mills, M. Milnik, R. Miquel, A. Mitra, G. Mitselmakher, A. Miyamoto, N. Moggi, B. Mohr, R. Moore, M. Morello, P. M. Fernandez, J. Mulmenstadt, A. Mukherjee, T. Muller, R. Mumford, P. Murat, J. Nachtman, J. Naganoma, S. Nahn, I. Nakano, A. Napier, D. Naumov, V. Necula, C. Neu, M. S. Neubauer, J. Nielsen, T. Nigmanov, L. Nodulman, O. Norriella, E. Nurse, T. Ogawa, S. H. Oh, Y. D. Oh, T. Okusawa, R. Oldeman, R. Orava, K. Osterberg, C. Pagliarone, E. Palencia, R. Paoletti, V. Papadimitriou, A. A. Paramonov, B. Parks, S. Pashapour, J. Patrick, G. Pauletta, M. Paulini, C. Paus, D. E. Pellett, A. Penzo, T. J. Phillips, G. Piacentino, J. Piedra, L. Pinera, K. Pitts, C. Plager, L. Pondrom, X. Portell, O. Poukhov, N. Pounder, F. Prakoshyn, A. Pronko, J. Proudfoot, F. Ptohos, G. Punzi, J. Pursley, J. Rademacker, A. Rahaman, A. Rakitin, S. Rappoccio, F. Ratnikov, B. Reisert, V. Rekovic, N. van Remortel, P. Renton, M. Rescigno, S. Richter, F. Rimondi, L. Ristori, W. J. Robertson, A. Robson, T. Rodrigo, E. Rogers, S. Rolli, R. Roser, M. Rossi, R. Rossin, C. Rott, A. Ruiz, J. Russ, V. Rusu, H. Saarikko, S. Sabik, A. Safonov, W. K. Sakumoto, G.

Salamanna, O. Salto, D. Saltzberg, C. Sanchez, L. Santi, S. Sarkar, L. Sartori, K. Sato, P. Savard, A. Savoy-Navarro, T. Scheidle, P. Schlabach, E. E. Schmidt, M. P. Schmidt, M. Schmitt, T. Schwarz, L. Scodellaro, A. L. Scott, A. Scribano, F. Scuri, A. Sedov, S. Seidel, Y. Seiya, A. Semenov, L. Sexton-Kennedy, I. Sfiligoi, M. D. Shapiro, T. Shears, P. F. Shepard, D. Sherman, M. Shimojima, M. Shochet, Y. Shon, I. Shreyber, A. Sidoti, P. Sinervo, A. Sisakyan, J. Sjolín, A. Skiba, A. J. Slaughter, K. Sliwa, J. R. Smith, F. D. Snider, R. Snihur, M. Soderberg, A. Soha, S. Somalwar, V. Sorin, J. Spalding, M. Spezziga, F. Spinella, T. Spreitzer, P. Squillacioti, M. Stanitzki, A. Staveris-Polykalas, R. S. Denis, B. Stelzer, O. Stelzer-Chilton, D. Stentz, J. Strologas, D. Stuart, J. S. Suh, A. Sukhanov, K. Sumorok, H. Sun, T. Suzuki, A. Taffard, R. Takashima, Y. Takeuchi, K. Takikawa, M. Tanaka, R. Tanaka, N. Tanimoto, M. Tecchio, P. K. Teng, K. Terashi, S. Tether, J. Thom, A. S. Thompson, E. Thomson, P. Tipton, V. Tiwari, S. Tkaczyk, D. Toback, S. Tokar, K. Tollefson, T. Tomura, D. Tonelli, M. Tonnesmann, S. Torre, D. Torretta, S. Tourneur, W. Trischuk, R. Tsuchiya, S. Tsuno, N. Turini, F. Ukegawa, T. Unverhau, S. Uozumi, D. Usynin, A. Vaiciulis, S. Vallecorsa, A. Varganov, E. Vataga, G. Velev, G. Veramendi, V. Veszpremi, R. Vidal, I. Vila, R. Vilar, T. Vine, I. Vollrath, I. Volobouev, G. Volpi, F. W. Rthwein, P. Wagner, R. G. Wagner, R. L. Wagner, W. Wagner, R. Wallny, T. Walter, Z. Wan, S. M. Wang, A. Warburton, S. Waschke, D. Waters, W. C. Wester, B. Whitehouse, D. Whiteson, A. B. Wicklund, E. Wicklund, G. Williams, H. H. Williams, P. Wilson, B. L. Winer, P. Wittich, S. Wolbers, C. Wolfe, T. Wright, X. Wu, S. M. Wynne, A. Yagil, K. Yamamoto, J. Yamaoka, T. Yamashita, C. Yang, U. K. Yang, Y. C. Yang, W. M. Yao, G. P. Yeh, J. Yoh, K. Yorita, T. Yoshida, G. B. Yu, I. Yu, S. S. Yu, J. C. Yun, L. Zanello, A. Zanetti, I. Zaw, F. Zetti, X. Zhang, J. Zhou, and S. Zucchelli

This paper is posted at Purdue e-Pubs.

[http://docs.lib.purdue.edu/physics\\_articles/15](http://docs.lib.purdue.edu/physics_articles/15)

## Measurement of the $B_s^0$ - $\bar{B}_s^0$ Oscillation Frequency

A. Abulencia,<sup>23</sup> D. Acosta,<sup>17</sup> J. Adelman,<sup>13</sup> T. Affolder,<sup>10</sup> T. Akimoto,<sup>55</sup> M. G. Albrow,<sup>16</sup> D. Ambrose,<sup>16</sup> S. Amerio,<sup>43</sup> D. Amidei,<sup>34</sup> A. Anastassov,<sup>52</sup> K. Anikeev,<sup>16</sup> A. Annovi,<sup>18</sup> J. Antos,<sup>1</sup> M. Aoki,<sup>55</sup> G. Apollinari,<sup>16</sup> J.-F. Arguin,<sup>33</sup> T. Arisawa,<sup>57</sup> A. Artikov,<sup>14</sup> W. Ashmanskas,<sup>16</sup> A. Attal,<sup>8</sup> F. Azfar,<sup>42</sup> P. Azzi-Bacchetta,<sup>43</sup> P. Azzurri,<sup>46</sup> N. Bacchetta,<sup>43</sup> H. Bachacou,<sup>28</sup> W. Badgett,<sup>16</sup> A. Barbaro-Galtieri,<sup>28</sup> V. E. Barnes,<sup>48</sup> B. A. Barnett,<sup>24</sup> S. Baroiant,<sup>7</sup> V. Bartsch,<sup>30</sup> G. Bauer,<sup>32</sup> F. Bedeschi,<sup>46</sup> S. Behari,<sup>24</sup> S. Belforte,<sup>54</sup> G. Bellettini,<sup>46</sup> J. Bellinger,<sup>59</sup> A. Belloni,<sup>32</sup> E. Ben Haim,<sup>44</sup> D. Benjamin,<sup>15</sup> A. Beretvas,<sup>16</sup> J. Beringer,<sup>28</sup> T. Berry,<sup>29</sup> A. Bhatti,<sup>50</sup> M. Binkley,<sup>16</sup> D. Bisello,<sup>43</sup> R. E. Blair,<sup>2</sup> C. Blocker,<sup>6</sup> B. Blumenfeld,<sup>24</sup> A. Bocci,<sup>15</sup> A. Bodek,<sup>49</sup> V. Boisvert,<sup>49</sup> G. Bolla,<sup>48</sup> A. Bolshov,<sup>32</sup> D. Bortoletto,<sup>48</sup> J. Boudreau,<sup>47</sup> A. Boveia,<sup>10</sup> B. Brau,<sup>10</sup> C. Bromberg,<sup>35</sup> E. Brubaker,<sup>13</sup> J. Budagov,<sup>14</sup> H. S. Budd,<sup>49</sup> S. Budd,<sup>23</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>43</sup> P. Bussey,<sup>20</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>15</sup> M. Campanelli,<sup>19</sup> M. Campbell,<sup>34</sup> F. Canelli,<sup>8</sup> A. Canepa,<sup>48</sup> D. Carlsmith,<sup>59</sup> R. Carosi,<sup>46</sup> S. Carron,<sup>15</sup> B. Casal,<sup>11</sup> M. Casarsa,<sup>54</sup> A. Castro,<sup>5</sup> P. Catastini,<sup>46</sup> D. Cauz,<sup>54</sup> M. Cavalli-Sforza,<sup>3</sup> A. Cerri,<sup>28</sup> L. Cerrito,<sup>42</sup> S. H. Chang,<sup>27</sup> J. Chapman,<sup>34</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>46</sup> G. Chlachidze,<sup>14</sup> F. Chlebana,<sup>16</sup> I. Cho,<sup>27</sup> K. Cho,<sup>27</sup> D. Chokheli,<sup>14</sup> J. P. Chou,<sup>21</sup> P. H. Chu,<sup>23</sup> S. H. Chuang,<sup>59</sup> K. Chung,<sup>12</sup> W. H. Chung,<sup>59</sup> Y. S. Chung,<sup>49</sup> M. Ciljak,<sup>46</sup> C. I. Ciobanu,<sup>23</sup> M. A. Ciocci,<sup>46</sup> A. Clark,<sup>19</sup> D. Clark,<sup>6</sup> M. Coca,<sup>15</sup> G. Compostella,<sup>43</sup> M. E. Convery,<sup>50</sup> J. Conway,<sup>7</sup> B. Cooper,<sup>30</sup> K. Copic,<sup>34</sup> M. Cordelli,<sup>18</sup> G. Cortiana,<sup>43</sup> F. Crescioli,<sup>46</sup> A. Cruz,<sup>17</sup> C. Cuenca Almenar,<sup>7</sup> J. Cuevas,<sup>11</sup> R. Culbertson,<sup>16</sup> D. Cyr,<sup>59</sup> S. DaRonco,<sup>43</sup> S. D'Auria,<sup>20</sup> M. D'Onofrio,<sup>3</sup> D. Dagenhart,<sup>6</sup> P. de Barbaro,<sup>49</sup> S. De Cecco,<sup>51</sup> A. Deisher,<sup>28</sup> G. De Lentdecker,<sup>49</sup> M. Dell'Orso,<sup>46</sup> F. Delli Paoli,<sup>43</sup> S. Demers,<sup>49</sup> L. Demortier,<sup>50</sup> J. Deng,<sup>15</sup> M. Deninno,<sup>5</sup> D. De Pedis,<sup>51</sup> P. F. Derwent,<sup>16</sup> G. P. Di Giovanni,<sup>44</sup> B. Di Ruzza,<sup>54</sup> C. Dionisi,<sup>51</sup> J. R. Dittmann,<sup>4</sup> P. DiTuro,<sup>52</sup> C. Dörr,<sup>25</sup> S. Donati,<sup>46</sup> M. Donega,<sup>19</sup> P. Dong,<sup>8</sup> J. Donini,<sup>43</sup> T. Dorigo,<sup>43</sup> S. Dube,<sup>52</sup> K. Ebina,<sup>57</sup> J. Efron,<sup>39</sup> J. Ehlers,<sup>19</sup> R. Erbacher,<sup>7</sup> D. Errede,<sup>23</sup> S. Errede,<sup>23</sup> R. Eusebi,<sup>16</sup> H. C. Fang,<sup>28</sup> S. Farrington,<sup>29</sup> I. Fedorko,<sup>46</sup> W. T. Fedorko,<sup>13</sup> R. G. Feild,<sup>60</sup> M. Feindt,<sup>25</sup> J. P. Fernandez,<sup>31</sup> R. Field,<sup>17</sup> G. Flanagan,<sup>48</sup> L. R. Flores-Castillo,<sup>47</sup> A. Foland,<sup>21</sup> S. Forrester,<sup>7</sup> G. W. Foster,<sup>28</sup> M. Franklin,<sup>21</sup> J. C. Freeman,<sup>28</sup> H. J. Frisch,<sup>13</sup> I. Furic,<sup>13</sup> M. Gallinaro,<sup>50</sup> J. Galyardt,<sup>12</sup> J. E. Garcia,<sup>46</sup> M. Garcia Sciveres,<sup>28</sup> A. F. Garfinkel,<sup>48</sup> C. Gay,<sup>60</sup> H. Gerberich,<sup>23</sup> D. Gerdes,<sup>34</sup> S. Giagu,<sup>51</sup> P. Giannetti,<sup>46</sup> A. Gibson,<sup>28</sup> K. Gibson,<sup>12</sup> C. Ginsburg,<sup>16</sup> N. Giokaris,<sup>14</sup> K. Giolo,<sup>48</sup> M. Giordani,<sup>54</sup> P. Giromini,<sup>18</sup> M. Giunta,<sup>46</sup> G. Giurgiu,<sup>12</sup> V. Glagolev,<sup>14</sup> D. Glenzinski,<sup>16</sup> M. Gold,<sup>37</sup> N. Goldschmidt,<sup>34</sup> J. Goldstein,<sup>42</sup> G. Gomez,<sup>11</sup> G. Gomez-Ceballos,<sup>11</sup> M. Goncharov,<sup>53</sup> O. González,<sup>31</sup> I. Gorelov,<sup>37</sup> A. T. Goshaw,<sup>15</sup> Y. Gotra,<sup>47</sup> K. Goulianos,<sup>50</sup> A. Gresele,<sup>43</sup> M. Griffiths,<sup>29</sup> S. Grinstein,<sup>21</sup> C. Grosso-Pilcher,<sup>13</sup> R. C. Group,<sup>17</sup> U. Grundler,<sup>23</sup> J. Guimaraes da Costa,<sup>21</sup> Z. Gunay-Unalan,<sup>35</sup> C. Haber,<sup>28</sup> S. R. Hahn,<sup>16</sup> K. Hahn,<sup>45</sup> E. Halkiadakis,<sup>52</sup> A. Hamilton,<sup>33</sup> B.-Y. Han,<sup>49</sup> J. Y. Han,<sup>49</sup> R. Handler,<sup>59</sup> F. Happacher,<sup>18</sup> K. Hara,<sup>55</sup> M. Hare,<sup>56</sup> S. Harper,<sup>42</sup> R. F. Harr,<sup>58</sup> R. M. Harris,<sup>16</sup> K. Hatakeyama,<sup>50</sup> J. Hauser,<sup>8</sup> C. Hays,<sup>15</sup> A. Heijboer,<sup>45</sup> B. Heinemann,<sup>29</sup> J. Heinrich,<sup>45</sup> M. Herndon,<sup>59</sup> D. Hidas,<sup>15</sup> C. S. Hill,<sup>10</sup> D. Hirschbuehl,<sup>25</sup> A. Hocker,<sup>16</sup> A. Holloway,<sup>21</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>29</sup> S.-C. Hsu,<sup>9</sup> B. T. Huffman,<sup>42</sup> R. E. Hughes,<sup>39</sup> J. Huston,<sup>35</sup> J. Incandela,<sup>10</sup> G. Introzzi,<sup>46</sup> M. Iori,<sup>51</sup> Y. Ishizawa,<sup>55</sup> A. Ivanov,<sup>7</sup> B. Iyutin,<sup>32</sup> E. James,<sup>16</sup> D. Jang,<sup>52</sup> B. Jayatilaka,<sup>34</sup> D. Jeans,<sup>51</sup> H. Jensen,<sup>16</sup> E. J. Jeon,<sup>27</sup> S. Jindariani,<sup>17</sup> M. Jones,<sup>48</sup> K. K. Joo,<sup>27</sup> S. Y. Jun,<sup>12</sup> T. R. Junk,<sup>23</sup> T. Kamon,<sup>53</sup> J. Kang,<sup>34</sup> P. E. Karchin,<sup>58</sup> Y. Kato,<sup>41</sup> Y. Kemp,<sup>25</sup> R. Kephart,<sup>16</sup> U. Kerzel,<sup>25</sup> V. Khotilovich,<sup>53</sup> B. Kilminster,<sup>39</sup> D. H. Kim,<sup>27</sup> H. S. Kim,<sup>27</sup> J. E. Kim,<sup>15</sup> M. J. Kim,<sup>12</sup> S. B. Kim,<sup>27</sup> S. H. Kim,<sup>55</sup> Y. K. Kim,<sup>13</sup> L. Kirsch,<sup>6</sup> S. Klimentenko,<sup>17</sup> M. Klute,<sup>32</sup> B. Knuteson,<sup>32</sup> B. R. Ko,<sup>15</sup> H. Kobayashi,<sup>55</sup> K. Kondo,<sup>57</sup> D. J. Kong,<sup>27</sup> J. Konigsberg,<sup>17</sup> A. Korytov,<sup>17</sup> A. V. Kotwal,<sup>15</sup> A. Kovalev,<sup>45</sup> A. Kraan,<sup>45</sup> J. Kraus,<sup>23</sup> I. Kravchenko,<sup>32</sup> M. Kreps,<sup>25</sup> J. Kroll,<sup>45</sup> N. Krumnack,<sup>4</sup> M. Kruse,<sup>15</sup> V. Krutelyov,<sup>53</sup> S. E. Kuhlmann,<sup>2</sup> Y. Kusakabe,<sup>57</sup> S. Kwang,<sup>13</sup> A. T. Laasanen,<sup>48</sup> S. Lai,<sup>33</sup> S. Lami,<sup>46</sup> S. Lammel,<sup>16</sup> M. Lancaster,<sup>30</sup> R. L. Lander,<sup>7</sup> K. Lannon,<sup>39</sup> A. Lath,<sup>52</sup> G. Latino,<sup>46</sup> I. Lazzizzera,<sup>43</sup> T. LeCompte,<sup>2</sup> J. Lee,<sup>49</sup> J. Lee,<sup>27</sup> Y. J. Lee,<sup>27</sup> S. W. Lee,<sup>53</sup> R. Lefèvre,<sup>3</sup> N. Leonardo,<sup>32</sup> S. Leone,<sup>46</sup> S. Levy,<sup>13</sup> J. D. Lewis,<sup>16</sup> C. Lin,<sup>60</sup> C. S. Lin,<sup>16</sup> M. Lindgren,<sup>16</sup> E. Lipeles,<sup>9</sup> T. M. Liss,<sup>23</sup> A. Lister,<sup>19</sup> D. O. Litvintsev,<sup>16</sup> T. Liu,<sup>16</sup> N. S. Lockyer,<sup>45</sup> A. Loginov,<sup>36</sup> M. Loreti,<sup>43</sup> P. Loverre,<sup>51</sup> R.-S. Lu,<sup>1</sup> D. Lucchesi,<sup>43</sup> P. Lujan,<sup>28</sup> P. Lukens,<sup>16</sup> G. Lungu,<sup>17</sup> L. Lyons,<sup>42</sup> J. Lys,<sup>28</sup> R. Lysak,<sup>1</sup> E. Lytken,<sup>48</sup> P. Mack,<sup>25</sup> D. MacQueen,<sup>33</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>16</sup> T. Maki,<sup>22</sup> P. Maksimovic,<sup>24</sup> S. Malde,<sup>42</sup> G. Manca,<sup>29</sup> F. Margaroli,<sup>5</sup> R. Marginean,<sup>16</sup> C. Marino,<sup>23</sup> A. Martin,<sup>60</sup> V. Martin,<sup>38</sup> M. Martínez,<sup>3</sup> T. Maruyama,<sup>55</sup> P. Mastrandrea,<sup>51</sup> H. Matsunaga,<sup>55</sup> M. E. Mattson,<sup>58</sup> R. Mazini,<sup>33</sup> P. Mazzanti,<sup>5</sup> K. S. McFarland,<sup>49</sup> P. McIntyre,<sup>53</sup> R. McNulty,<sup>29</sup> A. Mehta,<sup>29</sup> S. Menzemer,<sup>11</sup> A. Menzione,<sup>46</sup> P. Merkel,<sup>48</sup> C. Mesropian,<sup>50</sup> A. Messina,<sup>51</sup> M. von der Mey,<sup>8</sup> T. Miao,<sup>16</sup> N. Miladinovic,<sup>6</sup> J. Miles,<sup>32</sup> R. Miller,<sup>35</sup> J. S. Miller,<sup>34</sup> C. Mills,<sup>10</sup> M. Milnik,<sup>25</sup> R. Miquel,<sup>28</sup> A. Mitra,<sup>1</sup> G. Mitselmakher,<sup>17</sup> A. Miyamoto,<sup>26</sup> N. Moggi,<sup>5</sup> B. Mohr,<sup>8</sup> R. Moore,<sup>16</sup> M. Morello,<sup>46</sup> P. Movilla Fernandez,<sup>28</sup> J. Mülmenstädt,<sup>28</sup> A. Mukherjee,<sup>16</sup> Th. Müller,<sup>25</sup> R. Mumford,<sup>24</sup> P. Murat,<sup>16</sup> J. Nachtman,<sup>16</sup> J. Naganoma,<sup>57</sup> S. Nahn,<sup>32</sup> I. Nakano,<sup>40</sup> A. Napier,<sup>56</sup> D. Naumov,<sup>37</sup> V. Necula,<sup>17</sup> C. Neu,<sup>45</sup> M. S. Neubauer,<sup>9</sup>

J. Nielsen,<sup>28</sup> T. Nigmanov,<sup>47</sup> L. Nodulman,<sup>2</sup> O. Norriella,<sup>3</sup> E. Nurse,<sup>30</sup> T. Ogawa,<sup>57</sup> S. H. Oh,<sup>15</sup> Y. D. Oh,<sup>27</sup> T. Okusawa,<sup>41</sup> R. Oldeman,<sup>29</sup> R. Orava,<sup>22</sup> K. Osterberg,<sup>22</sup> C. Pagliarone,<sup>46</sup> E. Palencia,<sup>11</sup> R. Paoletti,<sup>46</sup> V. Papadimitriou,<sup>16</sup> A. A. Paramonov,<sup>13</sup> B. Parks,<sup>39</sup> S. Pashapour,<sup>33</sup> J. Patrick,<sup>16</sup> G. Pauletta,<sup>54</sup> M. Paulini,<sup>12</sup> C. Paus,<sup>32</sup> D. E. Pellett,<sup>7</sup> A. Penzo,<sup>54</sup> T. J. Phillips,<sup>15</sup> G. Piacentino,<sup>46</sup> J. Piedra,<sup>44</sup> L. Pinera,<sup>17</sup> K. Pitts,<sup>23</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>59</sup> X. Portell,<sup>3</sup> O. Poukhov,<sup>14</sup> N. Pounder,<sup>42</sup> F. Prakoshyn,<sup>14</sup> A. Pronko,<sup>16</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>18</sup> G. Punzi,<sup>46</sup> J. Pursley,<sup>24</sup> J. Rademacker,<sup>42</sup> A. Rahaman,<sup>47</sup> A. Rakitin,<sup>32</sup> S. Rappoccio,<sup>21</sup> F. Ratnikov,<sup>52</sup> B. Reiser,<sup>16</sup> V. Rekovic,<sup>37</sup> N. van Remortel,<sup>22</sup> P. Renton,<sup>42</sup> M. Rescigno,<sup>51</sup> S. Richter,<sup>25</sup> F. Rimondi,<sup>5</sup> L. Ristori,<sup>46</sup> W. J. Robertson,<sup>15</sup> A. Robson,<sup>20</sup> T. Rodrigo,<sup>11</sup> E. Rogers,<sup>23</sup> S. Rolli,<sup>56</sup> R. Roser,<sup>16</sup> M. Rossi,<sup>54</sup> R. Rossin,<sup>17</sup> C. Rott,<sup>48</sup> A. Ruiz,<sup>11</sup> J. Russ,<sup>12</sup> V. Rusu,<sup>13</sup> H. Saarikko,<sup>22</sup> S. Sabik,<sup>33</sup> A. Safonov,<sup>53</sup> W. K. Sakumoto,<sup>49</sup> G. Salamanna,<sup>51</sup> O. Saltó,<sup>3</sup> D. Saltzberg,<sup>8</sup> C. Sanchez,<sup>3</sup> L. Santi,<sup>54</sup> S. Sarkar,<sup>51</sup> L. Sartori,<sup>46</sup> K. Sato,<sup>55</sup> P. Savard,<sup>33</sup> A. Savoy-Navarro,<sup>44</sup> T. Scheidle,<sup>25</sup> P. Schlabach,<sup>16</sup> E. E. Schmidt,<sup>16</sup> M. P. Schmidt,<sup>60</sup> M. Schmitt,<sup>38</sup> T. Schwarz,<sup>34</sup> L. Scodellaro,<sup>11</sup> A. L. Scott,<sup>10</sup> A. Scribano,<sup>46</sup> F. Scuri,<sup>46</sup> A. Sedov,<sup>48</sup> S. Seidel,<sup>37</sup> Y. Seiya,<sup>41</sup> A. Semenov,<sup>14</sup> L. Sexton-Kennedy,<sup>16</sup> I. Sfiligoi,<sup>18</sup> M. D. Shapiro,<sup>28</sup> T. Shears,<sup>29</sup> P. F. Shepard,<sup>47</sup> D. Sherman,<sup>21</sup> M. Shimojima,<sup>55</sup> M. Shochet,<sup>13</sup> Y. Shon,<sup>59</sup> I. Shreyber,<sup>36</sup> A. Sidoti,<sup>44</sup> P. Sinervo,<sup>33</sup> A. Sisakyan,<sup>14</sup> J. Sjolín,<sup>42</sup> A. Skiba,<sup>25</sup> A. J. Slaughter,<sup>16</sup> K. Sliwa,<sup>56</sup> J. R. Smith,<sup>7</sup> F. D. Snider,<sup>16</sup> R. Snihur,<sup>33</sup> M. Soderberg,<sup>34</sup> A. Soha,<sup>7</sup> S. Somalwar,<sup>52</sup> V. Sorin,<sup>35</sup> J. Spalding,<sup>16</sup> M. Spezziga,<sup>16</sup> F. Spinella,<sup>46</sup> T. Spreitzer,<sup>33</sup> P. Squillacioti,<sup>46</sup> M. Stanitzki,<sup>60</sup> A. Staveris-Polykalas,<sup>46</sup> R. St. Denis,<sup>20</sup> B. Stelzer,<sup>8</sup> O. Stelzer-Chilton,<sup>42</sup> D. Stentz,<sup>38</sup> J. Strologas,<sup>37</sup> D. Stuart,<sup>10</sup> J. S. Suh,<sup>27</sup> A. Sukhanov,<sup>17</sup> K. Sumorok,<sup>32</sup> H. Sun,<sup>56</sup> T. Suzuki,<sup>55</sup> A. Taffard,<sup>23</sup> R. Takashima,<sup>40</sup> Y. Takeuchi,<sup>55</sup> K. Takikawa,<sup>55</sup> M. Tanaka,<sup>2</sup> R. Tanaka,<sup>40</sup> N. Tanimoto,<sup>40</sup> M. Tecchio,<sup>34</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>50</sup> S. Tether,<sup>32</sup> J. Thom,<sup>16</sup> A. S. Thompson,<sup>20</sup> E. Thomson,<sup>45</sup> P. Tipton,<sup>49</sup> V. Tiwari,<sup>12</sup> S. Tkaczyk,<sup>16</sup> D. Toback,<sup>53</sup> S. Tokar,<sup>14</sup> K. Tollefson,<sup>35</sup> T. Tomura,<sup>55</sup> D. Tonelli,<sup>46</sup> M. Tönnemann,<sup>35</sup> S. Torre,<sup>18</sup> D. Torretta,<sup>16</sup> S. Tourneur,<sup>44</sup> W. Trischuk,<sup>33</sup> R. Tsuchiya,<sup>57</sup> S. Tsuno,<sup>40</sup> N. Turini,<sup>46</sup> F. Ukegawa,<sup>55</sup> T. Unverhau,<sup>20</sup> S. Uozumi,<sup>55</sup> D. Usynin,<sup>45</sup> A. Vaiciulis,<sup>49</sup> S. Vallecorsa,<sup>19</sup> A. Varganov,<sup>34</sup> E. Vataga,<sup>37</sup> G. Velev,<sup>16</sup> G. Veramendi,<sup>23</sup> V. Veszpremi,<sup>48</sup> R. Vidal,<sup>16</sup> I. Vila,<sup>11</sup> R. Vilar,<sup>11</sup> T. Vine,<sup>30</sup> I. Vollrath,<sup>33</sup> I. Volobouev,<sup>28</sup> G. Volpi,<sup>46</sup> F. Würthwein,<sup>9</sup> P. Wagner,<sup>53</sup> R. G. Wagner,<sup>2</sup> R. L. Wagner,<sup>16</sup> W. Wagner,<sup>25</sup> R. Wallny,<sup>8</sup> T. Walter,<sup>25</sup> Z. Wan,<sup>52</sup> S. M. Wang,<sup>1</sup> A. Warburton,<sup>33</sup> S. Waschke,<sup>20</sup> D. Waters,<sup>30</sup> W. C. Wester III,<sup>16</sup> B. Whitehouse,<sup>56</sup> D. Whiteson,<sup>45</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>16</sup> G. Williams,<sup>33</sup> H. H. Williams,<sup>45</sup> P. Wilson,<sup>16</sup> B. L. Winer,<sup>39</sup> P. Wittich,<sup>16</sup> S. Wolbers,<sup>16</sup> C. Wolfe,<sup>13</sup> T. Wright,<sup>34</sup> X. Wu,<sup>19</sup> S. M. Wynne,<sup>29</sup> A. Yagil,<sup>16</sup> K. Yamamoto,<sup>41</sup> J. Yamaoka,<sup>52</sup> T. Yamashita,<sup>40</sup> C. Yang,<sup>60</sup> U. K. Yang,<sup>13</sup> Y. C. Yang,<sup>27</sup> W. M. Yao,<sup>28</sup> G. P. Yeh,<sup>16</sup> J. Yoh,<sup>16</sup> K. Yorita,<sup>13</sup> T. Yoshida,<sup>41</sup> G. B. Yu,<sup>49</sup> I. Yu,<sup>27</sup> S. S. Yu,<sup>16</sup> J. C. Yun,<sup>16</sup> L. Zanello,<sup>51</sup> A. Zanetti,<sup>54</sup> I. Zaw,<sup>21</sup> F. Zetti,<sup>46</sup> X. Zhang,<sup>23</sup> J. Zhou,<sup>52</sup> and S. Zucchelli<sup>5</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*<sup>3</sup>*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193 Bellaterra (Barcelona), Spain*<sup>4</sup>*Baylor University, Waco, Texas 76798, USA*<sup>5</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*<sup>6</sup>*Brandeis University, Waltham, Massachusetts 02254, USA*<sup>7</sup>*University of California, Davis, Davis, California 95616, USA*<sup>8</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*<sup>9</sup>*University of California, San Diego, La Jolla, California 92093, USA*<sup>10</sup>*University of California, Santa Barbara, Santa Barbara, California 93106, USA*<sup>11</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*<sup>12</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*<sup>13</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*<sup>14</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*<sup>15</sup>*Duke University, Durham, North Carolina 27708, USA*<sup>16</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*<sup>17</sup>*University of Florida, Gainesville, Florida 32611, USA*<sup>18</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*<sup>19</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*<sup>20</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*<sup>21</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*<sup>22</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014 Helsinki, Finland*

- <sup>23</sup>University of Illinois, Urbana, Illinois 61801, USA  
<sup>24</sup>The Johns Hopkins University, Baltimore, Maryland 21218, USA  
<sup>25</sup>Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany  
<sup>26</sup>High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan  
<sup>27</sup>Center for High Energy Physics: Kyungpook National University, Taegu 702-701, Korea;  
 Seoul National University, Seoul 151-742, Korea;  
 and SungKyunKwan University, Suwon 440-746, Korea  
<sup>28</sup>Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA  
<sup>29</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom  
<sup>30</sup>University College London, London WC1E 6BT, United Kingdom  
<sup>31</sup>Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain  
<sup>32</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA  
<sup>33</sup>Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8;  
 and University of Toronto, Toronto, Canada M5S 1A7  
<sup>34</sup>University of Michigan, Ann Arbor, Michigan 48109, USA  
<sup>35</sup>Michigan State University, East Lansing, Michigan 48824, USA  
<sup>36</sup>Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia  
<sup>37</sup>University of New Mexico, Albuquerque, New Mexico 87131, USA  
<sup>38</sup>Northwestern University, Evanston, Illinois 60208, USA  
<sup>39</sup>The Ohio State University, Columbus, Ohio 43210, USA  
<sup>40</sup>Okayama University, Okayama 700-8530, Japan  
<sup>41</sup>Osaka City University, Osaka 588, Japan  
<sup>42</sup>University of Oxford, Oxford OX1 3RH, United Kingdom  
<sup>43</sup>Istituto Nazionale di Fisica Nucleare, University of Padova, Sezione di Padova-Trento, I-35131 Padova, Italy  
<sup>44</sup>LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris F-75252, France  
<sup>45</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA  
<sup>46</sup>Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena, and Scuola Normale Superiore, I-56127 Pisa, Italy  
<sup>47</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA  
<sup>48</sup>Purdue University, West Lafayette, Indiana 47907, USA  
<sup>49</sup>University of Rochester, Rochester, New York 14627, USA  
<sup>50</sup>The Rockefeller University, New York, New York 10021, USA  
<sup>51</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Roma I, University of Rome "La Sapienza," I-00185 Roma, Italy  
<sup>52</sup>Rutgers University, Piscataway, New Jersey 08855, USA  
<sup>53</sup>Texas A&M University, College Station, Texas 77843, USA  
<sup>54</sup>Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy  
<sup>55</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan  
<sup>56</sup>Tufts University, Medford, Massachusetts 02155, USA  
<sup>57</sup>Waseda University, Tokyo 169, Japan  
<sup>58</sup>Wayne State University, Detroit, Michigan 48201, USA  
<sup>59</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>60</sup>Yale University, New Haven, Connecticut 06520, USA  
 (Received 13 June 2006; published 10 August 2006)

We present the first precise measurement of the  $B_s^0-\bar{B}_s^0$  oscillation frequency  $\Delta m_s$ . We use  $1 \text{ fb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  collected with the CDF II detector at the Fermilab Tevatron. The sample contains signals of 3600 fully reconstructed hadronic  $B_s$  decays and 37 000 partially reconstructed semileptonic  $B_s$  decays. We measure the probability as a function of proper decay time that the  $B_s$  decays with the same, or opposite, flavor as the flavor at production, and we find a signal consistent with  $B_s^0-\bar{B}_s^0$  oscillations. The probability that random fluctuations could produce a comparable signal is 0.2%. Under the hypothesis that the signal is due to  $B_s^0-\bar{B}_s^0$  oscillations, we measure  $\Delta m_s = 17.31^{+0.33}_{-0.18}(\text{stat}) \pm 0.07(\text{syst}) \text{ ps}^{-1}$  and determine  $|V_{td}/V_{ts}| = 0.208^{+0.001}_{-0.002}(\text{expt})^{+0.008}_{-0.006}(\text{theor})$ .

DOI: 10.1103/PhysRevLett.97.062003

PACS numbers: 14.40.Nd, 12.15.Ff, 12.15.Hh, 13.20.He

Neutral  $B$  mesons ( $b\bar{q}$ , with  $q = d, s$  for  $\bar{B}_d^0, \bar{B}_s^0$ ) oscillate from particle to antiparticle due to flavor-changing weak interactions. The probability density  $P_+$  ( $P_-$ ) for a  $\bar{B}_q^0$  meson produced at proper time  $t = 0$  to decay as a  $\bar{B}_q^0$  ( $B_q^0$ ) at time  $t$  is given by

$$P_{\pm}(t) = \frac{\Gamma_q}{2} e^{-\Gamma_q t} [1 \pm \cos(\Delta m_q t)],$$

where  $\Delta m_q$  is the mass difference between the two mass eigenstates  $B_{q,H}^0$  and  $B_{q,L}^0$  [1], and  $\Gamma_q$  is the decay width,

which is assumed to be equal for the two mass eigenstates. The mass differences  $\Delta m_d$  and  $\Delta m_s$  can be used to determine the fundamental parameters  $|V_{td}|$  and  $|V_{ts}|$ , respectively, of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [2], which relates the quark mass eigenstates to the flavor eigenstates. This determination, however, has large theoretical uncertainties. A measurement of  $\Delta m_s$  combined with  $\Delta m_d = 0.505 \pm 0.005 \text{ ps}^{-1}$  [3,4] would determine the ratio  $|V_{td}/V_{ts}|$  with a significantly smaller theoretical uncertainty, contributing to a stringent test of the unitarity of the CKM matrix. Earlier attempts to measure  $\Delta m_s$  have yielded a lower limit:  $\Delta m_s > 14.5 \text{ ps}^{-1}$  [3,5] at the 95% confidence level (C.L.). Recently the D0 Collaboration reported  $17 \text{ ps}^{-1} < \Delta m_s < 21 \text{ ps}^{-1}$  at 90% C.L. [6] using a large sample of semileptonic  $B_s$  [7] decays.

In this Letter we report a measurement of  $\Delta m_s$  using data from  $1 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  collected by the CDF II detector at the Fermilab Tevatron. We begin by reconstructing  $B_s$  decays in hadronic ( $\bar{B}_s^0 \rightarrow D_s^+ \pi^-, D_s^+ \pi^- \pi^+ \pi^-$ ) and semileptonic ( $\bar{B}_s^0 \rightarrow D_s^{+(*)} \ell^- \bar{\nu}_\ell$ ,  $\ell = e$  or  $\mu$ ) decay modes using charged particles only [8]. Using the method of maximum likelihood, we extract the value of  $\Delta m_s$  from the probability density functions (PDFs) that describe the measured time development of  $B_s$  mesons that decay with the same or opposite flavor as their flavor at production. The proper decay time for each  $B_s$  is calculated from the measured distance between the production and decay points, the measured momentum, and the  $B_s$  mass  $m_{B_s} = 5.3696 \text{ GeV}/c^2$  [3]. The  $B_s$  flavor ( $b$  or  $\bar{b}$ ) at decay is determined unambiguously by the charges of the decay products.

To identify the flavor of the  $B_s$  at production, we use characteristics of  $b$  quark production and fragmentation in  $p\bar{p}$  collisions. At the Tevatron, the dominant  $b$  quark production mechanisms produce  $b\bar{b}$  pairs. The  $b$  and  $\bar{b}$  are expected to fragment independently into hadrons. In a simple model of fragmentation, a  $b$  quark becomes a  $\bar{B}_s^0$  meson when some of the energy of the  $b$  quark is used to produce an  $s\bar{s}$  quark pair. The  $b$  and the  $\bar{s}$  bind to form a  $\bar{B}_s^0$ . The remaining  $s$  quark may form a  $K^-$ . Similarly, a  $\bar{b}$  that becomes a  $B_s^0$  is accompanied by a  $K^+$ . One of the two techniques used to identify the production flavor of the  $B_s$  is based on the charge of these kaons (same-side tag). The second technique uses the charge of the lepton from semileptonic decays or a momentum-weighted charge of the decay products of the second  $b$  hadron produced in the collision (opposite-side tag).

The hadronic and semileptonic decay modes are complementary. Because of the large branching ratio, the semileptonic decays provide a tenfold advantage in signal rate at the cost of significantly worsened decay-time resolution due to the unmeasured  $\nu$  momentum. Semileptonic decays dominate the sensitivity to oscillations at lower values of  $\Delta m_s$ . The fully reconstructed hadronic  $B_s$  decays have

superior decay-time resolution, and our large sample of these decays is the unique feature that makes CDF sensitive to much larger values of  $\Delta m_s$  than other experiments.

The CDF II detector [9] consists of a magnetic spectrometer surrounded by electromagnetic and hadronic calorimeters and muon detectors [10]. The key features for this measurement include precision vertex determination provided by the seven-layer double-sided inner silicon strip detector [11,12] supplemented with a single-sided layer of silicon [13] mounted directly on the beam pipe at an average radius of 1.5 cm. The 96-layer outer drift chamber [14] is used for both precision tracking and  $dE/dx$  particle identification. Time-of-flight (TOF) counters [15] located just outside the drift chamber are used to identify low momentum charged kaons.

Charm and bottom hadrons are selected using a three-level trigger system that exploits the kinematics of production and decay, and the long lifetimes of  $D$  and  $B$  mesons. A crucial component of the trigger system for this measurement is the Silicon Vertex Trigger [16], which selects events that contain  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  and  $D_s^+ \pi^- \pi^+ \pi^-$  decays. The trigger configuration used to collect the heavy flavor data sample is described in [17].

To reconstruct  $\bar{B}_s^0$  candidates, we first select  $D_s^+$  candidates. We use  $D_s^+ \rightarrow \phi \pi^+$ ,  $K^*(892)^0 K^+$ , and  $\pi^+ \pi^- \pi^+$ , with  $\phi \rightarrow K^+ K^-$  and  $K^{*0} \rightarrow K^+ \pi^-$ ; we require that  $\phi$  and  $K^{*0}$  candidates be consistent with the known masses and widths [3] of these two resonances. These  $D_s^+$  candidates are combined with one or three additional charged particles to form  $D_s^+ \ell^-$ ,  $D_s^+ \pi^-$ , or  $D_s^+ \pi^- \pi^+ \pi^-$  candidates. The  $D_s^+$  and other decay products of a  $\bar{B}_s^0$  candidate are constrained to originate from a common vertex in three dimensions. For the  $K^*(892)^0 K^+$  final state, we remove candidates that are consistent with the decay  $D^+ \rightarrow K^- \pi^+ \pi^+$ . We use a likelihood technique to identify muons [18] and electrons [19].

Backgrounds are suppressed by imposing a requirement on the minimum transverse momentum  $p_T$  [20] of the  $\bar{B}_s^0$  and by requiring that the  $\bar{B}_s^0$  and  $D_s^+$  decay vertices are displaced significantly from the  $p\bar{p}$  collision position. We find signals of 3600 hadronic  $B_s$  decays and 37 000 semileptonic  $B_s$  decays.

For the hadronic decays, the invariant mass distribution (see Fig. 1) has a signal centered close to  $m_{B_s} = 5.3696 \text{ GeV}/c^2$  with a width of 14 to 20  $\text{MeV}/c^2$ , depending on the decay mode. Candidates with masses greater than 5.5  $\text{GeV}/c^2$  are used to construct PDFs for combinatorial background. To remove contributions from  $\bar{B}_s^0 \rightarrow D_s^{*+} \pi^-$ ,  $\bar{B}_s^0 \rightarrow D_s^+ \rho^-$ , and semileptonic and other partially reconstructed decays, we require the mass of the decay candidates to be greater than 5.3  $\text{GeV}/c^2$ . For semileptonic decays we take into account several background contributions, including  $B$  meson decays to two charm mesons and real  $D_s$  mesons associated with a false lepton.

The decay time in the  $B_s$  rest frame is  $t = \kappa[L_T m_{B_s}/p_T]$ , where  $L_T$  is the displacement of the  $B_s$




FIG. 1. The invariant mass distributions for  $\bar{B}_s^0 \rightarrow D_s^+ \pi^-$  (left panel) and  $D_s^+ \pi^- \pi^+ \pi^-$  (right panel).

decay vertex with respect to the primary vertex projected onto the  $B_s$  transverse momentum vector. The factor  $\kappa$  corrects for missing momentum in the semileptonic decays ( $\kappa = 1$  for hadronic decays). To improve the decay-time resolution, we use event-by-event primary-vertex position measurements when computing the  $B_s$  vertex displacement. The signal decay-time distribution is modeled with  $P(t_i, \sigma_{t_i}) = \varepsilon(t_i) \int \Gamma_s e^{-\Gamma_s t'} \mathcal{G}(t' - t_i, \sigma_{t_i}) dt'$ , where  $t_i$  is the measured decay time of the  $i$ th candidate,  $\Gamma_s$  is the  $B_s$  decay width,  $\mathcal{G}(x - \mu, \sigma)$  is a Gaussian distribution of the random variable  $x$  with mean  $\mu$  and width  $\sigma$ , and  $\sigma_{t_i}$  is the estimated candidate decay-time resolution. The decay-time efficiency function  $\varepsilon(t)$  describes trigger and selection biases on the decay-time distribution and is determined from Monte Carlo simulation. For semileptonic decays, the  $\kappa$  distribution is determined from Monte Carlo simulation and is convoluted with the signal decay-time distribution. The missing transverse momentum from unreconstructed particles in the semileptonic decays is an important contribution to the decay-time resolution. To reduce this contribution and make optimal use of the semileptonic decays, we determine the  $\kappa$  distribution as a function of the invariant mass of the  $D_s \ell$  pair,  $m_{D_s \ell}$ . The rms width of the  $\kappa$  distribution is 3% (20%) for  $m_{D_s \ell} = 5.2 \text{ GeV}/c^2$  ( $3.0 \text{ GeV}/c^2$ ).

We estimate the decay-time resolution  $\sigma_{t_i}$  for each candidate using the measured track parameters and their estimated uncertainties. We calibrate this estimate using a large sample of prompt  $D^+$  mesons [21], which we combine with one or three charged particles from the primary vertex to mimic signal topologies. For hadronic decays, the average decay-time resolution is 87 fs, which corresponds to one-fifth of an oscillation period at the lower limit on  $\Delta m_s$  ( $14.5 \text{ ps}^{-1}$ ). For semileptonic decays, the decay-time resolution is worse due to decay topology and the missing momentum of unreconstructed decay products. For example, at  $t = 0$ ,  $\sigma_{t_i} = 100 \text{ fs}$  (200 fs) for  $m_{D_s \ell} = 5.2 \text{ GeV}/c^2$  ( $3.0 \text{ GeV}/c^2$ ) and increases to  $\sigma_{t_i} = 115 \text{ fs}$  (380 fs) at  $t = 1.5 \text{ ps}$ .

The flavor of the  $B_s$  at production is determined using both opposite-side and same-side flavor tagging tech-

niques. The effectiveness  $Q \equiv \varepsilon \mathcal{D}^2$  of these techniques is quantified with an efficiency  $\varepsilon$ , the fraction of signal candidates with a flavor tag, and a dilution  $\mathcal{D} \equiv 1 - 2w$ , where  $w$  is the probability that the tag is incorrect.

Opposite-side tags infer the production flavor of the  $B_s$  from the decay products of the  $b$  hadron produced from the other  $b$  quark in the event. We use lepton ( $e$  and  $\mu$ ) charge and jet charge as tags, building on techniques developed for a CDF run I measurement of  $\Delta m_d$  [22]. If both lepton and jet-charge tags are present, we use the lepton tag, which has a higher average dilution.

The dilution of opposite-side flavor tags is expected to be independent of the type of  $B$  meson that produces the hadronic or semileptonic decay. The dilution is measured in data using large samples of  $B^-$ , which do not change flavor, and  $\bar{B}^0$ , which can be used after accounting for their well-known oscillation frequency. The combined opposite-side tag effectiveness is  $Q = 1.5\% \pm 0.1\%$ , where the uncertainty is dominated by the statistics of the control samples.

Same-side flavor tags [23] are based on the charges of associated particles produced in the fragmentation of the  $b$  quark that produces the reconstructed  $B_s$ . In the simplest picture of fragmentation, a  $\pi^+$  ( $\pi^-$ ) accompanies the formation of a  $B^-$  ( $B^+$ ), a  $\pi^-$  ( $\pi^+$ ) accompanies a  $\bar{B}^0$  ( $B^0$ ), and a  $K^-$  ( $K^+$ ) accompanies a  $\bar{B}_s^0$  ( $B_s^0$ ). In run I, CDF established this method of production flavor identification in measurements of  $\Delta m_d$  [24] and the  $CP$  symmetry violating parameter  $\sin(2\beta)$  [25]. In this analysis, we use  $dE/dx$  [19] and TOF information in a combined particle identification likelihood to identify the kaons associated with  $B_s$  production. Tracks close in phase space to the  $B_s$  candidate are considered as same-side kaon tag candidates, and the track with the largest kaon likelihood is selected as the tagging track.

The performance of the same-side kaon tag for  $\bar{B}_s^0$  is expected to be different than for  $B^-$  and  $\bar{B}^0$ . We predict the dilution using simulated data samples generated with the PYTHIA Monte Carlo program [26]. Control samples of  $B^-$  and  $\bar{B}^0$  are used to validate the predictions of the simulation. The effectiveness of this flavor tag increases with the  $p_T$  of the  $\bar{B}_s^0$ ; we find  $Q = 3.5\%$  (4.0%) in the hadronic (semileptonic) decay sample. The fractional uncertainty on  $Q$  is approximately 25%. This uncertainty is dominated by the differences between data and simulation for kaons found close in phase space to the  $\bar{B}_s^0$  [27] and for the performance of the same-side kaon tag when applied to  $B^-$ .

If both a same-side tag and an opposite-side tag are present, we combine the information from both tags assuming they are uncorrelated. The addition of the same-side kaon tag increases the effective sample statistics by more than a factor of 3.

We use an unbinned maximum likelihood fit to search for  $B_s$  oscillations. The likelihood combines mass, decay-



time, decay-time resolution, and flavor tagging information for each candidate and includes terms for signal and each type of background. The fit is done in three stages. First, a combined mass and decay-time fit is performed to separate signal from background and to fix mass and decay-time models. Combined fits for  $B_s$  mass (Fig. 1) and decay width in hadronic samples and for decay width in the semileptonic samples yield measurements consistent with established values [3]. Second, flavor asymmetries are measured for background components. The third step is a fit for  $B_s^0\text{-}\bar{B}_s^0$  oscillations; the mass and decay-time models and background asymmetries are fixed from the previous two stages.

The signal PDF has the general form:

$$S_{\pm}(t_i, \sigma_{t_i}, \mathcal{D}_i) = \varepsilon(t_i) \int \frac{\Gamma_s}{2} e^{-\Gamma_s t'} [1 \pm \mathcal{A} \mathcal{D}_i \cos(\Delta m_s t')] \times \mathcal{G}(t_i - t', \sigma_{t_i}) dt', \quad (1)$$

where  $\mathcal{D}_i$  is the  $i$ th candidate dilution, and  $t_i$ ,  $\sigma_{t_i}$ ,  $\mathcal{G}$ , and  $\varepsilon(t)$  have been defined previously. Following the method described in [28], we fit for the oscillation amplitude  $\mathcal{A}$  while fixing  $\Delta m_s$  to a probe value. When all detector effects ( $\mathcal{D}_i$ ,  $\sigma_{t_i}$ ) are calibrated, the oscillation amplitude is expected to be consistent with  $\mathcal{A} = 1$  when the probe value is the true oscillation frequency, and consistent with  $\mathcal{A} = 0$  when the probe value is far from the true oscillation frequency. Figure 2 (upper panel) shows the fitted value of the amplitude as a function of the oscillation frequency. The sensitivity of the measurement is defined by the maximum value of  $\Delta m_s$  where  $\mathcal{A} = 1$  is excluded at 95% C.L. if the measured value of  $\mathcal{A}$  were zero. Our sensitivity is  $25.8 \text{ ps}^{-1}$  and exceeds the combined sensitivity of all previous experiments [3]. At  $\Delta m_s = 17.3 \text{ ps}^{-1}$ , the observed amplitude  $\mathcal{A} = 1.03 \pm 0.28(\text{stat})$  is consistent with unity, indicating that the data are compatible with  $B_s^0\text{-}\bar{B}_s^0$  oscillations with that frequency, while the amplitude is inconsistent with zero:  $\mathcal{A}/\sigma_{\mathcal{A}} = 3.7$ , where  $\sigma_{\mathcal{A}}$  is the uncertainty on  $\mathcal{A}$ . The negative amplitudes measured at frequencies slightly below and slightly above the peak frequency are expected and are due to the finite range in signal decay time that is imposed by the trigger and selection criteria. The systematic uncertainty on  $\mathcal{A}$  is mainly due to uncertainties on  $\sigma_{t_i}$  and  $\mathcal{D}_i$ . Since the effect of these uncertainties on  $\mathcal{A}$  and  $\sigma_{\mathcal{A}}$  are correlated, the ratio  $\mathcal{A}/\sigma_{\mathcal{A}}$  has negligible systematic uncertainty.

The significance of the potential signal is evaluated from  $\Lambda \equiv \log[\mathcal{L}^{\mathcal{A}=0}/\mathcal{L}^{\mathcal{A}=1}(\Delta m_s)]$ , which is the logarithm of the ratio of likelihoods for the hypothesis of oscillations ( $\mathcal{A} = 1$ ) at the probe value and the hypothesis that  $\mathcal{A} = 0$ , which is equivalent to random production flavor tags. Figure 2 (lower panel) shows  $\Lambda$  as a function of  $\Delta m_s$ . Separate curves are shown for the semileptonic data alone (dash-dotted line), the hadronic data alone (dotted line), and the combined data (solid line). A minimal value of




FIG. 2. (Upper panel) The measured amplitude values and uncertainties versus the  $B_s^0\text{-}\bar{B}_s^0$  oscillation frequency  $\Delta m_s$ . Shown in light gray and dark gray are the 95% one-sided confidence level bands for statistical uncertainties only and including systematic uncertainties, respectively. (Lower panel) The logarithm of the ratio of likelihoods for amplitude equal to zero and amplitude equal to one,  $\Lambda = \log[\mathcal{L}^{\mathcal{A}=0}/\mathcal{L}^{\mathcal{A}=1}(\Delta m_s)]$ , versus the oscillation frequency. The dashed horizontal line indicates the value of  $\Lambda$  that corresponds to a probability of 1% in the case of randomly tagged data.

$\Lambda = -6.75$  is observed at  $\Delta m_s = 17.3 \text{ ps}^{-1}$ . The significance of the signal is quantified by the probability that randomly tagged data would produce a value of  $\Lambda$  lower than  $-6.75$  at any value of  $\Delta m_s$ . We repeat the fit 50 000 times with random tagging decisions, and we find this probability is 0.2%.

Under the hypothesis that the signal is due to  $B_s^0\text{-}\bar{B}_s^0$  oscillations, we fix  $\mathcal{A} = 1$  and fit for the oscillation frequency. We find  $\Delta m_s = 17.31^{+0.33}_{-0.18}(\text{stat}) \pm 0.07(\text{syst}) \text{ ps}^{-1}$  and the range  $17.01 \text{ ps}^{-1} < \Delta m_s < 17.84 \text{ ps}^{-1}$  ( $16.96 \text{ ps}^{-1} < \Delta m_s < 17.91 \text{ ps}^{-1}$ ) at 90% (95%) C.L. All systematic uncertainties affecting  $\mathcal{A}$  are unimportant for  $\Delta m_s$ . The only non-negligible systematic uncertainty on  $\Delta m_s$  is from the uncertainty on the absolute scale of the decay-time measurement. Contributions to this uncertainty include biases in the primary-vertex reconstruction due to the presence of the opposite-side  $b$  hadron, uncertainties in the silicon-detector alignment, and biases in track fitting. The measured  $B_s^0\text{-}\bar{B}_s^0$  oscillation frequency is used to derive the ratio  $|V_{td}/V_{ts}| = \xi \sqrt{\frac{\Delta m_d}{\Delta m_s} \frac{m_{B^0}}{m_{B^0}}}$ . As inputs we use  $m_{B^0}/m_{B^0} = 0.98390$  [29] with negligible uncertainty,  $\Delta m_d = 0.505 \pm 0.005 \text{ ps}^{-1}$  [3], and  $\xi = 1.21^{+0.047}_{-0.035}$  [30]. We find  $|V_{td}/V_{ts}| = 0.208^{+0.001}_{-0.002}(\text{expt})^{+0.008}_{-0.006}(\text{theor})$ .

In conclusion, we present the first precise measurement of  $\Delta m_s$ . The value of  $\Delta m_s$  is consistent with standard model expectations [31] and with previous bounds. Our

measured value of  $\Delta m_s$  allows us to determine  $|V_{td}/V_{ts}|$  with unprecedented precision and can be used to improve constraints on the unitarity of the CKM matrix and on scenarios involving new physics.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, U.K.; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; in part by the European Community's Human Potential Programme under Contract No. HPRN-CT-2002-00292; and the Academy of Finland.

- 
- [1] C. Gay, *Annu. Rev. Nucl. Part. Sci.* **50**, 577 (2000). We set  $\hbar = c = 1$  and report  $\Delta m_q = m_{B_{q,H}^0} - m_{B_{q,L}^0}$  in inverse picoseconds.
- [2] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
- [3] S. Eidelman *et al.*, *Phys. Lett. B* **592**, 1 (2004), and 2005 partial update for the 2006 edition available on the PDG WWW page (<http://pdg.lbl.gov/>).
- [4] K. Abe *et al.* (BELLE Collaboration), *Phys. Rev. D* **71**, 072003 (2005); **71**, 079903(E) (2005); N. C. Hastings *et al.* (BELLE Collaboration), *Phys. Rev. D* **67**, 052004 (2003); B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **88**, 221803 (2002).
- [5] J. Abdallah *et al.* (DELPHI Collaboration), *Eur. Phys. J. C* **35**, 35 (2004); K. Abe *et al.* (SLD Collaboration), *Phys. Rev. D* **67**, 012006 (2003); A. Heister *et al.* (ALEPH Collaboration), *Eur. Phys. J. C* **29**, 143 (2003).
- [6] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **97**, 021802 (2006).
- [7] The symbol  $B_s$  refers to the combination of  $\bar{B}_s^0$  and  $B_s^0$  decays.
- [8] References to a particular process imply that the charge conjugate process is included as well.
- [9] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005); R. Blair *et al.* (CDF Collaboration), Fermilab Report No. FERMILAB-PUB-96-390-E, 1996.
- [10] A. Abulencia *et al.* (CDF Collaboration), hep-ex/0508029.
- [11] A. Sill *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **447**, 1 (2000).
- [12] A. Affolder *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **453**, 84 (2000).
- [13] C. S. Hill *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **530**, 1 (2004).
- [14] T. Affolder *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **526**, 249 (2004).
- [15] S. Cabrera *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **494**, 416 (2002).
- [16] W. Ashmanskas *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **518**, 532 (2004).
- [17] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **96**, 191801 (2006).
- [18] G. Giurgiu, Ph.D. thesis, Carnegie Mellon University (Fermilab Report No. FERMILAB-THESIS-2005-41, 2005).
- [19] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **97**, 012002 (2006).
- [20] The transverse momentum  $p_T$  is the magnitude of the component of the momentum perpendicular to the proton beam direction.
- [21] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **91**, 241804 (2003).
- [22] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **60**, 072003 (1999).
- [23] A. Ali and F. Barreiro, *Z. Phys. C* **30**, 635 (1986); M. Gronau, A. Nippe, and J. L. Rosner, *Phys. Rev. D* **47**, 1988 (1993); M. Gronau and J. L. Rosner, *Phys. Rev. D* **49**, 254 (1994).
- [24] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **59**, 032001 (1999).
- [25] T. Affolder *et al.* (CDF Collaboration), *Phys. Rev. D* **61**, 072005 (2000).
- [26] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001). We use version 6.216.
- [27] D. Usynin, Ph.D. thesis, University of Pennsylvania (Fermilab Report No. FERMILAB-THESIS-2005-68, 2005).
- [28] H. G. Moser and A. Roussarie, *Nucl. Instrum. Methods Phys. Res., Sect. A* **384**, 491 (1997).
- [29] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **96**, 202001 (2006).
- [30] M. Okamoto, *Proc. Sci. LAT2005* (2005) 013 [hep-lat/0510113].
- [31] M. Bona *et al.* (UTfit Collaboration), *J. High Energy Phys.* **07** (2005) 028; J. Charles *et al.* (CKMfitter Collaboration), *Eur. Phys. J. C* **41**, 1 (2005).