

## Measurement of the Proton and Deuteron Spin Structure Function $g_1$ in the Resonance Region\*

K. Abe,<sup>15</sup> T. Akagi,<sup>12,15</sup> P. L. Anthony,<sup>12</sup> R. Antonov,<sup>11</sup> R. G. Arnold,<sup>1</sup>  
 T. Averett,<sup>16,††</sup> H. R. Band,<sup>18</sup> J. M. Bauer,<sup>7</sup> H. Borel,<sup>5</sup> P. E. Bosted,<sup>1</sup>  
 V. Breton,<sup>3</sup> J. Button-Shafer,<sup>7</sup> J. P. Chen,<sup>16,8</sup> T. E. Chupp,<sup>8</sup> J. Clendenin,<sup>12</sup>  
 C. Comptour,<sup>3</sup> K. P. Coulter,<sup>8</sup> G. Court,<sup>12,\*</sup> D. Crabb,<sup>16</sup> M. Daoudi,<sup>12</sup>  
 D. Day,<sup>16</sup> F. S. Dietrich,<sup>6</sup> J. Dunne,<sup>1</sup> H. Dutz,<sup>12,\*\*</sup> R. Erbacher,<sup>12,13</sup>  
 J. Fellbaum,<sup>1</sup> A. Feltham,<sup>2</sup> H. Fonvieille,<sup>3</sup> E. Frlez,<sup>16</sup> D. Garvey,<sup>9</sup>  
 R. Gearhart,<sup>12</sup> J. Gomez,<sup>4</sup> P. Grenier,<sup>5</sup> K. A. Griffioen,<sup>11,17</sup>  
 S. Hoibraten,<sup>16,§</sup> E. W. Hughes,<sup>12,††</sup> C. E. Hyde-Wright,<sup>10</sup> J. R. Johnson,<sup>18</sup>  
 D. Kawall,<sup>13</sup> A. Klein,<sup>10</sup> S. E. Kuhn,<sup>10</sup> M. Kuriki,<sup>15</sup> R. Lindgren,<sup>16</sup>  
 T. J. Liu,<sup>16</sup> R. M. Lombard-Nelsen,<sup>5</sup> J. Marroncle,<sup>5</sup> T. Maruyama,<sup>12</sup>  
 X. K. Maruyama,<sup>9</sup> J. McCarthy,<sup>16</sup> W. Meyer,<sup>12,\*\*</sup> Z.-E. Meziani,<sup>13,14</sup>  
 R. Minehart,<sup>16</sup> J. Mitchell,<sup>4</sup> J. Morgenstern,<sup>5</sup> G. G. Petratos,<sup>12,†</sup>  
 R. Pitthan,<sup>12</sup> D. Pocanic,<sup>16</sup> C. Prescott,<sup>12</sup> R. Prepost,<sup>18</sup> P. Raines,<sup>11</sup>  
 B. A. Raue,<sup>10,†</sup> D. Reyna,<sup>1</sup> A. Rijllart,<sup>12,††</sup> Y. Roblin,<sup>3</sup> L. S. Rochester,<sup>12</sup>  
 S. E. Rock,<sup>1</sup> O. A. Rondon,<sup>16</sup> I. Sick,<sup>2</sup> L. C. Smith,<sup>16</sup> T. B. Smith,<sup>8</sup>  
 M. Spengos,<sup>1,11</sup> F. Staley,<sup>5</sup> P. Steiner,<sup>2</sup> S. St.Lorant,<sup>12</sup> L. M. Stuart,<sup>12</sup>  
 F. Suekane,<sup>15</sup> Z. M. Szalata,<sup>1</sup> H. Tang,<sup>12</sup> Y. Terrien,<sup>5</sup> T. Usher,<sup>12</sup> D. Walz,<sup>12</sup>  
 F. Wesselmann,<sup>10</sup> J. L. White,<sup>1,12</sup> K. Witte,<sup>12</sup> C. C. Young,<sup>12</sup>  
 B. Youngman,<sup>12</sup> H. Yuta,<sup>15</sup> G. Zapalac,<sup>18</sup> B. Zihlmann,<sup>2</sup> D. Zimmermann<sup>16</sup>

*(E143 Collaboration)*

<sup>1</sup>The American University, Washington, D.C. 20016

<sup>2</sup>Institut für Physik der Universität Basel, CH-4056 Basel, Switzerland

<sup>3</sup>LPC IN2P3/CNRS, University Blaise Pascal, F-63170 Aubiere Cedex,  
France

<sup>4</sup>TJNAF, Newport News, Virginia 23606

<sup>5</sup>DAPNIA-Service de Physique Nucleaire Centre d'Etudes de Saclay,  
F-91191 Gif/Yvette, France

<sup>6</sup>Lawrence Livermore National Laboratory, Livermore, California 94550

<sup>7</sup>University of Massachusetts, Amherst, Massachusetts 01003

<sup>8</sup>University of Michigan, Ann Arbor, Michigan 48109

---

\*Work supported by Department of Energy contract DE-AC03-76SF00515.

<sup>9</sup>Naval Postgraduate School, Monterey, California 93943

<sup>10</sup>Old Dominion University, Norfolk, Virginia 23529

<sup>11</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104

<sup>12</sup>Stanford Linear Accelerator Center, Stanford, California 94309

<sup>13</sup>Stanford University, Stanford, California 94305

<sup>14</sup>Temple University, Philadelphia, Pennsylvania 19122

<sup>15</sup>Tohoku University, Sendai 980, Japan

<sup>16</sup>University of Virginia, Charlottesville, Virginia 22901

<sup>17</sup>The College of William and Mary, Williamsburg, Virginia 23187

<sup>18</sup>University of Wisconsin, Madison, Wisconsin 53706

### Abstract

We have measured the proton and deuteron spin structure functions  $g_1^p$  and  $g_1^d$  in the region of the nucleon resonances for  $W^2 < 5 \text{ GeV}^2$  and  $Q^2 \approx 0.5$  and  $Q^2 \approx 1.2 \text{ GeV}^2$  by inelastically scattering 9.7 GeV polarized electrons off polarized  $^{15}\text{NH}_3$  and  $^{15}\text{ND}_3$  targets. We observe significant structure in  $g_1^p$  in the resonance region. We have used the present results, together with the deep-inelastic data at higher  $W^2$ , to extract  $\Gamma(Q^2) \equiv \int_0^1 g_1(x, Q^2) dx$ .

This is the first information on the low- $Q^2$  evolution of  $\Gamma$  toward the Gerasimov-Drell-Hearn limit at  $Q^2 = 0$ .

Submitted to *Physical Review Letters*

The nucleon spin structure functions  $g_1$  and  $g_2$  have been and continue to be intensively studied via deep-inelastic lepton scattering[1, 2, 3, 4, 5, 6]; however, they remain largely undetermined in the resonance region, where the squared invariant mass of the final state  $W^2$  is less than 4 GeV<sup>2</sup>. Here,  $g_1$  and  $g_2$  can be used to investigate the helicity structure of the resonance transition amplitudes. States of definite spin and parity are more easily understood in terms of the virtual photon asymmetries[7]

$$\begin{aligned} A_1(x, Q^2) &= \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}} = \frac{1}{F_1} \left[ g_1 - \frac{Q^2}{\nu^2} g_2 \right] \quad \text{and} \\ A_2(x, Q^2) &= \frac{\sigma_{LT}}{\sigma_T} = \frac{\sqrt{Q^2}}{\nu F_1} [g_1 + g_2], \end{aligned} \quad (1)$$

in which  $Q^2$  is the squared four-momentum transfer,  $\nu$  is the energy transfer,  $x = Q^2/2M\nu$ ,  $M$  is the nucleon mass, and the structure functions  $F_1$ ,  $g_1$  and  $g_2$  depend on both  $x$  (or  $W^2$ ) and  $Q^2$ . The cross sections  $\sigma_{1/2}$  and  $\sigma_{3/2}$  measure the strength of virtual transverse photon absorption leading to final-state spin projections of  $\frac{1}{2}$  and  $\frac{3}{2}$ . The cross sections  $\sigma_L$ ,  $\sigma_T \equiv (\sigma_{1/2} + \sigma_{3/2})/2$ , and  $\sigma_{LT}$  measure longitudinal, transverse and interference photon absorptions. Positivity limits require that  $|A_1| \leq 1$  and  $|A_2| \leq \sqrt{R(x, Q^2)}$ , in which  $R \equiv \sigma_L/\sigma_T$ . The excitation of the  $\Delta(1232)$  resonance (spin- $\frac{3}{2}$ ) allows for both  $\frac{1}{2}$  and  $\frac{3}{2}$  spin projections, and at low  $Q^2$  is expected to be primarily a magnetic dipole transition, for which  $\sigma_{3/2}/\sigma_{1/2} = 3$  and  $A_1 = -\frac{1}{2}$ . On the other hand, the  $S_{11}(1535)$  resonance has no spin- $\frac{3}{2}$  projection, so  $A_1$  should be unity. The observed values of  $A_1$  in the resonance region are a combination of asymmetries for individual resonances and for the nonresonant background. The goals of the present measurements are to gain a better understanding of the resonances and to determine their influence on the deep-inelastic results, both in terms of radiative corrections and the evolution of  $\Gamma(Q^2) \equiv \int_0^1 g_1(x, Q^2) dx$ . Although the resonant contribution to  $\Gamma$  is insignificant at high  $Q^2$ , it dominates the integral below about  $Q^2 = 0.5$  GeV<sup>2</sup>. In fact, the limits set by the Gerasimov-Drell-Hearn (GDH) sum rule[8] indicate that  $\Gamma(Q^2)$  should change sign in the region  $0 < Q^2 < 1$  GeV<sup>2</sup> and approach zero as  $Q^2 \rightarrow 0$ .

The data for the present analysis are part of the E143 data set[5, 6] taken with a 9.7 GeV polarized electron beam (average polarization 85%) and cryogenic <sup>15</sup>NH<sub>3</sub> and <sup>15</sup>ND<sub>3</sub> targets (average polarizations of 65% and

25%, respectively). The data taken with the two spectrometers at  $4.5^\circ$  and  $7^\circ$  corresponded to  $Q^2 \approx 0.5$  and  $1.2 \text{ GeV}^2$  in the resonance region.

Since these data were taken with longitudinal target polarization only, the determination of  $g_1$  or  $A_1$  requires additional assumptions about either  $g_2$  or  $A_2$ . We extract  $g_1$  rather than  $A_1$  because  $g_1$  is significantly less affected by our lack of knowledge of  $g_2$  or  $A_2$ . We set  $A_2 = 0$  (corresponding to  $g_1 = -g_2$ ) in our analysis. This is motivated by the fact that  $|A_2| \leq \sqrt{R}$  and existing data[9] indicate that  $R$  is small in the resonance region ( $R = 0.06 \pm 0.02$  for  $1 < Q^2 < 8 \text{ GeV}^2$  and  $W^2 < 3 \text{ GeV}^2$ ). We have explored the sensitivity to  $A_2$  by considering the alternate possibilities  $g_2 = 0$  and  $g_2 = g_2^{\text{WW}} = -g_1 + \int_x^1 g_1(x')/x' dx'$ , the Wandzura-Wilczek[10] twist-two form. Maximum deviations in  $g_1$  from the  $A_2 = 0$  case define the systematic errors due to uncertainty in  $A_2$ . Even if  $A_2$  were as large as 0.3, the extracted values of  $g_1$  would shift by less than 0.014, which is small compared to the statistical errors on each point.

We have extracted  $g_1$  from the absolute cross section differences for electron and nucleon spins aligned along the beam axis either parallel ( $\uparrow\uparrow$ ) or antiparallel ( $\uparrow\downarrow$ ) to each other[7]:

$$\frac{1}{\sigma_M} \frac{d\sigma^{\uparrow\downarrow(\uparrow\uparrow)}}{d\Omega dE'} = \frac{F_2}{\nu} + \frac{2}{M} \tan^2\left(\frac{\theta}{2}\right) F_1 + (-) \frac{2}{M\nu} \tan^2\left(\frac{\theta}{2}\right) \times \left[ (E + E' \cos \theta + Q^2/\nu) g_1 - \sqrt{Q^2} F_1 A_2 \right] \quad (2)$$

in which  $\sigma_M$  is the Mott scattering cross section,  $E$  ( $E'$ ) is the initial (final) electron energy,  $\theta$  is the laboratory scattering angle,  $\nu = E - E'$ , and  $F_1$  and  $F_2$  are the unpolarized structure functions.

This method requires good knowledge of spectrometer acceptances, the number density of polarizable protons or deuterons in the target, and detector efficiencies. Alternatively, one could extract  $g_1$  from the count rate asymmetry as in Ref. [5, 6]; however, the dilution factor (the fraction of scatterings coming from a polarizable nucleon in the target) is needed in this case, which is more difficult to obtain reliably in the resonance region. Nevertheless, when we tried both methods, we found that they agreed to within a fraction of the statistical errors on each point (typically better than 3%).

As the first step to determine the absolute cross section differences, we calculated the raw count differences per incident charge  $N^{\uparrow\downarrow}/q^{\uparrow\downarrow} - N^{\uparrow\uparrow}/q^{\uparrow\uparrow}$ , with each term corrected for dead-time in the trigger electronics. For each

electron,  $W^2$  was calculated using the momentum and scattering angle determined from tracking. The data were binned in  $W^2$ , normalized by the product of target and beam polarizations and corrected for absolute spectrometer efficiency. Each detector's efficiency was determined by making a strict cut to select good electron events without using one of the detectors and checking how often that detector registered the electron. The absolute spectrometer efficiency is the product of all of the individual detector efficiencies (no evidence for correlations between them was found)[11].

Fully corrected cross section differences  $d\Delta\sigma/dE'd\Omega$  were obtained with the help of a Monte Carlo simulation. This simulation was used to normalize the raw data for target density and spectrometer acceptance. The normalized data were then corrected for radiative and resolution effects by an additive term determined from the Monte Carlo routine. Small corrections for polarized  $^{15}\text{N}$  and  $^{14}\text{N}$  in the target (and  $^1\text{H}$  in the case of the deuteron) were applied as in Ref. [5].

The Monte Carlo code simulated all relevant aspects of the experiment and was able to predict total count rates and count rate differences based on a set of input tables of cross sections and asymmetries. Monte Carlo events were generated by picking the scattering point in the target, the scattering angle, and the final energy at random, properly weighted by tabulated cross sections. External radiation and multiple scattering were calculated depending on the target material traversed by each simulated electron. Parameter and geometry files were used to determine the trajectory of each simulated electron through the target and spectrometer magnetic fields and the detector elements. The Monte Carlo data were then analyzed in the same manner as the real data and corrected for target thickness and incident charge.

The unpolarized cross sections were calculated from a fit by Stuart *et al.*[12]. The polarized cross section and asymmetry tables were calculated from a combination of resonant and non-resonant contributions. The non-resonant part came from a parametrization of all existing deep inelastic data (Fit III of Ref. [6]), which was extrapolated into the resonance region. The contribution from the nucleon resonances was modeled by dividing the resonance region into three parts and adjusting the asymmetries for the dominant resonances in each part using three independent fit parameters.

The full model was used to calculate cross sections and asymmetries both in lowest order of QED (Born approximation) and with full radiative corrections following the prescription by Shumeiko *et al.*[13]. We used the peaking

approximation after convincing ourselves that the full code produced negligible differences. The radiated cross sections and asymmetries were tabulated as input for the Monte Carlo code, while the Born cross sections and asymmetries were compared with the Monte Carlo output to determine the normalization factor and the additive correction for the raw data.

Although the resonance parametrization is simplistic, it describes the data well, and provides radiative and resolution corrections with sufficient precision compared to the statistical accuracy of our data. The following additional tests showed that the Monte Carlo code and its input parameters were reasonable: Without any normalization factors, the generated unpolarized counts versus  $W^2$  agree with the data to better than 2.2% (3.4%) in the  $4.5^\circ$  ( $7^\circ$ ) spectrometer. The shape as a function of  $W^2$  is in even better agreement in the region of the resonances. This provides confidence that the acceptance and resolution of the spectrometer are properly modeled. In addition, the measured and simulated count-rate differences which were integrated over the (quasi-) elastic region ( $W^2 < 1 \text{ GeV}^2$ ) where model uncertainties are minimal because of reasonably good knowledge of elastic form factors and radiative corrections, agree well with each other. The overall  $\chi^2$  is 3.85 for 4 degrees of freedom corresponding to p( $4.5^\circ$ ), p( $7^\circ$ ), d( $4.5^\circ$ ) and d( $7^\circ$ ).

Fig. 1 shows  $g_1$  obtained from cross section differences for proton and deuteron (per nucleon) measured with the two spectrometers as a function of  $W^2$ . The full length of the error bars corresponds to the statistical and systematic uncertainties added in quadrature. The cross bars indicate statistical errors alone, which dominate the total errors. Plotted as triangles are the data of Baum *et al.*[14] taken at similar kinematics and converted to  $g_1$  for comparison by assuming  $A_2 = 0$ . Within errors, the two measurements agree well. Both data sets show a negative contribution in the region of the  $\Delta(1232)$  resonance at  $W^2 \approx 1.5 \text{ GeV}^2$ , and a strongly positive contribution just above  $W^2 = 2 \text{ GeV}^2$  where the  $S_{11}$  and  $D_{13}$  resonances are important. This peak is less pronounced for the deuteron. The solid lines show the Monte Carlo simulation, which agrees well with the data. The dashed curve is a calculation using the code AO by Burkert and Li[15]. One of the reasons this code does not accurately reproduce the data is that it does not include decay channels other than single- $\pi$  emission.

Fig. 2 shows the integrals  $\Gamma(Q^2)$  for proton and neutron. The integral was evaluated for the average  $Q^2$  of the resonance region ( $M^2 < W^2 < 4 \text{ GeV}^2$ )

by summing our results directly over the resonance region (where  $Q^2$  does not vary much) and by adding to it a contribution from smaller  $x$  (larger  $W^2$ ) at the same fixed  $Q^2$  taken from Fit III to the world's deep-inelastic  $g_1$  data[6]. The neutron integrals were derived assuming a 5% D-state probability for the deuteron. The statistical errors assigned to Fit III[6] at given values of  $x$  and  $Q^2$  corresponded to the kinematically closest E143 data points at 9.7 or 16 GeV, which dominated the fit in this region. Systematic errors were calculated using the systematic uncertainties for the measured  $g_1$  in the resonance region added linearly to the systematic errors for the deep-inelastic region, which are highly correlated with each other. Extrapolation errors for the region below the last measured datum at  $x = 0.03$  were taken to be as large as the value that Fit III yields for  $x < 0.03$ .

Although several models for the  $Q^2$  evolution of  $\Gamma(Q^2)$  exist[16, 17, 18, 19, 20], we show here only two representative ones, together with the evolution[21] of the world's deep-inelastic data due to the changing coupling constant  $\alpha_S$ . Although the GDH sum rule is strictly valid only at  $Q^2 = 0$  where  $\Gamma(Q^2)$  vanishes, it can be used to predict the slope of  $\Gamma(Q^2)$  for small  $Q^2$ . The solid line at low  $Q^2$  shows  $\Gamma = -\kappa^2 Q^2 / 8M^2$  in which  $\kappa$  is the anomalous magnetic moment of either the proton or neutron. Burkert and Ioffe[19] consider the contributions from the resonances using the code AO, and the nonresonant contributions using a simple higher-twist-type form fitted to the deep-inelastic data. Their model is constrained to fit both the GDH and the deep-inelastic limits, and it describes the data quite well. Soffer and Teryaev[20] assume that the integral over  $g_1 + g_2$  varies smoothly from high  $Q^2$  where  $g_2 \approx 0$  down to  $Q^2 = 0$ . Using their simple prediction for this integral and subtracting the contribution from  $g_2$  alone using the Burkhardt-Cottingham sum rule[22], gives the dashed curve in Fig. 2b, which also agrees quite well with our data.

The present spin structure function data in the region of the nucleon resonances allow us to determine the integrals  $\Gamma(Q^2)$  for the first time at  $Q^2$  below 2 GeV<sup>2</sup>. In contrast to the nearly flat behavior in the deep-inelastic region above  $Q^2 = 2$  GeV<sup>2</sup>,  $\Gamma$  varies rapidly below  $Q^2 = 2$  GeV<sup>2</sup>. Models that interpolate between the deep-inelastic and GDH limits describe the data quite well in this non-perturbative regime. More precise data over a larger range of  $Q^2$  are expected from several experiments at TJNAF in the future.

This work was supported by the Department of Energy; the National Science Foundation; the Schweizerische Nationalfonds; the Commonwealth

of Virginia; the Centre National de la Recherche Scientifique and the Commissariat a l'Energie Atomique (French groups); the Japanese Ministry of Education, Science, and Culture; and the Jeffress Memorial Trust (W&M).

## References

- [\*] Permanent address: Oliver Lodge Lab, University of Liverpool, Liverpool, U. K.
- [\*\*] Permanent address: University of Bonn, D-53113 Bonn, Germany.
- [§] Permanent address: FFIYM, P.O. Box 25, N-2007 Kjeller, Norway.
- [‡] Present address: Kent State University, Kent, Ohio 44242.
- [††] Permanent address: CERN, 1211 Geneva 23, Switzerland.
- [†] Permanent address: Florida International University, Miami, FL 33199.
- [‡‡] Present Address: California Institute of Technology, Pasadena, CA 91125
- [1] E80, M.J. Alguard, *et al.*, Phys. Rev. Lett. **37**, 1261 (1976); Phys. Rev. Lett. **41**, 70 (1978); E130, G. Baum, *et al.*, Phys. Rev. Lett. **51**, 1135 (1983);
- [2] EMC, J. Ashman *et al.*, Phys. Lett. **B206**, 364 (1988); Nucl. Phys. **B328**, 1 (1989).
- [3] E142, P. L. Anthony *et al.*, Phys. Rev. Lett. **71**, 959 (1993).
- [4] SMC, B. Adeva *et al.*, Phys. Lett. **B302**, 533 (1993); D. Adams *et al.*, Phys. Lett. **B329**, 399 (1994); **B336**, 125 (1994); **B357**, 248 (1995).
- [5] E143, K. Abe *et al.*, Phys. Rev. Lett. **74**, 346 (1995); Phys. Rev. Lett. **75**, 25 (1995); Phys. Rev. Lett. **76**, 587 (1996).
- [6] E143, K. Abe *et al.*, Phys. Lett. **B364**, 61 (1995).



- [7] R. D. Roberts, *The Structure of the Proton*, Cambridge Univ. Press (1990).
- [8] S. Gerasimov, *Sov. J. Nucl. Phys.* **2**, 430 (1966); S.D. Drell and A.C. Hearn, *Phys. Rev. Lett.* **16**, 908 (1966).
- [9] C. Keppel, PhD Thesis, American University, 1994; unpublished.
- [10] S. Wandzura and F. Wilczek, *Phys. Lett.* **B72**, 195 (1977).
- [11] P. Raines, PhD Thesis, University of Pennsylvania, 1996; unpublished.
- [12] L.M. Stuart, *et al.*, SLAC-PUB-6305 (1996).
- [13] T. V. Kukhto and N. M. Shumeiko, *Nucl. Phys.* **B219**, 412 (1983); I. V. Akusevich and N. M. Shumeiko, *J. Phys.* **G20**, 513 (1994).
- [14] G. Baum *et al.*, *Phys. Rev. Lett.* **45**, 2000 (1980).
- [15] V. Burkert and Z.-J. Li, *Phys. Rev. D* **47**, 46 (1993).
- [16] V. Bernard, N. Kaiser and U.-G. Meissner, *Phys. Rev. D* **48**, 3062 (1993).
- [17] X. Ji and P. Unrau, *Phys. Lett.* **B333**, 228 (1994).
- [18] Z.-P. Li and Zh. Li, *Phys. Rev. D* **50**, 3119 (1994).
- [19] V.D. Burkert and B.L. Ioffe, *Phys. Lett.* **B296** 223, (1992); CEBAF Preprint PR-93-034.
- [20] J. Soffer and O.V. Teryaev, *Phys. Rev. D* **51**, 25 (1995).
- [21] S.A. Larin and J.A.M. Vermaseren, *Phys. Lett.* **B259**, 4261 (1992).
- [22] H. Burkhardt and W. N. Cottingham, *Ann. Phys.* **56**, 453 (1970).

Table 1: Integrals  $\Gamma$  of the structure functions  $g_1$  for the proton (p), deuteron (d) and neutron (n). The measured sum  $\Gamma_{res}$  for the resonance region ( $W^2 < 4 \text{ GeV}^2$ ) is listed separately from the totals  $\Gamma_{tot}$ , which includes the deep-inelastic region as given by fits to the world's data.

$Q^2$ (GeV <sup>2</sup> )		$\Gamma_{res} (\pm\text{stat.}\pm\text{syst.})$	$\Gamma_{tot}(\pm\text{stat.}\pm\text{syst.})$
0.5	p	$0.026\pm 0.008\pm 0.008$	$0.049 \pm 0.008 \pm 0.013$
0.5	d	$0.000\pm 0.013\pm 0.008$	$0.003 \pm 0.013 \pm 0.010$
0.5	n		$-0.043 \pm 0.029 \pm 0.025$
1.2	p	$0.040\pm 0.003\pm 0.004$	$0.100 \pm 0.005 \pm 0.012$
1.2	d	$0.026\pm 0.006\pm 0.004$	$0.043 \pm 0.008 \pm 0.007$
1.2	n		$-0.006 \pm 0.019 \pm 0.020$

Table 2: Fully corrected cross section differences  $d(\sigma^{\uparrow\downarrow} - \sigma^{\uparrow\uparrow})/dE'd\Omega \equiv \Delta\sigma$  (nb/GeV-sr) and structure functions  $g_1$  for the proton in the resonance region. Listed are the data for the  $4.5^\circ$  and  $7.0^\circ$  spectrometers. The values of  $Q^2$  and  $W^2$  (GeV $^2$ ) are given at bin centers. The additive correction  $\Delta\sigma_R$  includes both radiative effects and resolution corrections.

	$W^2$	$Q^2$	$\Delta\sigma$	$\Delta\sigma_R$	$g_1$
$4.5^\circ$	1.31	0.55	$-74 \pm 133$	-714	$-0.013 \pm 0.023 \pm 0.023$
	1.69	0.54	$-289 \pm 116$	-244	$-0.070 \pm 0.028 \pm 0.006$
	2.06	0.53	$627 \pm 106$	282	$0.197 \pm 0.033 \pm 0.039$
	2.44	0.52	$828 \pm 109$	229	$0.320 \pm 0.042 \pm 0.046$
	2.81	0.50	$245 \pm 108$	-120	$0.113 \pm 0.050 \pm 0.025$
	3.19	0.49	$120 \pm 80$	-166	$0.064 \pm 0.043 \pm 0.034$
	3.56	0.48	$75 \pm 78$	-59	$0.046 \pm 0.048 \pm 0.015$
	3.94	0.47	$-2 \pm 79$	-29	$-0.001 \pm 0.055 \pm 0.010$
	4.31	0.46	$158 \pm 68$	-7	$0.123 \pm 0.053 \pm 0.015$
	4.69	0.45	$124 \pm 66$	-10	$0.106 \pm 0.057 \pm 0.010$
$7^\circ$	1.19	1.28	$-4 \pm 5$	-19	$-0.003 \pm 0.003 \pm 0.003$
	1.56	1.26	$-8 \pm 16$	-49	$-0.007 \pm 0.014 \pm 0.005$
	1.94	1.23	$43 \pm 12$	4	$0.043 \pm 0.012 \pm 0.015$
	2.31	1.20	$162 \pm 16$	45	$0.193 \pm 0.020 \pm 0.031$
	2.69	1.18	$121 \pm 16$	3	$0.165 \pm 0.022 \pm 0.016$
	3.06	1.15	$95 \pm 15$	-12	$0.147 \pm 0.024 \pm 0.019$
	3.44	1.12	$57 \pm 15$	1	$0.098 \pm 0.026 \pm 0.009$
	3.81	1.09	$70 \pm 15$	2	$0.134 \pm 0.029 \pm 0.012$
	4.19	1.07	$110 \pm 15$	5	$0.233 \pm 0.031 \pm 0.025$
	4.56	1.04	$83 \pm 14$	4	$0.193 \pm 0.033 \pm 0.020$
4.94	1.01	$63 \pm 14$	3	$0.158 \pm 0.036 \pm 0.022$	

Table 3: Same as Table 2, but for the deuteron.

	$W^2$	$Q^2$	$\Delta\sigma$	$\Delta\sigma_R$	$g_1$
4.5°	1.31	0.55	-195±207	-397	-0.034±0.036±0.021
	1.69	0.54	-190±185	-211	-0.046±0.045±0.015
	2.06	0.53	401±163	80	0.126±0.051±0.016
	2.44	0.52	127±165	43	0.049±0.064±0.007
	2.81	0.50	-41±145	-34	-0.019±0.067±0.005
	3.19	0.49	166±128	-32	0.089±0.069±0.013
	3.56	0.48	36±120	-17	0.022±0.074±0.004
	3.94	0.47	-8±112	-14	-0.006±0.078±0.002
	4.31	0.46	-27±101	-9	-0.021±0.078±0.003
	4.69	0.45	67±97	-9	0.058±0.084±0.007
7°	1.19	1.28	79±24	18	0.053±0.016±0.010
	1.56	1.26	1±21	-30	0.001±0.018±0.007
	1.94	1.23	30±19	1	0.030±0.019±0.007
	2.31	1.20	26±22	11	0.031±0.026±0.005
	2.69	1.18	28±22	0	0.038±0.030±0.004
	3.06	1.15	72±22	0	0.112±0.034±0.013
	3.44	1.12	19±22	3	0.034±0.038±0.004
	3.81	1.09	27±21	2	0.053±0.041±0.006
	4.19	1.07	42±21	2	0.089±0.044±0.010
	4.56	1.04	34±21	1	0.080±0.048±0.008
4.94	1.01	50±20	1	0.126±0.051±0.015	




Fig. 1. Measurements of  $g_1$  as a function of  $W^2$  for the proton at (a)  $4.5^\circ$  and (b)  $7^\circ$ ; and for the deuteron at (c)  $4.5^\circ$  and (d)  $7^\circ$ . The present data (circles) are plotted together with the data of Baum *et al.* (triangles), our Monte Carlo simulation (solid line), and the model AO of Burkert and Li[15] (dashed line). The full error bars correspond to statistical and systematic errors added in quadrature, whereas the cross bars indicate statistical errors only.




Fig. 2. Integrals of  $g_1$  at several fixed values of  $Q^2$  for (a) the neutron and (b) the proton. The present data (circles) are plotted together with data from CERN[4] (triangles), E143 deep-inelastic[5] (squares), and E142[3] (inverted triangle). The curves correspond to the evolution[21] of the deep-inelastic results due to changing  $\alpha_s$  (solid line), the predictions of Burkert and Ioffe[19] (dotted line), the model of Soffer[20] (dashed line), and the GDH approach to  $Q^2 = 0$  (solid line). Errors are indicated as in Fig. 1.