

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Measurements of Higgs boson properties in the diphoton decay channel with  $36 \text{ fb}^{-1}$  of pp collision data at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector

### Permalink

<https://escholarship.org/uc/item/3b61b813>

### Journal

Physical Review D, 98(5)

### ISSN

2470-0010

### Authors

Aaboud, M  
Aad, G  
Abbott, B  
et al.

### Publication Date

2018-09-18

### DOI

10.1103/PhysRevD.98.052005

Peer reviewed

# Measurements of Higgs boson properties in the diphoton decay channel with $36\text{ fb}^{-1}$ of $pp$ collision data at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS detector

M. Aaboud *et al.*<sup>\*</sup>  
(ATLAS Collaboration)



(Received 14 February 2018; published 18 September 2018)

Properties of the Higgs boson are measured in the two-photon final state using  $36.1\text{ fb}^{-1}$  of proton–proton collision data recorded at  $\sqrt{s} = 13\text{ TeV}$  by the ATLAS experiment at the Large Hadron Collider. Cross-section measurements for the production of a Higgs boson through gluon–gluon fusion, vector-boson fusion, and in association with a vector boson or a top-quark pair are reported. The signal strength, defined as the ratio of the observed to the expected signal yield, is measured for each of these production processes as well as inclusively. The global signal strength measurement of  $0.99 \pm 0.14$  improves on the precision of the ATLAS measurement at  $\sqrt{s} = 7$  and  $8\text{ TeV}$  by a factor of two. Measurements of gluon–gluon fusion and vector-boson fusion productions yield signal strengths compatible with the Standard Model prediction. Measurements of simplified template cross sections, designed to quantify the different Higgs boson production processes in specific regions of phase space, are reported. The cross section for the production of the Higgs boson decaying to two isolated photons in a fiducial region closely matching the experimental selection of the photons is measured to be  $55 \pm 10\text{ fb}$ , which is in good agreement with the Standard Model prediction of  $64 \pm 2\text{ fb}$ . Furthermore, cross sections in fiducial regions enriched in Higgs boson production in vector-boson fusion or in association with large missing transverse momentum, leptons or top-quark pairs are reported. Differential and double-differential measurements are performed for several variables related to the diphoton kinematics as well as the kinematics and multiplicity of the jets produced in association with a Higgs boson. These differential cross sections are sensitive to higher order QCD corrections and properties of the Higgs boson, such as its spin and  $CP$  quantum numbers. No significant deviations from a wide array of Standard Model predictions are observed. Finally, the strength and tensor structure of the Higgs boson interactions are investigated using an effective Lagrangian, which introduces additional  $CP$ -even and  $CP$ -odd interactions. No significant new physics contributions are observed.

DOI: [10.1103/PhysRevD.98.052005](https://doi.org/10.1103/PhysRevD.98.052005)

## I. INTRODUCTION

In July 2012, the ATLAS [1] and CMS [2] experiments announced the discovery of a Higgs boson [3,4] using proton–proton collisions collected at center-of-mass energies  $\sqrt{s} = 7\text{ TeV}$  and  $8\text{ TeV}$  at the CERN Large Hadron Collider (LHC). Subsequent measurements of its properties were found to be consistent with those expected for the Standard Model (SM) Higgs boson [5] with a mass  $m_H = 125.09 \pm 0.21(\text{stat}) \pm 0.11(\text{syst})\text{ GeV}$  [6].

Following the modifications of the LHC to provide proton–proton collisions at a center-of-mass energy of  $\sqrt{s} = 13\text{ TeV}$ , the Higgs sector can be probed more deeply:

the data set collected in 2015 and 2016 allows inclusive Higgs boson measurements to be repeated with about two times better precision than to those done at  $\sqrt{s} = 7$  and  $8\text{ TeV}$  with the Run 1 data set. The increased center-of-mass energy results in much larger cross sections for events at high partonic center-of-mass energy. This implies improved sensitivity to a variety of interesting physics processes, such as Higgs bosons produced at high transverse momentum or Higgs bosons produced in association with a top–antitop quark pair. The Higgs boson decay into two photons ( $H \rightarrow \gamma\gamma$ ) is a particularly attractive way to study the properties of the Higgs boson and to search for deviations from the Standard Model predictions due to beyond-Standard Model (BSM) processes. Despite the small branching ratio,  $(2.27 \pm 0.07) \times 10^{-3}$  for  $m_H = 125.09\text{ GeV}$  [7], a reasonably large signal yield can be obtained thanks to the high photon reconstruction and identification efficiency at the ATLAS experiment. Furthermore, due to the excellent photon energy resolution of the ATLAS calorimeter, the signal manifests itself as a

<sup>\*</sup>Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.

narrow peak in the diphoton invariant mass ( $m_{\gamma\gamma}$ ) spectrum on top of a smoothly falling background, and the Higgs boson signal yield can be measured using an appropriate fit to the  $m_{\gamma\gamma}$  distribution of the selected events.

In this paper, the results of measurements of the Higgs boson properties in the diphoton decay channel are presented using  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data collected at  $\sqrt{s} = 13 \text{ TeV}$  by the ATLAS detector in 2015 and 2016. All the measurements are performed under the assumption that the Higgs boson mass is  $125.09 \text{ GeV}$ , and are compared to Standard Model predictions. Three types of measurements are presented in this paper and are summarized in the remainder of this section: (i) measurements of the total Higgs boson production-mode cross sections and “signal strengths”; (ii) cross sections using the SM production modes as “templates” in simplified fiducial regions; and (iii) measurements of integrated or differential cross sections in fiducial phase-space regions closely matched to the experimental selection.

The rest of this paper is organized as follows. Section II provides a brief description of the ATLAS detector, and Sec. III describes the selected data set. The generation of simulated event samples is described in Sec. IV. Section V gives an overview of the event reconstruction and selection, and Sec. VI explains the signal and background modeling used in the measurement. The sources of systematic uncertainties are detailed in Sec. VII. Section VIII describes the measurement of the total Higgs boson production-mode cross sections, signal strengths, and simplified template cross sections (STXS). Similarly, Sec. IX describes the measurement of the fiducial and differential cross sections. Section X concludes with a brief summary of the main findings.

### A. Higgs boson production-mode cross sections and signal strengths

In this paper, cross sections times branching ratio of the Higgs to two photons  $B(H \rightarrow \gamma\gamma)$  are measured for inclusive Higgs boson production, as well as for several individual production processes: gluon–gluon fusion (ggH), vector-boson fusion (VBF), Higgs boson production in association with a vector boson (VH), and production of a Higgs boson in association with a top–antitop quark pair ( $t\bar{t}H$ ) or a single top quark ( $t$ -channel and W-associated, respectively denoted as  $tHq$  and  $tHW$ , or in their sum as “ $tH$ ”). In the SM, gluon–gluon fusion is the dominant production mechanism at the LHC, contributing to about 87% of the total cross section at  $\sqrt{s} = 13 \text{ TeV}$  [7]. Vector-boson fusion and associated production with either a vector boson, with a top–antitop quark pair or a bottom–antibottom quark pair correspond to 6.8%, 4.0%, 0.9%, and 0.9%, respectively, of the total Higgs boson production cross section.

The data are divided into 31 categories based on the reconstructed event properties to maximize the sensitivity

to different production modes and the different regions of the simplified template cross sections, which are further described in Sec. IB. The categories are defined using the expected properties of the different production mechanisms: 10 categories aimed to measure gluon–gluon fusion properties, 4 categories to measure vector-boson fusion, 8 categories that target associated production with vector bosons with different final states, and 9 categories that target associated production with a top–antitop quark pair or a single top-quark. The definition of each category was optimized using simulated events and a full summary of the categories can be found in Sec. VIII. In the sequence of the classification, priority is given to categories aimed at selecting signal events from processes with smaller cross sections.

In order to probe the production mechanisms independently of the  $H \rightarrow \gamma\gamma$  branching ratio, ratios of the different production-mode cross sections normalized to gluon–gluon fusion are also reported, with their full experimental correlations. In addition, measurements of the signal strength  $\mu$ , which is the ratio of the measured cross section to the SM prediction, are given for the different production processes as well as for the inclusive production. Finally, coupling-strength modifiers, which are scale factors of the tree-level Higgs boson couplings to the different particles or of the effective Higgs boson couplings to photons and gluons from loop-induced processes, are reported.

### B. Simplified template cross sections

The measurements of cross sections separated by the production mode as presented in the previous section are extended to measurements in specific regions of phase space using the framework of the “simplified template cross sections” introduced in Refs. [7,8]. These are reported as cross section times  $B(H \rightarrow \gamma\gamma)$  for a Higgs boson absolute rapidity<sup>1</sup>  $|y_H|$  less than 2.5 and with further particle-level requirements. The different production modes are separated in a theoretically motivated way using the SM modes ggH, VBF, VH and top-quark-associated production modes as “templates.” The fiducial regions are defined in a “simplified” way using the measured kinematics and topology of the final state, defined by the Higgs boson, the hadronic jets and the vector bosons or top quarks in the event, to avoid large model-dependent extrapolations. The Higgs

<sup>1</sup>The ATLAS experiment uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the center of the detector and the  $z$ -axis along the beam pipe. The  $x$ -axis points from the IP to the center of the LHC ring, and the  $y$ -axis points upward. Cylindrical coordinates  $(r, \phi)$  are used in the transverse plane,  $\phi$  being the azimuthal angle around the  $z$ -axis. The pseudorapidity is defined in terms of the polar angle  $\theta$  as  $\eta = -\ln \tan(\theta/2)$ . When dealing with massive particles, the rapidity  $y = 1/2 \ln[(E + p_z)/(E - p_z)]$  is used, where  $E$  is the energy and  $p_z$  is the  $z$ -component of the momentum. Angular separation is expressed in terms of  $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ .

boson is treated as a stable final-state particle, which allows an easy combination with other decay channels. Similarly, vector bosons or top quarks are treated as stable particles, but the cases of leptonic and hadronic decays of the vector boson are distinguished.

In this paper a merged version of the so-called “stage-1” simplified template cross-section measurements are investigated. These measurements provide more information for theoretical reinterpretation compared to the signal strength measurements used in Run 1 and are defined to reduce the theoretical uncertainties typically folded into the signal strength results. In the full stage-1 proposal, template cross sections would be measured in 31 regions of phase space for  $|y_H| < 2.5$ , where the latter requirement reflects the experimental acceptance. The experimental categories used in this study (the same as those used for the signal strength measurements) have been optimized to provide the maximum sensitivity to such regions [7,8].

Since the current data set is not large enough to probe all of the stage-1 cross sections with sufficiently small statistical uncertainties, regions with poor sensitivity or with large anticorrelations are merged together into ten regions: Six regions probe gluon-fusion Higgs boson production with zero, one, and two jets associated with them. Two regions probe VBF Higgs boson production and Higgs boson production associated with vector bosons that decay hadronically. A dedicated cross section is measured for Higgs boson production associated with vector bosons that decay via leptonic modes. The final cross section measures top-associated ( $t\bar{t}H$  and  $tH$ ) Higgs-boson production. To retain sensitivity to beyond the Standard Model Higgs boson production, the  $\geq 1$  jet,  $p_T^H > 200$  GeV gluon–gluon fusion and  $p_T^j > 200$  GeV VBF +  $VH$  regions are not merged with other regions. Here  $p_T^H$  and  $p_T^j$  denote the Higgs boson and leading jet transverse momenta, respectively, where the leading jet is the highest transverse momentum jet in a given event. However, due to their large anti-correlation, only the cross section for the summed yield of these two regions is quoted here, and thus a total of nine kinematic regions are reported. The experimental sensitivity to the difference in the yields of these two regions is expected to be small, and the corresponding result is treated as a nuisance parameter rather than a measurement.

Table I summarizes the ten probes merged stage-1 cross sections and details which of the full 31 stage-1 cross sections were merged (middle and last column). A detailed description of the full 31 cross section proposal can be found in Appendix A.

### C. Fiducial integrated and differential cross sections

Fiducial integrated and differential cross sections have previously been measured at  $\sqrt{s} = 8$  TeV in the  $H \rightarrow \gamma\gamma$  decay channel by both the ATLAS [9] and the CMS [10]

Collaborations. In this paper, fiducial cross sections are determined in a variety of phase-space regions sensitive to inclusive Higgs boson production and to explicit Higgs boson production mechanisms. The measurement of these cross sections provides an alternative way to study the properties of the Higgs boson and to search for physics beyond the Standard Model. For each fiducial region of an integrated cross-section measurement or bin of a differential distribution, the  $H \rightarrow \gamma\gamma$  signal is extracted using a fit to the corresponding diphoton invariant mass spectrum. The cross sections are determined by correcting these yields for experimental inefficiencies and resolution effects, and by taking into account the integrated luminosity of the data. No attempt is made to separate individual production modes in favor of presenting fiducial regions enriched with a given production mode.

The inclusive fiducial region is defined at the particle level by two photons, not originating from the decay of a hadron, that have absolute pseudorapidity  $|\eta| < 2.37$ , excluding the region  $1.37 < |\eta| < 1.52$ ,<sup>2</sup> with the leading (subleading) photon transverse momentum greater than 35% (25%) of  $m_{\gamma\gamma}$ . The two photons are required to be isolated from hadronic activity by imposing that the summed transverse momentum of charged stable particles (with a mean lifetime that satisfies  $c\tau > 10$  mm) with  $p_T > 1$  GeV, within a cone of  $\Delta R = 0.2$  centered on the photon direction, be less than 5% of the photon transverse momentum. This selection is applied to all the presented fiducial integrated and differential cross section results and the isolation criterion was tuned to mimic the detector level selection. One additional cross section and three cross-section limits are reported in smaller fiducial regions sensitive to specific Higgs boson production mechanisms:

- (i) a VBF-enhanced region with two jets with large invariant mass and rapidity separation,
- (ii) a region of events containing at least one charged lepton,<sup>3</sup>
- (iii) a region of events with large missing transverse momentum,
- (iv) and a region of events with a topology matching the presence of a top–antitop quark pair.

The fiducial cross section for different jet multiplicities are reported and compared to several predictions. Eleven fiducial differential cross sections are reported, for events belonging to the inclusive fiducial region as a function of the following observables:

- (i)  $p_T^{\gamma\gamma}$  and  $|y_{\gamma\gamma}|$ , the transverse momentum and rapidity of the diphoton system,
- (ii)  $p_T^j$  and  $|y_{j_1}|$ , the transverse momentum and rapidity of the leading jet,

<sup>2</sup>This pseudorapidity interval corresponds to the transition region between the barrel and endcap sections of the ATLAS electromagnetic calorimeter, see Sec. II.

<sup>3</sup>In this paper reconstructed charged leptons denote electrons and muons.

TABLE I. The particle-level kinematic regions of the stage-1 simplified template cross sections, along with the intermediate set of regions used for the measurements presented in this paper. All regions require  $|y_H| < 2.5$ . Jets are defined using the anti- $k_t$  algorithm with radius parameter  $R = 0.4$  and are required to have  $p_T > 30$  GeV. The leading jet and Higgs boson transverse momenta are denoted by  $p_T^j$  and  $p_T^H$ , respectively. The transverse momentum of the Higgs boson and the leading and subleading jet is denoted as  $p_T^{Hjj}$  with the subleading jet being the second highest momentum jet in a given event. Events are considered “VBF-like” if they contain at least two jets with an invariant mass of  $m_{jj} > 400$  GeV and a rapidity separation between the two jets of  $|\Delta y_{jj}| > 2.8$ . Events are considered “VH-like” if they contain at least two jets with an invariant mass of  $60 \text{ GeV} < m_{jj} < 120 \text{ GeV}$ . All  $qq' \rightarrow Hqq'$  VBF and VH events (with the vector boson  $V$  decaying hadronically) which are neither VBF nor VH-like are part of the “Rest” selection. For the  $p_T^H > 200$  GeV gluon–gluon fusion and  $p_T^j > 200$  GeV VBF + VH regions, only the sum of the corresponding cross sections is reported while the difference of the two is profiled in the fit. In total, the cross sections for nine kinematic regions are measured. The small contributions from  $b\bar{b}H$  are merged with  $ggH$ . The process  $gg \rightarrow ZH$  refers only to box and loop processes dominated by top and bottom quarks (see Sec. IV for more details).

Process	Measurement region	Particle-level stage-1 region
$ggH + gg \rightarrow Z(\rightarrow qq)H$	0-jet	0-jet
	1-jet, $p_T^H < 60$ GeV	1-jet, $p_T^H < 60$ GeV
	1-jet, $60 \leq p_T^H < 120$ GeV	1-jet, $60 \leq p_T^H < 120$ GeV
	1-jet, $120 \leq p_T^H < 200$ GeV	1-jet, $120 \leq p_T^H < 200$ GeV
	$\geq 1$ -jet, $p_T^H > 200$ GeV	1-jet, $p_T^H > 200$ GeV
	$\geq 2$ -jet, $p_T^H < 200$ GeV or VBF-like	$\geq 2$ -jet, $p_T^H > 200$ GeV
$qq' \rightarrow Hqq'$ (VBF + VH)	$p_T^j < 200$ GeV	$\geq 2$ -jet, $p_T^H < 60$ GeV
	$p_T^j > 200$ GeV	$\geq 2$ -jet, $60 \leq p_T^H < 120$ GeV
VH (leptonic decays)	VH leptonic	$\geq 2$ -jet, $120 \leq p_T^H < 200$ GeV
		VBF-like, $p_T^{Hjj} < 25$ GeV
		VBF-like, $p_T^{Hjj} \geq 25$ GeV
		$p_T^j < 200$ GeV, VBF-like, $p_T^{Hjj} < 25$ GeV
		$p_T^j < 200$ GeV, VBF-like, $p_T^{Hjj} \geq 25$ GeV
		$p_T^j < 200$ GeV, VH-like
		$p_T^j < 200$ GeV, Rest
		$p_T^j > 200$ GeV
		$q\bar{q} \rightarrow ZH$ , $p_T^Z < 150$ GeV
		$q\bar{q} \rightarrow ZH$ , $150 < p_T^Z < 250$ GeV, 0-jet
		$q\bar{q} \rightarrow ZH$ , $150 < p_T^Z < 250$ GeV, $\geq 1$ -jet
		$q\bar{q} \rightarrow ZH$ , $p_T^Z > 250$ GeV
		$q\bar{q} \rightarrow WH$ , $p_T^W < 150$ GeV
		$q\bar{q} \rightarrow WH$ , $150 < p_T^W < 250$ GeV, 0-jet
		$q\bar{q} \rightarrow WH$ , $150 < p_T^W < 250$ GeV, $\geq 1$ -jet
		$q\bar{q} \rightarrow WH$ , $p_T^W > 250$ GeV
		$gg \rightarrow ZH$ , $p_T^Z < 150$ GeV
		$gg \rightarrow ZH$ , $p_T^Z > 150$ GeV, 0-jet
		$gg \rightarrow ZH$ , $p_T^Z > 150$ GeV, $\geq 1$ -jet
		$t\bar{t}H$
Top-associated production	top	W-associated $tH$ ( $tHW$ )
		t-channel $tH$ ( $tHq$ )
$b\bar{b}H$	merged w/ $ggH$	$b\bar{b}H$

- (iii)  $p_T^{j_2}$  and  $|y_{j_2}|$ , the transverse momentum and rapidity of the subleading jet,
- (iv)  $|\cos \theta^*|$ , the cosine of the angle between the beam axis and the diphoton system in the Collins–Soper frame [11],
- (v)  $\Delta\phi_{jj}$  and  $|\Delta y_{jj}|$ , the difference in azimuthal angle and in rapidity between the leading and subleading jets,
- (vi)  $|\Delta\phi_{\gamma\gamma,jj}|$ , the difference in azimuthal angle between the dijet system formed by the leading and subleading jets and the diphoton system,

- (vii) and  $m_{jj}$ , the invariant mass of the leading and subleading jets.

Seven additional variables are reported in Appendix C. Inclusive Higgs boson production is dominated by gluon–gluon fusion, for which the transverse momentum of the Higgs boson is largely balanced by the emission of soft gluons and quarks. Measuring  $p_T^{\gamma\gamma}$  probes the perturbative QCD modeling of this production mechanism which is mildly sensitive to the bottom and charm quark Yukawa couplings of the Higgs boson [12]. The distribution at high



transverse momentum is sensitive to new heavy particles coupling to the Higgs boson and to the top quark Yukawa coupling. The rapidity distribution of the Higgs boson is also sensitive to the modeling of the gluon–gluon fusion production mechanism, as well as to the parton distribution functions (PDFs) of the colliding protons. The transverse momentum and absolute rapidity of the leading and subleading jets probe the perturbative QCD modeling and are sensitive to the relative contributions of the different Higgs production mechanisms. The angular variables  $|\cos\theta^*|$  and  $\Delta\phi_{jj}$  are sensitive to the spin and  $CP$  quantum numbers of the Higgs boson. The dijet rapidity separation  $|\Delta y_{jj}|$ , the dijet mass  $m_{jj}$  and the azimuthal difference between the dijet and diphoton system  $|\Delta\phi_{\gamma\gamma,jj}|$  are sensitive to the VBF production mechanism. All fiducial differential cross sections are reported with their full statistical and experimental correlations and are compared to several predictions.

The strength and tensor structure of the Higgs boson interactions are investigated using an effective Lagrangian, which introduces additional  $CP$ -even and  $CP$ -odd interactions that can lead to deviations in the kinematic properties and event rates of the Higgs boson and of the associated jets from those in the Standard Model. This is done by a simultaneous fit to five differential cross sections, which are sensitive to the Wilson coefficients of four dimension-six  $CP$ -even or  $CP$ -odd operators of the strongly interacting Light Higgs formulation [13]. A similar analysis was carried out at  $\sqrt{s} = 8$  TeV by the ATLAS Collaboration [14].

## II. ATLAS DETECTOR

The ATLAS detector [1] covers almost the entire solid angle about the proton–proton interaction point. It consists of an inner tracking detector, electromagnetic and hadronic calorimeters, and a muon spectrometer.

Charged-particle tracks and interaction vertices are reconstructed using information from the inner detector (ID). The ID consists of a silicon pixel detector (including the insertable B-layer [15] installed before the start of Run 2), of a silicon microstrip detector, and of a transition radiation tracker (TRT). The ID is immersed in a 2 T axial magnetic field provided by a thin superconducting solenoid. The silicon detectors provide precision tracking over the pseudorapidity interval  $|\eta| < 2.5$ , while the TRT offers additional tracking and substantial discrimination between electrons and charged hadrons for  $|\eta| < 2.0$ .

The solenoid is surrounded by electromagnetic (EM) and hadronic sampling calorimeters allowing energy measurements of photons, electrons and hadronic jets and discrimination between the different particle types. The EM calorimeter is a lead/liquid-argon (LAr) sampling calorimeter. It consists of a barrel section, covering the pseudorapidity region  $|\eta| < 1.475$ , and of two endcap sections, covering  $1.375 < |\eta| < 3.2$ . The EM calorimeter is divided

in three layers, longitudinally in depth, for  $|\eta| < 2.5$ , and in two layers for  $2.5 < |\eta| < 3.2$ . In the regions  $|\eta| < 1.4$  and  $1.5 < |\eta| < 2.4$ , the first layer has a fine  $\eta$  segmentation to discriminate isolated photons from neutral hadrons decaying to pairs of close-by photons. It also allows, together with the information from the cluster barycenter in the second layer, where most of the energy is collected, a measurement of the shower direction without assumptions on the photon production point. In the range of  $|\eta| < 1.8$  a presampler layer allows corrections to be made for energy losses upstream of the calorimeter. The hadronic calorimeter reconstructs hadronic showers using steel absorbers and scintillator tiles ( $|\eta| < 1.7$ ), or either copper ( $1.5 < |\eta| < 3.2$ ) or copper–tungsten ( $3.1 < |\eta| < 4.9$ ) absorbers immersed in a LAr active medium.

A muon spectrometer surrounds the calorimeter. It comprises separate trigger ( $|\eta| < 2.4$ ) and precision tracking chambers ( $|\eta| < 2.7$ ) in the magnetic field provided by three large air-core toroids.

A two-level trigger system [16] was used during the  $\sqrt{s} = 13$  TeV data-taking period. Dedicated hardware implements the first-level (L1) trigger selection, using only a subset of the detector information and reducing the event rate to at most 100 kHz. Events satisfying the L1 requirements are processed by a high-level trigger executing, on a computer farm, algorithms similar to the offline reconstruction software, in order to reduce the event rate to approximately 1 kHz.

## III. DATA SET

Events were selected using a diphoton trigger requiring the presence in the EM calorimeter of two clusters of energy depositions with transverse energy above 35 GeV and 25 GeV for the leading (highest transverse energy) and subleading (second highest transverse energy) cluster. In the high-level trigger the shape of the energy deposition of both clusters was required to be loosely consistent with that expected from an electromagnetic shower initiated by a photon. The diphoton trigger has an efficiency greater than 99% for events that satisfy the final event selection described in Sec. V.

After the application of data quality requirements, the data set amounts to an integrated luminosity of  $36.1 \text{ fb}^{-1}$ , of which  $3.2 \text{ fb}^{-1}$  were collected in 2015 and  $32.9 \text{ fb}^{-1}$  were collected in 2016. The mean number of proton–proton interactions per bunch crossing is 14 in the 2015 data set and 25 in the 2016 data set.

## IV. EVENT SIMULATION

Signal samples were generated for the main Higgs boson production modes using Monte Carlo event generators as described in the following. The mass and width of the Higgs boson were set in the simulation to  $m_H = 125 \text{ GeV}$  and  $\Gamma_H = 4.07 \text{ MeV}$  [17], respectively. The samples are normalized with the latest available theoretical calculations

of the corresponding SM production cross sections, as summarized in Ref. [7] and detailed below. The normalization of all Higgs boson samples also accounts for the  $H \rightarrow \gamma\gamma$  branching ratio of 0.227% calculated with HDECAY [18,19] and PROPHECY4F [20–22].

Higgs boson production via  $ggH$  is simulated at next-to-next-to-leading-order (NNLO) accuracy in QCD using the POWHEG NNLOPS program [23], with the PDF4LHC15 PDF set [24]. The simulation achieves NNLO accuracy for arbitrary inclusive  $gg \rightarrow H$  observables by reweighting the Higgs boson rapidity spectrum in HJ-MiNLO [25] to that of HNNLO [26]. The transverse momentum spectrum of the Higgs boson obtained with this sample was found to be compatible with the fixed-order HNNLO calculation [26] and the HRES 2.3 calculation [27,28] performing resummation at next-to-next-to-leading-logarithm accuracy matched to a NNLO fixed-order calculation (NNLL + NNLO). The HRES prediction includes the effects of the top and bottom quark masses up to NLO precision in QCD and uses dynamical renormalization ( $\mu_R$ ) and factorization ( $\mu_F$ ) scales,  $\mu_F = \mu_R = 0.5 \sqrt{m_H^2 + (p_T^H)^2}$ . The parton-level events produced by the POWHEG NNLOPS program are passed to PYTHIA8 [29] to provide parton showering, hadronization and underlying event, using the AZNLO set of parameters that are tuned to data [30]. The sample is normalized such that it reproduces the total cross section predicted by a next-to-next-to-next-to-leading-order ( $N^3$ LO) QCD calculation with NLO electroweak corrections applied [31–34].

Higgs boson production via VBF is generated at NLO accuracy in QCD using the POWHEG-BOX program [35–38] with the PDF4LHC15 PDF set. The parton-level events are passed to PYTHIA8 to provide parton showering, hadronization and the underlying event, using the AZNLO parameter set. The VBF sample is normalized with an approximate-NNLO QCD cross section with NLO electroweak corrections applied [39–41].

Higgs boson production via  $VH$  is generated at NLO accuracy in QCD through  $q\bar{q}/qg$ -initiated production, denoted as  $q\bar{q}' \rightarrow VH$ , and through  $gg \rightarrow ZH$  production using POWHEG-BOX [42] with the PDF4LHC15 PDF set. Higgs boson production through  $gg \rightarrow ZH$  has two distinct sources: a contribution with two additional partons,  $gg \rightarrow ZHq\bar{q}$ , and a contribution without any additional partons in the final state, including box and loop processes dominated by top and bottom quarks. In the following, the  $gg \rightarrow ZH$  notation refers only to this latter contribution. PYTHIA8 is used for parton showering, hadronization and the underlying event using the AZNLO parameter set. The samples are normalized with cross sections calculated at NNLO in QCD and NLO electroweak corrections for  $q\bar{q}' \rightarrow VH$  and at NLO and next-to-leading-logarithm accuracy in QCD for  $gg \rightarrow ZH$  [43–45].

Higgs boson production via  $t\bar{t}H$  is generated at NLO accuracy in QCD using MG5\_AMC@NLO [46] with the

NNPDF3.0 PDF set [47] and interfaced to PYTHIA8 to provide parton showering, hadronization and the underlying event, using the A14 parameter set [48]. The  $t\bar{t}H$  sample is normalized with a cross-section calculation accurate to NLO in QCD with NLO electroweak corrections applied [49–52].

Higgs boson production via  $b\bar{b}H$  is simulated using MG5\_AMC@NLO [53] interfaced to PYTHIA8 with the CT10 PDF set [54], and is normalized with the cross-section calculation obtained by matching, using the *Santander* scheme, the five-flavor scheme cross section accurate to NNLO in QCD with the four-flavor scheme cross section accurate to NLO in QCD [55–57]. The sample includes the effect of interference with the gluon–gluon fusion production mechanism.

Associated production of a Higgs boson with a single top-quark and a  $W$ -boson ( $tHW$ ) is generated at NLO accuracy, removing the overlap with the  $t\bar{t}H$  sample through a diagram regularization technique, using MG5\_AMC@NLO interfaced to HERWIG++ [58–60], with the HERWIG++ UEEE5 parameter set for the underlying event and the CT10 PDF set using the five-flavor scheme. Simulated Higgs boson events in association with a single top-quark, a  $b$ -quark and a light quark ( $tHq$ ) are produced at LO accuracy in QCD using MG5\_AMC@NLO interfaced to PYTHIA8 with the CT10 PDF set within the four-flavor scheme and using the A14 parameter set. The  $tHW$  and  $tHq$  samples are normalized with calculations accurate to NLO in QCD [61].

The generated Higgs boson events are passed through a GEANT4 [62] simulation of the ATLAS detector [63] and reconstructed with the same analysis software used for the data.

Background events from continuum  $\gamma\gamma$  production and  $V\gamma\gamma$  production are simulated using the SHERPA event generator [64], with the CT10 PDF set and the SHERPA default parameter set for the underlying-event activity. The corresponding matrix elements for  $\gamma\gamma$  and  $V\gamma\gamma$  are calculated at leading order (LO) in the strong coupling constant  $\alpha_s$  with the real emission of up to three or two additional partons, respectively, and are merged with the SHERPA parton shower [65] using the MEPS@LO prescription [66].

The very large sample size required for the modeling of the  $\gamma\gamma$  background processes is obtained through a fast parametric simulation of the ATLAS detector response [67]. For  $V\gamma\gamma$  events the same full detector simulation as for the signal samples is used.

Additional proton–proton interactions (pileup) are included in the simulation for all generated events such that the average number of interactions per bunch crossing reproduces that observed in the data. The inelastic proton–proton collisions were produced using PYTHIA8 with the A2 parameter set [68] that are tuned to data and the MSTW2008LO PDF set [69]. A summary of the used signal and background samples is shown in Table II.

TABLE II. Summary of the event generators and PDF sets used to model the signal and the main background processes. The SM cross sections  $\sigma$  for the Higgs production processes with  $m_H = 125.09$  GeV are also given separately for  $\sqrt{s} = 13$  TeV, together with the orders of the calculations corresponding to the quoted cross sections, which are used to normalize the samples, after multiplication by the Higgs boson branching ratio to diphotons, 0.227%. The following versions were used: PYTHIA8 version 8.212 (processes) and 8.186 (pile-up overlay); HERWIG++ version 2.7.1; POWHEG-BOX version 2; MG5\_AMC@NLO version 2.4.3; SHERPA version 2.2.1.

Process	Generator	Showering	PDF set	$\sigma$ [pb]	Order of calculation of $\sigma$
				$\sqrt{s} = 13$ TeV	
ggH	POWHEG NNLOPS	PYTHIA8	PDF4LHC15	48.52	N <sup>3</sup> LO(QCD) + NLO(EW)
VBF	POWHEG-BOX	PYTHIA8	PDF4LHC15	3.78	NNLO(QCD) + NLO(EW)
WH	POWHEG-BOX	PYTHIA8	PDF4LHC15	1.37	NNLO(QCD) + NLO(EW)
$q\bar{q}' \rightarrow ZH$	POWHEG-BOX	PYTHIA8	PDF4LHC15	0.76	NNLO(QCD) + NLO(EW)
$gg \rightarrow ZH$	POWHEG-BOX	PYTHIA8	PDF4LHC15	0.12	NLO + NLL(QCD)
$t\bar{t}H$	MG5_AMC@NLO	PYTHIA8	NNPDF3.0	0.51	NLO(QCD) + NLO(EW)
$b\bar{b}H$	MG5_AMC@NLO	PYTHIA8	CT10	0.49	5FS(NNLO) + 4FS(NLO)
$t$ -channel $tH$	MG5_AMC@NLO	PYTHIA8	CT10	0.07	4FS(LO)
$W$ -associated $tH$	MG5_AMC@NLO	HERWIG++	CT10	0.02	5FS(NLO)
$\gamma\gamma$	SHERPA	SHERPA	CT10		
$V\gamma\gamma$	SHERPA	SHERPA	CT10		

## V. EVENT RECONSTRUCTION AND SELECTION

### A. Photon reconstruction and identification

The reconstruction of photon candidates is seeded by energy clusters in the electromagnetic calorimeter with a size of  $\Delta\eta \times \Delta\phi = 0.075 \times 0.125$ , with transverse energy  $E_T$  greater than 2.5 GeV [70]. The reconstruction is designed to separate electron from photon candidates, and to classify the latter as *unconverted* or *converted* photon candidates. Converted photon candidates are associated with the conversion of photons into electron-positron pairs in the material upstream the electromagnetic calorimeter. Conversion vertex candidates are reconstructed from either two tracks consistent with originating from a photon conversion, or one track that does not have any hits in the innermost pixel layer. These tracks are required to induce transition radiation signals in the TRT consistent with the electron hypothesis, in order to suppress backgrounds from nonelectron tracks. Clusters without any matching track or conversion vertex are classified as unconverted photon candidates, while clusters with a matching conversion vertex are classified as converted photon candidates. In the simulation, the average reconstruction efficiency for photons with generated  $E_T$  above 20 GeV and generated pseudorapidity  $|\eta| < 2.37$  is 98%.

The energy from unconverted and converted photon candidates is measured from an electromagnetic cluster of size  $\Delta\eta \times \Delta\phi = 0.075 \times 0.175$  in the barrel region of the calorimeter, and  $\Delta\eta \times \Delta\phi = 0.125 \times 0.125$  in the calorimeter endcaps. The cluster size is chosen sufficiently large to optimize the collection of energy of the particles produced in the photon conversion. The cluster electromagnetic energy is corrected in four steps to obtain the calibrated energy of the photon candidate, using a combination of simulation-based and data-driven correction factors [71].

The simulation-based calibration procedure was re-optimized for the 13 TeV data. Its performance is found to be similar with that of Run 1 [71] in the full pseudorapidity range, and is improved in the barrel–endcap transition region, due to the use of information from additional scintillation detectors in this region [72]. The uniformity corrections and the intercalibration of the longitudinal calorimeter layers are unchanged compared to Run 1 [71], and the data-driven calibration factors used to set the absolute energy scale are determined from  $Z \rightarrow e^+e^-$  events in the full 2015 and 2016 data set. The energy response resolution is corrected in the simulation to match the resolution observed in data. This correction is derived simultaneously with the energy calibration factors using  $Z \rightarrow e^+e^-$  events by adjusting the electron energy resolution such that the width of the reconstructed  $Z$  boson peak in the simulation matches the width observed in data [72].

Photon candidates are required to satisfy a set of identification criteria to reduce the contamination from the background, primarily associated with neutral hadrons in jets decaying into photon pairs, based on the lateral and longitudinal shape of the electromagnetic shower in the calorimeter [73]. Photon candidates are required to deposit only a small fraction of their energy in the hadronic calorimeter, and to have a lateral shower shape consistent with that expected from a single electromagnetic shower. Two working points are used: a *loose* criterion, primarily used for triggering and preselection purposes, and a *tight* criterion. The tight selection requirements are tuned separately for unconverted and converted photon candidates. Corrections are applied to the electromagnetic shower shape variables of simulated photons, to account for small differences observed between data and simulation. The variation of the photon identification efficiency associated with the different reconstruction of converted photons in



the 2015 and 2016 data sets, due to the different TRT gas composition, has been studied with simulated samples and shown to be small. The efficiency of the tight identification criteria ranges from 84% to 94% (87% to 98%) for unconverted (converted) photons with transverse energy between 25 GeV and 200 GeV.

To reject the hadronic jet background, photon candidates are required to be isolated from any other activity in the calorimeter and the tracking detectors. The calorimeter isolation is computed as the sum of the transverse energies of positive-energy topological clusters [74] in the calorimeter within a cone of  $\Delta R = 0.2$  centered around the photon candidate. The transverse energy of the photon candidate is removed. The contributions of the underlying event and pileup are subtracted according to the method suggested in Ref. [75]. Candidates with a calorimeter isolation larger than 6.5% of the photon transverse energy are rejected. The track isolation is computed as the scalar sum of the transverse momenta of all tracks in a cone of  $\Delta R = 0.2$  with  $p_T > 1$  GeV which satisfy some loose track-quality criteria and originate from the diphoton primary vertex, i.e., the most likely production vertex of the diphoton pair (see Sec. VB). For converted photon candidates, the tracks associated with the conversion are removed. Candidates with a track isolation larger than 5% of the photon transverse energy are rejected.

### B. Event selection and selection of the diphoton primary vertex

Events are preselected by requiring at least two photon candidates with  $E_T > 25$  GeV and  $|\eta| < 2.37$  (excluding the transition region between the barrel and endcap calorimeters of  $1.37 < |\eta| < 1.52$ ) that fulfill the loose photon identification criteria [70]. The two photon candidates with the highest  $E_T$  are chosen as the diphoton candidate, and used to identify the *diphoton primary vertex* among all reconstructed vertices, using a neural-network algorithm based on track and primary vertex information, as well as the directions of the two photons measured in the calorimeter and inner detector [76]. The neural-network algorithm selects a diphoton vertex within 0.3 mm of the true  $H \rightarrow \gamma\gamma$  production vertex in 79% of simulated gluon-gluon fusion events. For the other Higgs production modes this fraction ranges from 84% to 97%, increasing with jet activity or the presence of charged leptons. The performance of the diphoton primary vertex neural-network algorithm is validated using  $Z \rightarrow e^+e^-$  events in data and simulation, by ignoring the tracks associated with the electron candidates and treating them as photon candidates. Sufficient agreement between the data and the simulation is found. The diphoton primary vertex is used to redefine the direction of the photon candidates, resulting in an improved diphoton invariant mass resolution. The invariant mass of the two photons is given by  $m_{\gamma\gamma} = \sqrt{2E_1E_2(1 - \cos\alpha)}$ , where  $E_1$  and  $E_2$  are the

energies of the leading and subleading photons and  $\alpha$  is the opening angle of the two photons with respect to the selected production vertex.

Following the identification of the diphoton primary vertex, the leading and subleading photon candidates in the diphoton candidate are respectively required to have  $E_T/m_{\gamma\gamma} > 0.35$  and 0.25, and to both satisfy the tight identification criteria as well as the calorimeter and track isolation requirements. Figure 1 compares the simulated per-event efficiency of the track- and calorimeter-based isolation requirement as a function of the number of primary vertex candidates with the per-event efficiency of the Run 1 algorithm described in Ref. [76], by using a MC sample of Higgs bosons produced by gluon-gluon fusion and decaying into two photons. The reoptimization of the thresholds applied to the transverse energy sum of the calorimeter energy deposits and to the transverse momentum scalar sum of the tracks in the isolation cone, as well as the reduction of the size of the isolation cone for the calorimeter-based isolation, greatly reduces the degradation of the efficiency as the number of reconstructed primary vertices increases in comparison to the Run 1 algorithm.

In total 332030 events are selected with diphoton candidates with invariant mass  $m_{\gamma\gamma}$  between 105 GeV and 160 GeV. The predicted signal efficiency, assuming the SM and including the acceptance of the kinematic selection, is 42% (with the acceptance of the kinematic selection being 52%).




FIG. 1. Efficiency for both photons to fulfill the isolation requirement as a function of the number of primary vertex candidates in each event, determined with a sample of simulated Higgs bosons with  $m_H = 125$  GeV, produced in gluon-gluon fusion and decaying into two photons. Events are required to satisfy the kinematic selection described in Sec. VB for the 8 TeV (violet squares) and 13 TeV (blue circles) simulated sample. The error bars show the statistical uncertainty of the generated samples. The Run 2 (Run 1) isolation requirement is based on the transverse energy deposited in the calorimeter in a  $\Delta R = 0.2$  ( $\Delta R = 0.4$ ) cone around the photon candidates. Both the Run 1 and Run 2 algorithms also use tracking information in a  $\Delta R = 0.2$  cone around the photon candidates.

### C. Reconstruction and selection of hadronic jets, $b$ -jets, leptons and missing transverse momentum

Jets are reconstructed using the anti- $k_t$  algorithm [77] with a radius parameter of 0.4 via the FASTJET package [78,79]. The inputs to the algorithm are three-dimensional topological clusters of energy deposits in the calorimeter cells [74]. Jets are corrected on an event-by-event basis for energy deposits originating from pileup [80], then calibrated using a combination of simulation-based and data-driven correction factors, which correct for different responses to electromagnetic and hadronic showers of the calorimeter and inactive regions of the calorimeter [81,82]. Jets are required to have  $p_T > 25$  GeV for  $|\eta| < 2.4$ . The jet selection is tightened to  $p_T > 30$  GeV within  $|\eta| < 4.4$  for most event reconstruction categories and the measurement of fiducial integrated and differential cross sections (with exceptions noted in Secs. VIII A and IX C). Jets that do not originate from the diphoton primary vertex are rejected, for  $|\eta| < 2.4$ , using the jet vertex tagging algorithm (JVT) [83], which combines tracking information into a multivariate likelihood. For jets with  $p_T < 60$  GeV and  $|\eta| < 2.4$  a *medium* working point is used, with an efficiency greater than 92% for nonpileup jets with  $p_T > 30$  GeV. The efficiency of the JVT algorithm is corrected in the simulation to match that observed in the data. Jets are discarded if they are within  $\Delta R = 0.4$  of an isolated photon candidate, or within  $\Delta R = 0.2$  of an isolated electron candidate.

Jets consistent with the decay of a  $b$ -hadron are identified using a multivariate discriminant, having as input information from track impact parameters and secondary vertices [84,85]. The chosen identification criterion has an efficiency of 70% for identifying jets originating from a  $b$ -quark. The efficiency is determined using a  $t\bar{t}$  control region, with rejection factors of about 12 and 380 for jets originating from  $c$ -quarks and light quarks, respectively. Data-driven correction factors are applied to the simulation such that the  $b$ -tagging efficiencies of jets originating from  $b$ -quarks,  $c$ -quarks and light quarks are consistent with the ones observed in the data.

The reconstruction and calibration of electron candidates proceeds similarly as for photon candidates. Electromagnetic calorimeter clusters with a matching track in the inner detector are reconstructed as electron candidates and calibrated using dedicated corrections from the simulation and from data control samples. Electron candidates are required to have  $p_T > 10$  GeV and  $|\eta| < 2.47$ , excluding the region  $1.37 < |\eta| < 1.52$ . Electrons must satisfy *medium* identification criteria [86] using a likelihood-based discriminant.

Muon candidates are primarily built from tracks reconstructed in the inner detector and the muon spectrometer, but are complemented by candidates reconstructed only in the muon spectrometer that are compatible with originating from the interaction point [87]. Muon candidates are required to have  $p_T > 10$  GeV and  $|\eta| < 2.7$ , and satisfy

*medium* identification criteria based on the number of hits in the silicon detectors, in the TRT and in the muon spectrometer. For the measurements of fiducial cross sections the electron and muon selections are tightened to  $p_T > 15$  GeV.

Lepton candidates are discarded if they are within  $\Delta R = 0.4$  of an isolated photon candidate or a jet. Isolation requirements are applied to all lepton candidates. Electron candidates are required to satisfy loose criteria for the calorimeter and track isolation, aimed at a combined efficiency of 99% independently of the candidate transverse momentum. Muon candidates are similarly required to satisfy loose criteria for the calorimeter and track isolation, in this case depending on the candidate transverse momentum, and aimed at a combined efficiency ranging from 95–97% at  $p_T = 10$ –60 GeV to 99% for  $p_T > 60$  GeV.

Tracks associated with both the electron and muon candidates are required to be consistent with originating from the diphoton primary vertex by requiring their longitudinal impact parameter  $z_0$  to satisfy  $|z_0 \sin \theta| < 0.5$  mm and their unsigned transverse impact parameter  $|d_0|$  relative to the beam axis to be respectively smaller than five or three times its uncertainty.

The lepton efficiency as well as energy/momentum scale and resolution are determined using the decays of  $Z$  bosons and  $J/\psi$  mesons in the full 2015 and 2016 data set using the methods described in Refs. [86,87]. Lepton efficiency correction factors are applied to the simulation to improve the agreement with the data.

The magnitude of the missing transverse momentum  $E_T^{\text{miss}}$  is measured from the negative vectorial sum of the transverse momenta of all photon, electron, and muon candidates and of all hadronic jets after accounting for overlaps between jets, photons, electrons, and muons, as well as an estimate of soft contributions based on tracks originating from the diphoton vertex which satisfy a set of quality criteria. A full description of this algorithm can be found in Refs. [88,89]. The  $E_T^{\text{miss}}$  significance is defined as  $E_T^{\text{miss}} / \sqrt{\sum E_T}$ , where  $\sum E_T$  is the sum of the transverse energies of all particles and jets used in the estimation of the missing transverse momentum in units of GeV.

## VI. SIGNAL AND BACKGROUND MODELING OF DIPHOTON MASS SPECTRUM

The Higgs boson signal yield is measured through an unbinned maximum-likelihood fit to the diphoton invariant mass spectrum in the range  $105 \text{ GeV} < m_{\gamma\gamma} < 160 \text{ GeV}$  for each event reconstruction category, fiducial region, or each bin of a fiducial differential cross section, as further discussed in Secs. VIII and IX. The mass range is chosen to be large enough to allow a reliable determination of the background from the data, and at the same time small enough to avoid large uncertainties from the choice of the background parametrization. The signal and background shapes are modeled as described below, and the

background model parameters are freely floated in the fit to the  $m_{\gamma\gamma}$  spectra.

### A. Signal model

The Higgs boson signal manifests itself as a narrow peak in the  $m_{\gamma\gamma}$  spectrum. The signal distribution is empirically modeled as a double-sided Crystal Ball function, consisting of a Gaussian central part and power-law tails on both sides. The Gaussian core of the

$$f_i^{\text{sig}}(m_{\gamma\gamma}; \Delta\mu_{\text{CB},i}, \sigma_{\text{CB},i}, \alpha_{\text{CB},i}^{\pm}, n_{\text{CB},i}^{\pm}) = \mathcal{N}_c \begin{cases} e^{-t^2/2} & -\alpha_{\text{CB},i}^- \leq t \leq \alpha_{\text{CB},i}^+ \\ \left(\frac{n_{\text{CB},i}^-}{|\alpha_{\text{CB},i}^-|}\right)^{n_{\text{CB},i}^-} e^{-|\alpha_{\text{CB},i}^-|^2/2} \left(\frac{n_{\text{CB},i}^-}{\alpha_{\text{CB},i}^-} - \alpha_{\text{CB},i}^- - t\right)^{-n_{\text{CB},i}^-} & t < -\alpha_{\text{CB},i}^- \\ \left(\frac{n_{\text{CB},i}^+}{|\alpha_{\text{CB},i}^+|}\right)^{n_{\text{CB},i}^+} e^{-|\alpha_{\text{CB},i}^+|^2/2} \left(\frac{n_{\text{CB},i}^+}{\alpha_{\text{CB},i}^+} - \alpha_{\text{CB},i}^+ - t\right)^{-n_{\text{CB},i}^+} & t > \alpha_{\text{CB},i}^+ \end{cases},$$

where  $t = (m_{\gamma\gamma} - m_H - \Delta\mu_{\text{CB},i})/\sigma_{\text{CB},i}$ , and  $\mathcal{N}_c$  is a normalization factor. The non-Gaussian parts are parametrized by  $\alpha_{\text{CB},i}^{\pm}$  and  $n_{\text{CB},i}^{\pm}$  separately for the low- (−) and high-mass (+) tails.

The parameters of the model that define the shape of the signal distribution are determined through fits to the simulated signal samples. The parametrization is derived separately for each reconstructed category or fiducial region of the integrated or differential cross-section measurement. Figure 2 shows an example for two categories with different mass resolution: the improved mass resolution in the central region of the detector (defined by requiring  $|\eta| \leq 0.95$  for both selected photons) with respect to the forward region (defined by requiring one photon with  $|\eta| \leq 0.95$  and one photon with  $0.95 < |\eta| < 2.37$ ) results in better discriminating power against the non-resonant background and in turn in a smaller statistical error of the extracted Higgs boson signal yield. The effective signal mass resolution of the two categories, defined as half the width containing 68% (90%) of the signal events, is 1.6 (2.7) GeV and 2.1 (3.8) GeV, respectively, and the mass resolution for all used categories can be found in Appendix E.

### B. Background composition and model

The diphoton invariant mass model for the background used to fit the data is determined from studies of the bias in the signal yield in signal + background fits to large control samples of data or simulated background events.

Continuum  $\gamma\gamma$  production is simulated with the SHERPA event generator as explained in Sec. IV, neglecting any interference effects with the  $H \rightarrow \gamma\gamma$  signal. The  $\gamma j$  and  $j j$  backgrounds are obtained by reweighting this sample using an  $m_{\gamma\gamma}$  dependent linear correction function obtained from the fraction of  $\gamma\gamma$  to  $\gamma j$  and  $\gamma\gamma$  to  $j j$  background events in data, respectively.

Crystal Ball function is parameterized by the peak position ( $m_H + \Delta\mu_{\text{CB}}$ ) and the width ( $\sigma_{\text{CB}}$ ). The non-Gaussian contributions to the mass resolution arise mostly from converted photons  $\gamma \rightarrow e^+e^-$  with at least one electron losing a significant fraction of its energy through bremsstrahlung in the inner detector material. The parametric form for a given reconstructed category or bin  $i$  of a fiducial cross section measurement, for a Higgs boson mass  $m_H$ , can be written as:

For very low rate categories targeting  $t\bar{t}H$  or  $tH$  events, in which the background simulation suffers from very large statistical uncertainties, various background-enriched control samples are directly obtained from the data by either reversing photon identification or isolation criteria, or by loosening or removing completely  $b$ -tagging identification requirements on the jets, and normalizing to the data in the  $m_{\gamma\gamma}$  sidebands of the events satisfying the nominal selection. For low rate categories targeting associated vector boson production, background control samples are obtained by summing the distributions from the main background




FIG. 2. The diphoton mass signal shapes of two gluon–gluon fusion categories that are later introduced in Sec. VIII A are shown: ggH 0J Fwd aims to select gluon–gluon fusion events with no additional jet and at least one photon in the pseudorapidity region  $|\eta| > 0.95$ ; ggH 0J Cen applies a similar selection, but requires both photons to have  $|\eta| \leq 0.95$  in order to have a better energy resolution. The simulated sample (labeled as MC) is compared to the fit model and contains simulated events from all Higgs boson production processes described in Sec. IV with  $m_H = 125$  GeV.




FIG. 3. The data-driven determination of (a) event yields and (b) event fractions for  $\gamma\gamma$ ,  $\gamma j$ , and  $jj$  events as a function of  $m_{\gamma\gamma}$  after the final selection outlined in Sec. V. The event fractions for two differential observables, (c)  $p_T^{\gamma\gamma}$  and (d)  $N_{\text{jets}}$  defined for jets with a  $p_T > 30$  GeV are shown as well. The shaded regions show the total uncertainty of the measured yield and fraction, and the error bars show the statistical uncertainties.

processes: those of  $\gamma\gamma$  and  $V\gamma\gamma$  events are obtained directly from the simulation, while the  $m_{\gamma\gamma}$  distributions of  $\gamma j$  and  $jj$  events are obtained from data control samples in which the nominal selection is applied except that at least one (for  $\gamma j$ ) or both (for  $jj$ ) of the two photon candidates fail to meet either the identification or isolation criteria. Except for the  $V\gamma\gamma$  component, which is normalized with its theoretical cross section, the other contributions are normalized according to their relative fractions determined in data as described in the following.

The measurement of the background fractions in data is performed for each category or fiducial region. The relative fractions of  $\gamma\gamma$ ,  $\gamma j$  and  $jj$  background events are determined using a double two-dimensional sideband method [90,91]. The nominal identification and isolation requirement are loosened for both photon candidates, and the data are split into 16 orthogonal regions defined by diphoton pairs in which one or both photons satisfy or fail to meet identification and/or isolation requirements. The region in which both photons satisfy the nominal identification and isolation requirements corresponds to the nominal selection of Sec. V, while the other 15 regions provide control regions,

whose  $\gamma\gamma$ ,  $\gamma j$ , and  $jj$  yields are related to those in the signal region via the efficiencies for photons and for hadronic jets to satisfy the photon identification and isolation requirements. The  $\gamma\gamma$ ,  $\gamma j$ , and  $jj$  yields in the signal region are thus obtained, together with the efficiencies for hadronic jets, by solving a linear system of equations using as inputs the observed yields in the 16 regions and the photon efficiencies predicted by the simulation. In the  $VH$  categories, a small extra contribution from  $V\gamma\gamma$  events with an electron originating from the decay of the vector boson  $V$  which is incorrectly reconstructed as a photon, is also estimated from the simulation and subtracted before applying the two-dimensional sideband method. The dominant systematic uncertainties in the measured background fractions are due to the definition of the background control regions. The yields and relative fractions of the  $\gamma\gamma$ ,  $\gamma j$ , and  $jj$  backgrounds are shown in Fig. 3 as a function of  $m_{\gamma\gamma}$  for the selected events. The fractions of these background sources in the inclusive diphoton sample are  $(78.7^{+1.8}_{-5.2})\%$ ,  $(18.6^{+4.2}_{-1.6})\%$ , and  $(2.6^{+0.5}_{-0.4})\%$ , respectively. The uncertainties in the measured background fractions are systematically dominated. These results are comparable to previous



results at  $\sqrt{s} = 7$  and 8 TeV [9,76]. In addition the purity is shown as a function of the  $p_T$  of the diphoton system, and the number of reconstructed jets with  $p_T > 30$  GeV.

The functional form used to model the background  $m_{\gamma\gamma}$  distribution in the fit to the data is chosen, in each region, to ensure a small bias in the extracted signal yield relative to its experimental precision, following the procedure described in Ref. [3]. The potential bias (*spurious signal*) is estimated as the maximum of the absolute value of the fitted signal yield, using a signal model with mass between 121 and 129 GeV, in fits to the background control regions described before.

The spurious signal is required, at 95% confidence level (CL), to be less than 10% of the expected SM signal yield or less than 20% of the expected statistical uncertainty in the SM signal yield. In the case when two or more functions satisfy those requirements, the background model with the least number of parameters is chosen.

Prior to the final fit to the data, the selected model is tested against a model from the same family of functions but with one more degree of freedom (d.o.f.) (for instance, the exponential of a second-order polynomial is tested against an exponential of a third-order polynomial) to check, using only events in the diphoton invariant mass sidebands (i.e., excluding the range  $121 \text{ GeV} < m_{\gamma\gamma} < 129 \text{ GeV}$ ), if the data favors a more complex model. A test statistic is built from the  $\chi^2$  values and number of d.o.f. of two binned fits to the data with the two background models. The expected distribution of the test statistic is built from pseudo-data assuming that the function with fewer d.o.f. is the true underlying model. The value of the test statistic obtained in the data is compared to such distribution, and the simpler model is rejected in favor of the more complex one if the  $p$ -value of such comparison is lower than 5%. The background distribution of all regions is found to be well modeled by at least one of the following functions: an exponential of a first- or second-order polynomial, a power law, or a third-order Bernstein polynomial.

### C. Statistical model

The data are interpreted following the statistical procedure summarized in Ref. [92] and described in detail in Ref. [93]. An extended likelihood function is built from the number of observed events and invariant diphoton mass values of the observed events using the analytic functions describing the distributions of  $m_{\gamma\gamma}$  in the range 105–160 GeV for the signal and the background.

The likelihood for a given reconstructed category, fiducial region, or differential bin  $i$  of the integrated or differential cross-section measurement is a marked Poisson probability distribution,

$$\mathcal{L}_i = \text{Pois}(n_i | N_i(\theta)) \cdot \prod_{j=1}^{n_i} f_i(m_{\gamma\gamma}^j, \theta) \cdot G(\theta),$$

where  $n_i$  ( $N_i$ ) is the observed (expected) number of selected candidates,  $f_i(m_{\gamma\gamma}^j)$  is the value of the probability density function (pdf) of the invariant mass distribution evaluated for each candidate  $j$ ,  $\theta$  are nuisance parameters and  $G(\theta)$  is a set of unit Gaussian constraints on a subset of the nuisance parameters, as described in the following. The likelihood for the measurements of the total Higgs boson production-mode cross sections and signal strengths is given by the product of the likelihood functions of each event reconstruction category. For the fiducial integrated and differential cross-section measurements the likelihood of all bins  $i$  in a spectrum is taken.

The number of expected candidates is the sum of the signal and background yields, denoted by  $N_i^{\text{sig}}$  and  $N_i^{\text{bkg}}$ , and the fitted spurious signal yield,  $N_i^{\text{spur}} \cdot \theta_i^{\text{spur}}$ ,

$$N_i = N_i^{\text{sig}} + N_i^{\text{bkg}} + N_i^{\text{spur}} \cdot \theta_i^{\text{spur}}.$$

In more detail, the invariant mass distribution for each category has signal and background components,

$$f_i(m_{\gamma\gamma}^j) = [(N_i^{\text{sig}} + N_i^{\text{spur}} \cdot \theta_i^{\text{spur}}) \cdot f_i^{\text{sig}}(m_{\gamma\gamma}^j, \theta_i^{\text{sig}}) + N_i^{\text{bkg}} \cdot f_i^{\text{bkg}}(m_{\gamma\gamma}^j, \theta_i^{\text{bkg}})] / N_i,$$

where  $\theta_i^{\text{sig}}$  and  $\theta_i^{\text{bkg}}$  are nuisance parameters associated with systematic uncertainties affecting the resolutions and positions (Sec. VII A) of the invariant mass distributions of the signal  $f_i^{\text{sig}}$  (further detailed in Sec. VI A) or the shape of the background  $f_i^{\text{bkg}}$  (as explained in Sec. VI B), respectively.

Systematic uncertainties are incorporated into the likelihood function by multiplying the relevant parameter of the statistical model by a factor

$$F_G(\sigma, \theta) = (1 + \sigma \cdot \theta)$$

in the case of a Gaussian pdf for the effect of an uncertainty of size  $\sigma$  or, for cases where a negative model parameter does not make physical sense (e.g., the uncertainty in the integrated luminosity), by a factor

$$F_{\text{LN}}(\sigma, \theta) = e^{\sqrt{\ln(1+\sigma^2)}\theta}$$

for a log-normal pdf. In both cases the corresponding component of the constraint product  $G(\theta)$  is a unit Gaussian centered at zero for the nuisance parameter  $\theta$ . The systematic uncertainties affecting the yield and mass resolution use the log-normal form while a Gaussian form is used for all others. When two uncertainties are considered fully correlated they share the same nuisance parameter. Systematic uncertainties with partial correlations are decomposed into their uncorrelated and fully correlated components before being assigned to nuisance parameters.

All measured Higgs boson signal yields are determined with the profile likelihood ratio test statistic

$$\Lambda(\nu) = -2 \ln \frac{\mathcal{L}(\nu, \hat{\theta}_\nu)}{\mathcal{L}(\hat{\nu}, \hat{\theta})}, \quad (1)$$

where  $\hat{\nu}$  and  $\hat{\theta}$  are the values of the parameter of interest (e.g., a signal strength or a simplified template cross section) and nuisance parameters that unconditionally maximize the likelihood while  $\hat{\theta}_\nu$  are the values of the nuisance parameters that maximize the likelihood on the condition that the parameter of interest is held fixed to a given value  $\nu$ . In the asymptotic approximation, which is valid for all the results presented here,  $\Lambda(\nu)$  may be interpreted as an increase in  $\chi^2$  from its minimum value [92] such that approximate confidence intervals are easily constructed. The total uncertainty in  $\nu$  is thus obtained from the  $\nu$  values such that  $\Lambda(\nu) = 1$  with all other parameters “profiled” (i.e., set to the values that maximize the likelihood for those values of  $\nu$ ). Theory uncertainties in the parameters of interest are found by fixing the nuisance parameters associated with experimental uncertainties and subtracting in quadrature the statistical uncertainty. The statistical uncertainty is similarly determined, by fixing all nuisance parameters to their best-fit values, except for those describing the background shape and normalization. The experimental uncertainty is found by subtracting in quadrature the theory and the statistical uncertainties from the total uncertainty.

#### D. Limit setting in the absence of a signal

In the absence of a significant signal yield in the measured production process categories or fiducial regions, upper limits on the corresponding signal strength or cross section are placed. For production-mode measurements, the limit is set by treating all other parameters of the fit as nuisance parameters. For the fiducial regions, each measurement is split into two orthogonal categories, one of which contains the events in the specified fiducial region and one that contains the events that are outside of it. The diphoton spectrum in both sets of events are simultaneously analyzed to extract the desired limit.

For category-based measurements the 95% CL upper limit on the parameter of interest  $\nu$  is determined using the CL<sub>s</sub> prescription [94]. For this, the agreement between data and the expected yield for the hypothesized value of the parameter of interest  $\nu$  is quantified by the test statistic,  $q_\nu$ , defined as

$$q_\nu = \begin{cases} \Lambda(\nu) & 0 < \hat{\nu} \leq \nu \\ 0 & \nu < \hat{\nu} \end{cases}, \quad (2)$$

where  $\hat{\nu} \geq 0$  is the fitted parameter of interest. The observed value of the test statistic,  $q_{\text{obs}}$ , is determined

from the ratio of the likelihood obtained by fixing the number of signal events to that predicted for a given value of the parameter of interest, to the likelihood normalized by allowing the number of signal events to float in the fit. The asymptotic behavior of Eq. (2) is well known [92]. For fiducial measurements the 95% CL upper limit are determined using a one-sided Gaussian interpretation of the observed cross section.

## VII. SYSTEMATIC UNCERTAINTIES

Several sources of systematic uncertainty are considered in this measurement. They can be grouped into three categories: (i) uncertainties associated with the parameterization of the signal and background when fitting the  $m_{\gamma\gamma}$  spectrum, (ii) experimental uncertainties arising either from the extraction of the signal in a given category or from migrations between categories, and (iii) theoretical and modeling uncertainties in each category, causing migrations between categories, or affect the fiducial acceptance.

The origin of the uncertainties and their treatment are discussed in detail below and summarized in Table III.<sup>4</sup>

The analysis based on event reconstruction categories and those of fiducial cross sections treat yield and migration uncertainties differently: whereas the former incorporate them directly into the likelihood function (cf. Sec. VIC), the latter incorporate them at a later stage as part of the correction factor (introduced in Sec. IX B) or the luminosity. Modeling uncertainties were also estimated with different approaches as discussed further in Secs. VII C and VII D. A summary of the impact of the uncertainties on the measurement is given in Secs. VIII B 2 and IX E 6.

#### A. Systematic uncertainties in the signal and background modeling from fitting the $m_{\gamma\gamma}$ spectrum

Systematic uncertainties associated with the signal and background parametrizations are treated in a similar way for all the measurements. These include systematic uncertainties in the photon energy scale and resolution, and the uncertainties due to the specific choice of background model.

The fit to the  $m_{\gamma\gamma}$  spectra is performed for a Higgs boson mass of  $m_H = 125.09 \pm 0.24$  GeV [6]. The uncertainties in the photon energy scale and resolution impact the signal model, as the photon energy scale shifts the position of the peak and the assumed energy resolution broadens or narrows the signal peak relative to its nominal width. Uncertainties in the photon energy scale are included as

<sup>4</sup>The breakdown of uncertainties differs from those used in the Run 1 measurement Ref. [76] as more updated recommendations for experimental and theory uncertainties are used.

TABLE III. Summary of the sources of systematic uncertainties for results based on event reconstruction categories or fiducial integrated and differential cross sections. The columns labels “Category Likelihood” and “Fiducial Likelihood” provide an overview about which terms are part of the Likelihood (✓) or incorporated at a later stage (⋯). Both sets of results incorporate uncertainties associated with the Higgs boson mass, photon energy scale and resolution, and uncertainties associated with the choice of the background function into the likelihood function, either using log normal ( $F_{\text{LN}}(\sigma_i, \theta_i)$ ) or Gaussian constraints ( $F_{\text{G}}(\sigma_i, \theta_i)$ ) with  $\sigma_i$  denoting the systematic uncertainty ( $i$  is the index to each of the unique nuisance parameters  $\theta$ ). When acting on  $N_{\text{S}}^{\text{tot}}$  the uncertainty value is the same for all processes, whereas the uncertainty has a different value for each signal process for the case denoted  $N_{\text{S}}^{\text{p}}$ . The number of nuisance parameters,  $N_{\text{NP}}$ , for the spurious signal uncertainty varies, e.g., for the category-based results 31 independent error sources are present and for the differential measurements one source per measured bin is included.

Systematic uncertainty source			$N_{\text{NP}}$	Constraint	Category likelihood	Fiducial likelihood
Theory	ggH QCD		9	$N_{\text{S}}^{\text{ggH}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	Missing higher orders (non-ggH)		6	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	$B(H \rightarrow \gamma\gamma)$		1	$N_{\text{S}}^{\text{tot}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	PDF		30	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	$\alpha_{\text{S}}$		1	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	UE/PS		5	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
Experimental	Yield	Heavy flavor content	1	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Luminosity	1	$N_{\text{S}}^{\text{tot}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Trigger	1	$N_{\text{S}}^{\text{tot}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Photon identification	1	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Photon isolation	2	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	Migration	Flavor tagging	14	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Jet	20	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Jet flavor composition	7	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Jet flavor response	7	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Electron	3	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Muon	11	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Missing transverse momentum	3	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Pileup	1	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
		Photon energy scale	40	$N_{\text{S}}^{\text{p}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	⋯
	Mass	ATLAS-CMS $m_H$	1	$\mu_{\text{CB}} F_{\text{G}}(\sigma_i, \theta_i)$	✓	✓
		Photon energy scale	40	$\mu_{\text{CB}} F_{\text{G}}(\sigma_i, \theta_i)$	✓	✓
		Photon energy resolution	9	$\sigma_{\text{CB}} F_{\text{LN}}(\sigma_i, \theta_i)$	✓	✓
	Background	Spurious signal	Varies	$N_{\text{spur},c} \theta_{\text{spur},c}$	✓	✓

nuisance parameters associated with Gaussian constraint terms in the likelihood functions. Uncertainties in the photon energy resolution are included as nuisance parameters, and are typically among the dominant sources of systematic uncertainty in the measurement. The systematic uncertainties in the photon energy resolution and scale follow those in Refs. [71,72]. The overall energy scale factors and their uncertainties have been determined using  $Z \rightarrow e^+e^-$  events collected during 2015 and 2016. Compared to Ref. [72], several systematic uncertainties were re-evaluated with the 13 TeV data, including uncertainties related to the observed LAr cell nonlinearity, the material simulation, the intercalibration of the first and second layer of the calorimeter, and the pedestal corrections. The typical impact of the photon energy scale

uncertainties is to shift the peak position by between  $\pm 0.21\%$  and  $\pm 0.36\%$  of the nominal peak position, whereas the typical impact of the photon energy resolution uncertainty is to change the width of the signal distribution by between  $\pm 6\%$  and  $\pm 13\%$  of the nominal width. The size of both uncertainties is dependent on the energy, rapidity and jet activity of the selected photon pair.

An additional uncertainty in the signal peak position is added as a nuisance parameter in the fit, reflecting the uncertainty in the measurement of the Higgs boson mass of 0.24 GeV [6]. The uncertainty in the Higgs boson mass is dominated by the statistical component, and the systematic component has contributions from both the ATLAS and the CMS muon momentum and electromagnetic energy scale

uncertainties. Therefore, the correlation between this uncertainty and the photon energy scale uncertainty in the measurements presented here is considered negligible. A variation of the signal mass by  $\pm 0.24$  GeV (without including this uncertainty in the fit) is found to impact the measured global signal strength or the diphoton fiducial cross section by less than  $\pm 0.1\%$ .

The uncertainty due to the choice of background function is taken to be the spurious signal yield obtained when fitting the  $m_{\gamma\gamma}$  spectrum reconstructed from background-only simulated samples (or signal-suppressed control regions in data), as discussed in Section VI.

### B. Experimental systematic uncertainties affecting the expected event yields

There are two categories of uncertainties: (1) those in the expected overall signal yield and (2) those that cause migrations of events between categories and bins, as well as into and out of the photon fiducial selection.

The sources of uncertainties in the expected signal yield consist of:

- (i) The luminosity delivered to the ATLAS experiment. The uncertainty in the combined 2015 + 2016 integrated luminosity is 3.2%. It is derived, following a methodology similar to that detailed in Ref. [95], from a calibration of the luminosity scale using  $x$ - $y$  beam-separation scans performed in August 2015 and May 2016.
- (ii) The efficiency of the diphoton trigger. Its uncertainty is estimated to be 0.4% by comparing the trigger efficiencies determined using a bootstrap method [16] in data and simulation.
- (iii) The photon identification efficiency. Its uncertainty is estimated to be 1.6% and is obtained by varying the efficiency scale factors within their uncertainties, derived from control samples of photons from radiative  $Z$  boson decays and from inclusive  $\gamma$  events, and of electrons from  $Z \rightarrow e^+e^-$  decays.
- (iv) The photon track isolation efficiency. Its uncertainty is estimated to be 0.8% and is derived from measurements of the efficiency correction factors using inclusive photon control samples.
- (v) The photon calorimeter isolation efficiency. Its uncertainty is estimated to be 0.1% and is obtained from the difference between applying and not applying corrections derived from inclusive photon events to the calorimeter isolation variable in the simulation.

Uncertainties which affect the calibration of photons, jets, and leptons cause migrations between categories and bins, as well as migrations into and out of the fiducial acceptance. These include:

- (i) The modeling of pileup in the simulation. The corresponding uncertainty is derived by varying

the average number of pileup events in the simulation by an amount consistent with data. The typical size ranges from 1.4% up to 5.6% depending on the category or fiducial cross section bin.

- (ii) Uncertainties in the photon energy scale and resolution. These uncertainties cause migrations into and out of the fiducial volume or between the event reconstruction categories and impact the expected number of events. The calibration of the absolute energy scale is derived using  $Z \rightarrow e^+e^-$  decays. The impact of the corresponding uncertainties is small, however, for all measurements, and ranges for instance between 0.2% for events with a low diphoton  $p_T$  up to 1.9% for events with a high diphoton  $p_T$ .
- (iii) Uncertainties in the jet energy calibration and the jet energy resolution. Uncertainties in the jet energy scale and resolution are estimated by varying the jet energies by an amount commensurate with the differences observed between 13 TeV data and simulation in the transverse momentum balance in dijet,  $\gamma$  + jet and  $Z$  + jet events [81,82,96]. The typical size of this uncertainty ranges from 2.8% to 15%.
- (iv) Uncertainties due to the efficiency of the jet vertex tagger. Such uncertainties are estimated by shifting the associated corrections applied to the simulation by an amount allowed by the data. For the measurement of the fiducial integrated and differential cross sections, uncertainties associated with the modeling of pileup jets in the simulation are estimated by recalculating the correction factor after removing 20% of pileup jets at random, which is commensurate with the observed differences in data and simulation for jets tagged as low-JVT (pileup) and high-JVT (hard scatter). Its typical size ranges from nil to 0.3%.
- (v) Uncertainties associated with the efficiency of the  $b$ -tagging algorithm. They have been estimated to be typically of the order of 3% and are determined using  $t\bar{t}$  events in 13 TeV data for jets containing the decay of a  $b$ -quark, using the method outlined in Ref. [97]. The corresponding uncertainties in the identification of jets originating from  $c$ -quarks, light quarks and gluons are taken directly from Run 1 studies [97], with additional uncertainties to cover the extrapolation to Run 2 conditions.
- (vi) Uncertainties in the electron [86] and muon [87] reconstruction, identification and isolation efficiencies. They have been obtained from dilepton decays of  $Z$  bosons and  $J/\psi$  mesons collected in Run 2, using a tag-and-probe technique. The typical size of these uncertainties is about 0.6% for electrons and about 0.5% for muons in the relevant categories or fiducial regions.



- (vii) Uncertainties in the electron [98] and muon [87] energy and momentum scale and resolution. They are determined from comparisons between the reconstructed invariant mass in data and simulation of dileptons from decays of  $Z$  bosons or  $J/\psi$  mesons. The impact is negligible for all measurements.
- (viii) Uncertainties associated with energy scales and resolutions of photons, jets and leptons are propagated to the  $E_T^{\text{miss}}$  uncertainty, together with the uncertainty in the contribution to  $E_T^{\text{miss}}$  from charged-particle tracks not associated with high- $p_T$  leptons, jets, or photon conversions [89]. This results in a typical migration uncertainties ranging from 4.0% to 4.8% for relevant categories or fiducial regions.

### C. Theoretical and modeling uncertainties for results based on event reconstruction categories

The overall theoretical cross-section uncertainties affect the signal strength measurements, which are ratios of the observed to predicted event yields, but not the cross-section measurements which do not rely on absolute predictions. Modeling uncertainties that alter the kinematic properties of the events, such as the Higgs boson transverse momentum or the jet multiplicity, have an impact on both types of measurements.

The theoretical modeling uncertainties in the per-category acceptance of each production process affect the measurement of production-mode cross sections and signal strengths. Uncertainties due to the choice of parton distribution functions and the value of  $\alpha_s$  are estimated using the PDF4LHC15 recommendations [24] with the nominal PDF4LHC\_nlo\_30\_as PDF set. Samples using the CT10 PDF set are reweighted to PDF4LHC15 to estimate these uncertainties. For the gluon–gluon fusion process, the total production-mode cross section has been calculated at N<sup>3</sup>LO precision in QCD and has an uncertainty of 3.9%, as determined by QCD-scale variations and including top, bottom, and charm quark mass effect uncertainties. However, the perturbative uncertainty becomes significantly larger in different kinematic regions, e.g. when requiring additional jets or high Higgs boson  $p_T$ . To take this effect into account nine uncertainty sources are included:

- (i) Four sources [7] account for uncertainties in the jet multiplicities due to missing higher-order corrections: two accounting for yield uncertainties (with uncertainties up to 8.9% in each STXS region) and two accounting for migrations between jet multiplicity bins (with uncertainties up to 18% in each STXS region), using the STWZ [99] and BLPTW [99–101] predictions as an input. For more details see Table 20 of Ref. [7].

- (ii) Three uncertainty sources parameterize modeling uncertainties in the Higgs boson  $p_T$ . The first two encapsulate the migration uncertainty between the intermediate and high  $p_T$  region with events with at least one jet. The third uncertainty parameterizes top-quark mass effects in the gluon–gluon fusion loop, where the difference between the LO and NLO predictions is taken as an uncertainty due to missing higher-order corrections. This introduces a negligible uncertainty at low Higgs boson  $p_T$  and a sizable uncertainty of the order of 30% at  $p_T > 500$  GeV.
- (iii) Two sources account for the uncertainty in the acceptance of gluon–gluon fusion events in the VBF categories, due to missing QCD higher-orders in the calculation. Such uncertainties are estimated by variations of the renormalization and factorization scales in MCFM [102] by a factor of two around the nominal scale of  $\mu_r = \mu_f = m_H$ . The two sources account for the uncertainty in the overall normalization of  $H + 2\text{jet}$  and  $H + \geq 3\text{jet}$  events as well as for the uncertainty due to the multivariate requirement on  $|\Delta\phi_{\gamma\gamma,jj}|$  (cf. Sec. VIII A 4), which suppresses additional jet activity. The uncertainty estimation uses an extension of the Stewart–Tackmann method [103,104] and typically ranges between 20% and 32%.

The applicability of these uncertainties to POWHEG NNLOPS was tested by comparing the STWZ+BLPTW and the MCFM cross section predictions in variables relevant for the definition of the simplified cross-section bins, and reasonable agreement was found. In addition, the ggH acceptance of POWHEG NNLOPS of all categories based on BDT classifiers is compared to the acceptance derived from the MG5\_AMC@NLO prediction or Refs. [46,105] which includes up to two jets at NLO accuracy using the FxFx merging scheme [106]. Sufficient agreement was found for all categories and no additional modeling uncertainties are assigned based on these comparisons.<sup>5</sup>

Finally, in the categories targeting production in association with top quarks, the normalization of each

<sup>5</sup>Recent measurements of QCD and electroweak (VBF)  $Z$ -boson production in association with two jets in Ref. [107] show large deviations between the data and the predictions for the QCD  $Zjj$  background at large  $m_{jj}$ . These differences are significantly larger than the 30–40% uncertainties assigned here to the ggH background in the experimental categories targeting VBF Higgs boson production. Increasing this uncertainty to 100% results in an increase in the theory uncertainty in the VBF signal strengths or simplified template cross sections by a factor of about two, while the increase in the total uncertainty is about 10%, as it is dominated by the statistical component.

of the  $ggH$ , VBF, and  $VH$  production mechanisms is assigned an uncertainty of 100%, motivated by comparisons of data with simulation in  $t\bar{t}b\bar{b}$  [108] and  $Vb$  [109,110] productions, but this has little impact on the final results.

The uncertainty in the modeling of the parton shower, underlying event and hadronization affects all measurements (labeled as “UE/PS” in the following). It is estimated by taking the relative difference in acceptance at particle level after switching the parton showering algorithm from PYTHIA8 to HERWIG7 in the  $ggH$ , VBF, and  $VH$  samples and from HERWIG++ to PYTHIA8 in the  $t\bar{t}H$  sample, respectively. These differences are treated as four independent uncertainty sources. Additionally, for  $ggH$  the effect of the eigenvector tunes from the AZNLO set are merged to provide one additional uncertainty component.

The theoretical modeling uncertainties in the measurement of signal strengths include all of the sources that affect the measurement of the production-mode cross sections, plus additional uncertainties in the overall normalization of each production mechanism. Uncertainties in the overall normalization of each production process from missing higher-order QCD effects and the choice of parton distribution function are specified as part of the theoretical calculations used to normalize the simulated samples. The normalization uncertainty from the  $H \rightarrow \gamma\gamma$  branching ratio is taken from HDECAY and PROPHECY4F.

#### D. Theoretical and modeling uncertainties for fiducial integrated and differential results

The theoretical modeling uncertainty in the detector correction factor (introduced in Sec. IX B) used to measure the fiducial integrated and differential cross sections is taken to be the envelope of the following three sources:

- (1) The uncertainty in the relative contributions of the different Higgs boson production mechanisms. This uncertainty is estimated by varying the fraction of the  $ggH$ , VBF,  $VH$  and  $t\bar{t}H$  processes by an amount commensurate with the 68% confidence levels of the measured production mode cross-section ratios [5]. The variations of each production mechanism are carried out simultaneously and include the known correlations between the measured production mode cross-section ratios. These uncertainties range from 0.1% to 31%, depending on the fiducial region or differential variable, increasing typically in bins and regions sensitive to  $t\bar{t}H$ -production.
- (2) The uncertainty in the detector correction factor due to a possible mismodeling of the Higgs boson transverse momentum and rapidity distributions is estimated by reweighting the Higgs boson

distributions in simulation to match those observed in the data. The resulting uncertainties range from 0.1% to 4.5%, increasing in fiducial regions and bins with high jet multiplicities.

- (3) The uncertainty in the modeling of the parton shower, underlying event, and hadronization. This uncertainty is derived as described in Sec. VII C and the size of this uncertainty ranges from from 0.1% up to 30%, with the highest uncertainties in fiducial regions with large missing transverse energy.

Typically differential measurements involving only the photon kinematics are less affected by these model uncertainties than measurements using selections on jets or missing transverse momentum.

#### E. Illustration of model errors for simplified template cross section and fiducial cross section measurements

To illustrate the difference between the two approaches of assigning theory and model errors used for category based results and the fiducial cross section results, the theoretical modeling uncertainties in the corresponding zero-jet  $ggF$ -dominated and VBF-dominated regions are compared.

The simplified template cross section defined as  $gg \rightarrow H$  events with  $|y_H| < 2.5$  and no jets derives its sensitivity from the two categories requiring no jet and either one or both photons reconstructed in the barrel region of the electromagnetic calorimeter (defined by  $|\eta| \leq 0.95$ ). The total theory uncertainty is dominated by the uncertainty in the choice of parton distribution functions (1.5%), in the value of  $\alpha_s$  (1.4%), and in the modeling of the parton shower, underlying event, and hadronization (1.7%), and amounts to a relative uncertainty of 2.7%. The fiducial zero-jet cross section, in contrast, has only a modeling uncertainty of 0.1%, dominated by the possible mismodeling of the Higgs boson transverse momentum and rapidity distributions.

The simplified template cross section defined as  $qq \rightarrow Hqq$  events with  $|y_H| < 2.5$  and no jets with  $p_T^j < 200$  GeV derives its sensitivity from four VBF categories employing multivariate methods to distinguish VBF events from background processes. The total theory uncertainty is dominated by the modeling uncertainties in the parton shower, underlying event, and hadronization (9%) and in the remaining  $ggH$  contamination (5%), and amounts to a relative uncertainty of 13%. In contrast, the fiducial VBF cross section, defined by a dijet mass  $m_{jj}$  of at least 400 GeV, a large rapidity separation  $|\Delta y_{jj}| > 2.8$ , and an azimuthal difference between the Higgs boson and the dijet pair of  $|\Delta\phi_{\gamma\gamma,jj}| > 2.6$ , has only a modeling uncertainty of 4.5%, dominated by the composition variation of the fractions of the  $ggH$ , VBF,  $VH$  and  $t\bar{t}H$  processes (4.5%).

## VIII. MEASUREMENT OF TOTAL PRODUCTION-MODE CROSS SECTIONS, SIGNAL STRENGTHS, AND STXS

### A. Event categorization

The events satisfying the diphoton selection discussed in Sec. VB are classified, in accord with the reconstructed event kinematics and topology, into 31 exclusive categories that are optimized for the best separation of the Higgs boson production processes and for the maximum sensitivity to the phase space regions defined by the stage 1 of the simplified template cross-section framework. A combined fit to the event reconstruction categories is then performed to determine nine simplified template cross sections (with  $|y_H| < 2.5$ ), as well as production-mode cross sections and signal strength interpretations of the data. The categorization proceeds from the production modes with the smallest expected cross sections to the production modes with largest expected cross sections, in the order described below. In categories with definitions based on jet properties, jets with transverse momenta greater than 30 GeV are used, unless explicitly stated otherwise.

#### 1. $t\bar{t}H$ and $tH$ enriched categories

Nine categories enriched in events produced in association with a top quark are defined to target the  $t\bar{t}H$ ,  $tHq$ , and  $tHW$  production modes. These categories are separated into a hadronic channel, where top quarks in the event decay to hadrons via  $t \rightarrow Wb \rightarrow qq'b$ ; and a leptonic channel, where at least one top quark decays to a charged lepton via  $t \rightarrow Wb \rightarrow \ell\nu b$ . The single top quark categories are optimized for sensitivity to SM  $tH$  production, and are expected to provide additional sensitivity to anomalous values of the top quark Yukawa coupling.

Three categories target the leptonic channel by requiring the presence of at least one prompt lepton and at least one  $b$ -tagged jet with transverse momentum greater than 25 GeV. Two of these categories target  $tH$  production while the third one is optimized for  $t\bar{t}H$  events. Both  $tH$  categories veto events with more than one prompt lepton. The first of these categories (“ $tH$  lep 0fwd”) contains events with at most three central jets ( $|\eta| < 2.5$ ) and a veto on forward jets ( $|\eta| > 2.5$ ). The second  $tH$  category (“ $tH$  lep 1fwd”) is defined by events with at most four central jets and at least 1 forward jet. The “ $t\bar{t}H$  lep” category includes events with at least two central jets, while no requirement is applied to the forward jets. To suppress  $ZH$  events with  $Z \rightarrow \ell\ell$ , same-flavor dilepton candidates with an invariant mass within 10 GeV of the  $Z$  boson mass are vetoed.

Six categories target the hadronic decay channel by selecting events with no prompt leptons and at least three jets, of which at least one is  $b$ -tagged. Four of

these categories (“ $t\bar{t}H$  had BDT1” to “ $t\bar{t}H$  had BDT4”) are defined by means of a boosted decision tree (BDT) trained to identify  $t\bar{t}H$  signal against  $ggH$  and multijet background. The BDT exploits five kinematic variables:  $H_T$ , the scalar sum of jet transverse momenta,  $m_{\text{all jets}}$ , the mass of all jets, as well as the number of all jets, central jets ( $|\eta| < 2.5$ ), and  $b$ -tagged jets. The training uses  $t\bar{t}H$  and  $ggH$  simulated events and a data-driven multijet background sample defined by diphoton events with at least three jets and in which at least one photon fails to meet either identification or isolation requirements. Using the BDT response as a discriminating variable, events are separated into four categories with an expected fraction of  $t\bar{t}H$  events (among all Higgs boson events in this category) of 95%, 89%, 86%, and 79%, respectively. Two additional hadronic categories enhanced in  $tH$  production (“ $tH$  had 4j1b” and “ $tH$  had 4j2b”) are included, defined by events with exactly four jets with transverse momentum greater than 25 GeV and split by events with exactly one or two  $b$ -tagged jets, respectively. The distributions of two of the discriminating variables are shown in Fig. 4.

#### 2. $VH$ leptonic enriched categories

Five categories are enriched in Higgs boson production in association with a vector boson, based on different decays of the vector bosons.

The  $VH$  dilepton category (“ $VH$  dilep”) targets  $ZH$  production with  $Z \rightarrow \ell\ell$  by requiring the presence of two same-flavor opposite-sign leptons with an invariant mass between 70 GeV and 110 GeV. Two additional  $VH$  one-lepton categories target  $WH$  production with  $W \rightarrow \ell\nu$ . Events are requested to contain exactly one selected electron or muon. To suppress  $ZH$  events with  $Z$  bosons decaying to  $ee$ , in which an electron is misidentified as a photon, a veto is applied to events in which the invariant mass of the selected electron and any of the two signal photons is between 84 GeV and 94 GeV. Events are then split into two regions, where the  $p_T$  of the lepton +  $E_T^{\text{miss}}$  system is higher (“ $VH$  lep High”) or lower (“ $VH$  lep Low”) than 150 GeV. An additional requirement on the  $E_T^{\text{miss}}$  significance, defined as  $E_T^{\text{miss}} / \sqrt{\sum E_T}$ , of at least 1.0 is applied to events in the low lepton +  $E_T^{\text{miss}}$   $p_T$  category.

Two  $VH$  missing transverse momentum categories target  $ZH$  production with  $Z \rightarrow \nu\nu$  and  $W \rightarrow \ell\nu$  where the lepton was not reconstructed or failed to meet the selection criteria. One category (“ $VH$  MET Low”) requires  $80 \text{ GeV} < E_T^{\text{miss}} < 150 \text{ GeV}$  and  $E_T^{\text{miss}}$  significance greater than 8. The other category (“ $VH$  MET High”) requires  $E_T^{\text{miss}} > 150 \text{ GeV}$  and  $E_T^{\text{miss}}$  significance greater than 9, or  $E_T^{\text{miss}} > 250 \text{ GeV}$ .






FIG. 4. The normalized distributions for the expected background of two kinematic variables used for the selection of the hadronic  $t\bar{t}H$  categories: (a)  $H_T$  and (b)  $m_{\text{all jets}}$  for events after the first step of the selection (see text) for simulated  $H \rightarrow \gamma\gamma$  events produced via  $t\bar{t}H$  (blue) and gluon–gluon fusion (red), for the expected background derived from the data control region (green) described in the text and events from data with  $105 \text{ GeV} < m_{\gamma\gamma} < 120 \text{ GeV}$  or  $130 \text{ GeV} < m_{\gamma\gamma} < 160 \text{ GeV}$  (black dots with error bars showing the statistical uncertainty).

### 3. BSM enriched and VH hadronic categories

To provide sensitivity to potential beyond SM contributions, a category (“jet BSM”) defined by events with a leading jet with transverse momentum greater than 200 GeV is included in the event selection. This category includes SM events in the typical VBF topology, boosted  $V(\rightarrow jj)H$  production where the vector boson is reconstructed as a single jet, as well as events produced in gluon–gluon fusion with an energetic jet.

Two  $VH$  hadronic categories target  $VH$  production with a hadronically decaying vector boson. Events are required to have at least two jets with  $60 < m_{jj} < 120 \text{ GeV}$ . A BDT classifies the events using the following information: the dijet invariant mass, the component  $p_{T\perp}^{\gamma\gamma}$  of the diphoton  $\vec{p}_T$  transverse to its thrust axis in the transverse plane, the rapidity difference between the dijet and the diphoton system, and the cosine  $\cos \theta_{\gamma\gamma,jj}^*$  where  $\theta_{\gamma\gamma,jj}^*$  is the angle between the diphoton system momentum and the direction of motion of the diphoton–dijet system in the Collins–Soper frame. The training uses  $VH$  events as signal, and a mixture of simulated signals (everything except  $VH$  events), simulated  $\gamma\gamma$  events, and  $\gamma j$  and  $jj$  data control samples as background. Using the BDT response as a discriminating variable, events are classified into two categories (“ $VH$  had tight” and “ $VH$  had loose”) with an expected fraction of signal events due to  $VH$  production of 42% and 25%, respectively.

Figure 5 shows the distributions of  $m_{jj}$  and  $p_{T\perp}^{\gamma\gamma}$  in signal and background events and in events selected in data from the  $m_{\gamma\gamma}$  sidebands. The variables show good separation between  $VH$  events and both the other signal events and background events.

### 4. VBF enriched categories

Four categories are defined to enhance the sensitivity to vector boson fusion production. Events are required to

contain at least two hadronic jets, and the selections applied are based on the two leading jets ( $j_1, j_2$ ) in the event. The pseudorapidity separation  $|\Delta\eta_{jj}|$  between the two leading jets is required to be greater than 2. In addition  $|\eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})|$  is required to be less than 5, with  $\eta_{\gamma\gamma}$  denoting the pseudorapidity of the diphoton system. The events are first split into two regions based on the value of the transverse momentum  $p_T^{Hjj}$  of the vector sum of the momenta of the reconstructed Higgs boson and of the two leading jets. This variable is highly correlated with the  $p_T$  of the third jet due to momentum balance. The signal in the  $p_T^{\gamma\gamma jj} < 25 \text{ GeV}$  “low  $p_T^{Hjj}$ ” region is dominated by exclusive two-jetlike events, while the signal in the  $p_T^{\gamma\gamma jj} > 25 \text{ GeV}$  “high  $p_T^{Hjj}$ ” region is dominated by inclusive  $\geq 3$ -jet like events. This choice minimizes the otherwise large ggH jet-migration uncertainties in this phase space and is similar to a central-jet veto that separates contributions from ggH and VBF.

A BDT is then used to classify events in each region, using six kinematic variables:  $m_{jj}$ ,  $|\Delta\eta_{jj}|$ ,  $p_{T\perp}^{\gamma\gamma}$ , the absolute azimuthal difference of the diphoton and the dijet system  $|\Delta\phi_{\gamma\gamma,jj}|$ , the minimum angular separation between either of the two signal photons and either of the two leading jets  $\Delta R_{\gamma j}^{\min}$ , and  $|\eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})|$ . A requirement of  $|\Delta\phi_{\gamma\gamma,jj}|$  to be near  $\pi$  effectively vetoes additional jets in the event by restricting the phase space for additional emissions and, to avoid large theoretical uncertainties, the BDT does not use shape information for events with  $|\Delta\phi_{\gamma\gamma,jj}| > 2.94$  by merging these events into one bin. The training of the BDT uses VBF events as signal, and a mixture of simulated gluon–gluon fusion and  $\gamma\gamma$  events and of  $\gamma j$  and  $jj$  data control samples as background. Four exclusive categories are defined with “loose” and “tight” requirements on the BDT classifier in the two  $p_T^{Hjj}$  regions. The “tight” category in the  $p_T^{Hjj} > 25 \text{ GeV}$  region has an






FIG. 5. The normalized distributions of two kinematic variables used for the selection of the  $VH$  hadronic categories: (a)  $m_{jj}$  and (b)  $p_{T\gamma\gamma}$  for events after the first step of the  $VH$  hadronic category preselection (see text) for simulated  $H \rightarrow \gamma\gamma$  events produced in association with hadronically decaying vector bosons (blue) or through  $ggH$ ,  $VBF$  or  $t\bar{t}H$  processes (red), for the expected background from data ( $\gamma j$ ,  $j j$ ) and simulation ( $\gamma\gamma$ ,  $V\gamma\gamma$ ) control samples (green, purple), and for events from data with  $105 \text{ GeV} < m_{\gamma\gamma} < 120 \text{ GeV}$  or  $130 \text{ GeV} < m_{\gamma\gamma} < 160 \text{ GeV}$  (black dots with error bars showing the statistical uncertainty).

expected fraction of VBF events among all Higgs boson events in this category of 49%, while the “loose” category has an expected fraction of VBF events of 20%. In the  $p_{T}^{Hjj} < 25 \text{ GeV}$  region the “tight” category has an expected fraction of VBF events of 85%, whereas the “loose” category has an expected fraction of 61%.

Figure 6 shows the distributions of  $|\Delta\eta_{jj}|$  and  $|\Delta\phi_{\gamma\gamma,jj}|$  in simulated  $H \rightarrow \gamma\gamma$  events, background events from simulated diphotons and data control samples of  $\gamma j$  and  $j j$  events, and events selected from the  $m_{\gamma\gamma}$  sidebands in data. The variables show good separation between VBF events and both gluon–gluon fusion events and background events.

### 5. Untagged categories

The remaining “untagged” events are dominated by events produced through gluon–gluon fusion and they are further split into ten categories. The untagged events are first separated by jet multiplicity into events with zero jets, exactly one jet, or at least two jets. The zero-jet events are split into two categories with either two photons in the “central” pseudorapidity region  $|\eta| < 0.95$ , in which the energy resolution is better (“ $ggH$  0J Cen”), or with at least one photon in the “forward” region  $|\eta| > 0.95$  which has worse energy resolution (“ $ggH$  0J FWD”). The exclusive one-jet (“ $ggH$  1J”) and inclusive two-jet (“ $ggH$  2J”)




FIG. 6. The normalized distributions for the expected background of two kinematic variables used for the selection of the VBF categories: (a)  $|\Delta\eta_{jj}|$  and (b)  $|\Delta\phi_{\gamma\gamma,jj}|$  for events after the first step of the selection (see text) for simulated  $H \rightarrow \gamma\gamma$  events produced via vector-boson fusion (blue) and gluon–gluon fusion (red), for the expected background from data ( $\gamma j$ ,  $j j$ ) and simulation ( $\gamma\gamma$ ) control samples (green), and for events from data with  $105 \text{ GeV} < m_{\gamma\gamma} < 120 \text{ GeV}$  or  $130 \text{ GeV} < m_{\gamma\gamma} < 160 \text{ GeV}$  (black dots with error bars showing the statistical uncertainty).

categories are further split into regions of diphoton transverse momentum with  $p_T^{\gamma\gamma} \in [0, 60]$  (“Low”),  $[60, 120]$  (“Med”),  $[120, 200]$  (“High”) or  $>200$  GeV (“BSM”), the latter of which is particularly sensitive to the presence of BSM physics in the loop diagrams associated with the gluon–gluon fusion production mode.

### 6. Categorization summary

A summary of the selection requirements defining each category is provided in Table IV. The predicted signal efficiencies times acceptance and the event fractions per production mode for each category are given in Table V. The

fractions of signal events in each reconstructed category originating from a given simplified template cross-section region are shown in Fig. 7. The defined ggH categories exhibit high purities as they are defined with a near one-to-one correspondence with the STXS regions despite small contaminations arising primarily from pileup and selection inefficiencies. The  $qq \rightarrow Hqq$  bins are more ambiguous however still retain much of the diagonal structure. On the other hand, the  $VH$  leptonic and top categories are not sensitive to all of the STXS regions of interest, necessitating a merging. Finally, the fractions of signal events in each category from a given production mode are shown in Figure 8.

TABLE IV. Shorthand label and event selection defining each of the 31 event reconstruction categories for the measurement of the signal strengths and simplified template cross sections. The labels denote the predominant production process or kinematic properties the category targets. Jets are required to have  $p_T > 30$  GeV unless otherwise noted. The categories are mutually exclusive and the criteria are applied in descending order of the shown categories.

Category	Selection
tH lep 0fwd	$N_{\text{lep}} = 1, N_{\text{jets}}^{\text{cen}} \leq 3, N_{b\text{-tag}} \geq 1, N_{\text{jets}}^{\text{fwd}} = 0 (p_T^{\text{jet}} > 25 \text{ GeV})$
tH lep 1fwd	$N_{\text{lep}} = 1, N_{\text{jets}}^{\text{cen}} \leq 4, N_{b\text{-tag}} \geq 1, N_{\text{jets}}^{\text{fwd}} \geq 1 (p_T^{\text{jet}} > 25 \text{ GeV})$
ttH lep	$N_{\text{lep}} \geq 1, N_{\text{jets}}^{\text{cen}} \geq 2, N_{b\text{-tag}} \geq 1, Z_{\ell\ell} \text{ veto } (p_T^{\text{jet}} > 25 \text{ GeV})$
ttH had BDT1	$N_{\text{lep}} = 0, N_{\text{jets}} \geq 3, N_{b\text{-tag}} \geq 1, \text{BDT}_{\text{ttH}} > 0.92$
ttH had BDT2	$N_{\text{lep}} = 0, N_{\text{jets}} \geq 3, N_{b\text{-tag}} \geq 1, 0.83 < \text{BDT}_{\text{ttH}} < 0.92$
ttH had BDT3	$N_{\text{lep}} = 0, N_{\text{jets}} \geq 3, N_{b\text{-tag}} \geq 1, 0.79 < \text{BDT}_{\text{ttH}} < 0.83$
ttH had BDT4	$N_{\text{lep}} = 0, N_{\text{jets}} \geq 3, N_{b\text{-tag}} \geq 1, 0.52 < \text{BDT}_{\text{ttH}} < 0.79$
tH had 4j1b	$N_{\text{lep}} = 0, N_{\text{jets}}^{\text{cen}} = 4, N_{b\text{-tag}} = 1 (p_T^{\text{jet}} > 25 \text{ GeV})$
tH had 4j2b	$N_{\text{lep}} = 0, N_{\text{jets}}^{\text{cen}} = 4, N_{b\text{-tag}} \geq 2 (p_T^{\text{jet}} > 25 \text{ GeV})$
VH dilep	$N_{\text{lep}} \geq 2, 70 \text{ GeV} \leq m_{\ell\ell} \leq 110 \text{ GeV}$
VH lep High	$N_{\text{lep}} = 1,  m_{e\gamma} - 89 \text{ GeV}  > 5 \text{ GeV}, p_T^{\ell + E_T^{\text{miss}}} > 150 \text{ GeV}$
VH lep Low	$N_{\text{lep}} = 1,  m_{e\gamma} - 89 \text{ GeV}  > 5 \text{ GeV}, p_T^{\ell + E_T^{\text{miss}}} < 150 \text{ GeV}, E_T^{\text{miss}} \text{ significance} > 1$
VH MET High	$150 \text{ GeV} < E_T^{\text{miss}} < 250 \text{ GeV}, E_T^{\text{miss}} \text{ significance} > 9 \text{ or } E_T^{\text{miss}} > 250 \text{ GeV}$
VH MET Low	$80 \text{ GeV} < E_T^{\text{miss}} < 150 \text{ GeV}, E_T^{\text{miss}} \text{ significance} > 8$
jet BSM	$p_{T,j1} > 200 \text{ GeV}$
VH had tight	$60 \text{ GeV} < m_{ij} < 120 \text{ GeV}, \text{BDT}_{\text{VH}} > 0.78$
VH had loose	$60 \text{ GeV} < m_{ij} < 120 \text{ GeV}, 0.35 < \text{BDT}_{\text{VH}} < 0.78$
VBF tight, high $p_T^{Hjj}$	$ \Delta\eta_{jj}  > 2,  \eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})  < 5, p_T^{Hjj} > 25 \text{ GeV}, \text{BDT}_{\text{VBF}} > 0.47$
VBF loose, high $p_T^{Hjj}$	$ \Delta\eta_{jj}  > 2,  \eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})  < 5, p_T^{Hjj} > 25 \text{ GeV}, -0.32 < \text{BDT}_{\text{VBF}} < 0.47$
VBF tight, low $p_T^{Hjj}$	$ \Delta\eta_{jj}  > 2,  \eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})  < 5, p_T^{Hjj} < 25 \text{ GeV}, \text{BDT}_{\text{VBF}} > 0.87$
VBF loose, low $p_T^{Hjj}$	$ \Delta\eta_{jj}  > 2,  \eta_{\gamma\gamma} - 0.5(\eta_{j1} + \eta_{j2})  < 5, p_T^{Hjj} < 25 \text{ GeV}, 0.26 < \text{BDT}_{\text{VBF}} < 0.87$
ggH 2J BSM	$\geq 2 \text{ jets}, p_T^{\gamma\gamma} \geq 200 \text{ GeV}$
ggH 2J High	$\geq 2 \text{ jets}, p_T^{\gamma\gamma} \in [120, 200] \text{ GeV}$
ggH 2J Med	$\geq 2 \text{ jets}, p_T^{\gamma\gamma} \in [60, 120] \text{ GeV}$
ggH 2J Low	$\geq 2 \text{ jets}, p_T^{\gamma\gamma} \in [0, 60] \text{ GeV}$
ggH 1J BSM	$= 1 \text{ jet}, p_T^{\gamma\gamma} \geq 200 \text{ GeV}$
ggH 1J High	$= 1 \text{ jet}, p_T^{\gamma\gamma} \in [120, 200] \text{ GeV}$
ggH 1J Med	$= 1 \text{ jet}, p_T^{\gamma\gamma} \in [60, 120] \text{ GeV}$
ggH 1J Low	$= 1 \text{ jet}, p_T^{\gamma\gamma} \in [0, 60] \text{ GeV}$
ggH 0J Fwd	$= 0 \text{ jets, one photon with }  \eta  > 0.95$
ggH 0J Cen	$= 0 \text{ jets, two photons with }  \eta  \leq 0.95$

TABLE V. Signal efficiencies times acceptance,  $\epsilon$ , and expected signal event fractions per production mode,  $f$ , in each category for  $\sqrt{s} = 13$  TeV and  $m_H = 125.09$  GeV. The second-to-last row shows the total efficiency per production process summed over the categories. Values labeled as “nil” correspond to efficiencies or fractions that are smaller than 0.05%. The total number of expected signal events,  $N_S$ , in the last row corresponds to an integrated luminosity of  $36.1 \text{ fb}^{-1}$ .

Category	ggH		VBF		WH		ZH		ttH		$b\bar{b}H$		$tHq$		$tHW$		All
	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$\epsilon$ [%]	$f$ [%]	$N_S$
ggH 0J Cen	8.9	97.3	1.2	1.1	1.4	0.4	1.9	0.4	nil	nil	8.2	0.9	nil	nil	nil	nil	333.5
ggH 0J Fwd	15.5	97.0	2.4	1.2	3.0	0.5	3.7	0.4	nil	nil	14.7	0.9	0.2	nil	0.1	nil	579.5
ggH 1J Low	7.2	90.5	5.7	5.7	5.0	1.7	4.4	1.0	0.1	nil	9.1	1.1	0.5	nil	0.2	nil	289.9
ggH 1J Med	3.6	83.5	6.4	11.7	4.2	2.6	4.1	1.6	0.1	nil	1.9	0.4	0.6	nil	0.3	nil	156.2
ggH 1J High	0.7	76.0	1.9	17.5	1.1	3.4	1.4	2.7	0.1	0.1	0.3	0.3	0.2	nil	0.1	nil	31.5
ggH 1J BSM	nil	72.4	0.1	16.9	0.1	6.0	0.2	4.2	nil	0.3	nil	nil	nil	0.1	nil	nil	2.2
ggH 2J Low	1.8	79.1	2.7	9.6	3.7	4.5	4.1	3.1	2.2	1.1	5.4	2.3	3.9	0.3	1.9	nil	81.1
ggH 2J Med	1.5	77.6	3.1	12.2	3.2	4.4	3.8	3.2	2.6	1.5	1.6	0.7	4.5	0.4	2.4	nil	72.4
ggH 2J High	0.6	75.8	1.3	12.8	1.4	4.9	1.9	4.0	1.4	2.0	0.1	0.1	2.2	0.4	1.6	0.1	29.2
ggH 2J BSM	0.2	76.2	0.3	10.3	0.4	4.9	0.6	4.6	0.6	3.0	0.1	0.2	0.8	0.6	1.3	0.2	7.6
VBF Hjj Low loose	0.2	32.3	4.5	66.7	0.1	0.3	0.1	0.3	nil	nil	0.1	0.2	0.3	0.1	nil	nil	19.4
VBF Hjj Low tight	nil	12.9	4.2	86.7	nil	0.1	nil	0.1	nil	nil	nil	nil	0.3	0.1	nil	nil	13.8
VBF Hjj High loose	0.3	69.9	1.4	23.8	0.4	2.2	0.5	1.8	0.4	0.9	0.4	0.7	1.8	0.6	0.5	nil	16.5
VBF Hjj High tight	0.3	47.0	3.4	48.2	0.2	1.2	0.4	1.3	0.4	0.8	0.2	0.3	4.4	1.2	0.6	nil	20.2
VHhad loose	0.3	67.2	0.3	4.9	2.4	14.6	2.9	11.0	0.6	1.6	0.2	0.4	0.8	0.3	0.8	0.1	16.5
VHhad tight	0.2	52.4	0.1	3.4	3.0	23.8	3.5	18.0	0.6	1.9	nil	0.1	0.5	0.2	1.0	0.1	12.3
jet BSM	0.4	59.9	2.4	25.8	1.6	5.9	1.9	4.4	2.0	3.0	0.1	0.1	3.1	0.6	5.1	0.2	26.7
VHMET Low	nil	11.9	nil	0.4	0.1	23.4	0.6	63.2	nil	0.5	nil	0.3	nil	0.2	nil	nil	0.6
VHMET High	nil	1.3	nil	0.1	0.3	22.8	1.4	66.2	0.3	8.3	nil	nil	0.1	0.6	0.8	0.7	1.3
VHlep Low	nil	11.4	nil	1.1	4.4	68.0	0.8	8.1	1.3	8.5	0.2	0.9	1.8	1.6	2.2	0.4	6.4
VHlep High	nil	0.2	nil	nil	1.2	76.5	0.1	4.6	0.6	16.2	nil	nil	0.3	1.2	1.6	1.3	1.5
VHdilep	nil	nil	nil	nil	nil	nil	1.4	95.8	0.1	4.0	nil	nil	nil	nil	0.1	0.2	0.9
tHhad 4j2b	nil	23.8	nil	2.8	nil	1.6	0.1	13.5	0.6	39.0	0.1	8.2	1.2	10.5	0.3	0.6	0.6
tHhad 4j1b	nil	35.4	nil	4.0	0.1	4.3	0.3	7.9	2.2	36.3	0.2	2.2	3.8	8.5	2.6	1.3	2.5
ttHhadBDT4	nil	7.0	nil	0.8	nil	1.4	0.2	4.5	4.8	79.4	nil	0.3	1.9	4.3	4.7	2.4	2.5
ttHhadBDT3	nil	3.5	nil	0.5	nil	1.0	nil	3.1	1.3	86.1	nil	0.5	0.3	3.1	1.1	2.2	0.6
ttHhadBDT2	nil	3.6	nil	0.3	nil	0.8	nil	1.6	3.8	89.3	nil	0.2	0.6	1.8	3.4	2.4	1.8
ttHhadBDT1	nil	1.2	nil	0.1	nil	0.1	nil	0.7	3.4	95.0	nil	0.1	0.2	0.7	2.5	2.1	1.4
ttHlep	nil	nil	nil	nil	nil	0.2	nil	0.1	5.6	96.0	nil	0.1	0.4	1.0	5.0	2.6	2.4
tHlep 1fwd	nil	1.8	nil	0.2	nil	1.4	nil	0.9	2.1	79.4	nil	0.2	2.6	13.5	2.3	2.6	1.1
tHlep 0fwd	nil	4.1	nil	0.2	0.1	5.6	nil	2.8	1.9	75.7	nil	0.9	1.5	8.2	2.1	2.5	1.0
Total $\epsilon$ [%]	41.8	...	41.3	...	37.6	...	40.5	...	39.1	...	42.8	...	38.9	...	44.5	...	41.8
Events	1518.4		119.1		37.1		25.2		16.0		14.8		2.2		0.5		1733.2

More information about the number of background events, the purity and the SM signal composition can be found in Tables XXVII and XXVIII in Appendix E.

### B. Production mode measurements

Using the 31 categories, total and production mode specific signal strength measurements are carried out. Measurements of total production cross sections and simplified template cross sections are reported. The simplified template cross sections are measured in a merged scheme introduced in Sec. IB and summarized in Table I. In addition, the result of coupling-strength fits are reported.

### 1. Observed data

The observed invariant mass distribution of the selected diphoton pairs of all categories as defined in Table IV, is shown in Fig. 9. Figure 10 shows the invariant mass distributions for the sums of the categories most sensitive to the different production modes. In all cases, for illustration purposes, events in each category are weighted according to the expected signal ( $S_{90}$ ) to background ( $B_{90}$ ) ratio in a  $m_{\gamma\gamma}$  region containing 90% of the expected signal yield, using a weight of the form  $\ln(1 + S_{90}/B_{90})$ . The results of signal-plus-background fits to these spectra, displaying both the total sum and the background-only components, are shown, as well as the residuals between the data and the background




FIG. 7. The fraction of signal events assigned to each reconstructed category ( $x$  axis and listed in Table IV) and originating from a given region (listed in Table I) of the stage-1 simplified template cross section framework ( $y$  axis). The black lines separate the  $t\bar{t}H$  and  $tH$ ,  $VH$  leptonic,  $VH$  hadronic and VBF enriched, and untagged categories, along with the simplified template cross-section regions they are most sensitive to. The color shows the purity of the region per category.

component. Both the signal-plus-background and background-only distributions shown are obtained from the sum of the individual distributions in each category weighted in the same way as the data points. In the fit of Fig. 9 a single signal strength  $\mu$  affecting simultaneously all production modes has been assumed, while in the fits of Fig. 10 the four signal strengths  $\mu_{\text{ggH}}$ ,  $\mu_{\text{VBF}}$ ,  $\mu_{\text{VH}}$  and  $\mu_{t\bar{t}H+tH}$  are allowed to vary separately, as described in the following section. The observed mass peak of the Higgs boson, constrained in the fit as  $m_H = 125.09 \pm 0.24$  GeV, is well within 68% CL of the Run 1 ATLAS + CMS combined measurement.

## 2. Signal strengths

The signal strengths, i.e., the ratios of the measured Higgs boson production-mode cross sections times diphoton branching ratio to the SM predictions for each production mode, are measured with the extended likelihood analysis described in Sec. VIC. In the likelihood the signal yield  $N_{\text{sig},m}^i$  in each category  $i$  for a particular production mode  $m$  is expressed as the product of the integrated luminosity  $\int L dt$ , the signal strength  $\mu_m$  for that production mode, the expected SM Higgs boson production mode cross section times branching ratio to diphotons, and the acceptance times efficiency  $\epsilon$  (Table V) for signal events from that production mode in the selected category ( $N_{\text{sig},m}^i = \mu_m \times \int L dt \times \sigma_m^{\text{SM}} \times B(H \rightarrow \gamma\gamma) \times \epsilon_m^i$ ).

A global signal strength  $\mu$  is measured assuming the ratios between different production processes to be as predicted by the SM. The profile of the negative log-likelihood ratio  $\lambda(\mu)$  of the global signal strength of all Higgs processes  $\mu$  for  $m_H = 125.09 \pm 0.24$  GeV is shown in Fig. 11.

The measured central value and 68% CL interval for  $\mu$  is found to be

$$\mu = 0.99^{+0.15}_{-0.14} = 0.99 \pm 0.12(\text{stat})^{+0.06}_{-0.05}(\text{exp})^{+0.07}_{-0.05}(\text{theo}),$$

well compatible with the SM prediction ( $\mu = 1$ ). This result confirms the ATLAS Run-1 diphoton signal strength measurement of  $\mu = 1.17 \pm 0.23(\text{stat})^{+0.10}_{-0.08}(\text{exp})^{+0.12}_{-0.08}(\text{theo})$  with around a factor of two improvement in each component of the uncertainty. The Run-1 result was obtained using the NNLO SM prediction for ggH production [17,111], which is about 10% lower than the N<sup>3</sup>LO calculation used here (see Sec. IV). The impact of the main sources of systematic uncertainty (presented in Table III and Sec. VII) in the measured global signal strength is summarized in Table VI. The distinction between yield and migration uncertainties adopted in Table III is used and the uncertainties are grouped into theory uncertainties, experimental uncertainties, mass resolution and scale, background shape, and luminosity.






FIG. 8. The expected composition of the selected Higgs boson events, in terms of the different production modes, for each reconstructed category.

In addition to the global signal strength, the signal strengths of the primary production processes are evaluated by exploiting the sensitivities of the analysis categories of Table IV to specific production processes. The measured signal strengths are shown together with the global signal strengths discussed above in Fig. 12 and found to be

$$\mu_{\text{ggH}} = 0.81^{+0.19}_{-0.18} = 0.81 \pm 0.16(\text{stat})^{+0.07}_{-0.06}(\text{exp})^{+0.07}_{-0.05}(\text{theo})$$

$$\mu_{\text{VBF}} = 2.0^{+0.6}_{-0.5} = 2.0 \pm 0.5(\text{stat})^{+0.3}_{-0.2}(\text{exp})^{+0.3}_{-0.2}(\text{theo})$$

$$\mu_{\text{VH}} = 0.7^{+0.9}_{-0.8} = 0.7 \pm 0.8(\text{stat})^{+0.2}_{-0.2}(\text{exp})^{+0.2}_{-0.1}(\text{theo})$$

$$\mu_{\text{top}} = 0.5^{+0.6}_{-0.6} = 0.5^{+0.6}_{-0.5}(\text{stat})^{+0.1}_{-0.1}(\text{exp})^{+0.1}_{-0.0}(\text{theo})$$

For Higgs boson production via  $VH$  the signal strength is assumed to be scaled by a single parameter (i.e.,

$\mu_{\text{VH}} = \mu_{\text{ZH}} = \mu_{\text{WH}}$ ). The  $b\bar{b}H$  contributions are scaled with  $\text{ggH}$  (i.e.,  $\mu_{\text{bbH}} = \mu_{\text{ggH}}$ ), and the  $tH$  and  $\bar{t}H$  productions are measured together rather than separately (i.e.,  $\mu_{\text{top}} = \mu_{\text{tH}+\bar{t}H}$ ).

The  $\text{ggH}$  signal strength is  $1\sigma$  below the Standard Model prediction, while the VBF signal strength is  $2.2\sigma$  above the prediction. The expected and observed significances  $Z_0$  of VBF production are reported in Table VII: the significance of the observed VBF signal is close to  $5\sigma$ .

Since no significant evidence is observed for  $VH$  and top-associated Higgs boson production, upper limits at 95% CL are reported for their signal strengths, as shown in Table VIII and Fig. 13. The accuracy of the asymptotic approximation was validated using ensembles of pseudoexperiments. Appendix F provides separate limits on  $\mu_{\text{ZH}}$  and  $\mu_{\text{WH}}$ , and Appendix G 1 shows the expected uncertainties for the inclusive and production-mode specific signal strengths reported in Fig. 12.




FIG. 9. Weighted diphoton invariant mass spectrum observed in the 2015 and 2016 data at 13 TeV. Each event is weighted by the  $\ln(1 + S_{90}/B_{90})$  ratio of the expected signal ( $S_{90}$ ) and background ( $B_{90}$ ) of the 90% signal quantile in the category to which it belongs to. The values of  $S_{90}$  and  $B_{90}$  used for each category are shown in Table XXVII of Appendix E. The error bars represent 68% confidence intervals of the weighted sums. The solid red curve shows the fitted signal-plus-background model when the Higgs boson mass is constrained to be  $125.09 \pm 0.24$  GeV. The background component of the fit is shown with the dotted blue curve. The signal component of the fit is shown with the solid black curve. Both the signal-plus-background and background-only curves reported here are obtained from the sum of the individual curves in each category weighted by the logarithm of unity plus the signal-to-background ratio. The bottom plot shows the residuals between the data and the background component of the fitted model.

### 3. Production-mode cross sections

The production-mode cross sections for  $m_H = 125.09 \pm 0.24$  GeV in a region with Higgs-boson rapidity  $|y_H| < 2.5$ , multiplied by the branching ratio of the Higgs boson decay to diphotons, are evaluated in the following way. The fitted value of  $\sigma_{\text{top}}$  corresponds to the sum of  $t\bar{t}H$ ,  $tHq$ , and  $tHW$  production-mode cross sections under the assumption that their relative ratios are as predicted by the SM. The  $VH$  production-mode cross section value is fitted under the assumption that the ratio of the  $WH$  and  $ZH$  production mode cross sections is as predicted by the SM and includes both production from quark and gluon initial states. Such results are obtained through signal + background fits to the diphoton invariant mass distribution in each category by expressing, in the likelihood, the signal yield  $N_{\text{sig},m}^i$  in each category  $i$  for a particular production mode  $m$  as  $N_{\text{sig},m}^i = \int L dt \times \sigma_m^{\text{SM}} \times B^{\text{SM}}(H \rightarrow \gamma\gamma) \times \epsilon_m^i$  using the same notation as in Sec. VIII B 2.

The production-mode cross sections are summarized in Fig. 14 and Table IX.

The 68% and 95% CL two-dimensional contours of  $\sigma_{\text{ggH}} \times B(H \rightarrow \gamma\gamma)$  and  $\sigma_{\text{VBF}} \times B(H \rightarrow \gamma\gamma)$  are shown in Fig. 15, profiling  $\sigma_{\text{VH}} \times B(H \rightarrow \gamma\gamma)$  and  $\sigma_{\text{top}} \times B(H \rightarrow \gamma\gamma)$  in the fits. The SM expectation of  $\sigma_{\text{ggH}} \times B(H \rightarrow \gamma\gamma)$  vs  $\sigma_{\text{VBF}} \times B(H \rightarrow \gamma\gamma)$  is within the 95% CL contour of this measurement.

To remove the impact of possible deviations in the  $H \rightarrow \gamma\gamma$  branching ratio, ratios of the production-mode cross sections to the ggH cross section are also extracted. Such ratios, normalized for convenience of presentation to the central values of their SM predictions, are<sup>6</sup>

$$\frac{\sigma_{\text{VBF}}/\sigma_{\text{ggH}}}{(\sigma_{\text{VBF}}/\sigma_{\text{ggH}})^{\text{SM}}} = 2.5^{+1.3}_{-0.9} = 2.5^{+1.1}_{-0.8}(\text{stat})^{+0.5}_{-0.3}(\text{exp})^{+0.5}_{-0.3}(\text{theo})$$

$$\frac{\sigma_{\text{VH}}/\sigma_{\text{ggH}}}{(\sigma_{\text{VH}}/\sigma_{\text{ggH}})^{\text{SM}}} = 0.9^{+1.3}_{-1.0} = 0.9^{+1.2}_{-0.9}(\text{stat})^{+0.3}_{-0.3}(\text{exp})^{+0.2}_{-0.1}(\text{theo})$$

$$\frac{\sigma_{\text{top}}/\sigma_{\text{ggH}}}{(\sigma_{\text{top}}/\sigma_{\text{ggH}})^{\text{SM}}} = 0.7^{+0.8}_{-0.7} = 0.7^{+0.8}_{-0.7}(\text{stat})^{+0.2}_{-0.1}(\text{exp})^{+0.2}_{-0.0}(\text{theo})$$

The ratios are also given in Table X, along with their statistical, experimental and theoretical uncertainties without the normalization to the central values of the SM predictions. Both the measurements of the ggH and VBF production modes and the evaluations of the  $VH$  and top production modes agree within  $1-2\sigma$  with the SM expectations. Appendix G 2 provides the expected uncertainties for the production mode cross sections.

### 4. Simplified template cross sections

As the current data are not yet sensitive to all of the 31 regions with  $|y_H| < 2.5$  (assuming SM acceptance) of the “stage-1” scheme of the simplified template cross-section framework, simplified template cross sections are reported for 10 phase space regions obtained from merging the initial 31 as described in Sec. IB and Table I. To retain sensitivity to BSM Higgs boson production, the  $p_T^H > 200$  GeV gluon-gluon fusion and  $p_T^j > 200$  GeV VBF regions are not merged with other regions. This scheme has been chosen to reduce strong anti-correlations between the measured cross sections and to keep measurements near or below 100% total uncertainty. In the likelihood, the signal yield  $N_{\text{sig}}^i$  in each category  $i$  is the sum over the yields  $N_{\text{sig},r}^i$  expected from each of the 9 regions  $r$  of phase space, where  $N_{\text{sig},r}^i = \int L dt \times \sigma_r^{\text{SM}} \times B^{\text{SM}}(H \rightarrow \gamma\gamma) \times \epsilon_r^i$  and the additional region corresponds to the difference of the cross sections for the

<sup>6</sup>The quoted theory uncertainty only accounts for the uncertainty in the acceptance. The production cross-section uncertainties are not included in the uncertainty budget. Uncertainties smaller than 0.05 are displayed as 0.0.




FIG. 10. Weighted diphoton invariant mass spectra observed in the 13 TeV data for events belonging to: (a) “untagged” categories and the “jet BSM” category, in which the expected signal is produced mainly through gluon–gluon fusion, (b) VBF categories, (c)  $VH$  categories and (d)  $t\bar{t}H$  categories. Each event is weighted by the  $\ln(1 + S_{90}/B_{90})$  ratio of the expected signal ( $S_{90}$ ) and background ( $B_{90}$ ) of the 90% signal quantile in the category it belongs to. The values of  $S_{90}$  and  $B_{90}$  used for each category are shown in Table XXVII of Appendix E. The error bars represent 68% confidence intervals of the weighted sums. The solid red curve shows the fitted signal-plus-background model when the Higgs boson mass is constrained to be  $125.09 \pm 0.24$  GeV. The background component of the fit is shown with the dotted blue curve. The signal component of the fit is shown with the solid black curve. Both the signal-plus-background and background-only curves reported here are obtained from the sum of the individual curves in each category weighted by the logarithm of unity plus the signal-to-background ratio. The bottom plot shows the residuals between the data and the background component of the fitted model.




FIG. 11. Observed negative log-profile likelihood  $\Lambda$  of the global signal strength  $\mu$ . The three likelihood contours shown correspond to all theory and experimental nuisance parameters fixed (Stat.), all experimental nuisance parameters fixed (Theo.), and with all nuisance parameters floating (Total). The intersections of the solid curves and horizontal lines at  $\Lambda = 1$  and  $\Lambda = 4$  indicate the  $1\sigma$  and  $2\sigma$  confidence intervals of the corresponding result.

$p_T^H > 200$  GeV gluon–gluon fusion and  $p_T^j > 200$  GeV VBF regions. The observed cross sections are reported in Table XI. These measurements have been defined to minimize theoretical uncertainties and are strongly dominated by experimental uncertainty, hence only the total systematic uncertainty is reported.

The evaluated cross sections including their correlations are summarized in Figs. 16 and 17. The expected Standard Model correlations can be found in Appendix H. All observed cross sections are in agreement with the Standard Model values. The Standard Model prediction is determined using the generators in Sec. IV and the theory

TABLE VI. Main systematic uncertainties  $\sigma_\mu^{\text{syst}}$  in the combined signal strength parameter  $\mu$ . The values for each group of uncertainties are determined by subtracting in quadrature from the total uncertainty the change in the 68% CL range of  $\mu$  when the corresponding nuisance parameters are fixed to their best fit values. The experimental uncertainty in the yield does not include the luminosity contribution, which is accounted for separately. The uncertainties correspond to the sources detailed in Table III.

Uncertainty group	$\sigma_\mu^{\text{syst}}$
Theory (QCD)	0.041
Theory ( $B(H \rightarrow \gamma\gamma)$ )	0.028
Theory (PDF + $\alpha_s$ )	0.021
Theory (UE/PS)	0.026
Luminosity	0.031
Experimental (yield)	0.017
Experimental (migrations)	0.015
Mass resolution	0.029
Mass scale	0.006
Background shape	0.027




FIG. 12. Summary of the signal strengths measured for the different production processes (ggH, VBF, VH and top) and globally ( $\mu_{\text{Run-2}}$ ), compared to the global signal strength measured at 7 and 8 TeV ( $\mu_{\text{Run-1}}$ ) [76]. The black and orange error bars show the total and statistical uncertainties. The signal strength  $\mu_{\text{Run-1}}$  was derived assuming the Higgs production-mode cross section based on Refs. [17,111]. Uncertainties smaller than 0.05 are displayed as 0.0. In the more recent theoretical predictions used in this analysis [7,32], the gluon–gluon fusion production-mode cross section is larger by approximately 10%. In this measurement, the  $b\bar{b}H$  contributions are scaled with ggH ( $\mu_{b\bar{b}H} = \mu_{\text{ggH}}$ ), and the  $tH$  and  $\bar{t}H$  productions are measured together ( $\mu_{\text{top}} = \mu_{tH+\bar{t}H}$ ). Associated production with Z or W bosons is assumed to be scaled by a single signal strength parameter ( $\mu_{\text{VH}} = \mu_{\text{ZH}} = \mu_{\text{WH}}$ ).

uncertainties due to missing higher-order corrections and due to the chosen PDF set are constructed as described in Sec. VII C. The largest deviation ( $1.7\sigma$ ) from the SM prediction is found in the ggH, 0 jet bin. The difference of

TABLE VII. Expected and observed significances of the VBF, VH and top quark associated production mode signal strengths.

Measurement	Expected $Z_0$	Observed $Z_0$
$\mu_{\text{VBF}}$	$2.6\sigma$	$4.9\sigma$
$\mu_{\text{VH}}$	$1.4\sigma$	$0.8\sigma$
$\mu_{\text{top}}$	$1.8\sigma$	$1.0\sigma$

TABLE VIII. Observed and expected upper limits at 95% CL on the signal strengths  $\mu_{\text{VH}}$  and  $\mu_{\text{top}}$ . The median expected limits are given for either the case when the true value of the signal strength under study is the SM value ( $\mu_i = 1$ ) or zero. The  $\pm 1\sigma$  and  $\pm 2\sigma$  intervals for the expected upper limit in the case  $\mu_i = 0$  are also reported.

Measurement	Observed	Expected Limit					
		$(\mu_i = 1)$	$(\mu_i = 0)$	$+2\sigma$	$+1\sigma$	$-1\sigma$	$-2\sigma$
$\mu_{\text{VH}}$	2.3	2.5	1.5	3.1	2.2	1.1	0.8
$\mu_{\text{top}}$	1.7	2.3	1.2	2.6	1.8	0.9	0.6






FIG. 13. Summary of asymptotic limits for the signal strengths of the associated production processes ( $VH$  and  $top$ ).




FIG. 14. Summary plot of the measured production-mode cross sections times the Higgs to diphoton branching ratio. For illustration purposes the central values have been divided by their SM expectations but no additional theory uncertainties have been added to the uncertainty of the ratio. The uncertainties in the predicted SM cross sections are shown in gray bands in the plot. The fitted value of  $\sigma_{top}$  corresponds to the sum of  $t\bar{t}H$ ,  $tHq$ , and  $tHW$  production-mode cross sections under the assumption that their relative ratios are as predicted by the SM. The  $VH$  production mode cross-section values are determined under the assumption that the ratio of the  $WH$  and  $ZH$  production-mode cross sections is as predicted by the SM and includes production from both the quark and gluon initial states. The  $b\bar{b}H$  contributions are merged with  $ggH$ .

TABLE IX. Best-fit values and uncertainties of the production-mode cross sections times branching ratio. The SM predictions [7] with their uncertainties are shown for each production process. Uncertainties smaller than 0.05 are displayed as 0.0.

Process	Result ( $ y_H  < 2.5$ )	Uncertainty [fb]				SM prediction [fb]
		Total	Statistical	Experimental	Theoretical	
$ggH$	82	$^{+19}_{-18}$	$(\pm 16)$	$^{+7}_{-6}$	$^{+5}_{-4}$	$102^{+5}_{-7}$
$VBF$	16	$^{+5}_{-4}$	$(\pm 4)$	$\pm 2$	$^{+3}_{-2}$	$8.0 \pm 0.2$
$VH$	3	$\pm 4$	$(^{+4}_{-3})$	$\pm 1$	$^{+1}_{-0}$	$4.5 \pm 0.2$
$Top$	0.7	$^{+0.9}_{-0.7}$	$(^{+0.8}_{-0.7})$	$^{+0.2}_{-0.1}$	$^{+0.2}_{-0.0}$	$1.3 \pm 0.1$

the cross sections for the  $p_T^H > 200$  GeV  $ggH$  and  $p_T^j > 200$  GeV  $VBF$  regions is found to be  $4.8^{+2.9}_{-2.7}$  fb.

Limits at 95% CL on the  $ggH + qq \rightarrow Hqq$  BSM-like ( $p_T^j > 200$  GeV) bin are set, profiling all other parameters, as shown in Table XII. Appendix G 3 provides the expected uncertainties for all quoted simplified template cross sections.

In Appendix B additional measurements are reported for a “minimally merged” set of 15 cross sections of kinematic regions defined by the requirement that the fits to expected event yields be stable even in the presence of large uncertainties or correlations.

### 5. Coupling-strength fits

Following the tree-level-motivated framework and benchmark models recommended in Ref. [17], measurements of Higgs boson coupling-strength modifiers  $\kappa_j$  are implemented. In the narrow width approximation for the Higgs boson, the cross section  $\sigma(i \rightarrow H \rightarrow \gamma\gamma)$  can be parametrized as




FIG. 15. Likelihood contours in the  $(\sigma_{ggH} \times B(H \rightarrow \gamma\gamma), \sigma_{VBF} \times B(H \rightarrow \gamma\gamma))$  plane, compared to the Standard Model prediction (red cross) for a Higgs boson mass  $m_H = 125.09$  GeV.

TABLE X. Ratios of the production-mode cross sections with respect to the ggH cross section and uncertainties are shown. The SM predictions [7] with their uncertainties are shown for each production process.

Process ( $ y_H  < 2.5$ )	Result	Uncertainty				SM prediction
		Total	Statistical	Experimental	Theoretical	
$\sigma_{\text{VBF}}/\sigma_{\text{ggH}}$	0.20	+0.10 -0.07	(+0.09 -0.06)	+0.04 -0.02	+0.04 -0.02	$0.078^{+0.005}_{-0.006}$
$\sigma_{\text{VH}}/\sigma_{\text{ggH}}$	0.04	+0.06 -0.05	(+0.06 -0.04)	+0.01 -0.01	+0.01 -0.01	$0.045^{+0.004}_{-0.005}$
$\sigma_{\text{top}}/\sigma_{\text{ggH}}$	0.009	+0.010 -0.009	(+0.010 -0.009)	+0.002 -0.001	+0.002 -0.001	$0.012^{+0.001}_{-0.002}$

$$\sigma(i \rightarrow H \rightarrow \gamma\gamma) = \frac{\sigma_i(\vec{\kappa})\Gamma^{\gamma\gamma}(\vec{\kappa})}{\Gamma_H},$$

where  $\Gamma_H$  is the total width of the Higgs boson and  $\Gamma^{\gamma\gamma}$  is the partial decay width to two photons. A set of coupling-strength modifiers,  $\vec{\kappa}$ , is introduced to parametrize possible deviations from the SM predictions of the Higgs boson coupling to SM bosons and fermions. For a given production process or decay mode  $j$ , a coupling-strength modifier  $\kappa_j$  is defined such that:

$$\kappa_j^2 = \sigma_j/\sigma_{j,\text{SM}} \quad \text{or} \quad \kappa_\gamma^2 = \Gamma^{\gamma\gamma}/\Gamma_{\text{SM}}^{\gamma\gamma},$$

where all  $\kappa_j$  values equal unity in the SM. Here, by construction, the SM cross sections and branching ratio include the best available higher-order QCD and EW corrections. This higher-order accuracy is not necessarily preserved for  $\kappa_j$  values different from unity, but the dominant higher-order QCD corrections factorize to a large extent from any rescaling of the coupling strengths and are therefore assumed to remain valid over the entire range of  $\kappa_j$  values considered.

TABLE XI. Best-fit values and uncertainties of the simplified template cross sections times branching ratio. The SM predictions [7] are shown for each region.

Measurement region ( $ y_H  < 2.5$ )	Result	Uncertainty			SM prediction
		Total	Statistical	Systematic	
ggH, 0 jet	37	$^{+16}_{-15}$	( $\pm 14$ )	$^{+6}_{-5}$ fb	$63 \pm 5$ fb
ggH, 1 jet, $p_T^H < 60$ GeV	13	$^{+13}_{-12}$	( $\pm 12$ )	$^{+5}_{-4}$ fb	$15 \pm 2$ fb
ggH, 1 jet, $60 \leq p_T^H < 120$ GeV	5	$\pm 6$	( $\pm 6$ )	$^{+2}_{-1}$ fb	$10 \pm 2$ fb
ggH, 1 jet $120 \leq p_T^H < 200$ GeV	2.8	$^{+1.7}_{-1.6}$	( $^{+1.6}_{-1.5}$ )	$^{+0.7}_{-0.5}$ fb	$1.7 \pm 0.3$ fb
ggH, $\geq 2$ jet	20	$^{+9}_{-8}$	( $\pm 8$ )	$^{+4}_{-3}$ fb	$11 \pm 2$ fb
$qq \rightarrow Hqq$ , $p_T^j < 200$ GeV	15	$^{+6}_{-5}$	( $\pm 5$ )	$^{+3}_{-2}$ fb	$10 \pm 0.5$ fb
ggH + $qq \rightarrow Hqq$ , BSM-like	2.0	$\pm 1.4$	( $\pm 1.3$ )	$\pm 0.6$ fb	$1.8 \pm 0.4$ fb
VH, leptonic	0.7	$^{+1.4}_{-1.3}$	( $^{+1.4}_{-1.2}$ )	$^{+0.4}_{-0.3}$ fb	$1.4 \pm 0.1$ fb
Top	0.7	$^{+0.8}_{-0.7}$	( $^{+0.8}_{-0.7}$ )	$^{+0.2}_{-0.1}$ fb	$1.3 \pm 0.1$ fb




FIG. 16. Summary plot of the measured simplified template cross sections times the Higgs to diphoton branching ratio. For illustration purposes the central values have been divided by their SM expectations but no additional theory uncertainties have been included in the uncertainty of the ratio due to this. The uncertainties in the predicted SM cross sections are shown in gray in the plot. The definition of the measured regions can be found in Table I. The fitted value of  $\sigma(\text{top})$  corresponds to the sum of  $t\bar{t}H$  and  $tH$  production-mode cross sections under the assumption that their relative ratios are as predicted by the SM. The  $\sigma(\text{VH, leptonic})$  cross-section values are determined under the assumption that the ratio of the  $WH$  and  $ZH$  production mode cross sections is as predicted by the SM and includes production from both the quark and gluon initial states. The  $b\bar{b}H$  contributions are merged with ggH.




FIG. 17. Observed correlations between the measured simplified template cross sections, including both the statistical and systematic uncertainties. The color indicates the size of the correlation.

Individual coupling-strength modifiers corresponding to tree-level Higgs boson couplings to different particles are introduced as well as two effective coupling-strength modifiers,  $\kappa_g$  and  $\kappa_\gamma$ , which describe the loop processes for  $ggH$  production and  $H \rightarrow \gamma\gamma$  decay. This is possible because BSM particles that might be present in these loops are not expected to appreciably change the kinematics of the corresponding process. The  $gg \rightarrow H$  and  $H \rightarrow \gamma\gamma$  loop processes can thus be studied through these effective coupling-strength modifiers, providing sensitivity to potential BSM particles in the loops. In contrast, the  $gg \rightarrow ZH$  process, which occurs at LO through box and triangular loop diagrams, is always taken into account by resolving the loop in terms of the corresponding coupling-strength

modifiers ( $\kappa_Z$  and  $\kappa_t$ ). No decays to particles other than those predicted in the SM are assumed to take place. These considerations and the limited sensitivity of the data available in this analysis lead to introducing two distinct models.

In the first model, the two parameters  $\kappa_g$  and  $\kappa_\gamma$  introduced above are tested assuming that all other couplings are as in the SM. The 68% and 95% CL two-dimensional contours of both effective couplings are shown in Fig. 18(a) and the best fit values and uncertainties are  $\kappa_g = 0.76^{+0.17}_{-0.14}$  and  $\kappa_\gamma = 1.16^{+0.14}_{-0.14}$ .

In a second model, universal coupling-strength modifiers,  $\kappa_F$  (for all fermions) and  $\kappa_V$  (for all bosons), are defined that resolve the  $gg \rightarrow H$  and  $H \rightarrow \gamma\gamma$  loops:

TABLE XII. Observed and expected upper limits at 95% CL on the simplified template cross section times the Higgs to diphoton branching ratio in the BSM sensitive phase space with  $p_T^j > 200$  GeV. The median expected limits are given for either the case when the true value of the cross section under study is SM-like ( $\sigma = \sigma_{\text{SM}}$ ) or zero. The  $\pm 1\sigma$  and  $\pm 2\sigma$  intervals for the expected upper limit ( $\sigma = 0$  fb) are also reported.

Measurement	Observed	Expected Limit ( $\sigma = \sigma_{\text{SM}}$ )	Expected Limit ( $\sigma = 0$ fb)	+2 $\sigma$	+1 $\sigma$	-1 $\sigma$	-2 $\sigma$
$ggH + qq \rightarrow Hqq$ , BSM-like	4.4 fb	4.3 fb	2.7 fb	5.3 fb	3.8 fb	2.0 fb	1.5 fb




FIG. 18. Likelihood contours in (a) the  $(\kappa_g, \kappa_\gamma)$  plane, and (b) the  $(\kappa_V, \kappa_F)$  plane, compared to the Standard Model prediction (red star) for a Higgs boson mass  $m_H = 125.09$  GeV. In (a), all coupling-strength modifiers other than  $\kappa_g$  and  $\kappa_\gamma$  are fixed to their SM value. In (b), the  $gg \rightarrow H$  and  $H \rightarrow \gamma\gamma$  loops are resolved in terms of two universal coupling-strength modifiers  $\kappa_F$  and  $\kappa_V$ , under the assumption that  $\kappa_V = \kappa_W = \kappa_Z$  and  $\kappa_F = \kappa_t = \kappa_b = \kappa_\tau = \kappa_\mu$ .




FIG. 19. The profile of negative log-likelihood  $\Lambda$  of the observed and expected coupling-strength modifier ratio  $\lambda_{tg} = \kappa_t/\kappa_g$ . The parameters  $\kappa_{g\gamma}$  and  $\lambda_{Vg}$  are also profiled within the fit. The intersections of the solid and dashed curves with the horizontal dashed line at  $\Lambda = 1$  and  $\Lambda = 4$  indicate the 1 and  $2\sigma$  confidence intervals of the observed and expected results, respectively.

$$\kappa_F = \kappa_t = \kappa_b = \kappa_\tau = \kappa_\mu,$$

$$\kappa_V = \kappa_W = \kappa_Z.$$

The 68% and 95% CL two-dimensional contours of both parameters are shown in Fig. 18(b) and the best fit values

TABLE XIII. Best-fit values and uncertainties of  $\kappa_{g\gamma}$ ,  $\lambda_{Vg}$ , and  $\lambda_{tg}$ .

Parameter	Result	Uncertainty			
		Total	Statistical	Experimental	Theoretical
$\kappa_{g\gamma}$	0.90	$\pm 0.10$	$(\pm 0.09)$	$\pm 0.04$	$^{+0.04}_{-0.03}$
$\lambda_{Vg}$	1.41	$^{+0.31}_{-0.26}$	$(^{+0.28}_{-0.23})$	$^{+0.10}_{-0.07}$	$^{+0.04}_{-0.03}$
$\lambda_{tg}$	0.8	$^{+0.4}_{-0.6}$	$(^{+0.4}_{-0.6})$	$\pm 0.1$	$^{+0.1}_{-0.0}$

and uncertainties are  $\kappa_F = 0.64^{+0.18}_{-0.14}$  and  $\kappa_V = 0.92^{+0.08}_{-0.07}$ . Due to the very limited sensitivity to  $\kappa_b$ ,  $\kappa_\tau$  and  $\kappa_\mu$ , the shown CLs would not change if these coupling-strength modifiers would be fixed to the SM expectation.

The SM prediction is found within the 68% CL contour for the first model and within the 95% CL contour for the second model.

Finally, a set of three ratios is constructed to probe the loop vertices ( $\kappa_g$ ,  $\kappa_\gamma$ ), total width ( $\kappa_H$ ), and the vector and top couplings ( $\kappa_t$  and  $\kappa_V$  respectively):  $\kappa_{g\gamma} = \kappa_g \kappa_\gamma / \kappa_H$ ,  $\lambda_{Vg} = \kappa_V / \kappa_g$ , and  $\lambda_{tg} = \kappa_t / \kappa_g$ . The parameter  $\lambda_{tg}$  is allowed to be negative to exploit the sensitivity to the relative sign from the  $tH$  and  $gg \rightarrow ZH$  processes. The expected and observed sensitivities to the relative sign are illustrated in Fig. 19. The bottom quark Yukawa coupling strength is kept fixed to the top quark Yukawa coupling strength ( $\lambda_{bg} = \lambda_{tg}$ ); this contribution is irrelevant to the  $\lambda_{tg}$  measurement as there is no sensitivity to  $b\bar{b}H$  in the analysis. All other parameters are assumed to be positive without losing generality. The inclusion of  $\kappa_H$  in the parametrization allows for non-SM decays of the Higgs boson, but this parameter is not determined directly. The best fit values of these coupling ratios are summarized in Table XIII.

## IX. MEASUREMENT OF FIDUCIAL INTEGRATED AND DIFFERENTIAL CROSS SECTIONS

The measurement of fiducial integrated and differential cross sections provides an alternative way to study the properties of the Higgs boson and to search for physics beyond the Standard Model. The fiducial volumes are defined to closely mimic the detector-level photon and object selections described in Sec. V. This reduces the model-dependence of the quoted cross sections in contrast to the per production mode simplified template cross-section measurements of Sec. VIII B 4. The cross sections are determined by correcting measured signal yields for experimental inefficiencies and



TABLE XIV. Summary of the particle-level definitions of the five fiducial integrated regions described in the text. The photon isolation  $p_T^{\text{iso},0.2}$  is defined analogously to the reconstructed-level track isolation as the transverse momentum of the system of charged particles within  $\Delta R < 0.2$  of the photon.

Objects	Definition
Photons	$ \eta  < 1.37$ or $1.52 <  \eta  < 2.37$ , $p_T^{\text{iso},0.2}/p_T^\gamma < 0.05$
Jets	anti- $k_t$ , $R = 0.4$ , $p_T > 30$ GeV, $ y  < 4.4$
Leptons, $\ell$	$e$ or $\mu$ , $p_T > 15$ GeV, $ \eta  < 2.47$ for $e$ (excluding $1.37 <  \eta  < 1.52$ ) and $ \eta  < 2.7$ for $\mu$
Fiducial region	Definition
Diphoton fiducial	$N_\gamma \geq 2$ , $p_T^{\gamma 1} > 0.35m_{\gamma\gamma} = 43.8$ GeV, $p_T^{\gamma 2} > 0.25m_{\gamma\gamma} = 31.3$ GeV
VBF-enhanced	Diphoton fiducial, $N_j \geq 2$ with $p_T^{\text{jet}} > 25$ GeV, $m_{jj} > 400$ GeV, $ \Delta y_{jj}  > 2.8$ , $ \Delta\phi_{\gamma\gamma,jj}  > 2.6$
$N_{\text{lepton}} \geq 1$	Diphoton fiducial, $N_\ell \geq 1$
High $E_T^{\text{miss}}$	Diphoton fiducial, $E_T^{\text{miss}} > 80$ GeV, $p_T^{\gamma\gamma} > 80$ GeV
$t\bar{t}H$ -enhanced	Diphoton fiducial, $(N_j \geq 4, N_{b\text{-jets}} \geq 1)$ or $(N_j \geq 3, N_{b\text{-jets}} \geq 1, N_\ell \geq 1)$

resolution effects, and by taking into account the integrated luminosity of the data. Rather than separating individual production modes, fiducial regions are defined such that they are enriched with a given production mode: Fiducial cross sections are measured in a variety of phase space regions, sensitive to for instance gluon–gluon fusion Higgs boson production, vector-boson fusion production, but also to production of the Higgs boson in association with charged leptons, top quarks, and neutrinos. Differential and double-differential cross sections are reported for variables related to the diphoton kinematics and the jet activity produced in the Higgs boson events. The observed signal yields are corrected for detector effects resulting in cross sections measured at the particle level. The full statistical and systematic correlations between measured distributions are determined and are available in HEPDATA along with the central values of the measured fiducial and differential cross sections to allow future comparisons and interpretations.

### A. Particle-level fiducial definition of the Higgs boson diphoton cross sections

The fiducial volume at particle level is defined using particles with a mean lifetime  $c\tau > 10$  mm. Only photons and leptons which do not originate from the decay of hadrons are considered.<sup>7</sup> The two highest- $p_T$  photons with  $|\eta| < 2.37$ —excluding  $1.37 < |\eta| < 1.52$ —are selected as the diphoton system. The leading (subleading) photon is required to satisfy  $p_T/m_{\gamma\gamma} > 0.35(0.25)$ , where  $m_{\gamma\gamma} = m_H = 125.09$  GeV. Furthermore, for each photon the scalar  $p_T$ -sum of charged particles with  $p_T > 1$  GeV within a cone of  $\Delta R = 0.2$  around the photon is required to be less than 5% of the photon  $p_T$ . The lepton four-momentum is

<sup>7</sup>Leptons originating from the decay of  $\tau$  leptons are only considered if the  $\tau$  lepton itself did not originate from the decay of hadrons.

defined as the combination of an electron (or muon) and all nearby photons within  $\Delta R < 0.1$  that do not originate from the decay of a hadron. Muons are required to have  $p_T > 15$  GeV and  $|\eta| < 2.7$ . Electrons are required to have  $p_T > 15$  GeV and  $|\eta| < 2.47$ , excluding the region  $1.37 < |\eta| < 1.52$ , and are rejected if the distance  $\Delta R$  to a photon with  $p_T > 15$  GeV is less than 0.4. Jets are reconstructed from all particles, excluding muons and neutrinos, using the anti- $k_t$  algorithm with a radius parameter of 0.4. Unless stated otherwise, jets are required to have  $p_T > 30$  GeV,  $|y| < 4.4$  and to be well separated from photons with  $p_T > 15$  GeV ( $\Delta R > 0.4$ ) and electrons ( $\Delta R > 0.2$ ). The acceptance for the VBF-enhanced fiducial region (introduced in Sec. IX B) is increased by loosening the  $p_T$  cut to 25 GeV. Jets are considered to originate from a  $b$ -hadron if there is a  $b$ -hadron with  $p_T > 5$  GeV within a cone of size  $\Delta R = 0.4$  around the jet.

The missing transverse momentum is defined as the vector sum of neutrino transverse momenta, for neutrinos that do not originate from the decay of a hadron. The particle-level fiducial definition is summarized in Table XIV.

### B. Fiducial integrated and differential cross sections

The cross section ( $\sigma_i$ ) in a fiducial integrated region, and the differential cross section ( $d\sigma_i/dx$ ) in a bin of variable  $x$ , are given by

$$\sigma_i = \frac{N_i^{\text{sig}}}{c_i \int L dt} \quad \text{and} \quad \frac{d\sigma_i}{dx} = \frac{N_i^{\text{sig}}}{c_i \Delta x_i \int L dt},$$

where  $N_i^{\text{sig}}$  is the number of signal events as introduced in Sec. VI C,  $\int L dt$  is the integrated luminosity of the data set,  $c_i$  is a correction factor that accounts for detector inefficiency and resolution, and  $\Delta x_i$  is the bin width. The correction factors are determined using the simulated samples discussed in Sec. IV. This bin-by-bin method




FIG. 20. The observed statistical correlations between  $p_T^{\gamma\gamma}$ ,  $N_{\text{jets}}$ ,  $m_{jj}$ ,  $|\Delta\phi_{jj}|$ , and  $p_T^{j1}$  are shown. These correlations were determined from an ensemble of 100,000 bootstrapped data sets which are each reanalyzed using an unbinned maximum-likelihood fit of the diphoton invariant mass spectrum to extract the correlations.

showed similar performance to that of the non-regularized inversion of the full migration matrix and of regularized methods [112–114] within the current statistical accuracy and systematic uncertainties.

The correction factor is  $0.75 \pm 0.03$  in the diphoton fiducial region, defined to unfold all signal events to the fiducial definition of Sec. IX A, which is dominated by the photon identification and isolation efficiency. The correction factor also accounts for migrations caused by detector energy resolution and migration in and out of the fiducial phase space due to detector effects. In addition, the correction factor removes a small fraction (0.5% for the diphoton fiducial region) of reconstructed  $H \rightarrow f\bar{f}\gamma$  Dalitz decays.<sup>8</sup>

The correction factor is different in fiducial regions defined by associated jet activity, for example, taking values of 0.66 and 0.87 for the  $t\bar{t}H$  and VBF fiducial regions defined in the next section, respectively. For the diphoton fiducial region the uncertainty in the correction factor is dominated by the theoretical modeling uncertainty. For the  $t\bar{t}H$  and VBF fiducial regions the uncertainties in

the correction factors are dominated by uncertainties associated with the knowledge of the jet energy scale and energy resolution, as well as the theoretical modeling. A more complete breakdown is given in Sec. IX E 6 and Table XVI.

The measured differential cross sections in different observables are partially statistically correlated, since they correspond to the same data set in a given fiducial region. These correlations are obtained using a random sampling with replacement method on the detector-level data, often referred to as “bootstrapping” [115]. Bootstrapped event samples are constructed from the data by assigning each event a weight pulled from a Poisson distribution with unit mean. All measured differential distributions are then reconstructed using the weighted events, and the signal yields in each bin of a differential distribution are determined using an unbinned maximum-likelihood fit of the diphoton invariant mass spectrum. The procedure is repeated with statistically independent weights and the correlation between two bins of different distributions is determined from the obtained cross sections. Figure 20 shows as an illustration the determined correlations between  $p_T^{\gamma\gamma}$ ,  $N_{\text{jets}}$ ,  $m_{jj}$ ,  $|\Delta\phi_{jj}|$ , and  $p_T^{j1}$ : the lowest  $p_T^{\gamma\gamma}$  bin, reconstructing events with a Higgs boson  $p_T$  between

<sup>8</sup>Here  $f$  denotes any fermion but the top quark.

0 and 20 GeV, is highly correlated with the zero-jet bin. The lowest  $p_T^{j1}$  bin, reconstructing events with a jet  $p_T$  between 30 and 55 GeV, is strongly correlated with the one-jet bin. And the lowest  $m_{jj}$  bin, reconstructing events with at least two jets and a dijet mass between 0 and 170 GeV, is strongly correlated with the two jet bin. The systematic correlations are obtained by fully correlating identical error sources described in Sec. VII across bins and observables to construct the corresponding systematic covariance matrix. Knowledge of these correlations allows to simultaneously analyze all fiducial regions, differential and double differential cross sections. This is illustrated later in Sec. IX E 8 with a simultaneous fit of the shown five variables of Fig. 20 to set limits on new physics contributions.

### C. Measurements of cross sections of fiducial integrated regions

Cross sections in five fiducial integrated regions are measured that target either specific Higgs boson production mechanisms or are sensitive to the presence of physics beyond the Standard Model. The selection criteria defining these regions are summarized in Table XIV and a description of each region follows:

- (1) Diphoton fiducial: This region unfolds all signal events after the selection presented in Section V.
- (2) VBF-enhanced: This region retains all events with at least two jets and with an invariant dijet mass  $m_{jj}$  of at least 400 GeV, a large rapidity separation  $|\Delta y_{jj}| > 2.8$ , and an azimuthal difference between the Higgs boson and the dijet pair of  $|\Delta\phi_{\gamma\gamma,jj}| > 2.6$ . All variables are computed using the two highest- $p_T$  jets in the event with  $p_T > 25$  GeV with matching detector-level cuts.
- (3)  $N_{\text{lepton}} \geq 1$ : This region retains events that contain at least one electron or one muon with  $p_T > 15$  GeV. For electrons the pseudo-rapidity needs to satisfy  $|\eta| < 2.47$  (excluding  $1.37 < |\eta| < 1.52$ ) and for muons  $|\eta| < 2.7$  is required. Such events are enriched in Higgs bosons produced in association with a vector boson.
- (4) High  $E_T^{\text{miss}}$ : This region retains events with missing transverse momentum  $E_T^{\text{miss}} > 80$  GeV and  $p_T^{\gamma\gamma} > 80$  GeV is defined to study  $VH$  production and possible contributions of Higgs boson production with dark matter particles. The simultaneous requirement that the Higgs boson system balances the missing transverse momentum reduces the fraction of selected events at detector level without particle-level  $E_T^{\text{miss}} > 80$  GeV.
- (5)  $t\bar{t}H$ -enhanced: This region retains events with either at least one lepton and three jets or no leptons and four jets to study Higgs boson production in association with top quarks. In addition, one of the jets needs to be identified as originating from a bottom quark.

The expected composition of Higgs boson events in the Standard Model after reconstruction and at particle level is summarized in Fig. 21. At particle level the VBF-enhanced fiducial region contains about 65% VBF and 32% ggH events. The particle-level  $N_{\text{lepton}} \geq 1$  region is dominated by  $WH$  (47%),  $t\bar{t}H$  (37%), and  $ZH$  (13%) production. The particle-level high  $E_T^{\text{miss}}$  region is populated by about equal amounts of  $WH$ ,  $ZH$ , and  $t\bar{t}H$  (32%, 30%, and 35%). Finally, the particle-level  $t\bar{t}H$ -enhanced region contains about 80%  $t\bar{t}H$  events.

The fitted invariant mass spectra for all regions are shown in Figs. 22 and 23. The results of signal-plus-background fits to these spectra is shown, displaying both the total sum and the background-only component as well as the residuals between the data and the background. In the diphoton fiducial region, the Higgs boson signal is clearly visible on the falling nonresonant background. In total,  $1491 \pm 248(\text{stat}) \pm 64(\text{syst})$  Higgs boson signal events are extracted. Clear evidence for Higgs boson production is observed in the VBF-enhanced region with




FIG. 21. The expected composition of Higgs boson events in each fiducial region (a) after the reconstruction and (b) at particle-level. Details about the reconstruction can be found in Sec. V and the definition of the particle-level fiducial volume is given in Sec. IX A.




FIG. 22. Diphoton invariant mass  $m_{\gamma\gamma}$  spectrum observed in the 2015 and 2016 data at  $\sqrt{s} = 13$  TeV for events in the diphoton fiducial region. The solid red curve shows the fitted signal-plus-background model when the Higgs boson mass is constrained to be  $125.09 \pm 0.24$  GeV. The background component of the fit is shown with the dotted blue curve. The signal component of the fit is shown with the solid black curve. The bottom plot shows the residuals between the data and the background component of the fitted model.

$117 \pm 26(\text{stat}) \pm 4(\text{syst})$  signal events, corresponding to an observed significance of 4.2 standard deviations.

The remaining three regions all show positive signal yields with large, predominantly statistical, uncertainties:  $14 \pm 11$ ,  $19 \pm 11$ ,  $6 \pm 15$  for the  $N_{\text{lepton}} \geq 1$ , high  $E_T^{\text{miss}}$ , and  $t\bar{t}H$ -enhanced fiducial regions, respectively, and the error corresponds to the sum of the statistical and systematic uncertainties.

The cross section for  $pp \rightarrow H \rightarrow \gamma\gamma$  measured in the diphoton fiducial region is

$$\sigma_{\text{fid}} = 55 \pm 9(\text{stat}) \pm 4(\text{exp}) \pm 0.1(\text{theo}) \text{ fb},$$

which is to be compared with the Standard Model prediction of  $64 \pm 2$  fb. The gluon–gluon fusion contribution to the Standard Model prediction and its uncertainty are taken to be the  $\text{N}^3\text{LO}$  QCD and NLO EW prediction of Refs. [7,24,31–34] corrected for the  $H \rightarrow \gamma\gamma$  branching ratio and the fiducial acceptance. The fiducial acceptance is defined using the POWHEG NNLOPS prediction for gluon–gluon fusion [23]. The contributions to the Standard Model prediction from the VBF,  $VH$ ,  $b\bar{b}H$ , and  $t\bar{t}H$  production mechanisms are determined using the particle-level predictions normalized with theoretical calculations as discussed in Sec. IV, and are collectively referred to as  $XH$ .

The measured cross section is compatible with the Standard Model prediction and the observed  $ggH$  coupling strength measured in Sec. VIII, as the diphoton fiducial region is dominated by gluon–gluon fusion production.

The cross section of the VBF-enhanced region is measured to be

$$\sigma_{\text{VBF-enhanced}} = 3.7 \pm 0.8(\text{stat}) \pm 0.5(\text{exp}) \pm 0.2(\text{theo}) \text{ fb},$$

which is to be compared with the Standard Model prediction of  $2.3 \pm 0.1$  fb. The gluon–gluon fusion part of the SM prediction is constructed from the POWHEG NNLOPS prediction for gluon–gluon fusion normalized with the  $\text{N}^3\text{LO}$  in QCD and NLO EW prediction of Refs. [7,24,31–34]. This prediction is labeled as “default MC” in the following and includes all theory uncertainties related to gluon–gluon fusion as discussed in Sec. VII C.

For the  $N_{\text{lepton}} \geq 1$ , high  $E_T^{\text{miss}}$ , and  $t\bar{t}H$ -enhanced fiducial regions, limits on the cross sections are reported at the 95% CL.<sup>9</sup>

Figure 24 and Table XV summarize measured cross sections of the fiducial regions and limits, and compare both to the Standard Model expectations, constructed as outlined above. The POWHEG NNLOPS prediction, without any additional corrections, is also shown. The uncertainty band is estimated using a set of scale variations and includes PDF uncertainties from eigenvector variations. The Standard Model predictions of all fiducial regions are in agreement with the corresponding measured cross sections.

#### D. Measurements of cross sections of inclusive and exclusive jet multiplicities

The production of Higgs bosons in association with jets is sensitive to the theoretical modeling in QCD and to the contribution of different Higgs boson production mechanisms. In the SM, events with zero or one jet are dominated by gluon–gluon fusion production. In events with two jets the contributions from VBF and  $VH$  production modes become more important. Higgs boson production in association with top quarks ( $t\bar{t}H$ ) can be probed in events with the highest jet multiplicities. In BSM scenarios, the jet multiplicity distribution is sensitive to new heavy particles coupling to the Higgs boson and vector bosons. For the measurements presented here, jet multiplicity bins with zero, one, two, and at least three jets with  $p_T$  larger than 30 GeV and absolute rapidity  $|y| < 4.4$  are defined. In addition, jet multiplicity bins with a  $p_T$  larger than 50 GeV are defined for zero, one, or at least two jets. The measured cross sections are compared to a range of predictions of gluon–gluon fusion production:

<sup>9</sup>The quoted CL values were obtained using the unfolded cross sections and their corresponding uncertainties assuming Gaussian errors.






FIG. 23. Diphoton invariant mass  $m_{\gamma\gamma}$  spectra observed in the 2015 and 2016 data at  $\sqrt{s} = 13$  TeV for events in the (a) VBF-enhanced, (b)  $N_{\text{lepton}} \geq 1$ , (c) high  $E_T^{\text{miss}}$ , and (d)  $t\bar{t}H$ -enhanced fiducial regions. The solid red curve shows the fitted signal-plus-background model when the Higgs boson mass is constrained to be  $125.09 \pm 0.24$  GeV. The background component of the fit is shown with the dotted blue curve. The signal component of the fit is shown with the solid black curve. The bottom plot shows the residuals between the data and the background component of the fitted model.

- (i) The parton-level N<sup>3</sup>LO QCD and NLO EW prediction of Refs. [7,24,31–34]. This prediction is shown for the inclusive zero-jet cross section.
- (ii) The parton-level JVE + N<sup>3</sup>LO prediction of Ref. [116], which includes NNLL resummation in QCD of the  $p_T$  of the leading jet which is matched to

the N<sup>3</sup>LO total cross section. This prediction is shown for the inclusive one-jet cross section.

- (iii) The parton-level STWZ-BLPTW predictions of Refs. [99,101], which include NNLL' + NNLO resummation for the  $p_T$  of the leading jet in QCD, combined with a NLL' + NLO resummation in QCD




FIG. 24. The measured cross sections or cross-section upper limits of the diphoton, VBF-enhanced,  $N_{\text{lepton}} \geq 1$ , high  $E_T^{\text{miss}}$ , and  $t\bar{t}H$ -enhanced fiducial regions are shown. The intervals on the vertical axis each represent one of these fiducial regions. The data are shown as filled (black) circles. The error bar on each measured cross section represents the total uncertainty in the measurement, with the systematic uncertainty shown as a dark gray rectangle. Each cross section limit is shown at the 95% confidence level. The measured cross sections are compared to a range of predictions and a detailed description of each prediction can be found in the text. All comparisons include the SM predictions arising from VBF,  $VH$ ,  $t\bar{t}H$ , and  $b\bar{b}H$ , which are collectively labeled as  $XH$ .

for the subleading jet.<sup>10</sup> The numerical predictions for  $\sqrt{s} = 13$  TeV are taken from Ref. [7]. This prediction is shown for the inclusive zero-, one- and two-jet cross sections as well as for the exclusive zero- and one-jet cross sections.

- (iv) The parton-level NNLOJET prediction of Refs. [117,118] is a fixed-order NNLO prediction in QCD for inclusive  $H$  + one-jet production. This prediction is shown for the inclusive one-, two-jet, and three-jet cross sections as well as for the exclusive one- and two-jet cross sections.
- (v) The parton-level GOSAM prediction of Refs. [119, 120], which provides the fixed-order loop contributions accurate at NLO in QCD in the inclusive  $H$  + zero-jet,  $H$  + one-jet,  $H$  + two-jet, and  $H$  + three-jet regions. The real-emission contributions at fixed order in QCD are provided by SHERPA [64]. This prediction is shown for the inclusive one-, two-jet, and three-jet cross sections as well as for the exclusive one- and two-jet cross sections.
- (vi) The default MC prediction (POWHEG NNLOPS normalized with the  $N^3\text{LO}$  in QCD and NLO EW cross

section) introduced in Sec. IX C. This prediction is shown for all measured inclusive and exclusive jet cross sections.

- (vii) The POWHEG NNLOPS prediction which is already described in Sec. IV. This prediction is shown for all measured inclusive and exclusive jet cross sections.
- (viii) The SHERPA (MEPS@NLO) prediction of Refs. [64,65, 120–129] is accurate to NLO in QCD in the inclusive  $H$  + zero-jet,  $H$  + one-jet,  $H$  + two-jet, and  $H$  + three-jet regions and includes top-quark mass effects. The one-loop corrections are incorporated from GoSAM [119,120] and the different jet multiplicity regions are merged using the MEPS@NLO multijet merging technique. This prediction is shown for all measured inclusive and exclusive jet cross sections.
- (ix) The MG5\_AMC@NLO prediction of Refs. [46,105], which includes up to two jets at NLO accuracy using the FxFx merging scheme [106]. The central merging scale is taken to be 30 GeV. The generated events are passed to PYTHIA8 [29] to provide parton showering and hadronization to create the full final state (without underlying event). This prediction is shown for all measured inclusive and exclusive jet cross sections.

All predictions but NNLOJET and SHERPA (MEPS@NLO) use the NNLO PDF set following the PDF4LHC15 recommendations. The NNLOJET prediction uses the CT14 NNLO PDF set [130] and SHERPA (MEPS@NLO) uses the NNPDF3.0 PDF set [47]. GOSAM, SHERPA (MEPS@NLO), and NNLOJET apply the kinematic selection on the final-state photons. For all other predictions, the fiducial acceptance is determined using POWHEG NNLOPS. The cross sections of all parton-level predictions are multiplied with isolation correction factors to account for the efficiency of the fiducial photon isolation criterion. The additional uncertainties in the isolation correction are determined by studying multiple event generators and/or event generator tunes, and are included in the uncertainty bands of the parton-level predictions. No correction factors nor additional uncertainties to account for the impact of hadronization and the underlying event activity are applied, so the theory uncertainties in the parton-level predictions may be incomplete, but example values for such corrections and their uncertainties can be found in Table XXIV in Appendix D. All other acceptance and correction factors along with their associated uncertainties can also be found in Appendix D.

No  $K$ -factors are applied to the predictions and the contributions from  $XH$  are also included in the comparison using the corresponding generators and cross sections described in Section IV.

Figure 25(a) shows exclusive and inclusive zero-, one- and two-jet cross sections and the inclusive three-jet cross section for jets defined with  $p_T > 30$  GeV. Figure 25(b) shows the exclusive zero- and one- and the inclusive two-jet cross section with  $p_T > 50$  GeV. The measured cross

<sup>10</sup>The prime indicates that the leading contributions from  $N^3\text{LL}$  (resp. NNLL) are included along with the full NNLL (resp. NLL) corrections.

TABLE XV. The measured cross sections in the diphoton, VBF-enhanced,  $N_{\text{lepton}} \geq 1$ , high  $E_T^{\text{miss}}$ , and  $t\bar{t}H$ -enhanced fiducial regions. The gluon–gluon fusion contribution to the Standard Model prediction of the diphoton fiducial region is taken to be the N<sup>3</sup>LO prediction of Refs. [7,24,31–34] corrected for the  $H \rightarrow \gamma\gamma$  branching ratio and the fiducial acceptance. The gluon–gluon fusion contribution to the Standard Model for all the other regions is taken from the POWHEG NNLOPS prediction normalized with the N<sup>3</sup>LO prediction and includes all theory uncertainties related to gluon–gluon fusion as discussed in Sec. VII C. The contributions to the Standard Model prediction from VBF,  $VH$ ,  $t\bar{t}H$  and  $b\bar{b}H$  production mechanisms are determined using the particle-level predictions described in Sec. IV normalized with theoretical calculations.

Fiducial region	Measured cross section	SM prediction	
Diphoton fiducial	$55 \pm 9(\text{stat}) \pm 4(\text{exp}) \pm 0.1(\text{theo})$ fb	$64 \pm 2$ fb	[N <sup>3</sup> LO + $XH$ ]
VBF-enhanced	$3.7 \pm 0.8(\text{stat}) \pm 0.5(\text{exp}) \pm 0.2(\text{theo})$ fb	$2.3 \pm 0.1$ fb	[default MC + $XH$ ]
$N_{\text{lepton}} \geq 1$	$\leq 1.39$ fb 95% CL	$0.57 \pm 0.03$ fb	[default MC + $XH$ ]
High $E_T^{\text{miss}}$	$\leq 1.00$ fb 95% CL	$0.30 \pm 0.02$ fb	[default MC + $XH$ ]
$t\bar{t}H$ -enhanced	$\leq 1.27$ fb 95% CL	$0.55 \pm 0.06$ fb	[default MC + $XH$ ]




FIG. 25. Cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of inclusive and exclusive jet multiplicities for jets with (a)  $p_T > 30$  GeV and (b)  $p_T > 50$  GeV. The data are shown as filled (black) circles. The vertical error bar on each data point represents the total uncertainty in the measured cross section and the shaded (gray) band is the systematic component. The measured differential cross sections are compared to a range of predictions and details can be found in the text. The width of the bands of each prediction reflects the total theoretical uncertainty. The small contribution from VBF,  $VH$ ,  $t\bar{t}H$ , and  $b\bar{b}H$  is also shown as a (green) histogram and denoted by  $XH$ .

sections are in agreement with the Standard Model predictions, although there is a  $1.5\sigma$  deficit in the  $N_{\text{jet}} = 0$  cross section for jets defined with  $p_T > 30$  GeV. As shown in Fig. 20, there is a sizeable positive correlation between zero-jet and low- $p_T^{\gamma\gamma}$  events, and a similar deficit is observed there (cf. Sec. IX E 1).

### E. Measurements of differential and double-differential cross sections

Eleven fiducial differential cross sections are measured that characterize the Higgs boson production kinematics, the kinematics of jets produced in association with the Higgs boson, the spin and  $CP$  quantum numbers of the Higgs boson and variables sensitive to the VBF production mechanism. In addition, two double-differential cross sections are reported. The measurement of seven additional variables can be found in Appendix C.

#### 1. Measurements of cross sections probing the Higgs boson production kinematics

Measuring the transverse momentum of the diphoton system,  $p_T^{\gamma\gamma}$ , probes the perturbative QCD modeling of the  $ggH$  production mechanism which is mildly sensitive to the bottom- and charm-quark Yukawa couplings [12]. The distribution at high transverse momentum is sensitive to new heavy particles coupling to the Higgs boson and to the top-quark Yukawa coupling. The rapidity distribution of the diphoton system,  $|y_{\gamma\gamma}|$ , is also sensitive to the modeling of the  $ggH$  production mechanism. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of  $p_T^{\gamma\gamma}$  and  $|y_{\gamma\gamma}|$  are shown in Fig. 26. The chosen bin widths are a compromise between retaining a sufficiently significant signal and providing spectra with good granularity. Each bin is chosen such that it retains an expected significance of at least two standard deviations, estimated using the POWHEG NNLOPS and additional predictions described in Sec. IV as well as using a fit to  $m_{\gamma\gamma}$  sidebands. The measurements are compared to several predictions of gluon–gluon fusion:




FIG. 26. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $p_T^{\gamma\gamma}$  and (b)  $|y_{\gamma\gamma}|$  are shown and compared to the SM expectations.

- (i) The default MC prediction (POWHEG NNLOPS normalized with the N<sup>3</sup>LO in QCD and NLO EW cross section) introduced in Sec. IX C.
- (ii) HRES [131,132], which provides predictions at NNLO with  $p_T^H$  resummation at NNLL and differentially in  $p_T^{\gamma\gamma}$ . Finite top-, bottom-, and charm-quark masses are included at NLO accuracy. The renormalization and factorization scales are chosen to be  $\frac{1}{2}\sqrt{m_H^2 + (p_T^H)^2}$ , and the two resummation scales are chosen to be  $m_H/2$  and  $2m_b$ .
- (iii) RADISH+NNLOJET [133], which provides predictions using a  $p_T^H$  resummation to NNLL and matching to the one-jet NNLO differential spectrum from NNLOJET [117,118]. The shown  $p_T^{\gamma\gamma}$  RADISH+NNLOJET prediction does include corrections from the finite top and bottom quark masses.
- (iv) SCETLIB+MCFM8 which provides predictions at NNLO + NNLL <sub>$\phi$</sub>  accuracy derived by applying a resummation of the virtual corrections to the gluon form factor [134,135] and differentially in  $|y_{\gamma\gamma}|$  and  $|\cos\theta^*|$ .<sup>11</sup> The underlying NNLO predictions are obtained using MCFM8 with zero-jettiness subtractions [136,137].

No additional  $K$ -factors are applied to the predictions, which all use the NNLO PDF set following the PDF4LHC15 recommendations, and the fiducial acceptance for RADISH+NNLOJET is determined using POWHEG NNLOPS. The SCETLIB+MCFM8 and HRES predictions include the

<sup>11</sup>The subscript  $\phi$  refers to the fact that the applied resummation is to the gluon form factor.

kinematic acceptance and are corrected and apply correction factors accounting for the photon isolation efficiency as described in Sec. IX D. As also mentioned in Sec. IX D, no correction factors to account for the impact of hadronization and the underlying-event activity are used. The SM prediction shows a slight excess at low transverse momentum and low rapidity, and shows a slight deficit at large transverse momentum. The slightly harder Higgs boson transverse momentum shown in Fig. 26 is consistent with the ATLAS Run 1 measurements in both the  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4\ell$  decay channels [9,138] and the measured zero-jet cross section. The Standard Model prediction is in agreement with the measured distributions.

## 2. Measurements of cross sections probing the jet kinematics

The transverse momentum and absolute rapidity of the leading jet,  $p_T^{j_1}$  and  $|y_{j_1}|$ , as well as the transverse momentum and absolute rapidity of the subleading jet,  $p_T^{j_2}$  and  $|y_{j_2}|$ , are sensitive to the theoretical modeling and to the relative contributions of the different Higgs boson production mechanisms. The transverse momentum distribution of the leading jet probes the emission of energetic quarks and gluons. In events with two jets, the contributions of VBF and  $VH$  productions become more important. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of  $p_T^{j_1}$ ,  $|y_{j_1}|$ ,  $p_T^{j_2}$ , and  $|y_{j_2}|$  are shown in Fig. 27. The chosen bin widths are a compromise between keeping migrations between bins small whilst retaining enough statistical power to measure the differential spectra. The measured  $p_T^{j_1}$  spectrum shown in Fig. 27(a) is compared to the default






FIG. 27. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $p_T^{j1}$ , (b)  $|y_{j1}|$ , (c)  $p_T^{j2}$ , and (d)  $|y_{j2}|$  are shown and compared to the SM expectations. The data and theoretical predictions are presented in the same way as in Fig. 26. In addition, the NNLOJET and SCETLIB(STWZ) predictions, the NNLOJET prediction, and the SHERPA (MEPS@NLO) and GoSAM predictions, described in the text, are displayed in (a), (b) and (c+d), respectively.

MC prediction as introduced in the previous section as well as to the NNLOJET and SCETLIB(STWZ) [99,135] predictions. Both the NNLOJET and SCETLIB predictions are corrected using isolation correction factors to account for the impact of the isolation efficiency. In addition, the NNLOJET prediction is corrected for the kinematic acceptance and the uncertainties in these corrections is included in the uncertainty bands of both NNLOJET and SCETLIB. The first bin of the leading jet  $p_T$  spectrum represents zero-jet events that do

not contain any jet with  $p_T > 30$  GeV. The predicted  $p_T$  distributions slightly exceed the measured distribution at low transverse momentum and all show a slight deficit at large transverse momentum. Both are compatible with the observed slightly harder Higgs boson transverse momentum distribution. The measured  $|y_{j1}|$  distribution shown in Fig. 27(b) is compared to the default MC and the NNLOJET predictions: Both show a slight excess at low rapidity. In Fig. 27(c) the measured subleading jet  $p_T$




FIG. 28. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $|\cos \theta^*|$  and (b)  $\Delta\phi_{jj}$  are shown and compared to the SM expectations. The data and theoretical predictions are presented in the same way as in Fig. 26. In addition, the SCETLIB+MCFM8 prediction and the SHERPA (MEPS@NLO) and GoSAM predictions, described in the text, are displayed in (a) and (b), respectively.

distribution is shown. The first bin of  $p_T^{j_2}$  represents one-jet events that do not contain two or more jets with  $p_T > 30$  GeV. The measured distribution is compared to the default MC, SHERPA (MEPS@NLO), and GoSAM predictions, as introduced in Section IX D. Finally, in Fig. 27(d) the subleading jet rapidity distribution,  $|y_{j_2}|$ , is shown and compared to the expectation from the default MC, SHERPA (MEPS@NLO), and GoSAM predictions. The SM predictions are in agreement with the measured distributions and no significant deviations are seen.

### 3. Measurements of cross sections probing spin and CP

The absolute value of the cosine of the angle between the beam axis and the photons in the Collins–Soper frame [11] of the Higgs boson,  $|\cos \theta^*|$ , can be used to study the spin of the Higgs boson. The azimuthal angle between the two leading jets,  $\Delta\phi_{jj}$ ,<sup>12</sup> in events containing two or more jets is sensitive to the charge conjugation and parity properties of the Higgs boson interactions with gluons and weak bosons in the gluon–gluon fusion and the VBF production channels, respectively [139–142].

The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of  $|\cos \theta^*|$  and  $\Delta\phi_{jj}$  are shown in Fig. 28. For a scalar particle  $|\cos \theta^*|$ , shows a strong drop around 0.6 due to the fiducial requirement on the photon system, whereas

for a spin-2 particle, an enhancement would be present in precisely this region. The charge conjugation and parity properties of the Higgs boson are encoded in the azimuthal angle between the jets: For example, in gluon–gluon fusion, its distribution for a  $CP$ -even coupling has a dip at  $\pm\frac{\pi}{2}$  and present peaks at 0 and  $\pm\pi$ , whereas for a purely  $CP$ -odd coupling it would present as peaks at  $\pm\frac{\pi}{2}$  and dips at 0 and  $\pm\pi$ . For VBF the SM prediction for  $\Delta\phi_{jj}$  is approximately constant with a slight rise towards  $\Delta\phi_{jj} = \pm\pi$ . Any additional anomalous  $CP$ -even or  $CP$ -odd contribution to the interaction between the Higgs boson and weak bosons would manifest itself as an additional oscillatory component, and any interference between the SM and anomalous couplings can produce distributions peaked at either  $\Delta\phi_{jj} = 0$  or  $\Delta\phi_{jj} = \pm\pi$  [139,141,142]. The shape of the distribution is therefore sensitive to the relative contribution of gluon–gluon fusion and vector-boson fusion, as well as to the tensor structure of the interactions between the Higgs boson and gluons or weak bosons. This is exploited in Sec. IX E 8 to set limits on new physics contributions. To quantify the structure of the azimuthal angle between the two jets, a ratio is defined as

$$A_{|\Delta\phi_{jj}|} = \frac{\sigma(|\Delta\phi_{jj}| < \frac{\pi}{3}) - \sigma(\frac{\pi}{3} < |\Delta\phi_{jj}| < \frac{2\pi}{3}) + \sigma(|\Delta\phi_{jj}| > \frac{2\pi}{3})}{\sigma(|\Delta\phi_{jj}| < \frac{\pi}{3}) + \sigma(\frac{\pi}{3} < |\Delta\phi_{jj}| < \frac{2\pi}{3}) + \sigma(|\Delta\phi_{jj}| > \frac{2\pi}{3})},$$

<sup>12</sup>To preserve the sign information, the azimuthal angles of the jets are ordered according to the jet with the highest rapidity. This definition of  $\Delta\phi_{jj}$  is invariant under a redefinition of the ordering by choosing the opposite beam axis, as explained in Ref. [139].

which is motivated by a similar ratio presented in Ref. [141]. The measured ratio in data as determined by measuring  $|\Delta\phi_{jj}|$  in three bins is

$$A_{|\Delta\phi_{jj}|}^{\text{meas}} = 0.45_{-0.24}^{+0.18}(\text{stat})_{-0.11}^{+0.10}(\text{syst}).$$

This value can be compared to the SM prediction from the default MC simulation. The predicted value is  $A_{|\Delta\phi_{jj}|}^{\text{SM}} = 0.44 \pm 0.01$ , consistent with the measured ratio.

In summary, the measured  $|\cos\theta^*|$  and  $\Delta\phi_{jj}$  distributions are consistent with Standard Model predictions for a  $CP$ -even scalar particle.

#### 4. Cross sections probing the VBF production mode

The distribution of the dijet rapidity separation,  $|\Delta y_{jj}|$ , the azimuthal angle between the dijet and diphoton systems,  $|\Delta\phi_{\gamma\gamma,jj}|$ , and the invariant mass of the leading and subleading jets,  $m_{jj}$  for events with two or more jets are sensitive to the differences between the gluon–gluon fusion and VBF production mechanisms. In vector-boson fusion, the  $t$ -channel exchange of a  $W/Z$  boson typically results in two moderate- $p_T$  jets that are well separated in rapidity. Furthermore, quark/gluon radiation in the rapidity interval between the two jets is suppressed in the VBF process when compared to the gluon–gluon fusion process, because there is no color flow between the two jets. The  $|\Delta\phi_{\gamma\gamma,jj}|$  distribution for VBF production is therefore expected to be steeper and more peaked towards  $|\Delta\phi_{\gamma\gamma,jj}| = \pi$  than for gluon–gluon fusion.

The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of  $|\Delta y_{jj}|$ ,  $|\Delta\phi_{\gamma\gamma,jj}|$ , and  $m_{jj}$  are shown for events with at least two jets with  $p_T > 30$  GeV in Fig. 29. These variables are used to discriminate between gluon–gluon fusion and the VBF production of the Higgs boson and enter the multivariate classifier introduced in Sec. VIII A 4 that defines the categories used for the simplified template cross-section and coupling measurements. The measured distributions are in agreement to the default MC, SHERPA (MEPS@NLO), and the GoSAM predictions. The accuracy of the fixed-order parton-level prediction from GoSAM breaks down in the lowest bin of  $\pi - |\Delta\phi_{\gamma\gamma,jj}|$  and the measured cross section moderately exceeds the SM predictions at high  $m_{jj}$  values.

#### 5. Double-differential cross sections

The double-differential cross section for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of  $p_T^{\gamma\gamma}$  and  $N_{\text{jet}}$ , for jets with  $p_T > 30$  GeV, and  $p_T^{\gamma\gamma}$  and  $|\cos\theta^*|$  are shown in Fig. 30. These cross sections are sensitive to the modeling of the Higgs boson kinematic, its production mechanisms, and its spin- $CP$  properties. Both double-differential cross sections are in agreement with the Standard Model expectation.

#### 6. Impact of systematic uncertainties on results

A summary of the uncertainties in the measured cross sections of the fiducial regions are shown in Table XVI. As an example concerning the differential measurements, a

breakdown of the systematic uncertainties in the differential cross sections as a function of  $p_T^{\gamma\gamma}$  and  $N_{\text{jet}}$  is shown in Fig. 31. The measurements are dominated by the statistical uncertainties. For the systematic uncertainties, the uncertainty in the fitted signal yield, due to the background modeling and the photon energy resolution, is typically more important than the uncertainty in the correction factor due to the theoretical modeling. The jet energy scale and resolution uncertainties become increasingly important for high-jet multiplicities and in the  $t\bar{t}H$ - and VBF-enhanced phase space.

#### 7. Compatibility of measured distributions with the Standard Model

The compatibility between the measured distributions and the Standard Model is tested by comparing the first and second moments of the measured distributions. Figure 32 shows the first and second moments (mean and RMS) of the distributions and compares them to the moments of the default MC prediction, as calculated from the measured and predicted cross-section bins. The theory uncertainties are constructed as outlined in Sec. VII C. The measured Higgs boson transverse momentum has somewhat higher first and second moments than the Standard Model prediction, which is consistent with the previous observations [9,138]. The leading-jet  $p_T$  spectrum shows the same feature. In addition a  $\chi^2$  test is carried out for all distributions reported in Sec. IX: The resulting  $p$ -values are reported in Table XVII, which confirms that within the current uncertainties the data are in agreement with the SM predictions.

#### 8. Search for anomalous Higgs-boson interactions using an effective field theory approach

The strength and tensor structure of the Higgs-boson interactions can be investigated using an effective field theory approach, which introduces additional  $CP$ -even and  $CP$ -odd interactions that change the event rates and the kinematic properties of the Higgs boson and associated jet spectra from those in the Standard Model. The parameters of the effective field theory are probed using a fit to five of the most relevant differential cross sections. The effective Lagrangian of Ref. [143] is used which adds dimension-six operators of the strongly interacting light Higgs formulation [13] to the Standard Model interactions. The  $H \rightarrow \gamma\gamma$  differential cross sections are mainly sensitive to the operators that affect the Higgs boson interactions with gauge bosons and the relevant terms in the Lagrangian can be specified by

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \bar{c}_g \mathcal{O}_g + \bar{c}_{HW} \mathcal{O}_{HW} + \bar{c}_{HB} \mathcal{O}_{HB} \\ & + \tilde{c}_g \tilde{\mathcal{O}}_g + \tilde{c}_{HW} \tilde{\mathcal{O}}_{HW} + \tilde{c}_{HB} \tilde{\mathcal{O}}_{HB}, \end{aligned}$$




FIG. 29. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $|\Delta y_{jj}|$ , (b)  $\pi - |\Delta\phi_{\gamma\gamma,jj}|$ , and (c)  $m_{jj}$  are shown and compared to the SM expectations. The data and theoretical predictions are presented in the same way as in Fig. 26. In addition, the SHERPA (MEPS@NLO) and GoSAM predictions are shown for all three cross sections.

where  $\tilde{c}_i$  and  $\tilde{c}_i$  are dimensionless Wilson coefficients specifying the strength of the new  $CP$ -even and  $CP$ -odd interactions, respectively, and the dimension-six operators  $\mathcal{O}_i$  and  $\tilde{\mathcal{O}}_i$  are those described in Refs. [143,144]. In the SM, all of the Wilson coefficients are equal to zero. The  $\mathcal{O}_g$  and  $\tilde{\mathcal{O}}_g$  operators introduce new interactions between the Higgs boson and two gluons and can be probed through the gluon–gluon fusion Higgs production mechanism. The  $\mathcal{O}_{HW}$  and  $\tilde{\mathcal{O}}_{HW}$  operators introduce new  $HWW$ ,  $HZZ$ , and  $HZ\gamma$  interactions.

The  $HZZ$  and  $HZ\gamma$  interactions are also impacted by  $\mathcal{O}_{HB}$  and  $\tilde{\mathcal{O}}_{HB}$ . The  $\mathcal{O}_{HW}$ ,  $\tilde{\mathcal{O}}_{HW}$ ,  $\mathcal{O}_{HB}$ , and  $\tilde{\mathcal{O}}_{HB}$  operators can be probed through vector-boson fusion and associated production. Other operators in the full effective Lagrangian of Ref. [143] can also modify Higgs-boson interactions but are not considered here due to the lack of sensitivity of the  $H \rightarrow \gamma\gamma$  decay channel. Combinations of some of the  $CP$ -even operators have been constrained using global fits to experimental data from LEP and the LHC [143,145,146].






FIG. 30. The double-differential cross section for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $p_T^{\gamma\gamma}$  and  $N_{\text{jet}}$ , for jets with  $p_T > 30$  GeV, and (b)  $p_T^{\gamma\gamma}$  and  $|\cos \theta^*|$  separating the two regions of  $|\cos \theta^*| < 0.5$  and  $|\cos \theta^*| > 0.5$  from each other. The data and theoretical predictions are presented in the same way as in Fig. 26.

The effective Lagrangian has been implemented in FEYNRULES [144].<sup>13</sup> Parton-level event samples are produced for specific values of Wilson coefficients by interfacing the universal file output from FEYNRULES to the MG5\_AMC@NLO event generator [46]. Higgs bosons are produced via gluon–gluon fusion with up to two additional partons in the final state using leading-order matrix elements.

The generated events are passed to PYTHIA8 [29] to provide parton showering, hadronization and underlying event and the zero-, one- and two-parton events are merged using the MLM matching scheme [148] to create the full final state. Event samples containing a Higgs boson produced either in association with a vector boson or via vector-boson fusion are produced using leading-order matrix elements and passed through the PYTHIA8 generator. For each production mode, the Higgs boson mass is set to 125 GeV and events are generated using the NNPDF2.3LO PDF set [47] and the A14 parameter set [48]. All other Higgs boson production modes are assumed to occur as predicted by the SM.

Event samples are produced for different values of a given Wilson coefficient. The particle-level differential cross sections are produced using RIVET [149]. The PROFESSOR method [150] is used to interpolate between these samples,

<sup>13</sup>The implementation in Ref. [144] involves a redefinition of the gauge boson propagators that results in unphysical amplitudes unless certain physical constants are also redefined. The original implementation did not include the redefinition of these physical constants. However, the impact of redefining the physical constants is found to be negligible on the predicted cross sections across the range of Wilson coefficients studied. The relative change in the predicted Higgs boson cross sections as functions of the different Wilson coefficients is also found to agree with that predicted by the Higgs characterization framework [147], with less than 2% variation across the parameter ranges studied.

for each bin of each distribution, to provide a parameterization of the effective Lagrangian prediction. The parameterization function is determined using 11 samples when studying a single Wilson coefficient, whereas 25 samples are used when studying two Wilson coefficients simultaneously. As the Wilson coefficients enter the effective Lagrangian in a linear fashion, second-order polynomials are used to predict the cross sections in each bin. The method was validated by comparing the differential cross sections obtained with the parameterization function to the predictions obtained with dedicated event samples generated at the specific point in parameter space.

The model implemented in FeynRules fixes the Higgs boson width to be that of the SM,  $\Gamma_H = 4.07 \text{ MeV}$  [17]. The cross sections are scaled by  $\Gamma_H/(\Gamma_H + \Delta\Gamma)$ , where  $\Delta\Gamma$  is the change in partial widths due to a specific choice of Wilson coefficient. The change in partial widths is determined for each Higgs coupling using the partial-width calculator in MG5\_AMC@NLO and normalized to reproduce the SM prediction from HDECAY [18].

The leading-order predictions obtained from MG5\_AMC@NLO are reweighted to account for higher-order QCD and electroweak corrections to the SM process, assuming that these corrections factorize from the new physics effects. The differential cross section as a function of variable  $X$  for a specific choice of Wilson coefficient,  $c_i$ , is given by

$$\frac{d\sigma}{dX} = \sum_j \left( \frac{d\sigma_j}{dX} \right)^{\text{ref}} \cdot \left( \frac{d\sigma_j}{dX} \right)_{c_i}^{\text{MG5}} / \left( \frac{d\sigma_j}{dX} \right)_{c_i=0}^{\text{MG5}},$$

where the summation  $j$  is over the different Higgs boson production mechanisms, “MG5” labels the interpolated MG5\_AMC@NLO prediction and “ref” labels a reference

TABLE XVI. The expected uncertainties, expressed in percent, in the cross sections measured in the diphoton fiducial, VBF-enhanced,  $N_{\text{lepton}} \geq 1$ ,  $t\bar{t}H$ -enhanced, and high  $E_T^{\text{miss}}$  regions. The fit systematic uncertainty includes the effect of the photon energy scale and resolution, and the impact of the background modeling on the signal yield. The theoretical modeling uncertainty is defined as the envelope of the signal composition, the modeling of Higgs boson transverse momentum and rapidity distribution, and the uncertainty of parton shower and the underlying event (labeled as “UE/PS”) as described in Sec. VII D.

Source	Uncertainty in fiducial cross section				
	Diphoton	VBF-enhanced	$N_{\text{lepton}} \geq 1$	$t\bar{t}H$ -enhanced	High $E_T^{\text{miss}}$
Fit (stat.)	17%	22%	72%	176%	53%
Fit (syst.)	6%	9%	27%	138%	13%
Photon energy scale & resolution	4.3%	3.5%	3.1%	10%	4.1%
Background modeling	4.2%	7.8%	26.7%	138%	12.2%
Photon efficiency	1.8%	1.8%	1.8%	1.8%	1.9%
Jet energy scale/resolution	...	8.9%	...	4.5%	6.9%
$b$ -jet flavor tagging	...	...	...	3%	...
Lepton selection	...	...	0.7%	0.2%	...
Pileup	1.1%	2.9%	1.3%	2.5%	2.5%
Theoretical modeling	0.1%	4.5%	4.0%	8.1%	31%
Signal composition	0.1%	4.5%	3.1%	8.1%	25%
Higgs boson $p_T^H$ & $ y_H $	0.1%	0.9%	0.2%	0.7%	0.1%
UE/PS	...	0.3%	0.7%	1.1%	31%
Luminosity	3.2%	3.2%	3.2%	3.2%	3.2%
Total	18%	26%	77%	224%	63%




FIG. 31. The relative size of systematic uncertainties associated with the signal extraction, the correction factors (experimental and theoretical modeling) and the luminosity on the differential cross sections are shown as a function of (a)  $p_T^{\gamma\gamma}$  and (b)  $N_{\text{jets}}$ . The statistical uncertainty associated with the signal extraction is also shown as a gray band. For completeness, the relevant components of the uncertainties in the correction factors are shown as a function of (c)  $p_T^{\gamma\gamma}$  and (d)  $N_{\text{jets}}$ .




FIG. 32. (a) The ratio of the first moment (mean) of each differential distribution predicted by the Standard Model to that observed in the data. The SM moment is calculated by using the default MC distributions for gluon–gluon fusion and the other production mechanisms. (b) The ratio of the second moment (RMS) of each differential distribution predicted by the Standard Model to that observed in the data. The intervals on the vertical axes each represent one of the differential distributions. The band for the theoretical prediction represents the corresponding uncertainty in that prediction (see text for details). The error bar on the data represents the total uncertainty in the measurement, with the gray band representing only the systematic uncertainty.

sample for SM Higgs boson production. For the reference sample the default MC simulation is used.

The measured differential cross sections of  $p_T^{\gamma\gamma}$ ,  $N_{\text{jet}}$ ,  $m_{jj}$ ,  $|\Delta\phi_{jj}|$ , and  $p_T^{j1}$  are compared in Fig. 33(a) to the SM hypothesis and to two non-SM hypotheses, specified by  $\tilde{c}_g = 2 \times 10^{-4}$  and  $\tilde{c}_{HW} = 0.05$ , respectively. The new  $CP$ -odd gluon–gluon fusion operator results in a large increase in rate and the additional  $CP$ -even  $WH$  operator leads to a

larger number of Higgs boson with sizeable  $p_T$  and an increased number of zero-jet events.

The ratios of the expected differential cross sections to the SM predictions for some representative values of the Wilson coefficients are shown in Fig. 33(b). The impact of the  $\tilde{c}_g$  and  $\tilde{c}_g$  coefficients are presented for the gluon–gluon fusion production: it displays a large change in the overall cross-section normalization. The  $\tilde{c}_g$  coefficient also changes the shape of the  $\Delta\phi_{jj}$  distribution, which is expected from consideration of the tensor structure of  $CP$ -even and  $CP$ -odd interactions [139,141]. In contrast, the impact of the  $\tilde{c}_{HW}$  and  $\tilde{c}_{HW}$  coefficients are presented specifically for the VBF + VH production channel: one expects large shape changes in all of the studied distributions and the  $\Delta\phi_{jj}$  distribution is known to discriminate between  $CP$ -even and  $CP$ -odd interactions in the VBF production channel [140].

Limits on Wilson coefficients are set by constructing a likelihood function

$$\mathcal{L} = \frac{1}{\sqrt{(2\pi)^k |C|}} \exp \left( -\frac{1}{2} (\vec{\sigma}_{\text{data}} - \vec{\sigma}_{\text{pred}})^T C^{-1} (\vec{\sigma}_{\text{data}} - \vec{\sigma}_{\text{pred}}) \right),$$

where  $\vec{\sigma}_{\text{data}}$  and  $\vec{\sigma}_{\text{pred}}$  are  $k$ -dimensional vectors from the measured and predicted differential cross sections of the five analyzed observables,  $C = C_{\text{stat}} + C_{\text{syst}} + C_{\text{theo}}$  is the  $k \times k$  total covariance matrix defined by the sum of the statistical, systematic and theoretical covariances, and  $|C|$  denotes its determinant. The theory covariance is constructed as outlined in Sec. VII C and includes no

TABLE XVII. Probabilities from a  $\chi^2$  test for the comparison between data and the default SM prediction.

Distribution	Default MC prediction
$p_T^{\gamma\gamma}$	51%
$ y_{\gamma\gamma} $	57%
$p_T^{j1}$	32%
$ y_{j1} $	66%
$p_T^{j2}$	61%
$ y_{j2} $	56%
$ \cos \theta^* $	47%
$\Delta\phi_{jj}$	64%
$ \Delta y_{jj} $	53%
$ \Delta\phi_{\gamma\gamma,jj} $	43%
$m_{jj}$	54%
$N_{\text{jet}} (p_T > 30 \text{ GeV})$	56%
$N_{\text{jet}} (p_T > 50 \text{ GeV})$	19%




FIG. 33. (a) The measured differential cross sections as a function of  $p_T^{\gamma\gamma}$ ,  $N_{\text{jet}}$ ,  $m_{jj}$ ,  $|\Delta\phi_{jj}|$ , and  $p_T^{j1}$  are compared to the SM hypothesis and two non-SM hypotheses with  $\bar{c}_g = 1 \times 10^{-4}$  and  $\bar{c}_{HW} = 0.05$ , respectively. (b) Ratios of differential cross sections, as predicted for specific choices of Wilson coefficient, to the differential cross sections predicted by the SM: the impact of non-zero  $\bar{c}_g$  and  $\bar{c}_g$  is shown relative to the SM ggH prediction, while the impact of nonzero  $\bar{c}_{HW}$  and  $\bar{c}_{HW}$  is shown relative to the SM VBF + VH prediction.

additional uncertainty to account for the factorization assumption in Sec. IX E 8. Based on this likelihood, one can construct a  $\chi^2$  test to test the compatibility of the five distributions with the SM and a probability of 93% is found. In what follows, the likelihood function is numerically maximized to determine  $\mathcal{L}_{\text{max}}$  and confidence limits for one or several Wilson coefficients are determined via

$$1 - \text{CL} = \int_{-2 \ln \mathcal{L}(c_i) + 2 \ln \mathcal{L}_{\text{max}}}^{\infty} dx f(x),$$

with  $\mathcal{L}(c_i)$  denoting the likelihood value evaluated for a given Wilson coefficient value  $c_i$ , and  $f(x)$  denoting the distribution of the test statistic. The coverage of the confidence limit is determined using ensembles of pseudo-experiments. Form factors are sometimes used to regularize the change of the cross section above a momentum scale  $\Lambda_{\text{FF}}$ . This was investigated by reweighting the VBF + VH samples using form-factor predictions from VBFNLO [151]. The impact on the  $\bar{c}_{HW}$  and  $\bar{c}_{HW}$  limits is negligible for  $\Lambda_{\text{FF}} > 1$  TeV.



TABLE XVIII. Observed allowed ranges at 95% CL for the  $\bar{c}_g$  and  $\bar{c}_{HW}$  Wilson coefficients and the  $CP$ -conjugate coefficients. Limits on  $\bar{c}_g$  and  $\tilde{c}_g$  are each derived with all other Wilson coefficients set to zero. Limits on  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$  are derived with  $\bar{c}_{HB} = \bar{c}_{HW}$  and  $\tilde{c}_{HB} = \tilde{c}_{HW}$ , respectively.

Coefficient	Observed 95% CL limit	Expected 95% CL limit
$\bar{c}_g$	$[-0.8, 0.1] \times 10^{-4} \cup [-4.6, -3.8] \times 10^{-4}$	$[-0.4, 0.5] \times 10^{-4} \cup [-4.9, -4.1] \times 10^{-4}$
$\tilde{c}_g$	$[-1.0, 0.9] \times 10^{-4}$	$[-1.4, 1.3] \times 10^{-4}$
$\bar{c}_{HW}$	$[-5.7, 5.1] \times 10^{-2}$	$[-5.0, 5.0] \times 10^{-2}$
$\tilde{c}_{HW}$	$[-0.16, 0.16]$	$[-0.14, 0.14]$

In Table XVIII, the observed and expected 95% CL limits for four Wilson coefficient fits are given. The limit for  $\bar{c}_g$  is derived by fixing all other Wilson coefficients to zero. This additional interaction can interfere with the corresponding SM interaction and destructive interference causes the gluon–gluon fusion production-mode cross section to be zero at  $\bar{c}_g \sim -2.2 \times 10^{-4}$ . The  $\tilde{c}_g$  coefficient is also derived after setting all Wilson coefficients to zero. Due to the  $CP$  conjugate structure of the interaction, no interference with the SM process is possible. The 95% CL limit for  $\bar{c}_{HW}$  is obtained after setting  $\bar{c}_{HB} = \bar{c}_{HW}$  to ensure that the partial width for  $H \rightarrow Z\gamma$  is unchanged from the SM prediction (Values of  $|\bar{c}_{HW} - \bar{c}_{HB}| > 0.03$  lead to a very large decay rate for the  $H \rightarrow Z\gamma$  process that is contradicted by the experimental constraints reported by ATLAS [152,153]) and setting all other Wilson coefficients to zero. Finally, the 95% CL limit for  $\tilde{c}_{HW}$  is given after setting  $\tilde{c}_{HB} = \tilde{c}_{HW}$  to ensure a SM decay rate for  $H \rightarrow Z\gamma$  and all other Wilson coefficients to zero. The observed limits are improved by about a factor of two compared to the Run 1 analysis of Ref. [14].




FIG. 34. The observed 68% (dark) and 95% (light) confidence level regions from the simultaneous fit to the  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$  Wilson coefficients. The values of  $\bar{c}_{HB}$  and  $\tilde{c}_{HB}$  are set to be equal to  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$ , respectively, and all other Wilson coefficients are set to zero, except for  $\bar{c}_{HB}$  and  $\tilde{c}_{HB}$  which are set to be equal to  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$ , respectively. The SM expectation at (0,0) is also shown, together with the Run-1 confidence regions reported in Ref. [14].

Figure 34 shows the 68% and 95% confidence regions obtained from scanning  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$  simultaneously, with the other two Wilson coefficients set to  $\bar{c}_{HB} = \bar{c}_{HW}$  and  $\tilde{c}_{HB} = \tilde{c}_{HW}$ . All other Wilson coefficients are fixed at zero. The  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$  Wilson coefficients produce large shape changes in all distributions, as shown in Fig. 33, and the obtained limits are strongest when fitting all five distributions simultaneously. The shape of the observed 68% confidence regions thus results from both shape and yield differences between data and expectations: the operators proportional to  $\bar{c}_{HW}$  can destructively interfere with the SM contributions, a negative value of  $\bar{c}_{HW}$  reduces the overall predicted cross section in the zero-jet and the lowest  $m_{jj}$  bins, where deficits are observed in the data. The operators proportional to  $\tilde{c}_{HW}$  can only increase the cross section from its SM value and can increase the predicted cross sections in the higher jet bins and the tails of the distributions (cf. Fig. 33). If only shape information is used to constrain the Wilson coefficients, the reported limits on  $\bar{c}_{HW}$  and  $\tilde{c}_{HW}$  weaken by about 20% and 50%, respectively. As also shown in Fig. 34, these results display significant improvements on similar limits obtained from the Run-1 analysis [14]. All reported results assume that QCD effects and new physics effects factorize. This assumption cannot be avoided with the current state-of-the-art implementation of the effective Lagrangian of Ref. [143]. The full statistical and systematic correlations between measured distributions and all measured fiducial and differential cross sections are available in HEPDATA to allow future interpretations with better models.

## X. SUMMARY CONCLUSIONS

Measurements of Higgs boson cross sections in the Higgs boson to diphoton decay channel are performed using  $pp$  collision data recorded by the ATLAS experiment at the LHC. The data were taken at a center-of-mass energy of  $\sqrt{s} = 13$  TeV and correspond to an integrated luminosity of  $36.1 \text{ fb}^{-1}$ . All measurements assume a Higgs boson mass of  $125.09 \pm 0.24$  GeV. The measured signal strength relative to the Standard Model expectation is found to be:

$$\mu = 0.99^{+0.15}_{-0.14}.$$

Signal strengths of the main production modes are measured separately via event reconstruction categories that are designed to be sensitive to the specific production modes. They are found to be:

$$\begin{aligned}\mu_{\text{ggH}} &= 0.81^{+0.19}_{-0.18}, & \mu_{\text{VBF}} &= 2.0^{+0.6}_{-0.5}, \\ \mu_{\text{VH}} &= 0.7^{+0.9}_{-0.8}, & \mu_{\text{top}} &= 0.5^{+0.6}_{-0.6}.\end{aligned}$$

The total uncertainties of both the global and the production mode signal strengths is dominated by their respective statistical uncertainties. The global signal strength measurement improves on the precision of the previous ATLAS measurement in the diphoton channel by a factor of two [154]. The ggH (VBF) signal strength is measured to be  $1\sigma$  below ( $2\sigma$  above) the Standard Model expectation. The precision of the coupling-strength measurements involving top quarks improves by about a factor of three compared to the previous ATLAS measurement in the diphoton channel [154]. These improvements result from a combination of the larger Higgs boson sample collected at  $\sqrt{s} = 13$  TeV, from the use of multivariate techniques to target the VBF, VH, and top-quark associated production modes more efficiently, from the improved precision of the ggH Standard Model theory predictions, and from a significant reduction of some of the experimental uncertainties such as the photon energy resolution. Production mode cross-section measurements for a Higgs boson of rapidity  $|y_H| < 2.5$  for gluon–gluon fusion, vector-boson fusion, and Higgs boson production in association with vector bosons or a top quark pair are reported:

$$\begin{aligned}\sigma_{\text{ggH}} &= 82^{+19}_{-18} \text{ fb}, & \sigma_{\text{VBF}} &= 16^{+5}_{-4} \text{ fb}, \\ \sigma_{\text{VH}} &= 3 \pm 4 \text{ fb}, & \sigma_{\text{top}} &= 0.7^{+0.9}_{-0.7} \text{ fb}.\end{aligned}$$

These values can be compared to the Standard Model expectations of

$$\begin{aligned}\sigma_{\text{ggH}}^{\text{SM}} &= 102^{+5}_{-7} \text{ fb}, & \sigma_{\text{VBF}}^{\text{SM}} &= 8 \pm 0.2 \text{ fb}, \\ \sigma_{\text{VH}}^{\text{SM}} &= 5 \pm 0.2 \text{ fb}, & \sigma_{\text{top}}^{\text{SM}} &= 1.3 \pm 0.1 \text{ fb},\end{aligned}$$

and show a similar level of agreement as that obtained with the coupling-strength measurements.

Nine measurements of so-called simplified template cross sections, designed to measure the different Higgs boson production processes in specific regions of phase space, are reported:

$$\begin{aligned}\sigma(\text{ggH}, 0 \text{ jet}) &= 37^{+16}_{-15} \text{ fb}, \\ \sigma(\text{ggH}, 1 \text{ jet}, p_T^H < 60 \text{ GeV}) &= 13^{+13}_{-12} \text{ fb}, \\ \sigma(\text{ggH}, 1 \text{ jet}, 60 \leq p_T^H < 120 \text{ GeV}) &= 5 \pm 6 \text{ fb}, \\ \sigma(\text{ggH}, 1 \text{ jet}, 120 \leq p_T^H < 200 \text{ GeV}) &= 2.8^{+1.7}_{-1.6} \text{ fb}, \\ \sigma(\text{ggH}, \geq 2 \text{ jet}) &= 20^{+9}_{-8} \text{ fb}, \\ \sigma(qq \rightarrow Hqq, p_T^j < 200 \text{ GeV}) &= 15^{+6}_{-5} \text{ fb}, \\ \sigma(\text{ggH} + qq \rightarrow Hqq, \text{BSM-like}) &= 2.0 \pm 1.4 \text{ fb}, \\ \sigma(\text{VH}, \text{leptonic}) &= 0.7^{+1.4}_{-1.3} \text{ fb}, \\ \sigma(\text{top}) &= 0.7^{+0.8}_{-0.7} \text{ fb}.\end{aligned}$$

All reported results show agreement with the Standard Model expectation.

Higgs boson coupling-strength modifiers are reported and two models are investigated: the first one reports results on effective coupling-strength modifiers for Higgs boson production in gluon–gluon fusion and decay,  $\kappa_g$  and  $\kappa_\gamma$ , respectively. They are found to be

$$\kappa_g = 0.76^{+0.17}_{-0.14}, \quad \text{and} \quad \kappa_\gamma = 1.16^{+0.14}_{-0.14}.$$

The second model resolves the Higgs boson production and decay loops in terms of the more fundamental fermionic and vector boson couplings under the assumption of universal coupling-strength modifiers for all fermions and vector bosons, namely  $\kappa_V$  and  $\kappa_F$ , respectively. They are found to be

$$\kappa_V = 0.92^{+0.08}_{-0.07}, \quad \text{and} \quad \kappa_F = 0.64^{+0.18}_{-0.14}.$$

Fiducial cross-section measurements are reported for a Higgs boson decaying into two isolated photons with transverse momentum greater than 35% and 25% of the diphoton invariant mass (corresponding to a photon  $p_T$  of 43.8 GeV and 31.3 GeV), and with  $|\eta| < 2.37$ , excluding the region of  $1.37 < |\eta| < 1.52$ . The total fiducial cross section is measured to be

$$\sigma_{\text{fid}} = 55 \pm 9(\text{stat}) \pm 4(\text{exp}) \pm 0.1(\text{theo}) \text{ fb},$$

and is in agreement with the Standard Model expectation of  $64 \pm 2$  fb. Additional cross sections in fiducial regions probing Higgs boson production from vector-boson fusion or associated with large missing transverse momentum, leptons or top quarks are reported. The cross section for the VBF-enhanced region is measured to be

$$\sigma_{\text{VBF-enhanced}} = 3.7 \pm 0.8(\text{stat}) \pm 0.5(\text{exp}) \pm 0.2(\text{theo}) \text{ fb},$$

which is to be compared with the Standard Model prediction of  $2.3 \pm 0.1$  fb. The larger measured cross section is consistent with the VBF signal-strength

measurement reported above, if one scales the expected SM contributions from VBF (about 65%) and ggH (about 35%) in this fiducial region with the corresponding measured signal strengths. For the remaining fiducial regions, limits at 95% CL are reported

$$\sigma_{N_{\text{lepton}} \geq 1} < 1.39 \text{ fb}, \quad \sigma_{\text{High} E_T^{\text{miss}}} < 1.00 \text{ fb},$$

$$\sigma_{\text{H}H\text{-enhanced}} < 1.27 \text{ fb},$$

which can be compared with the Standard Model expectations of  $0.57 \pm 0.03 \text{ fb}$ ,  $0.30 \pm 0.02 \text{ fb}$ , and  $0.55 \pm 0.06 \text{ fb}$ , respectively.

The fiducial cross sections for different jet multiplicities is reported and compared to several theoretical predictions. Eleven differential cross sections and two double-differential cross sections are reported for events belonging to the inclusive diphoton fiducial region, as a function of kinematic variables of the diphoton system or of jets produced in association with the Higgs boson. The reported cross sections are sensitive to the Higgs boson production kinematics, the jet kinematics, the spin and  $CP$  quantum numbers of the Higgs boson, and the VBF production mechanism. All measured differential cross sections are compared to predictions and no significant deviation from the Standard Model expectation is observed. The full statistical and systematic correlations between measured distributions are determined and are available in HEPDATA along with the central values of the measured fiducial and differential cross sections to allow future comparisons and interpretations.

The strength and tensor structure of the Higgs boson interactions is investigated using five differential variables and an effective Lagrangian, which introduces additional  $CP$ -even and  $CP$ -odd interactions. No significant new physics contributions are observed and the reported 68% and 95% limits on such contributions have improved by a factor of two in comparison to the previous ATLAS measurement.

The measurements presented in this paper lay the foundation for further studies. All reported results are statistically limited and their precision will further improve with the full data set to be recorded during Run 2 of the LHC.

## ACKNOWLEDGMENTS

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently. We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTC, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; CONICYT, Chile; CAS, MOST and NSFC, China; COLCIENCIAS, Colombia; MSMT CR, MPO CR and VSC CR, Czech

Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS, CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF, and MPG, Germany; GSRT, Greece; RGC, Hong Kong SAR, China; ISF, I-CORE and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW and NCN, Poland; FCT, Portugal; MNE/IFA, Romania; MES of Russia and NRC KI, Russian Federation; JINR; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DST/NRF, South Africa; MINECO, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TAEK, Turkey; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, the Canada Council, CANARIE, CRC, Compute Canada, FQRNT, and the Ontario Innovation Trust, Canada; EPLANET, ERC, ERDF, FP7, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex and Idex, ANR, Région Auvergne and Fondation Partager le Savoir, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF; BSF, GIF and Minerva, Israel; BRF, Norway; CERCA Programme Generalitat de Catalunya, Generalitat Valenciana, Spain; the Royal Society and Leverhulme Trust, United Kingdom. The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [155].

## APPENDIX A: SIMPLIFIED TEMPLATE CROSS-SECTION FRAMEWORK

As introduced in Sec. IB, this paper includes cross-section measurements using the so called “stage-1” of the simplified template cross-section framework [7,8]. In the full stage-1 proposal, template cross sections are defined in 31 regions of phase space with  $|y_H| < 2.5$ . These regions have been chosen to minimize the dependence on theoretical uncertainties and isolate possible BSM effects, while maximizing the experimental sensitivity.

The 31 regions of particle-level phase space corresponding to the stage 1 of the template cross-section approach are the following [7,8]:

- (i) Gluon–gluon fusion (11 regions). Gluon–gluon fusion events, together with  $gg \rightarrow ZH$  events followed by hadronic decays of the  $Z$  boson, are split according to the number of jets in the event in 0, 1, and  $\geq 2$ -jet events. Jets are reconstructed from all

stable particles<sup>14</sup> with lifetime greater than 10 ps using the anti- $k_t$  algorithm [77] with a jet radius parameter  $R = 0.4$  and must have  $p_T > 30$  GeV. The region containing two or more jets is split into two, with one of the two subregions (“VBF-like”) containing events with a topology similar to vector-boson fusion events (invariant mass of the leading- $p_T$  jet pair  $m_{jj} > 400$  GeV, and rapidity separation between the two jets  $|\Delta y_{jj}| > 2.8$ ). The one-jet and non-VBF-like two-jet regions are further split according to the transverse momentum of the Higgs boson in “low” (0–60 GeV), “medium” (60–120 GeV), “high” (120–200 GeV) and “BSM” (>200 GeV) regions. The VBF-like events are further split into exclusive two-jetlike and inclusive three-jetlike events through a requirement on the transverse momentum  $p_T^{Hjj}$  of the system formed by the two photons and the two leading- $p_T$  jets ( $p_T^{Hjj} < 25$  GeV or  $p_T^{Hjj} > 25$  GeV, respectively). The separation between events with zero, one, or two or more jets probes perturbative QCD predictions. Events containing a very high transverse momentum Higgs-boson of more than 200 GeV are sensitive to BSM contributions, such as those from loop-induced amplitudes mediated by hypothetical particles heavier than the top-quark.

- (ii) Vector-boson fusion (5 regions). Vector-boson fusion events, and  $VH$  events followed by hadronic  $V$ -boson decays, are first split according to the  $p_T$  of the leading jet. Events that contain at least one jet with a transverse momentum greater than 200 GeV, which are sensitive to BSM contributions, are measured separately in a “VBF BSM” category. The remaining events are separated into VBF-like events,  $VH$ -like events, and events that have a  $ggH$ -like topology (referred to as “Rest”). VBF-like events satisfy the same  $m_{jj}$  and  $|\Delta y_{jj}|$  requirements as for the gluon–gluon fusion VBF-like category and are similarly split into “two-jet” and “ $\geq 3$ -jet” events by requiring  $p_T^{Hjj} < 25$  GeV or  $p_T^{Hjj} > 25$  GeV, respectively.  $VH$ -like events are selected by requiring that they have at least two jets and an invariant mass of  $60 \text{ GeV} < m_{jj} < 120 \text{ GeV}$ .
- (iii) Associated production with vector bosons decaying to leptons (11 regions).  $VH$  events are first split according to their production mode ( $q\bar{q}' \rightarrow WH$ ,  $q\bar{q} \rightarrow ZH$ , or  $gg \rightarrow ZH$ ). Events are separated further into regions of the vector boson transverse momentum  $p_T^V$ , and of jet multiplicity. For  $gg \rightarrow ZH$ , two regions are defined with  $p_T^V$  (“low”: 0–150 GeV, and “high”: >150 GeV). The “high- $p_T^V$ ”  $gg \rightarrow ZH$

region is further split into zero-jet and  $\geq 1$ -jet regions. Regions sensitive to BSM contributions with  $p_T^V > 250$  GeV are defined for the  $q\bar{q} \rightarrow VH$  production modes and two further  $p_T^V$  regions are defined (“low”: 0–150 GeV, and “high”: 150–250 GeV). The “high- $p_T^V$ ”  $q\bar{q} \rightarrow VH$  region is finally split into zero-jet and  $\geq 1$ -jet regions.

- (iv) Associated production with top and bottom quarks (4 regions).  $t\bar{t}H$ ,  $t$ -channel  $tH$ ,  $W$ -associated  $tH$ , and  $b\bar{b}H$  events are classified according to their production mode, with no further separation into specific regions of phase space.

Table XIX summarizes the acceptances for each of the stage-1 STXS  $gg \rightarrow H$  regions, and for five  $qq \rightarrow Hqq$  regions, split into their VBF,  $WH$ , and  $ZH$  respective

TABLE XIX. The SM acceptances of stage-1 STXS regions useful to the results presented in this paper. For the  $gg \rightarrow H$  regions each acceptance is relative to inclusive  $gg \rightarrow H$  production; for all other regions, each acceptance is relative to the inclusive process shown at the top of the column. All regions require  $|y_H| < 2.5$ .

$gg \rightarrow H$ regions	0-jet	1-jet	$\geq 2$ -jet
$p_T^H < 60$ GeV	0.562	0.134	0.025
$60 \text{ GeV} \leq p_T^H < 120$ GeV	...	0.093	0.038
$120 \text{ GeV} \leq p_T^H < 200$ GeV	...	0.015	0.020
$p_T^H \geq 200$ GeV	...	0.003	0.009
VBF-like			
$p_T^{Hjj} < 25$ GeV	...	...	0.006
$p_T^{Hjj} \geq 25$ GeV	...	...	0.007
$qq \rightarrow Hqq$ regions	VBF	$q\bar{q}' \rightarrow WH$	$q\bar{q} \rightarrow ZH$
$p_T^j \geq 200$ GeV	0.043	0.027	0.029
$p_T^j < 200$ GeV			
VH-like	0.023	0.189	0.224
Rest	0.556	0.368	0.363
VBF-like			
$p_T^{Hjj} < 25$ GeV	0.235	0.002	0.002
$p_T^{Hjj} \geq 25$ GeV	0.074	0.007	0.008
VH, leptonic region	$gg \rightarrow ZH$	$q\bar{q}' \rightarrow WH$	$q\bar{q} \rightarrow ZH$
	0.289	0.286	0.265
Top region	$t\bar{t}H$	$t$ -channel $tH$	$W$ -associated $tH$
	0.987	0.921	0.989
Beauty region			$b\bar{b}H$
			0.945

<sup>14</sup>The Higgs boson is treated as stable and consequently its decay products are removed from the jet finding.



contributions. The table also lists the summed acceptance for the 11  $VH$  leptonic regions, separately for the  $gg \rightarrow ZH$ ,  $q\bar{q}' \rightarrow WH$  and  $q\bar{q} \rightarrow ZH$  processes. Finally, the acceptances are shown for the rarer production processes:  $t\bar{t}H$ ,  $t$ -channel  $tH$ ,  $W$ -associated  $tH$ , and  $b\bar{b}H$ . All STXS regions require  $|y_H| < 2.5$  and are determined using the samples summarized in Table II.

TABLE XX. The kinematic regions of the stage 1 of the simplified template cross sections, along with the intermediate (minimally merged set of) regions used for the measurements presented in this appendix. The  $VH$ -like, VBF-like, and rest regions are defined as in Table I and Appendix A. All regions require  $|y_H| < 2.5$ . The leading jet transverse momentum is denoted by  $p_T^j$ . In total, the cross sections for fifteen kinematic regions are measured.

Process	Measurement region	Particle-level stage 1 region
$ggH + gg \rightarrow Z(\rightarrow qq)H$	0-jet	0-jet
	1-jet, $p_T^H < 60$ GeV	1-jet, $p_T^H < 60$ GeV
	1-jet, $60 \leq p_T^H < 120$ GeV	1-jet, $60 \leq p_T^H < 120$ GeV
	1-jet, $120 \leq p_T^H < 200$ GeV	1-jet, $120 \leq p_T^H < 200$ GeV
	1-jet, $p_T^H > 200$ GeV	1-jet, $p_T^H > 200$ GeV
	$\geq 2$ -jet, $p_T^H < 60$ GeV	$\geq 2$ -jet, $p_T^H < 60$ GeV
	$\geq 2$ -jet, $60 \leq p_T^H < 120$ GeV	$\geq 2$ -jet, $60 \leq p_T^H < 120$ GeV
	$\geq 2$ -jet, $120 \leq p_T^H < 200$ GeV	$\geq 2$ -jet, $120 \leq p_T^H < 200$ GeV
	$\geq 2$ -jet, $p_T^H > 200$ GeV	$\geq 2$ -jet, $p_T^H > 200$ GeV
	VBF-like	VBF-like, $p_T^{Hjj} < 25$ GeV
$qq' \rightarrow Hqq'$ (VBF + $VH$ )	$p_T^j < 200$ GeV, VBF-like	$p_T^j < 200$ GeV, VBF-like, $p_T^{Hjj} < 25$ GeV
	$p_T^j < 200$ GeV, $VH$ + Rest	$p_T^j < 200$ GeV, VBF-like, $p_T^{Hjj} \geq 25$ GeV
	$p_T^j > 200$ GeV, BSM-like	$p_T^j < 200$ GeV, $VH$ -like
		$p_T^j < 200$ GeV, Rest
$VH$ (leptonic decays)	$VH$ leptonic	$p_T^j > 200$ GeV
		$q\bar{q} \rightarrow ZH$ , $p_T^Z < 150$ GeV
		$q\bar{q} \rightarrow ZH$ , $150 < p_T^Z < 250$ GeV, 0-jet
		$q\bar{q} \rightarrow ZH$ , $150 < p_T^Z < 250$ GeV, $\geq 1$ -jet
		$q\bar{q} \rightarrow ZH$ , $p_T^Z > 250$ GeV
		$q\bar{q} \rightarrow WH$ , $p_T^W < 150$ GeV
		$q\bar{q} \rightarrow WH$ , $150 < p_T^W < 250$ GeV, 0-jet
		$q\bar{q} \rightarrow WH$ , $150 < p_T^W < 250$ GeV, $\geq 1$ -jet
		$q\bar{q} \rightarrow WH$ , $p_T^W > 250$ GeV
		$gg \rightarrow ZH$ , $p_T^Z < 150$ GeV
Top-associated production	top	$gg \rightarrow ZH$ , $p_T^Z > 150$ GeV, 0-jet
		$gg \rightarrow ZH$ , $p_T^Z > 150$ GeV, $\geq 1$ -jet
		$t\bar{t}H$
$b\bar{b}H$	merged w/ $ggH$	$tHW$
		$tHq$
		$b\bar{b}H$

TABLE XXI. Best-fit values and uncertainties of the simplified template cross sections times branching ratio, as defined in Table XX. The SM predictions [7] are shown for each region.

Measurement region ( $ y_H  < 2.5$ )	Result	Uncertainty			SM prediction
		Total	Statistical	Systematic	
$ggH$ , 0 jet	38	$^{+16}_{-15}$	$(\pm 14)$	$^{+6}_{-5}$ fb	$63 \pm 5$ fb
$ggH$ , 1 jet, $p_T^H < 60$ GeV	23	$^{+14}_{-13}$	$(\pm 13)$	$^{+5}_{-4}$ fb	$15 \pm 2$ fb
$ggH$ , 1 jet, $60 \leq p_T^H < 120$ GeV	11	$\pm 8$	$(\pm 7)$	$^{+3}_{-2}$ fb	$10 \pm 2$ fb

(Table continued)



TABLE XXI. (Continued)

Measurement region ( $ y_H  < 2.5$ )	Result	Uncertainty			SM prediction
		Total	Statistical	Systematic	
$ggH$ , 1 jet, $120 \leq p_T^H < 200$ GeV	4.0	$^{+2.1}_{-1.9}$	$(\pm 1.8)$	$^{+0.9}_{-0.6}$ fb	$1.7 \pm 0.3$ fb
$ggH$ , 1 jet, $p_T^H \geq 200$ GeV	2.6	$^{+1.6}_{-1.2}$	$(^{+1.3}_{-1.1})$	$^{+0.8}_{-0.5}$ fb	$0.4 \pm 0.1$ fb
$ggH$ , $\geq 2$ jet, $p_T^H < 60$ GeV	0	$\pm 8$	$(\pm 8)$	$^{+3}_{-2}$ fb	$3 \pm 1$ fb
$ggH$ , $\geq 2$ jet, $60 \leq p_T^H < 120$ GeV	12	$^{+8}_{-7}$	$(\pm 7)$	$^{+3}_{-2}$ fb	$4 \pm 1$ fb
$ggH$ , $\geq 2$ jet, $120 \leq p_T^H < 200$ GeV	7.9	$^{+3.5}_{-3.4}$	$(\pm 3.3)$	$^{+1.1}_{-0.9}$ fb	$2.3 \pm 0.6$ fb
$ggH$ , $\geq 2$ jet, $p_T^H \geq 200$ GeV	2.6	$^{+1.6}_{-1.4}$	$(^{+1.5}_{-1.4})$	$^{+0.6}_{-0.5}$ fb	$1.0 \pm 0.3$ fb
$ggH$ , VBF – like	6.2	$^{+5.0}_{-4.5}$	$(\pm 4.1)$	$\pm 1.2$ fb	$1.5 \pm 0.3$ fb
$qq \rightarrow Hqq$ , VBF – like	3.8	$^{+2.5}_{-2.3}$	$(^{+2.2}_{-2.0})$	$\pm 1.2$ fb	$2.7 \pm 0.2$ fb
$qq \rightarrow Hqq$ , VH + Rest	-19	$\pm 22$	$(^{+21}_{-20})$	$^{+6}_{-7}$ fb	$7.7 \pm 0.4$ fb
$qq \rightarrow Hqq$ , $p_T^j \geq 200$ GeV	-3.2	$^{+1.9}_{-2.0}$	$(\pm 1.7)$	$^{+0.7}_{-0.9}$ fb	$0.5 \pm 0.1$ fb
VH, leptonic	0.7	$^{+1.4}_{-1.2}$	$(^{+1.4}_{-1.2})$	$^{+0.4}_{-0.3}$ fb	$1.4 \pm 0.1$ fb
Top	0.7	$^{+0.8}_{-0.7}$	$(^{+0.8}_{-0.7})$	$^{+0.2}_{-0.1}$ fb	$1.3 \pm 0.1$ fb




FIG. 35. Summary plot of the measured simplified template cross sections times the Higgs to diphoton branching ratio, as defined in Table XX. For illustration purposes, the central values have been divided by their SM expectations but no additional SM uncertainties have been folded into the measurement. The uncertainties in the SM predicted cross sections are shown in gray in the plot. The fitted value of  $\sigma(\text{top})$  corresponds to the sum of the  $t\bar{t}H$ ,  $tHq$ , and  $tHW$  production-mode cross sections under the assumption that their relative ratios are as predicted by the SM. The  $\sigma(\text{VH, leptonic})$  cross-section values are determined under the assumption that the ratio of the  $WH$  and  $ZH$  production mode cross sections is as predicted by the SM and includes production from both the quark and gluon initial states. The  $b\bar{b}H$  contributions are merged with  $ggH$ .

### APPENDIX C: ADDITIONAL UNFOLDED DIFFERENTIAL CROSS SECTIONS

This appendix presents additional measurements and comparisons to theoretical predictions to those discussed in Sec. IX E.

Figure 36 shows differential cross sections as a function of  $p_T^{\gamma\gamma}$ , the orthogonal component of the diphoton momentum when projected on the axis given by the difference of the 3-momenta of the two photons, as well as  $|\Delta y_{\gamma\gamma}|$ , the rapidity separation of the two photons.

Figure 37 shows differential cross sections as a function of  $H_T$ , the scalar sum of all reconstructed jets in a given event with  $p_T > 30$  GeV, the absolute value of the azimuthal difference  $|\Delta\phi_{jj}|$  between the leading and subleading jet in events with at least two jets, and the vectorial sum of the transverse momentum of the diphoton system and the leading and subleading jet system,  $p_T^{\gamma jj}$ , in events with at least two jets.

Figure 38 displays measurements of the beam-thrust-like variables  $\tau_{C,j1}$  and  $\sum \tau_{C,j}$ . For a given jet,  $\tau$  is defined by

$$\tau = \frac{m_T}{2 \cosh y^*}, \quad y^* = y - y_{\gamma\gamma}, \quad m_T = \sqrt{p_T^2 + m^2}, \quad (\text{C1})$$

where  $y$  is the jet rapidity and  $m$  is the jet mass. The variable  $\tau_{C,j1}$  refers to the highest- $\tau$  jet, and  $\sum \tau_{C,j}$  is the scalar sum of  $\tau$  for all jets with  $\tau > 8$  GeV. For large jet rapidities,  $\tau$  corresponds to the small light-cone component of the jet,  $p_{\text{jet}}^+ = E_{\text{jet}} - |p_{z,\text{jet}}|$ , while the sum is closely related to the beam-thrust global event shape [156], as measured in the diphoton rest frame.




FIG. 36. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $p_{T}^{\gamma\gamma}$  and (b)  $|\Delta y_{\gamma\gamma}|$  are shown and compared to the SM expectations. The data are shown as filled (black) circles. The vertical error bar on each data point represents the total uncertainty in the measured cross section and the shaded (gray) band is the systematic component. The SM prediction, defined using the POWHEG NNLOPS prediction for gluon–gluon fusion and the default MC samples for the other production mechanisms, is presented as a hatched (blue) band, with the width of the band reflecting the total theoretical uncertainty (see text for details). The small contribution from VBF,  $VH$ ,  $t\bar{t}H$  and  $b\bar{b}H$  is also shown as a (green) histogram and denoted by  $XH$ . The default MC has been normalized with the N<sup>3</sup>LO prediction of Refs. [7,24,31–34]. In addition, the HRES and SCETLIB+MCFM8 predictions, described in Sec. IX E, are displayed in (a) and (b), respectively.

Figure 39 shows the first and second moments of each of the additional differential distributions. The data are compared to a variety of theoretical predictions. In general, the SM predictions are in agreement with the measured distributions.

#### APPENDIX D: DIPHOTON ACCEPTANCE, PHOTON ISOLATION, AND NONPERTURBATIVE CORRECTION FACTORS FOR PARTON-LEVEL GLUON–GLUON FUSION CALCULATIONS

This appendix presents the diphoton acceptance factors that are applied to parton-level calculations of Higgs production via gluon–gluon fusion, in order to correctly account for the diphoton selection criteria applied to the Higgs-boson decay products, are shown in Table XXII for the fiducial and differential cross sections presented in Section IX E and Appendix C. Multiplicative isolation efficiency and nonperturbative correction factors that account for the efficiency of the photon isolation criterion and the impact of hadronization and underlying-event activity are presented in Tables XXIII and XXIV, respectively. The isolation efficiency is defined as the fraction of selected diphoton events (i.e., within the kinematic

acceptance) that also satisfy the isolation criteria, and is determined using samples before including hadronization and the underlying-event activity. The non-perturbative correction factors are defined as the ratios of cross sections produced with and without hadronization and the underlying event. The default nonperturbative correction is taken as the central value of an envelope formed from multiple event generators and/or event generator tunes, with the uncertainty taken to be the maximal deviation observed in the envelope. Table XXV also provides the combined non-perturbative and isolation correction with a total uncertainty that takes into account the correlations between the uncertainties of both factors. Note though that no non-perturbative correction factors are applied to the SM predictions presented in this paper.

A summary of the binning of all differential variables is given in Table XXVI.

#### APPENDIX E: SUPPLEMENT TO EVENT CATEGORIZATION

Table XXVII summarizes the number of expected signal events and measured background events in the smallest interval expected to contain 90% of the expected SM signal events, together with the expected signal purity and local significance in the same interval, for each of the event




FIG. 37. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $H_T$ , (b)  $|\Delta\phi_{jj}|$ , and (c)  $p_T^{\gamma\gamma}$  are shown and compared to the SM expectations. The data and theoretical predictions are presented in the same way as in Fig. 36. In addition, the NNLOJET prediction is displayed in (a), and the SHERPA and GOSAM predictions are displayed in (b) and (c). More details of these predictions can be found in Sec. IX E 1.

reconstruction categories. The definition of the categories can be found in Table IV in Sec. VIII A 6.

Table XXVIII summarizes the fractions of signal events from the different production modes expected in each reconstruction category, as illustrated in Fig. 8.

Table XXIX summarizes the chosen background function used in each reconstruction category.

## APPENDIX F: LIMITS ON $\mu_{ZH}$ AND $\mu_{WH}$ USING PSEUDOEXPERIMENTS

As discussed in Sec. VIII B 2, Table XXX shows the observed and expected limits for  $\mu_{VH}$ , and separately for  $\mu_{ZH}$  and  $\mu_{WH}$ , as obtained using the asymptotic approximation and ensembles of pseudo-experiments.




FIG. 38. The differential cross sections for  $pp \rightarrow H \rightarrow \gamma\gamma$  as a function of (a)  $\tau_{C,j1}$  and (b)  $\sum \tau_{C,j}$  are shown and compared to the SM expectations. The data and theoretical predictions are presented in the same way as in Fig. 36. In addition, the NNLOJET prediction is displayed in (b).




FIG. 39. (a) The ratio of the first moment (mean) of each differential distribution predicted by the Standard Model to that observed in the data. The SM moment is calculated by using the POWHEG NNLOPS prediction for gluon–gluon fusion and the default MC samples for the other production mechanisms. (b) The ratio of the second moment (RMS) of each differential distribution predicted by the Standard Model to that observed in the data. The intervals on the vertical axes each represent one of the differential distributions. The band for the theoretical prediction represents the corresponding uncertainty in that prediction (see text for details). The error bar on the data represents the total uncertainty in the measurement, with the gray band representing only the systematic uncertainty.

TABLE XXII. Diphoton kinematic acceptances in percent for gluon–gluon fusion for the diphoton fiducial region and all differential variable bins studied in this paper, defined as the probability to fulfill the diphoton kinematic criteria:  $p_T/m_{\gamma\gamma} < 0.35$  (0.25) for the leading (subleading) photon and  $|\eta_\gamma| < 2.37$ . The factors are evaluated using the POWHEG NNLOPS event generator. Uncertainties are taken from PDF variations. QCD scale variations have a negligible impact on these factors. The range of each bin is given in Table XXVI.

	Bin 1	Bin 2	Bin 3	Bin 4	Bin 5	Bin 6	Bin 7	Bin 8	Bin 9	Bin 10	Bin 11	Bin 12	Bin 13
Diphoton fiducial	51.9 ± 0.4	...	...	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma}$	52.3 ± 0.5	52.9 ± 0.5	52.0 ± 0.4	50.8 ± 0.4	50.0 ± 0.3	48.6 ± 0.3	51.0 ± 0.3	57.7 ± 0.3	65.2 ± 0.3	...	...	...	...
$ y_{\gamma\gamma} $	74.0 ± 0.2	73.6 ± 0.2	72.2 ± 0.2	69.3 ± 0.1	67.1 ± 0.1	64.8 ± 0.1	61.5 ± 0.1	56.0 ± 0.2	31.4 ± 0.1	...	...	...	...
$p_{Tl}^{\gamma\gamma}$	49.5 ± 0.6	49.4 ± 0.6	49.2 ± 0.6	48.8 ± 0.6	49.0 ± 0.6	49.8 ± 0.6	52.8 ± 0.7	59.7 ± 0.5	69.5 ± 0.5	73.2 ± 0.5	75.7 ± 0.5	78.3 ± 0.4	82.9 ± 0.4
$ \cos\theta^* $	78.8 ± 0.6	74.9 ± 0.6	73.3 ± 0.6	71.7 ± 0.6	69.6 ± 0.6	67.6 ± 0.7	65.8 ± 0.7	60.4 ± 0.7	23.6 ± 0.5	...	...	...	...
$ \Delta y_{\gamma\gamma} $	80.0 ± 0.5	76.1 ± 0.5	74.9 ± 0.5	73.7 ± 0.6	72.4 ± 0.6	70.6 ± 0.6	68.6 ± 0.6	67.4 ± 0.6	65.3 ± 0.7	50.8 ± 0.7	...	...	...
$N_{\text{jet}}, p_T > 30 \text{ GeV}$	51.8 ± 0.5	51.8 ± 0.4	52.3 ± 0.4	54.2 ± 0.7	...	...	...	...	...	...	...	...	...
$N_{\text{jet}}, p_T > 50 \text{ GeV}$	51.2 ± 0.5	54.2 ± 0.4	55.8 ± 0.7	...	...	...	...	...	...	...	...	...	...
$p_T^j$	51.8 ± 0.5	49.8 ± 0.4	56.1 ± 0.4	52.7 ± 0.4	55.5 ± 0.6	...	...	...	...	...	...	...	...
$p_T^j$	51.8 ± 0.4	51.9 ± 0.4	57.1 ± 0.8	...	...	...	...	...	...	...	...	...	...
$H_T$	51.8 ± 0.5	51.1 ± 0.4	52.1 ± 0.4	53.8 ± 0.5	57.3 ± 0.9	...	...	...	...	...	...	...	...
$ y_{j_1} $	55.2 ± 0.4	54.6 ± 0.4	53.1 ± 0.4	53.0 ± 0.4	52.3 ± 0.4	50.4 ± 0.4	48.2 ± 0.4	...	...	...	...	...	...
$ y_{j_2} $	54.1 ± 0.5	53.5 ± 0.5	52.3 ± 0.5	...	...	...	...	...	...	...	...	...	...
$m_{jj}$	52.8 ± 0.5	53.3 ± 0.6	51.6 ± 0.6	...	...	...	...	...	...	...	...	...	...
$ \Delta y_{jj} $	70.2 ± 3.7	68.7 ± 3.4	67.2 ± 3.4	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{jj} $	53.8 ± 0.6	53.3 ± 0.5	51.6 ± 0.4	...	...	...	...	...	...	...	...	...	...
$\Delta\phi_{jj}$	51.9 ± 0.4	54.1 ± 0.6	53.4 ± 0.6	52.0 ± 0.4	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma jj}$	47.6 ± 0.4	54.4 ± 0.5	...	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{\gamma\gamma jj} $	50.9 ± 0.6	54.0 ± 0.4	54.0 ± 0.4	...	...	...	...	...	...	...	...	...	...
$\tau_{C,j}$	51.0 ± 0.5	50.9 ± 0.4	55.9 ± 0.4	63.0 ± 0.5	59.0 ± 0.7	...	...	...	...	...	...	...	...
$\sum \tau_{C,j}$	51.0 ± 0.5	50.5 ± 0.4	52.3 ± 0.4	59.5 ± 0.4	57.6 ± 0.5	61.7 ± 0.9	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [ \cos\theta^*  < 0.5]$	72.5 ± 0.6	71.1 ± 0.5	54.0 ± 0.3	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [0.5 \leq  \cos\theta^*  < 1.0]$	32.4 ± 0.4	30.4 ± 0.3	55.1 ± 0.6	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 0]$	51.2 ± 0.5	52.1 ± 0.5	52.2 ± 0.5	59.9 ± 1.9	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 1]$	58.4 ± 0.5	49.6 ± 0.4	48.8 ± 0.4	51.6 ± 0.5	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 2]$	52.5 ± 0.4	50.1 ± 0.4	63.1 ± 0.3	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} \geq 3]$	52.2 ± 0.4	63.0 ± 0.3	...	...	...	...	...	...	...	...	...	...	...



TABLE XXIII. Isolation efficiencies in percent for gluon–gluon fusion  $H \rightarrow \gamma\gamma$  for the diphoton fiducial region and all differential variable bins in this analysis. The isolation efficiency is defined as the probability for both photons to fulfill the isolation criteria (as described in Sec. IX A) for events that satisfy the diphoton kinematic criteria. Regions of phase space where no reliable estimate could be obtained are listed as “100” without uncertainties. Uncertainties are assigned in the same way as for the non-perturbative correction factors: by varying the fragmentation and underlying-event modeling. These factors can be multiplied by the kinematic acceptance factors (see Table XXII) to extrapolate an inclusive gluon–gluon fusion Higgs prediction to the fiducial volume used in this analysis. The range of each bin is given in Table XXVI.

	Bin 1	Bin 2	Bin 3	Bin 4	Bin 5	Bin 6	Bin 7	Bin 8	Bin 9	Bin 10	Bin 11	Bin 12	Bin 13
Diphoton fiducial	97.7 ± 0.1	...	...	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma}$	98.4 ± 0.1	97.6 ± 0.1	97.2 ± 0.0	96.9 ± 0.0	97.0 ± 0.0	97.7 ± 0.1	98.5 ± 0.0	98.7 ± 0.1	98.9 ± 0.2	...	...	...	...
$ y_{\gamma\gamma} $	97.6 ± 0.1	97.6 ± 0.1	97.7 ± 0.1	97.6 ± 0.2	97.7 ± 0.2	97.7 ± 0.0	97.7 ± 0.1	97.8 ± 0.0	98.0 ± 0.0	...	...	...	...
$p_{Tl}^{\gamma\gamma}$	97.2 ± 0.2	97.4 ± 0.0	97.5 ± 0.1	97.7 ± 0.0	97.9 ± 0.0	97.9 ± 0.0	98.0 ± 0.1	98.0 ± 0.1	98.2 ± 0.1	98.5 ± 0.1	98.9 ± 0.0	99.1 ± 0.0	99.2 ± 0.2
$ \cos\theta^* $	97.9 ± 0.0	97.9 ± 0.1	97.9 ± 0.0	97.8 ± 0.1	97.8 ± 0.0	97.7 ± 0.2	97.7 ± 0.1	97.6 ± 0.1	97.6 ± 0.1	...	...	...	...
$ \Delta y_{\gamma\gamma} $	97.9 ± 0.1	98.0 ± 0.1	97.9 ± 0.1	97.9 ± 0.1	97.9 ± 0.2	97.9 ± 0.1	97.8 ± 0.0	97.7 ± 0.1	97.7 ± 0.1	97.4 ± 0.1	...	...	...
$N_{\text{jet}}, p_T > 30 \text{ GeV}$	97.7 ± 0.1	97.8 ± 0.1	97.8 ± 0.1	97.8 ± 0.1	...	...	...	...	...	...	...	...	...
$N_{\text{jet}}, p_T > 50 \text{ GeV}$	97.7 ± 0.1	97.8 ± 0.1	98.1 ± 0.0	...	...	...	...	...	...	...	...	...	...
$p_T^l$	97.7 ± 0.1	97.6 ± 0.1	97.8 ± 0.0	97.8 ± 0.2	98.3 ± 0.0	...	...	...	...	...	...	...	...
$p_T^2$	97.8 ± 0.1	97.7 ± 0.1	98.3 ± 0.0	...	...	...	...	...	...	...	...	...	...
$H_T$	97.7 ± 0.1	97.7 ± 0.1	97.8 ± 0.1	98.0 ± 0.2	98.2 ± 0.0	...	...	...	...	...	...	...	...
$ y_{l1} $	97.7 ± 0.1	97.7 ± 0.1	97.8 ± 0.1	97.8 ± 0.1	97.8 ± 0.1	97.7 ± 0.4	97.8 ± 0.1	...	...	...	...	...	...
$ y_{l2} $	97.8 ± 0.1	97.8 ± 0.1	97.8 ± 0.1	...	...	...	...	...	...	...	...	...	...
$m_{jj}$	97.7 ± 0.0	97.8 ± 0.1	98.2 ± 0.1	...	...	...	...	...	...	...	...	...	...
$ \Delta y_{jj} $	97.8 ± 0.1	97.8 ± 0.0	97.8 ± 0.3	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{jj} $	98.1 ± 0.1	97.9 ± 0.1	97.4 ± 0.1	...	...	...	...	...	...	...	...	...	...
$\Delta\phi_{jj}$	97.6 ± 0.1	98.0 ± 0.2	98.1 ± 0.0	97.5 ± 0.1	...	...	...	...	...	...	...	...	...
$p_T^{llj}$	98.3 ± 0.2	97.7 ± 0.1	...	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{\gamma\gamma,jj} $	98.4 ± 0.1	97.8 ± 0.2	96.4 ± 0.2	...	...	...	...	...	...	...	...	...	...
$\tau_{C,j1}$	97.7 ± 0.1	97.8 ± 0.1	97.7 ± 0.1	97.6 ± 0.1	97.9 ± 0.0	98.1 ± 0.2	...	...	...	...	...	...	...
$\sum \tau_{C,j}$	97.7 ± 0.2	97.7 ± 0.1	97.6 ± 0.0	97.7 ± 0.1	97.9 ± 0.2	98.3 ± 0.1	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [ \cos\theta^*  < 0.5]$	98.3 ± 0.1	97.0 ± 0.0	98.4 ± 0.2	98.9 ± 0.4	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [0.5 \leq  \cos\theta^*  < 1.0]$	97.6 ± 0.1	97.4 ± 0.1	98.9 ± 0.0	99.0 ± 1.0	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 0]$	98.6 ± 0.1	97.8 ± 0.1	96.4 ± 0.0	91.2 ± 1.3	100	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 1]$	97.5 ± 0.3	97.6 ± 0.2	97.7 ± 0.0	98.4 ± 0.1	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 2]$	97.0 ± 0.3	98.4 ± 0.1	98.8 ± 0.1	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} \geq 3]$	97.4 ± 0.2	98.5 ± 0.7	...	...	...	...	...	...	...	...	...	...	...

TABLE XXIV. Nonperturbative correction factors in percent accounting for the impact of hadronization and the underlying-event activity for the diphoton fiducial region and all differential variable bins. Uncertainties are evaluated by deriving these factors using different generators and tunes as described in the text. The range of each bin is given in Table XXVI.

	Bin 1	Bin 2	Bin 3	Bin 4	Bin 5	Bin 6	Bin 7	Bin 8	Bin 9	Bin 10	Bin 11	Bin 12	Bin 13
Diphoton fiducial	100 ± 2	...	...	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma}$	99.7 ± 0.4	100.3 ± 0.3	100.2 ± 0.4	99.7 ± 0.6	100.0 ± 0.7	99.8 ± 0.5	100.4 ± 0.6	99.9 ± 0.9	100.1 ± 0.6	...	...	...	...
$ y_{\gamma\gamma} $	99.8 ± 0.4	99.9 ± 0.3	100.1 ± 0.2	100.0 ± 0.5	100.0 ± 0.3	100.2 ± 0.4	99.9 ± 0.5	100.2 ± 0.3	100.0 ± 0.4	...	...	...	...
$p_T^{\gamma\gamma}$	100.0 ± 0.5	100.1 ± 0.3	100.0 ± 0.4	100.3 ± 0.3	100.0 ± 0.8	100.3 ± 0.7	99.8 ± 0.7	100.0 ± 0.3	99.9 ± 1.1	99.8 ± 0.8	99.8 ± 1.1	100.6 ± 0.9	100.4 ± 1.5
$ \cos\theta^* $	100.1 ± 0.5	100.0 ± 0.5	99.8 ± 0.5	99.7 ± 0.4	99.9 ± 0.4	100.0 ± 0.4	100.2 ± 0.5	100.0 ± 0.4	100.2 ± 0.2	...	...	...	...
$ \Delta y_{\gamma\gamma} $	99.9 ± 0.7	100.0 ± 0.5	100.0 ± 0.9	99.7 ± 0.5	99.7 ± 0.4	100.1 ± 0.4	99.9 ± 0.3	100.0 ± 0.5	100.1 ± 0.7	100.1 ± 0.2	...	...	...
$N_{\text{jet}}, p_T > 30 \text{ GeV}$	102 ± 4	99.2 ± 2.8	98.2 ± 7.3	94.4 ± 11.9	...	...	...	...	...	...	...	...	...
$N_{\text{jet}}, p_T > 50 \text{ GeV}$	100.7 ± 2.2	98.7 ± 3.6	96.4 ± 6.5	...	...	...	...	...	...	...	...	...	...
$p_T^j$	102 ± 4	98.7 ± 5.2	98.0 ± 4.7	98.1 ± 4.3	97.4 ± 3.7	...	...	...	...	...	...	...	...
$p_T^b$	99.2 ± 2.8	97.2 ± 9.2	97.3 ± 6.2	...	...	...	...	...	...	...	...	...	...
$H_T$	102 ± 4	99.1 ± 3.8	98.1 ± 5.3	97.0 ± 6.7	96.7 ± 6.5	...	...	...	...	...	...	...	...
$ y_{j_1} $	98.5 ± 4.6	98.9 ± 4.4	98.5 ± 4.7	98.0 ± 4.5	98.4 ± 4.3	99.3 ± 4.9	97.9 ± 4.9	...	...	...	...	...	...
$ y_{j_2} $	96.7 ± 8.6	97.6 ± 8.4	97.0 ± 8.9	...	...	...	...	...	...	...	...	...	...
$m_{jj}$	96.1 ± 9.4	98.6 ± 8.0	98.4 ± 8.3	...	...	...	...	...	...	...	...	...	...
$ \Delta y_{jj} $	96.5 ± 8.3	98.5 ± 9.0	96.5 ± 10.4	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{jj} $	95.2 ± 8.3	97.0 ± 7.9	100.3 ± 10.6	...	...	...	...	...	...	...	...	...	...
$\Delta\phi_{jj}$	99.0 ± 9.6	95.8 ± 8.0	95.5 ± 8.3	98.7 ± 9.5	...	...	...	...	...	...	...	...	...
$p_T^{\gamma jj}$	94.3 ± 12.4	97.8 ± 7.5	...	...	...	...	...	...	...	...	...	...	...
$ \Delta\phi_{\gamma jj} $	96.6 ± 7.1	96.3 ± 7.7	102.5 ± 14.3	...	...	...	...	...	...	...	...	...	...
$\tau_{C,j1}$	102 ± 4	98.1 ± 4.3	98.0 ± 5.1	99.2 ± 5.0	99.9 ± 4.9	98.8 ± 4.1	...	...	...	...	...	...	...
$\sum \tau_{C,j}$	100.9 ± 2.9	98.0 ± 4.0	100.1 ± 4.2	98.8 ± 5.3	98.8 ± 6.0	97.8 ± 5.3	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [ \cos\theta^*  < 0.5]$	100.1 ± 1.0	100.2 ± 0.7	100.0 ± 0.7	100.8 ± 6.8	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [0.5 \leq  \cos\theta^*  < 1.0]$	100.2 ± 0.9	99.8 ± 0.8	100.2 ± 1.3	100.3 ± 4.1	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 0]$	98.6 ± 2.3	99.8 ± 0.7	103.1 ± 4.7	110.1 ± 23.7	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 1]$	98.2 ± 9.4	99.1 ± 0.9	103.2 ± 5.8	104.9 ± 8.1	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 2]$	99.9 ± 5.2	102.5 ± 7.9	106.1 ± 12.3	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} \geq 3]$	100.1 ± 2.5	102 ± 11	...	...	...	...	...	...	...	...	...	...	...

TABLE XXV. Combined nonperturbative (Table XXIV) and particle-level isolation correction factors (Table XXIII) in percent accounting for the impact of hadronization and the underlying-event activity for the diphoton fiducial region and all differential variable bins. The uncertainties in the combined values properly take into account the correlations between both multiplicative factors.

	Bin 1	Bin 2	Bin 3	Bin 4	Bin 5	Bin 6	Bin 7	Bin 8	Bin 9	Bin 10	Bin 11	Bin 12	Bin 13
Diphoton fiducial	$97.7 \pm 2.4$	...	...	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma}$	$97.8 \pm 0.5$	$97.7 \pm 0.6$	$97.2 \pm 1.0$	$96.3 \pm 1.0$	$96.5 \pm 0.9$	$96.9 \pm 0.7$	$98.4 \pm 1.0$	$98.3 \pm 1.1$	$99.1 \pm 0.7$	...	...	...	...
$ y_{\gamma\gamma} $	$97.2 \pm 0.7$	$97.3 \pm 0.5$	$97.4 \pm 0.6$	$97.4 \pm 0.8$	$97.5 \pm 0.6$	$97.5 \pm 0.7$	$97.5 \pm 0.6$	$97.5 \pm 0.7$	$97.8 \pm 0.8$	...	...	...	...
$p_{Tl}^{\gamma\gamma}$	$97.0 \pm 0.6$	$97.4 \pm 0.7$	$97.2 \pm 0.6$	$97.6 \pm 0.7$	$97.1 \pm 0.9$	$97.8 \pm 1.0$	$97.5 \pm 0.9$	$97.6 \pm 0.6$	$97.8 \pm 1.4$	$98.0 \pm 0.8$	$98.1 \pm 1.2$	$99.6 \pm 1.0$	$99.5 \pm 1.6$
$ \cos \theta^* $	$97.7 \pm 0.8$	$97.7 \pm 0.7$	$97.3 \pm 0.8$	$97.1 \pm 0.7$	$97.4 \pm 0.6$	$97.5 \pm 0.5$	$97.6 \pm 0.7$	$97.3 \pm 0.6$	$97.5 \pm 0.7$	...	...	...	...
$ \Delta y_{\gamma\gamma} $	$97.6 \pm 0.9$	$97.9 \pm 0.5$	$97.6 \pm 1.1$	$97.4 \pm 0.7$	$97.1 \pm 0.6$	$97.7 \pm 0.7$	$97.4 \pm 0.7$	$97.5 \pm 0.9$	$97.3 \pm 0.6$	$97.3 \pm 0.8$	...	...	...
$N_{\text{jet}}, p_T > 30 \text{ GeV}$	$99.3 \pm 3.8$	$96.5 \pm 3.1$	$95.5 \pm 7.6$	$91.9 \pm 12.2$	...	...	...	...	...	...	...	...	...
$N_{\text{jet}}, p_T > 50 \text{ GeV}$	$98.4 \pm 1.5$	$96.1 \pm 3.9$	$94.2 \pm 6.9$	...	...	...	...	...	...	...	...	...	...
$p_T^j$	$99.3 \pm 3.8$	$95.9 \pm 5.5$	$95.3 \pm 5.0$	$95.5 \pm 4.8$	$95.1 \pm 4.3$	...	...	...	...	...	...	...	...
$p_T^2$	$96.5 \pm 3.1$	$94.4 \pm 9.5$	$95.1 \pm 6.5$	...	...	...	...	...	...	...	...	...	...
$H_T$	$99.3 \pm 3.8$	$96.4 \pm 4.1$	$95.4 \pm 5.7$	$94.5 \pm 7.0$	$94.5 \pm 6.8$	...	...	...	...	...	...	...	...
$ y_{j_1} $	$95.8 \pm 5.0$	$96.2 \pm 4.7$	$95.9 \pm 5.0$	$95.3 \pm 4.9$	$95.6 \pm 4.7$	$96.8 \pm 5.4$	$95.2 \pm 5.3$	...	...	...	...	...	...
$ y_{j_2} $	$94.1 \pm 9.0$	$94.8 \pm 8.7$	$94.3 \pm 9.2$	...	...	...	...	...	...	...	...	...	...
$m_{jj}$	$93.3 \pm 9.6$	$95.9 \pm 8.3$	$96.0 \pm 8.7$	...	...	...	...	...	...	...	...	...	...
$ \Delta y_{jj} $	$93.8 \pm 8.6$	$95.8 \pm 9.3$	$93.8 \pm 10.8$	...	...	...	...	...	...	...	...	...	...
$ \Delta \phi_{jj} $	$92.9 \pm 8.7$	$94.4 \pm 8.3$	$97.2 \pm 11.0$	...	...	...	...	...	...	...	...	...	...
$\Delta \phi_{jj}$	$96.0 \pm 9.9$	$93.4 \pm 8.3$	$93.0 \pm 8.5$	$95.7 \pm 9.7$	...	...	...	...	...	...	...	...	...
$p_T^{\gamma jj}$	$92.3 \pm 12.6$	$94.9 \pm 7.8$	...	...	...	...	...	...	...	...	...	...	...
$ \Delta \phi_{\gamma\gamma jj} $	$94.6 \pm 7.4$	$93.6 \pm 8.0$	$97.9 \pm 14.7$	...	...	...	...	...	...	...	...	...	...
$\tau_{C,j1}$	$99.3 \pm 3.8$	$95.4 \pm 4.8$	$95.2 \pm 5.5$	$96.3 \pm 5.2$	$97.1 \pm 5.4$	$96.5 \pm 4.5$	...	...	...	...	...	...	...
$\sum \tau_{C,j}$	$98.7 \pm 2.3$	$95.3 \pm 4.5$	$97.2 \pm 4.6$	$96.0 \pm 5.6$	$96.3 \pm 6.2$	$95.7 \pm 5.8$	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [ \cos \theta^*  < 0.5]$	$98.1 \pm 0.7$	$97.0 \pm 0.7$	$98.2 \pm 1.0$	$99.8 \pm 6.8$	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [0.5 \leq  \cos \theta^*  < 1.0]$	$97.6 \pm 0.6$	$96.9 \pm 1.1$	$98.8 \pm 1.4$	$99.7 \pm 4.5$	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 0]$	$96.7 \pm 2.4$	$97.3 \pm 1.1$	$99.0 \pm 4.6$	$102.0 \pm 24.5$	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 1]$	$95.4 \pm 9.7$	$96.3 \pm 1.3$	$100.4 \pm 5.6$	$103.2 \pm 7.9$	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} = 2]$	$96.5 \pm 5.4$	$100.5 \pm 7.7$	$104.5 \pm 12.2$	...	...	...	...	...	...	...	...	...	...
$p_T^{\gamma\gamma} [N_{\text{jet}} \geq 3]$	$97.3 \pm 3.1$	$99.9 \pm 10.6$	...	...	...	...	...	...	...	...	...	...	...

TABLE XXVI. Bin ranges for each of the studied variables.

Bin	1	2	3	4	5	6	7	8	9	10	11	12	13
$p_T^{\gamma\gamma}$ [GeV]	0–20	20–30	30–45	45–60	60–80	80–120	120–170	170–220	220–350				
$ y_{\gamma\gamma} $	0–0.15	0.15–0.3	0.3–0.45	0.45–0.6	0.6–0.75	0.75–0.9	0.9–1.2	1.2–1.6	1.6–2.4				
$p_T^{\tau\tau}$ [GeV]	0–5	5–10	10–15	15–22	22–30	30–40	40–50	50–65	65–80	80–100	100–125	125–160	160–250
$ \cos\theta^* $	0–0.0625	0.0625–0.125	0.125–0.1875	0.1875–0.25	0.25–0.3125	0.3125–0.375	0.375–0.5	0.5–0.625	0.625–1				
$ \Delta y_{\tau\tau} $	0–0.1	0.1–0.2	0.2–0.3	0.3–0.4	0.4–0.5	0.5–0.65	0.65–0.8	0.8–1	1–1.25	1.25–2			
$N_{\text{jet}}, p_T > 30$ GeV	0	1	2	$\geq 3$									
$N_{\text{jet}}, p_T > 50$ GeV	0	1	$\geq 2$										
$p_T^1$ [GeV]	0–30	30–55	55–75	75–120	120–350								
$p_T^2$ [GeV]	0–30	30–70	70–120										
$H_T$ [GeV]	0–30	30–70	70–140	140–200	200–500								
$ y_{t_1} $	0–0.5	0.5–1	1–1.5	1.5–1.9	1.9–2.3	2.3–2.5	2.5–4.4						
$ y_{t_2} $	0–1.2	1.2–2	2–4.4										
$m_{jj}$ [GeV]	0–170	170–500	500–1500										
$ \Delta y_{jj} $	0–2	2–4	4–8.8										
$ \Delta\phi_{jj} $	$0-\frac{\pi}{3}$	$\frac{\pi}{3}-\frac{2\pi}{3}$	$\frac{2\pi}{3}-\pi$										
$\Delta\phi_{jj}$	$-\pi$	$-\frac{\pi}{2}$	$0-\frac{\pi}{2}$	$\frac{\pi}{2}-\pi$									
$p_T^{ij}$ [GeV]	0–15	15–200											
$ \Delta\phi_{\tau\tau,ij} $	0–3.01	3.01–3.1	3.1– $\pi$										
$\tau_{C,j}$ [GeV]	0–10	10–20	20–30	30–40	40–150								
$\sum \tau_{C,j}$ [GeV]	0–8	8–17	17–25	25–40	40–80	80–150							
$p_T^{\tau\tau} [ \cos\theta^*  < 0.5]$ [GeV]	0–30	30–120	120–350										
$p_T^{\tau\tau} [0.5 \leq  \cos\theta^*  < 1.0]$ [GeV]	0–30	30–120	120–350										
$p_T^{\tau\tau} [N_{\text{jet}} = 0]$ [GeV]	0–15	15–30	30–75	75–350									
$p_T^{\tau\tau} [N_{\text{jet}} = 1]$ [GeV]	0–40	40–60	60–100	100–350									
$p_T^{\tau\tau} [N_{\text{jet}} = 2]$ [GeV]	0–100	100–200	200–350										
$p_T^{\tau\tau} [N_{\text{jet}} \geq 3]$ [GeV]	0–200	200–350											

TABLE XXVII. The effective signal mass resolutions  $\sigma_{68}$  ( $\sigma_{90}$ ) in GeV defined as half the width containing 68% (90%) of the signal events for listed for each reconstructed category. Further, the numbers of background events  $B_{90}$ , measured by fits to the data, in the smallest interval expected to contain 90% of the SM signal events  $S_{90}$  are given, accompanied by the expected purities  $f_{90} \equiv S_{90}/(S_{90} + B_{90})$  and expected significances  $Z_{90} \equiv \sqrt{2((S_{90} + B_{90}) \log(1 + S_{90}/B_{90}) - S_{90})}$ .

Category	$\sigma_{68}$ [GeV]	$\sigma_{90}$ [GeV]	$S_{90}$	$B_{90}$	$f_{90}$	$Z_{90}$
ttH lep 0fwd	1.7	3.0	0.93	3.6	0.21	0.47
ttH lep 1fwd	1.7	3.0	0.99	1.9	0.34	0.67
ttH lep	1.6	2.9	2.1	2.7	0.44	1.16
ttH had BDT1	1.6	2.8	1.3	2.0	0.40	0.85
ttH had BDT2	1.6	2.9	1.6	3.9	0.29	0.75
ttH had BDT3	1.6	2.9	0.54	2.3	0.19	0.35
ttH had BDT4	1.6	2.9	2.2	14.0	0.14	0.58
tH had 4j1b	1.7	3.0	2.3	48	0.05	0.32
tH had 4j2b	1.7	3.1	0.56	6.8	0.08	0.21
VH dilep	1.7	3.0	0.84	1.1	0.43	0.72
VH lep High	1.5	2.8	1.4	2.4	0.37	0.82
VH lep Low	1.8	3.3	5.8	52	0.10	0.79
VH MET High	1.6	2.8	1.2	2.3	0.34	0.72
VH MET Low	1.8	3.3	0.56	3.4	0.14	0.30
jet BSM	1.4	2.6	24	280	0.08	1.41
VH had tight	1.5	2.8	11	47	0.19	1.55
VH had loose	1.7	3.1	15	220	0.06	0.98
VBF tight, high $p_T^{Hjj}$	1.7	2.8	18	120	0.13	1.62
VBF loose, high $p_T^{Hjj}$	1.8	3.1	15	250	0.06	0.93
VBF tight, low $p_T^{Hjj}$	1.6	2.9	12	12	0.50	3.12
VBF loose, low $p_T^{Hjj}$	1.8	3.3	17	110	0.14	1.62
ggH 2J BSM	1.4	2.6	6.8	26	0.21	1.29
ggH 2J High	1.6	2.9	26	280	0.08	1.53
ggH 2J Med	1.8	3.2	65	1700	0.04	1.56
ggH 2J Low	1.9	3.4	73	3100	0.02	1.30
ggH 1J BSM	1.4	2.6	2.0	7.1	0.22	0.72
ggH 1J High	1.6	2.9	28	240	0.11	1.80
ggH 1J Med	1.8	3.2	140	2900	0.05	2.61
ggH 1J Low	1.9	3.4	260	8000	0.03	2.89
ggH 0J Fwd	2.1	3.8	520	21000	0.02	3.62
ggH 0J Cen	1.6	2.7	300	5300	0.05	4.07



TABLE XXVIII. Composition of the selected Higgs boson events, in terms of the different production modes, as expected for each reconstructed category. The total expected numbers of Higgs boson events are given in the column labeled  $N_H$ .

Category	$N_H$	Composition [%]								
		ggH	VBF	WH	ZH	ggZH	$t\bar{t}H$	$b\bar{b}H$	$tHq$	$tHW$
tH lep 0fwd	1.0	4.1	0.2	5.6	2.2	0.6	75.7	0.9	8.2	2.5
tH lep 1fwd	1.1	1.8	0.2	1.4	0.8	0.2	79.4	0.2	13.5	2.6
ttH lep	2.4	...	...	0.2	0.1	...	96.0	0.1	1.0	2.6
ttH had BDT1	1.4	1.2	0.1	0.1	0.5	0.2	95.0	0.1	0.7	2.1
ttH had BDT2	1.8	3.6	0.3	0.8	1.2	0.4	89.3	0.2	1.8	2.4
ttH had BDT3	0.6	3.5	0.5	1.0	2.0	1.1	86.1	0.5	3.1	2.2
ttH had BDT4	2.5	7.0	0.8	1.4	2.7	1.7	79.4	0.3	4.3	2.4
tH had 4j1b	2.5	35.4	4.0	4.3	5.7	2.2	36.4	2.2	8.5	1.3
tH had 4j2b	0.62	23.8	2.8	1.6	9.8	3.6	39.0	8.3	10.5	0.6
VH dilep	0.93	...	...	...	76.9	18.9	4.0	...	...	0.2
VH lep High	1.5	0.2	...	76.2	3.5	1.2	16.4	...	1.2	1.3
VH lep Low	6.4	11.4	1.1	68.0	6.8	1.3	8.5	0.9	1.6	0.4
VH MET High	1.3	1.3	0.1	22.4	48.1	18.5	8.3	...	0.6	0.7
VH MET Low	0.62	11.9	0.4	23.4	48.0	15.2	0.5	0.3	0.2	...
jet BSM	27	59.9	25.8	5.9	3.3	1.1	3.0	0.1	0.6	0.2
VH had tight	12	52.4	3.5	23.8	13.5	4.4	1.9	0.1	0.2	0.1
VH had loose	16	67.3	4.9	14.6	8.8	2.2	1.6	0.4	0.3	0.1
VBF tight, high $p_T^{Hjj}$	20	46.9	48.3	1.2	0.7	0.6	0.8	0.3	1.2	...
VBF loose, high $p_T^{Hjj}$	17	69.9	23.8	2.2	1.3	0.6	0.9	0.8	0.6	...
VBF tight, low $p_T^{Hjj}$	14	13.0	86.7	0.1	0.1	...	...	...	0.1	...
VBF loose, low $p_T^{Hjj}$	19	32.5	66.6	0.3	0.2	0.1	...	0.2	0.1	...
ggH 2J BSM	7.5	76.1	10.3	4.9	2.8	1.8	3.0	0.2	0.6	0.2
ggH 2J High	29	75.8	12.8	4.8	2.6	1.3	2.0	0.1	0.4	0.1
ggH 2J Med	72	77.6	12.2	4.4	2.6	0.6	1.5	0.7	0.4	...
ggH 2J Low	81	79.1	9.5	4.5	2.9	0.3	1.1	2.3	0.3	...
ggH 1J BSM	2.2	72.4	16.9	6.0	2.7	1.5	0.3	...	0.1	...
ggH 1J High	32	76.0	17.5	3.4	1.9	0.8	0.1	0.3	...	...
ggH 1J Med	160	83.6	11.7	2.6	1.5	0.2	...	0.4	...	...
ggH 1J Low	290	90.5	5.7	1.7	0.9	...	...	1.1	...	...
ggH 0J Fwd	580	97.0	1.2	0.5	0.4	...	...	0.9	...	...
ggH 0J Cen	330	97.3	1.1	0.4	0.3	...	...	0.9	...	...

TABLE XXIX. The used background functions are listed: Either a power law ( $m_{\tau\tau}^\alpha$ ), exponential function of a first order polynomial ( $e^{m_{\tau\tau}^\alpha}$ ) or a second order polynomial ( $e^{m_{\tau\tau}^\alpha + m_{\tau\tau}^2 \beta}$ ) are used to describe the nonresonant diphoton background.

Category	Background function
ttH lep 0fwd	Power law
ttH lep 1fwd	Power law
ttH lep	Power law
ttH had BDT1	Exponential of a first order polynomial
ttH had BDT2	Exponential of a first order polynomial
ttH had BDT3	Exponential of a first order polynomial
ttH had BDT4	Exponential of a first order polynomial
tH had 4j1b	Power law
tH had 4j2b	Power law
VH dilep	Power law
VH lep High	Exponential of a first order polynomial
VH lep Low	Exponential of a first order polynomial
VH MET High	Exponential of a first order polynomial
VH MET Low	Exponential of a first order polynomial
jet BSM	Exponential of a first order polynomial
VH had tight	Exponential of a first order polynomial
VH had loose	Exponential of a first order polynomial
VBF tight, high $p_T^{Hjj}$	Exponential of a first order polynomial
VBF loose, high $p_T^{Hjj}$	Exponential of a first order polynomial
VBF tight, low $p_T^{Hjj}$	Exponential of a first order polynomial
VBF loose, low $p_T^{Hjj}$	Exponential of a first order polynomial
ggH 2J BSM	Power law
ggH 2J High	Power law
ggH 2J Med	Exponential of a second order polynomial
ggH 2J Low	Exponential of a second order polynomial
ggH 1J BSM	Exponential of a first order polynomial
ggH 1J High	Power law
ggH 1J Med	Exponential of a second order polynomial
ggH 1J Low	Exponential of a second order polynomial
ggH 0J Fwd	Exponential of a second order polynomial
ggH 0J Cen	Exponential of a second order polynomial

TABLE XXX. Observed and expected 95% CL limits for the signal strengths of the VH associated production processes. The observed asymptotic limit on  $\mu_{VH}$  is compared to that obtained using an ensemble of pseudoexperiments (PEs). Separate observed limits obtained from toys are reported for  $\mu_{ZH}$  and  $\mu_{WH}$ . These are shown for the background-only case ( $\mu_i = 0$ ), together with the  $\pm 1\sigma$  and  $\pm 2\sigma$  intervals.

Measurement	Observed	Exp. Limit ( $\mu = 1$ )	Exp. Limit ( $\mu = 0$ )	$+2\sigma$	$+1\sigma$	$-1\sigma$	$-2\sigma$
$\mu_{VH}$	2.3	2.5	1.5	3.1	2.2	1.1	0.8
$\mu_{VH}$ (PE)	2.2		1.5	3.1	2.2	1.1	1.0
$\mu_{ZH}$ (PE)	2.3		3.1	6.2	4.4	2.2	1.9
$\mu_{WH}$ (PE)	4.5		2.7	4.9	3.8	1.8	1.4

## APPENDIX G: SUMMARY OF COUPLINGS RESULTS

In this Appendix the expected and observed central values and uncertainties of signal strength measurements, production mode cross section measurements, and simplified template cross section measurements from Sec. VIII B and Appendix B are summarized.

### 1. Signal strengths

Table XXXI summarizes the observed and expected signal strengths for inclusive production and for various production modes.

TABLE XXXI. Observed and expected signal strengths for inclusive production and for various production modes. Uncertainties smaller than 0.5 (0.05) are displayed as 0 (0.0).

Production mode	Observed $\mu$				Expected $\mu$			
	Result	Statistical	Experimental	Theoretical	Result	Statistical	Experimental	Theoretical
Inclusive	0.99	+0.12 -0.12	+0.06 -0.05	+0.07 -0.05	1.00	+0.12 -0.12	+0.07 -0.06	+0.07 -0.05
ggH	0.81	+0.16 -0.16	+0.07 -0.06	+0.07 -0.05	1.00	+0.16 -0.17	+0.08 -0.06	+0.08 -0.06
VBF	2.0	+0.5 -0.5	+0.3 -0.2	+0.3 -0.2	1.0	+0.4 -0.4	+0.2 -0.1	+0.2 -0.1
VH	0.7	+0.8 -0.8	+0.2 -0.2	+0.2 -0.1	1.0	+0.8 -0.7	+0.2 -0.2	+0.1 -0.1
$ttH + tH$	0.5	+0.6 -0.5	+0.1 -0.1	+0.1 -0.0	1.0	+0.7 -0.6	+0.1 -0.1	+0.2 -0.0

### 2. Production mode cross sections

Table XXXII summarizes the observed and expected cross sections times diphoton branching ratio for various production modes, in the fiducial region  $|y_H| < 2.5$ .

### 3. Simplified template cross sections

Table XXXIII summarizes the observed and expected simplified template cross sections times diphoton branching ratio, in the fiducial region  $|y_H| < 2.5$ .

TABLE XXXII. Observed and expected cross sections times diphoton branching ratio for various production modes, in the fiducial region  $|y_H| < 2.5$ . Uncertainties smaller than 0.5 (0.05) are displayed as 0 (0.0).

Production mode ( $ y_H  < 2.5$ )	Observed $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]				Expected $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]			
	Result	Statistical	Experimental	Theoretical	Result	Statistical	Experimental	Theoretical
ggH	82	+16 -16	+7 -6	+5 -4	102	+17 -17	+8 -6	+5 -4
VBF	17	+5 -4	+2 -2	+3 -2	8	+3 -3	+1 -1	+2 -1
VH	3	+4 -3	+1 -1	+1 -0	5	+4 -3	+1 -1	+0 -0
$ttH + tH$	0.7	+0.8 -0.7	+0.2 -0.1	+0.2 -0.0	1.3	+0.9 -0.8	+0.2 -0.1	+0.3 -0.1

TABLE XXXIII. Observed and expected simplified template cross sections times diphoton branching ratio, in the fiducial region  $|y_H| < 2.5$ .

Simplified fiducial region ( $ y_H  < 2.5$ )	Observed $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]			Expected $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]		
	Result	Statistical	Systematic	Result	Statistical	Systematic
ggH, 0 jet	37	+14 -14	+6 -5	63	+15 -15	+8 -6
ggH, 1 jet, $p_T^H < 60$ GeV	13	+12 -12	+5 -4	15	+12 -12	+6 -4
ggH, 1 jet, $60 \leq p_T^H < 120$ GeV	5	+6 -6	+2 -1	10	+6 -6	+2 -1
ggH, 1 jet, $120 \leq p_T^H < 200$ GeV	2.8	+1.6 -1.5	+0.7 -0.5	1.7	+1.6 -1.6	+0.5 -0.4
ggH, $\geq 2$ jet	20	+8 -8	+4 -3	11	+8 -8	+3 -2
$qq \rightarrow Hqq$ , $p_T^j < 200$ GeV	15	+5 -5	+3 -2	10	+5 -5	+2 -1
ggH + $qq \rightarrow Hqq$ , BSM – like	2.0	+1.3 -1.3	+0.6 -0.6	1.8	+1.3 -1.3	+0.5 -0.5
VH, leptonic	0.7	+1.4 -1.2	+0.4 -0.3	1.4	+1.3 -1.2	+0.3 -0.3
$ttH + tH$	0.7	+0.8 -0.7	+0.2 -0.1	1.3	+0.9 -0.8	+0.3 -0.1

#### 4. Minimally merged simplified template cross sections

Table XXXIV summarizes the observed and expected minimally merged simplified template cross sections times diphoton branching ratio, in the fiducial region  $|y_H| < 2.5$ .

#### APPENDIX H: OBSERVED AND EXPECTED CORRELATION MAPS

This Appendix summaries the observed and expected correlations between the parameters of interest of each of the measurements presented in Sec. VIII B are given. The observed and expected correlations for the

TABLE XXXIV. Observed and expected simplified template cross sections times diphoton branching ratio, in the fiducial region  $|y_H| < 2.5$ .

Simplified fiducial region ( $ y_H  < 2.5$ )	Observed $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]			Expected $\sigma \times B(H \rightarrow \gamma\gamma)$ [fb]		
	Result	Statistical	Systematic	Result	Statistical	Systematic
ggH, 0 jet	38	+14 -14	+6 -5	63	+15 -15	+8 -6
ggH, 1 jet, $p_T^H < 60$ GeV	23	+13 -13	+5 -4	15	+12 -13	+6 -4
ggH, 1 jet, $60 \leq p_T^H < 120$ GeV	11	+7 -7	+3 -2	10	+8 -8	+2 -2
ggH, 1 jet, $120 \leq p_T^H < 200$ GeV	4.0	+1.8 -1.8	+0.9 -0.6	1.7	+1.9 -1.8	+0.6 -0.4
ggH, 1 jet, $p_T^H \geq 200$ GeV	2.6	+1.3 -1.1	+0.8 -0.5	0.4	+1.1 -0.9	+0.5 -0.4
ggH, $\geq 2$ jet, $p_T^H < 60$ GeV	0	+8 -8	+3 -2	3	+8 -8	+4 -2
ggH, $\geq 2$ jet, $60 \leq p_T^H < 120$ GeV	12	+7 -7	+3 -2	4	+7 -7	+2 -1
ggH, $\geq 2$ jet, $120 \leq p_T^H < 200$ GeV	7.9	+3.3 -3.3	+1.1 -0.9	2.3	+3.4 -3.3	+0.8 -0.7
ggH, $\geq 2$ jet, $p_T^H \geq 200$ GeV	2.6	+1.5 -1.4	+0.6 -0.5	1.0	+1.4 -1.3	+0.5 -0.4
ggH, VBF – like	6.2	+4.1 -4.1	+1.2 -1.2	1.5	+3.9 -3.8	+1.4 -1.0
$qq \rightarrow Hqq$ , VBF – like	3.8	+2.2 -2.0	+1.2 -1.2	2.7	+2.0 -1.8	+0.8 -0.5
$qq \rightarrow Hqq$ , VH + Rest	-19	+21 -20	+6 -7	8	+22 -21	+6 -5
$qq \rightarrow Hqq$ , $p_T^j > 200$ GeV	-3.2	+1.7 -1.7	+0.7 -0.9	0.5	+1.7 -1.7	+0.6 -0.6
VH, leptonic	0.7	+1.4 -1.2	+0.4 -0.3	1.4	+1.3 -1.2	+0.3 -0.3
$ttH + tH$	0.7	+0.8 -0.7	+0.2 -0.1	1.3	+0.9 -0.8	+0.3 -0.1

production-mode cross sections and production mode cross-section ratios are shown in Figs. 40 and 41. The observed and expected correlations for the simplified

template cross sections and minimally merged simplified template cross sections are shown in Figs. 42 and 43.




FIG. 40. Observed (left) and expected (right) correlations between the measured simplified template cross sections, including both the statistical and systematic uncertainties. The color indicates the size of the correlation.




FIG. 41. Observed (left) and expected (right) correlations between the measured simplified template cross section ratios, including both the statistical and systematic uncertainties. The color indicates the size of the correlation.






FIG. 42. Observed (top) and expected (bottom) correlations between the measured simplified template cross sections, including both the statistical and systematic uncertainties. The color indicates the size of the correlation.




FIG. 43. Observed (top) and expected (bottom) correlations between the measured simplified template cross sections, including both the statistical and systematic uncertainties. The color indicates the size of the correlation.

- [1] ATLAS Collaboration, The ATLAS experiment at the CERN Large Hadron Collider, *J. Instrum.* **3**, S08003 (2008).
- [2] CMS Collaboration, The CMS experiment at the CERN LHC, *J. Instrum.* **3**, S08004 (2008).
- [3] ATLAS Collaboration, Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012).
- [4] CMS Collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
- [5] ATLAS and CMS Collaborations, Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC  $pp$  collision data at  $\sqrt{s} = 7$  and 8 TeV, *J. High Energy Phys.* **08** (2016) 045.
- [6] ATLAS and CMS Collaborations, Combined Measurement of the Higgs Boson Mass in  $pp$  Collisions at  $\sqrt{s} = 7$  and 8 TeV with the ATLAS and CMS Experiments, *Phys. Rev. Lett.* **114**, 191803 (2015).
- [7] D. de Florian *et al.* (LHC Higgs Cross Section Working Group), Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector, [arXiv:1610.07922](https://arxiv.org/abs/1610.07922).
- [8] J. R. Andersen *et al.*, Les Houches 2015: Physics at TeV colliders standard model working group report, [arXiv:1605.04692](https://arxiv.org/abs/1605.04692).
- [9] ATLAS Collaboration, Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at  $\sqrt{s} = 8$  TeV with ATLAS, *J. High Energy Phys.* **09** (2014) 112.
- [10] CMS Collaboration, Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in  $pp$  collisions at  $\sqrt{s} = 8$  TeV, *Eur. Phys. J. C* **76**, 13 (2016).
- [11] J. C. Collins, D. E. Soper, and G. F. Sterman, Transverse momentum distribution in Drell-Yan pair and W and Z boson production, *Nucl. Phys.* **B250**, 199 (1985).
- [12] F. Bishara, U. Haisch, P. F. Monni, and E. Re, Constraining Light-Quark Yukawa Couplings from Higgs Distributions, *Phys. Rev. Lett.* **118**, 121801 (2017).
- [13] G. F. Giudice, C. Grojean, A. Pomarol, and R. Rattazzi, The strongly-interacting light Higgs, *J. High Energy Phys.* **06** (2007) 045.
- [14] ATLAS Collaboration, Constraints on non-standard model Higgs boson interactions in an effective Lagrangian using differential cross sections measured in the  $H \rightarrow \gamma\gamma$  decay channel at  $\sqrt{s} = 8$  TeV with the ATLAS detector, *Phys. Lett. B* **753**, 69 (2016).
- [15] ATLAS Collaboration, ATLAS insertable B-layer technical design report, Report No. ATLAS-TDR-19, 2010, <https://cds.cern.ch/record/1291633>; ATLAS insertable B-layer technical design report, Addendum No. ATLAS-TDR-19-ADD-1, 2012, <https://cds.cern.ch/record/1451888>.
- [16] ATLAS Collaboration, Performance of the ATLAS trigger system in 2015, *Eur. Phys. J. C* **77**, 317 (2017).
- [17] S. Heinemeyer *et al.* (LHC Higgs Cross Section Working Group), Handbook of LHC Higgs cross sections: 3. Higgs properties, [arXiv:1307.1347](https://arxiv.org/abs/1307.1347).
- [18] A. Djouadi, J. Kalinowski, and M. Spira, HDECAY: a program for Higgs boson decays in the standard model and its supersymmetric extension, *Comput. Phys. Commun.* **108**, 56 (1998).
- [19] A. Djouadi, M. M. Mühlleitner, and M. Spira, Decays of supersymmetric particles: The Program SUSY-HIT (SUSpect-SdecaY-Hdecay-InTeface), *Acta Phys. Polon. B* **38**, 635 (2007).
- [20] A. Bredenstein, A. Denner, S. Dittmaier, and M. M. Weber, Radiative corrections to the semileptonic and hadronic Higgs-boson decays  $H \rightarrow WW/ZZ \rightarrow 4$  fermions, *J. High Energy Phys.* **02** (2007) 080.
- [21] A. Bredenstein, A. Denner, S. Dittmaier, and M. M. Weber, Precise predictions for the Higgs-boson decay  $H \rightarrow WW/ZZ \rightarrow 4$  leptons, *Phys. Rev. D* **74**, 013004 (2006).
- [22] A. Bredenstein, A. Denner, S. Dittmaier, and M. M. Weber, Precision calculations for the Higgs decays  $H \rightarrow ZZ/WW \rightarrow 4$  leptons, *Nucl. Phys. B, Proc. Suppl.* **160**, 131 (2006).
- [23] K. Hamilton, P. Nason, E. Re, and G. Zanderighi, NNLOPS simulation of Higgs boson production, *J. High Energy Phys.* **10** (2013) 222.
- [24] J. Butterworth *et al.*, PDF4LHC recommendations for LHC Run II, *J. Phys. G* **43**, 023001 (2016).
- [25] K. Hamilton, P. Nason, C. Oleari, and G. Zanderighi, Merging  $H/W/Z + 0$  and 1 jet at NLO with no merging scale: a path to parton shower + NNLO matching, *J. High Energy Phys.* **05** (2013) 082.
- [26] S. Catani and M. Grazzini, An NNLO Subtraction Formalism in Hadron Collisions and Its Application to Higgs Boson Production at the LHC, *Phys. Rev. Lett.* **98**, 222002 (2007).
- [27] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC, *Nucl. Phys.* **B737**, 73 (2006).
- [28] D. de Florian, G. Ferrera, M. Grazzini, and D. Tommasini, Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC, *J. High Energy Phys.* **11** (2011) 064.
- [29] T. Sjöstrand, S. Mrenna, and P. Z. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178** (2008) 852.
- [30] ATLAS Collaboration, Measurement of the  $Z/\gamma^*$  boson transverse momentum distribution in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *J. High Energy Phys.* **09** (2014) 145.
- [31] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger, Higgs Boson Gluon-Fusion Production in QCD at Three Loops, *Phys. Rev. Lett.* **114**, 212001 (2015).
- [32] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, A. Lazopoulos, and B. Mistlberger, High precision determination of the gluon fusion Higgs boson cross-section at the LHC, *J. High Energy Phys.* **05** (2016) 058.
- [33] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, NLO electroweak corrections to Higgs boson production at hadron colliders, *Phys. Lett. B* **670**, 12 (2008).
- [34] C. Anastasiou, R. Boughezal, and F. Petriello, Mixed QCD-electroweak corrections to Higgs boson production in gluon fusion, *J. High Energy Phys.* **04** (2009) 003.
- [35] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, *J. High Energy Phys.* **11** (2004) 040.

- [36] S. Frixione, P. Nason, and C. Oleari, Matching NLO QCD computations with parton shower simulations: The POWHEG method, *J. High Energy Phys.* **11** (2007) 070.
- [37] S. Alioli, P. Nason, C. Oleari, and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: The POWHEG BOX, *J. High Energy Phys.* **06** (2010) 043.
- [38] P. Nason and C. Oleari, NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG, *J. High Energy Phys.* **02** (2010) 037.
- [39] M. Ciccolini, A. Denner, and S. Dittmaier, Strong and Electroweak Corrections to the Production of Higgs +2 Jets via Weak Interactions at the LHC, *Phys. Rev. Lett.* **99**, 161803 (2007).
- [40] M. Ciccolini, A. Denner, and S. Dittmaier, Electroweak and QCD corrections to Higgs production via vector-boson fusion at the LHC, *Phys. Rev. D* **77**, 013002 (2008).
- [41] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, Higgs Production via Vector-Boson Fusion at NNLO in QCD, *Phys. Rev. Lett.* **105**, 011801 (2010).
- [42] K. Mimasu, V. Sanz, and C. Williams, Higher order QCD predictions for associated Higgs production with anomalous couplings to gauge bosons, *J. High Energy Phys.* **08** (2016) 039.
- [43] O. Brein, A. Djouadi, and R. Harlander, NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders, *Phys. Lett. B* **579**, 149 (2004).
- [44] L. Altenkamp, S. Dittmaier, R. V. Harlander, H. Rzehak, and T. J. E. Zirke, Gluon-induced Higgs-strahlung at next-to-leading order QCD, *J. High Energy Phys.* **02** (2013) 078.
- [45] A. Denner, S. Dittmaier, S. Kallweit, and A. Mück, Electroweak corrections to Higgs-strahlung off W/Z bosons at the Tevatron and the LHC with HAWK, *J. High Energy Phys.* **03** (2012) 075.
- [46] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [47] NNPDF Collaboration, Parton distributions for the LHC Run II, *J. High Energy Phys.* **04** (2015) 040.
- [48] ATLAS Collaboration, ATLAS Run 1 Pythia8 tunes, Report No. ATL-PHYS-PUB-2014-021, 2014; <https://cds.cern.ch/record/1966419>.
- [49] W. Beenakker, S. Dittmaier, M. Krämer, B. Plümper, M. Spira, and P. M. Zerwas, NLO QCD corrections to  $t\bar{t}H$  production in hadron collisions, *Nucl. Phys.* **B653**, 151 (2003).
- [50] S. Dawson, C. Jackson, L. Orr, L. Reina, and D. Wackeroth, Associated Higgs production with top quarks at the large hadron collider: NLO QCD corrections, *Phys. Rev. D* **68**, 034022 (2003).
- [51] Y. Zhang, W.-G. Ma, R.-Y. Zhang, C. Chen, and L. Guo, QCD NLO and EW NLO corrections to  $t\bar{t}H$  production with top quark decays at hadron collider, *Phys. Lett. B* **738**, 1 (2014).
- [52] S. Frixione, V. Hirschi, D. Pagani, H.-S. Shao, and M. Zaro, Electroweak and QCD corrections to top-pair hadroproduction in association with heavy bosons, *J. High Energy Phys.* **06** (2015) 184.
- [53] M. Wiesemann, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, and P. Torrielli, Higgs production in association with bottom quarks, *J. High Energy Phys.* **02** (2015) 132.
- [54] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin, and C.-P. Yuan, New parton distributions for collider physics, *Phys. Rev. D* **82**, 074024 (2010).
- [55] S. Dawson, C. Jackson, L. Reina, and D. Wackeroth, Exclusive Higgs boson production with bottom quarks at hadron colliders, *Phys. Rev. D* **69**, 074027 (2004).
- [56] S. Dittmaier, M. Krämer, and M. Spira, Higgs radiation off bottom quarks at the Tevatron and the CERN LHC, *Phys. Rev. D* **70**, 074010 (2004).
- [57] R. Harlander, M. Kramer, and M. Schumacher, Bottom-quark associated Higgs-boson production: reconciling the four- and five-flavour scheme approach, [arXiv:1112.3478](https://arxiv.org/abs/1112.3478).
- [58] S. Gieseke, A. Ribon, M. H. Seymour, P. Stephens, and B. Webber, Herwig++ 1.0: an event generator for  $e^+e^-$  annihilation, *J. High Energy Phys.* **02** (2004) 005.
- [59] J. Bellm *et al.*, Herwig++ 2.7 release note, [arXiv:1310.6877](https://arxiv.org/abs/1310.6877).
- [60] M. Bahr *et al.*, Herwig++ physics and manual, *Eur. Phys. J. C* **58**, 693 (2008).
- [61] F. Demartin, F. Maltoni, K. Mawatari, and M. Zaro, Higgs production in association with a single top quark at the LHC, *Eur. Phys. J. C* **75**, 267 (2015).
- [62] GEANT4 Collaboration, GEANT4: a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [63] ATLAS Collaboration, The ATLAS Simulation Infrastructure, *Eur. Phys. J. C* **70**, 823 (2010).
- [64] T. Gleisberg, S. Höche, F. Krauss, M. Schönherr, S. Schumann, F. Siegert, and J. Winter, Event generation with SHERPA 1.1, *J. High Energy Phys.* **02** (2009) 007.
- [65] S. Schumann and F. Krauss, A parton shower algorithm based on Catani-Seymour dipole factorisation, *J. High Energy Phys.* **03** (2008) 038.
- [66] S. Höche, F. Krauss, S. Schumann, and F. Siegert, QCD matrix elements and truncated showers, *J. High Energy Phys.* **05** (2009) 053.
- [67] ATLAS Collaboration, The ATLAS Simulation Infrastructure, *Eur. Phys. J. C* **70**, 823 (2010).
- [68] ATLAS Collaboration, Summary of ATLAS Pythia 8 tunes, Report No. ATL-PHYS-PUB-2012-003, 2012; <https://cds.cern.ch/record/1474107>.
- [69] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, Parton distributions for the LHC, *Eur. Phys. J. C* **63**, 189 (2009).
- [70] ATLAS Collaboration, Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run-1 data, *Eur. Phys. J. C* **76**, 666 (2016).
- [71] ATLAS Collaboration, Electron and photon energy calibration with the ATLAS detector using LHC Run 1 data, *Eur. Phys. J. C* **74**, 3071 (2014).
- [72] ATLAS Collaboration, Electron and photon energy calibration with the ATLAS detector using data collected in 2015 at  $\sqrt{s} = 13$  TeV, Report No. ATL-PHYS-PUB-2016-015, 2016; <https://cds.cern.ch/record/2203514>.
- [73] ATLAS Collaboration, Photon identification in 2015 ATLAS data, Report No. ATL-PHYS-PUB-2016-014, 2016; <https://cds.cern.ch/record/2203125>.



- [74] ATLAS Collaboration, Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1, *Eur. Phys. J. C* **77** (2017) 490.
- [75] M. Cacciari, G. P. Salam, and S. Sapeta, On the characterisation of the underlying event, *J. High Energy Phys.* **04** (2010) 065.
- [76] ATLAS Collaboration, Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector, *Phys. Rev. D* **90**, 112015 (2014).
- [77] M. Cacciari, G. P. Salam, and G. Soyez, The anti- $k_t$  jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063.
- [78] M. Cacciari and G. P. Salam, Dispelling the  $N^3$  myth for the  $k_t$  jet-finder, *Phys. Lett. B* **641**, 57 (2006).
- [79] M. Cacciari, G. P. Salam, and G. Soyez, FastJet user manual, *Eur. Phys. J. C* **72**, 1896 (2012).
- [80] ATLAS Collaboration, Performance of pile-up mitigation techniques for jets in  $pp$  collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector, *Eur. Phys. J. C* **76**, 581 (2016).
- [81] ATLAS Collaboration, Jet calibration and systematic uncertainties for jets reconstructed in the ATLAS detector at  $\sqrt{s} = 13$  TeV, Report No. ATLAS-PHYS-PUB-2015-015, 2015; <https://cds.cern.ch/record/2037613>.
- [82] ATLAS Collaboration, Jet energy measurement and its systematic uncertainty in proton-proton collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Eur. Phys. J. C* **75**, 17 (2015).
- [83] ATLAS Collaboration, Tagging and suppression of pileup jets with the ATLAS detector, Report No. ATLAS-CONF-2014-018, 2014; <https://cds.cern.ch/record/1700870>.
- [84] ATLAS Collaboration, Expected performance of the ATLAS b-tagging algorithms in Run-2, Report No. ATL-PHYS-PUB-2015-022, 2015; <https://cds.cern.ch/record/2037697>.
- [85] ATLAS Collaboration, Optimisation of the ATLAS b-tagging performance for the 2016 LHC Run, Report No. ATL-PHYS-PUB-2016-012, 2016; <https://cds.cern.ch/record/2160731>.
- [86] ATLAS Collaboration, Electron efficiency measurements with the ATLAS detector using the 2015 LHC proton-proton collision data, Report No. ATLAS-CONF-2016-024, 2016; <https://cds.cern.ch/record/2157687>.
- [87] ATLAS Collaboration, Muon reconstruction performance of the ATLAS detector in proton-proton collision data at  $\sqrt{s} = 13$  TeV, *Eur. Phys. J. C* **76**, 292 (2016).
- [88] ATLAS Collaboration, Expected performance of missing transverse momentum reconstruction for the ATLAS detector at  $\sqrt{s} = 13$  TeV, Report No. ATLAS-PHYS-PUB-2015-023, 2015; <https://cds.cern.ch/record/2037700>.
- [89] ATLAS Collaboration, Performance of missing transverse momentum reconstruction for the ATLAS detector in the first proton-proton collisions at  $\sqrt{s} = 13$  TeV, Report No. ATLAS-PHYS-PUB-2015-027, 2015; <https://cds.cern.ch/record/2037904>.
- [90] ATLAS Collaboration, Measurement of the isolated diphoton cross-section in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Phys. Rev. D* **85**, 012003 (2012).
- [91] ATLAS Collaboration, Measurement of the inclusive isolated prompt photon cross section in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Phys. Rev. D* **83**, 052005 (2011).
- [92] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011); Erratum **73**, 2501(E) (2013).
- [93] ATLAS Collaboration, Combined search for the standard model Higgs boson in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Phys. Rev. D* **86**, 032003 (2012).
- [94] A. L. Read, Presentation of search results: The CLs technique, *J. Phys. G* **28**, 2693 (2002).
- [95] ATLAS Collaboration, Luminosity determination in  $pp$  collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector at the LHC, *Eur. Phys. J. C* **76**, 653 (2016).
- [96] ATLAS Collaboration, Jet energy scale measurements and their systematic uncertainties in proton-proton collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, *Phys. Rev. D* **96**, 072002 (2017).
- [97] ATLAS Collaboration, Performance of  $b$ -jet identification in the ATLAS experiment, *J. Instrum.* **11**, P04008 (2016).
- [98] Electron and photon energy calibration with the ATLAS detector using data collected in 2015 at  $\sqrt{s} = 13$  TeV, <https://cds.cern.ch/record/2203514>.
- [99] I. W. Stewart, F. J. Tackmann, J. R. Walsh, and S. Zuberi, Jet  $p_T$  Resummation in Higgs production at  $NNLL' + NNLO$ , *Phys. Rev. D* **89**, 054001 (2014).
- [100] X. Liu and F. Petriello, Reducing theoretical uncertainties for exclusive Higgs-boson plus one-jet production at the LHC, *Phys. Rev. D* **87**, 094027 (2013).
- [101] R. Boughezal, X. Liu, F. Petriello, F. J. Tackmann, and J. R. Walsh, Combining resummed Higgs predictions across jet bins, *Phys. Rev. D* **89**, 074044 (2014).
- [102] J. M. Campbell and R. K. Ellis, MCFM for the Tevatron and the LHC, *Nucl. Phys. B, Proc. Suppl.* **205–206**, 10 (2010).
- [103] I. W. Stewart and F. J. Tackmann, Theory uncertainties for Higgs and other searches using jet bins, *Phys. Rev. D* **85**, 034011 (2012).
- [104] S. Gangal and F. J. Tackmann, Next-to-leading-order uncertainties in Higgs + 2 jets from gluon fusion, *Phys. Rev. D* **87**, 093008 (2013).
- [105] R. Frederix, S. Frixione, E. Vryonidou, and M. Wiesemann, Heavy-quark mass effects in Higgs plus jets production, *J. High Energy Phys.* **08** (2016) 006.
- [106] R. Frederix and S. Frixione, Merging meets matching in MC@NLO, *J. High Energy Phys.* **12** (2012) 061.
- [107] ATLAS Collaboration, Measurement of the cross-section for electroweak production of dijets in association with a Z boson in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, *Phys. Lett. B* **775**, 206 (2017).
- [108] ATLAS Collaboration, Measurements of fiducial cross-sections for  $t\bar{t}$  production with one or two additional  $b$ -jets in pp collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector, *Eur. Phys. J. C* **76**, 11 (2016).
- [109] ATLAS Collaboration, Measurement of differential production cross-sections for a Z boson in association with  $b$ -jets in 7 TeV proton-proton collisions with the ATLAS detector, *J. High Energy Phys.* **10** (2014) 141.
- [110] ATLAS Collaboration, Measurement of the cross-section for W boson production in association with  $b$ -jets in pp



- collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *J. High Energy Phys.* **06** (2013) 084.
- [111] D. de Florian and M. Grazzini, Higgs production at the LHC: Updated cross sections at  $\sqrt{s} = 8$  TeV, *Phys. Lett. B* **718**, 117 (2012).
- [112] G. D'Agostini, A multidimensional unfolding method based on Bayes' theorem, *Nucl. Instrum. Methods Phys. Res., Sect. A* **362**, 487 (1995).
- [113] A. Höcker and V. Kartvelishvili, SVD approach to data unfolding, *Nucl. Instrum. Methods Phys. Res., Sect. A* **372**, 469 (1996).
- [114] B. Malaescu, An iterative, dynamically stabilized method of data unfolding, [arXiv:0907.3791](https://arxiv.org/abs/0907.3791).
- [115] K. G. Hayes, M. L. Perl, and B. Efron, Application of the bootstrap statistical method to the tau-decay-mode problem, *Phys. Rev. D* **39**, 274 (1989).
- [116] A. Banfi *et al.*, Jet-vetoed Higgs cross section in gluon fusion at N<sup>3</sup>LO + NNLL with small- $R$  resummation, *J. High Energy Phys.* **04** (2016) 049.
- [117] X. Chen, T. Gehrmann, E. W. N. Glover, and M. Jaquier, Precise QCD predictions for the production of Higgs + jet final states, *Phys. Lett. B* **740**, 147 (2015).
- [118] X. Chen, J. Cruz-Martinez, T. Gehrmann, E. W. N. Glover, and M. Jaquier, NNLO QCD corrections to Higgs boson production at large transverse momentum, *J. High Energy Phys.* **10** (2016) 066.
- [119] G. Cullen, N. Greiner, G. Heinrich, G. Luisoni, P. Mastrolia, G. Ossola, T. Reiter, and F. Tramontano, Automated one-loop calculations with GoSam, *Eur. Phys. J. C* **72**, 1889 (2012).
- [120] G. Cullen *et al.*, GOSAM-2.0: a tool for automated one-loop calculations within the standard model and beyond, *Eur. Phys. J. C* **74**, 3001 (2014).
- [121] N. Greiner, S. Höche, G. Luisoni, M. Schönherr, J.-C. Winter, and V. Yundin, Phenomenological analysis of Higgs boson production through gluon fusion in association with jets, *J. High Energy Phys.* **01** (2016) 169.
- [122] S. Höche, F. Krauss, M. Schönherr, and F. Siegert, A critical appraisal of NLO + PS matching methods, *J. High Energy Phys.* **09** (2012) 049.
- [123] S. Höche, F. Krauss, M. Schönherr, and F. Siegert, QCD matrix elements + parton showers: The NLO case, *J. High Energy Phys.* **04** (2013) 027.
- [124] M. Buschmann, D. Gonçalves, S. Kuttimalai, M. Schönherr, F. Krauss, and T. Plehn, Mass effects in the Higgs-gluon coupling: Boosted vs off-shell production, *J. High Energy Phys.* **02** (2015) 038.
- [125] E. Bothmann, M. Schönherr, and S. Schumann, Reweighting QCD matrix-element and parton-shower calculations, *Eur. Phys. J. C* **76**, 590 (2016).
- [126] T. Gleisberg and S. Höche, Comix, a new matrix element generator, *J. High Energy Phys.* **12** (2008) 039.
- [127] F. Krauss, R. Kuhn, and G. Soff, AMEGIC++ 1.0: A matrix element generator in C++, *J. High Energy Phys.* **02** (2002) 044.
- [128] S. Höche and M. Schönherr, Uncertainties in next-to-leading order plus parton shower matched simulations of inclusive jet and dijet production, *Phys. Rev. D* **86**, 094042 (2012).
- [129] S. Höche, F. Krauss, and M. Schönherr, Uncertainties in MEPS@NLO calculations of  $h + \text{jets}$ , *Phys. Rev. D* **90**, 014012 (2014).
- [130] S. Dulat, T.-J. Hou, J. Gao, M. Guzzi, J. Huston, P. Nadolsky, J. Pumplin, C. Schmidt, D. Stump, and C.-P. Yuan, New parton distribution functions from a global analysis of quantum chromodynamics, *Phys. Rev. D* **93**, 033006 (2016).
- [131] D. de Florian, G. Ferrera, M. Grazzini, and D. Tommasini, Higgs boson production at the LHC: transverse momentum resummation effects in the  $H \rightarrow \gamma\gamma$ ,  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$  and  $H \rightarrow ZZ \rightarrow 4\ell$  decay modes, *J. High Energy Phys.* **06** (2012) 132.
- [132] M. Grazzini and H. Sargsyan, Heavy-quark mass effects in Higgs boson production at the LHC, *J. High Energy Phys.* **09** (2013) 129.
- [133] P. F. Monni, E. Re, and P. Torrielli, Higgs Transverse-Momentum Resummation in Direct Space, *Phys. Rev. Lett.* **116**, 242001 (2016).
- [134] M. A. Ebert, J. K. L. Michel, and F. J. Tackmann, Resummation improved rapidity spectrum for gluon fusion Higgs production, *J. High Energy Phys.* **05** (2017) 088.
- [135] M. A. Ebert *et al.*, SCETlib: A C++ package for numerical calculations in QCD and soft-collinear effective theory, Report No. DESY-17-099; <http://scetlib.desy.de>.
- [136] R. Boughezal, J. M. Campbell, R. K. Ellis, C. Focke, W. Giele, X. Liu, F. Petriello, and C. Williams, Color singlet production at NNLO in MCFM, *Eur. Phys. J. C* **77**, 7 (2017).
- [137] J. Gaunt, M. Stahlhofen, F. J. Tackmann, and J. R. Walsh, N-jettiness Subtractions for NNLO QCD calculations, *J. High Energy Phys.* **09** (2015) 058.
- [138] ATLAS Collaboration, Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in  $pp$  collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector, *Phys. Lett. B* **738**, 234 (2014).
- [139] G. Klamke and D. Zeppenfeld, Higgs plus two jet production via gluon fusion as a signal at the CERN LHC, *J. High Energy Phys.* **04** (2007) 052.
- [140] T. Plehn, D. L. Rainwater, and D. Zeppenfeld, Determining the Structure of Higgs Couplings at the LHC, *Phys. Rev. Lett.* **88**, 051801 (2002).
- [141] J. R. Andersen, K. Arnold, and D. Zeppenfeld, Azimuthal angle correlations for Higgs boson plus multi-jet events, *J. High Energy Phys.* **06** (2010) 091.
- [142] M. J. Dolan, P. Harris, M. Jankowiak, and M. Spannowsky, Constraining  $CP$ -violating Higgs Sectors at the LHC using gluon fusion, *Phys. Rev. D* **90**, 073008 (2014).
- [143] R. Contino, M. Ghezzi, C. Grojean, M. Muhlleitner, and M. Spira, Effective Lagrangian for a light Higgs-like scalar, *J. High Energy Phys.* **07** (2013) 035.
- [144] A. Alloul, B. Fuks, and V. Sanz, Phenomenology of the Higgs Effective Lagrangian via FEYNRULES, *J. High Energy Phys.* **04** (2014) 110.
- [145] A. Pomarol and F. Riva, Towards the ultimate SM fit to close in on Higgs physics, *J. High Energy Phys.* **01** (2014) 151.
- [146] J. Ellis, V. Sanz, and T. You, The effective Standard Model after LHC Run I, *J. High Energy Phys.* **03** (2015) 157.

- [147] P. Artoisenet *et al.*, A framework for Higgs characterisation, *J. High Energy Phys.* **11** (2013) 043.
- [148] M. L. Mangano, M. Moretti, F. Piccinini, and M. Treccani, Matching matrix elements and shower evolution for top-quark production in hadronic collisions, *J. High Energy Phys.* **01** (2007) 013.
- [149] A. Buckley, J. Butterworth, D. Grellscheid, H. Hoeth, L. Lönnblad, J. Monk, H. Schulz, and F. Siegert, Rivet user manual, *Comput. Phys. Commun.* **184**, 2803 (2013).
- [150] A. Buckley, H. Hoeth, H. Lacker, H. Schulz, and J. E. von Seggern, Systematic event generator tuning for the LHC, *Eur. Phys. J. C* **65**, 331 (2010).
- [151] K. Arnold *et al.*, VBFNLO: A parton level Monte Carlo for processes with electroweak bosons, *Comput. Phys. Commun.* **180**, 1661 (2009).
- [152] ATLAS Collaboration, Search for Higgs boson decays to a photon and a Z boson in  $pp$  collisions at  $\sqrt{s} = 7$  and 8 TeV with the ATLAS detector, *Phys. Lett. B* **732**, 8 (2014).
- [153] ATLAS Collaboration, Searches for the  $Z\gamma$  decay mode of the Higgs boson and for new high-mass resonances in  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, *J. High Energy Phys.* **10** (2017) 112.
- [154] ATLAS Collaboration, Measurement of Higgs boson production in the diphoton decay channel in  $pp$  collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector, *Phys. Rev. D* **90**, 112015 (2014).
- [155] ATLAS Collaboration, ATLAS Computing Acknowledgements, Report No. ATL-GEN-PUB-2016-002; <https://cds.cern.ch/record/2202407>.
- [156] I. W. Stewart, F. J. Tackmann, and W. J. Waalewijn, Factorization at the LHC: From PDFs to initial state jets, *Phys. Rev. D* **81**, 094035 (2010).

M. Aaboud,<sup>137c</sup> G. Aad,<sup>88</sup> B. Abbott,<sup>115</sup> O. Abidinov,<sup>12,a</sup> B. Abeloos,<sup>119</sup> S. H. Abidi,<sup>161</sup> O. S. AbouZeid,<sup>139</sup> N. L. Abraham,<sup>151</sup> H. Abramowicz,<sup>155</sup> H. Abreu,<sup>154</sup> Y. Abulaiti,<sup>148a,148b</sup> B. S. Acharya,<sup>167a,167b,b</sup> S. Adachi,<sup>157</sup> L. Adamczyk,<sup>41a</sup> J. Adelman,<sup>110</sup> M. Adersberger,<sup>102</sup> T. Adye,<sup>133</sup> A. A. Affolder,<sup>139</sup> Y. Afik,<sup>154</sup> C. Agheorghiesei,<sup>28c</sup> J. A. Aguilar-Saavedra,<sup>128a,128f</sup> S. P. Ahlen,<sup>24</sup> F. Ahmadov,<sup>68,c</sup> G. Aielli,<sup>135a,135b</sup> S. Akatsuka,<sup>71</sup> H. Akerstedt,<sup>148a,148b</sup> T. P. A. Åkesson,<sup>84</sup> E. Akilli,<sup>52</sup> A. V. Akimov,<sup>98</sup> G. L. Alberghi,<sup>22a,22b</sup> J. Albert,<sup>172</sup> P. Albicocco,<sup>50</sup> M. J. Alconada Verzini,<sup>74</sup> S. C. Alderweireldt,<sup>108</sup> M. Aleksa,<sup>32</sup> I. N. Aleksandrov,<sup>68</sup> C. Alexa,<sup>28b</sup> G. Alexander,<sup>155</sup> T. Alexopoulos,<sup>10</sup> M. Alhroob,<sup>115</sup> B. Ali,<sup>130</sup> M. Aliev,<sup>76a,76b</sup> G. Alimonti,<sup>94a</sup> J. Alison,<sup>33</sup> S. P. Alkire,<sup>38</sup> B. M. M. Allbrooke,<sup>151</sup> B. W. Allen,<sup>118</sup> P. P. Allport,<sup>19</sup> A. Aloisio,<sup>106a,106b</sup> A. Alonso,<sup>39</sup> F. Alonso,<sup>74</sup> C. Alpigiani,<sup>140</sup> A. A. Alshehri,<sup>56</sup> M. I. Alstady,<sup>88</sup> B. Alvarez Gonzalez,<sup>32</sup> D. Álvarez Piqueras,<sup>170</sup> M. G. Alvigi,<sup>106a,106b</sup> B. T. Amadio,<sup>16</sup> Y. Amaral Coutinho,<sup>26a</sup> C. Amelung,<sup>25</sup> D. Amidei,<sup>92</sup> S. P. Amor Dos Santos,<sup>128a,128c</sup> S. Amoroso,<sup>32</sup> C. Anastopoulos,<sup>141</sup> L. S. Ancu,<sup>52</sup> N. Andari,<sup>19</sup> T. Andeen,<sup>11</sup> C. F. Anders,<sup>60b</sup> J. K. Anders,<sup>18</sup> K. J. Anderson,<sup>33</sup> A. Andreazza,<sup>94a,94b</sup> V. Andrei,<sup>60a</sup> S. Angelidakis,<sup>37</sup> I. Angelozzi,<sup>109</sup> A. Angerami,<sup>38</sup> A. V. Anisenkov,<sup>111,d</sup> N. Anjos,<sup>13</sup> A. Annovi,<sup>126a</sup> C. Antel,<sup>60a</sup> M. Antonelli,<sup>50</sup> A. Antonov,<sup>100,a</sup> D. J. Antrim,<sup>166</sup> F. Anulli,<sup>134a</sup> M. Aoki,<sup>69</sup> L. Aperio Bella,<sup>32</sup> G. Arabidze,<sup>93</sup> Y. Arai,<sup>69</sup> J. P. Araque,<sup>128a</sup> V. Araujo Ferraz,<sup>26a</sup> A. T. H. Arce,<sup>48</sup> R. E. Ardell,<sup>80</sup> F. A. Arduh,<sup>74</sup> J.-F. Arguin,<sup>97</sup> S. Argyropoulos,<sup>66</sup> M. Arik,<sup>20a</sup> A. J. Armbruster,<sup>32</sup> L. J. Armitage,<sup>79</sup> O. Arnaez,<sup>161</sup> H. Arnold,<sup>51</sup> M. Arratia,<sup>30</sup> O. Arslan,<sup>23</sup> A. Artamonov,<sup>99,a</sup> G. Artoni,<sup>122</sup> S. Artz,<sup>86</sup> S. Asai,<sup>157</sup> N. Asbah,<sup>45</sup> A. Ashkenazi,<sup>155</sup> L. Asquith,<sup>151</sup> K. Assamagan,<sup>27</sup> R. Astalos,<sup>146a</sup> M. Atkinson,<sup>169</sup> N. B. Atlay,<sup>143</sup> K. Augsten,<sup>130</sup> G. Avolio,<sup>32</sup> B. Axen,<sup>16</sup> M. K. Ayoub,<sup>35a</sup> G. Azuelos,<sup>97,e</sup> A. E. Baas,<sup>60a</sup> M. J. Baca,<sup>19</sup> H. Bachacou,<sup>138</sup> K. Bachas,<sup>76a,76b</sup> M. Backes,<sup>122</sup> P. Bagnaia,<sup>134a,134b</sup> M. Bahmani,<sup>42</sup> H. Bahraseramani,<sup>144</sup> J. T. Baines,<sup>133</sup> M. Bajic,<sup>39</sup> O. K. Baker,<sup>179</sup> P. J. Bakker,<sup>109</sup> D. Bakshi Gupta,<sup>82</sup> E. M. Baldin,<sup>111,d</sup> P. Balek,<sup>175</sup> F. Balli,<sup>138</sup> W. K. Balunas,<sup>124</sup> E. Banas,<sup>42</sup> A. Bandyopadhyay,<sup>23</sup> Sw. Banerjee,<sup>176,f</sup> A. A. E. Bannoura,<sup>178</sup> L. Barak,<sup>155</sup> E. L. Barberio,<sup>91</sup> D. Barberis,<sup>53a,53b</sup> M. Barbero,<sup>88</sup> T. Barillari,<sup>103</sup> M.-S. Barisits,<sup>65</sup> J. T. Barkeloo,<sup>118</sup> T. Barklow,<sup>145</sup> N. Barlow,<sup>30</sup> S. L. Barnes,<sup>36b</sup> B. M. Barnett,<sup>133</sup> R. M. Barnett,<sup>16</sup> Z. Barnovska-Blenessy,<sup>36c</sup> A. Baroncelli,<sup>136a</sup> G. Barone,<sup>25</sup> A. J. Barr,<sup>122</sup> L. Barranco Navarro,<sup>170</sup> F. Barreiro,<sup>85</sup> J. Barreiro Guimarães da Costa,<sup>35a</sup> R. Bartoldus,<sup>145</sup> A. E. Barton,<sup>75</sup> P. Bartos,<sup>146a</sup> A. Basalaeu,<sup>125</sup> A. Bassalat,<sup>119,g</sup> R. L. Bates,<sup>56</sup> S. J. Batista,<sup>161</sup> J. R. Batley,<sup>30</sup> M. Battaglia,<sup>139</sup> M. Bauce,<sup>134a,134b</sup> F. Bauer,<sup>138</sup> K. T. Bauer,<sup>166</sup> H. S. Bawa,<sup>145,h</sup> J. B. Beacham,<sup>113</sup> M. D. Beattie,<sup>75</sup> T. Beau,<sup>83</sup> P. H. Beauchemin,<sup>165</sup> P. Bechtel,<sup>23</sup> H. P. Beck,<sup>18,i</sup> H. C. Beck,<sup>57</sup> K. Becker,<sup>122</sup> M. Becker,<sup>86</sup> C. Becot,<sup>112</sup> A. J. Beddall,<sup>20d</sup> A. Beddall,<sup>20b</sup> V. A. Bednyakov,<sup>68</sup> M. Bedognetti,<sup>109</sup> C. P. Bee,<sup>150</sup> T. A. Beermann,<sup>32</sup> M. Begalli,<sup>26a</sup> M. Begel,<sup>27</sup> J. K. Behr,<sup>45</sup> A. S. Bell,<sup>81</sup> G. Bella,<sup>155</sup> L. Bellagamba,<sup>22a</sup> A. Bellerive,<sup>31</sup> M. Bellomo,<sup>154</sup> K. Belotskiy,<sup>100</sup> O. Beltramello,<sup>32</sup> N. L. Belyaev,<sup>100</sup> O. Benary,<sup>155,a</sup> D. Bencheikroun,<sup>137a</sup> M. Bender,<sup>102</sup> N. Benekos,<sup>10</sup> Y. Benhammou,<sup>155</sup> E. Benhar Nocchioli,<sup>179</sup> J. Benitez,<sup>66</sup> D. P. Benjamin,<sup>48</sup> M. Benoit,<sup>52</sup> J. R. Bensinger,<sup>25</sup> S. Bentvelsen,<sup>109</sup> L. Beresford,<sup>122</sup> M. Beretta,<sup>50</sup> D. Berge,<sup>109</sup> E. Bergeaas Kuutmann,<sup>168</sup> N. Berger,<sup>5</sup> L. J. Bergsten,<sup>25</sup> J. Beringer,<sup>16</sup> S. Berlendis,<sup>58</sup> N. R. Bernard,<sup>89</sup> G. Bernardi,<sup>83</sup> C. Bernius,<sup>145</sup> F. U. Bernlochner,<sup>23</sup> T. Berry,<sup>80</sup> P. Berta,<sup>86</sup> C. Bertella,<sup>35a</sup> G. Bertoli,<sup>148a,148b</sup> I. A. Bertram,<sup>75</sup> C. Bertsche,<sup>45</sup> G. J. Besjes,<sup>39</sup> O. Bessidskaia Bylund,<sup>148a,148b</sup> M. Bessner,<sup>45</sup> N. Besson,<sup>138</sup> A. Bethani,<sup>87</sup>

S. Bethke,<sup>103</sup> A. Betti,<sup>23</sup> A. J. Bevan,<sup>79</sup> J. Beyer,<sup>103</sup> R. M. Bianchi,<sup>127</sup> O. Biebel,<sup>102</sup> D. Biedermann,<sup>17</sup> R. Bielski,<sup>87</sup> K. Bierwagen,<sup>86</sup> N. V. Biesuz,<sup>126a,126b</sup> M. Biglietti,<sup>136a</sup> T. R. V. Billoud,<sup>97</sup> H. Bilokon,<sup>50</sup> M. Bindi,<sup>57</sup> A. Bingul,<sup>20b</sup> C. Bini,<sup>134a,134b</sup> S. Biondi,<sup>22a,22b</sup> T. Bisanz,<sup>57</sup> C. Bittrich,<sup>47</sup> D. M. Bjergaard,<sup>48</sup> J. E. Black,<sup>145</sup> K. M. Black,<sup>24</sup> R. E. Blair,<sup>6</sup> T. Blazek,<sup>146a</sup> I. Bloch,<sup>45</sup> C. Blocker,<sup>25</sup> A. Blue,<sup>56</sup> U. Blumenschein,<sup>79</sup> Dr. Blunier,<sup>34a</sup> G. J. Bobbink,<sup>109</sup> V. S. Bobrovnikov,<sup>111,d</sup> S. S. Bocchetta,<sup>84</sup> A. Bocci,<sup>48</sup> C. Bock,<sup>102</sup> M. Boehler,<sup>51</sup> D. Boerner,<sup>178</sup> D. Bogavac,<sup>102</sup> A. G. Bogdanchikov,<sup>111</sup> C. Bohm,<sup>148a</sup> V. Boisvert,<sup>80</sup> P. Bokan,<sup>168j</sup> T. Bold,<sup>41a</sup> A. S. Boldyrev,<sup>101</sup> A. E. Bolz,<sup>60b</sup> M. Bomben,<sup>83</sup> M. Bona,<sup>79</sup> J. S. Bonilla,<sup>118</sup> M. Boonekamp,<sup>138</sup> A. Borisov,<sup>132</sup> G. Borissov,<sup>75</sup> J. Bortfeldt,<sup>32</sup> D. Bortoletto,<sup>122</sup> V. Bortolotto,<sup>62a</sup> D. Boscherini,<sup>22a</sup> M. Bosman,<sup>13</sup> J. D. Bossio Sola,<sup>29</sup> J. Boudreau,<sup>127</sup> E. V. Bouhova-Thacker,<sup>75</sup> D. Boumediene,<sup>37</sup> C. Bourdarios,<sup>119</sup> S. K. Boutle,<sup>56</sup> A. Boveia,<sup>113</sup> J. Boyd,<sup>32</sup> I. R. Boyko,<sup>68</sup> A. J. Bozson,<sup>80</sup> J. Bracinik,<sup>19</sup> A. Brandt,<sup>8</sup> G. Brandt,<sup>178</sup> O. Brandt,<sup>60a</sup> F. Braren,<sup>45</sup> U. Bratzler,<sup>158</sup> B. Brau,<sup>89</sup> J. E. Brau,<sup>118</sup> W. D. Breaden Madden,<sup>56</sup> K. Brendlinger,<sup>45</sup> A. J. Brennan,<sup>91</sup> L. Brenner,<sup>109</sup> R. Brenner,<sup>168</sup> S. Bressler,<sup>175</sup> D. L. Briglin,<sup>19</sup> T. M. Bristow,<sup>49</sup> D. Britton,<sup>56</sup> D. Britzger,<sup>60b</sup> F. M. Brochu,<sup>30</sup> I. Brock,<sup>23</sup> R. Brock,<sup>93</sup> G. Brooijmans,<sup>38</sup> T. Brooks,<sup>80</sup> W. K. Brooks,<sup>34b</sup> E. Brost,<sup>110</sup> J. H. Broughton,<sup>19</sup> P. A. Bruckman de Renstrom,<sup>42</sup> D. Bruncko,<sup>146b</sup> A. Bruni,<sup>22a</sup> G. Bruni,<sup>22a</sup> L. S. Bruni,<sup>109</sup> S. Bruno,<sup>135a,135b</sup> B. H. Brunt,<sup>30</sup> M. Bruschi,<sup>22a</sup> N. Bruscino,<sup>127</sup> P. Bryant,<sup>33</sup> L. Bryngemark,<sup>45</sup> T. Buanes,<sup>15</sup> Q. Buat,<sup>144</sup> P. Buchholz,<sup>143</sup> A. G. Buckley,<sup>56</sup> I. A. Budagov,<sup>68</sup> F. Buehrer,<sup>51</sup> M. K. Bugge,<sup>121</sup> O. Bulekov,<sup>100</sup> D. Bullock,<sup>8</sup> T. J. Burch,<sup>110</sup> S. Burdin,<sup>77</sup> C. D. Burgard,<sup>109</sup> A. M. Burger,<sup>5</sup> B. Burghgrave,<sup>110</sup> K. Burka,<sup>42</sup> S. Burke,<sup>133</sup> I. Burmeister,<sup>46</sup> J. T. P. Burr,<sup>122</sup> D. Büscher,<sup>51</sup> V. Büscher,<sup>86</sup> E. Buschmann,<sup>57</sup> P. Bussey,<sup>56</sup> J. M. Butler,<sup>24</sup> C. M. Buttar,<sup>56</sup> J. M. Butterworth,<sup>81</sup> P. Butti,<sup>32</sup> W. Buttinger,<sup>27</sup> A. Buzatu,<sup>153</sup> A. R. Buzykaev,<sup>111,d</sup> S. Cabrera Urbán,<sup>170</sup> D. Caforio,<sup>130</sup> H. Cai,<sup>169</sup> V. M. M. Cairo,<sup>2</sup> O. Cakir,<sup>4a</sup> N. Calace,<sup>52</sup> P. Calafiura,<sup>16</sup> A. Calandri,<sup>88</sup> G. Calderini,<sup>83</sup> P. Calfayan,<sup>64</sup> G. Callea,<sup>40a,40b</sup> L. P. Caloba,<sup>26a</sup> S. Calvente Lopez,<sup>85</sup> D. Calvet,<sup>37</sup> S. Calvet,<sup>37</sup> T. P. Calvet,<sup>88</sup> R. Camacho Toro,<sup>33</sup> S. Camarda,<sup>32</sup> P. Camarri,<sup>135a,135b</sup> D. Cameron,<sup>121</sup> R. Caminal Armadans,<sup>169</sup> C. Camincher,<sup>58</sup> S. Campana,<sup>32</sup> M. Campanelli,<sup>81</sup> A. Camplani,<sup>94a,94b</sup> A. Campoverde,<sup>143</sup> V. Canale,<sup>106a,106b</sup> M. Cano Bret,<sup>36b</sup> J. Cantero,<sup>116</sup> T. Cao,<sup>155</sup> M. D. M. Capeans Garrido,<sup>32</sup> I. Caprini,<sup>28b</sup> M. Caprini,<sup>28b</sup> M. Capua,<sup>40a,40b</sup> R. M. Carbone,<sup>38</sup> R. Cardarelli,<sup>135a</sup> F. Cardillo,<sup>51</sup> I. Carli,<sup>131</sup> T. Carli,<sup>32</sup> G. Carlino,<sup>106a</sup> B. T. Carlson,<sup>127</sup> L. Carminati,<sup>94a,94b</sup> R. M. D. Carney,<sup>148a,148b</sup> S. Caron,<sup>108</sup> E. Carquin,<sup>34b</sup> S. Carrá,<sup>94a,94b</sup> G. D. Carrillo-Montoya,<sup>32</sup> D. Casadei,<sup>19</sup> M. P. Casado,<sup>13,k</sup> A. F. Casha,<sup>161</sup> M. Casolino,<sup>13</sup> D. W. Casper,<sup>166</sup> R. Castelijin,<sup>109</sup> V. Castillo Gimenez,<sup>170</sup> N. F. Castro,<sup>128a,1</sup> A. Catinaccio,<sup>32</sup> J. R. Catmore,<sup>121</sup> A. Cattai,<sup>32</sup> J. Caudron,<sup>23</sup> V. Cavaliere,<sup>169</sup> E. Cavallaro,<sup>13</sup> D. Cavalli,<sup>94a</sup> M. Cavalli-Sforza,<sup>13</sup> V. Cavasinni,<sup>126a,126b</sup> E. Celebi,<sup>20c</sup> F. Ceradini,<sup>136a,136b</sup> L. Cerda Alberich,<sup>170</sup> A. S. Cerqueira,<sup>26b</sup> A. Cerri,<sup>151</sup> L. Cerrito,<sup>135a,135b</sup> F. Cerutti,<sup>16</sup> A. Cervelli,<sup>22a,22b</sup> S. A. Cetin,<sup>20c</sup> A. Chafaq,<sup>137a</sup> D. Chakraborty,<sup>110</sup> S. K. Chan,<sup>59</sup> W. S. Chan,<sup>109</sup> Y. L. Chan,<sup>62a</sup> P. Chang,<sup>169</sup> J. D. Chapman,<sup>30</sup> D. G. Charlton,<sup>19</sup> C. C. Chau,<sup>31</sup> C. A. Chavez Barajas,<sup>151</sup> S. Che,<sup>113</sup> S. Cheatham,<sup>167a,167c</sup> A. Chegwidan,<sup>93</sup> S. Chekanov,<sup>6</sup> S. V. Chekulaev,<sup>163a</sup> G. A. Chelkov,<sup>68,m</sup> M. A. Chelstowska,<sup>32</sup> C. Chen,<sup>36c</sup> C. Chen,<sup>67</sup> H. Chen,<sup>27</sup> J. Chen,<sup>36c</sup> J. Chen,<sup>38</sup> S. Chen,<sup>35b</sup> S. Chen,<sup>157</sup> X. Chen,<sup>35c,n</sup> Y. Chen,<sup>70</sup> H. C. Cheng,<sup>92</sup> H. J. Cheng,<sup>35a,35d</sup> A. Cheplakov,<sup>68</sup> E. Cheremushkina,<sup>132</sup> R. Cherkaoui El Moursli,<sup>137d</sup> E. Cheu,<sup>7</sup> K. Cheung,<sup>63</sup> L. Chevalier,<sup>138</sup> V. Chiarella,<sup>50</sup> G. Chiarelli,<sup>126a</sup> G. Chiodini,<sup>76a</sup> A. S. Chisholm,<sup>32</sup> A. Chitan,<sup>28b</sup> Y. H. Chiu,<sup>172</sup> M. V. Chizhov,<sup>68</sup> K. Choi,<sup>64</sup> A. R. Chomont,<sup>37</sup> S. Chouridou,<sup>156</sup> Y. S. Chow,<sup>62a</sup> V. Christodoulou,<sup>81</sup> M. C. Chu,<sup>62a</sup> J. Chudoba,<sup>129</sup> A. J. Chuinard,<sup>90</sup> J. J. Chwastowski,<sup>42</sup> L. Chytka,<sup>117</sup> A. K. Ciftci,<sup>4a</sup> D. Cinca,<sup>46</sup> V. Cindro,<sup>78</sup> I. A. Cioară,<sup>23</sup> A. Ciocio,<sup>16</sup> F. Ciotto,<sup>106a,106b</sup> Z. H. Citron,<sup>175</sup> M. Citterio,<sup>94a</sup> M. Ciubancan,<sup>28b</sup> A. Clark,<sup>52</sup> M. R. Clark,<sup>38</sup> P. J. Clark,<sup>49</sup> R. N. Clarke,<sup>16</sup> C. Clement,<sup>148a,148b</sup> Y. Coadou,<sup>88</sup> M. Cobal,<sup>167a,167c</sup> A. Coccaro,<sup>52</sup> J. Cochran,<sup>67</sup> L. Colasurdo,<sup>108</sup> B. Cole,<sup>38</sup> A. P. Colijn,<sup>109</sup> J. Collot,<sup>58</sup> T. Colombo,<sup>166</sup> P. Conde Muño,<sup>128a,128b</sup> E. Coniavitis,<sup>51</sup> S. H. Connell,<sup>147b</sup> I. A. Connelly,<sup>87</sup> S. Constantinescu,<sup>28b</sup> G. Conti,<sup>32</sup> F. Conventi,<sup>106a,o</sup> A. M. Cooper-Sarkar,<sup>122</sup> F. Cormier,<sup>171</sup> K. J. R. Cormier,<sup>161</sup> M. Corradi,<sup>134a,134b</sup> E. E. Corrigan,<sup>84</sup> F. Corriveau,<sup>90,p</sup> A. Cortes-Gonzalez,<sup>32</sup> M. J. Costa,<sup>170</sup> D. Costanzo,<sup>141</sup> G. Cottin,<sup>30</sup> G. Cowan,<sup>80</sup> B. E. Cox,<sup>87</sup> K. Cranmer,<sup>112</sup> S. J. Crawley,<sup>56</sup> R. A. Creager,<sup>124</sup> G. Cree,<sup>31</sup> S. Crépe-Renaudin,<sup>58</sup> F. Crescioli,<sup>83</sup> W. A. Cribbs,<sup>148a,148b</sup> M. Cristinziani,<sup>23</sup> V. Croft,<sup>112</sup> G. Crosetti,<sup>40a,40b</sup> A. Cueto,<sup>85</sup> T. Cuhadar Donszelmann,<sup>141</sup> A. R. Cukierman,<sup>145</sup> J. Cummings,<sup>179</sup> M. Curatolo,<sup>50</sup> J. Cúth,<sup>86</sup> S. Czekierda,<sup>42</sup> P. Czodrowski,<sup>32</sup> G. D'amen,<sup>22a,22b</sup> S. D'Auria,<sup>56</sup> L. D'eraimo,<sup>83</sup> M. D'Onofrio,<sup>77</sup> M. J. Da Cunha Sargedas De Sousa,<sup>128a,128b</sup> C. Da Via,<sup>87</sup> W. Dabrowski,<sup>41a</sup> T. Dado,<sup>146a</sup> S. Dahbi,<sup>137d</sup> T. Dai,<sup>92</sup> O. Dale,<sup>15</sup> F. Dallaire,<sup>97</sup> C. Dallapiccola,<sup>89</sup> M. Dam,<sup>39</sup> J. R. Dandoy,<sup>124</sup> M. F. Daneri,<sup>29</sup> N. P. Dang,<sup>176,f</sup> N. S. Dann,<sup>87</sup> M. Danninger,<sup>171</sup> M. Dano Hoffmann,<sup>138</sup> V. Dao,<sup>150</sup> G. Darbo,<sup>53a</sup> S. Darmora,<sup>8</sup> J. Dassoulas,<sup>3</sup> A. Dattagupta,<sup>118</sup> T. Daubney,<sup>45</sup> W. Davey,<sup>23</sup> C. David,<sup>45</sup> T. Davidek,<sup>131</sup> D. R. Davis,<sup>48</sup> P. Davison,<sup>81</sup> E. Dawe,<sup>91</sup> I. Dawson,<sup>141</sup> K. De,<sup>8</sup> R. de Asmundis,<sup>106a</sup> A. De Benedetti,<sup>115</sup> S. De Castro,<sup>22a,22b</sup> S. De Cecco,<sup>83</sup> N. De Groot,<sup>108</sup> P. de Jong,<sup>109</sup> H. De la Torre,<sup>93</sup> F. De Lorenzi,<sup>67</sup> A. De Maria,<sup>57</sup> D. De Pedis,<sup>134a</sup> A. De Salvo,<sup>134a</sup> U. De Sanctis,<sup>135a,135b</sup>



- A. De Santo,<sup>151</sup> K. De Vasconcelos Corga,<sup>88</sup> J. B. De Vivie De Regie,<sup>119</sup> R. Debbe,<sup>27</sup> C. Debenedetti,<sup>139</sup> D. V. Dedovich,<sup>68</sup> N. Dehghanian,<sup>3</sup> I. Deigaard,<sup>109</sup> M. Del Gaudio,<sup>40a,40b</sup> J. Del Peso,<sup>85</sup> D. Delgove,<sup>119</sup> F. Deliot,<sup>138</sup> C. M. Delitzsch,<sup>7</sup> A. Dell'Acqua,<sup>32</sup> L. Dell'Asta,<sup>24</sup> M. Della Pietra,<sup>106a,106b</sup> D. della Volpe,<sup>52</sup> M. Delmastro,<sup>5</sup> C. Delporte,<sup>119</sup> P. A. Delsart,<sup>58</sup> D. A. DeMarco,<sup>161</sup> S. Demers,<sup>179</sup> M. Demichev,<sup>68</sup> A. Demilly,<sup>83</sup> S. P. Denisov,<sup>132</sup> D. Denysiuk,<sup>138</sup> D. Derendarz,<sup>42</sup> J. E. Derkaoui,<sup>137c</sup> F. Derue,<sup>83</sup> P. Dervan,<sup>77</sup> K. Desch,<sup>23</sup> C. Deterre,<sup>45</sup> K. Dette,<sup>161</sup> M. R. Devesa,<sup>29</sup> P. O. Deviveiros,<sup>32</sup> A. Dewhurst,<sup>133</sup> S. Dhaliwal,<sup>25</sup> F. A. Di Bello,<sup>52</sup> A. Di Ciaccio,<sup>135a,135b</sup> L. Di Ciaccio,<sup>5</sup> W. K. Di Clemente,<sup>124</sup> C. Di Donato,<sup>106a,106b</sup> A. Di Girolamo,<sup>32</sup> B. Di Girolamo,<sup>32</sup> B. Di Micco,<sup>136a,136b</sup> R. Di Nardo,<sup>32</sup> K. F. Di Petrillo,<sup>59</sup> A. Di Simone,<sup>51</sup> R. Di Sipio,<sup>161</sup> D. Di Valentino,<sup>31</sup> C. Diaconu,<sup>88</sup> M. Diamond,<sup>161</sup> F. A. Dias,<sup>39</sup> M. A. Diaz,<sup>34a</sup> J. Dickinson,<sup>16</sup> E. B. Diehl,<sup>92</sup> J. Dietrich,<sup>17</sup> S. Díez Cornell,<sup>45</sup> A. Dimitrievska,<sup>16</sup> J. Dingfelder,<sup>23</sup> P. Dita,<sup>28b</sup> S. Dita,<sup>28b</sup> F. Dittus,<sup>32</sup> F. Djama,<sup>88</sup> T. Djobava,<sup>54b</sup> J. I. Djuvsland,<sup>60a</sup> M. A. B. do Vale,<sup>26c</sup> M. Dobre,<sup>28b</sup> D. Dodsworth,<sup>25</sup> C. Doglioni,<sup>84</sup> J. Dolejsi,<sup>131</sup> Z. Dolezal,<sup>131</sup> M. Donadelli,<sup>26d</sup> S. Donati,<sup>126a,126b</sup> J. Donini,<sup>37</sup> J. Dopke,<sup>133</sup> A. Doria,<sup>106a</sup> M. T. Dova,<sup>74</sup> A. T. Doyle,<sup>56</sup> E. Drechsler,<sup>57</sup> M. Dris,<sup>10</sup> Y. Du,<sup>36a</sup> J. Duarte-Campderros,<sup>155</sup> F. Dubinin,<sup>98</sup> A. Dubreuil,<sup>52</sup> E. Duchovni,<sup>175</sup> G. Duckeck,<sup>102</sup> A. Ducourthial,<sup>83</sup> O. A. Ducu,<sup>97,q</sup> D. Duda,<sup>109</sup> A. Dudarev,<sup>32</sup> A. Chr. Dudder,<sup>86</sup> E. M. Duffield,<sup>16</sup> L. Dufлот,<sup>119</sup> M. Dührssen,<sup>32</sup> C. Dulsen,<sup>178</sup> M. Dumancic,<sup>175</sup> A. E. Dumitriu,<sup>28b</sup> A. K. Duncan,<sup>56</sup> M. Dunford,<sup>60a</sup> A. Duperrin,<sup>88</sup> H. Duran Yildiz,<sup>4a</sup> M. Düren,<sup>55</sup> A. Durglishvili,<sup>54b</sup> D. Duschinger,<sup>47</sup> B. Dutta,<sup>45</sup> D. Duvnjak,<sup>1</sup> M. Dyndal,<sup>45</sup> B. S. Dziedzic,<sup>42</sup> C. Eckardt,<sup>45</sup> K. M. Ecker,<sup>103</sup> R. C. Edgar,<sup>92</sup> T. Eifert,<sup>32</sup> G. Eigen,<sup>15</sup> K. Einsweiler,<sup>16</sup> T. Ekelof,<sup>168</sup> M. El Kacimi,<sup>137b</sup> R. El Kosseifi,<sup>88</sup> V. Ellajosyula,<sup>88</sup> M. Ellert,<sup>168</sup> S. Elles,<sup>5</sup> F. Ellinghaus,<sup>178</sup> A. A. Elliot,<sup>172</sup> N. Ellis,<sup>32</sup> J. Elmsheuser,<sup>27</sup> M. Elsing,<sup>32</sup> D. Emelianov,<sup>133</sup> Y. Enari,<sup>157</sup> J. S. Ennis,<sup>173</sup> M. B. Epland,<sup>48</sup> J. Erdmann,<sup>46</sup> A. Ereditato,<sup>18</sup> M. Ernst,<sup>27</sup> S. Errede,<sup>169</sup> M. Escalier,<sup>119</sup> C. Escobar,<sup>170</sup> B. Esposito,<sup>50</sup> O. Estrada Pastor,<sup>170</sup> A. I. Etienvre,<sup>138</sup> E. Etzion,<sup>155</sup> H. Evans,<sup>64</sup> A. Ezhilov,<sup>125</sup> M. Ezzi,<sup>137d</sup> F. Fabbri,<sup>22a,22b</sup> L. Fabbri,<sup>22a,22b</sup> V. Fabiani,<sup>108</sup> G. Facini,<sup>81</sup> R. M. Fakhruddinov,<sup>132</sup> S. Falciano,<sup>134a</sup> R. J. Falla,<sup>81</sup> J. Faltova,<sup>32</sup> Y. Fang,<sup>35a</sup> M. Fanti,<sup>94a,94b</sup> A. Farbin,<sup>8</sup> A. Farilla,<sup>136a</sup> E. M. Farina,<sup>123a,123b</sup> T. Farooque,<sup>93</sup> S. Farrell,<sup>16</sup> S. M. Farrington,<sup>173</sup> P. Farthouat,<sup>32</sup> F. Fassi,<sup>137d</sup> P. Fassnacht,<sup>32</sup> D. Fassouliotis,<sup>9</sup> M. Fauci Giannelli,<sup>49</sup> A. Favareto,<sup>53a,53b</sup> W. J. Fawcett,<sup>122</sup> L. Fayard,<sup>119</sup> O. L. Fedin,<sup>125,r</sup> W. Fedorko,<sup>171</sup> S. Feigl,<sup>121</sup> L. Feligioni,<sup>88</sup> C. Feng,<sup>36a</sup> E. J. Feng,<sup>32</sup> M. Feng,<sup>48</sup> M. J. Fenton,<sup>56</sup> A. B. Fenyuk,<sup>132</sup> L. Feremenga,<sup>8</sup> P. Fernandez Martinez,<sup>170</sup> J. Ferrando,<sup>45</sup> A. Ferrari,<sup>168</sup> P. Ferrari,<sup>109</sup> R. Ferrari,<sup>123a</sup> D. E. Ferreira de Lima,<sup>60b</sup> A. Ferrer,<sup>170</sup> D. Ferrere,<sup>52</sup> C. Ferretti,<sup>92</sup> F. Fiedler,<sup>86</sup> A. Filipčič,<sup>78</sup> M. Filipuzzi,<sup>45</sup> F. Filthaut,<sup>108</sup> M. Fincke-Keeler,<sup>172</sup> K. D. Finelli,<sup>24</sup> M. C. N. Fiolhais,<sup>128a,128c,s</sup> L. Fiorini,<sup>170</sup> C. Fischer,<sup>13</sup> J. Fischer,<sup>178</sup> W. C. Fisher,<sup>93</sup> N. Flaschel,<sup>45</sup> I. Fleck,<sup>143</sup> P. Fleischmann,<sup>92</sup> R. R. M. Fletcher,<sup>124</sup> T. Flick,<sup>178</sup> B. M. Flierl,<sup>102</sup> L. R. Flores Castillo,<sup>62a</sup> N. Fomin,<sup>15</sup> G. T. Forcolin,<sup>87</sup> A. Formica,<sup>138</sup> F. A. Förster,<sup>13</sup> A. Forti,<sup>87</sup> A. G. Foster,<sup>19</sup> D. Fournier,<sup>119</sup> H. Fox,<sup>75</sup> S. Fracchia,<sup>141</sup> P. Francavilla,<sup>126a,126b</sup> M. Franchini,<sup>22a,22b</sup> S. Franchino,<sup>60a</sup> D. Francis,<sup>32</sup> L. Franconi,<sup>121</sup> M. Franklin,<sup>59</sup> M. Frate,<sup>166</sup> M. Fraternali,<sup>123a,123b</sup> D. Freeborn,<sup>81</sup> S. M. Fressard-Batraneanu,<sup>32</sup> B. Freund,<sup>97</sup> W. S. Freund,<sup>26a</sup> D. Froidevaux,<sup>32</sup> J. A. Frost,<sup>122</sup> C. Fukunaga,<sup>158</sup> T. Fusayasu,<sup>104</sup> J. Fuster,<sup>170</sup> O. Gabizon,<sup>154</sup> A. Gabrielli,<sup>22a,22b</sup> A. Gabrielli,<sup>16</sup> G. P. Gach,<sup>41a</sup> S. Gadatsch,<sup>32</sup> S. Gadomski,<sup>80</sup> G. Gagliardi,<sup>53a,53b</sup> L. G. Gagnon,<sup>97</sup> C. Galea,<sup>108</sup> B. Galhardo,<sup>128a,128c</sup> E. J. Gallas,<sup>122</sup> B. J. Gallop,<sup>133</sup> P. Gallus,<sup>130</sup> G. Galster,<sup>39</sup> K. K. Gan,<sup>113</sup> S. Ganguly,<sup>175</sup> Y. Gao,<sup>77</sup> Y. S. Gao,<sup>145,h</sup> F. M. Garay Walls,<sup>34a</sup> C. García,<sup>170</sup> J. E. García Navarro,<sup>170</sup> J. A. García Pascual,<sup>35a</sup> M. Garcia-Sciveres,<sup>16</sup> R. W. Gardner,<sup>33</sup> N. Garelli,<sup>145</sup> V. Garonne,<sup>121</sup> A. Gascon Bravo,<sup>45</sup> K. Gasnikova,<sup>45</sup> C. Gatti,<sup>50</sup> A. Gaudiello,<sup>53a,53b</sup> G. Gaudio,<sup>123a</sup> I. L. Gavrilenko,<sup>98</sup> C. Gay,<sup>171</sup> G. Gaycken,<sup>23</sup> E. N. Gazis,<sup>10</sup> C. N. P. Gee,<sup>133</sup> J. Geisen,<sup>57</sup> M. Geisen,<sup>86</sup> M. P. Geisler,<sup>60a</sup> K. Gellerstedt,<sup>148a,148b</sup> C. Gemme,<sup>53a</sup> M. H. Genest,<sup>58</sup> C. Geng,<sup>92</sup> S. Gentile,<sup>134a,134b</sup> C. Gentsos,<sup>156</sup> S. George,<sup>80</sup> D. Gerbaudo,<sup>13</sup> G. Geßner,<sup>46</sup> S. Ghasemi,<sup>143</sup> M. Ghneimat,<sup>23</sup> B. Giacobbe,<sup>22a</sup> S. Giagu,<sup>134a,134b</sup> N. Giangiacomi,<sup>22a,22b</sup> P. Giannetti,<sup>126a</sup> S. M. Gibson,<sup>80</sup> M. Gignac,<sup>171</sup> M. Gilchriese,<sup>16</sup> D. Gillberg,<sup>31</sup> G. Gilles,<sup>178</sup> D. M. Gingrich,<sup>3,e</sup> M. P. Giordani,<sup>167a,167c</sup> F. M. Giorgi,<sup>22a</sup> P. F. Giraud,<sup>138</sup> P. Giromini,<sup>59</sup> G. Giugliarelli,<sup>167a,167c</sup> D. Giugni,<sup>94a</sup> F. Giulì,<sup>122</sup> M. Giuliani,<sup>60b</sup> B. K. Gjølsten,<sup>121</sup> S. Gkaitatzis,<sup>156</sup> I. Gkialas,<sup>9,t</sup> E. L. Gkougkousis,<sup>13</sup> P. Gkoutoumis,<sup>10</sup> L. K. Gladilin,<sup>101</sup> C. Glasman,<sup>85</sup> J. Glatzer,<sup>13</sup> P. C. F. Glaysheer,<sup>45</sup> A. Glazov,<sup>45</sup> M. Goblirsch-Kolb,<sup>25</sup> J. Godlewski,<sup>42</sup> S. Goldfarb,<sup>91</sup> T. Golling,<sup>52</sup> D. Golubkov,<sup>132</sup> A. Gomes,<sup>128a,128b,128d</sup> R. Gonçalves,<sup>128a</sup> R. Goncalves Gama,<sup>26a</sup> J. Goncalves Pinto Firmino Da Costa,<sup>138</sup> G. Gonella,<sup>51</sup> L. Gonella,<sup>19</sup> A. Gongadze,<sup>68</sup> F. Gonnella,<sup>19</sup> J. L. Gonski,<sup>59</sup> S. González de la Hoz,<sup>170</sup> S. Gonzalez-Sevilla,<sup>52</sup> L. Goossens,<sup>32</sup> P. A. Gorbounov,<sup>99</sup> H. A. Gordon,<sup>27</sup> B. Gorini,<sup>32</sup> E. Gorini,<sup>76a,76b</sup> A. Gorišek,<sup>78</sup> A. T. Goshaw,<sup>48</sup> C. Gössling,<sup>46</sup> M. I. Gostkin,<sup>68</sup> C. A. Gottardo,<sup>23</sup> C. R. Goudet,<sup>119</sup> D. Goujdami,<sup>137b</sup> A. G. Goussiou,<sup>140</sup> N. Govender,<sup>147b,u</sup> C. Goy,<sup>5</sup> E. Gozani,<sup>154</sup> I. Grabowska-Bold,<sup>41a</sup> P. O. J. Gradin,<sup>168</sup> E. C. Graham,<sup>77</sup> J. Gramling,<sup>166</sup> E. Gramstad,<sup>121</sup> S. Grancagnolo,<sup>17</sup> V. Gratchev,<sup>125</sup> P. M. Gravila,<sup>28e</sup> C. Gray,<sup>56</sup> H. M. Gray,<sup>16</sup> Z. D. Greenwood,<sup>82,v</sup> C. Greife,<sup>23</sup> K. Gregersen,<sup>81</sup> I. M. Gregor,<sup>45</sup> P. Grenier,<sup>145</sup> K. Grevtsov,<sup>5</sup> J. Griffiths,<sup>8</sup> A. A. Grillo,<sup>139</sup>

K. Grimm,<sup>75</sup> S. Grinstein,<sup>13,w</sup> Ph. Gris,<sup>37</sup> J.-F. Grivaz,<sup>119</sup> S. Groh,<sup>86</sup> E. Gross,<sup>175</sup> J. Grosse-Knetter,<sup>57</sup> G. C. Grossi,<sup>82</sup> Z. J. Grout,<sup>81</sup> A. Grummer,<sup>107</sup> L. Guan,<sup>92</sup> W. Guan,<sup>176</sup> J. Guenther,<sup>32</sup> F. Guescini,<sup>163a</sup> D. Guest,<sup>166</sup> O. Gueta,<sup>155</sup> B. Gui,<sup>113</sup> E. Guido,<sup>53a,53b</sup> T. Guillemain,<sup>5</sup> S. Guindon,<sup>32</sup> U. Gul,<sup>56</sup> C. Gumpert,<sup>32</sup> J. Guo,<sup>36b</sup> W. Guo,<sup>92</sup> Y. Guo,<sup>36c,x</sup> R. Gupta,<sup>43</sup> S. Gurbuz,<sup>20a</sup> G. Gustavino,<sup>115</sup> B. J. Gutelman,<sup>154</sup> P. Gutierrez,<sup>115</sup> N. G. Gutierrez Ortiz,<sup>81</sup> C. Gutsche,<sup>81</sup> C. Guyot,<sup>138</sup> M. P. Guzik,<sup>41a</sup> C. Gwenlan,<sup>122</sup> C. B. Gwilliam,<sup>77</sup> A. Haas,<sup>112</sup> C. Haber,<sup>16</sup> H. K. Hadavand,<sup>8</sup> N. Haddad,<sup>137d</sup> A. Hadeef,<sup>88</sup> S. Hageböck,<sup>23</sup> M. Hagihara,<sup>164</sup> H. Hakobyan,<sup>180,a</sup> M. Haleem,<sup>45</sup> J. Haley,<sup>116</sup> G. Halladjian,<sup>93</sup> G. D. Hallerwell,<sup>88</sup> K. Hamacher,<sup>178</sup> P. Hamal,<sup>117</sup> K. Hamano,<sup>172</sup> A. Hamilton,<sup>147a</sup> G. N. Hamity,<sup>141</sup> P. G. Hamnett,<sup>45</sup> K. Han,<sup>36c,y</sup> L. Han,<sup>36c</sup> S. Han,<sup>35a,35d</sup> K. Hanagaki,<sup>69,z</sup> K. Hanawa,<sup>157</sup> M. Hance,<sup>139</sup> D. M. Handl,<sup>102</sup> B. Haney,<sup>124</sup> P. Hanke,<sup>60a</sup> J. B. Hansen,<sup>39</sup> J. D. Hansen,<sup>39</sup> M. C. Hansen,<sup>23</sup> P. H. Hansen,<sup>39</sup> K. Hara,<sup>164</sup> A. S. Hard,<sup>176</sup> T. Harenberg,<sup>178</sup> F. Hariri,<sup>119</sup> S. Harkusha,<sup>95</sup> P. F. Harrison,<sup>173</sup> N. M. Hartmann,<sup>102</sup> Y. Hasegawa,<sup>142</sup> A. Hasib,<sup>49</sup> S. Hassani,<sup>138</sup> S. Haug,<sup>18</sup> R. Hauser,<sup>93</sup> L. Hauswald,<sup>47</sup> L. B. Havener,<sup>38</sup> M. Havranek,<sup>130</sup> C. M. Hawkes,<sup>19</sup> R. J. Hawkings,<sup>32</sup> D. Hayden,<sup>93</sup> C. P. Hays,<sup>122</sup> J. M. Hays,<sup>79</sup> H. S. Hayward,<sup>77</sup> S. J. Haywood,<sup>133</sup> T. Heck,<sup>86</sup> V. Hedberg,<sup>84</sup> L. Heelan,<sup>8</sup> S. Heer,<sup>23</sup> K. K. Heidegger,<sup>51</sup> S. Heim,<sup>45</sup> T. Heim,<sup>16</sup> B. Heinemann,<sup>45,aa</sup> J. J. Heinrich,<sup>102</sup> L. Heinrich,<sup>112</sup> C. Heinz,<sup>55</sup> J. Hejbal,<sup>129</sup> L. Helary,<sup>32</sup> A. Held,<sup>171</sup> S. Hellman,<sup>148a,148b</sup> C. Helsens,<sup>32</sup> R. C. W. Henderson,<sup>75</sup> Y. Heng,<sup>176</sup> S. Henkelmann,<sup>171</sup> A. M. Henriques Correia,<sup>32</sup> S. Henrot-Versille,<sup>119</sup> G. H. Herbert,<sup>17</sup> H. Herde,<sup>25</sup> V. Herget,<sup>177</sup> Y. Hernández Jiménez,<sup>147c</sup> H. Herr,<sup>86</sup> G. Herten,<sup>51</sup> R. Hertenberger,<sup>102</sup> L. Hervas,<sup>32</sup> T. C. Herwig,<sup>124</sup> G. G. Hesketh,<sup>81</sup> N. P. Hessey,<sup>163a</sup> J. W. Hetherly,<sup>43</sup> S. Higashino,<sup>69</sup> E. Higón-Rodríguez,<sup>170</sup> K. Hildebrand,<sup>33</sup> E. Hill,<sup>172</sup> J. C. Hill,<sup>30</sup> K. H. Hiller,<sup>45</sup> S. J. Hillier,<sup>19</sup> M. Hils,<sup>47</sup> I. Hinchliffe,<sup>16</sup> M. Hirose,<sup>51</sup> D. Hirschbuehl,<sup>178</sup> B. Hiti,<sup>78</sup> O. Hladik,<sup>129</sup> D. R. Hlaluku,<sup>147c</sup> X. Hoad,<sup>49</sup> J. Hobbs,<sup>150</sup> N. Hod,<sup>163a</sup> M. C. Hodgkinson,<sup>141</sup> P. Hodgson,<sup>141</sup> A. Hoecker,<sup>32</sup> M. R. Hoferkamp,<sup>107</sup> F. Hoenig,<sup>102</sup> D. Hohn,<sup>23</sup> T. R. Holmes,<sup>33</sup> M. Holzbock,<sup>102</sup> M. Homann,<sup>46</sup> S. Honda,<sup>164</sup> T. Honda,<sup>69</sup> T. M. Hong,<sup>127</sup> B. H. Hooberman,<sup>169</sup> W. H. Hopkins,<sup>118</sup> Y. Horii,<sup>105</sup> A. J. Horton,<sup>144</sup> J.-Y. Hostachy,<sup>58</sup> A. Hostiuc,<sup>140</sup> S. Hou,<sup>153</sup> A. Hoummada,<sup>137a</sup> J. Howarth,<sup>87</sup> J. Hoya,<sup>74</sup> M. Hrabovsky,<sup>117</sup> J. Hrdinka,<sup>32</sup> I. Hristova,<sup>17</sup> J. Hrivnac,<sup>119</sup> T. Hryn'ova,<sup>5</sup> A. Hrynevich,<sup>96</sup> P. J. Hsu,<sup>63</sup> S.-C. Hsu,<sup>140</sup> Q. Hu,<sup>27</sup> S. Hu,<sup>36b</sup> Y. Huang,<sup>35a</sup> Z. Hubacek,<sup>130</sup> F. Hubaut,<sup>88</sup> F. Huegging,<sup>23</sup> T. B. Huffman,<sup>122</sup> E. W. Hughes,<sup>38</sup> M. Huhtinen,<sup>32</sup> R. F. H. Hunter,<sup>31</sup> P. Huo,<sup>150</sup> N. Huseynov,<sup>68,c</sup> J. Huston,<sup>93</sup> J. Huth,<sup>59</sup> R. Hyneman,<sup>92</sup> G. Iacobucci,<sup>52</sup> G. Iakovidis,<sup>27</sup> I. Ibragimov,<sup>143</sup> L. Iconomidou-Fayard,<sup>119</sup> Z. Idrissi,<sup>137d</sup> P. Iengo,<sup>32</sup> O. Igonkina,<sup>109,bb</sup> T. Iizawa,<sup>174</sup> Y. Ikegami,<sup>69</sup> M. Ikeno,<sup>69</sup> Y. Ilchenko,<sup>11,cc</sup> D. Iliadis,<sup>156</sup> N. Ilic,<sup>145</sup> F. Iltzsche,<sup>47</sup> G. Introzzi,<sup>123a,123b</sup> P. Ioannou,<sup>9,a</sup> M. Iodice,<sup>136a</sup> K. Iordanidou,<sup>38</sup> V. Ippolito,<sup>59</sup> M. F. Isacson,<sup>168</sup> N. Ishijima,<sup>120</sup> M. Ishino,<sup>157</sup> M. Ishitsuka,<sup>159</sup> C. Issever,<sup>122</sup> S. Istin,<sup>20a</sup> F. Ito,<sup>164</sup> J. M. Iturbe Ponce,<sup>62a</sup> R. Iuppa,<sup>162a,162b</sup> H. Iwasaki,<sup>69</sup> J. M. Izen,<sup>44</sup> V. Izzo,<sup>106a</sup> S. Jabbar,<sup>3</sup> P. Jackson,<sup>1</sup> R. M. Jacobs,<sup>23</sup> V. Jain,<sup>2</sup> G. Jakel,<sup>178</sup> K. B. Jakobi,<sup>86</sup> K. Jakobs,<sup>51</sup> S. Jakobsen,<sup>65</sup> T. Jakoubek,<sup>129</sup> D. O. Jamin,<sup>116</sup> D. K. Jana,<sup>82</sup> R. Jansky,<sup>52</sup> J. Janssen,<sup>23</sup> M. Janus,<sup>57</sup> P. A. Janus,<sup>41a</sup> G. Jarlskog,<sup>84</sup> N. Javadov,<sup>68,c</sup> T. Javůrek,<sup>51</sup> M. Javurkova,<sup>51</sup> F. Jeanneau,<sup>138</sup> L. Jeanty,<sup>16</sup> J. Jejelava,<sup>54a,dd</sup> A. Jelinskas,<sup>173</sup> P. Jenni,<sup>51,ee</sup> C. Jeske,<sup>173</sup> S. Jézéquel,<sup>5</sup> H. Ji,<sup>176</sup> J. Jia,<sup>150</sup> H. Jiang,<sup>67</sup> Y. Jiang,<sup>36c</sup> Z. Jiang,<sup>145</sup> S. Jiggins,<sup>81</sup> J. Jimenez Pena,<sup>170</sup> S. Jin,<sup>35b</sup> A. Jinaru,<sup>28b</sup> O. Jinnouchi,<sup>159</sup> H. Jivan,<sup>147c</sup> P. Johansson,<sup>141</sup> K. A. Johns,<sup>7</sup> C. A. Johnson,<sup>64</sup> W. J. Johnson,<sup>140</sup> K. Jon-And,<sup>148a,148b</sup> R. W. L. Jones,<sup>75</sup> S. D. Jones,<sup>151</sup> S. Jones,<sup>7</sup> T. J. Jones,<sup>77</sup> J. Jongmanns,<sup>60a</sup> P. M. Jorge,<sup>128a,128b</sup> J. Jovicevic,<sup>163a</sup> X. Ju,<sup>176</sup> A. Juste Rozas,<sup>13,w</sup> A. Kaczmarska,<sup>42</sup> M. Kado,<sup>119</sup> H. Kagan,<sup>113</sup> M. Kagan,<sup>145</sup> S. J. Kahn,<sup>88</sup> T. Kaji,<sup>174</sup> E. Kajomovitz,<sup>154</sup> C. W. Kalderon,<sup>84</sup> A. Kaluza,<sup>86</sup> S. Kama,<sup>43</sup> A. Kamenshchikov,<sup>132</sup> N. Kanaya,<sup>157</sup> L. Kanjir,<sup>78</sup> Y. Kano,<sup>157</sup> V. A. Kantserov,<sup>100</sup> J. Kanzaki,<sup>69</sup> B. Kaplan,<sup>112</sup> L. S. Kaplan,<sup>176</sup> D. Kar,<sup>147c</sup> K. Karakostas,<sup>10</sup> N. Karastathis,<sup>10</sup> M. J. Kareem,<sup>163b</sup> E. Karentzos,<sup>10</sup> S. N. Karpov,<sup>68</sup> Z. M. Karpova,<sup>68</sup> V. Kartvelishvili,<sup>75</sup> A. N. Karyukhin,<sup>132</sup> K. Kasahara,<sup>164</sup> L. Kashif,<sup>176</sup> R. D. Kass,<sup>113</sup> A. Kastanas,<sup>149</sup> Y. Kataoka,<sup>157</sup> C. Kato,<sup>157</sup> A. Katre,<sup>52</sup> J. Katzy,<sup>45</sup> K. Kawade,<sup>70</sup> K. Kawagoe,<sup>73</sup> T. Kawamoto,<sup>157</sup> G. Kawamura,<sup>57</sup> E. F. Kay,<sup>77</sup> V. F. Kazanin,<sup>111,d</sup> R. Keeler,<sup>172</sup> R. Kehoe,<sup>43</sup> J. S. Keller,<sup>31</sup> E. Kellermann,<sup>84</sup> J. J. Kempster,<sup>80</sup> J. Kendrick,<sup>19</sup> H. Keoshkerian,<sup>161</sup> O. Kepka,<sup>129</sup> B. P. Kerševan,<sup>78</sup> S. Kersten,<sup>178</sup> R. A. Keyes,<sup>90</sup> M. Khader,<sup>169</sup> F. Khalil-zada,<sup>12</sup> A. Khanov,<sup>116</sup> A. G. Kharlamov,<sup>111,d</sup> T. Kharlamova,<sup>111,d</sup> A. Khodinov,<sup>160</sup> T. J. Khoo,<sup>52</sup> V. Khovanskii,<sup>99,a</sup> E. Khramov,<sup>68</sup> J. Khubua,<sup>54b,ff</sup> S. Kido,<sup>70</sup> M. Kiehn,<sup>52</sup> C. R. Kilby,<sup>80</sup> H. Y. Kim,<sup>8</sup> S. H. Kim,<sup>164</sup> Y. K. Kim,<sup>33</sup> N. Kimura,<sup>167a,167c</sup> O. M. Kind,<sup>17</sup> B. T. King,<sup>77</sup> D. Kirchmeier,<sup>47</sup> J. Kirk,<sup>133</sup> A. E. Kiryunin,<sup>103</sup> T. Kishimoto,<sup>157</sup> D. Kisielewska,<sup>41a</sup> V. Kitali,<sup>45</sup> O. Kivernyk,<sup>5</sup> E. Kladiva,<sup>146b</sup> T. Klapdor-Kleingrothaus,<sup>51</sup> M. H. Klein,<sup>92</sup> M. Klein,<sup>77</sup> U. Klein,<sup>77</sup> K. Kleinknecht,<sup>86</sup> P. Klimek,<sup>110</sup> A. Klimentov,<sup>27</sup> R. Klingenberg,<sup>46,a</sup> T. Klingl,<sup>23</sup> T. Klioutchnikova,<sup>32</sup> F. F. Klitzner,<sup>102</sup> E.-E. Kluge,<sup>60a</sup> P. Kluit,<sup>109</sup> S. Kluth,<sup>103</sup> E. Kneringer,<sup>65</sup> E. B. F. G. Knoops,<sup>88</sup> A. Knue,<sup>51</sup> A. Kobayashi,<sup>157</sup> D. Kobayashi,<sup>73</sup> T. Kobayashi,<sup>157</sup> M. Kobel,<sup>47</sup> M. Kocian,<sup>145</sup> P. Kodys,<sup>131</sup> T. Koffas,<sup>31</sup> E. Koffman,<sup>109</sup> N. M. Köhler,<sup>103</sup> T. Koi,<sup>145</sup> M. Kolb,<sup>60b</sup> I. Koletsou,<sup>5</sup> T. Kondo,<sup>69</sup> N. Kondrashova,<sup>36b</sup> K. Köneke,<sup>51</sup> A. C. König,<sup>108</sup> T. Kono,<sup>69,gg</sup> R. Konoplich,<sup>112,hh</sup> N. Konstantinidis,<sup>81</sup>



- B. Konya,<sup>84</sup> R. Kopeliansky,<sup>64</sup> S. Koperny,<sup>41a</sup> K. Korcyl,<sup>42</sup> K. Kordas,<sup>156</sup> A. Korn,<sup>81</sup> I. Korolkov,<sup>13</sup> E. V. Korolkova,<sup>141</sup> O. Kortner,<sup>103</sup> S. Kortner,<sup>103</sup> T. Kosek,<sup>131</sup> V. V. Kostyukhin,<sup>23</sup> A. Kotwal,<sup>48</sup> A. Koulouris,<sup>10</sup> A. Kourkouveli-Charalampidi,<sup>123a,123b</sup> C. Kourkouvelis,<sup>9</sup> E. Kourlitis,<sup>141</sup> V. Kouskoura,<sup>27</sup> A. B. Kowalewska,<sup>42</sup> R. Kowalewski,<sup>172</sup> T. Z. Kowalski,<sup>41a</sup> C. Kozakai,<sup>157</sup> W. Kozanecki,<sup>138</sup> A. S. Kozhin,<sup>132</sup> V. A. Kramarenko,<sup>101</sup> G. Kramberger,<sup>78</sup> D. Krasnopevtsev,<sup>100</sup> M. W. Krasny,<sup>83</sup> A. Krasznahorkay,<sup>32</sup> D. Krauss,<sup>103</sup> J. A. Kremer,<sup>41a</sup> J. Kretschmar,<sup>77</sup> K. Kreutzfeldt,<sup>55</sup> P. Krieger,<sup>161</sup> K. Krizka,<sup>16</sup> K. Kroeninger,<sup>46</sup> H. Kroha,<sup>103</sup> J. Kroll,<sup>129</sup> J. Kroll,<sup>124</sup> J. Kroseberg,<sup>23</sup> J. Krstic,<sup>14</sup> U. Kruchonak,<sup>68</sup> H. Krüger,<sup>23</sup> N. Krumnack,<sup>67</sup> M. C. Kruse,<sup>48</sup> T. Kubota,<sup>91</sup> H. Kucuk,<sup>81</sup> S. Kudah,<sup>4b</sup> J. T. Kuechler,<sup>178</sup> S. Kuehn,<sup>32</sup> A. Kugel,<sup>60a</sup> F. Kuger,<sup>177</sup> T. Kuhl,<sup>45</sup> V. Kukhtin,<sup>68</sup> R. Kukla,<sup>88</sup> Y. Kulchitsky,<sup>95</sup> S. Kuleshov,<sup>34b</sup> Y. P. Kulinich,<sup>169</sup> M. Kuna,<sup>11</sup> T. Kunigo,<sup>71</sup> A. Kupco,<sup>129</sup> T. Kupfer,<sup>46</sup> O. Kuprash,<sup>155</sup> H. Kurashige,<sup>70</sup> L. L. Kurchaninov,<sup>163a</sup> Y. A. Kurochkin,<sup>95</sup> M. G. Kurth,<sup>35a,35d</sup> E. S. Kuwertz,<sup>172</sup> M. Kuze,<sup>159</sup> J. Kvita,<sup>117</sup> T. Kwan,<sup>172</sup> A. La Rosa,<sup>103</sup> J. L. La Rosa Navarro,<sup>26d</sup> L. La Rotonda,<sup>40a,40b</sup> F. La Ruffa,<sup>40a,40b</sup> C. Lacasta,<sup>170</sup> F. Lacava,<sup>134a,134b</sup> J. Lacey,<sup>45</sup> D. P. J. Lack,<sup>87</sup> H. Lacker,<sup>17</sup> D. Lacour,<sup>83</sup> E. Ladygin,<sup>68</sup> R. Lafaye,<sup>5</sup> B. Laforge,<sup>83</sup> S. Lai,<sup>57</sup> S. Lammers,<sup>64</sup> W. Lampl,<sup>7</sup> E. Lançon,<sup>27</sup> U. Landgraf,<sup>51</sup> M. P. J. Landon,<sup>79</sup> M. C. Lanfermann,<sup>52</sup> V. S. Lang,<sup>45</sup> J. C. Lange,<sup>13</sup> R. J. Langenberg,<sup>32</sup> A. J. Lankford,<sup>166</sup> F. Lanni,<sup>27</sup> K. Lantzsch,<sup>23</sup> A. Lanza,<sup>123a</sup> A. Lapertosa,<sup>53a,53b</sup> S. Laplace,<sup>83</sup> J. F. Laporte,<sup>138</sup> T. Lari,<sup>94a</sup> F. Lasagni Manghi,<sup>22a,22b</sup> M. Lassnig,<sup>32</sup> T. S. Lau,<sup>62a</sup> P. Laurelli,<sup>50</sup> W. Lavrijsen,<sup>16</sup> A. T. Law,<sup>139</sup> P. Laycock,<sup>77</sup> T. Lazovich,<sup>59</sup> M. Lazzaroni,<sup>94a,94b</sup> B. Le,<sup>91</sup> O. Le Dortz,<sup>83</sup> E. Le Guirriec,<sup>88</sup> E. P. Le Quilleuc,<sup>138</sup> M. LeBlanc,<sup>7</sup> T. LeCompte,<sup>6</sup> F. Ledroit-Guillon,<sup>58</sup> C. A. Lee,<sup>27</sup> G. R. Lee,<sup>34a</sup> S. C. Lee,<sup>153</sup> L. Lee,<sup>59</sup> B. Lefebvre,<sup>90</sup> G. Lefebvre,<sup>83</sup> M. Lefebvre,<sup>172</sup> F. Legger,<sup>102</sup> C. Leggett,<sup>16</sup> G. Lehmann Miotto,<sup>32</sup> X. Lei,<sup>7</sup> W. A. Leight,<sup>45</sup> M. A. L. Leite,<sup>26d</sup> R. Leitner,<sup>131</sup> D. Lellouch,<sup>175</sup> B. Lemmer,<sup>57</sup> K. J. C. Leney,<sup>81</sup> T. Lenz,<sup>23</sup> B. Lenzi,<sup>32</sup> R. Leone,<sup>7</sup> S. Leone,<sup>126a</sup> C. Leonidopoulos,<sup>49</sup> G. Lerner,<sup>151</sup> C. Leroy,<sup>97</sup> R. Les,<sup>161</sup> A. A. J. Lesage,<sup>138</sup> C. G. Lester,<sup>30</sup> M. Levchenko,<sup>125</sup> J. Levêque,<sup>5</sup> D. Levin,<sup>92</sup> L. J. Levinson,<sup>175</sup> M. Levy,<sup>19</sup> D. Lewis,<sup>79</sup> B. Li,<sup>36c,x</sup> Changqiao Li,<sup>36c</sup> H. Li,<sup>150</sup> L. Li,<sup>36b</sup> Q. Li,<sup>35a,35d</sup> Q. Li,<sup>36c</sup> S. Li,<sup>48</sup> X. Li,<sup>36b</sup> Y. Li,<sup>143</sup> Z. Liang,<sup>35a</sup> B. Liberti,<sup>135a</sup> A. Liblong,<sup>161</sup> K. Lie,<sup>62c</sup> A. Limosani,<sup>152</sup> C. Y. Lin,<sup>30</sup> K. Lin,<sup>93</sup> S. C. Lin,<sup>182</sup> T. H. Lin,<sup>86</sup> R. A. Linck,<sup>64</sup> B. E. Lindquist,<sup>150</sup> A. E. Lioni,<sup>52</sup> E. Lipeles,<sup>124</sup> A. Lipniacka,<sup>15</sup> M. Lisovsky,<sup>60b</sup> T. M. Liss,<sup>169,ii</sup> A. Lister,<sup>171</sup> A. M. Litke,<sup>139</sup> B. Liu,<sup>67</sup> H. Liu,<sup>92</sup> H. Liu,<sup>27</sup> J. K. K. Liu,<sup>122</sup> J. Liu,<sup>36a</sup> J. B. Liu,<sup>36c</sup> K. Liu,<sup>88</sup> L. Liu,<sup>169</sup> M. Liu,<sup>36c</sup> Y. L. Liu,<sup>36c</sup> Y. Liu,<sup>36c</sup> M. Livan,<sup>123a,123b</sup> A. Lleres,<sup>58</sup> J. Llorente Merino,<sup>35a</sup> S. L. Lloyd,<sup>79</sup> C. Y. Lo,<sup>62b</sup> F. Lo Sterzo,<sup>43</sup> E. M. Lobodzinska,<sup>45</sup> P. Loch,<sup>7</sup> F. K. Loebinger,<sup>87</sup> A. Loesle,<sup>51</sup> K. M. Loew,<sup>25</sup> T. Lohse,<sup>17</sup> K. Lohwasser,<sup>141</sup> M. Lokajicek,<sup>129</sup> B. A. Long,<sup>24</sup> J. D. Long,<sup>169</sup> R. E. Long,<sup>75</sup> L. Longo,<sup>76a,76b</sup> K. A. Looper,<sup>113</sup> J. A. Lopez,<sup>34b</sup> I. Lopez Paz,<sup>13</sup> A. Lopez Solis,<sup>83</sup> J. Lorenz,<sup>102</sup> N. Lorenzo Martinez,<sup>5</sup> M. Losada,<sup>21</sup> P. J. Lösel,<sup>102</sup> X. Lou,<sup>35a</sup> A. Lounis,<sup>119</sup> J. Love,<sup>6</sup> P. A. Love,<sup>75</sup> H. Lu,<sup>62a</sup> N. Lu,<sup>92</sup> Y. J. Lu,<sup>63</sup> H. J. Lubatti,<sup>140</sup> C. Luci,<sup>134a,134b</sup> A. Lucotte,<sup>58</sup> C. Luedtke,<sup>51</sup> F. Luehring,<sup>64</sup> W. Lukas,<sup>65</sup> L. Luminari,<sup>134a</sup> B. Lund-Jensen,<sup>149</sup> M. S. Lutz,<sup>89</sup> P. M. Luzi,<sup>83</sup> D. Lynn,<sup>27</sup> R. Lysak,<sup>129</sup> E. Lytken,<sup>84</sup> F. Lyu,<sup>35a</sup> V. Lyubushkin,<sup>68</sup> H. Ma,<sup>27</sup> L. L. Ma,<sup>36a</sup> Y. Ma,<sup>36a</sup> G. Maccarrone,<sup>50</sup> A. Macchiolo,<sup>103</sup> C. M. Macdonald,<sup>141</sup> B. Maček,<sup>78</sup> J. Machado Miguens,<sup>124,128b</sup> D. Madaffari,<sup>170</sup> R. Madar,<sup>37</sup> W. F. Mader,<sup>47</sup> A. Madsen,<sup>45</sup> N. Madysa,<sup>47</sup> J. Maeda,<sup>70</sup> S. Maeland,<sup>15</sup> T. Maeno,<sup>27</sup> A. S. Maevskiy,<sup>101</sup> V. Magerl,<sup>51</sup> C. Maiani,<sup>119</sup> C. Maidantchik,<sup>26a</sup> T. Maier,<sup>102</sup> A. Maio,<sup>128a,128b,128d</sup> O. Majersky,<sup>146a</sup> S. Majewski,<sup>118</sup> Y. Makida,<sup>69</sup> N. Makovec,<sup>119</sup> B. Malaescu,<sup>83</sup> Pa. Malecki,<sup>42</sup> V. P. Maleev,<sup>125</sup> F. Malek,<sup>58</sup> U. Mallik,<sup>66</sup> D. Malon,<sup>6</sup> C. Malone,<sup>30</sup> S. Maltezos,<sup>10</sup> S. Malyukov,<sup>32</sup> J. Mamuzic,<sup>170</sup> G. Mancini,<sup>50</sup> I. Mandić,<sup>78</sup> J. Maneira,<sup>128a,128b</sup> L. Manhaes de Andrade Filho,<sup>26b</sup> J. Manjarres Ramos,<sup>47</sup> K. H. Mankinen,<sup>84</sup> A. Mann,<sup>102</sup> A. Manousos,<sup>32</sup> B. Mansoulie,<sup>138</sup> J. D. Mansour,<sup>35a</sup> R. Mantifel,<sup>90</sup> M. Mantoani,<sup>57</sup> S. Manzoni,<sup>94a,94b</sup> L. Mapelli,<sup>32</sup> G. Marceca,<sup>29</sup> L. March,<sup>52</sup> L. Marchese,<sup>122</sup> G. Marchiori,<sup>83</sup> M. Marcisovsky,<sup>129</sup> C. A. Marin Tobon,<sup>32</sup> M. Marjanovic,<sup>37</sup> D. E. Marley,<sup>92</sup> F. Marroquim,<sup>26a</sup> S. P. Marsden,<sup>87</sup> Z. Marshall,<sup>16</sup> M. U. F. Martensson,<sup>168</sup> S. Marti-Garcia,<sup>170</sup> C. B. Martin,<sup>113</sup> T. A. Martin,<sup>173</sup> V. J. Martin,<sup>49</sup> B. Martin dit Latour,<sup>15</sup> M. Martinez,<sup>13,w</sup> V. I. Martinez Outschoorn,<sup>169</sup> S. Martin-Haugh,<sup>133</sup> V. S. Martoiu,<sup>28b</sup> A. C. Martyniuk,<sup>81</sup> A. Marzin,<sup>32</sup> L. Masetti,<sup>86</sup> T. Mashimo,<sup>157</sup> R. Mashinistov,<sup>98</sup> J. Masik,<sup>87</sup> A. L. Maslennikov,<sup>111,d</sup> L. H. Mason,<sup>91</sup> L. Massa,<sup>135a,135b</sup> P. Mastrandrea,<sup>5</sup> A. Mastroberardino,<sup>40a,40b</sup> T. Masubuchi,<sup>157</sup> P. Mättig,<sup>178</sup> J. Maurer,<sup>28b</sup> S. J. Maxfield,<sup>77</sup> D. A. Maximov,<sup>111,d</sup> R. Mazini,<sup>153</sup> I. Maznas,<sup>156</sup> S. M. Mazza,<sup>94a,94b</sup> N. C. Mc Fadden,<sup>107</sup> G. Mc Goldrick,<sup>161</sup> S. P. Mc Kee,<sup>92</sup> A. McCarn,<sup>92</sup> R. L. McCarthy,<sup>150</sup> T. G. McCarthy,<sup>103</sup> L. I. McClymont,<sup>81</sup> E. F. McDonald,<sup>91</sup> J. A. Mcfayden,<sup>32</sup> G. Mchedlidze,<sup>57</sup> S. J. McMahon,<sup>133</sup> P. C. McNamara,<sup>91</sup> C. J. McNicol,<sup>173</sup> R. A. McPherson,<sup>172,p</sup> Z. A. Meadows,<sup>89</sup> S. Meehan,<sup>140</sup> T. J. Megy,<sup>51</sup> S. Mehlhase,<sup>102</sup> A. Mehta,<sup>77</sup> T. Meideck,<sup>58</sup> K. Meier,<sup>60a</sup> B. Meirose,<sup>44</sup> D. Melini,<sup>170,jj</sup> B. R. Mellado Garcia,<sup>147c</sup> J. D. Mellenthin,<sup>57</sup> M. Melo,<sup>146a</sup> F. Meloni,<sup>18</sup> A. Melzer,<sup>23</sup> S. B. Menary,<sup>87</sup> L. Meng,<sup>77</sup> X. T. Meng,<sup>92</sup> A. Mengarelli,<sup>22a,22b</sup> S. Menke,<sup>103</sup> E. Meoni,<sup>40a,40b</sup> S. Mergelmeyer,<sup>17</sup> C. Merlassino,<sup>18</sup> P. Mermod,<sup>52</sup> L. Merola,<sup>106a,106b</sup> C. Meroni,<sup>94a</sup> F. S. Merritt,<sup>33</sup> A. Messina,<sup>134a,134b</sup> J. Metcalfe,<sup>6</sup>

A. S. Mete,<sup>166</sup> C. Meyer,<sup>124</sup> J.-P. Meyer,<sup>138</sup> J. Meyer,<sup>109</sup> H. Meyer Zu Theenhausen,<sup>60a</sup> F. Miano,<sup>151</sup> R. P. Middleton,<sup>133</sup> S. Miglioranza,<sup>53a,53b</sup> L. Mijović,<sup>49</sup> G. Mikenberg,<sup>175</sup> M. Mikesstikova,<sup>129</sup> M. Mikuž,<sup>78</sup> M. Milesi,<sup>91</sup> A. Milic,<sup>161</sup> D. A. Millar,<sup>79</sup> D. W. Miller,<sup>33</sup> A. Milov,<sup>175</sup> D. A. Milstead,<sup>148a,148b</sup> A. A. Minaenko,<sup>132</sup> Y. Minami,<sup>157</sup> I. A. Minashvili,<sup>54b</sup> A. I. Mincer,<sup>112</sup> B. Mindur,<sup>41a</sup> M. Mineev,<sup>68</sup> Y. Minegishi,<sup>157</sup> Y. Ming,<sup>176</sup> L. M. Mir,<sup>13</sup> A. Mirto,<sup>76a,76b</sup> K. P. Mistry,<sup>124</sup> T. Mitani,<sup>174</sup> J. Mitrevski,<sup>102</sup> V. A. Mitsou,<sup>170</sup> A. Miucci,<sup>18</sup> P. S. Miyagawa,<sup>141</sup> A. Mizukami,<sup>69</sup> J. U. Mjörnmark,<sup>84</sup> T. Mkrchyan,<sup>180</sup> M. Mlynarikova,<sup>131</sup> T. Moa,<sup>148a,148b</sup> K. Mochizuki,<sup>97</sup> P. Mogg,<sup>51</sup> S. Mohapatra,<sup>38</sup> S. Molander,<sup>148a,148b</sup> R. Moles-Valls,<sup>23</sup> M. C. Mondragon,<sup>93</sup> K. Mönig,<sup>45</sup> J. Monk,<sup>39</sup> E. Monnier,<sup>88</sup> A. Montalbano,<sup>150</sup> J. Montejo Berlingen,<sup>32</sup> F. Monticelli,<sup>74</sup> S. Monzani,<sup>94a</sup> R. W. Moore,<sup>3</sup> N. Morange,<sup>119</sup> D. Moreno,<sup>21</sup> M. Moreno Llacer,<sup>32</sup> P. Morettini,<sup>53a</sup> M. Morgenstern,<sup>109</sup> S. Morgenstern,<sup>32</sup> D. Mori,<sup>144</sup> T. Mori,<sup>157</sup> M. Morii,<sup>59</sup> M. Morinaga,<sup>174</sup> V. Morisbak,<sup>121</sup> A. K. Morley,<sup>32</sup> G. Mornacchi,<sup>32</sup> J. D. Morris,<sup>79</sup> L. Morvaj,<sup>150</sup> P. Moschovakos,<sup>10</sup> M. Mosidze,<sup>54b</sup> H. J. Moss,<sup>141</sup> J. Moss,<sup>145,kk</sup> K. Motohashi,<sup>159</sup> R. Mount,<sup>145</sup> E. Mountricha,<sup>27</sup> E. J. W. Moyse,<sup>89</sup> S. Muanza,<sup>88</sup> F. Mueller,<sup>103</sup> J. Mueller,<sup>127</sup> R. S. P. Mueller,<sup>102</sup> D. Muenstermann,<sup>75</sup> P. Mullen,<sup>56</sup> G. A. Mullier,<sup>18</sup> F. J. Munoz Sanchez,<sup>87</sup> W. J. Murray,<sup>173,133</sup> H. Musheghyan,<sup>32</sup> M. Muškinja,<sup>78</sup> C. Mwewa,<sup>147a</sup> A. G. Myagkov,<sup>132,ll</sup> J. Myers,<sup>118</sup> M. Myska,<sup>130</sup> B. P. Nachman,<sup>16</sup> O. Nackenhorst,<sup>46</sup> K. Nagai,<sup>122</sup> R. Nagai,<sup>69,gg</sup> K. Nagano,<sup>69</sup> Y. Nagasaka,<sup>61</sup> K. Nagata,<sup>164</sup> M. Nagel,<sup>51</sup> E. Nagy,<sup>88</sup> A. M. Nairz,<sup>32</sup> Y. Nakahama,<sup>105</sup> K. Nakamura,<sup>69</sup> T. Nakamura,<sup>157</sup> I. Nakano,<sup>114</sup> R. F. Naranjo Garcia,<sup>45</sup> R. Narayan,<sup>11</sup> D. I. Narrias Villar,<sup>60a</sup> I. Naryshkin,<sup>125</sup> T. Naumann,<sup>45</sup> G. Navarro,<sup>21</sup> R. Nayyar,<sup>7</sup> H. A. Neal,<sup>92</sup> P. Yu. Nechaeva,<sup>98</sup> T. J. Neep,<sup>138</sup> A. Negri,<sup>123a,123b</sup> M. Negrini,<sup>22a</sup> S. Nektarijevic,<sup>108</sup> C. Nellist,<sup>57</sup> A. Nelson,<sup>166</sup> M. E. Nelson,<sup>122</sup> S. Nemecek,<sup>129</sup> P. Nemethy,<sup>112</sup> M. Nessi,<sup>32,mm</sup> M. S. Neubauer,<sup>169</sup> M. Neumann,<sup>178</sup> P. R. Newman,<sup>19</sup> T. Y. Ng,<sup>62c</sup> Y. S. Ng,<sup>17</sup> T. Nguyen Manh,<sup>97</sup> R. B. Nickerson,<sup>122</sup> R. Nicolaidou,<sup>138</sup> J. Nielsen,<sup>139</sup> N. Nikiforou,<sup>11</sup> V. Nikolaenko,<sup>132,ll</sup> I. Nikolic-Audit,<sup>83</sup> K. Nikolopoulos,<sup>19</sup> P. Nilsson,<sup>27</sup> Y. Ninomiya,<sup>69</sup> A. Nisati,<sup>134a</sup> N. Nishu,<sup>36b</sup> R. Nisius,<sup>103</sup> I. Nitsche,<sup>46</sup> T. Nitta,<sup>174</sup> T. Nobe,<sup>157</sup> Y. Noguchi,<sup>71</sup> M. Nomachi,<sup>120</sup> I. Nomidis,<sup>31</sup> M. A. Nomura,<sup>27</sup> T. Nooney,<sup>79</sup> M. Nordberg,<sup>32</sup> N. Norjoharuddeen,<sup>122</sup> O. Novgorodova,<sup>47</sup> R. Novotny,<sup>130</sup> M. Nozaki,<sup>69</sup> L. Nozka,<sup>117</sup> K. Ntekas,<sup>166</sup> E. Nurse,<sup>81</sup> F. Nuti,<sup>91</sup> K. O'connor,<sup>25</sup> D. C. O'Neil,<sup>144</sup> A. A. O'Rourke,<sup>45</sup> V. O'Shea,<sup>56</sup> F. G. Oakham,<sup>31,e</sup> H. Oberlack,<sup>103</sup> T. Obermann,<sup>23</sup> J. Ocariz,<sup>83</sup> A. Ochi,<sup>70</sup> I. Ochoa,<sup>38</sup> J. P. Ochoa-Ricoux,<sup>34a</sup> S. Oda,<sup>73</sup> S. Odaka,<sup>69</sup> A. Oh,<sup>87</sup> S. H. Oh,<sup>48</sup> C. C. Ohm,<sup>149</sup> H. Ohman,<sup>168</sup> H. Oide,<sup>53a,53b</sup> H. Okawa,<sup>164</sup> Y. Okumura,<sup>157</sup> T. Okuyama,<sup>69</sup> A. Olariu,<sup>28b</sup> L. F. Oleiro Seabra,<sup>128a</sup> S. A. Olivares Pino,<sup>34a</sup> D. Oliveira Damazio,<sup>27</sup> J. L. Oliver,<sup>1</sup> M. J. R. Olsson,<sup>33</sup> A. Olszewski,<sup>42</sup> J. Olszowska,<sup>42</sup> A. Onofre,<sup>128a,128e</sup> K. Onogi,<sup>105</sup> P. U. E. Onyisi,<sup>11,cc</sup> H. Oppen,<sup>121</sup> M. J. Oreglia,<sup>33</sup> Y. Oren,<sup>155</sup> D. Orestano,<sup>136a,136b</sup> E. C. Orgill,<sup>87</sup> N. Orlando,<sup>62b</sup> R. S. Orr,<sup>161</sup> B. Osculati,<sup>53a,53b,a</sup> R. Ospanov,<sup>36c</sup> G. Otero y Garzon,<sup>29</sup> H. Otono,<sup>73</sup> M. Ouchrif,<sup>137c</sup> F. Ould-Saada,<sup>121</sup> A. Ouraou,<sup>138</sup> K. P. Oussoren,<sup>109</sup> Q. Ouyang,<sup>35a</sup> M. Owen,<sup>56</sup> R. E. Owen,<sup>19</sup> V. E. Ozcan,<sup>20a</sup> N. Ozturk,<sup>8</sup> K. Pachal,<sup>144</sup> A. Pacheco Pages,<sup>13</sup> L. Pacheco Rodriguez,<sup>138</sup> C. Padilla Aranda,<sup>13</sup> S. Pagan Griso,<sup>16</sup> M. Paganini,<sup>179</sup> F. Paige,<sup>27</sup> G. Palacino,<sup>64</sup> S. Palazzo,<sup>40a,40b</sup> S. Palestini,<sup>32</sup> M. Palka,<sup>41b</sup> D. Pallin,<sup>37</sup> E. St. Panagiotopoulou,<sup>10</sup> I. Panagoulas,<sup>10</sup> C. E. Pandini,<sup>52</sup> J. G. Panduro Vazquez,<sup>80</sup> P. Pani,<sup>32</sup> S. Panitkin,<sup>27</sup> D. Pantea,<sup>28b</sup> L. Paolozzi,<sup>52</sup> Th. D. Papadopoulos,<sup>10</sup> K. Papageorgiou,<sup>9,t</sup> A. Paramonov,<sup>6</sup> D. Paredes Hernandez,<sup>179</sup> T. H. Park,<sup>31</sup> A. J. Parker,<sup>75</sup> M. A. Parker,<sup>30</sup> K. A. Parker,<sup>45</sup> F. Parodi,<sup>53a,53b</sup> J. A. Parsons,<sup>38</sup> U. Parzefall,<sup>51</sup> V. R. Pascuzzi,<sup>161</sup> J. M. Pasner,<sup>139</sup> E. Pasqualucci,<sup>134a</sup> S. Passaggio,<sup>53a</sup> Fr. Pastore,<sup>80</sup> S. Pataria,<sup>86</sup> J. R. Pater,<sup>87</sup> T. Pauly,<sup>32</sup> B. Pearson,<sup>103</sup> S. Pedraza Lopez,<sup>170</sup> R. Pedro,<sup>128a,128b</sup> S. V. Peleganchuk,<sup>111,d</sup> O. Penc,<sup>129</sup> C. Peng,<sup>35a,35d</sup> H. Peng,<sup>36c</sup> J. Penwell,<sup>64</sup> B. S. Peralva,<sup>26b</sup> M. M. Perego,<sup>138</sup> D. V. Perepelitsa,<sup>27</sup> F. Peri,<sup>17</sup> L. Perini,<sup>94a,94b</sup> H. Pernegger,<sup>32</sup> S. Perrella,<sup>106a,106b</sup> R. Peschke,<sup>45</sup> V. D. Peshekhonov,<sup>68,a</sup> K. Peters,<sup>45</sup> R. F. Y. Peters,<sup>87</sup> B. A. Petersen,<sup>32</sup> T. C. Petersen,<sup>39</sup> E. Petit,<sup>58</sup> A. Petridis,<sup>1</sup> C. Petridou,<sup>156</sup> P. Petroff,<sup>119</sup> E. Petrolo,<sup>134a</sup> M. Petrov,<sup>122</sup> F. Petrucci,<sup>136a,136b</sup> N. E. Pettersson,<sup>89</sup> A. Peyaud,<sup>138</sup> R. Pezoa,<sup>34b</sup> T. Pham,<sup>91</sup> F. H. Phillips,<sup>93</sup> P. W. Phillips,<sup>133</sup> G. Piacquadio,<sup>150</sup> E. Pianori,<sup>173</sup> A. Picazio,<sup>89</sup> M. A. Pickering,<sup>122</sup> R. Piegaia,<sup>29</sup> J. E. Pilcher,<sup>33</sup> A. D. Pilkington,<sup>87</sup> M. Pinamonti,<sup>135a,135b</sup> J. L. Pinfold,<sup>3</sup> H. Pirumov,<sup>45</sup> M. Pitt,<sup>175</sup> L. Plazak,<sup>146a</sup> M.-A. Pleier,<sup>27</sup> V. Pleskot,<sup>86</sup> E. Plotnikova,<sup>68</sup> D. Pluth,<sup>67</sup> P. Podberezko,<sup>111</sup> R. Poettgen,<sup>84</sup> R. Poggi,<sup>123a,123b</sup> L. Poggioli,<sup>119</sup> I. Pogrebnyak,<sup>93</sup> D. Pohl,<sup>23</sup> I. Pokharel,<sup>57</sup> G. Polesello,<sup>123a</sup> A. Poley,<sup>45</sup> A. Policicchio,<sup>40a,40b</sup> R. Polifka,<sup>32</sup> A. Polini,<sup>22a</sup> C. S. Pollard,<sup>45</sup> V. Polychronakos,<sup>27</sup> K. Pommès,<sup>32</sup> D. Ponomarenko,<sup>100</sup> L. Pontecorvo,<sup>134a</sup> G. A. Popeneciu,<sup>28d</sup> D. M. Portillo Quintero,<sup>83</sup> S. Pospisil,<sup>130</sup> K. Potamianos,<sup>45</sup> I. N. Potrap,<sup>68</sup> C. J. Potter,<sup>30</sup> H. Potti,<sup>11</sup> T. Poulsen,<sup>84</sup> J. Poveda,<sup>32</sup> M. E. Pozo Astigarraga,<sup>32</sup> P. Pralavorio,<sup>88</sup> A. Pranko,<sup>16</sup> S. Prell,<sup>67</sup> D. Price,<sup>87</sup> M. Primavera,<sup>76a</sup> S. Prince,<sup>90</sup> N. Proklova,<sup>100</sup> K. Prokofiev,<sup>62c</sup> F. Prokoshin,<sup>34b</sup> S. Protopopescu,<sup>27</sup> J. Proudfoot,<sup>6</sup> M. Przybycien,<sup>41a</sup> A. Puri,<sup>169</sup> P. Puzo,<sup>119</sup> J. Qian,<sup>92</sup> Y. Qin,<sup>87</sup> A. Quadri,<sup>57</sup> M. Queitsch-Maitland,<sup>45</sup> D. Quilty,<sup>56</sup> S. Raddum,<sup>121</sup> V. Radeka,<sup>27</sup> V. Radescu,<sup>122</sup> S. K. Radhakrishnan,<sup>150</sup> P. Radloff,<sup>118</sup> P. Rados,<sup>91</sup> F. Ragusa,<sup>94a,94b</sup> G. Rahal,<sup>181</sup> J. A. Raine,<sup>87</sup> S. Rajagopalan,<sup>27</sup> T. Rashid,<sup>119</sup>

- S. Raspopov,<sup>5</sup> M. G. Ratti,<sup>94a,94b</sup> D. M. Rauch,<sup>45</sup> F. Rauscher,<sup>102</sup> S. Rave,<sup>86</sup> I. Ravinovich,<sup>175</sup> J. H. Rawling,<sup>87</sup> M. Raymond,<sup>32</sup> A. L. Read,<sup>121</sup> N. P. Readioff,<sup>58</sup> M. Reale,<sup>76a,76b</sup> D. M. Rebutti,<sup>123a,123b</sup> A. Redelbach,<sup>177</sup> G. Redlinger,<sup>27</sup> R. Reece,<sup>139</sup> R. G. Reed,<sup>147c</sup> K. Reeves,<sup>44</sup> L. Rehnisch,<sup>17</sup> J. Reichert,<sup>124</sup> A. Reiss,<sup>86</sup> C. Rembser,<sup>32</sup> H. Ren,<sup>35a,35d</sup> M. Rescigno,<sup>134a</sup> S. Resconi,<sup>94a</sup> E. D. Resseguie,<sup>124</sup> S. Rettie,<sup>171</sup> E. Reynolds,<sup>19</sup> O. L. Rezanova,<sup>111,d</sup> P. Reznicek,<sup>131</sup> R. Rezvani,<sup>97</sup> R. Richter,<sup>103</sup> S. Richter,<sup>81</sup> E. Richter-Was,<sup>41b</sup> O. Ricken,<sup>23</sup> M. Ridel,<sup>83</sup> P. Rieck,<sup>103</sup> C. J. Riegel,<sup>178</sup> J. Rieger,<sup>57</sup> O. Rifki,<sup>115</sup> M. Rijssenbeek,<sup>150</sup> A. Rimoldi,<sup>123a,123b</sup> M. Rimoldi,<sup>18</sup> L. Rinaldi,<sup>22a</sup> G. Ripellino,<sup>149</sup> B. Ristić,<sup>32</sup> E. Ritsch,<sup>32</sup> I. Riu,<sup>13</sup> F. Rizatdinova,<sup>116</sup> E. Rizvi,<sup>79</sup> C. Rizzi,<sup>13</sup> R. T. Roberts,<sup>87</sup> S. H. Robertson,<sup>90,p</sup> A. Robichaud-Veronneau,<sup>90</sup> D. Robinson,<sup>30</sup> J. E. M. Robinson,<sup>45</sup> A. Robson,<sup>56</sup> E. Rocco,<sup>86</sup> C. Roda,<sup>126a,126b</sup> Y. Rodina,<sup>88,nm</sup> S. Rodriguez Bosca,<sup>170</sup> A. Rodriguez Perez,<sup>13</sup> D. Rodriguez Rodriguez,<sup>170</sup> A. M. Rodríguez Vera,<sup>163b</sup> S. Roe,<sup>32</sup> C. S. Rogan,<sup>59</sup> O. Röhne,<sup>121</sup> J. Roloff,<sup>59</sup> A. Romaniouk,<sup>100</sup> M. Romano,<sup>22a,22b</sup> S. M. Romano Saez,<sup>37</sup> E. Romero Adam,<sup>170</sup> N. Rompotis,<sup>77</sup> M. Ronzani,<sup>51</sup> L. Roos,<sup>83</sup> S. Rosati,<sup>134a</sup> K. Rosbach,<sup>51</sup> P. Rose,<sup>139</sup> N.-A. Rosien,<sup>57</sup> E. Rossi,<sup>106a,106b</sup> L. P. Rossi,<sup>53a</sup> J. H. N. Rosten,<sup>30</sup> R. Rosten,<sup>140</sup> M. Rotaru,<sup>28b</sup> J. Rothberg,<sup>140</sup> D. Rousseau,<sup>119</sup> D. Roy,<sup>147c</sup> A. Rozanov,<sup>88</sup> Y. Rozen,<sup>154</sup> X. Ruan,<sup>147c</sup> F. Rubbo,<sup>145</sup> F. Rühr,<sup>51</sup> A. Ruiz-Martinez,<sup>31</sup> Z. Rurikova,<sup>51</sup> N. A. Rusakovich,<sup>68</sup> H. L. Russell,<sup>90</sup> J. P. Rutherford,<sup>7</sup> N. Ruthmann,<sup>32</sup> E. M. Rüttinger,<sup>45</sup> Y. F. Ryabov,<sup>125</sup> M. Rybar,<sup>169</sup> G. Rybkin,<sup>119</sup> S. Ryu,<sup>6</sup> A. Ryzhov,<sup>132</sup> G. F. Rzehorz,<sup>57</sup> A. F. Saavedra,<sup>152</sup> G. Sabato,<sup>109</sup> S. Sacerdoti,<sup>29</sup> H. F.-W. Sadrozinski,<sup>139</sup> R. Sadykov,<sup>68</sup> F. Safai Tehrani,<sup>134a</sup> P. Saha,<sup>110</sup> M. Sahinsoy,<sup>60a</sup> M. Saimpert,<sup>45</sup> M. Saito,<sup>157</sup> T. Saito,<sup>157</sup> H. Sakamoto,<sup>157</sup> Y. Sakurai,<sup>174</sup> G. Salamanna,<sup>136a,136b</sup> J. E. Salazar Loyola,<sup>34b</sup> D. Salek,<sup>109</sup> P. H. Sales De Bruin,<sup>168</sup> D. Salihagic,<sup>103</sup> A. Salnikov,<sup>145</sup> J. Salt,<sup>170</sup> D. Salvatore,<sup>40a,40b</sup> F. Salvatore,<sup>151</sup> A. Salvucci,<sup>62a,62b,62c</sup> A. Salzburger,<sup>32</sup> D. Sammel,<sup>51</sup> D. Sampsonidis,<sup>156</sup> D. Sampsonidou,<sup>156</sup> J. Sánchez,<sup>170</sup> A. Sanchez Pineda,<sup>167a,167c</sup> H. Sandaker,<sup>121</sup> R. L. Sandbach,<sup>79</sup> C. O. Sander,<sup>45</sup> M. Sandhoff,<sup>178</sup> C. Sandoval,<sup>21</sup> D. P. C. Sankey,<sup>133</sup> M. Sannino,<sup>53a,53b</sup> Y. Sano,<sup>105</sup> A. Sansoni,<sup>50</sup> C. Santoni,<sup>37</sup> H. Santos,<sup>128a</sup> I. Santoyo Castillo,<sup>151</sup> A. Saponov,<sup>68</sup> J. G. Saraiva,<sup>128a,128d</sup> O. Sasaki,<sup>69</sup> K. Sato,<sup>164</sup> E. Sauvan,<sup>5</sup> G. Savage,<sup>80</sup> P. Savard,<sup>161,e</sup> N. Savic,<sup>103</sup> C. Sawyer,<sup>133</sup> L. Sawyer,<sup>82,v</sup> C. Sbarra,<sup>22a</sup> A. Sbrizzi,<sup>22a,22b</sup> T. Scanlon,<sup>81</sup> D. A. Scannicchio,<sup>166</sup> J. Schaarschmidt,<sup>140</sup> P. Schacht,<sup>103</sup> B. M. Schachtner,<sup>102</sup> D. Schaefer,<sup>33</sup> L. Schaefer,<sup>124</sup> J. Schaeffer,<sup>86</sup> S. Schaepe,<sup>32</sup> U. Schäfer,<sup>86</sup> A. C. Schaffer,<sup>119</sup> D. Schaile,<sup>102</sup> R. D. Schamberger,<sup>150</sup> V. A. Schegelsky,<sup>125</sup> D. Scheirich,<sup>131</sup> F. Schenck,<sup>17</sup> M. Schernau,<sup>166</sup> C. Schiavi,<sup>53a,53b</sup> S. Schier,<sup>139</sup> L. K. Schildgen,<sup>23</sup> C. Schillo,<sup>51</sup> E. J. Schioppa,<sup>32</sup> M. Schioppa,<sup>40a,40b</sup> S. Schlenker,<sup>32</sup> K. R. Schmidt-Sommerfeld,<sup>103</sup> K. Schmieden,<sup>32</sup> C. Schmitt,<sup>86</sup> S. Schmitt,<sup>45</sup> S. Schmitz,<sup>86</sup> U. Schnoor,<sup>51</sup> L. Schoeffel,<sup>138</sup> A. Schoening,<sup>60b</sup> B. D. Schoenrock,<sup>93</sup> E. Schopf,<sup>23</sup> M. Schott,<sup>86</sup> J. F. P. Schouwenberg,<sup>108</sup> J. Schovancova,<sup>32</sup> S. Schramm,<sup>52</sup> N. Schuh,<sup>86</sup> A. Schulte,<sup>86</sup> M. J. Schultens,<sup>23</sup> H.-C. Schultz-Coulon,<sup>60a</sup> M. Schumacher,<sup>51</sup> B. A. Schumm,<sup>139</sup> Ph. Schune,<sup>138</sup> A. Schwartzman,<sup>145</sup> T. A. Schwarz,<sup>92</sup> H. Schweiger,<sup>87</sup> Ph. Schwemling,<sup>138</sup> R. Schwienhorst,<sup>93</sup> J. Schwindling,<sup>138</sup> A. Sciandra,<sup>23</sup> G. Sciolla,<sup>25</sup> M. Scornajenghi,<sup>40a,40b</sup> F. Scuri,<sup>126a</sup> F. Scutti,<sup>91</sup> J. Searcy,<sup>92</sup> P. Seema,<sup>23</sup> S. C. Seidel,<sup>107</sup> A. Seiden,<sup>139</sup> J. M. Seixas,<sup>26a</sup> G. Sekhniaidze,<sup>106a</sup> K. Sekhon,<sup>92</sup> S. J. Sekula,<sup>43</sup> N. Semprini-Cesari,<sup>22a,22b</sup> S. Senkin,<sup>37</sup> C. Serfon,<sup>121</sup> L. Serin,<sup>119</sup> L. Serkin,<sup>167a,167b</sup> M. Sessa,<sup>136a,136b</sup> R. Seuster,<sup>172</sup> H. Severini,<sup>115</sup> T. Šfiligoj,<sup>78</sup> F. Sforza,<sup>165</sup> A. Sfyrila,<sup>52</sup> E. Shabalina,<sup>57</sup> N. W. Shaikh,<sup>148a,148b</sup> L. Y. Shan,<sup>35a</sup> R. Shang,<sup>169</sup> J. T. Shank,<sup>24</sup> M. Shapiro,<sup>16</sup> P. B. Shatalov,<sup>99</sup> K. Shaw,<sup>167a,167b</sup> S. M. Shaw,<sup>87</sup> A. Shcherbakova,<sup>148a,148b</sup> C. Y. Shehu,<sup>151</sup> Y. Shen,<sup>115</sup> N. Sherafati,<sup>31</sup> A. D. Sherman,<sup>24</sup> P. Sherwood,<sup>81</sup> L. Shi,<sup>153,oo</sup> S. Shimizu,<sup>70</sup> C. O. Shimmin,<sup>179</sup> M. Shimojima,<sup>104</sup> I. P. J. Shipsey,<sup>122</sup> S. Shirabe,<sup>73</sup> M. Shiyakova,<sup>68,pp</sup> J. Shlomi,<sup>175</sup> A. Shmeleva,<sup>98</sup> D. Shoaleh Saadi,<sup>97</sup> M. J. Shochet,<sup>33</sup> S. Shojaii,<sup>94a,94b</sup> D. R. Shope,<sup>115</sup> S. Shrestha,<sup>113</sup> E. Shulga,<sup>100</sup> M. A. Shupe,<sup>7</sup> P. Sicho,<sup>129</sup> A. M. Sickles,<sup>169</sup> P. E. Sidebo,<sup>149</sup> E. Sideras Haddad,<sup>147c</sup> O. Sidiropoulou,<sup>177</sup> A. Sidoti,<sup>22a,22b</sup> F. Siegert,<sup>47</sup> Dj. Sijacki,<sup>14</sup> J. Silva,<sup>128a,128d</sup> M. Silva Jr.,<sup>176</sup> S. B. Silverstein,<sup>148a</sup> V. Simak,<sup>130</sup> L. Simic,<sup>68</sup> S. Simion,<sup>119</sup> E. Simioni,<sup>86</sup> B. Simmons,<sup>81</sup> M. Simon,<sup>86</sup> P. Sinervo,<sup>161</sup> N. B. Sinev,<sup>118</sup> M. Sioli,<sup>22a,22b</sup> G. Siragusa,<sup>177</sup> I. Siral,<sup>92</sup> S. Yu. Sivoklokov,<sup>101</sup> J. Sjölin,<sup>148a,148b</sup> M. B. Skinner,<sup>75</sup> P. Skubic,<sup>115</sup> M. Slater,<sup>19</sup> T. Slavicek,<sup>130</sup> M. Slawinska,<sup>42</sup> K. Sliwa,<sup>165</sup> R. Slovak,<sup>131</sup> V. Smakhtin,<sup>175</sup> B. H. Smart,<sup>5</sup> J. Smiesko,<sup>146a</sup> N. Smirnov,<sup>100</sup> S. Yu. Smirnov,<sup>100</sup> Y. Smirnov,<sup>100</sup> L. N. Smirnova,<sup>101,qq</sup> O. Smirnova,<sup>84</sup> J. W. Smith,<sup>57</sup> M. N. K. Smith,<sup>38</sup> R. W. Smith,<sup>38</sup> M. Smizanska,<sup>75</sup> K. Smolek,<sup>130</sup> A. A. Snesarev,<sup>98</sup> I. M. Snyder,<sup>118</sup> S. Snyder,<sup>27</sup> R. Sobie,<sup>172,p</sup> F. Socher,<sup>47</sup> A. M. Sofa,<sup>166</sup> A. Soffer,<sup>155</sup> A. Sogaard,<sup>49</sup> D. A. Soh,<sup>153</sup> G. Sokhrannyi,<sup>78</sup> C. A. Solans Sanchez,<sup>32</sup> M. Solar,<sup>130</sup> E. Yu. Soldatov,<sup>100</sup> U. Soldevila,<sup>170</sup> A. A. Solodkov,<sup>132</sup> A. Soloshenko,<sup>68</sup> O. V. Solovyanov,<sup>132</sup> V. Solov'yev,<sup>125</sup> P. Sommer,<sup>141</sup> H. Son,<sup>165</sup> W. Song,<sup>133</sup> A. Sopczak,<sup>130</sup> D. Sosa,<sup>60b</sup> C. L. Sotiropoulou,<sup>126a,126b</sup> S. Sottocornola,<sup>123a,123b</sup> R. Soualah,<sup>167a,167c</sup> A. M. Soukharev,<sup>111,d</sup> D. South,<sup>45</sup> B. C. Sowden,<sup>80</sup> S. Spagnolo,<sup>76a,76b</sup> M. Spalla,<sup>126a,126b</sup> M. Spangenberg,<sup>173</sup> F. Spanò,<sup>80</sup> D. Sperlich,<sup>17</sup> F. Spettel,<sup>103</sup> T. M. Spieker,<sup>60a</sup> R. Spighi,<sup>22a</sup> G. Spigo,<sup>32</sup> L. A. Spiller,<sup>91</sup> M. Spousta,<sup>131</sup> R. D. St. Denis,<sup>56,a</sup> A. Stabile,<sup>94a,94b</sup> R. Stamen,<sup>60a</sup> S. Stamm,<sup>17</sup> E. Stanecka,<sup>42</sup> R. W. Stanek,<sup>6</sup> C. Stanescu,<sup>136a</sup> M. M. Stanitzki,<sup>45</sup> B. S. Stapf,<sup>109</sup> S. Stapnes,<sup>121</sup>

E. A. Starchenko,<sup>132</sup> G. H. Stark,<sup>33</sup> J. Stark,<sup>58</sup> S. H. Stark,<sup>39</sup> P. Staroba,<sup>129</sup> P. Starovoitov,<sup>60a</sup> S. Starz,<sup>32</sup> R. Staszewski,<sup>42</sup> M. Stegler,<sup>45</sup> P. Steinberg,<sup>27</sup> B. Stelzer,<sup>144</sup> H. J. Stelzer,<sup>32</sup> O. Stelzer-Chilton,<sup>163a</sup> H. Stenzel,<sup>55</sup> T. J. Stevenson,<sup>79</sup> G. A. Stewart,<sup>56</sup> M. C. Stockton,<sup>118</sup> M. Stoebe,<sup>90</sup> G. Stoicea,<sup>28b</sup> P. Stolte,<sup>57</sup> S. Stonjek,<sup>103</sup> A. R. Stradling,<sup>8</sup> A. Straessner,<sup>47</sup> M. E. Stramaglia,<sup>18</sup> J. Strandberg,<sup>149</sup> S. Strandberg,<sup>148a,148b</sup> M. Strauss,<sup>115</sup> P. Strizenec,<sup>146b</sup> R. Strohmer,<sup>177</sup> D. M. Strom,<sup>118</sup> R. Stroynowski,<sup>43</sup> A. Strubig,<sup>49</sup> S. A. Stucci,<sup>27</sup> B. Stugu,<sup>15</sup> N. A. Styles,<sup>45</sup> D. Su,<sup>145</sup> J. Su,<sup>127</sup> S. Suchek,<sup>60a</sup> Y. Sugaya,<sup>120</sup> M. Suk,<sup>130</sup> V. V. Sulin,<sup>98</sup> DMS Sultan,<sup>52</sup> S. Sultansoy,<sup>4c</sup> T. Sumida,<sup>71</sup> S. Sun,<sup>59</sup> X. Sun,<sup>3</sup> K. Suruliz,<sup>151</sup> C. J. E. Suster,<sup>152</sup> M. R. Sutton,<sup>151</sup> S. Suzuki,<sup>69</sup> M. Svatos,<sup>129</sup> M. Swiatkowski,<sup>33</sup> S. P. Swift,<sup>2</sup> A. Sydorenko,<sup>86</sup> I. Sykora,<sup>146a</sup> T. Sykora,<sup>131</sup> D. Ta,<sup>51</sup> K. Tackmann,<sup>45</sup> J. Taenzer,<sup>155</sup> A. Taffard,<sup>166</sup> R. Tafirout,<sup>163a</sup> E. Tahirovic,<sup>79</sup> N. Taiblum,<sup>155</sup> H. Takai,<sup>27</sup> R. Takashima,<sup>72</sup> E. H. Takasugi,<sup>103</sup> K. Takeda,<sup>70</sup> T. Takeshita,<sup>142</sup> Y. Takubo,<sup>69</sup> M. Talby,<sup>88</sup> A. A. Talyshev,<sup>111,d</sup> J. Tanaka,<sup>157</sup> M. Tanaka,<sup>159</sup> R. Tanaka,<sup>119</sup> R. Tanioka,<sup>70</sup> B. B. Tannenwald,<sup>113</sup> S. Tapia Araya,<sup>34b</sup> S. Tapprogge,<sup>86</sup> A. T. Tarek Abouelfadl Mohamed,<sup>83</sup> S. Tarem,<sup>154</sup> G. F. Tartarelli,<sup>94a</sup> P. Tas,<sup>131</sup> M. Tasevsky,<sup>129</sup> T. Tashiro,<sup>71</sup> E. Tassi,<sup>40a,40b</sup> A. Tavares Delgado,<sup>128a,128b</sup> Y. Tayalati,<sup>137d</sup> A. C. Taylor,<sup>107</sup> A. J. Taylor,<sup>49</sup> G. N. Taylor,<sup>91</sup> P. T. E. Taylor,<sup>91</sup> W. Taylor,<sup>163b</sup> P. Teixeira-Dias,<sup>80</sup> D. Temple,<sup>144</sup> H. Ten Kate,<sup>32</sup> P. K. Teng,<sup>153</sup> J. J. Teoh,<sup>120</sup> F. Tepel,<sup>178</sup> S. Terada,<sup>69</sup> K. Terashi,<sup>157</sup> J. Terron,<sup>85</sup> S. Terzo,<sup>13</sup> M. Testa,<sup>50</sup> R. J. Teuscher,<sup>161,p</sup> S. J. Thais,<sup>179</sup> T. Thevenaux-Pelzer,<sup>88</sup> F. Thiele,<sup>39</sup> J. P. Thomas,<sup>19</sup> J. Thomas-Wilsker,<sup>80</sup> P. D. Thompson,<sup>19</sup> A. S. Thompson,<sup>56</sup> L. A. Thomsen,<sup>179</sup> E. Thomson,<sup>124</sup> Y. Tian,<sup>38</sup> R. E. Ticse Torres,<sup>57</sup> V. O. Tikhomirov,<sup>98,r</sup> Yu. A. Tikhonov,<sup>111,d</sup> S. Timoshenko,<sup>100</sup> P. Tipton,<sup>179</sup> S. Tisserant,<sup>88</sup> K. Todome,<sup>159</sup> S. Todorova-Nova,<sup>5</sup> S. Todt,<sup>47</sup> J. Tojo,<sup>73</sup> S. Tokar,<sup>146a</sup> K. Tokushuku,<sup>69</sup> E. Tolley,<sup>113</sup> L. Tomlinson,<sup>87</sup> M. Tomoto,<sup>105</sup> L. Tompkins,<sup>145,ss</sup> K. Toms,<sup>107</sup> B. Tong,<sup>59</sup> P. Tornambe,<sup>51</sup> E. Torrence,<sup>118</sup> H. Torres,<sup>47</sup> E. Torro Pastor,<sup>140</sup> J. Toth,<sup>88,t</sup> F. Touchard,<sup>88</sup> D. R. Tovey,<sup>141</sup> C. J. Treado,<sup>112</sup> T. Trefzger,<sup>177</sup> F. Tresoldi,<sup>151</sup> A. Tricoli,<sup>27</sup> I. M. Trigger,<sup>163a</sup> S. Trincas-Duvoid,<sup>83</sup> M. F. Tripiana,<sup>13</sup> W. Trischuk,<sup>161</sup> B. Trocme,<sup>58</sup> A. Trofymov,<sup>45</sup> C. Troncon,<sup>94a</sup> M. Trovatelli,<sup>172</sup> L. Truong,<sup>147b</sup> M. Trzebinski,<sup>42</sup> A. Trzupek,<sup>42</sup> K. W. Tsang,<sup>62a</sup> J. C-L. Tseng,<sup>122</sup> P. V. Tsiarashka,<sup>95</sup> N. Tsirintanis,<sup>9</sup> S. Tsiskaridze,<sup>13</sup> V. Tsiskaridze,<sup>51</sup> E. G. Tskhadadze,<sup>54a</sup> I. I. Tsukerman,<sup>99</sup> V. Tsulaia,<sup>16</sup> S. Tsuno,<sup>69</sup> D. Tsybychev,<sup>150</sup> Y. Tu,<sup>62b</sup> A. Tudorache,<sup>28b</sup> V. Tudorache,<sup>28b</sup> T. T. Tulbure,<sup>28a</sup> A. N. Tuna,<sup>59</sup> S. Turchikhin,<sup>68</sup> D. Turgeman,<sup>175</sup> I. Turk Cakir,<sup>4b,uu</sup> R. Turra,<sup>94a</sup> P. M. Tuts,<sup>38</sup> G. Uccelli,<sup>22a,22b</sup> I. Ueda,<sup>69</sup> M. Ughetto,<sup>148a,148b</sup> F. Ukegawa,<sup>164</sup> G. Unal,<sup>32</sup> A. Undrus,<sup>27</sup> G. Unel,<sup>166</sup> F. C. Ungaro,<sup>91</sup> Y. Unno,<sup>69</sup> K. Uno,<sup>157</sup> J. Urban,<sup>146b</sup> P. Urquijo,<sup>91</sup> P. Urrejola,<sup>86</sup> G. Usai,<sup>8</sup> J. Usui,<sup>69</sup> L. Vacavant,<sup>88</sup> V. Vacek,<sup>130</sup> B. Vachon,<sup>90</sup> K. O. H. Vadla,<sup>121</sup> A. Vaidya,<sup>81</sup> C. Valderanis,<sup>102</sup> E. Valdes Santurio,<sup>148a,148b</sup> M. Valente,<sup>52</sup> S. Valentinetti,<sup>22a,22b</sup> A. Valero,<sup>170</sup> L. Valery,<sup>13</sup> A. Vallier,<sup>5</sup> J. A. Valls Ferrer,<sup>170</sup> W. Van Den Wollenberg,<sup>109</sup> H. van der Graaf,<sup>109</sup> P. van Gemmeren,<sup>6</sup> J. Van Nieuwkoop,<sup>144</sup> I. van Vulpen,<sup>109</sup> M. C. van Woerden,<sup>109</sup> M. Vanadia,<sup>135a,135b</sup> W. Vandelli,<sup>32</sup> A. Vaniachine,<sup>160</sup> P. Vankov,<sup>109</sup> G. Vardanyan,<sup>180</sup> R. Vari,<sup>134a</sup> E. W. Varnes,<sup>7</sup> C. Varni,<sup>53a,53b</sup> T. Varol,<sup>43</sup> D. Varouchas,<sup>119</sup> A. Vartapetian,<sup>8</sup> K. E. Varvell,<sup>152</sup> J. G. Vasquez,<sup>179</sup> G. A. Vasquez,<sup>34b</sup> F. Vazeille,<sup>37</sup> D. Vazquez Furelos,<sup>13</sup> T. Vazquez Schroeder,<sup>90</sup> J. Veatch,<sup>57</sup> V. Veeraraghavan,<sup>7</sup> L. M. Veloce,<sup>161</sup> F. Veloso,<sup>128a,128c</sup> S. Veneziano,<sup>134a</sup> A. Ventura,<sup>76a,76b</sup> M. Venturi,<sup>172</sup> N. Venturi,<sup>32</sup> V. Vercesi,<sup>123a</sup> M. Verducci,<sup>136a,136b</sup> W. Verkerke,<sup>109</sup> A. T. Vermeulen,<sup>109</sup> J. C. Vermeulen,<sup>109</sup> M. C. Vetterli,<sup>144,e</sup> N. Viaux Maira,<sup>34b</sup> O. Viazlo,<sup>84</sup> I. Vichou,<sup>169,a</sup> T. Vickey,<sup>141</sup> O. E. Vickey Boeriu,<sup>141</sup> G. H. A. Viehhauser,<sup>122</sup> S. Viel,<sup>16</sup> L. Vigani,<sup>122</sup> M. Villa,<sup>22a,22b</sup> M. Villaplana Perez,<sup>94a,94b</sup> E. Vilucchi,<sup>50</sup> M. G. Vincet,<sup>31</sup> V. B. Vinogradov,<sup>68</sup> A. Vishwakarma,<sup>45</sup> C. Vittori,<sup>22a,22b</sup> I. Vivarelli,<sup>151</sup> S. Vlachos,<sup>10</sup> M. Vogel,<sup>178</sup> P. Vokac,<sup>130</sup> G. Volpi,<sup>13</sup> S. E. von Buddenbrock,<sup>147c</sup> H. von der Schmitt,<sup>103</sup> E. von Toerne,<sup>23</sup> V. Vorobel,<sup>131</sup> K. Vorobev,<sup>100</sup> M. Vos,<sup>170</sup> R. Voss,<sup>32</sup> J. H. Vossebeld,<sup>77</sup> N. Vranjes,<sup>14</sup> M. Vranjes Milosavljevic,<sup>14</sup> V. Vrba,<sup>130</sup> M. Vreeswijk,<sup>109</sup> R. Vuillermet,<sup>32</sup> I. Vukotic,<sup>33</sup> P. Wagner,<sup>23</sup> W. Wagner,<sup>178</sup> J. Wagner-Kuhr,<sup>102</sup> H. Wahlberg,<sup>74</sup> S. WAhrmund,<sup>47</sup> K. Wakamiya,<sup>70</sup> J. Walder,<sup>75</sup> R. Walker,<sup>102</sup> W. Walkowiak,<sup>143</sup> V. Wallangen,<sup>148a,148b</sup> A. M. Wang,<sup>59</sup> C. Wang,<sup>36a,vv</sup> F. Wang,<sup>176</sup> H. Wang,<sup>16</sup> H. Wang,<sup>3</sup> J. Wang,<sup>60b</sup> J. Wang,<sup>152</sup> Q. Wang,<sup>115</sup> R.-J. Wang,<sup>83</sup> R. Wang,<sup>6</sup> S. M. Wang,<sup>153</sup> T. Wang,<sup>38</sup> W. Wang,<sup>153,ww</sup> W. Wang,<sup>36c,xx</sup> Z. Wang,<sup>36b</sup> C. Wanotayaraj,<sup>45</sup> A. Warburton,<sup>90</sup> C. P. Ward,<sup>30</sup> D. R. Wardrope,<sup>81</sup> A. Washbrook,<sup>49</sup> P. M. Watkins,<sup>19</sup> A. T. Watson,<sup>19</sup> M. F. Watson,<sup>19</sup> G. Watts,<sup>140</sup> S. Watts,<sup>87</sup> B. M. Waugh,<sup>81</sup> A. F. Webb,<sup>11</sup> S. Webb,<sup>86</sup> M. S. Weber,<sup>18</sup> S. M. Weber,<sup>60a</sup> S. A. Weber,<sup>31</sup> J. S. Webster,<sup>6</sup> A. R. Weidberg,<sup>122</sup> B. Weinert,<sup>64</sup> J. Weingarten,<sup>57</sup> M. Weirich,<sup>86</sup> C. Weiser,<sup>51</sup> P. S. Wells,<sup>32</sup> T. Wenaus,<sup>27</sup> T. Wengler,<sup>32</sup> S. Wenig,<sup>32</sup> N. Wermes,<sup>23</sup> M. D. Werner,<sup>67</sup> P. Werner,<sup>32</sup> M. Wessels,<sup>60a</sup> T. D. Weston,<sup>18</sup> K. Whalen,<sup>118</sup> N. L. Whallon,<sup>140</sup> A. M. Wharton,<sup>75</sup> A. S. White,<sup>92</sup> A. White,<sup>8</sup> M. J. White,<sup>1</sup> R. White,<sup>34b</sup> D. Whiteson,<sup>166</sup> B. W. Whitmore,<sup>75</sup> F. J. Wickens,<sup>133</sup> W. Wiedenmann,<sup>176</sup> M. WIELERS,<sup>133</sup> C. WIGLESWORTH,<sup>39</sup> L. A. M. Wiik-Fuchs,<sup>51</sup> A. Wildauer,<sup>103</sup> F. Wilk,<sup>87</sup> H. G. Wilkens,<sup>32</sup> H. H. Williams,<sup>124</sup> S. Williams,<sup>30</sup> C. Willis,<sup>93</sup> S. Willocq,<sup>89</sup> J. A. Wilson,<sup>19</sup> I. Wingerter-Seez,<sup>5</sup> E. Winkels,<sup>151</sup> F. Winklmeier,<sup>118</sup> O. J. Winston,<sup>151</sup> B. T. Winter,<sup>23</sup> M. Wittgen,<sup>145</sup> M. Wobisch,<sup>82,v</sup> A. Wolf,<sup>86</sup> T. M. H. Wolf,<sup>109</sup> R. Wolff,<sup>88</sup> M. W. Wolter,<sup>42</sup> H. Wolters,<sup>128a,128c</sup> V. W. S. Wong,<sup>171</sup> N. L. Woods,<sup>139</sup>



S. D. Worm,<sup>19</sup> B. K. Wosiek,<sup>42</sup> J. Wotschack,<sup>32</sup> K. W. Wozniak,<sup>42</sup> M. Wu,<sup>33</sup> S. L. Wu,<sup>176</sup> X. Wu,<sup>52</sup> Y. Wu,<sup>92</sup> T. R. Wyatt,<sup>87</sup>  
 B. M. Wynne,<sup>49</sup> S. Xella,<sup>39</sup> Z. Xi,<sup>92</sup> L. Xia,<sup>35c</sup> D. Xu,<sup>35a</sup> L. Xu,<sup>27</sup> T. Xu,<sup>138</sup> W. Xu,<sup>92</sup> B. Yabsley,<sup>152</sup> S. Yacoob,<sup>147a</sup>  
 K. Yajima,<sup>120</sup> D. P. Yallup,<sup>81</sup> D. Yamaguchi,<sup>159</sup> Y. Yamaguchi,<sup>159</sup> A. Yamamoto,<sup>69</sup> S. Yamamoto,<sup>157</sup> T. Yamanaka,<sup>157</sup>  
 F. Yamane,<sup>70</sup> M. Yamatani,<sup>157</sup> T. Yamazaki,<sup>157</sup> Y. Yamazaki,<sup>70</sup> Z. Yan,<sup>24</sup> H. Yang,<sup>36b</sup> H. Yang,<sup>16</sup> S. Yang,<sup>66</sup> Y. Yang,<sup>153</sup>  
 Z. Yang,<sup>15</sup> W.-M. Yao,<sup>16</sup> Y. C. Yap,<sup>45</sup> Y. Yasu,<sup>69</sup> E. Yatsenko,<sup>5</sup> K. H. Yau Wong,<sup>23</sup> J. Ye,<sup>43</sup> S. Ye,<sup>27</sup> I. Yeletsikh,<sup>68</sup>  
 E. Yigitbasi,<sup>24</sup> E. Yildirim,<sup>86</sup> K. Yorita,<sup>174</sup> K. Yoshihara,<sup>124</sup> C. Young,<sup>145</sup> C. J. S. Young,<sup>32</sup> J. Yu,<sup>8</sup> J. Yu,<sup>67</sup> S. P. Y. Yuen,<sup>23</sup>  
 I. Yusuff,<sup>30,yy</sup> B. Zabinski,<sup>42</sup> G. Zacharis,<sup>10</sup> R. Zaidan,<sup>13</sup> A. M. Zaitsev,<sup>132,II</sup> N. Zakharchuk,<sup>45</sup> J. Zalieckas,<sup>15</sup> A. Zaman,<sup>150</sup>  
 S. Zambito,<sup>59</sup> D. Zanzi,<sup>32</sup> C. Zeitnitz,<sup>178</sup> G. Zemaityte,<sup>122</sup> J. C. Zeng,<sup>169</sup> Q. Zeng,<sup>145</sup> O. Zenin,<sup>132</sup> T. Ženiš,<sup>146a</sup> D. Zerwas,<sup>119</sup>  
 D. Zhang,<sup>36a</sup> D. Zhang,<sup>92</sup> F. Zhang,<sup>176</sup> G. Zhang,<sup>36c,xx</sup> H. Zhang,<sup>119</sup> J. Zhang,<sup>6</sup> L. Zhang,<sup>51</sup> L. Zhang,<sup>36c</sup> M. Zhang,<sup>169</sup>  
 P. Zhang,<sup>35b</sup> R. Zhang,<sup>23</sup> R. Zhang,<sup>36c,vv</sup> X. Zhang,<sup>36a</sup> Y. Zhang,<sup>35a,35d</sup> Z. Zhang,<sup>119</sup> X. Zhao,<sup>43</sup> Y. Zhao,<sup>36a,y</sup> Z. Zhao,<sup>36c</sup>  
 A. Zhemchugov,<sup>68</sup> B. Zhou,<sup>92</sup> C. Zhou,<sup>176</sup> L. Zhou,<sup>43</sup> M. Zhou,<sup>35a,35d</sup> M. Zhou,<sup>150</sup> N. Zhou,<sup>36b</sup> Y. Zhou,<sup>7</sup> C. G. Zhu,<sup>36a</sup>  
 H. Zhu,<sup>35a</sup> J. Zhu,<sup>92</sup> Y. Zhu,<sup>36c</sup> X. Zhuang,<sup>35a</sup> K. Zhukov,<sup>98</sup> A. Zibell,<sup>177</sup> D. Zieminska,<sup>64</sup> N. I. Zimine,<sup>68</sup> S. Zimmermann,<sup>51</sup>  
 Z. Zinonos,<sup>103</sup> M. Zinser,<sup>86</sup> M. Ziolkowski,<sup>143</sup> L. Živković,<sup>14</sup> G. Zobernig,<sup>176</sup> A. Zoccoli,<sup>22a,22b</sup> R. Zou,<sup>33</sup>  
 M. zur Nedden,<sup>17</sup> and L. Zwalinski<sup>32</sup>

(ATLAS Collaboration)

<sup>1</sup>*Department of Physics, University of Adelaide, Adelaide, Australia*

<sup>2</sup>*Physics Department, SUNY Albany, Albany, New York, USA*

<sup>3</sup>*Department of Physics, University of Alberta, Edmonton, Alberta, Canada*

<sup>4a</sup>*Department of Physics, Ankara University, Ankara, Turkey*

<sup>4b</sup>*Istanbul Aydin University, Istanbul, Turkey*

<sup>4c</sup>*Division of Physics, TOBB University of Economics and Technology, Ankara, Turkey*

<sup>5</sup>*LAPP, CNRS/IN2P3 and Université Savoie Mont Blanc, Annecy-le-Vieux, France*

<sup>6</sup>*High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois, USA*

<sup>7</sup>*Department of Physics, University of Arizona, Tucson, Arizona, USA*

<sup>8</sup>*Department of Physics, The University of Texas at Arlington, Arlington, Texas, USA*

<sup>9</sup>*Physics Department, National and Kapodistrian University of Athens, Athens, Greece*

<sup>10</sup>*Physics Department, National Technical University of Athens, Zografou, Greece*

<sup>11</sup>*Department of Physics, The University of Texas at Austin, Austin, Texas, USA*

<sup>12</sup>*Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan*

<sup>13</sup>*Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Barcelona, Spain*

<sup>14</sup>*Institute of Physics, University of Belgrade, Belgrade, Serbia*

<sup>15</sup>*Department for Physics and Technology, University of Bergen, Bergen, Norway*

<sup>16</sup>*Physics Division, Lawrence Berkeley National Laboratory and University of California, Berkeley, California, USA*

<sup>17</sup>*Department of Physics, Humboldt University, Berlin, Germany*

<sup>18</sup>*Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern, Switzerland*

<sup>19</sup>*School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*

<sup>20a</sup>*Department of Physics, Bogazici University, Istanbul, Turkey*

<sup>20b</sup>*Department of Physics Engineering, Gaziantep University, Gaziantep, Turkey*

<sup>20c</sup>*Istanbul Bilgi University, Faculty of Engineering and Natural Sciences, Istanbul, Turkey*

<sup>20d</sup>*Bahcesehir University, Faculty of Engineering and Natural Sciences, Istanbul, Turkey*

<sup>21</sup>*Centro de Investigaciones, Universidad Antonio Narino, Bogota, Colombia*

<sup>22a</sup>*INFN Sezione di Bologna, Italy*

<sup>22b</sup>*Dipartimento di Fisica e Astronomia, Università di Bologna, Bologna, Italy*

<sup>23</sup>*Physikalisches Institut, University of Bonn, Bonn, Germany*

<sup>24</sup>*Department of Physics, Boston University, Boston, Massachusetts, USA*

<sup>25</sup>*Department of Physics, Brandeis University, Waltham, Massachusetts, USA*

<sup>26a</sup>*Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro, Brazil*

<sup>26b</sup>*Electrical Circuits Department, Federal University of Juiz de Fora (UFJF), Juiz de Fora, Brazil*

<sup>26c</sup>*Federal University of Sao Joao del Rei (UFSJ), Sao Joao del Rei, Brazil*

<sup>26d</sup>*Instituto de Física, Universidade de Sao Paulo, Sao Paulo, Brazil*

<sup>27</sup>*Physics Department, Brookhaven National Laboratory, Upton, New York, USA*



- <sup>28a</sup>Transilvania University of Brasov, Brasov, Romania
- <sup>28b</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- <sup>28c</sup>Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi, Romania
- <sup>28d</sup>National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj Napoca, Romania
- <sup>28e</sup>West University in Timisoara, Timisoara, Romania
- <sup>29</sup>Departamento de Física, Universidad de Buenos Aires, Buenos Aires, Argentina
- <sup>30</sup>Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- <sup>31</sup>Department of Physics, Carleton University, Ottawa, Ontario, Canada
- <sup>32</sup>CERN, Geneva, Switzerland
- <sup>33</sup>Enrico Fermi Institute, University of Chicago, Chicago, Illinois, USA
- <sup>34a</sup>Departamento de Física, Pontificia Universidad Católica de Chile, Santiago, Chile
- <sup>34b</sup>Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso, Chile
- <sup>35a</sup>Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China
- <sup>35b</sup>Department of Physics, Nanjing University, Jiangsu, China
- <sup>35c</sup>Physics Department, Tsinghua University, Beijing 100084, China
- <sup>35d</sup>University of Chinese Academy of Science (UCAS), Beijing, China
- <sup>36a</sup>School of Physics, Shandong University, Shandong, China
- <sup>36b</sup>School of Physics and Astronomy, Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology, Tsung-Dao Lee Institute, Shanghai Jiao Tong University, China
- <sup>36c</sup>Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Anhui, China
- <sup>37</sup>Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
- <sup>38</sup>Nevis Laboratory, Columbia University, Irvington, New York, USA
- <sup>39</sup>Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- <sup>40a</sup>INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati, Italy
- <sup>40b</sup>Dipartimento di Fisica, Università della Calabria, Rende, Italy
- <sup>41a</sup>AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow, Poland
- <sup>41b</sup>Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow, Poland
- <sup>42</sup>Institute of Nuclear Physics Polish Academy of Sciences, Krakow, Poland
- <sup>43</sup>Physics Department, Southern Methodist University, Dallas, Texas, USA
- <sup>44</sup>Physics Department, University of Texas at Dallas, Richardson, Texas, USA
- <sup>45</sup>DESY, Hamburg and Zeuthen, Germany
- <sup>46</sup>Lehrstuhl für Experimentelle Physik IV, Technische Universität Dortmund, Dortmund, Germany
- <sup>47</sup>Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden, Germany
- <sup>48</sup>Department of Physics, Duke University, Durham, North Carolina, USA
- <sup>49</sup>SUPA—School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- <sup>50</sup>INFN e Laboratori Nazionali di Frascati, Frascati, Italy
- <sup>51</sup>Fakultät für Mathematik und Physik, Albert-Ludwigs-Universität, Freiburg, Germany
- <sup>52</sup>Département de Physique Nucleaire et Corpusculaire, Université de Genève, Geneva, Switzerland
- <sup>53a</sup>INFN Sezione di Genova, Italy
- <sup>53b</sup>Dipartimento di Fisica, Università di Genova, Genova, Italy
- <sup>54a</sup>E. Andronikashvili Institute of Physics, Iv. Javakishvili Tbilisi State University, Tbilisi, Georgia
- <sup>54b</sup>High Energy Physics Institute, Tbilisi State University, Tbilisi, Georgia
- <sup>55</sup>II Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen, Germany
- <sup>56</sup>SUPA—School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- <sup>57</sup>II Physikalisches Institut, Georg-August-Universität, Göttingen, Germany
- <sup>58</sup>Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS/IN2P3, Grenoble, France
- <sup>59</sup>Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, Massachusetts, USA
- <sup>60a</sup>Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- <sup>60b</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- <sup>61</sup>Faculty of Applied Information Science, Hiroshima Institute of Technology, Hiroshima, Japan
- <sup>62a</sup>Department of Physics, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China
- <sup>62b</sup>Department of Physics, The University of Hong Kong, Hong Kong, China
- <sup>62c</sup>Department of Physics and Institute for Advanced Study, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China

- <sup>63</sup>*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan*  
<sup>64</sup>*Department of Physics, Indiana University, Bloomington, Indiana, USA*  
<sup>65</sup>*Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität, Innsbruck, Austria*  
<sup>66</sup>*University of Iowa, Iowa City, Iowa, USA*  
<sup>67</sup>*Department of Physics and Astronomy, Iowa State University, Ames, Iowa, USA*  
<sup>68</sup>*Joint Institute for Nuclear Research, JINR Dubna, Dubna, Russia*  
<sup>69</sup>*KEK, High Energy Accelerator Research Organization, Tsukuba, Japan*  
<sup>70</sup>*Graduate School of Science, Kobe University, Kobe, Japan*  
<sup>71</sup>*Faculty of Science, Kyoto University, Kyoto, Japan*  
<sup>72</sup>*Kyoto University of Education, Kyoto, Japan*  
<sup>73</sup>*Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka, Japan*  
<sup>74</sup>*Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata, Argentina*  
<sup>75</sup>*Physics Department, Lancaster University, Lancaster, United Kingdom*  
<sup>76a</sup>*INFN Sezione di Lecce, Italy*  
<sup>76b</sup>*Dipartimento di Matematica e Fisica, Università del Salento, Lecce, Italy*  
<sup>77</sup>*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*  
<sup>78</sup>*Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana, Slovenia*  
<sup>79</sup>*School of Physics and Astronomy, Queen Mary University of London, London, United Kingdom*  
<sup>80</sup>*Department of Physics, Royal Holloway University of London, Surrey, United Kingdom*  
<sup>81</sup>*Department of Physics and Astronomy, University College London, London, United Kingdom*  
<sup>82</sup>*Louisiana Tech University, Ruston, Louisiana, USA*  
<sup>83</sup>*Laboratoire de Physique Nucléaire et de Hautes Energies, UPMC and Université Paris-Diderot and CNRS/IN2P3, Paris, France*  
<sup>84</sup>*Fysiska institutionen, Lunds universitet, Lund, Sweden*  
<sup>85</sup>*Departamento de Física Teórica C-15, Universidad Autónoma de Madrid, Madrid, Spain*  
<sup>86</sup>*Institut für Physik, Universität Mainz, Mainz, Germany*  
<sup>87</sup>*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*  
<sup>88</sup>*CPPM, Aix-Marseille Université and CNRS/IN2P3, Marseille, France*  
<sup>89</sup>*Department of Physics, University of Massachusetts, Amherst, Massachusetts, USA*  
<sup>90</sup>*Department of Physics, McGill University, Montreal, Quebec, Canada*  
<sup>91</sup>*School of Physics, University of Melbourne, Victoria, Australia*  
<sup>92</sup>*Department of Physics, The University of Michigan, Ann Arbor, Michigan, USA*  
<sup>93</sup>*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan, USA*  
<sup>94a</sup>*INFN Sezione di Milano, Italy*  
<sup>94b</sup>*Dipartimento di Fisica, Università di Milano, Milano, Italy*  
<sup>95</sup>*B.I. Stepanov Institute of Physics, National Academy of Sciences of Belarus, Minsk, Republic of Belarus*  
<sup>96</sup>*Research Institute for Nuclear Problems of Byelorussian State University, Minsk, Republic of Belarus*  
<sup>97</sup>*Group of Particle Physics, University of Montreal, Montreal, Quebec, Canada*  
<sup>98</sup>*P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow, Russia*  
<sup>99</sup>*Institute for Theoretical and Experimental Physics (ITEP), Moscow, Russia*  
<sup>100</sup>*National Research Nuclear University MEPhI, Moscow, Russia*  
<sup>101</sup>*D.V. Skobeltsyn Institute of Nuclear Physics, M.V. Lomonosov Moscow State University, Moscow, Russia*  
<sup>102</sup>*Fakultät für Physik, Ludwig-Maximilians-Universität München, München, Germany*  
<sup>103</sup>*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München, Germany*  
<sup>104</sup>*Nagasaki Institute of Applied Science, Nagasaki, Japan*  
<sup>105</sup>*Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya, Japan*  
<sup>106a</sup>*INFN Sezione di Napoli, Italy*  
<sup>106b</sup>*Dipartimento di Fisica, Università di Napoli, Napoli, Italy*  
<sup>107</sup>*Department of Physics and Astronomy, University of New Mexico, Albuquerque, New Mexico, USA*  
<sup>108</sup>*Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands*  
<sup>109</sup>*Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam, Netherlands*  
<sup>110</sup>*Department of Physics, Northern Illinois University, DeKalb, Illinois, USA*  
<sup>111</sup>*Budker Institute of Nuclear Physics, SB RAS, Novosibirsk, Russia*  
<sup>112</sup>*Department of Physics, New York University, New York, New York, USA*  
<sup>113</sup>*Ohio State University, Columbus, Ohio, USA*  
<sup>114</sup>*Faculty of Science, Okayama University, Okayama, Japan*

- <sup>115</sup>*Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman, Oklahoma, USA*
- <sup>116</sup>*Department of Physics, Oklahoma State University, Stillwater, Oklahoma, USA*
- <sup>117</sup>*Palacký University, RCPTM, Olomouc, Czech Republic*
- <sup>118</sup>*Center for High Energy Physics, University of Oregon, Eugene, Oregon, USA*
- <sup>119</sup>*LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France*
- <sup>120</sup>*Graduate School of Science, Osaka University, Osaka, Japan*
- <sup>121</sup>*Department of Physics, University of Oslo, Oslo, Norway*
- <sup>122</sup>*Department of Physics, Oxford University, Oxford, United Kingdom*
- <sup>123a</sup>*INFN Sezione di Pavia, Italy*
- <sup>123b</sup>*Dipartimento di Fisica, Università di Pavia, Pavia, Italy*
- <sup>124</sup>*Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania, USA*
- <sup>125</sup>*National Research Centre “Kurchatov Institute” B.P.Konstantinov Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
- <sup>126a</sup>*INFN Sezione di Pisa, Italy*
- <sup>126b</sup>*Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa, Italy*
- <sup>127</sup>*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania, USA*
- <sup>128a</sup>*Laboratório de Instrumentação e Física Experimental de Partículas—LIP, Lisboa, Portugal*
- <sup>128b</sup>*Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal*
- <sup>128c</sup>*Department of Physics, University of Coimbra, Coimbra, Portugal*
- <sup>128d</sup>*Centro de Física Nuclear da Universidade de Lisboa, Lisboa, Portugal*
- <sup>128e</sup>*Departamento de Física, Universidade do Minho, Braga, Portugal*
- <sup>128f</sup>*Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada, Spain*
- <sup>129</sup>*Institute of Physics, Academy of Sciences of the Czech Republic, Praha, Czech Republic*
- <sup>130</sup>*Czech Technical University in Prague, Praha, Czech Republic*
- <sup>131</sup>*Charles University, Faculty of Mathematics and Physics, Prague, Czech Republic*
- <sup>132</sup>*State Research Center Institute for High Energy Physics (Protvino), NRC KI, Russia*
- <sup>133</sup>*Particle Physics Department, Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>134a</sup>*INFN Sezione di Roma, Italy*
- <sup>134b</sup>*Dipartimento di Fisica, Sapienza Università di Roma, Roma, Italy*
- <sup>135a</sup>*INFN Sezione di Roma Tor Vergata, Italy*
- <sup>135b</sup>*Dipartimento di Fisica, Università di Roma Tor Vergata, Roma, Italy*
- <sup>136a</sup>*INFN Sezione di Roma Tre, Italy*
- <sup>136b</sup>*Dipartimento di Matematica e Fisica, Università Roma Tre, Roma, Italy*
- <sup>137a</sup>*Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies—Université Hassan II, Casablanca, Morocco*
- <sup>137b</sup>*Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech, Morocco*
- <sup>137c</sup>*Faculté des Sciences, Université Mohamed Premier and LPTPM, Oujda, Morocco*
- <sup>137d</sup>*Faculté des sciences, Université Mohammed V, Rabat, Morocco*
- <sup>138</sup>*DSM/IRFU (Institut de Recherches sur les Lois Fondamentales de l’Univers), CEA Saclay (Commissariat à l’Energie Atomique et aux Energies Alternatives), Gif-sur-Yvette, France*
- <sup>139</sup>*Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz, California, USA*
- <sup>140</sup>*Department of Physics, University of Washington, Seattle, Washington, USA*
- <sup>141</sup>*Department of Physics and Astronomy, University of Sheffield, Sheffield, United Kingdom*
- <sup>142</sup>*Department of Physics, Shinshu University, Nagano, Japan*
- <sup>143</sup>*Department Physik, Universität Siegen, Siegen, Germany*
- <sup>144</sup>*Department of Physics, Simon Fraser University, Burnaby, British Columbia, Canada*
- <sup>145</sup>*SLAC National Accelerator Laboratory, Stanford, California, USA*
- <sup>146a</sup>*Faculty of Mathematics, Physics & Informatics, Comenius University, Bratislava, Slovak Republic*
- <sup>146b</sup>*Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice, Slovak Republic*
- <sup>147a</sup>*Department of Physics, University of Cape Town, Cape Town, South Africa*
- <sup>147b</sup>*Department of Physics, University of Johannesburg, Johannesburg, South Africa*
- <sup>147c</sup>*School of Physics, University of the Witwatersrand, Johannesburg, South Africa*
- <sup>148a</sup>*Department of Physics, Stockholm University, Sweden*
- <sup>148b</sup>*The Oskar Klein Centre, Stockholm, Sweden*
- <sup>149</sup>*Physics Department, Royal Institute of Technology, Stockholm, Sweden*

- <sup>150</sup>*Departments of Physics & Astronomy and Chemistry, Stony Brook University, Stony Brook, New York, USA*
- <sup>151</sup>*Department of Physics and Astronomy, University of Sussex, Brighton, United Kingdom*
- <sup>152</sup>*School of Physics, University of Sydney, Sydney, Australia*
- <sup>153</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan*
- <sup>154</sup>*Department of Physics, Technion: Israel Institute of Technology, Haifa, Israel*
- <sup>155</sup>*Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv, Israel*
- <sup>156</sup>*Department of Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece*
- <sup>157</sup>*International Center for Elementary Particle Physics and Department of Physics, The University of Tokyo, Tokyo, Japan*
- <sup>158</sup>*Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo, Japan*
- <sup>159</sup>*Department of Physics, Tokyo Institute of Technology, Tokyo, Japan*
- <sup>160</sup>*Tomsk State University, Tomsk, Russia*
- <sup>161</sup>*Department of Physics, University of Toronto, Toronto, Ontario, Canada*
- <sup>162a</sup>*INFN-TIFPA, Italy*
- <sup>162b</sup>*University of Trento, Trento, Italy*
- <sup>163a</sup>*TRIUMF, Vancouver, British Columbia, Canada*
- <sup>163b</sup>*Department of Physics and Astronomy, York University, Toronto, Ontario, Canada*
- <sup>164</sup>*Faculty of Pure and Applied Sciences, and Center for Integrated Research in Fundamental Science and Engineering, University of Tsukuba, Tsukuba, Japan*
- <sup>165</sup>*Department of Physics and Astronomy, Tufts University, Medford, Massachusetts, USA*
- <sup>166</sup>*Department of Physics and Astronomy, University of California Irvine, Irvine, California, USA*
- <sup>167a</sup>*INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine, Italy*
- <sup>167b</sup>*ICTP, Trieste, Italy*
- <sup>167c</sup>*Dipartimento di Chimica, Fisica e Ambiente, Università di Udine, Udine, Italy*
- <sup>168</sup>*Department of Physics and Astronomy, University of Uppsala, Uppsala, Sweden*
- <sup>169</sup>*Department of Physics, University of Illinois, Urbana, Illinois, USA*
- <sup>170</sup>*Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia—CSIC, Spain*
- <sup>171</sup>*Department of Physics, University of British Columbia, Vancouver, British Columbia, Canada*
- <sup>172</sup>*Department of Physics and Astronomy, University of Victoria, Victoria, British Columbia, Canada*
- <sup>173</sup>*Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>174</sup>*Waseda University, Tokyo, Japan*
- <sup>175</sup>*Department of Particle Physics, The Weizmann Institute of Science, Rehovot, Israel*
- <sup>176</sup>*Department of Physics, University of Wisconsin, Madison, Wisconsin, USA*
- <sup>177</sup>*Fakultät für Physik und Astronomie, Julius-Maximilians-Universität, Würzburg, Germany*
- <sup>178</sup>*Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal, Germany*
- <sup>179</sup>*Department of Physics, Yale University, New Haven, Connecticut, USA*
- <sup>180</sup>*Yerevan Physics Institute, Yerevan, Armenia*
- <sup>181</sup>*Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules (IN2P3), Villeurbanne, France*
- <sup>182</sup>*Academia Sinica Grid Computing, Institute of Physics, Academia Sinica, Taipei, Taiwan*

<sup>a</sup>Deceased.

<sup>b</sup>Also at Department of Physics, King's College London, London, United Kingdom.

<sup>c</sup>Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku, Azerbaijan.

<sup>d</sup>Also at Novosibirsk State University, Novosibirsk, Russia.

<sup>e</sup>Also at TRIUMF, Vancouver, British Columbia, Canada.

<sup>f</sup>Also at Department of Physics & Astronomy, University of Louisville, Louisville, Kentucky, USA.

<sup>g</sup>Also at Physics Department, An-Najah National University, Nablus, Palestine.

<sup>h</sup>Also at Department of Physics, California State University, Fresno, California, USA.

<sup>i</sup>Also at Department of Physics, University of Fribourg, Fribourg, Switzerland.

<sup>j</sup>Also at II Physikalisches Institut, Georg-August-Universität, Göttingen, Germany.

<sup>k</sup>Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona, Spain.

<sup>l</sup>Also at Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Portugal.

<sup>m</sup>Also at Tomsk State University, Tomsk, and Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia.

<sup>n</sup>Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing, China.

<sup>o</sup>Also at Università di Napoli Parthenope, Napoli, Italy.

<sup>p</sup>Also at Institute of Particle Physics (IPP), Canada.

<sup>q</sup>Also at Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania.



- <sup>r</sup> Also at Department of Physics, St. Petersburg State Polytechnical University, St. Petersburg, Russia.
- <sup>s</sup> Also at Borough of Manhattan Community College, City University of New York, New York City, New York, USA.
- <sup>t</sup> Also at Department of Financial and Management Engineering, University of the Aegean, Chios, Greece.
- <sup>u</sup> Also at Centre for High Performance Computing, CSIR Campus, Rosebank, Cape Town, South Africa.
- <sup>v</sup> Also at Louisiana Tech University, Ruston, Louisiana, USA.
- <sup>w</sup> Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona, Spain.
- <sup>x</sup> Also at Department of Physics, The University of Michigan, Ann Arbor, Michigan, USA.
- <sup>y</sup> Also at LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France.
- <sup>z</sup> Also at Graduate School of Science, Osaka University, Osaka, Japan.
- <sup>aa</sup> Also at Fakultät für Mathematik und Physik, Albert-Ludwigs-Universität, Freiburg, Germany.
- <sup>bb</sup> Also at Institute for Mathematics, Astrophysics and Particle Physics, Radboud University Nijmegen/Nikhef, Nijmegen, Netherlands.
- <sup>cc</sup> Also at Department of Physics, The University of Texas at Austin, Austin, Texas, USA.
- <sup>dd</sup> Also at Institute of Theoretical Physics, Ilia State University, Tbilisi, Georgia.
- <sup>ee</sup> Also at CERN, Geneva, Switzerland.
- <sup>ff</sup> Also at Georgian Technical University (GTU), Tbilisi, Georgia.
- <sup>gg</sup> Also at Ochadai Academic Production, Ochanomizu University, Tokyo, Japan.
- <sup>hh</sup> Also at Manhattan College, New York, New York, USA.
- <sup>ii</sup> Also at The City College of New York, New York, New York, USA.
- <sup>jj</sup> Also at Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada, Spain.
- <sup>kk</sup> Also at Department of Physics, California State University, Sacramento, California, USA.
- <sup>ll</sup> Also at Moscow Institute of Physics and Technology State University, Dolgoprudny, Russia.
- <sup>mm</sup> Also at Departement de Physique Nucleaire et Corpusculaire, Université de Genève, Geneva, Switzerland.
- <sup>nn</sup> Also at Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Barcelona, Spain.
- <sup>oo</sup> Also at School of Physics, Sun Yat-sen University, Guangzhou, China.
- <sup>pp</sup> Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia, Bulgaria.
- <sup>qq</sup> Also at Faculty of Physics, M.V.Lomonosov Moscow State University, Moscow, Russia.
- <sup>rr</sup> Also at National Research Nuclear University MEPhI, Moscow, Russia.
- <sup>ss</sup> Also at Department of Physics, Stanford University, Stanford, California, USA.
- <sup>tt</sup> Also at Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Budapest, Hungary.
- <sup>uu</sup> Also at Giresun University, Faculty of Engineering, Turkey.
- <sup>vv</sup> Also at CPPM, Aix-Marseille Université and CNRS/IN2P3, Marseille, France.
- <sup>ww</sup> Also at Department of Physics, Nanjing University, Jiangsu, China.
- <sup>xx</sup> Also at Institute of Physics, Academia Sinica, Taipei, Taiwan.
- <sup>yy</sup> Also at University of Malaya, Department of Physics, Kuala Lumpur, Malaysia.