

# Mechanism of Action of *Melaleuca alternifolia* (Tea Tree) Oil on *Staphylococcus aureus* Determined by Time-Kill, Lysis, Leakage, and Salt Tolerance Assays and Electron Microscopy

Christine F. Carson,<sup>1\*</sup> Brian J. Mee,<sup>1</sup> and Thomas V. Riley<sup>1,2</sup>

Department of Microbiology, The University of Western Australia, Crawley,<sup>1</sup> and Division of Microbiology and Infectious Diseases, The Western Australian Centre for Pathology and Medical Research, Nedlands,<sup>2</sup> Western Australia 6009, Australia

Received 24 April 2001/Returned for modification 19 August 2001/Accepted 6 February 2002

**The essential oil of *Melaleuca alternifolia* (tea tree) has broad-spectrum antimicrobial activity. The mechanisms of action of tea tree oil and three of its components, 1,8-cineole, terpinen-4-ol, and  $\alpha$ -terpineol, against *Staphylococcus aureus* ATCC 9144 were investigated. Treatment with these agents at their MICs and two times their MICs, particularly treatment with terpinen-4-ol and  $\alpha$ -terpineol, reduced the viability of *S. aureus*. None of the agents caused lysis, as determined by measurement of the optical density at 620 nm, although cells became disproportionately sensitive to subsequent autolysis. Loss of 260-nm-absorbing material occurred after treatment with concentrations equivalent to the MIC, particularly after treatment with 1,8-cineole and  $\alpha$ -terpineol. *S. aureus* organisms treated with tea tree oil or its components at the MIC or two times the MIC showed a significant loss of tolerance to NaCl. When the agents were tested at one-half the MIC, only 1,8-cineole significantly reduced the tolerance of *S. aureus* to NaCl. Electron microscopy of terpinen-4-ol-treated cells showed the formation of mesosomes and the loss of cytoplasmic contents. The predisposition to lysis, the loss of 260-nm-absorbing material, the loss of tolerance to NaCl, and the altered morphology seen by electron microscopy all suggest that tea tree oil and its components compromise the cytoplasmic membrane.**

The essential oil derived by steam distillation from the leaves of *Melaleuca alternifolia* is also known as tea tree oil (TTO) or *Melaleuca* oil. TTO is well characterized and contains approximately 100 terpenes and their related alcohols (6). The physical and chemical properties of commercial TTO are regulated by an international standard (23). TTO has antibacterial (8, 17), antifungal (18, 19), antiviral (4), and anti-inflammatory (5) properties in vitro, suggesting that it may have a role in the treatment of cutaneous infection. Clinical trials have demonstrated that TTO may be efficacious in the treatment of acne (2) and oral candidiasis (24) and in the decolonization of methicillin-resistant *Staphylococcus aureus* carriers (7). Although the in vitro antimicrobial activity and in vivo efficacy of TTO have been reported, less is known about its mechanism of action. Since this will have implications for its spectrum of activity, selective toxicity, and the development of resistance, we examined the mechanism of action of TTO and its components against *S. aureus*.

## MATERIALS AND METHODS

**TTO and components.** TTO (batch 971) was kindly provided by Australian Plantations Pty. Ltd., Wyrallah, New South Wales, Australia. The levels of the components, determined by gas chromatographic analysis according to the international standard (23), were as follows: 41.5% terpinen-4-ol, 21.2%  $\gamma$ -terpinene, 10.2%  $\alpha$ -terpinene, 3.5% terpinolene, 2.9%  $\alpha$ -terpineol, 2.5%  $\alpha$ -pinene, 2.1% 1,8-cineole, 1.5%  $p$ -cymene, 1% aromadendrene, 1%  $\delta$ -cadinene, 0.9% ledene, 0.9% limonene, 0.6% globulol, 0.4% sabinene, and 0.3% viridiflorol. Terpinen-4-ol and  $\alpha$ -terpineol (Aldrich Chemical Co. Inc., Milwaukee, Wis.)

were 97 and 98% pure, respectively. 1,8-Cineole (Sigma Chemical Co., St. Louis, Mo.) was at least 99% pure. Terpinen-4-ol and  $\alpha$ -terpineol were chosen on the basis of their antimicrobial activities (10). 1,8-Cineole was selected since it has some antimicrobial activity and its presence in TTO has been contentious, due mainly to its erroneous reputation as a skin irritant (10).

**Bacterium.** *S. aureus* ATCC 9144 was used in all experiments and was obtained from the culture collection of the Department of Microbiology, The University of Western Australia, Crawley, Western Australia, Australia.

**Inoculum effect.** The MICs and minimal bactericidal concentrations (MBCs) of TTO for *S. aureus* were determined by a previously described broth microdilution method (8), except that the test medium was Mueller-Hinton broth (Oxoid Ltd., Basingstoke, United Kingdom), the Tween 80 concentration was reduced to 0.001% (vol/vol), and several inoculum concentrations were used. Final inoculum concentrations were  $5 \times 10^3$ ,  $5 \times 10^5$ ,  $5 \times 10^7$ , and  $5 \times 10^9$  CFU/ml.

**Preparation of bacterial suspensions.** Unless stated otherwise, suspensions of organisms in the stationary phase of growth were prepared by inoculating two colonies of *S. aureus* from overnight cultures on blood agar into 400 ml of Mueller-Hinton broth, which was incubated at 37°C for 18 h with shaking. After incubation, the bacteria were separated from the growth medium by centrifugation at  $10,000 \times g$  for 12 min at 4°C, washed twice with phosphate-buffered saline (PBS; pH 7.4), and resuspended in PBS supplemented with 0.001% Tween 80 (Sigma) (PBS-T). Suspensions were adjusted so that the optical density at 620 nm ( $OD_{620}$ ) of a 1-in-100 dilution was  $\sim 0.3$ , which was  $\sim 3 \times 10^{10}$  CFU/ml.

**General experimental conditions.** Experiments were conducted at room temperature (22°C). The concentrations of TTO and its components used for treatment were derived from MICs previously determined by broth microdilution methods and ranged from one-half to two times the MICs. The MICs of TTO, terpinen-4-ol, and  $\alpha$ -terpineol for *S. aureus* were 0.25%; and the MIC of 1,8-cineole was 0.5% (9, 10). Stock concentrations of TTO or its components were prepared in PBS-T at 10-fold the desired final concentration (in percent [vol/vol]) and were added to bacterial suspensions at a ratio of 1:9. PBS-T was added to the control suspensions. No inactivating agents have been established for TTO or its components. In the absence of an inoculum effect, dilution was used to arrest treatment and reduce carryover. The minimum dilution used was 1 in 10. All serial dilutions were done in PBS-T. All experiments except time-kill assays and inoculum effect assays were conducted four to six times; time-kill assays were done in triplicate, and inoculum effect assays were done in duplicate. Viable counts were plated in duplicate by using a spiral plater (Don Whitley Scientific Ltd., Shipley, United Kingdom) and nutrient agar (NA) (Oxoid) in all experi-

\* Corresponding author. Mailing address: Department of Microbiology, The University of Western Australia, Crawley, Western Australia 6009, Australia. Phone: 61 8 9346 3288. Fax: 61 8 9346 2912. E-mail: carson@cyllene.uwa.edu.au.




FIG. 1. Time-kill curves of *S. aureus* ATCC 9144 in control suspensions ( $\blacklozenge$ ) and after treatment with TTO (A), 1,8-cineole (B), terpinen-4-ol (C), or  $\alpha$ -terpineol (D) at one-half the MICs ( $\triangle$ ), the MICs ( $\bullet$ ), and two times the MICs ( $\square$ ). The MICs of TTO, terpinen-4-ol, and  $\alpha$ -terpineol were 0.25% (vol/vol); and that of 1,8-cineole was 0.5%. The organisms were suspended in PBS-T. Each symbol indicates the means  $\pm$  SEs for at least three replicates. The lower detection threshold was  $10^3$  CFU/ml.

ments with the exception of those investigating an inoculum effect. After incubation at 37°C for 24 to 72 h, the colonies were counted manually with the aid of a counting template (Don Whitley Scientific). The detection threshold was  $10^3$  CFU/ml. For inoculum effect assays, 10  $\mu$ l of serial 10-fold dilutions of inocula was plated onto blood agar in quadruplicate, the plates were incubated, and the resulting colonies were counted. In assays for bacterial killing, bacteriolysis, and loss of 260-nm-absorbing material, samples taken 5 min before treatment addition were adjusted to account for dilution and used to determine the concentrations and ODs at 0 min.

**Bacterial killing assays.** The activities of TTO and its components against *S. aureus* were evaluated by measuring the reduction in the numbers of CFU per milliliter over 2 h. Bacterial suspensions (5 ml) were prepared as described above. After a pretreatment sample (0.5 ml) was taken, TTO or one of its components was added to yield final concentrations of one-half, one, and two times the respective MICs. The suspensions were mixed for 20 s with a vortex mixer; and a sample (0.5 ml) was removed at 30 s, serially diluted, and plated onto NA. Additional samples were taken at 30, 60, and 120 min. When rapid killing occurred, samples were also taken at 3, 6, 9, 12, and 15 min. Samples were

diluted, plated onto NA, and incubated overnight. The mean number of survivors was determined.

**Bacteriolysis.** Suspensions of *S. aureus* were prepared as described above. After retrieval of the pretreatment sample, the agents were added so that the suspensions contained TTO or one of its components at concentrations equivalent to the MIC and two times the MIC. The suspensions were mixed for 20 s with a vortex mixer, a sample was removed at 30 s and diluted 1 in 100, and the OD<sub>620</sub> was measured. Additional samples were taken at 30, 60, 90, and 120 min; and the ODs were measured immediately. Corresponding dilutions of test agents were used as blanks. Preliminary work indicated that the ODs of undiluted samples and samples diluted 1 in 10 were inconsistent (data not shown). The 1-in-100 dilution reduced the level of carryover of TTO or its components to levels which contributed negligibly to the absorbance. The results were expressed as a ratio of the OD<sub>620</sub> at each time interval versus the OD<sub>620</sub> at 0 min (in percent).

In two separate experiments, after the initial OD<sub>620</sub> measurement, the samples were retained and the OD<sub>620</sub> was remeasured 6.5 or 23 h later. The OD<sub>620</sub>s of the control suspensions diminished over these time intervals; the OD<sub>620</sub>s of the

samples obtained at 0 min were measured again 6.5 h later and were 91.5% of the original OD<sub>620</sub>, while the OD<sub>620</sub>s measured 23 h later were 84.4% of the original OD<sub>620</sub>. Since there was an inherent progressive reduction in the OD<sub>620</sub> over time, when the OD<sub>620</sub>s of the samples were measured 6.5 or 23 h later, their OD<sub>620</sub>s were compared to the OD<sub>620</sub>s of the 0-min samples, which were also remeasured 6.5 or 23 h later.

**Loss of 260-nm-absorbing material.** Suspensions of bacteria were prepared as described above. A pretreatment sample was taken, diluted 1 in 100, and filtered through a 0.2- $\mu$ m-pore-size filter (Gelman Sciences, Ann Arbor, Mich.). The treatment agents were added at final concentrations equivalent to their MICs. Additional samples were removed after 30 and 60 min, diluted, and filtered as described above. Filtrates of the appropriate dilution of each agent were prepared, and 200- $\mu$ l volumes were used to blank the wells of a UV-transparent 96-well tray (Molecular Devices, Sunnyvale, Calif.) at 260 nm. The blanks were discarded, the wells were loaded with the corresponding test filtrates, and the OD<sub>260</sub>s were determined. The OD<sub>260</sub>s of four to eight samples obtained at each time point were measured. The OD<sub>260</sub> at each time point was expressed as a proportion of the initial OD<sub>260</sub>. Mean ratios for each treatment agent and time were calculated and compared to the means for the corresponding untreated samples by using the two-tailed Student *t* test ( $P < 0.05$ ).

**Loss of salt tolerance.** The ability of *S. aureus* cells treated with TTO or its components to grow on NA supplemented with NaCl (Merck Pty. Ltd., Kilsyth, Victoria, Australia) was investigated. In preliminary experiments, untreated suspensions of *S. aureus* were plated onto NA and NA containing NaCl at 5 to 100 g/liter (NA-NaCl) and incubated, and the resulting colonies were counted. Concentrations of NaCl that modestly compromised the colony-forming abilities of untreated organisms were selected. These were 50 and 75 g/liter. Suspensions of bacteria were prepared as described above and treated with TTO and 1,8-cineole at one-half, one, and two times their MICs and with terpinen-4-ol and  $\alpha$ -terpineol at one-half and one time their MICs. After 30 min, samples were removed, serially diluted, and inoculated onto NA and the selective medium NA-NaCl. After incubation, the numbers of CFU per milliliter on each NA-NaCl plate were compared to those on the NA plate, and the result was expressed as a percentage. The mean proportions of survivors from treated suspensions were compared to the corresponding means for the untreated controls by the two-tailed Student *t* test ( $P < 0.05$ ).

**Electron microscopy of terpinen-4-ol-treated bacteria.** Suspensions of *S. aureus* in the stationary phase of growth were prepared by inoculating and incubating 30 ml of heart infusion broth (HIB; Gibco Diagnostics, Madison, Wis.). Organisms were harvested by centrifugation at 1,500  $\times$  g for 10 min, and the pellet was resuspended in HIB supplemented with Tween 80 (0.5% [vol/vol]) (HIB-T). Suspensions of *S. aureus* were treated with 0.3% terpinen-4-ol for 10 min. Controls in HIB-T stood for 10 min. After centrifugation at 1,500  $\times$  g for 5 min, the pellets were fixed overnight in 2.5% glutaraldehyde in 0.1 M cacodylate buffer at room temperature. Fixed microbial pellets were processed in graded alcohols, propylene oxide, and araldite and cured for 48 h at 60°C. Ultrathin sections were stained with uranyl acetate and lead citrate and were examined with a Philips 410 transmission electron microscope at an accelerating voltage of 80 kV.

## RESULTS

**Inoculum effect.** Inoculum concentrations over the range 10<sup>3</sup> to 10<sup>9</sup> CFU/ml did not significantly alter the MICs or MBCs of TTO. At an inoculum of 5  $\times$  10<sup>3</sup> CFU/ml, the MIC was 0.12% and the MBC was 0.25%. At inocula of 5  $\times$  10<sup>5</sup> and 5  $\times$  10<sup>7</sup> CFU/ml, the MICs were both 0.25% and the MBCs were both 0.5%. At an inoculum of 5  $\times$  10<sup>9</sup> CFU/ml, the MIC was 0.5% and the MBC was 1%.

**Bacterial killing assays.** Treatment of *S. aureus* with TTO at two times the MIC reduced the viability of *S. aureus* by  $\sim$ 5 log<sub>10</sub> over 120 min, while treatment with TTO at the MIC resulted in an  $\sim$ 1 log<sub>10</sub> reduction (Fig. 1A). Treatment with 1,8-cineole at the MIC or two times the MIC resulted in greater than 3 log<sub>10</sub> reductions over 120 min (Fig. 1B). Terpinen-4-ol or  $\alpha$ -terpineol at one-half times the MIC had little effect on the viability of *S. aureus*. In contrast, treatment with either of these components at two times the MIC effected reductions of  $\sim$ 7.5 and  $\sim$ 6.5 log<sub>10</sub> in 9 and 30 min, respectively

TABLE 1. Proportion of initial OD<sub>620</sub> of suspensions of *S. aureus* remaining after treatment with TTO or its components for 120 min

Treatment agent and concn (fold MIC) <sup>a</sup>	OD <sub>620</sub> at 120 min/OD <sub>620</sub> at 0 min (%) measured <sup>b</sup> :		
	Immediately	After 6.5 h	After 23 h
Control	96.9 $\pm$ 1.3 <sup>c</sup>	101.9	98.9
TTO			
One time	95.1 $\pm$ 3.1	89.6	ND <sup>d</sup>
Two times	96.8 $\pm$ 1.4	85.5	ND
1,8-Cineole			
One time	94.8 $\pm$ 0.6	ND	40.8
Two times	95.7 $\pm$ 1.0	ND	33.3
Terpinen-4-ol			
One time	91.0 $\pm$ 3.9	75.6	27.2
Two times	91.6 $\pm$ 1.3	53.4	28.5
$\alpha$ -Terpineol			
One time	91.3 $\pm$ 1.5	78.9	28.4
Two times	89.4 $\pm$ 1.6	55.8	29.5

<sup>a</sup> The MICs of TTO, terpinen-4-ol, and  $\alpha$ -terpineol were 0.25% (vol/vol); and that of 1,8-cineole was 0.5%.

<sup>b</sup> In two separate experiments, after the initial immediate measurement, the samples were retained and the OD<sub>620</sub>s were remeasured 6.5 or 23 h later.

<sup>c</sup> Values are means  $\pm$  SEs.

<sup>d</sup> ND, not done.

(Fig. 1C and D). Treatment with terpinen-4-ol at the MIC consistently reduced the viability by 6.5 log<sub>10</sub> over 120 min, while treatment with  $\alpha$ -terpineol at the MIC gave inconsistent results. Representative results are shown in Fig. 1D.

**Bacteriolysis.** The mean OD<sub>620</sub> of untreated *S. aureus* suspensions after 120 min was 96.9% ( $n = 9$ ; standard error [SE] = 1.3%) of the original. Table 1 shows the OD<sub>620</sub>s of suspensions treated with TTO or its components. Treatment with TTO or 1,8-cineole at the MIC or two times the MIC had no effect (OD<sub>620</sub> range, 95 to 97% of the original). Treatment with terpinen-4-ol or  $\alpha$ -terpineol reduced the OD<sub>620</sub> slightly. Table




FIG. 2. Appearance of 260-nm-absorbing material in the filtrates of *S. aureus* control suspensions (white bars) and after treatment with the MICs of TTO (0.25%; bars with diagonal stripes), 1,8-cineole (0.5%; grey bars), terpinen-4-ol (0.25%; bars with horizontal stripes), or  $\alpha$ -terpineol (0.25%; black bars). The organisms were suspended in PBS-T. The means  $\pm$  SEs for at least three replicates are illustrated.




FIG. 3. Proportion of *S. aureus* cells able to form colonies on NA (white bars), NA supplemented with 50 g of NaCl per liter (black bars), and NA supplemented with 75 g of NaCl per liter (grey bars) after 30 min of treatment with TTO (A), 1,8-cineole (B), terpinen-4-ol (C), or  $\alpha$ -terpineol (D) at one-half the MICs, the MICs, and two times the MICs. The MICs of TTO, terpinen-4-ol, and  $\alpha$ -terpineol were 0.25% (vol/vol); and that of 1,8-cineole was 0.5%. The organisms were suspended in PBS-T. The means  $\pm$  SEs for at least four replicates are illustrated.

1 also shows the results obtained when the OD<sub>620s</sub> of dilutions of *S. aureus* suspensions was remeasured either 6.5 or 23 h after the initial OD<sub>620s</sub> were measured. Modest reductions in the OD<sub>620s</sub> of the TTO-treated suspensions were noted when they were measured again 6.5 h later. In contrast, large reductions were noted in the OD<sub>620s</sub> of the suspensions treated with the components of TTO, particularly those left for 23 h, in which only ~30% of the original OD<sub>620s</sub> remained. Notably, the OD<sub>620s</sub> of the suspensions treated with terpinen-4-ol or  $\alpha$ -terpineol at two times the MIC for 30 s were reduced to 44.9 and 67.4%, respectively, when they were read again 23 h later (data not shown).

**Loss of 260-nm-absorbing material.** The OD<sub>260s</sub> of filtrates from control suspensions were not significantly different after 30 min but were significantly different ( $P = 0.020$ ) after 60 min (Fig. 2). Significant increases in the OD<sub>260s</sub> occurred after 60 min of treatment with TTO ( $P = 0.019$ ), 1,8-cineole ( $P = 4.3 \times 10^{-4}$ ), terpinen-4-ol ( $P = 0.038$ ), or  $\alpha$ -terpineol ( $P = 0.005$ ) at the MIC. Only 1,8-cineole treatment resulted in significant ( $P = 1.4 \times 10^{-7}$ ) increases after 30 min.

**Loss of salt tolerance.** The results of tests of salt tolerance loss are shown in Fig. 3A to D. The addition of NaCl to NA reduced the colony-forming ability of untreated *S. aureus* cells to 81.4% ( $n = 22$ ; SE = 2.5%) with NaCl at 50 g/liter and






FIG. 4. Electron micrographs of *S. aureus* cells stained with uranyl acetate after no treatment (A) and after treatment with 0.3% terpinen-4-ol for 10 min (B and C). Magnifications:  $\times 11,500$  (A) and  $\times 14,200$  (B and C).

0.032). Of the four treatments at one-half the MIC, only 1,8-cineole significantly reduced the colony-forming ability of the survivors, and this was only when the higher salt concentration was used ( $P = 0.030$ ).

**Electron microscopy of terpinen-4-ol-treated bacteria.** Terpinen-4-ol-treated *S. aureus* cells contained multilamellar, mesosomelike structures (Fig. 4B and C) that were not seen in untreated cells (Fig. 4A). In addition, the contents of some treated cells appeared depleted and amorphous (Fig. 4C).

## DISCUSSION

Interactions with the hydrophobic structures of bacteria play a key role in the antimicrobial actions of hydrocarbons (32). Consequently, assumptions regarding the mechanism of action of TTO have been based on the nature of its components. In the present investigation, *S. aureus* cells in the stationary phase of growth were killed by TTO and its components. Organisms in this growth phase are generally less sensitive to injury than those in the exponential phase (11), and this has been shown for *Escherichia coli* treated with TTO (16). Since antimicrobial agents that affect cellular synthetic processes often have little effect on organisms in the stationary phase of growth (15, 28), these results suggest that the principal target of TTO is not a macromolecular synthetic process.

Some antimicrobial agents cause gross membrane damage and provoke whole-cell lysis (14, 28); and this has been reported previously for essential oils from oregano, rosewood, and thyme (20). However, the major components of these oils, including carvacrol, citronellol, geraniol, and thymol (38), are not found in TTO. The failure of TTO or its components to lyse *S. aureus* cells suggests that their primary mechanism of

73.3% ( $n = 22$ ; SE = 2.8%) with NaCl at 75 g/liter. Treatment with TTO or 1,8-cineole at two times their MICs significantly reduced the ability of survivors to form colonies on NA-NaCl (with NaCl at 50 or 75 g/liter), with only 1.5% or less of the survivors able to form colonies. 1,8-Cineole at the MIC had a similar effect, with 0.7% or less of the survivors able to form colonies on NA-NaCl. Treatment with TTO and terpinen-4-ol at their MICs also reduced the ability of survivors to form colonies on NA-NaCl, but to a lesser extent (<30%). By treatment with  $\alpha$ -terpineol at the MIC, the proportion of survivors able to form colonies on NA-NaCl was not significantly reduced when NaCl was used at 50 g/liter ( $P = 0.072$ ) but was significantly reduced when NaCl was used at 75 g/liter ( $P =$

action is not gross cell wall damage. Treatment-induced release of membrane-bound cell wall autolytic enzymes will induce lysis eventually (15), and this may explain the delayed lysis of *S. aureus* seen when suspensions were reexamined after several hours; *S. aureus* cells treated for as little as 30 s were disproportionately sensitive to lysis. While the activation of autolytic enzymes may have been responsible for this effect, the lysis may also have been due to weakening of the cell wall and the subsequent rupture of the cytoplasmic membrane due to osmotic pressure (rather than a specific action on the cytoplasmic membrane).

Marked leakage of cytoplasmic material is considered indicative of gross and irreversible damage to the cytoplasmic membrane (21). Many antimicrobial compounds that act on the bacterial cytoplasmic membrane induce the loss of 260-nm-absorbing material, including chlorhexidine (21), hexachlorophene (25), phenethyl alcohol (34), tetracyclines, polymyxin (12),  $\alpha$ -pinene (1), and lemongrass oil (27). *S. aureus* suspensions treated with TTO or its components lost significant 260-nm-absorbing material, suggesting that nucleic acids were lost through a damaged cytoplasmic membrane.

Sublethal injury of microbial cell membranes may alter their permeability and affect the membrane's ability to osmoregulate the cell adequately or to exclude toxic materials (15). Consequently, the loss of tolerance to salts or other potentially toxic compounds may be exploited to reveal membrane damage (22) in sublethally injured bacteria. Treatment of *S. aureus* with TTO or its components significantly reduced the ability of the survivors to form colonies on media containing NaCl. This effect was most marked with 1,8-cineole, which was also the only compound with which treatment with one-half the MIC resulted in a significant loss of salt tolerance. These results correlate well with the OD<sub>260</sub> results since, in each case, treatment with TTO or its components at the MICs induced the loss of salt tolerance and 260-nm-absorbing material. Notably, treatment with TTO, terpinen-4-ol, or  $\alpha$ -terpineol at one-half the MIC had no effect on salt tolerance or actually increased it. Treatment with these agents at one-half their MICs may have killed the most susceptible cells, leaving the more robust and, possibly, more salt-tolerant cells. However, of the three treatments with which an increase in salt tolerance was noted, only treatment with TTO at one-half the MIC for 30 min reduced the viability of *S. aureus*. Alternatively, the treatments may have induced the expression of stress tolerance mechanisms.

Electron microscopy of terpinen-4-ol-treated cells corroborated the inability of terpinen-4-ol to lyse *S. aureus* and suggested membrane damage by the appearance of mesosomes and a loss of cytoplasmic material. These lesions have been reported previously after treatment with antimicrobial agents including vancomycin (29), phenethyl alcohol (11), defensins (30), and betane (3). Similar cytoplasmic losses, as well as bleb formation, were reported for *E. coli* treated with lemongrass oil (26), and both effects were noted when *E. coli* was treated with TTO (16).

The original premise was that TTO and/or its components act on microbial membranes. The loss of 260-nm-absorbing material, increased susceptibility to NaCl, the formation of mesosomes, and the loss of cytoplasmic material suggest that the cytoplasmic membrane is compromised by treatment with TTO and its components. Additional support for this idea

comes from work showing that TTO treatment of *E. coli* or *S. aureus* induces the loss of potassium ions, inhibits respiration, and promotes the uptake of propidium iodide (13). In other work done with essential oils and their components,  $\alpha$ -pinene inhibited respiratory activity in yeast mitochondria (1) and  $\beta$ -pinene and limonene had similar effects in intact yeast cells and isolated mitochondria (35). Collectively, these observations suggest that further work could examine the effect of treatment on microbial energy transduction, in particular, the transmembrane proton motive force.

When the mechanisms of action of lipophilic biocides have been examined, effects on the cytoplasmic membrane and/or the enzymes embedded in it have been demonstrated (1, 3, 31, 33, 35–37). The data on TTO and its components support these observations. However, the possibility remains that sites of action other than the cytoplasmic membrane exist. Furthermore, it is possible that other oil components, not examined thus far, may contribute to the antimicrobial activity of the oil by other mechanisms. Given the heterogeneous composition of TTO and the antimicrobial activities of many of its components, it seems unlikely that there is only one mechanism of action or that only one component is responsible for the antimicrobial action. Further work is required to understand fully the mechanisms involved.

#### ACKNOWLEDGMENTS

This work was supported by a grant (grant UWA-24A) from the Rural Industries Research & Development Corporation, Canberra, Australian Capital Territory, Australia.

#### REFERENCES

1. Andrews, R. E., L. W. Parks, and K. D. Spence. 1980. Some effects of Douglas fir terpenes on certain microorganisms. *Appl. Environ. Microbiol.* **40**:301–304.
2. Bassett, I. B., D. L. Pannowitz, and R. S. C. Barnetson. 1990. A comparative study of tea tree oil versus benzoylperoxide in the treatment of acne. *Med. J. Aust.* **153**:455–458.
3. Beveridge, E. G., I. Boyd, I. Dew, M. Haswell, and C. W. G. Lowe. 1991. Electron and light microscopy of damaged bacteria, p. 135–153. *In* S. P. Denyer and W. B. Hugo (ed.), *Mechanisms of action of chemical biocides: their study and exploitation*. Blackwell Scientific Publications, Oxford, United Kingdom.
4. Bishop, C. D. 1995. Antiviral activity of the essential oil of *Melaleuca alternifolia* (Maiden and Betche) Cheel (tea tree) against tobacco mosaic virus. *J. Essent. Oil Res.* **7**:641–644.
5. Brand, C., A. Ferrante, R. H. Prager, T. V. Riley, C. F. Carson, J. J. Finlay-Jones, and P. H. Hart. 2001. The water-soluble components of the essential oil of *Melaleuca alternifolia* (tea tree oil) suppress the production of superoxide by human monocytes, but not neutrophils, activated in vitro. *Inflamm. Res.* **50**:213–219.
6. Brophy, J. J., N. W. Davies, I. A. Southwell, I. A. Stiff, and L. R. Williams. 1989. Gas chromatographic quality control for oil of *Melaleuca terpinen-4-ol* type (Australian tea tree). *J. Agric. Food Chem.* **37**:1330–1335.
7. Caelli, M., J. Porteous, C. F. Carson, R. Heller, and T. V. Riley. 2000. Tea tree oil as an alternative topical decolonisation agent for methicillin-resistant *Staphylococcus aureus*. *J. Hosp. Infect.* **46**:236–237.
8. Carson, C. F., B. D. Cookson, H. D. Farrelly, and T. V. Riley. 1995. Susceptibility of methicillin-resistant *Staphylococcus aureus* to the essential oil of *Melaleuca alternifolia*. *J. Antimicrob. Chemother.* **35**:421–424.
9. Carson, C. F., K. A. Hammer, and T. V. Riley. 1995. Broth micro-dilution method for determining the susceptibility of *Escherichia coli* and *Staphylococcus aureus* to the essential oil of *Melaleuca alternifolia* (tea tree oil). *Microbios* **82**:181–185.
10. Carson, C. F., and T. V. Riley. 1995. Antimicrobial activity of the major components of the essential oil of *Melaleuca alternifolia*. *J. Appl. Bacteriol.* **78**:264–269.
11. Corre, J., J. J. Lucchini, G. M. Mercier, and A. Cremieux. 1990. Antibacterial activity of phenethyl alcohol and resulting membrane alterations. *Res. Microbiol.* **141**:483–497.
12. Corry, J. E. L., H. Van Doorne, and D. A. A. Mossel. 1977. Recovery and revival of microbial cells, especially those from environments containing

- antibiotics, p. 174–196. In M. Woodbine (ed.), *Antibiotics in agriculture*. Proceedings of the 25th Easter School, University of Nottingham. Butterworths, London, United Kingdom.
13. Cox, S. D., C. M. Mann, J. L. Markham, H. C. Bell, J. E. Gustafson, J. R. Warmington, and S. G. Wyllie. 2000. The mode of antimicrobial action of the essential oil of *Melaleuca alternifolia* (tea tree oil). *J. Appl. Microbiol.* **88**:170–175.
  14. Denyer, S. P., and W. B. Hugo. 1991. Biocide-induced damage to the bacterial cytoplasmic membrane, p. 171–187. In S. P. Denyer and W. B. Hugo (ed.), *Mechanisms of action of chemical biocides: their study and exploitation*. Blackwell Scientific Publications, Oxford, United Kingdom.
  15. Gilbert, P. 1984. The revival of microorganisms sublethally injured by chemical inhibitors, p. 175–197. In M. H. E. Andrews and A. D. Russell (ed.), *The revival of injured microbes*. Academic Press, London, United Kingdom.
  16. Gustafson, J. E., Y. C. Liew, S. Chew, J. Markham, H. C. Bell, S. G. Wyllie, and J. R. Warmington. 1998. Effects of tea tree oil on *Escherichia coli*. *Lett. Appl. Microbiol.* **26**:194–198.
  17. Hammer, K. A., C. F. Carson, and T. V. Riley. 1996. Susceptibility of transient and commensal skin flora to the essential oil of *Melaleuca alternifolia* (tea tree oil). *Am. J. Infect. Control* **24**:186–189.
  18. Hammer, K. A., C. F. Carson, and T. V. Riley. 1997. In vitro susceptibility of *Malassezia furfur* to the essential oil of *Melaleuca alternifolia*. *J. Med. Vet. Mycol.* **35**:375–377.
  19. Hammer, K. A., C. F. Carson, and T. V. Riley. 1998. In vitro activity of essential oils, in particular *Melaleuca alternifolia* (tea tree) oil and tea tree oil products, against *Candida* spp. *J. Antimicrob. Chemother.* **42**:591–595.
  20. Horne, D. S., M. Holm, C. Oberg, S. Chao, and D. G. Young. 2001. Antimicrobial effects of essential oils on *Streptococcus pneumoniae*. *J. Essent. Oil Res.* **13**:387–392.
  21. Hugo, W. B., and R. Longworth. 1964. Some aspects of the mode of action of chlorhexidine. *J. Pharm. Pharmacol.* **16**:655–662.
  22. Iandolo, J. J., and Z. J. Ordal. 1966. Repair of thermal injury of *Staphylococcus aureus*. *J. Bacteriol.* **91**:134–142.
  23. International Organisation for Standardisation. 1996. Essential oils—oil of *Melaleuca*, terpinen-4-ol type (tea tree oil). ISO-4730. International Organisation for Standardisation, Geneva, Switzerland.
  24. Jandourek, A., J. K. Vaishampayan, and J. A. Vazquez. 1998. Efficacy of melaleuca oral solution for the treatment of fluconazole refractory oral candidiasis in AIDS patients. *AIDS* **12**:1033–1037.
  25. Joswick, H. L., T. R. Corner, J. N. Silvernale, and P. Gerhardt. 1971. Antimicrobial actions of hexachlorophene: release of cytoplasmic materials. *J. Bacteriol.* **108**:492–500.
  26. Ogunlana, E. O., S. Höglund, G. Onawunmi, and O. Sköld. 1987. Effects of lemongrass oil on the morphological characteristics and peptidoglycan synthesis of *Escherichia coli* cells. *Microbios* **50**:43–59.
  27. Onawunmi, G. O., and E. O. Ogunlana. 1985. Effects of lemon grass oil on the cells and spheroplasts of *Escherichia coli* NCTC 9001. *Microbios Lett.* **28**:63–68.
  28. Russell, A. D., A. Morris, and M. C. Allwood. 1973. Methods for assessing damage to bacteria induced by chemical and physical agents. *Methods Microbiol.* **8**:95–182.
  29. Sanyal, D., and D. Greenwood. 1993. An electron microscope study of glycopeptide antibiotic-resistant strains of *Staphylococcus epidermidis*. *J. Med. Microbiol.* **39**:204–210.
  30. Shimoda, M., K. Ohki, Y. Shimamoto, and O. Kohashi. 1995. Morphology of defensin-treated *Staphylococcus aureus*. *Infect. Immun.* **63**:2886–2891.
  31. Sikkema, J., J. A. M. de Bont, and B. Poolman. 1994. Interactions of cyclic hydrocarbons with biological membranes. *J. Biol. Chem.* **269**:8022–8028.
  32. Sikkema, J., J. A. M. de Bont, and B. Poolman. 1995. Mechanisms of membrane toxicity of hydrocarbons. *Microbiol. Rev.* **59**:201–222.
  33. Sikkema, J., B. Poolman, W. N. Konings, and J. A. M. de Bont. 1992. Effects of the membrane action of tetralin on the functional and structural properties of artificial and bacterial membranes. *J. Bacteriol.* **174**:2986–2992.
  34. Silver, S., and L. Wendt. 1967. Mechanism of action of phenethyl alcohol: breakdown of the cellular permeability barrier. *J. Bacteriol.* **93**:560–566.
  35. Uribe, S., J. Ramirez, and A. Peña. 1985. Effects of  $\beta$ -pinene on yeast membrane functions. *J. Bacteriol.* **161**:1195–1200.
  36. Uribe, S., P. Rangel, G. Espínola, and G. Aguirre. 1990. Effects of cyclohexane, an industrial solvent, on the yeast *Saccharomyces cerevisiae* and on isolated yeast mitochondria. *Appl. Environ. Microbiol.* **56**:2114–2119.
  37. Williams, D. E., L. J. Swango, G. R. Wilt, and S. D. Worley. 1991. Effect of organic *N*-halamines on selected membrane functions in intact *Staphylococcus aureus* cells. *Appl. Environ. Microbiol.* **57**:1121–1127.
  38. Windholz, M., S. Budavari, R. F. Blumetti, and E. S. Otterbein (ed.). 1983. *The Merck index*, 10th ed. Merck & Co., Inc., Rahway, N.J.