



Published in final edited form as:

J Membr Biol. 2006 ; 211(3): 139–150. doi:10.1007/s00232-006-0861-0.

Mechanisms of Acid and Base Secretion by the Airway Epithelium

Horst Fischer and Jonathan H. Widdicombe

Children's Hospital Oakland Research Institute, Oakland CA, and University of California, Davis CA

SUMMARY

One of the main functions of the airway epithelium is to inactivate and remove infectious particles from inhaled air and thereby prevent infection of the distal lung. This function is achieved by mucociliary and cough clearance and by antimicrobial factors present in the airway surface liquid (ASL). There are indications that airway defenses are affected by the pH of the ASL and historically, acidification of the airway surfaces has been suggested as a measure of airway disease. However, even in health, the ASL is slightly acidic, and this acidity might be part of normal airway defense. Only recently research has focused on the mechanisms responsible for acid and base secretion into the ASL. Advances resulted from research into the airway disease associated with cystic fibrosis (CF) after it was found that the CFTR Cl⁻ channel conducts HCO₃⁻ and, therefore, may contribute to ASL pH. However, the acidity of the ASL indicated parallel mechanisms for H⁺ secretion. Recent investigations identified several H⁺ transporters in the apical membrane of the airway epithelium. These include H⁺ channels and ATP-driven H⁺ pumps, including a non-gastric isoform of the H⁺-K⁺ ATPase and a vacuolar-type H⁺ ATPase. Current knowledge of acid and base transporters and their potential roles in airway mucosal pH regulation is reviewed here.

AIRWAY EPITHELIUM AND AIRWAY SURFACE LIQUID

In the human lung, there are 23 generations of dichotomously branching tubes. From generations 0 (trachea) to between 8 and 11 (depending on location within the lung), the airways contain both cartilage and mucous glands, and are termed bronchi. From about generation 10 to 16 the airways lack cartilage, glands and alveoli, and are known as conducting bronchioles. From generation 17 on, the airways become respiratory bronchioles with increasing numbers of alveoli budding off them. By generation 20, the entire wall of the airway is occupied by alveolar openings and the airway has become an “alveolar duct”, which will end in an alveolar sac at generation 23 [97].

In the human trachea, the epithelium is pseudostratified and about 50 μm in height. Numerically its three main cell types are ciliated, basal and goblet (i.e. mucus-secreting). With increasing airway generations, the height of the epithelium decreases progressively until it reaches about ~10 μm in the respiratory bronchioles, by which point it has assumed a columnar morphology. Not only does the form of the epithelium change along the airways, but so also does its cellular composition. Though ciliated cells account for ~50% of the cells in the epithelia of all airway generations [42], basal cells are lost completely before the respiratory bronchioles are reached. Similarly, the number of goblet cells declines progressively to reach zero by the terminal bronchioles (i.e. the smallest bronchioles lacking alveoli). After this, in the respiratory bronchioles, they are replaced by Clara cells [9]. This switch from goblet to Clara cells may reflect the fact that mucus is primarily a defense against particles, and these are deposited predominantly in the larger airways. Clara cells, by contrast, defend mainly against toxic vapors

[75]. Occasional within the surface epithelium of human bronchioles are serous cells [81], specialized for water secretion [92].

On the apical surface of the epithelium is a film of liquid, varying from 5 to 40 μm in depth [101], corresponding to volumes of 0.5 to 4 μl per cm^2 of surface. Visual inspection of the surface of monkey nasal mucosa led Lucas and Douglas [65] to propose that this airway surface liquid (ASL) consisted of two layers: a watery sol between the cilia and a mucous blanket over their tips. They proposed that the low viscosity of the periciliary sol allowed the cilia to beat, and that during their effective strokes the tips of the cilia contacted the underside of the mucous blanket, propelling it, together with entrapped particles, to the mouth, where it was swallowed or expectorated. This process of “mucociliary clearance” thus kept the surface of the airways clean [85,96]. Subsequently, many ultrastructural studies have confirmed the presence of two layers of ASL [101]. In most cases, the depth of the putative sol is approximately the same as the length of the cilia.

ASL is about 96% water, 1% salts, 1% lipids, 1% proteins and 1% mucus [7] and leukocytes are also present. Both ASL composition and volume are considered critical for its function [16,54,88,93,99,100]. How does pH affect this mixture, and thus, its function? Firstly, pH-dependent changes in transepithelial salt and water movement by way of regulation of ion conductances [4,13,79] could alter the concentrations of all components. Secondly, the function of several components is pH-sensitive. Mucus, produced mainly by glands in the larger airways, and by goblet cells in the smaller, contains acidic groups that are involved in the interactions between individual molecules. Its properties are well known to be pH-sensitive. A number of proteins involved in antimicrobial defense are secreted by airway surface epithelium and glands [33] whose functions are potentially sensitive to pH. For example, the function of the antibacterial lactoperoxidase secreted by airway glands has a pH optimum at a slightly acidic pH [14,17,25]. In a recent review, Ricciardolo et al [80] examined the role of excessive acidity on airway function, including 1) effects of H^+ on sensory neurons and associated bronchoconstriction, 2) induction of cough by acid, 3) microvascular leakage and mucus production caused by excessive acidity, 4) effects on leukocyte function, and 5) interaction with nitric oxide and superoxide metabolism of the airways (reviewed in [80]). Thus, the regulation of the pH of the ASL is critical for its normal function.

pH OF THE AIRWAY SURFACE LIQUID

Normal mucosal airway pH

Normal ASL pH is considered acidic compared to a blood and interstitial pH of 7.4. There are indications that in airway diseases ASL pH is comparably alkaline (e.g. chronic bronchitis) or more acidic (e.g. CF or pneumonia) suggesting breakdowns in the pH regulatory mechanisms. Table 1 summarizes studies performed over the past 70 years in which ASL pH was determined using different techniques. The commonest means of measuring ASL pH has been by direct placement of a pH electrode on the nasal mucosa. pH electrodes of various sizes and materials have been used, but measurements are confounded by electrode design, immersion depth, placement of the reference electrode, and possibly mechanical disturbance of the epithelial layer. Measurements have also been made on expectorated sputum or in situ with the use of pH-sensing fluorescent probes. Astoundingly, despite improvements in the measurement techniques over the years, reported pH values measured on the airway surface are quite consistent and repeatable averaging ~ 6.6 in healthy airways (Table 1).

In 1941, Fabricant [29] placed a pH electrode in the nostril of healthy subjects to contact the mucosa directly and found a pH range of 5.5 to 6.5. Because in this study the external reference electrode was placed on the skin of the subject, the acidity of these measurements was possibly overestimated (owing to the higher negative skin potential compared to the nasal potential

[3]). Nevertheless, this study established early on the acidity of the normal nasal mucosa. Using a similar electrode placement, England et al. [28] recently measured the same pH range for the nasal mucosa using an antimony pH probe. Because of the confounding effect of an externally placed reference electrode, McShane et al. [66] used a glass pH electrode with an internal reference electrode which was placed together with the pH sensor on the nasal mucosa. Normal nasal pH was 6.6, and the pH was similar along the respiratory mucosa down to the 3rd generation bronchus. Using an endoscopically guided pH electrode Steinmann [90] measured endobronchial pH values of 5.7 to 6.6. In a more extensive study, where many segments of the bronchial tree were compared, Bodem et al. [8] found an average endobronchial pH of 6.6 and, again, no difference was found between various distal bronchial segments. A number of additional studies found similar tracheal and bronchial mucosal pH values. Although there is some variability, which may in part be due to technical differences between studies, these measurements show consistently that the normal mucosal pH of the airways is quite acidic averaging pH 6.6 for all investigations listed in Table 1. Interestingly, one study grouped measurements of nasal pH by ethnicity [52] and found that the nasal pH in people of African descent was significantly more acidic (pH 6.4) than in Caucasians (pH 6.9). The potentially confounding factor of ethnicity had not been considered in earlier studies.

The relative contribution of different resident or mobile pH buffers has not been well investigated but both mucus and HCO_3^- are likely significant contributors to ASL pH and its buffer capacity. Healthy airways have very little mucus and therefore its role in ASL pH has not been analyzed. In contrast, inflamed airways consistently secrete large amounts of mucus. Airway mucins are acidic and show low isoelectric point values [82] and thus may contribute to low pH buffering. Holma et al. [46,47] used induced sputum from smokers to determine the contribution of airway mucus to ASL pH. Sputum reflects in large parts mucous secretions from the upper airways with variable contamination of saliva. In an initial study average sputum pH was 7.25 and the buffer capacity, usually expressed as the amount of acid needed to achieve a change of one pH unit, was ~ 7 mM/pH in the pH range from pH 5 to 6.5 [46] and increased to ~ 15 mM/pH at increasing pH values up to 7.5. (calculated from Table 1 in ref. [46]). In a second study by these investigators it was found that the average sputum pH (6.82) and buffer capacity (6 mM/pH) was somewhat lower and declined with increasing pH to 2.5 mM/pH at pH 7.5 (calculated from Table 1 in ref. [47]). These two studies established the buffer capacity of human airway mucus using measurements of sputum pH. The differences between these two studies are consistent with differences in HCO_3^- levels. Specifically, the increased baseline pH and increased buffer capacity at high pH values in the initial study [46] is consistent with elevated levels of HCO_3^- or additional buffers. More recent estimates of buffer capacity are similar to Holma's. The buffer capacity of ASL of bovine tracheal cultures was 14 mM/pH at pH 7 [55]. In human gland secretions induced by pilocarpine the buffer capacity was found to be 12 mM/pH at pH 7. This value was determined in the presence of HCO_3^- . When the buffer capacity of pig gland secretions was determined in the absence of HCO_3^- it was found to be substantially lower, 3.7 mM/pH, [88] indicating HCO_3^- as a substantial contributor to the buffer capacity of airway gland secretions.

Mucosal airway pH in disease

As early as the 1930s, nasal secretions were sampled and their pH determined using indicator dyes. In one early study, nasal secretions were found to be acidic during acute inflammation [69], while pH alkalinized over the course of the common cold to pH 7.2 to 8.3 [45]. This resulted in the proposition that the nasal pH could be used as an indicator for airway disease [29]. More recent measurements also reveal changes in ASL pH in disease (Table 1). Induced mucoid or purulent sputum from chronic bronchitis patients were found to be alkaline (7.6 and 7.8, respectively [2]), tracheobronchial mucus samples suctioned from lungs of acutely ill patients showed an average pH of 7.7 [68], endobronchial pH in pneumonia averaged 7.2 –

7.4 [90], and nasal pH from rhinitis patients measured by placing an electrode onto the nasal mucosa resulted in pH values of 7.2 – 8.3 [28]. These studies indicated quite consistently that pH of inflamed or infected airways is alkaline supporting the notion that airway infection activates a mechanism that alkalizes the mucosal pH. In contrast, one study of endobronchial pH during active bacterial infections resulted in quite acidic values (5.6 – 6.2, ref. [40]). Also measurements on CF primary cell cultures using a capillary sampling technique resulted in ASL pH values of 6.0 [16], while in nasal pH recordings in CF patients, McShane et al. [66] found no increased acidity compared to normal. The wide variation of ASL pH in disease supports the notion that the activity of epithelial acid and base transporters is differentially affected in disease.

Recently, a number of cellular mechanisms of acid and base transport have been identified in the airway epithelium and apical localization suggests a direct role in ASL pH regulation. Although their relative importance in health and disease are as yet uncertain, the measured changes in ASL pH likely reflects a selective regulation of specific transporters. These pH regulating transport mechanisms are discussed individually in the next section.

MECHANISMS OF BASE SECRETION

CFTR

The cAMP-activated Cl⁻ channel CFTR (cystic fibrosis transmembrane conductance regulator) is expressed in the apical membrane of ciliated surface cells and gland ducts [59]. The permeability of CFTR to HCO₃⁻, the relative acidity of the pH of CF gland secretions, and its apical localization support the notion that CFTR activity alkalizes ASL pH. In a seminal study using normal and CF primary human cultures, Smith and Welsh [86] described cAMP-stimulated HCO₃⁻ transport that was inhibited by the CFTR blocker DPC (diphenylaminecarboxylate) and was absent in CF airways implicating CFTR in HCO₃⁻ secretion. In addition, block of carbonic anhydrase by acetazolamide effectively blocked HCO₃⁻ secretion. These data suggested a model where intracellular carbonic anhydrase generated HCO₃⁻ that was released apically through CFTR.

Conduction of HCO₃⁻ by CFTR has been debated intensely [78]. Initial observations by Gray et al. [38] in pancreatic duct cells showed a small-conductance Cl⁻ channel expressing a HCO₃⁻ to-Cl⁻ selectivity of 0.13. Firm evidence for conduction of HCO₃⁻ across CFTR was presented by Poulsen et al. [76] in a patch clamp study of recombinantly expressed CFTR, and a number of further studies investigating the relative permeability of CFTR to HCO₃⁻ with respect to Cl⁻ resulted in ratios of 0.13 to 0.25 [38,48,49,63,76].

However, the actual rates of Cl⁻ and HCO₃⁻ secretion across CFTR are determined not only by their relative permeabilities but also by the electrochemical driving forces for both anions. Interestingly, under unstimulated conditions driving forces for Cl⁻ movement are small. With an apical membrane potential of airway cells of -25 mV [15,18,102,103], and intracellular [Cl⁻] of 43 mM [103] and ASL [Cl⁻] of 140 mM [56] a net electrochemical driving force for inward Cl⁻ movement of 5 mV can be estimated. By comparison, HCO₃⁻ concentrations in the cytosol (8 mM, [74]) and the ASL (1.5 mM; at pH 6.6 and pCO₂ = 2.2 kPa) result in a large outwardly directed electrochemical driving force for HCO₃⁻ across the apical membrane of -68 mV. Therefore, in presence of an acidic ASL pH, HCO₃⁻ is predicted as the major secreted anion through CFTR despite its relatively low permeability compared to Cl⁻. In support of this notion, it has been shown that primary airway epithelia or the Calu-3 cell line (a human airway cell line with characteristics of submucosal gland serous cells [26,84]) secrete significant amounts of HCO₃⁻ and the relative contribution of Cl⁻ and HCO₃⁻ to CFTR-mediated current is governed by the respective driving forces [23,24,49,60,62,91].

Cl⁻-HCO₃⁻ exchange

The Cl⁻-HCO₃⁻ exchanger (anion exchanger, AE) isoform 2 (AE2) is broadly expressed in epithelia and has been consistently found in human trachea, bronchi and Calu-3 cells using PCR techniques [27,50]. When human primary airway epithelial cultures were investigated using separate perfusion of the mucosal and serosal compartments [74] Cl⁻-HCO₃⁻ exchange activity was found to be present in the basolateral membrane, whereas HCO₃⁻ release across the apical membrane was well explained by CFTR activity, based on ion replacement studies, block by DIDS, and use of CF epithelia [23,74]. Basolateral localization of AE2 was found in the Calu-3 cell line and rat tracheal submucosal glands, and ciliated surface cells (showing less expression) using immunocytochemistry and sensitivity to serosal DIDS and DNDS in Ussing chambers [64]. Interestingly, basolateral localization of AE2 is expected to result in an acidification of the ASL. The direction of transport by AE2 is determined by the respective gradients for Cl⁻ and HCO₃⁻. As long as the blood-to-cell directed Cl⁻ gradient is larger than the cell-to-blood directed HCO₃⁻ gradient, HCO₃⁻ is moved out of the cell and, thus, AE2 activity tends to acidify the cytosol. This reduces the electrochemical driving force for HCO₃⁻ exit across apical CFTR. Thus a basolaterally localized AE may reduce apical HCO₃⁻ release mediated by CFTR and tends to acidify the ASL.

In addition, Cl⁻-HCO₃⁻ exchange activity based on DRA (downregulated in adenoma [67]), has been found in mouse trachea and CFTR-corrected CFT1 airway cells using Northern hybridization [98]. DRA is widely expressed in intestinal epithelia [95] and apical localization of DRA has been shown by immunocytochemistry in mouse pancreas [39]. In analogy to its function in HCO₃⁻ secretion by the pancreatic duct epithelium [87], a model has been proposed by Wheat et al. [98] for the airway epithelium in which Cl⁻-HCO₃⁻ exchange activity governed by DRA in the apical membrane secretes HCO₃⁻ into the ASL with Cl⁻ recycling through CFTR [98]. However, that study was done under conditions that did not allow distinction of apical from basolaterally located transporters, and studies in polarized airway epithelia have not found Cl⁻-HCO₃⁻ exchange activity in the apical membrane [74]. Therefore, at present it appears that the function of Cl⁻-HCO₃⁻ exchange in airway epithelia is not base secretion into the ASL, but is instead the maintenance of intracellular pH during acute cellular alkalosis [74].

Na⁺-HCO₃⁻ cotransport

In the Calu-3 cell line, the Na⁺-HCO₃⁻ cotransporter (NBC) isoform 1 (NBC1) has been detected by RT-PCR [50]. These cells secrete HCO₃⁻ and contain high levels of apical membrane CFTR, and a basolaterally localized NBC has been identified using the sided effects of anion replacements and small-molecule blockers in Ussing chambers [24]. Thus, movement of HCO₃⁻ into Calu-3 cells on NBC is followed by efflux through apical CFTR [24]. In human primary airway cultures, NBC activity was identified by blocking HCO₃⁻-dependent transepithelial currents with high concentrations of DNDS [23]. The same study introduced the notion that basolateral NBC activity coupled with an acetazolamide-sensitive carbonic anhydrase each provide fractions of the apically secreted HCO₃⁻. This extended the initial model of HCO₃⁻ secretion by Smith and Welsh [86] by adding basolateral NBC as an additional source of intracellular HCO₃⁻. A role for carbonic anhydrase activity as a source of HCO₃⁻ was later also suggested by Krouse et al [60]. Similarly, in isolated pig bronchi [6] both NBC and carbonic anhydrase were determined to be HCO₃⁻ sources supporting HCO₃⁻ secretion into the lumen via CFTR. Obviously, when carbonic anhydrase generates intracellular HCO₃⁻ it generates equal amounts of intracellular H⁺, and the resulting effects on ASL pH will therefore be determined by the localizations (apical vs. basolateral) and rates of turn-over of both HCO₃⁻ and H⁺ transporters.

Paracellular pathway

The airway epithelium is moderately tight, the paracellular pathway expresses a relatively high permeability, and ion conduction across the paracellular pathway supports secretion or absorption of approximately isotonic salt solution [10]. With a transepithelial potential of -32 mV (lumen negative) in normal human trachea *in vivo* [57] and a serosal pH of 7.4, HCO_3^- can be calculated to be in equilibrium across the epithelium at an ASL pH of 6.85, i.e., at more acidic ASL pH, there is a net driving force for HCO_3^- flux from the serosa into the ASL across the paracellular pathway. A paracellular HCO_3^- permeability has been suggested in CF airway epithelial cultures [16] by measuring the effects of transepithelial HCO_3^- gradients on transepithelial potentials. Using Calu-3 epithelia the characteristics of anion permeation across the paracellular pathway was investigated and the paracellular HCO_3^- permeability was found to be quite high (0.94×10^{-6} cm/s, compared to the paracellular Cl^- permeability of 6.28×10^{-6} cm/s, calculated from data in [48]). For this permeability, an average ASL pH of 6.6 in normal airways (i.e., the average of measurements in humans listed in Table 1) can be calculated to drive a HCO_3^- current of $3.2 \mu\text{A}/\text{cm}^2$ across the paracellular pathway (using the Goldman-Hodgkin-Katz equation and $[\text{HCO}_3^-]_{\text{serosal}} = 21$ mM and $[\text{HCO}_3^-]_{\text{mucosal}} = 1.5$ mM at a time-averaged $p\text{CO}_2$ of 2.2 kPa in the airway lumen). In comparison, in forskolin-stimulated primary airway cultures, a transcellular HCO_3^- current of $3.7 \mu\text{A}/\text{cm}^2$ was measured, identified by its sensitivity to acetazolamide and serosal DNDS (4,4'-dinitrostilben-2,2'-disulfonate, a blocker of NBC) [23]. Although the driving forces for HCO_3^- transport are quite different in this comparison it suggests that in presence of an acidic ASL pH a substantial part of transepithelial HCO_3^- secretion may be via the paracellular pathway.

A paracellular HCO_3^- permeability has considerable implications for airway acidification, which is predicted to be limited by the gradient-driven paracellular HCO_3^- flux into an acidic ASL. This is consistent with the finding that CO_2 -induced acidification of ASL in ferret trachea resulted in minute changes of ASL pH which could not be explained by the buffering capacity of the ASL [61]. On the other hand, in similar measurements using bovine tracheal cultures, the ASL was readily acidified with HCl and the pH recovery was largely blocked by DIDS [55] indicating that in these cells the transcellular pathway is the major route for HCO_3^- . Bovine tracheal cultures express very high transepithelial resistances [58], which is expected to limit the paracellular HCO_3^- permeation.

MECHANISMS OF ACID SECRETION

Initially, Smith and Welsh [86] measured HCO_3^- secretion across primary cultures of human tracheal epithelium and predicted that it would contribute to alkalinization of ASL *in vivo*. Then Poulsen et al. [76] showed that CFTR was permeable to HCO_3^- and concluded that CFTR activity might contribute to an alkalinization of ASL pH. However, normal steady-state ASL pH is consistently found to be acidic (Table 1) and mucosal acidification of airway epithelial cultures or native tissues is apparent. When the mucosa of human airway epithelial cultures was mildly alkalinized to pH 7.5, it re-acidifies at initial rates of ~ 0.2 pH/hour [16] which predicts a rate of net acid secretion of $3.5 \text{ nmole}\cdot\text{h}^{-1}\cdot\text{cm}^{-2}$ (assuming an ASL volume of $2.5 \mu\text{l}/\text{cm}^2$ and a buffer capacity of 7 mM/pH). On the other hand, basal rates of epithelial acid secretion by human primary airway cultures in HCO_3^- free conditions are $170 \text{ nmole}\cdot\text{h}^{-1}\cdot\text{cm}^{-2}$ [30] indicating that the airways constitutively secrete both acid and base in parallel at substantial rates. The investigation of the cellular mechanisms involved in acid secretion into the ASL has only recently focused on acid transporters known from other epithelial tissues. Several acid transporters have been identified and localized in airways using both molecular and functional approaches. However, at this point we still lack a concise understanding of the regulation of the acidification of the ASL which necessarily includes the activity of both acid and base transporters. Below we discuss the roles in acid secretion of the major acid transporters that have been identified in airway epithelial cells.

Na⁺-H⁺ exchange

The ubiquitously expressed Na⁺-H⁺ exchanger (NHE) releases H⁺ from cells across the plasma membrane driven by the large electrochemical gradient for Na⁺ entry. Its major function is the maintenance of intracellular pH by protecting the cytosol from acidification. NHE might participate in epithelial acid secretion if expressed in the apical membrane. For example, in the intestinal epithelium NHE activity has been found in the apical brush border membrane where it functions in Na⁺ absorption from and H⁺ secretion into the intestinal lumen [5]. The NHE1 isoform was detected by RT-PCR in all bronchial regions of the human lung [27]. Amiloride-sensitive NHE activity has been identified in isolated primary human airway epithelial cells using intracellular pH measurements [72]. In a study using bilateral perfusion of intact epithelia amiloride-sensitive NHE activity was localized to the basolateral membrane [73], without any measurable apical activity. Basolateral localization of NHE indicates its role in intracellular pH regulation, rather than in H⁺ secretion.

Interestingly, when using mucosal amiloride or its analogs as tools to identify possible apical NHE activity, a number of studies found small transient effects. Acid secretion across intact tracheal tubes isolated from sheep recorded using the pH stat technique was transiently reduced by mucosal EIPA (5-(N-ethyl-N-isopropyl)-amiloride) by 12%, which was interpreted as NHE-mediated acid secretion present in the apical membrane [1]. Similarly, in pH stat recordings of acid secretion by primary human airway epithelia addition of amiloride to the mucosal bath transiently inhibited acid secretion by 34% in a fraction of investigated cultures [30]. In intact distal airways isolated from pigs mucosal acidification was slightly reduced by mucosal dimethylamiloride [51], and using pH-sensitive microelectrodes intracellular pH slightly and transiently acidified when human airway epithelia were treated with mucosal amiloride [102]. In most of these studies, these effects were considered insignificant and not NHE-mediated [30,51,102] and the transient nature of the effects is inconsistent with a block of apical NHE activity. However, these surprisingly similar observations support an alternative notion: acid secretion may have been blocked by amiloride (or its analogs) owing to its effect on apical amiloride-sensitive Na⁺ channels and the concomitant hyperpolarization of the apical membrane potential. This is expected to inhibit H⁺ channels in the apical membrane (which are blocked by hyperpolarization and activated by intracellular acidity, see below) and reduce H⁺ channel-mediated acid secretion. Block of H⁺ channels, in turn, acidifies intracellular pH resulting in the activation of H⁺ channels, and thus, a transient blocker effect on H⁺ secretion. This explanation, although hypothetical, explains the transient effects of mucosal amiloride on H⁺ secretion observed in a number of studies in absence of an apical NHE.

H⁺-K⁺ ATPase

Activity of the gastric parietal cell H⁺-K⁺ ATPase is experimentally readily identified by its sensitivity to highly selective probes, such as SCH28080. In cultured bovine tracheal epithelial cells, Poulsen and Machen [77] found variable block of intracellular alkalization by SCH28080, suggesting the expression of a gastric-type H⁺-K⁺ ATPase in these cells. However the localization was not determined in that study. In other studies, addition of SCH28080 to the mucosal side of polarized human airway cultures [16,30] or intact glands [88] had no effects on acid secretion arguing against a role of the gastric-type H⁺-K⁺ ATPase in acid secretion by airways. By contrast, a non-gastric isoform of the H⁺-K⁺ ATPase, which was initially identified in the apical membrane of rat distal colon epithelium [22], was detected by RT-PCR in cultured bronchial epithelia, and immunocytochemistry localized it to the apical membrane pole [16]. Transport function of the colonic H⁺-K⁺ ATPase is blocked by ouabain, but not by SCH28080. Mucosal ouabain entirely inhibited ASL acidification of bronchial cultures when treated for several hours [16] and in pH stat experiments in Calu-3 cells mucosal ouabain caused effects consistent with block of an apical H⁺-K⁺ ATPase [60]. On the other hand acute effects of

mucosal ouabain on H^+ secretion measured in pH stat experiments were small in tracheo-bronchial cultures [30,83] or in pig tracheal surface epithelium [88].

Vacuolar H^+ ATPase

The vacuolar-type H^+ ATPase (v-ATPase) is an ATP-dependent H^+ pump that is responsible for the acidification of intracellular compartments [31]. In addition v-ATPases have been shown in the plasma membranes of epithelial cells (such as the sweat duct [37]) where v-ATPases pump H^+ across the plasma membrane and acidify the extracellular space. Identification of v-ATPase activity in functional experiments is aided by the highly selective inhibitor bafilomycin A1 [11]. A role of v-ATPase activity in airway acidification is supported by the work of Inglis et al [51] in isolated perfused pig bronchi, which acidify their luminal surface at high rates of $2.3 \mu\text{mole}\cdot\text{h}^{-1}\cdot\text{cm}^2$ [51] of which ~60% is inhibited by addition of luminal bafilomycin A1. In contrast, in a number of other studies, inhibition of acid secretion by bafilomycin A1 showed comparably small effects, including in primary human airway epithelia [30,83], or no measurable effects in cow tracheal cells [77] or pig airway surface epithelia [88].

H^+ channels

Voltage-gated, highly selective H^+ channels were first described in snail neurons and since have been found in many other species and cell types including epithelia [20,83,94]. H^+ channels are typically activated by intracellular acidification, extracellular alkalinization, and/or membrane depolarization. These characteristics result functionally in a strong outward rectification of H^+ currents because H^+ currents are both driven and activated by an outwardly directed H^+ -motive force suggesting a general role for H^+ channels as relief valves that permit effective H^+ release from cells during intracellular acidification [20]. In rat alveolar type II cells, for example, the biophysical characteristics have been studied extensively and localization in the apical membrane has been demonstrated using apical patch clamp recordings [19]. Using intracellular pH measurements in single cells it was found that H^+ channels in alveolar type II cells are active during intracellular acidification [70].

In human primary airway cells or in the JME CF airway cell line [30,83] we identified a whole cell H^+ conductance with typical characteristics of H^+ channels as described in alveolar type II cells, including slow current activation at depolarizing potentials, outward rectification, H^+ selectivity, and block by Zn^{2+} . By using $ZnCl_2$ as a probe for H^+ channel activity, this conductance was localized to the apical membrane using measurements of H^+ secretion by airway epithelia in Ussing chambers. H^+ secretion was blocked 50-70% by $ZnCl_2$ [30,83], and the remaining unblocked fraction was partially inhibited by mucosal bafilomycin A1 (11%) or ouabain (15%, [83]), suggesting that in these studies the primary mechanism was channel-mediated H^+ secretion. Currently it is unclear why different investigations found different acid transport mechanisms as the primary mediator of H^+ secretion in airways, such as the H^+ - K^+ ATPase in primary airway cultures [16], v-ATPase in intact pig bronchi [51] or H^+ channels in primary airway cultures or an airway cell line [30,83]. We surmise that this might be affected by the H^+ gradients across the plasma membrane, which will particularly affect the participation of H^+ channels in H^+ secretion.

ATP-driven H^+ transporters such as the v-ATPase or the H^+ - K^+ ATPase can transport H^+ against a considerable electrochemical gradient. In contrast, H^+ channel-mediated H^+ secretion occurs only in presence of an outwardly directed electrochemical H^+ gradient owing to the outwardly rectifying characteristics of H^+ currents, which preclude inward movement of H^+ under most physiological conditions.

H⁺ channels are evidently present and active in airway cultures, but what is the mechanism responsible for generating an outwardly directed electrochemical H⁺ gradient? For comparison, in professional phagocytes such a transmembrane H⁺ gradient is generated by NADPH oxidase activity. NADPH oxidase is a multicomponent enzyme complex that oxidizes intracellular NADPH to NADP⁺ + H⁺ and the resulting electrons are transferred across the plasma membrane where extracellular O₂ is reduced to superoxide (O₂^{•-}). Thus NADPH oxidase activity both acidifies the cytosol by NADPH oxidation [53] and depolarizes the plasma membrane potential by the outward electron current [21], which creates an outwardly directed H⁺ gradient and activates H⁺ channels in phagocytes. We hypothesized that a similar process was active in the apical membrane of airway epithelial cells and we set out to investigate the expression and function of NADPH oxidase in airway epithelia.

A number of isoforms of the transmembrane, electron-transporting β subunits of NADPH oxidase have been described. The two large homologs DUOX1 and DUOX2 (for dual oxidase; 83% similarity between isoforms) have been recently identified as the major isoforms expressed in human airway epithelial cells [36,43,83] and DUOX1 and/or DUOX2 are present in the apical membrane [83]. NADPH oxidase-mediated superoxide production by airway cells was identified [32,36] and intracellular H⁺ production and secretion was correlated with DUOX1 expression in two airway cell types and inhibited by NADPH oxidase blockers [83]. These observations suggested that the airway epithelium expresses a DUOX-based NADPH oxidase that produces superoxide and H⁺. Thus, in analogy to the phagocytic NADPH oxidase, DUOX activity in the apical membrane of airway cells is a candidate to both depolarize the membrane and acidify the cytosol to drive H⁺ currents across channels. In cultured human airway epithelia the resting apical membrane potential ranges from -19 mV to -29 mV [15, 18,102,103] and the intracellular pH is 7.1, [73,74,102], resulting in an inwardly directed driving force for H⁺, which essentially rules out a contribution of H⁺ channels to H⁺ secretion and likely ATP-driven transporters dominate basal H⁺ secretion. However, H⁺ secretion is greatly increased by mucosal histamine or ATP [30,88], two agonists that elevate intracellular Ca²⁺. Both DUOX isozymes express Ca²⁺-binding EF-hands, and NADPH oxidase activity in airways has been shown to be regulated by the Ca²⁺-elevating agonists ATP, thapsigargin [32], and ionomycin [36]. Further, DUOX1 expression in airways has been shown to be correlated with intracellular H⁺ production [83]. The sum of these observations suggests a role for a DUOX-based NADPH oxidase in apical H⁺ production by the airway epithelium, which might support gradient-driven, channel-mediated H⁺ secretion. The presence of NADPH oxidase activity in the apical membrane tightly links acid and oxidant production by the airway epithelium. As a corollary to its function in leukocytes, there are indications that NADPH oxidase-derived H⁺ and oxidants are part of innate airway defenses [32,36,83].

CONCLUSION

Despite a long history of measurements of airway mucosal pH and its suggested role in airway disease, acid and base transporters expressed by the airway epithelium have been investigated and identified only very recently. Figure 1 summarizes the distribution of currently known transporters to the apical and basolateral poles of airway epithelial cells, and intracellular sources of H⁺ and HCO₃⁻. Parallel mechanisms for the independent secretion of both H⁺ and HCO₃⁻ result in a sensitive way to tightly regulate ASL pH. There are a number of indications that link ASL pH to airway function and airway disease, including roles of pH in innate defense and airway stress. Changes in ASL pH have been found in airway diseases resulting in epithelial stress and increased susceptibility to bacterial airway infections. Currently, the relations between ASL pH, epithelial integrity and function, and antibacterial activity has not been worked out in detail, however, there are indications that a tightly regulated ASL pH is important for normal airway health.

Acknowledgments

Many thanks to Sarah K. Inglis (Dundee, Scotland) and Mauri Krouse (Stanford, California) for reading and commenting on the manuscript. The authors' laboratories are supported by NIH HL071829.

References

1. Acevedo M, Steele LW. Na⁺-H⁺ exchanger in isolated epithelial tracheal cells from sheep. Involvement in tracheal proton secretion. *Exp Physiol* 1993;78:383–94. [PubMed: 8392357]
2. Adler K, Wooten O, Philippoff W, Lerner E, Dulfano M. Physical properties of sputum. 3. Rheologic variability and intrinsic relationships. *Am Rev Respir Dis* 1972;106:86–96. [PubMed: 5036835]
3. Alton EFWF, Currie AD, Logan-Sinclair R, Warner JO, Hodson ME, Geddes DM. Nasal potential difference: a clinical test for cystic fibrosis. *Eur Resp J* 1990;3:922–926.
4. Awayda MS, Boudreaux MJ, Reger RL, Hamm LL. Regulation of the epithelial Na⁺ channel by extracellular acidification. *Am J Physiol Cell Physiol* 2000;279:C1896–1905. [PubMed: 11078705]
5. Bachmann O, Riederer B, Rossmann H, Groos S, Schultheis PJ, Shull GE, Gregor M, Manns MP, Seidler U. The Na⁺/H⁺ exchanger isoform 2 is the predominant NHE isoform in murine colonic crypts and its lack causes NHE3 upregulation. *Am J Physiol Gastrointest Liver Physiol* 2004;287:G125–133. [PubMed: 14962844]
6. Ballard ST, Trout L, Garrison J, Inglis SK. Ionic mechanism of forskolin-induced liquid secretion by porcine bronchi. *Am J Physiol Lung Cell Mol Physiol* 2005;00159–2005.
7. Boat TF, Cheng PW. Biochemistry of airway mucus secretions. *Fed Proc* 1980;39:3067–74. [PubMed: 7428951]
8. Bodem C, Lampton L, Miller D, Tarka E, Everett E. Endobronchial pH. Relevance of aminoglycoside activity in gram-negative bacillary pneumonia. *Am Rev Respir Dis* 1983;127:39–41. [PubMed: 6849547]
9. Boers JE, Ambergen AW, Thunnissen FBJM. Number and Proliferation of Clara Cells in Normal Human Airway Epithelium. *Am J Respir Crit Care Med* 1999;159:1585–1591. [PubMed: 10228131]
10. Boucher RC. Molecular insights into the physiology of the 'thin film' of airway surface liquid. *J Physiol Lond* 1999;516:631–638. [PubMed: 10200413]
11. Bowman EJ, Siebers A, Altendorf K. Bafilomycins: A Class of Inhibitors of Membrane ATPases from Microorganisms, Animal Cells, and Plant Cells. *PNAS* 1988;85:7972–7976. [PubMed: 2973058]
12. Breuninger H. Über das physikalisch-chemische Verhalten des Nasenschleims. *Eur Arch Oto-Rhino-Laryng* 1964;184:133–138.
13. Chalfant ML, Denton JS, Berdiev BK, Ismailov II, Benos DJ, Stanton BA. Intracellular H⁺ regulates the alpha-subunit of ENaC, the epithelial Na⁺ channel. *Am J Physiol Cell Physiol* 1999;276:C477–86.
14. Ciaccio C, De Sanctis G, Marini S, Sinibaldi F, Santucci R, Arcovito A, Bellelli A, Ghibaudi E, Ferrari Rosa P, Coletta M. Proton Linkage for CO Binding and Redox Properties of Bovine Lactoperoxidase. *Biophys J* 2004;86:448–454. [PubMed: 14695287]
15. Clarke LL, Paradiso AM, Boucher RC. Histamine-induced Cl⁻ secretion in human nasal epithelium: responses of apical and basolateral membranes. *Am J Physiol Cell Physiol* 1992;263:C1190–9.
16. Coakley RD, Grubb BR, Paradiso AM, Gatzky JT, Johnson LG, Kreda SM, O'Neal WK, Boucher RC. Abnormal surface liquid pH regulation by cultured cystic fibrosis bronchial epithelium. *Proc Natl Acad Sci USA* 2003;100:16083–16088. [PubMed: 14668433]
17. Conner GE, Salathe M, Forteza R. Lactoperoxidase and Hydrogen Peroxide Metabolism in the Airway. *Am J Respir Crit Care Med* 2002;166:57S–61.
18. Cotton CU, Stutts MJ, Knowles MR, Gatzky JT, Boucher RC. Abnormal apical cell membrane in cystic fibrosis respiratory epithelium. An in vitro electrophysiologic analysis. *J Clin Invest* 1987;79:80–5. [PubMed: 3793933]
19. DeCoursey T, Cherny V. Voltage-activated proton currents in membrane patches of rat alveolar epithelial cells. *J Physiol (Lond)* 1995;489:299–307. [PubMed: 8847626]

20. DeCoursey TE. Voltage-gated proton channels and other proton transfer pathways. *Physiol Rev* 2003;83:475–579. [PubMed: 12663866]
21. DeCoursey TE. During the Respiratory Burst, Do Phagocytes Need Proton Channels or Potassium Channels or Both? *Sci STKE* 2004;2004:pe21. [PubMed: 15150421]
22. Del Castillo JR, Rajendran VM, Binder HJ. Apical membrane localization of ouabain-sensitive K⁺-activated ATPase activities in rat distal colon. *Am J Physiol Cell Physiol* 1991;261:G1005–11.
23. Devor DC, Bridges RJ, Pilewski JM. Pharmacological modulation of ion transport across wild-type and Δ F508 CFTR-expressing human bronchial epithelia. *Am J Physiol Cell Physiol* 2000;279:C461–79. [PubMed: 10913013]
24. Devor DC, Singh AK, Lambert LC, DeLuca A, Frizzell RA, Bridges RJ. Bicarbonate and chloride secretion in Calu-3 human airway epithelial cells. *J Gen Physiol* 1999;113:743–60. [PubMed: 10228185]
25. DiPaola M, Maxfield F. Conformational changes in the receptors for epidermal growth factor and asialoglycoproteins induced by the mildly acidic pH found in endocytic vesicles. *J Biol Chem* 1984;259:9163–9171. [PubMed: 6086623]
26. Dubin RF, Robinson SK, Widdicombe JH. Secretion of lactoferrin and lysozyme by cultures of human airway epithelium. *Am J Physiol Lung Cell Mol Physiol* 2004;286:L750–755. [PubMed: 15003937]
27. Dudeja PK, Hafez N, Tyagi S, Gailey CA, Toofanfard M, Alrefai WA, Nazir TM, Ramaswamy K, Al-Bazzaz FJ. Expression of the Na⁺/H⁺ and Cl⁻/HCO₃⁻ exchanger isoforms in proximal and distal human airways. *Am J Physiol Lung Cell Mol Physiol* 1999;276:L971–978.
28. England R, Homer J, Knight L, Ell S. Nasal pH measurement: a reliable and repeatable parameter. *Clin Otolaryngol Allied Sci* 1999;24:67–8. [PubMed: 10196653]
29. Fabricant ND. Significance of the pH of nasal secretions in situ. *Arch Otolaryngology* 1941;34:150–163.
30. Fischer H, Widdicombe JH, Illek B. Acid secretion and proton conductance in human airway epithelium. *Am J Physiol Cell Physiol* 2002;282:C736–743. [PubMed: 11880261]
31. Forgac M. Structure and properties of the vacuolar H⁺-ATPases. *J Biol Chem* 1999;274:12951–4. [PubMed: 10224039]
32. Forteza R, Salathe M, Miot F, Forteza R, Conner GE. Regulated Hydrogen Peroxide Production by Duox in Human Airway Epithelial Cells. *Am J Respir Cell Mol Biol* 2005;32:462–469. [PubMed: 15677770]
33. Ganz T. Antimicrobial polypeptides in host defense of the respiratory tract. *J Clin Invest* 2002;109:693–697. [PubMed: 11901174]
34. Gatto L. pH of mucus in rat trachea. *J Appl Physiol* 1981;50:1224–1226. [PubMed: 7263382]
35. Gatto L. pH of mucus in rabbit trachea: cholinergic stimulation and block. *Lung* 1985;163:109–15. [PubMed: 3927070]
36. Geiszt M, Witta J, Baffi J, Lekstrom K, Leto TL. Dual oxidases represent novel hydrogen peroxide sources supporting mucosal surface host defense. *Faseb J* 2003;17:1502–1504. [PubMed: 12824283]
37. Granger D, Marsolais M, Burry J, Laprade R. V-type H⁺-ATPase in the human eccrine sweat duct: immunolocalization and functional demonstration. *Am J Physiol Cell Physiol* 2002;282:C1454–1460. [PubMed: 11997260]
38. Gray MA, Pollard CE, Harris A, Coleman L, Greenwell JR, Argent BE. Anion selectivity and block of the small-conductance chloride channel on pancreatic duct cells. *Am J Physiol Cell Physiol* 1990;259:C752–761.
39. Greeley T, Shumaker H, Wang Z, Schweinfest CW, Soleimani M. Downregulated in adenoma and putative anion transporter are regulated by CFTR in cultured pancreatic duct cells. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G1301–1308. [PubMed: 11668039]
40. Guerrin F, Voisin C, Macquet V, Robin H, Lequien P. Apport de la pH metrie bronchique in situ. *Progr Respir Res* 1971;6:372–383.
41. Guerrin, F.; Voisin, C.; Macquet, V.; Robin, H.; Wattel, F.; Boulanger, J. Hypersecretion Bronchique. *Clinchy: Poincot; 1969. Possibilities de la pH metrie bronchique in situ; p. 249-256.*

42. Harkema, JR.; Mariassy, A.; St. George, JA.; Hyde, D.; Plopper, CG. Epithelial cells of the conducting airways. A species comparison. In: Farmer, SG.; Hay, DWP., editors. *The Airway Epithelium: Physiology, Pathology, and Pharmacology*. Marcel Dekker, Inc; New York: 1991. p. 3-39.
43. Harper RW, Xu C, Eiserich J, Chen Y, Kao C-Y, Thai P, Setiadi H, Wu R. Differential regulation of dual NADPH oxidases/peroxidases, Duox1 and Duox2, by Th1 and Th2 cytokines in respiratory tract epithelium. *Febs Lett* 2005;579:4911–4917. [PubMed: 16111680]
44. Hehar S, Mason J, Stephen A, Washington N, Jones N, Jackson S, Bush D. Twenty-four hour ambulatory nasal pH monitoring. *Clin Otolaryngol Allied Sci* 1999;24:24–5. [PubMed: 10196643]
45. Hilding A. The common cold. *Arch Otolaryng* 1930;12:133.
46. Holma B. Influence of buffer capacity and pH-dependent rheological properties of respiratory mucus on health effects due to acidic pollution. *Sci Total Environ* 1985;41:101–23. [PubMed: 2984768]
47. Holma B, Hegg PO. pH and protein-dependent buffer capacity and viscosity of respiratory mucus: their interrelationships and influence on health. *Sci Total Environ* 1989;84:71–82. [PubMed: 2772626]
48. Illek B, Tam AW-K, Fischer H, Machen TE. Anion selectivity of apical membrane conductance of Calu 3 human airway epithelia. *Pflugers Arch* 1999;437:812–822. [PubMed: 10370058]
49. Illek B, Yankaskas JR, Machen TE. cAMP and genistein stimulate HCO₃⁻ conductance through CFTR in human airway epithelia. *Am J Physiol Lung Cell Mol Physiol* 1997;272:L752–61.
50. Inglis SK, Finlay L, Ramminger SJ, Richard K, Ward MR, Wilson SM, Olver RE. Regulation of intracellular pH in Calu-3 human airway cells. *J Physiol (Lond)* 2002;538:527–539. [PubMed: 11790817]
51. Inglis SK, Wilson SM, Olver RE. Secretion of acid and base equivalents by intact distal airways. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L855–862. [PubMed: 12676770]
52. Ireson NJ, Tait JS, MacGregor GA, Baker EH. Comparison of nasal pH values in black and white individuals with normal and high blood pressure. *Clin Sci* 2001;100:327–333. [PubMed: 11222120]
53. Jankowski A, Grinstein S. Modulation of the cytosolic and phagosomal pH by the NADPH oxidase. *Antioxid Redox Signal* 2002;4:61–8. [PubMed: 11970844]
54. Jayaraman S, Song Y, Verkman AS. Airway surface liquid osmolality measured using fluorophore-encapsulated liposomes. *J Gen Physiol* 2001;117:423–30. [PubMed: 11331352]
55. Jayaraman S, Song Y, Verkman AS. Airway surface liquid pH in well-differentiated airway epithelial cell cultures and mouse trachea. *Am J Physiol Cell Physiol* 2001;281:C1504–1511. [PubMed: 11600413]
56. Jayaraman S, Song Y, Vetrivel L, Shankar L, Verkman AS. Noninvasive in vivo fluorescence measurement of airway-surface liquid depth, salt concentration, and pH. *J Clin Invest* 2001;107:317–324. [PubMed: 11160155]
57. Knowles MR, Buntin WH, Bromberg PA, Gatzky JT, Boucher RC. Measurements of transepithelial electric potential differences in the trachea and bronchi of human subjects in vivo. *Am Rev Respir Dis* 1982;126:108–12. [PubMed: 7091895]
58. Kondo M, Finkbeiner WE, Widdicombe JH. Cultures of bovine tracheal epithelium with differentiated ultrastructure and ion transport. *In Vitro Cell Dev Biol* 1992;29A:19–24. [PubMed: 8444742]
59. Kreda SM, Mall M, Mengos A, Rochelle L, Yankaskas J, Riordan JR, Boucher RC. Characterization of wild-type and $\Delta F508$ cystic fibrosis transmembrane regulator in human respiratory epithelia. *Mol Biol Cell* 2005;16:2154–2167. [PubMed: 15716351]
60. Krouse ME, Talbott JF, Lee MM, Joo NS, Wine JJ. Acid and base secretion in the Calu-3 model of human serous cells. *Am J Physiol Lung Cell Mol Physiol* 2004;287:L1274–1283. [PubMed: 15310554]
61. Kyle H, Ward JP, Widdicombe JG. Control of pH of airway surface liquid of the ferret trachea in vitro. *J Appl Physiol* 1990;68:135–40. [PubMed: 2107164]
62. Lee MC, Penland CM, Widdicombe JH, Wine JJ. Evidence that Calu-3 human airway cells secrete bicarbonate. *Am J Physiol Lung Cell Mol Physiol* 1998;274:L450–3.
63. Linsdell P, Tabcharani JA, Rommens JM, Hou YX, Chang XB, Tsui LC, Riordan JR, Hanrahan JW. Permeability of wild-type and mutant cystic fibrosis transmembrane conductance regulator chloride channels to polyatomic anions. *J Gen Physiol* 1997;110:355–64. [PubMed: 9379168]

64. Loffing J, Moyer BD, Reynolds D, Shmukler BE, Alper SL, Stanton BA. Functional and molecular characterization of an anion exchanger in airway serous epithelial cells. *Am J Physiol Cell Physiol* 2000;279:C1016–1023. [PubMed: 11003582]
65. Lucas A, Douglas L. Principles underlying ciliary activity in the respiratory tract. II. A comparison of nasal clearance in man, monkey and other mammals. *Arch Otolaryngol* 1934;20:518–541.
66. McShane D, Davies JC, Davies MG, Bush A, Geddes DM, Alton EFWF. Airway surface pH in subjects with cystic fibrosis. *Eur Respir J* 2003;21:37–42. [PubMed: 12570106]
67. Melvin JE, Park K, Richardson L, Schultheis PJ, Shull GE. Mouse down-regulated in adenoma (DRA) Is an Intestinal $\text{Cl}^-/\text{HCO}_3^-$ exchanger and is up-regulated in colon of mice lacking the NHE3 Na^+/H^+ exchanger. *J Biol Chem* 1999;274:22855–22861. [PubMed: 10428871]
68. Metheny N, Stewart B, Smith L, Yan H, Diebold M, Clouse R. pH and concentration of bilirubin in feeding tube aspirates as predictors of tube placement. *Nurs Res* 1999;48:189–97. [PubMed: 10414681]
69. Mittermaier R. Untersuchungen über die Wasserstoffionenkonzentration an Sekreten und Schleimhäuten, im besonderen bei chronischen Nebenhöhlenerkrankungen. *Eur Arch Oto-Rhino-Laryngol* 1930;127:149–172.
70. Murphy R, Cherny VV, Morgan D, DeCoursey TE. Voltage-gated proton channels help regulate pHi in rat alveolar epithelium. *Am J Physiol Lung Cell Mol Physiol* 2005;288:L398–408. [PubMed: 15516489]
71. Palmer L, Merrill W, Niederman M, Ferranti R, Reynolds H. Bacterial adherence to respiratory tract cells. Relationships between in vivo and in vitro pH and bacterial attachment. *Am Rev Respir Dis* 1986;133:784–8. [PubMed: 3706886]
72. Paradiso AM. Identification of Na^+/H^+ exchange in human normal and cystic fibrotic ciliated airway epithelium. *Am J Physiol Lung Cell Mol Physiol* 1992;262:L757–64.
73. Paradiso AM. ATP-activated basolateral Na^+/H^+ exchange in human normal and cystic fibrosis airway epithelium. *Am J Physiol Lung Cell Mol Physiol* 1997;273:L148–58.
74. Paradiso AM, Coakley RD, Boucher RC. Polarized distribution of HCO_3^- -transport in human normal and cystic fibrosis nasal epithelia. *J Physiol (Lond)* 2003;548:203–218. [PubMed: 12562898]
75. Plopper, CG.; Hyde, DM.; Buckpitt, AR. Clara cells. In: Crystal, RG.; West, JB.; Weibel, ER.; Barnes, PJ., editors. *The Lung: Scientific Foundations*. Lippincott-Raven; Philadelphia: 1997. p. 517-534.
76. Poulsen JH, Fischer H, Illek B, Machen TE. Bicarbonate conductance and pH regulatory capability of cystic fibrosis transmembrane conductance regulator. *Proc Natl Acad Sci USA* 1994;91:5340–4. [PubMed: 7515498]
77. Poulsen JH, Machen TE. HCO_3^- -dependent pHi regulation in tracheal epithelial cells. *Pflugers Arch* 1996;432:546–54. [PubMed: 8766016]
78. Quinton PM. The neglected ion: HCO_3^- . *Nat Med* 2001;7:292–3. [PubMed: 11231624]
79. Reddy MM, Kopito RR, Quinton PM. Cytosolic pH regulates GCl through control of phosphorylation states of CFTR. *Am J Physiol Cell Physiol* 1998;275:C1040–7.
80. Ricciardolo FLM, Gaston B, Hunt J. Acid stress in the pathology of asthma. *J Allergy Clin Immunol* 2004;113:610–619. [PubMed: 15100663]
81. Rogers AV, Dewar A, Corrin B, Jeffery PK. Identification of serous-like cells in the surface epithelium of human bronchioles. *Eur Respir J* 1993;6:498–504. [PubMed: 8491299]
82. Rose MC. Mucins: structure, function, and role in pulmonary diseases. *Am J Physiol Lung Cell Mol Physiol* 1992;263:L413–429.
83. Schwarzer C, Machen TE, Illek B, Fischer H. NADPH oxidase-dependent acid production in airway epithelial cells. *J Biol Chem* 2004;279:36454–36461. [PubMed: 15210697]
84. Shen BQ, Finkbeiner WE, Wine JJ, Mrsny RJ, Widdicombe JH. Calu-3: a human airway epithelial cell line that shows cAMP-dependent Cl^- secretion. *Am J Physiol Lung Cell Mol Physiol* 1994;266:L493–501.
85. Sleight MA, Blake JR, Liron N. The propulsion of mucus by cilia. *Am Rev Respir Dis* 1988;137:726–41. [PubMed: 3278666]
86. Smith JJ, Welsh MJ. cAMP stimulates bicarbonate secretion across normal, but not cystic fibrosis airway epithelia. *J Clin Invest* 1992;89:1148–53. [PubMed: 1313448]

87. Sohma Y, Gray M, Imai Y, Argent B. 150 mM HCO_3^- - how does the pancreas do it? Clues from computer modelling of the duct cell. *J Pancreas (Online)* 2001;2:198–202.
88. Song Y, Salinas D, Nielson DW, Verkman AS. Hyperacidity of secreted fluid from submucosal glands in early cystic fibrosis. *Am J Physiol Cell Physiol* 2005;00379–2005.
89. Song Y, Thiagarajah J, Verkman AS. Sodium and Chloride Concentrations, pH, and Depth of Airway Surface Liquid in Distal Airways. *J Gen Physiol* 2003;122:511–519. [PubMed: 14557401]
90. Steinmann E. La secretion bronchique et le pH. *Bronches* 1956;6:126–129.
91. Tamada T, Hug MJ, Frizzell RA, Bridges RJ. Microelectrode and impedance analysis of anion secretion in Calu-3 cells. *J Pancreas (Online)* 2001;2:219–228.
92. Tandler B. Ultrastructure of the human submaxillary gland. I. Architecture and histological relationships of the secretory cells. *Am J Anat* 1962;111:287–307. [PubMed: 13984839]
93. Tarran R, Grubb BR, Gatzky JT, Davis CW, Boucher RC. The Relative Roles of Passive Surface Forces and Active Ion Transport in the Modulation of Airway Surface Liquid Volume and Composition. *J Gen Physiol* 2001;118:223–236. [PubMed: 11479349]
94. Thomas RC, Meech RW. Hydrogen ion currents and intracellular pH in depolarized voltage-clamped snail neurones. *Nature* 1982;299:826–8. [PubMed: 7133121]
95. Wang Z, Petrovic S, Mann E, Soleimani M. Identification of an apical $\text{Cl}^-/\text{HCO}_3^-$ exchanger in the small intestine. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G573–579. [PubMed: 11842009]
96. Wanner A, Salathe M, O’Riordan TG. Mucociliary clearance in the airways. *Am J Respir Crit Care Med* 1996;154:1868–1902. [PubMed: 8970383]
97. Weibel, ER. *Morphometry of the Human Lung*. Heidelberg: Springer-Verlag; 1963.
98. Wheat VJ, Shumaker H, Burnham C, Shull GE, Yankaskas JR, Soleimani M. CFTR induces the expression of DRA along with $\text{Cl}^-/\text{HCO}_3^-$ exchange activity in tracheal epithelial cells. *Am J Physiol Cell Physiol* 2000;279:C62–71. [PubMed: 10898717]
99. Widdicombe JG. Relationships among the composition of mucus, epithelial lining liquid, and adhesion of microorganisms. *Am J Respir Crit Care Med* 1995;151:2088–2092. [PubMed: 7767562]
100. Widdicombe JH, Bastacky SJ, Wu DX, Lee CY. Regulation of depth and composition of airway surface liquid. *Eur Respir J* 1997;10:2892–7. [PubMed: 9493680]
101. Widdicombe JH, Widdicombe JG. Regulation of human airway surface liquid. *Respir Physiol* 1995;99:3–12. [PubMed: 7740210]
102. Willumsen NJ, Boucher RC. Intracellular pH and its relationship to regulation of ion transport in normal and cystic fibrosis human nasal epithelia. *J Physiol Lond* 1992;455:247–269. [PubMed: 1336551]
103. Willumsen NJ, Davis CW, Boucher RC. Intracellular Cl^- activity and cellular Cl^- pathways in cultured human airway epithelium. *Am J Physiol Cell Physiol* 1989;256:C1033–C1044.
104. Zippel R, Meyer P, Schubel F. Eine Methode zur Messung der Wasserstoffionenkonzentration und ihre Ergebnisse an Schleimhautoberflächen in Nase und Mundrachen. *Eur Arch Oto-Rhino-Laryng* 1965;186:115–127.




Figure 1. Major acid and base transporters of airway epithelium

HCO_3^- is generated both intracellularly by carbonic anhydrase (CA) activity and enters over the basolateral membrane by way of NBC1, whose Na^+ -to- HCO_3^- stoichiometry may be 1 : 2 or 1 : 3 [60,91]. HCO_3^- is released apically via CFTR and basolaterally via an anion exchanger. Intracellular H^+ is produced by CA and an apical DUOX-based NADPH oxidase. Three apical H^+ secreting mechanisms have been identified: H^+ channels, v-ATPase, and $\text{H}^+\text{-K}^+$ ATPase. The role of the basolateral NHE1 is likely to maintain intracellular pH. For abbreviations see text.

Table 1

pH of the airway mucosa

	pH	Method	Citation
Human			
Nasal			
	5.5 - 6.5	pH electrode	[29]
	6.6	pH electrode	[44]
	6.44 / 6.91	pH electrode, measurements grouped by ethnicity (African descent vs. Caucasians)	[52]
	5.5 - 6.5	pH electrode	[28]
	6.6	pH electrode	[66]
	6.4 - 7.9	pH electrode	[12]
	6.4 - 6.5	pH electrode	[104]
Tracheal			
	6.1 - 7.9	pH electrode	[40]
	6.71	micropipette sampling	[41]
Bronchial			
	5.7 - 6.6	pH electrode	[90]
	7 - 7.5	pH electrode	[40]
	6.7	pH electrode	[71]
	7.1	pH electrode	[66]
	6.25	micropipette sampling, primary cell cultures	[16]
	6.78	BCECF dextran	[56]
	6.58	pH electrode	[8]
Human, disease			
Smokers			
	7.25	Sputum	[46]
	6.82	Sputum	[47]
Chronic bronchitis			
	7.59	Sputum, mucoid	[2]
	7.83	Sputum, purulent	[2]
Rhinitis			
	7.2 - 8.3	pH electrode, nasal	[28]
Common cold			
	7.2 - 8.3	pH indicators	[45]
Cystic fibrosis			
	6	micropipette, primary cell cultures	[16]
	6.8	pH electrode, nasal	[66]
Bacterially infected			
	6.48	pH electrode, bronchial	[8]
	5.6 - 6.2	pH electrode, bronchial	[40]
	7.2 - 7.4	pH electrode, bronchial	[90]
Non-human			
Rabbit			

	pH	Method	Citation
	6.73	pH electrode	[40]
	7.7	pH electrode	[35]
Rat			
	7.52	pH electrode	[34]
Ferret			
	6.85	pH electrode	[61]
Mouse			
	7.28	BCECF dextran, isolated lungs	[89]
	7.14	BCECF dextran	[55]
	6.95	BCECF dextran, in situ	[54]
Pig			
	6.93	pH of bronchial perfusate	[51]
Cow			
	6.81	BCECF dextran, primary cell cultures	[55]
	6.94	BCECF dextran, primary cell cultures	[54]
Dog			
	7.09	pH electrode, bronchial	[8]

"pH electrode" refers to measurements where an electrode was placed on the airway mucosa.