



# Mechanisms Underlying the Anti-inflammatory and Immunosuppressive Activity of Ruxolitinib

Elena Maria Elli<sup>1\*</sup>, Claudia Baratè<sup>2</sup>, Francesco Mendicino<sup>3</sup>, Francesca Palandri<sup>4</sup> and Giuseppe Alberto Palumbo<sup>5\*</sup>

<sup>1</sup> Hematology Division and Bone Marrow Transplant Unit, Ospedale San Gerardo, ASST Monza, Monza, Italy, <sup>2</sup> Department of Clinical and Experimental Medicine, Section of Hematology, University of Pisa, Pisa, Italy, <sup>3</sup> Hematology Unit, Department of Hemato-Oncology, Ospedale Annunziata, Cosenza, Italy, <sup>4</sup> Institute of Hematology "L. and A. Seràgnoli", Sant'Orsola-Malpighi University Hospital, Bologna, Italy, <sup>5</sup> Dipartimento di Scienze Mediche, Chirurgiche e Tecnologie Avanzate "G.F. Ingrassia", University of Catania, Catania, Italy

## OPEN ACCESS

### Edited by:

J. Luis Espinoza,  
Kindai University, Japan

### Reviewed by:

Belinda Pinto Simoes,  
University of São Paulo, Brazil  
Vemika Chandra,  
Children's Hospital of Philadelphia,  
United States

### \*Correspondence:

Elena Maria Elli  
elena.elli@libero.it  
Giuseppe Alberto Palumbo  
palumbo.ga@gmail.com

### Specialty section:

This article was submitted to  
Hematologic Malignancies,  
a section of the journal  
Frontiers in Oncology

**Received:** 07 June 2019

**Accepted:** 21 October 2019

**Published:** 07 November 2019

### Citation:

Elli EM, Baratè C, Mendicino F,  
Palandri F and Palumbo GA (2019)  
Mechanisms Underlying the  
Anti-inflammatory and  
Immunosuppressive Activity of  
Ruxolitinib. *Front. Oncol.* 9:1186.  
doi: 10.3389/fonc.2019.01186

The JAK-STAT signaling pathway plays a central role in signal transduction in hematopoietic cells, as well as in cells of the immune system. The occurrence in most patients affected by myeloproliferative neoplasms (MPNs) of driver mutations resulting in the constitutive activation of JAK2-dependent signaling identified the deregulated JAK-STAT signal transduction pathway as the major pathogenic mechanism of MPNs. It also prompted the development of targeted drugs for MPNs. Ruxolitinib is a potent and selective oral inhibitor of both JAK2 and JAK1 protein kinases. Its use in patients with myelofibrosis is associated with a substantial reduction in spleen volume, attenuation of symptoms and decreased mortality. With growing clinical experience, concerns about infectious complications, and increased risk of B-cell lymphoma, presumably caused by the effects of JAK1/2 inhibition on immune response and immunosurveillance, have been raised. Evidence shows that ruxolitinib exerts potent anti-inflammatory and immunosuppressive effects. Cellular targets of ruxolitinib include various components of both the innate and adaptive immune system, such as natural killer cells, dendritic cells, T helper, and regulatory T cells. On the other hand, immunomodulatory properties have proven beneficial in some instances, as highlighted by the successful use of ruxolitinib in corticosteroid-resistant graft vs. host disease. The objective of this article is to provide an overview of published evidence addressing the key question of the mechanisms underlying ruxolitinib-induced immunosuppression.

**Keywords:** ruxolitinib, JAK inhibitors, immune system, immunosuppression, myeloproliferative neoplasms (MPNs), natural killer (NK) cells, dendritic cells, T cells

## INTRODUCTION

The majority of patients affected by classic Philadelphia chromosome-negative (or *BCR-ABL1*-negative) myeloproliferative neoplasms (MPNs), including polycythemia vera (PV), essential thrombocythemia (ET), and myelofibrosis (MF), harbor mutations of the genes encoding for Janus kinase 2 (JAK2), thrombopoietin receptor gene (MPL), or calreticulin (CALR) (1, 2). Among patients with primary myelofibrosis (PMF), ~50% carry a *JAK2* mutation, 35% a *CALR* mutation,

and <10% an *MPL* mutation (3, 4). By contrast, among patients with PV, *JAK2* mutations are predominant (>95%) (4). These so-called “driver mutations” which are mutually exclusive, result in constitutively activated *JAK2* signaling and upregulation of *JAK*-signal transducer and activator of transcription (*STAT*) target genes (2, 4). The *JAK*-*STAT* signaling pathway plays a central role in normal hematopoiesis by mediating signals from a variety of cytokines and hematopoietic growth factors, in hematopoietic stem cells (1). It is also crucial for cytokine activation and signaling in the immune system (5, 6). As a consequence, patients with MPNs, and particularly MF, exhibit both uncontrolled myeloproliferation and abnormally elevated levels of circulating proinflammatory cytokines causing disease-related systemic symptoms (7).

Four members of the *JAK* family of kinases are recognized (*JAK1*, *JAK2*, *JAK3*, and *TYK2*). The *JAK* family plays a pivotal role in myeloid and lymphoid cell proliferation and differentiation and are differentially activated by different cytokines and growth factors. For example, the *JAK2* protein kinase is primarily associated with the hematopoietic growth factors erythropoietin and thrombopoietin, and mediates the process of differentiation, proliferation, and avoidance of apoptosis; the *JAK1* isoform is involved in the signaling pathway of proinflammatory cytokines such as *IL-2*, *IL-6*, and *TNF-alpha* (1). Interdependence between *JAK* family members is common (5).

The discovery in MPNs patients of mutations affecting *JAK2* signaling has led to the identification of deregulated signaling through the *JAK*-*STAT* pathway as a major pathogenic mechanism of MPNs and prompted the development of drugs targeting *JAK2* (8–11). Ruxolitinib (SB1518) was the first *JAK2* inhibitor to be granted approval and reached the market in the US and Europe in 2011. This drug is a potent and selective oral inhibitor of both *JAK2* and *JAK1* protein kinases (IC<sub>50</sub>, 2.8 nM, and 3.3 nM, respectively). It is currently indicated for the treatment of patients with intermediate- or high-risk MF, including PMF, post-polycythemia vera, and post-essential thrombocythemia MF (PPV/PET-MF). Ruxolitinib is also indicated for the treatment of patients with PV who have had an inadequate response or are intolerant to cytoreductive therapy with hydroxyurea (12, 13). The drug is not specific for the mutated form of *JAK2* and inhibits both the wild-type and *JAK2V617F*.

**Abbreviations:** APC, Antigen-presenting cells; BCCs, Basal cell carcinomas; CALR, Calreticulin; COMFORT, Controlled Myelofibrosis Study with Oral *JAK* Inhibitor Treatment; DCs, Dendritic cells; EMA, European Medicines Agency; ET, Essential thrombocythemia; FDA, U.S. Food and Drug Administration; GVHD, Graft versus host disease; HBV, Hepatitis B virus; HCV, Hepatitis C virus; IC<sub>50</sub>, Half maximal inhibitory concentration; IFN, Interferon; IL, Interleukin; *JAK*, Janus kinase; JUMP, *JAK* Inhibitor Ruxolitinib in Myelofibrosis Patients; MF, Myelofibrosis; *MPL*, Thrombopoietin receptor gene; MPNs, Myeloproliferative neoplasms; NKs, Natural killer cells; NMSC, Nonmelanoma skin cancer; PPV/PET-MF, Post-polycythemia vera, and post-essential thrombocythemia MF; PV, Polycythemia vera; RESPONSE, Randomized Study of Efficacy and Safety in Polycythemia Vera With *JAK* Inhibitor INCB018424 Versus Best Supportive Care; ROCK, Rho-associated coiled-coil kinase; SCC, Squamous cell carcinoma; *STAT*, Signal transducer and activator of transcription; Th, T helper; *TNF*, Tumor necrosis factor; Treg, T regulatory cells.

The efficacy of this compound in a larger cohort of patients with MF was demonstrated at first on 2 phase III randomized studies (14, 15), Controlled Myelofibrosis Study with Oral *JAK* Inhibitor Treatment (COMFORT-I and COMFORT-II). The anti-*JAK2* inhibitory action is responsible for the efficacy of ruxolitinib on the control of myeloproliferation, reducing splenomegaly and, in some cases, the *JAK2V617F* allele burden (16–18). The inhibition of wild-type *JAK2* protein results also in myelosuppression, primarily expressed as anemia and thrombocytopenia, and less frequently by neutropenia, which rarely leads to drug discontinuation. The anti-*JAK1* inhibitory action is responsible for the reduction of proinflammatory cytokines, with a consequent improvement of symptoms, quality of life and, ultimately, bone marrow fibrosis (19, 20). At the same time, the anti-cytokine action could potentially cause an immunosuppressive effect of the drug, since the immune system and the hematopoietic system share intracellular signaling pathways, mediated by common receptors for cytokines and growth factors which, by acting on the *JAK*-*STAT* pathway, are important for the proliferation, differentiation, and activation of cells involved in innate and adaptive immune responses. Indeed, relatively high rates of infectious complications and of hematological and solid tumors have been observed in ruxolitinib-treated patients (21–23). Immunomodulatory properties, on the other hand, may be beneficial in some instances as highlighted by the use of ruxolitinib in graft vs. host disease (GVHD) (24–28).

The key question as to what the immunological targets of *JAK1/2* inhibition are is far from being solved, and data elucidating the mechanisms of ruxolitinib-induced immunosuppression are still limited. The objective of this article is to provide an overview of published evidence addressing this question. We will first discuss the data that have suggested and confirmed an immunosuppressive action exerted by ruxolitinib; we will then briefly review the available studies that have attempted to dissect the mechanisms underlying ruxolitinib-induced immunomodulation.

## IMMUNOSUPPRESSIVE ACTIVITY OF RUXOLITINIB

In the registrative COMFORT I trial, grade 3 and 4 neutropenia were recognized in 7.1 and 2% of patients in the ruxolitinib and placebo arm of the study (15). More recently, 5-year follow-up of COMFORT II trial noted grade 3 or 4 neutropenia or leukopenia in 8.9 and 6.3% of ruxolitinib-treated patients (29). Unfortunately, leukocyte subpopulations and functions or antibody deficiency were not documented in both studies. Theocharides et al. (30) recognized that homozygous calreticulin mutations in PMF lead to acquired myeloperoxidase deficiency and consequential neutrophil reduced efficiency.

The possibility that ruxolitinib may increase the susceptibility to infections in patients with MF was first suggested by the results of the COMFORT-I and COMFORT-II trials. In the COMFORT-I study (15), patients treated with ruxolitinib had higher rates of bacterial and herpes zoster infections than patients treated with

placebo. The incidence of urinary tract infections and herpes zoster was 10.5 and 2.1%, respectively in the first 12 weeks of therapy, 6.7 and 3.5% between 12 and 24 weeks, 7.7 and 3.4% between 24 and 36 weeks. At 5 years of follow-up (18), the most severe infections were pneumonia and sepsis, which occurred at similar rates in patients treated with ruxolitinib or placebo. Over time, herpes zoster infections occurred at a higher rate in patients treated with ruxolitinib compared with placebo, but in the majority of cases these were single episodes of grade 1 or 2. Septic events were reported as a cause of death in 6 ruxolitinib arm patients, 1 in the placebo arm and 4 placebo arm patients with subsequent cross-over.

In the COMFORT-II study (29), most infections were grade 1 or 2. Pneumonia was the only serious infectious adverse event reported (1% in the ruxolitinib group vs. 5% in the best available therapy group). In the final 5-year analysis, with a median ruxolitinib exposure time of 2.6 years, herpes zoster infections (11.5%), pneumonia (13.1%), sepsis (7.9%), and urinary tract infections (24.6%) were reported, with only two confirmed cases of tuberculosis (1%). In both COMFORT studies, the incidence of severe infections in patients treated with ruxolitinib was similar to that observed in patients in the control group. A summary of exposure-adjusted rates of immunosuppressive events during long-term treatment with ruxolitinib in the COMFORT studies is presented in **Table 1**.

In the JUMP study (an open-label phase 3b expanded-access trial which enrolled 2,233 patients from 26 countries, without access to ruxolitinib outside of a clinical study), a preliminary analysis of 1,144 intermediate and high-risk patients showed that the incidence of infections was low: pneumonia (5.3%), urinary tract infections (6%), and nasopharyngitis (6.3%). Tuberculosis was reported in five patients (0.3%); the reactivation of hepatitis B virus (HBV) in 1 patient (0.1%), while the incidence of herpes zoster infections was 3.6% (31). Possible explanations for the occurrence of potentially severe and/or recurrent viral infections include impairment of natural killer cells (NKs), which have a key role in controlling herpes infections, especially when T cells are low and a reduction of T regulatory cells (Tregs) protective against virus occurs.

Also, in the PV setting, two randomized phase 3 trials which compared ruxolitinib vs. best Supportive Care (RESPONSE and RESPONSE-2 studies) (32, 33) showed that the rates of herpes zoster infections during extended treatment were higher in patients receiving ruxolitinib as respect to best available therapy. In RESPONSE at 80 weeks, the rate per 100 patient-years of exposure was 5.4 vs. none (34). Most herpes zoster infections were grade 1 or 2 and were resolved without sequelae. Similarly, week 80 data in RESPONSE-2 (35) showed that the all-grade exposure-adjusted rate of herpes zoster infection per 100 patient-years of exposure was 3.8 in patients originally randomized to ruxolitinib, 7.5 in patients receiving ruxolitinib after crossover, and none in the best available therapy arm. No pneumonia or tuberculosis reactivation was observed in the ruxolitinib group. A summary of exposure-adjusted rates of immunosuppressive events during long-term treatment with ruxolitinib in the RESPONSE studies is presented in **Table 2**.

**TABLE 1 |** Common immunosuppressive events during long-term treatment with ruxolitinib: Data from the final 5-year analyses of the COMFORT I and COMFORT II trials.

	Ruxolitinib, randomized (n = 155)		Ruxolitinib, crossover* (n = 111)		Best available therapy (n = 151)	
	All grades	Grade 3 or 4	All grades	Grade 3 or 4	All grades	Grade 3 or 4
<b>COMFORT I (Verstovsek S et al. J Hematol Oncol 2017; 10:55)</b>						
Upper respiratory tract infection	8.5	0	9.5	0	15.5	1.0
Urinary tract infection	7.5	1.0	6.7	1.2	6.9	1.0
Pneumonia	7.2	5.1	7.1	3.2	10.7	7.8
Herpes zoster	3.5	0	5.8	0.4	1.0	0
Bronchitis	3.1	0	4.5	1.2	1.9	0
Nasopharyngitis	3.1	0	3.5	0	9.1	0
Sinusitis	2.6	0.2	2.8	0	2.9	1.0
Cellulitis	2.1	0.4	1.1	0	1.9	0
Influenza	1.7	0	1.1	0.4	0	0
Sepsis	1.7	1.7	1.5	1.5	1.9	1.0
Tooth abscess	1.5	0.2	1.5	0	0	0
Oral herpes	1.3	0	0.7	0	1.9	0
Skin infection	1.1	0	1.1	0	1.0	0
Viral infection	1.1	0	0.8	0	0	0
Viral gastroenteritis	0.9	0	0.4	0	1.9	0
Diverticulitis	0.8	0.2	1.1	0.4	1.9	0
Ear infection	0.8	0	1.5	0	0	0
Fungal infection	0.8	0	0.7	0.4	1.9	0
Localized infection	0.8	0	0.4	0.4	1.0	0
Lower respiratory tract infection	0.8	0	0.4	0	1.9	1.0
Septic shock	0.4	0.4	1.1	1.1	0	0
	Ruxolitinib, randomized (n = 146)		Ruxolitinib, crossover* (n = 45)		Best available therapy (n = 73)	
	All grades	Grade 3 or 4	All grades	Grade 3 or 4	All grades	Grade 3 or 4
<b>COMFORT II (Harrison CN et al. Leukemia 2016; 30:1701-7)</b>						
Bronchitis	10.0	NR	3.8	NR	9.0	NR
Nasopharyngitis	9.8	NR	5.0	NR	13.4	NR
Pyrexia	9.5	NR	10.0	NR	10.5	NR
Anemia	NR	7.6	NR	15.1	NR	7.5
Thrombocytopenia	NR	4.9	NR	11.3	NR	6.0
Pneumonia	NR	2.4	NR	1.3	NR	6.0
Herpes zoster	3.9	NR	6.3	NR	0	NR
Gastroenteritis	4.2	NR	1.3	NR	1.5	NR
Urinary tract infection	4.6	NR	8.8	NR	3.0	NR
Cystitis	3.7	NR	1.3	NR	4.5	NR

Data are exposure-adjusted rates per 100 patient years, regardless of relationship to study drug.

\*Patients randomized to best available therapy were allowed to crossover to receive ruxolitinib after 6 (COMFORT I) or 12 (COMFORT II) months in response to protocol-defined disease progression.

COMFORT, Controlled Myelofibrosis Study with Oral JAK Inhibitor Treatment; NR not available.

**TABLE 2 |** Common immunosuppressive events during long-term treatment with ruxolitinib: Data from week 80 analyses of the RESPONSE and RESPONSE-2 trials in patients with polycythemia vera.

	Ruxolitinib, randomized (n = 110)		Ruxolitinib, crossover* (n = 98)		Best available therapy (n = 111)**	
	All grades	Grade 3 or 4	All grades	Grade 3 or 4	All grades	Grade 3 or 4
<b>RESPONSE (Verstovsek S et al. Haematologica 2016; 101:821-9)</b>						
All infections	29.4	4.0	27.8	5.4	58.4	4.1
Herpes zoster	5.3	0.9	5.4	0.7	0	0
Pyrexia	5.3	0	5.4	0.7	6.8	0
Nasopharyngitis	5.7	0	6.1	0	12.2	0
	Ruxolitinib, randomized (n = 74)		Ruxolitinib, crossover* (n = 58)		Best available therapy (n = 75)	
	All grades	Grade 3 or 4	All grades	Grade 3 or 4	All grades	Grade 3 or 4
<b>RESPONSE-2 (Griesshammer M et al. Ann Hematol 2018; 97:1591-1600)</b>						
Infections and infestations	24.9	2.3	29.9	1.5	33.7	3.7
Upper respiratory tract infection	2.3	0	3.0	0	13.1	0
Nasopharyngitis	3.8	0	9.0	0	3.7	0
Influenza	4.5	0.8	1.5	0	9.4	1.9
Anemia	14.3	0	17.9	0	3.7	1.9
Thrombocytopenia	1.5	0	4.5	0	15.0	5.6
Pneumonia	0.8	0	0	0	1.9	1.9
Herpes zoster	3.8	0	7.5	0	0	0
Urinary tract infection	1.5	0	1.5	0	0	0
Urosepsis	0.8	0	0	0	0	0
Septic shock	0	0	0	0	1.9	1.9

Data are exposure-adjusted rates per 100 patient years, regardless of relationship to study drug.

\*Patients randomized to best available therapy were allowed to crossover to receive ruxolitinib after 32 (RESPONSE) or 28 (RESPONSE-2) weeks if they did not meet the primary end or for safety-related reasons.

\*\*One patient was randomized to best available therapy but did not receive study treatment.

RESPONSE, Randomized Study of Efficacy and Safety in Polycythemia Vera With JAK Inhibitor INCB018424 vs. Best Supportive Care.

Details of the characteristics of the randomized phase III trials of ruxolitinib in MF or PV can be found in **Supplementary Table S1**.

With the increasing use of ruxolitinib in clinical practice, there have also been reports of HBV reactivation and severe and uncommon infections including *Cryptococcus neoformans* pneumonia, toxoplasmosis retinitis, disseminated tuberculosis, and progressive multifocal leukoencephalopathy (36–42). Particularly, tuberculosis reactivation during ruxolitinib may be due to the impairment of dendritic cells (DCs) and IL-12 production, a key cytokine involved in the transcription of interferon (IFN)-gamma. In addition, ruxolitinib induces depression of T helper (Th)1 lymphocyte responses and

production of IFN-gamma and tumor necrosis factor (TNF)-alpha (43).


A recent multicenter retrospective study on a large cohort of 446 patients with MF treated with ruxolitinib (44) showed that almost 30% of patients, with a median exposure to ruxolitinib of 23.5 months, experienced infectious events (incidence rate of 17 cases per 100 patients/year). Infections involving the respiratory tract were predominant and accounted for 50% of all infections reported. The rate of infections tended to decrease with time. Respiratory tract infections were more frequently observed (81 events, 50%), and bacteria were the most frequent etiological agents (68.9%). However, viral (14.9%) and fungal infections (2.5%) were also observed. Multivariate analysis found that a previous infectious event and a high-risk score according to the International Prognostic Score System (45) correlated significantly with a greater risk of infections. Of note, splenomegaly reduction by  $\geq 50\%$  from baseline to 3 months was significantly associated with longer infection-free survival. **Figure 1** summarizes the incidence of the most important infectious events, regardless of relationship to study drug, in the principal studies of ruxolitinib in MF.

An increased risk of lymphoma during treatment with ruxolitinib or discontinuation of ruxolitinib has been suggested by sporadic reports (46, 47). Recently, four cases of aggressive lymphoma were reported during treatment among 69 MPNs patients receiving ruxolitinib (5.8% rate of lymphoma development) (22). Among MPNs patients from the same center who did not receive ruxolitinib ( $n = 557$ ), two (0.36%) developed lymphoma. The analysis of an independent MPNs cohort ( $n = 929$ ) found similar rates of lymphoma development (3.51% in patients receiving JAK1/2 inhibitor therapy and 0.23% in patients receiving conventional therapy). The lymphomas were found to have developed from pre-existing B-cell clones. In mice lacking STAT1 and treated with ruxolitinib, the phenotype of coexisting MPNs and B-cell transformation was reproduced, suggesting that the reduced ability of T lymphocytes and NKs to eliminate malignant hematopoietic cells, caused by STAT1 deficiency, could facilitate the development of a B-cell lymphoma (22). However, in a recent large database review, that included 2,583 patients with MPNs (comprised of 1,617 patients with MF) no statistically significant difference in the incidence of a subsequent lymphoma diagnosis in patients with MPNs when comparing those who received prior JAK inhibitor therapy vs. those who did not, was found (48).

Five-year efficacy data on ruxolitinib in MF showed 17.1% of patients on ruxolitinib went on to have basal cell carcinomas (BCCs) or squamous cell carcinomas (SCCs) compared with only 2.7% of patients on best available therapy (29). Skin cancers occurring in patients treated with ruxolitinib have been reported in the dermatology literature to display more aggressive biological behavior and metastatic potential (49). In a recent large international case-control study, including 1,881 patients with MPNs, a significantly higher risk of non-melanoma skin cancer (NMSC) was documented for ruxolitinib, suggesting that JAK1-JAK2 inhibitor may act as immunosuppressive agent (23).

In the PV setting, in the RESPONSE study, the rates of NMSC per 100 patient-years of exposure seemed to be higher in





patients receiving ruxolitinib, as respect to best available therapy (4.4 vs. 2.7). NMSC cancers were observed in the originally randomized ruxolitinib arm, mainly in patients with a history of NMSC or precancer; however, exposure-adjusted rates at the time of this analysis were generally similar between the originally randomized ruxolitinib and best available therapy arms (34).

Finally, the potent anti-inflammatory and immunosuppressive properties of ruxolitinib can be exploited for the management of severe conditions including autoimmune or inflammatory diseases (50, 51) and corticosteroid-resistant GVHD (24–28). Preliminary studies with topical ruxolitinib demonstrated improvement in psoriasis compared to treatment with placebo or other topical approved therapies, but it was not a sustained improvement after discontinuation (52); importantly, systemic absorption was minimal, and there was no evidence of systemic toxicity. The potential efficacy of ruxolitinib is also being investigated in the treatment of alopecia areata and vitiligo, but these studies are still in their infancy (51, 53). A recent multicenter retrospective analysis of ruxolitinib as salvage treatment in patients with steroid-resistant acute or chronic GVHD found elevated overall response rates (>80%) and 6-month survival rates ranging from 79% (acute disease) to 97% (chronic disease), despite the fact that the patients had been heavily pretreated for GVHD, and their condition was generally severe (26). Based on the evidence supporting a role in preventing GVHD, ruxolitinib received in 2016 Breakthrough Therapy Designation from the US FDA for the treatment of GVHD (54).

## CELLULAR TARGETS OF THE IMMUNOSUPPRESSIVE ACTIVITY OF RUXOLITINIB


Available evidence suggests that ruxolitinib acts on cellular components of both innate and adaptive immunity (5). The

innate immune system ensures front-line host defense and includes anatomic barriers, antimicrobial molecules, such as complement and cellular components: eosinophils, neutrophils, mast cells, basophils, NKs, monocytes/macrophages, and DCs; the adaptive immune system comprises CD4+ T-helper lymphocytes (Th1, Th2, Th17) and Tregs, CD8+ cytotoxic lymphocytes, and B lymphocytes (55). **Figure 2** summarizes the complex relationship between ruxolitinib and the immune “orchestra.” In the following paragraphs, we will briefly discuss studies reporting the effects of ruxolitinib on NKs, DCs, and T cells. Potential additional immunological targets of ruxolitinib are also briefly discussed.

### Natural Killer Cells

NKs are the main effectors of the innate immune system. They are large lymphocyte-like cells with distinctive cytotoxic granules and lack antigen-specific receptors (55). They can detect and exert lytic activity on certain virus-infected cells, as well as on malignant cells, and are therefore important components also of cancer immunosurveillance mechanisms. NKs also produce cytokines, mainly IFN-gamma, and TNF-alpha, which modulate the differentiation of cells involved in adaptive immunity (Th1) and induce the maturation of DCs. The process leading to the development and differentiation of NKs from lymphoid precursors is mainly regulated by IL-15 and IL-2 with the involvement of the JAK1-JAK3/STAT5 pathway (5).

A recent study provided a detailed analysis of the influence of ruxolitinib on the biology of NKs by comparing the effects of JAK1/2 inhibition on this cell type in a cohort of 28 MPNs patients with or without ruxolitinib treatment and 24 healthy individuals (56). The analysis included cell frequency, receptor expression, proliferation, immune synapse formation, and cytokine signaling. The study found a reduction in NKs number in ruxolitinib-treated patients that was linked to the appearance of clinically relevant infections. An increase in the



ratio of immature/mature NKs suggested that the observed reduction was due to impaired NKs maturation. *In vitro* experiments showed that the observed reduction in killing activity was caused by impairment in lytic synapse formation with target cells. The effect was reversible, as NKs function was restored upon ruxolitinib discontinuation. The authors suggested that ruxolitinib was likely inhibiting the JAK1-JAK3/STAT5 pathway downstream of the IL-2- and IL-15 receptors. In a study that compared the effects of ruxolitinib and the JAK2-specific inhibitor pacritinib (TG101348) on the function and activation of NKs, ruxolitinib was shown to completely block IL-2, IL-15, and the phosphorylation of STAT5 mediated by DCs, along with the capacity of NKs to secrete IFN-gamma or lyse target cells (57). In contrast, pacritinib treatment of stimulated NKs resulted in substantially less functional impairment. Therefore, selective JAK2 blockade ensures greater NKs activity than non-selective targeting of both JAK1 and JAK2.

Lastly, Ankathatti and Hu (58) showed, in a study with human cell lines, that small molecule blockers, including ruxolitinib, for JAK-STAT pathway, significantly inhibited cytokine secretion by macrophages activated by viral infectious trigger. Usually,

macrophages respond to viral infection by releasing soluble mediators, helping in the recruitment of innate, and adaptive effector cells. In this process, they recruit NK cells to the site of infection and with DCs help, regulate NK maturation, and NK killing activity through production of type I interferons, IL-12, and IL-15. The authors demonstrated that NK cells activated by cytokines produced by activated macrophages could be inhibited in their killing activity by ruxolitinib, because of block of IL-12 and IL-15 production by macrophages.

## Dendritic Cells

DCs are important antigen-presenting cells (APC) and phagocytic cells; as macrophages, they are responsible for presenting antigens to T lymphocytes and initiating adaptive immunity (55). DCs are the unique propriety of inducing the differentiation of naive T cells in helper and effector T cells. More in detail, the induction of an adaptive immune response initiates following the ingestion of a pathogen by an immature DCs in the infected tissue. Upon ingestion of a pathogen, DCs become activated and migrate to a nearby lymph node where they prime T lymphocytes toward effector or regulatory responses. Activated

DCs secrete cytokines that influence both innate and adaptive immune responses and play, therefore, a key role in determining whether, and how, the immune system responds to an infectious agent. In particular, DCs control the adaptive response that leads to the production of IL-12 and IL-23, cytokines that drive Th1 and Th17 lymphocytes phenotypes. More specifically, Th1 cells produce cytokine IFN-gamma, IL-2, and TNF-alpha. TNF-alpha improves Th1 against intracellular pathogens. Th17 secrete IL-17 and IL-22, key cytokines against extracellular bacteria. More mature DCs mediate the induction of T-regs.

The important study by Heine and coworkers on the effects of ruxolitinib on DCs biology demonstrated *in vitro* and *in vivo* that JAK1/2 inhibitor therapy affects cell differentiation, tissue migration and function of DCs resulting in impaired induction of allogenic or antigen-specific T-cell responses, including viral clearance. These findings were among the first to clearly show that ruxolitinib has potent anti-inflammatory and immunomodulating activity, and to provide a possible explanation for the increased infection rates reported in MPNs patients exposed to ruxolitinib (59).

A more recent study analyzed the impact of ruxolitinib on the migration of DCs with a focus on short-term effects and the identification of potential target molecules mediating these effects (60). The study identified Rho-associated coiled-coil kinase 1 (ROCK) as a target of ruxolitinib. Notably, this protein regulates the reorganization and contraction of cellular actin-myosin filaments and plays an important role in the migration of DCs. Via interference with ROCK activation, ruxolitinib profoundly impairs DCs migration; subsequently, the loss of trafficking DCs may lead to reduced activation of T cells in draining lymph nodes and might, therefore, explain the abundance of these proinflammatory cells from the blood of MPNs patients after ruxolitinib treatment (61).

## CD4+ T Cells

CD4+ T cells, which differentiate from naïve T cells, are a heterogeneous cell population with a central role in adaptive immunity. They mainly act via the secretion of cytokines and chemokines that induce and/or recruit target cells and are able to adopt a series of distinct differentiated states including T-helper and Treg cell types (61, 62). Tregs suppress autoreactive lymphocytes, prevent autoimmune disease, and control innate and adaptive immune responses (62). Tregs also control viral, fungal, and protozoan infections. They produce inhibitory cytokines such as IL-10 and TGF-beta that promote fibrosis, affect the function and induce apoptosis of T-effector lymphocytes. Given the importance of JAK-STAT signaling in regulating the fate and function of CD4+ T cells, a study attempted to define the effect of ruxolitinib on the function of this cell subset in MPNs patients (61). The study, which involved both *in vitro* and *in vivo* experiments, highlighted a reduction in total CD3+ cells after 3 weeks of ruxolitinib treatment in MPNs patients. The number of Th1, Th17, and Tregs were also reduced, a result that was validated also *in vitro*. In line with these findings, inflammatory cytokines, including TNF-alpha, IL-5, IL-6, and IL-1B, were found to be significantly downregulated in T cells isolated from the patients. Notably, ruxolitinib did

not interfere with the T cell receptor signaling pathway, but inhibited IL2-dependent STAT5 activation. The reduction in the number of Tregs induced by ruxolitinib was in line with the results of a previous study conducted by Massa et al. in 18 patients with MF to define the impact of JAK1/2 inhibition on immunosurveillance (63). Massa et al. showed that the treatment with ruxolitinib resulted in a long-lasting reduction in circulating Tregs. Dose reduction or withdrawal of the drug failed to restore the Treg subset to values comparable to those of the control, suggesting that the effect of ruxolitinib was irreversible. The authors suggested that the severe reduction in circulating Tregs observed in their patients exposed to ruxolitinib may constitute a major cause of immunosurveillance disruption.

A recent study investigated the frequency and function of CD4+ T cell subsets in 50 MPN patients at baseline as well as during treatment with either ruxolitinib or fedratinib (SAR302503; TG101348), a selective JAK2 inhibitor (64). The study showed, for the first time, that Tregs are reduced in MPNs patients compared to healthy controls and that this decrease becomes more pronounced following JAK inhibitor therapy. However, after 6 months, responders to treatment with JAK inhibitor therapy displayed an increased number of Th17, which are known to secrete IL-17 and to play a key role in autoimmune disease, as well as in tumor immunosurveillance. This result could suggest a potential polarization from a Treg/Th1 to a Th17 phenotype. Functional "silencing" of Th cells, both *in vivo* and *in vitro*, and the blockade of proinflammatory cytokines from these cells were also observed. Thus, treatment with JAK inhibitor therapy had both short- and long-term effects on CD4+ T cells in MPNs. The short-term effect involved a reduction in Treg number, Th silencing, and reduction of cytokine secretion. The long-term effect, on the other hand, resulted in immune response polarization toward a Th17-type response. Thanks to this repolarization effect, according to the authors, immunosurveillance against the malignant clone could be resumed.

## Additional Immunological Targets of Ruxolitinib

Given the complexity of the signaling pathways modulated by JAK1 and JAK2, and the wide range of immune system components affected, it can reasonably be expected that JAK1/2 inhibitor therapy with ruxolitinib will have other cellular targets beside those described in the previous sections. An important open question concerns the effect of ruxolitinib on the B cell population and, in particular, on antibody production. Optimal B cell activity requires a complex interplay of inputs from a variety of T cells, including T effector cells and Tregs (5). Establishing the impact of JAK inhibition on antibody production is currently difficult to predict yet crucial when considering the need for optimal defense against pathogenic infections. No evidence of antibody deficiency has emerged from the COMFORT-I and II studies (15, 18, 29). With regard to possible abnormalities in B cells associated with MPNs, available evidence shows that the mutant clone may also include lymphoid-derived cells (5). A number of reports in small groups of patients have shown that a

few patients with chronic MPNs carry the *JAK2* V617F mutation in both B- and T-lymphocytes, suggesting that the mutation occurred early, in lympho-myeloid progenitor cells (65, 66).

Other potential targets of ruxolitinib include neutrophils and macrophages, two important components of innate immunity. Neutrophil activation is mediated by *JAK1/2* signaling (26). Furthermore, it has been shown that neutrophils have a key role in the pathogenesis of acute GVHD (26, 67, 68). As for macrophages, ruxolitinib was shown to prevent the up-regulation of various proinflammatory cytokines in human macrophages (69). Ruxolitinib impairs cytokine expression by inhibiting LPS/TLR4/IFN beta signaling pathway. Cytokine repression contributes to the anti-inflammatory effect of ruxolitinib. However, earlier data suggested that the inhibition of *JAKs* may increase the inflammatory potential of macrophages exposed to TLR4 agonists, which potently stimulate cytokine production (70).

## CONCLUSIONS

The substantial clinical benefits and efficacy associated with the use of *JAK1/2* inhibitor therapy need to be balanced against the multiple effects on components of both innate and adaptive immunity, including NKs, DCs, and T cells (Th and Tregs). Ruxolitinib impairs several cytokines, modulates DCs function and T cell response and reduces NKs levels in MPNs patients, which may lead to increased risk of opportunistic infection and reactivation of latent infections. Age and comorbidities, treatments (such as steroids) and environmental exposure may influence the risk of infections, that may occur early and late after treatment.

Thus, prior to initiation of therapy with ruxolitinib and other *JAK* inhibitors, patient counseling, assessment of risk factors for tuberculosis reactivation, screening for previous hepatitis B and C exposure, anti-infective prophylaxis, along with particular caution in patients who are already immunocompromised, are highly recommended (12, 13, 43).

Surveillance of HBV markers and viral load are important due to the high incidence of latent HBV and reactivation during steroid or immunosuppressive drugs. Prompt antiviral prophylaxis should be considered in hematological patients with a high risk of HBV reactivation according to treatment guidelines (71). Similarly, HCV carriers must be identified and treated according to viral load.

Also, active surveillance of second solid neoplasia and hematological diseases, including high-grade lymphomas, should

always be performed during therapy. Accordingly, current ruxolitinib prescriber information labels warn of the risk of NMSC, and performing periodic skin examinations is recommended patients (23). Also, the detection of a preexisting B-cell clone may identify individuals at risk for lymphoma development (22).

More robust data are necessary to answer the question of possible immune derangement of ruxolitinib treatment in MPNs patients. Although its immunosuppressive properties should not be forgotten, ruxolitinib remains a cornerstone in the treatment of BCR-ABL negative MPNs, where its anti-inflammatory activity is a benefit for the patient and especially in MF where symptom control, reduction of splenomegaly and possibly improvement of survival may be obtained. Finally, the potent anti-inflammatory and immunosuppressive activity of ruxolitinib is proving valuable for the management of severe and life-threatening conditions like GVHD in the allogeneic stem cell transplantation setting.

## AUTHOR CONTRIBUTIONS

EE and GP conceived the manuscript and prepared the first draft. All authors critically revised the article for important intellectual content. All authors gave final approval of the version to be published, and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## FUNDING

Medical writing assistance for the preparation of this article was funded by Novartis Farma, Italy.

## ACKNOWLEDGMENTS

The authors thank Dr. Lorenza Lanini for providing editorial assistance during manuscript preparation on behalf of Health Publishing & Services Srl, Milan, Italy.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fonc.2019.01186/full#supplementary-material>

## REFERENCES

- Quintàs-Cardama A, Kantarjian H, Cortes J, Verstovsek S. Janus kinase inhibitors for the treatment of myeloproliferative neoplasias and beyond. *Nat Rev Drug Discov*. (2011) 10:127–40. doi: 10.1038/nrd3264
- Griesshammer M, Sadjadian P. The BCR-ABL1-negative myeloproliferative neoplasms: a review of *JAK* inhibitors in the therapeutic armamentarium. *Expert Opin Pharmacother*. (2017) 18:1929–38. doi: 10.1080/14656566.2017.1404574
- Klampfl T, Gisslinger H, Harutyunyan AS, Nivarthi H, Rumi E, Milosevic JD, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. (2013) 369:2379–90. doi: 10.1056/NEJMoa1311347
- Palumbo GA, Stella S, Pennisi MS, Piroso MC, Fermo E, Fabris S, et al. The role of new technologies in myeloproliferative neoplasms. *Front Oncol*. (2019) 9:321. doi: 10.3389/fonc.2019.00321
- McLornan DP, Khan AA, Harrison CN. Immunological consequences of *JAK* inhibition: friend or foe? *Curr Hematol Malig Rep*. (2015) 10:370–9. doi: 10.1007/s11899-015-0284-z



6. Senkevitch E, Durum S. The promise of Janus kinase inhibitors in the treatment of hematological malignancies. *Cytokine*. (2017) 98:33–41. doi: 10.1016/j.cyto.2016.10.012
7. Bryan JC, Verstovsek S. Overcoming treatment challenges in myelofibrosis and polycythemia vera: the role of ruxolitinib. *Cancer Chemother Pharmacol*. (2016) 77:1125–42. doi: 10.1007/s00280-016-3012-z
8. Baxter EJ, Scott LM, Campbell PJ, East C, Fourouclas N, Swanton S, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet*. (2005) 365:1054–61. doi: 10.1016/s0140-6736(05)71142-9
9. James C, Ugo V, Le Couedic JP, Staerk J, Delhommeau F, Lacout C, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature*. (2005) 434:1144–8. doi: 10.1038/nature03546
10. Kralovics R, Passamonti F, Buser AS, Teo SS, Tiedt R, Passweg JR, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med*. (2005) 352:1779–90. doi: 10.1056/NEJMoa051113
11. Levine RL, Wadleigh M, Cools J, Ebert BL, Wernig G, Huntly BJ, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell*. (2005) 7:387–97. doi: 10.1016/j.ccr.2005.03.023
12. European Medicines Agency. *Jakavi (ruxolitinib): Summary of Product Characteristics*. (2017). Available online at: [https://www.ema.europa.eu/documents/product-information/jakavi-epar-product-information\\_en.pdf](https://www.ema.europa.eu/documents/product-information/jakavi-epar-product-information_en.pdf) (accessed May 31, 2019).
13. U.S. Food and Drug Administration (2017). (FDA)JAKAFI (ruxolitinib): Prescribing Information. Available online at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2017/202192s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/202192s015lbl.pdf) (accessed May 31, 2019).
14. Harrison C, Kiladjan JJ, Al-Ali HK, Gisslinger H, Waltzman R, Stalbovska V, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. (2012) 366:787–98. doi: 10.1056/NEJMoa1110556
15. Verstovsek S, Mesa RA, Gotlib J, Levy RS, Gupta V, DiPersio JF, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. (2012) 366:799–807. doi: 10.1056/NEJMoa1110557
16. Deininger M, Radich J, Burn TC, Huber R, Paranagama D, Verstovsek S. The effect of long-term ruxolitinib treatment on JAK2p.V617F allele burden in patients with myelofibrosis. *Blood*. (2015) 126:1551–4. doi: 10.1182/blood-2015-03-635235
17. Harrison C. JAK inhibitors and myelofibrosis, Einstein and ruxolitinib. *Haematologica*. (2015) 100:409–11. doi: 10.3324/haematol.2015.124099
18. Verstovsek S, Mesa RA, Gotlib J, Gupta V, DiPersio JF, Catalano JV, et al. Long-term treatment with ruxolitinib for patients with myelofibrosis: 5-year update from the randomized, double-blind, placebo-controlled, phase 3 COMFORT-I trial. *J Hematol Oncol*. (2017) 10:55. doi: 10.1186/s13045-017-0417-z
19. Verstovsek S, Mesa RA, Gotlib J, Levy RS, Gupta V, DiPersio JF, et al. Efficacy, safety and survival with ruxolitinib in patients with myelofibrosis: results of a median 2-year follow-up of COMFORT-I. *Haematologica*. (2013) 98:1865–71. doi: 10.3324/haematol.2013.092155
20. Kvasnicka HM, Thiele J, Bueso-Ramos CE, Sun W, Cortes J, Kantarjian HM, et al. Long-term effects of ruxolitinib versus best available therapy on bone marrow fibrosis in patients with myelofibrosis. *J Hematol Oncol*. (2018) 11:42. doi: 10.1186/s13045-018-0585-5
21. Lussana F, Cattaneo M, Rambaldi A, Squizzato A. Ruxolitinib-associated infections: a systematic review and meta-analysis. *Am J Hematol*. (2018) 93:339–47. doi: 10.1002/ajh.24976
22. Porpaczy E, Tripolt S, Hoelbl-Kovacic A, Gisslinger B, Bago-Horvath Z, Casanova-Hevia E, et al. Aggressive B-cell lymphomas in patients with myelofibrosis receiving JAK1/2 inhibitor therapy. *Blood*. (2018) 132:694–706. doi: 10.1182/blood-2017-10-810739
23. Barbui T, Ghirardi A, Masciulli A, Carrobbio A, Palandri F, Vianelli N, et al. Second cancer in Philadelphia negative myeloproliferative neoplasms (MPN-K): a nested case-control study. *Leukemia*. (2019) 33:1996–2005. doi: 10.1038/s41375-41019-40487-41378
24. Choi J, Cooper ML, Alahmari B, Ritchey J, Collins L, Holt M, et al. Pharmacologic blockade of JAK1/JAK2 reduces GvHD and preserves the graft-versus-leukemia effect. *PLoS ONE*. (2014) 9:e109799. doi: 10.1371/journal.pone.0109799
25. Spoerl S, Mathew NR, Bscheider M, Schmitt-Graeff A, Chen S, Mueller T, et al. Activity of therapeutic JAK 1/2 blockade in graft-versus-host disease. *Blood*. (2014) 123:3832–42. doi: 10.1182/blood-2013-12-543736
26. Zeiser R, Burchert A, Lengerke C, Verbeek M, Maas-Bauer K, Metzelder SK, et al. Ruxolitinib in corticosteroid-refractory graft-versus-host disease after allogeneic stem cell transplantation: a multicenter survey. *Leukemia*. (2015) 29:2062–8. doi: 10.1038/leu.2015.212
27. Khandelwal P, Teusink-Cross A, Davies SM, Nelson AS, Dandoy CE, El-Bietar J, et al. Ruxolitinib as salvage therapy in steroid-refractory acute graft-versus-host disease in pediatric hematopoietic stem cell transplant patients. *Biol Blood Marrow Transplant*. (2017) 23:1122–7. doi: 10.1016/j.bbmt.2017.03.029
28. Sarmiento Maldonado M, Ramirez Villanueva P, Bertin Cortes-Monroy P, Jara Arias V, Soto Donoso K, Uribe Gonzalez P, et al. Compassionate use of ruxolitinib in acute and chronic graft versus host disease refractory both to corticosteroids and extracorporeal photopheresis. *Exp Hematol Oncol*. (2017) 6:32. doi: 10.1186/s40164-017-0092-3
29. Harrison CN, Vannucchi AM, Kiladjan JJ, Al-Ali HK, Gisslinger H, Knoops L, et al. Long-term findings from COMFORT-II, a phase 3 study of ruxolitinib vs best available therapy for myelofibrosis. *Leukemia*. (2016) 30:1701–7. doi: 10.1038/leu.2016.148
30. Theocharides AP, Lundberg P, Lakkaraju AK, Lysenko V, Myburgh R, Aguzzi A, et al. Homozygous calreticulin mutations in patients with myelofibrosis lead to acquired myeloperoxidase deficiency. *Blood*. (2016) 127:3253–9. doi: 10.1182/blood-2016-02-696310
31. Al-Ali HK, Griesshammer M, le Coutre P, Waller CF, Liberati AM, Schafhausen P, et al. Safety and efficacy of ruxolitinib in an open-label, multicenter, single-arm phase 3b expanded-access study in patients with myelofibrosis: a snapshot of 1144 patients in the JUMP trial. *Haematologica*. (2016) 101:1065–73. doi: 10.3324/haematol.2016.143677
32. Vannucchi AM, Kiladjan JJ, Griesshammer M, Masszi T, Durrant S, Passamonti F, et al. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. *N Engl J Med*. (2015) 372:426–35. doi: 10.1056/NEJMoa1409002
33. Passamonti F, Griesshammer M, Palandri F, Egyed M, Benevolo G, Devos T, et al. Ruxolitinib for the treatment of inadequately controlled polycythemia vera without splenomegaly (RESPONSE-2): a randomised, open-label, phase 3b study. *Lancet Oncol*. (2017) 18:88–99. doi: 10.1016/S1470-2045(16)30558-7
34. Verstovsek S, Vannucchi AM, Griesshammer M, Masszi T, Durrant S, Passamonti F, et al. Ruxolitinib versus best available therapy in patients with polycythemia vera: 80-week follow-up from the RESPONSE trial. *Haematologica*. (2016) 101:821–9. doi: 10.3324/haematol.2016.143644
35. Griesshammer M, Saydam G, Palandri F, Benevolo G, Egyed M, Callum J, et al. Ruxolitinib for the treatment of inadequately controlled polycythemia vera without splenomegaly: 80-week follow-up from the RESPONSE-2 trial. *Ann Hematol*. (2018) 97:1591–600. doi: 10.1007/s00277-018-3365-y
36. Colomba C, Rubino R, Siracusa L, Lalicata F, Trizzino M, Titone L, et al. Disseminated tuberculosis in a patient treated with a JAK2 selective inhibitor: a case report. *BMC Res Notes*. (2012) 5:552. doi: 10.1186/1756-0500-5-552
37. Goldberg RA, Reichel E, Oshry LJ. Bilateral toxoplasmosis retinitis associated with ruxolitinib. *N Engl J Med*. (2013) 369:681–3. doi: 10.1056/NEJMc1302895
38. Wathes R, Moule S, Milojkovic D. Progressive multifocal leukoencephalopathy associated with ruxolitinib. *N Engl J Med*. (2013) 369:197–8. doi: 10.1056/NEJMc1302135
39. Wysham NG, Sullivan DR, Allada G. An opportunistic infection associated with ruxolitinib, a novel Janus kinase 1,2 inhibitor. *Chest*. (2013) 143:1478–9. doi: 10.1378/chest.12-1604
40. Caocci G, Murgia F, Podda L, Solinas A, Atzeni S, La Nasa G. Reactivation of hepatitis B virus infection following ruxolitinib treatment in a patient with myelofibrosis. *Leukemia*. (2014) 28:225–7. doi: 10.1038/leu.2013.235
41. Palandri F, Polverelli N, Catani L, Vianelli N. Ruxolitinib-associated tuberculosis: a case of successful ruxolitinib rechallenge. *Ann Hematol*. (2015) 94:519–20. doi: 10.1007/s00277-014-2183-0
42. Manduzio P. Ruxolitinib in myelofibrosis: to be or not to be an immune disruptor. *Ther Clin Risk Manag*. (2017) 13:169–77. doi: 10.2147/TCRM.S121683
43. Heine A, Brossart P, Wolf D. Ruxolitinib is a potent immunosuppressive compound: is it time for anti-infective prophylaxis? *Blood*. (2013) 122:3843–4. doi: 10.1182/blood-2013-10-531103

44. Polverelli N, Palumbo GA, Binotto G, Abruzzese E, Benevolo G, Bergamaschi M, et al. Epidemiology, outcome, and risk factors for infectious complications in myelofibrosis patients receiving ruxolitinib: a multicenter study on 446 patients. *Hematol Oncol.* (2018) 36:561–9. doi: 10.1002/hon.2509
45. Cervantes F, Dupriez B, Pereira A, Passamonti F, Reilly JT, Morra E, et al. New prognostic scoring system for primary myelofibrosis based on a study of the international working group for myelofibrosis research and treatment. *Blood.* (2009) 113:2895–901. doi: 10.1182/blood-2008-07-170449
46. Tefferi A, Pardanani A. Serious adverse events during ruxolitinib treatment discontinuation in patients with myelofibrosis. *Mayo Clin Proc.* (2011) 86:1188–91. doi: 10.4065/mcp.2011.0518
47. Bhatt VR, Bociek RG, Yuan J, Fu K, Greiner TC, Dave BJ, et al. Leukemic diffuse large B-cell lymphoma in a patient with myeloproliferative disorder. *J Natl Compr Canc Netw.* (2015) 13:281–7. doi: 10.6004/jnccn.2015.0039
48. Pemmaraju N, Kantarjian H, Nastoupil L, Dupuis M, Zhou L, Pierce S, et al. Characteristics of patients with myeloproliferative neoplasms with lymphoma, with or without JAK inhibitor therapy. *Blood.* (2019) 133:2348–51. doi: 10.1182/blood-2019-01-897637
49. Blechman AB, Cabell CE, Weinberger CH, Duckworth A, Leitenberger JJ, Zwald FO, et al. Aggressive skin cancers occurring in patients treated with the Janus kinase inhibitor ruxolitinib. *J Drugs Dermatol.* (2017) 16:508–11. Available online at: <https://jddonline.com/articles/dermatology/S1545961617P0508X/>
50. Ghoreschi K, Gadina M. Jakpot! New small molecules in autoimmune and inflammatory diseases. *Exp Dermatol.* (2014) 23:7–11. doi: 10.1111/exd.12265
51. Schwartz DM, Kanno Y, Villarino A, Ward M, Gadina M, O’Shea JJ. JAK inhibition as a therapeutic strategy for immune and inflammatory diseases. *Nat Rev Drug Discov.* (2017) 16:843–62. doi: 10.1038/nrd.2017.201
52. Punwani N, Scherle P, Flores R, Shi J, Liang J, Yeleswaram S, et al. Preliminary clinical activity of a topical JAK1/2 inhibitor in the treatment of psoriasis. *J Am Acad Dermatol.* (2012) 67:658–64. doi: 10.1016/j.jaad.2011.12.018
53. Craiglow BG, Tavares D, King BA. Topical ruxolitinib for the treatment of alopecia universalis. *JAMA Dermatol.* (2016) 152:490–1. doi: 10.1001/jamadermatol.2015.4445
54. Im A, Hakim FT, Pavletic SZ. Novel targets in the treatment of chronic graft-versus-host disease. *Leukemia.* (2017) 31:543–54. doi: 10.1038/leu.2016.367
55. Janeway CA, Travers P, Walport M, Shlomchik M. *Immunobiology: The Immune System in Health and Disease*. 5th ed. New York, NY: Garland Science (2001).
56. Schönberg K, Rudolph J, Vonnahme M, Parampalli Yajnanarayana S, Cornez I, Hejazi M, et al. JAK inhibition impairs NK cell function in myeloproliferative neoplasms. *Cancer Res.* (2015) 75:2187–99. doi: 10.1158/0008-5472.CAN-14-3198
57. Curran SA, Shyer JA, St Angelo ET, Talbot LR, Sharma S, Chung DJ, et al. Human dendritic cells mitigate NK-cell dysfunction mediated by nonselective JAK1/2 blockade. *Cancer Immunol Res.* (2017) 5:52–60. doi: 10.1158/2326-6066.CIR-16-0233
58. Ankathatti Munegowda M, Hu J. Transient blocking of NK cell function with small molecule inhibitors for helper dependant adenoviral vector-mediated gene delivery. *Cell Biosci.* (2015) 5:29. doi: 10.1186/s13578-015-0023-0
59. Heine A, Held SA, Daecke SN, Wallner S, Yajnanarayana SP, Kurts C, et al. The JAK-inhibitor ruxolitinib impairs dendritic cell function *in vitro* and *in vivo*. *Blood.* (2013) 122:1192–202. doi: 10.1182/blood-2013-03-484642
60. Rudolph J, Heine A, Quast T, Kolanus W, Trebicka J, Brossart P, et al. The JAK inhibitor ruxolitinib impairs dendritic cell migration via off-target inhibition of ROCK. *Leukemia.* (2016) 30:2119–23. doi: 10.1038/leu.2016.155
61. Parampalli Yajnanarayana S, Stübig T, Cornez I, Alchalby H, Schonberg K, Rudolph J, et al. JAK1/2 inhibition impairs T cell function *in vitro* and in patients with myeloproliferative neoplasms. *Br J Haematol.* (2015) 169:824–33. doi: 10.1111/bjh.13373
62. Wing K, Sakaguchi S. Regulatory T cells exert checks and balances on self tolerance and autoimmunity. *Nat Immunol.* (2010) 11:7–13. doi: 10.1038/ni.1818
63. Massa M, Rosti V, Campanelli R, Fois G, Barosi G. Rapid and long-lasting decrease of T-regulatory cells in patients with myelofibrosis treated with ruxolitinib. *Leukemia.* (2014) 28:449–51. doi: 10.1038/leu.2013.296
64. Keohane C, Kordasti SY, Seidl T, Abellan PP, Thomas NSB, Harrison CN, et al. JAK inhibition induces silencing of T helper cytokine secretion and a profound reduction in T regulatory cells. *BJM.* (2015) 171:60–73. doi: 10.1111/bjh.13519
65. Delhommeau F, Dupont S, Tonetti C, Masse A, Godin I, Le Couedic JP, et al. Evidence that the JAK2 G1849T (V617F) mutation occurs in a lymphomyeloid progenitor in polycythemia vera and idiopathic myelofibrosis. *Blood.* (2007) 109:71–7. doi: 10.1182/blood-2006-03-007146
66. Larsen TS, Christensen JH, Hasselbalch HC, Pallisgaard N. The JAK2 V617F mutation involves B- and T-lymphocyte lineages in a subgroup of patients with Philadelphia-chromosome negative chronic myeloproliferative disorders. *Br J Haematol.* (2007) 136:745–51. doi: 10.1111/j.1365-2141.2007.06497.x
67. Giroux M, Delisle JS, Gauthier SD, Heinonen KM, Hinsinger J, Houde B, et al. SMAD3 prevents graft-versus-host disease by restraining Th1 differentiation and granulocyte-mediated tissue damage. *Blood.* (2011) 117:1734–44. doi: 10.1182/blood-2010-05-287649
68. Schwab L, Goroncy L, Palaniyandi S, Gautam S, Triantafyllopoulou A, Mocsai A, et al. Neutrophil granulocytes recruited upon translocation of intestinal bacteria enhance graft-versus-host disease via tissue damage. *Nat Med.* (2014) 20:648–54. doi: 10.1038/nm.3517
69. Febvre-James M, Lecureur V, Augagneur Y, Mayati A, Fardel O. Repression of interferon beta-regulated cytokines by the JAK1/2 inhibitor ruxolitinib in inflammatory human macrophages. *Int Immunopharmacol.* (2018) 54:354–65. doi: 10.1016/j.intimp.2017.11.032
70. Pattison MJ, Mackenzie KE, Arthur JS. Inhibition of JAKs in macrophages increases lipopolysaccharide-induced cytokine production by blocking IL-10-mediated feedback. *J Immunol.* (2012) 189:2784–92. doi: 10.4049/jimmunol.1200310
71. Perrillo RP, Gish R, Falck-Ytter YT. American gastroenterological association institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology.* (2015) 148:221–44 e223. doi: 10.1053/j.gastro.2014.10.038

**Conflict of Interest:** EE has participated at advisory boards for Novartis. CB has participated at advisory boards for Abbvie, BMS, Celgene, Incyte, Janssen, Novartis. FP has received honoraria from Novartis. GP has received honoraria from Amgen, Celgene, Novartis and has participated at advisory boards for Celgene, Janssen, Novartis.

The remaining author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Elli, Baratè, Mendicino, Palandri and Palumbo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.