



Provided by the author(s) and University College Dublin Library in accordance with publisher policies. Please cite the published version when available.

Title	Mesenchymal Stem Cells and Osteoarthritis: Remedy or Accomplice?
Authors(s)	Coleman, Cynthia M.; Curtin, Caroline; Barry, Frank P.; et al.
Publication date	2010-10
Publication information	Human Gene Therapy, 21 (10): 1239-1250
Publisher	Mary Ann Liebert
Item record/more information	http://hdl.handle.net/10197/5030
Publisher's statement	This is a copy of an article published in Human Gene Therapy 2010 Mary Ann Liebert, Inc. Human Gene Therapy is available online at: http://www.liebertonline.com .
Publisher's version (DOI)	10.1089/hum.2010.138

Downloaded 2022-08-25T11:36:46Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



Mesenchymal Stem Cells and Osteoarthritis: Remedy or Accomplice?

Cynthia M. Coleman, Caroline Curtin, Frank P. Barry, Cathal O'Flatharta, and J. Mary Murphy

Abstract

Multipotent mesenchymal stromal or stem cells (MSCs) are likely to be agents of connective tissue homeostasis and repair. Because the hallmark of osteoarthritis (OA) is degeneration and failure to repair connective tissues it is compelling to think that these cells have a role to play in OA. Indeed, MSCs have been implicated in the pathogenesis of OA and, in turn, progression of the disease has been shown to be therapeutically modulated by MSCs. This review discusses current knowledge on the potential of both marrow- and local joint-derived MSCs in OA, the mode of action of the cells, and possible effects of the osteoarthritic niche on the function of MSCs. The use of stem cells for repair of isolated cartilage lesions and strategies for modulation of OA using local cell delivery are discussed as well as therapeutic options for the future to recruit and appropriately activate endogenous progenitors and/or locally systemically administered MSCs in the early stages of the disease. The use of gene therapy protocols, particularly as they pertain to modulation of inflammation associated with the osteoarthritic niche, offer an additional option in the treatment of this chronic disease. In summary, elucidation of the etiology of OA and development of technologies to detect early disease, allied to an increased understanding of the role MSCs in aging and OA, should lead to more targeted and efficacious treatments for this debilitating chronic disease in the future.

Osteoarthritis

OSTEoarthritis (OA), a degenerative disease of joints, is the most common musculoskeletal disorder (Buckwalter and Martin, 2006). Assessment of disease burden in the United States, using disability-adjusted life-years as a measure, pinpointed OA as a major cause of premature death and disability (Michaud *et al.*, 2006). Moreover, the societal impact of OA is anticipated to increase exponentially with aging populations accompanied by the rising prevalence of obesity, the principle nongenetic risk factor for the disease (Dunlop *et al.*, 2003). Despite this, the etiology of OA is still unknown. OA is a complex condition of broad pathology with loss of or damage to articular cartilage being consistent elements, accompanied by changes to the subchondral bone and synovium (Findlay, 2010; Goldring and Goldring, 2010). Synovial inflammation in particular can disturb joint homeostasis (Chen and Tuan, 2008; Attur *et al.*, 2010) and is associated with pain and OA disease progression (Scanzello *et al.*, 2008). Current treatments for OA are not regenerative and have little impact on the progressive degeneration of joint tissues. Clinical interventions are primarily symptomatic with a focus on pain reduction and

control of inflammation with nonsteroidal antiinflammatory drugs and ultimately through total joint replacement (Buckwalter *et al.*, 2004).

Stem Cells

Stem cells represent an important element in regenerative strategies for tissue repair by virtue of their availability in large numbers and relative ease of preparation and delivery.

Embryonic stem cells (ESCs), although pluripotent with the potential to address regeneration of tissues in all germ lines, have ethical issues associated with their use. These issues may be overcome by the use of induced pluripotent stem cells (iPSCs) generated by reprogramming of human somatic cells (Jaenisch and Young, 2008; Nishikawa *et al.*, 2008; Tweedell, 2008). iPSCs have a similar, although not identical, phenotype to ESCs. Reports highlight major differences between both cell lines: ESCs generated from human embryos carrying the fragile X mutation demonstrated silencing of the *FMR1* gene after differentiation, whereas this gene was not activated in iPSCs derived from fibroblasts of affected individuals carrying the mutation despite successful

reprogramming to pluripotency (Urbach *et al.*, 2010). Human iPSCs, although capable of differentiating to hematopoietic derivatives, did so with significantly decreased efficiency compared with ESCs. Furthermore, the iPSC-derived hemangioblasts demonstrated increased apoptosis, limited growth and expansion capability, and rapid onset of senescence (Feng *et al.*, 2010). Clinical use of both cell types may require the development of predifferentiation protocols and depletion of the residual parent population before introduction into patients. Alternatively, there is increasing evidence that most adult tissues have a resident stem or progenitor cell population with a critical role in homeostasis and tissue repair. Adult stem cells have a more restricted differentiation potential than pluripotent stem cells, but are currently used in clinical practice (e.g., hematopoietic stem cells for marrow transplantation) and clinical testing of other adult stem cells, including multipotent mesenchymal stromal or stem cell (MSC) preparations, has progressed rapidly (Ankrum and Karp, 2010).

MSCs were first described by Friedenstein, who identified a subpopulation of cells within the stromal compartment of bone marrow with osteogenic potential (Friedenstein *et al.*, 1966, 1970, 1974). These adherent, fibroblastic-like cells were shown to be capable of forming colonies from a single cell (colony-forming unit fibroblastic [CFU-F]) and had the capacity to form multiple skeletal tissues *in vivo* (Owen and Friedenstein, 1988; Friedenstein, 1995). Subsequently, individual, clonal populations of stem cells in human bone marrow were identified that retained the potential to differentiate into chondrocytes, osteocytes, and adipocytes (Pittenger *et al.*, 1999). Since 1999, MSCs have been isolated and characterized from many other human sources including adipose tissue (Gimble *et al.*, 2007; Meliga *et al.*, 2007; Bunnell *et al.*, 2008), umbilical cord blood, and Wharton's jelly (Weiss and Troyer, 2006; Weiss *et al.*, 2006; Flynn *et al.*, 2007; Troyer and Weiss, 2008). They have the capacity to differentiate into cells of connective tissue lineages, including bone, fat, cartilage, and muscle. Bone marrow-derived MSCs have an additional potential in providing the stromal support system for hematopoietic stem cells (Barry and Murphy, 2004; Delorme and Charbord, 2007; Bianco *et al.*, 2008). It has also been suggested that stromal cells derived from bone marrow may be capable of vascular smooth muscle differentiation in long-term cultures (Dennis and Charbord, 2002) and cells with MSC characteristics isolated from arteries and microvessels have been described as multipotential "pericyte-like" stem cells (Mody *et al.*, 2003; Tintut *et al.*, 2003; Abedin *et al.*, 2004; Tavian *et al.*, 2005; da Silva Meirelles *et al.*, 2006). These cells have the capacity to differentiate to multiple lineages (Shi and Gronthos, 2003; Tintut *et al.*, 2003) and various reports have suggested a perivascular niche as the source of MSCs derived from bone marrow, skeletal muscle, brain, and fat (Shi and Gronthos, 2003; Tavian and Peault, 2005; Sacchetti *et al.*, 2007; Zannettino *et al.*, 2008). Prospective isolation techniques identified progenitor cells from blood vessels in various tissues including skeletal muscle and adipose tissue that exhibited multipotentiality at a clonal level and expressed MSC markers in both cultured and noncultured cells. Perivascular cells were proposed as the precursors for MSCs and other adult stem cells (Crisan *et al.*, 2008), leading to the hypothesis that all MSCs are pericytes (Caplan, 2008).

MSCs from joint tissues

Isolation of MSCs from joint tissue was first reported in 2001 when multipotent mesenchymal cells with the capacity for chondrogenesis, osteogenesis, adipogenesis, and sporadic myogenesis were grown from a digest of adult human synovial membrane (De Bari *et al.*, 2001b). As for marrow-derived MSCs, clonal heterogeneity has been demonstrated for MSCs isolated from synovium with progenitors of varying proliferative capacity and distinct mesenchymal differentiation potency described (Karystinou *et al.*, 2009). Interestingly, we have shown that marrow-derived MSCs, whether injected into a healthy or injured joint, engrafted primarily to synovial tissue or perichondrium with no cells detected at the surface of normal or damaged articular cartilage (Murphy *et al.*, 2003). De Bari and coworkers also demonstrated *in vivo* myogenesis with synovium-derived MSCs contributing to myofibers and functional satellite cells in regenerating nude mouse muscle (De Bari *et al.*, 2003). However, a more recent report suggested that synovium-derived MSCs had a limited capacity to undergo myogenic differentiation *in vitro* or to contribute to muscle regeneration *in vivo* (Meng *et al.*, 2010). MSCs have also been isolated from normal and early osteoarthritic synovial fluid; these authors acknowledged that the source of the cells was likely degrading synovium. Interestingly, in comparison with matched bone marrow-derived cells, synovial fluid cells showed greater clonogenicity and chondrogenic differentiation capacity (Jones *et al.*, 2008). In general, synovium-derived MSCs seem to have a more chondrogenic phenotype than those derived from bone marrow or infrapatellar fat pad (Sakaguchi *et al.*, 2005; Lee *et al.*, 2010).

Progenitor cells at the surface of healthy articular cartilage were originally described in bovine tissue (Dowthwaite *et al.*, 2004). When prepared as clonal populations these cells had greater growth potential and higher telomerase activity than dedifferentiated chondrocytes isolated from the same source (Khan *et al.*, 2009). Mesenchymal-like cells with progenitor characteristics were subsequently identified in normal and osteoarthritic human articular cartilage (Alsalameh *et al.*, 2004; Fickert *et al.*, 2004) and were shown to undergo chondrogenic and osteogenic, but not adipogenic, differentiation (Grogan *et al.*, 2009). Pacifici and co-workers have linked *Gdf5* expression with progenitor potential by crossing a ROSA-LacZ-reporter with *Gdf5-Cre* mice to track events at prospective joint sites. LacZ-positive cells initially constituted the interzone and persisted in joint-forming areas throughout development. These progenitor cells formed articular cartilage and synovial lining as well as other joint tissues. However, they contributed minimally if at all to underlying growth plate cartilage and bone growth, indicating that a population of *Gdf5*-expressing mesenchymal progenitors uniquely contributed to articular cartilage (Koyama *et al.*, 2008).

Mode of action

MSCs remain at the forefront of current translational efforts in cellular therapy for a broad spectrum of diseases. However, some controversy exists as to the primary effect of the cells in the injured environment: do MSCs undergo tissue-specific differentiation or act in a paracrine manner to produce soluble reparative factors, or is the therapeutic effect

a consequence of both mechanisms? Data suggest that the therapeutic potential of these cells, at least in some applications, is related to paracrine effects such as the release of factors that (1) modulate the immune response, (2) mobilize or promote host cell survival, (3) recruit and induce mitosis of endogenous tissue progenitor cells at the site of injury while stimulating an angiogenic response, or (4) prevent an inappropriate fibrotic response (for reviews see Caplan, 2009; Oh *et al.*, 2010). However, studies in the area of cardiac repair question the concept that MSCs act solely through paracrine mechanisms. A study in pigs using allogeneic MSCs for the treatment of chronic ischemic cardiomyopathy suggested that long-term engraftment and trilineage differentiation to cardiomyocytes, vascular smooth muscle cells, and endothelial cells contributed at least in part to an improvement in cardiac function (Quevedo *et al.*, 2009). However, the trilineage differentiation shown was minimal.

Immunomodulatory effects

Factors produced by MSCs in response to the inflammatory environment include interleukin (IL)-10, IL-1 receptor antagonist (IL-1Ra), and transforming growth factor (TGF)- β (Pittenger, 2009). Indeed, the establishment of an immunosuppressive local milieu has been proposed as a major factor in the immune privilege of MSCs (English *et al.*, 2008, 2009). There is evidence that hematopoietic cells may provide the "licensing" signal for MSCs (English *et al.*, 2007; Polchert *et al.*, 2008) to deliver immunosuppressive signals including IL-10 and prostaglandin E₂ (PGE₂). Upregulation of IL-10 in the brain has been associated with the therapeutic potential of MSCs in reducing neuronal injury after an ischemic injury in rats (Liu *et al.*, 2009) and increased IL-10 production by host macrophages in response to MSCs has been described to occur through a PGE₂-dependent mechanism in treating sepsis in mice (Németh *et al.*, 2009). These results and others have led to studies assessing gene therapeutic strategies. In particular, systemic administration of genetically modified MSCs overexpressing IL-10 in a model of experimental arthritis resulted in the inhibition of symptoms through suppression of the autoimmune response as well as the production of cytokines such as IL-4 (Choi *et al.*, 2008).

Paracrine effects

The beneficial paracrine effects of MSCs have been demonstrated in the articular joint via administration of cells after surgically induced injury. Local delivery of autologous caprine MSCs in a solution of hyaluronan to a meniscectomized stifle joint resulted in significantly increased regeneration of meniscal tissue and chondroprotection when compared with meniscectomized joints treated with hyaluronan alone (Murphy *et al.*, 2003). However, it was readily apparent that the green fluorescent protein (GFP)-labeled MSCs used in the study colonized just a small proportion of the regenerated meniscus. It was therefore concluded that the implanted MSCs induced a host repair response through the release of paracrine factors to replace the resected medial meniscus. Similarly, a model involving subcutaneous implantation of hydroxyapatite scaffolds loaded with MSCs into syngeneic, allogeneic, or immunocompromised mice was used to assess the origin of bone routinely found in the scaffold after an

extended period *in vivo*. Interestingly, no bone was formed in the allogeneic setting as a result of rapid destruction of the cells. However, in syngeneic and immunocompromised recipients the implanted cells were found to be pivotal to bone formation, but tissue formation was dependent on recruited recipient osteoprogenitors (Tasso *et al.*, 2009). The accumulation of human MSCs has also been associated with an increased number of oligodendrocytes in lesion areas of mice with experimental allergic encephalomyelitis (Bai *et al.*, 2009).

The antifibrotic action of MSCs was first described in a bleomycin-induced lung injury model in which murine MSCs, administered after exposure to the bleomycin, homed to the injured lung and reduced not only the inflammatory response but also collagen deposition (Ortiz *et al.*, 2003). IL-1Ra, secreted by the MSCs, was found to be integral to this process and acted by blocking tumor necrosis factor (TNF)- α and IL-1 in the lung (Ortiz *et al.*, 2007). An immunosuppressive mechanism of action for MSCs was also demonstrated in carbon tetrachloride-induced liver injury resulting in suppressed fibrosis, an effect specific for MSCs as administration of hematopoietic stem cells resulted in acute inflammation and subsequent fibrosis (Pulavendran *et al.*, 2010).

Stimulation of angiogenesis

Increased angiogenesis has been shown to occur in late OA. MSCs can promote early angiogenic events, a mechanism known to contribute to tissue repair, by increasing endothelial cell (EC) proliferation and migration *in vitro* as well as significantly increasing the stability of vessels formed by ECs through a cell contact-mediated mechanism (Duffy *et al.*, 2009). Similarly, MSC differentiation to endothelial and smooth muscle cells was found to occur in support of new blood vessel formation in the kidney (J. Chen *et al.*, 2008). Other studies support a paracrine mechanism for MSC reconstitution of the microcirculation or collateral perfusion (Kinnaird *et al.*, 2004; Ladage *et al.*, 2007) through molecules such as secreted frizzled-related protein-1 or Cyr61 (Dufourcq *et al.*, 2008; Estrada *et al.*, 2009).

In contrast to the paradigm of MSCs influencing healing or repair through paracrine mechanisms, early tissue engineering principles of cell therapy were based on the ability of the delivered cells to engraft, differentiate, and contribute to the formation of repair tissue. Differentiation remains an important avenue of research for tissue engineering applications and MSCs remain at the forefront of current translational efforts in cellular therapy for a broad spectrum of diseases. Because our understanding of the mode of action of MSCs in specific applications is incomplete, it is not possible to rule out cell differentiation, whether overt as may be the case for connective tissue applications or minimal in other scenarios, as a factor in therapeutic efficacy.

MSCs in osteoarthritis

Inherent differences between MSCs derived from healthy or osteoarthritic bone marrow have been well documented. A number of years ago MSCs isolated from the marrow of patients undergoing joint replacement surgery were compared with cells from healthy, age-matched control subjects with no evidence of OA. The proliferative rates of MSCs

derived from osteoarthritic marrow were significantly reduced compared with healthy controls. Furthermore, the differentiation profile of the osteoarthritic MSCs was altered, with reduced chondrogenic and adipogenic activity but increased osteogenesis (Murphy *et al.*, 2002). These changes were systemic in that osteoarthritic MSCs isolated from iliac crest or tibial and femoral components were comparable. Although these cells were cultured in selected fetal calf serum without supplemental growth factors, studies using growth factor-complemented medium for the establishment and growth of MSC populations did not show decreased proliferation or chondrogenesis by cells isolated from the patients with OA (Scharstuhl *et al.*, 2007), although it must be noted that quantitation of proteoglycan deposition per cell was not performed in this study. Indeed, the continuing controversy about the effects of age and disease on proliferation and functional properties of MSC populations may be attributed to culture conditions and the inability to recapitulate either the aging or indeed the osteoarthritic niche *in vitro*. For example, human periosteal MSCs have spontaneous chondrogenic activity in cultures from donors younger than 30 years that is lost with passaging and absent from older donors (De Bari *et al.*, 2001a), a deficiency overcome with fibroblast growth factor (FGF)-2 supplementation (Im *et al.*, 2006).

Synovial fluid, the milieu to which cells within the joint are exposed, changes dramatically with OA onset (for review see Goldring and Goldring, 2007). Chondrocytes, synovial cells, and perhaps even joint progenitor cells contribute and respond to cytokines, chemokines, and antioxidant levels in the diseased synovial fluid, resulting in oxidative stress. For example, the inflammatory mediators IL-1 β and TNF- α are increased in OA synovial fluid and have dramatic effects on chondrogenesis of stem cells. Figure 1 shows the effect of 5% osteoarthritic synovial fluid on chondrogenesis of goat bone marrow MSCs. Chondrogenesis induced by TGF- β_3 was significantly inhibited by the addition of OA synovial fluid

and preexposure of the early micromass culture to the fluid before addition of TGF- β_3 had an even more dramatic effect. Inhibition of Sox9, the putative master chondrogenic transcription factor, by inflammatory cytokines present in synovial fluid was first described in 2000 (Murakami *et al.*, 2000). More recently, the inhibitory effect of the cytokines on MSC chondrogenesis was shown to be mediated through NF- κ B-dependent pathways (Wehling *et al.*, 2009). Interestingly, the NF- κ B pathway was also implicated in inflammation-associated responses of fibroblast-like synoviocytes (FLSs) in rheumatoid arthritic pannus. A major proportion of FLSs was shown to be marrow-derived, with the capacity to differentiate to mesenchymal lineages. These researchers showed increased proliferation and repression of the osteogenic and adipogenic differentiation of FLSs with inhibition of NF- κ B, and proposed that arthritic FLSs were MSCs with arrested differentiation in response to the arthritic inflammatory environment (Li and Makarov, 2006).

Aberrant progenitor function has also been suggested as pertinent to the osteoarthritic changes that occur in articular cartilage. Notch-1-positive cartilage progenitor cells have been isolated from immature cartilage and a number of groups have described an increased incidence of these cells in osteoarthritic cartilage (Alsalamah *et al.*, 2004; Fickert *et al.*, 2004; Hiraoka *et al.*, 2006). However, one study found that more than 45% of cells in normal and osteoarthritic cartilage were positive for the putative progenitor markers Notch-1, Stro-1, and vascular cell adhesion molecule (VCAM)-1. In contrast, the progenitor cartilage side population isolated by fluorescence-activated cell sorting (FACS) represented just 0.14% of the cellular complement in both tissues and this population had chondrogenic and osteogenic capacity. These data led to the suggestion that the markers may not be useful for identification of cartilage progenitors but rather contribute to the abnormal cell activation and differentiation process characteristic of OA (Grogan *et al.*, 2009).




FIG. 1. Potential effects of the osteoarthritic joint niche on chondrogenesis of MSCs. MSCs, isolated from goat bone marrow, were placed in pellet culture and induced to differentiate in the presence of TGF- β_3 for 7 days (*top*) and 14 days (*bottom*) as described previously (Murphy *et al.*, 2003). Synovial fluid was harvested from osteoarthritic goat knees and maintained at 5% (v/v) throughout the culture with 10-ng/ml TGF- β_3 (**B** and **E**) or added to the cultures for 2 days before addition of growth factor (**C** and **F**). Control pellets were treated with 10-ng/ml TGF- β_3 (**A** and **D**) for 7 and 14 days, respectively. Original magnification, $\times 40$. Color images available online at www.liebertonline.com/hum.

Intriguingly, migratory cells not normally found in healthy articular cartilage add to the complexity of progenitor populations in degenerating cartilage. A clonal, multipotent population in repair tissue associated with late-stage OA (Koelling *et al.*, 2009) was hypothesized to migrate from blood vessels occupying fissures or breaks in the tidemark of vascularized cartilage tissue. Although maintaining an osteochondroprogenitor phenotype, they do not differentiate into fully committed chondrocytes *in situ*, but instead are possibly restricted to a fibrocartilaginous phenotype providing temporary repair due to the absence of appropriate differentiation cues (Khan *et al.*, 2009).

MSCs may also play a role in osteophyte formation, perhaps another attempt at repair to counteract the altered architecture of the injured or OA joint. Osteophytes are thought to be derived from precursors in the periosteum or perichondrium in response to growth factors such as TGF- β and develop through endochondral ossification (van der Kraan and van den Berg, 2007). Mice with a targeted disruption of mitogen-inducible gene 6 develop early-onset OA characterized by significant joint enlargement and deformity associated with osteophyte formation. Furthermore, proliferation of mesenchymal progenitors followed by chondrogenesis was shown to be associated with the osteophyte formation (Zhang *et al.*, 2005).

Stem Cell Therapy for Osteoarthritis

Maintenance or restoration of a fully functional joint with biomechanically stable articular cartilage remains the holy grail of therapeutic or regenerative strategies in OA. In reality, complete cartilage degeneration requires total joint replacement through artificial implants and clinical intervention for cartilage repair is beneficial only when focal lesions are small enough to be reconstituted with cells or reparative tissue from various sources (Redman *et al.*, 2005). Operative interventions for the treatment of cartilage lesions include the transplantation of osteochondral grafts (mosaicplasty), microfracture, and autologous chondrocyte implantation (ACI) with or without a scaffold matrix (MACI; matrix-induced autologous chondrocyte implantation) to deliver reparative cells (Bentley *et al.*, 2003; Hangody and Fules, 2003; Knutsen *et al.*, 2004; Bartlett *et al.*, 2005). A major limitation of all these strategies is the inability to treat large defects (Steinert *et al.*, 2007), thus excluding patients with OA.

Much of the early work on the preclinical evaluation of MSCs in musculoskeletal applications also focused on repair of articular cartilage and bone. Cells were delivered to defects in cartilage or bone in procedures requiring the application of a cell-loaded three-dimensional solid scaffold implanted through an open surgical procedure. As early as 1994, researchers described repair of defects in articular cartilage in rabbits (Wakitani *et al.*, 1994) and humans (Wakitani *et al.*, 2002) and segmental-defect bone repair (Bruder *et al.*, 1998). The first human trial assessing the ability of MSCs to promote repair a large musculoskeletal defect was reported in 2001 with successful healing of a large bone defect (Quarto *et al.*, 2001). This group performed a 6- to 7-year follow-up on four patients and reported integration of implants and no evidence of late fractures in the implant zone, indicating long-term durability of this tissue engi-

neering approach with autologous bone marrow MSCs loaded on a macroporous bioceramic scaffold (Marcacci *et al.*, 2007).

Efforts have focused on the use of MSCs as a therapy for traumatic OA. Murphy and colleagues have used a somewhat simpler scaffold-free approach for the treatment of OA associated with meniscal injury. Delivery of autologous MSCs, retrovirally transduced to express green fluorescent protein, to caprine joints subjected to total meniscectomy and resection of the anterior cruciate ligament resulted in regeneration of meniscal tissue and significant chondroprotection (Murphy *et al.*, 2003). Evidence of cell engraftment to the synovium and the perichondrium as well as the surface lining of other joint tissues including the lateral meniscus and fat pad was noted. Although a significant number of labeled cells was detected at the surface of regenerated meniscus, and to a much lesser extent embedded in the repair tissue, the primary reparative response was host-derived and presumably resulted from interaction between the implanted stem cells and synoviocytes or endogenous progenitors at the site of injury. MSCs derived from synovium have been similarly used to treat meniscal defects. These cells adhered to meniscal lesions and differentiated into meniscal or collagen type II-expressing fibrocartilage cells. In one study the cells promoted meniscal regeneration (Horie *et al.*, 2009) but no therapeutic benefit was found in an earlier study (Mizuno *et al.*, 2008). The same strategy of direct intraarticular injection of MSCs resulted in improved repair of chondral defects in a pig model (Lee *et al.*, 2007).

Delivery of allogeneic MSCs to the destabilized joint after total or partial meniscectomy also resulted in protection against the development of osteoarthritic symptoms (our unpublished results). However, host protection by transplanted MSCs in the OA models was not complete and all treated joints displayed some degree of damage to the articular cartilage.

Stem cell targeting

Adequate tissue repair strategies may require specific cellular targeting to the site of injury as retention and engraftment of transplanted cells are inadequate. Although cartilage has an easily accessible repertoire of chondrogenic progenitor cells that are capable of mitotic division and chondrogenic differentiation (Dowthwaite *et al.*, 2004; Koelling *et al.*, 2009), it is incapable of self-regeneration in adults (Wei and Messner, 1999). Conceivably, these progenitor cells that localize to the surface zone in normal cartilage are compromised or lost early in the osteoarthritic process. We therefore proposed that targeted delivery of progenitor cells in early OA may increase MSC engraftment and impact progression of the osteoarthritic process. Although no studies have been performed on cell targeting to osteoarthritic cartilage to date, prechondrocytes have been targeted to an osteochondral defect in a rabbit model using "cell-painting" techniques (Dennis *et al.*, 2004) whereby the cells were coated with lipidated protein G to enable attachment of selected antibodies that could in turn bind specific cartilage matrix molecules. In another study, CD44 present on MSCs was enzymatically converted to confer potent E-selectin/L-selectin-like binding affinity to enhance targeting of MSCs to




FIG. 2. Targeting of MSCs to human osteoarthritic cartilage. **(A)** Fluorescence microscopy analysis of a human osteoarthritic cartilage explant exposed to 1×10^6 hMSCs/ml of Cell Tracker red and DAPI-stained hMSCs for 20 min with agitation. Scale bar: $60 \mu\text{m}$. **(B)** Hematoxylin-stained osteoarthritic cartilage explant treated as in **(A)** for hMSC binding for 20 min and cultured for 14 days in serum-free chondrogenic medium containing TGF- β_3 . Arrows indicate MSCs embedded in the resurfaced cartilage. Scale bar: $100 \mu\text{m}$. Color images available online at www.liebertonline.com/hum.

bone marrow (Sackstein *et al.*, 2008). Although MSCs introduced into the normal or osteoarthritic joint do not engraft to the cartilage surface (Murphy *et al.*, 2003), these cells will attach and populate deep fissures of fibrillated cartilage after a 20-min incubation with gentle agitation, using an *ex vivo* cartilage explant model. The attached MSCs will ultimately form a regenerated surface after exposure to chondrogenic differentiation medium (Fig. 2).

Gene Therapy for Osteoarthritis

As OA is not a systemic disorder, but is instead limited to enclosed joints, it is uniquely suited to therapeutic interventions delivered via gene therapy. Gene delivery therefore offers a method for localized, continued overexpression of a therapeutic agent with the aim of suppressing OA-associated chondrocyte apoptosis, supporting cell viability and stimulating the deposition of a healthy cartilage-like extracellular matrix (ECM) containing collagen type II and sulfated glycosaminoglycans (GAGs).

Genetic modification of therapeutic cells, or of host tissue directly, to continually overexpress anabolic factors may support the adaptation of therapeutic progenitor cells to a chondrocyte-like phenotype, as well as enhanced deposition of a cartilaginous ECM. For example, virally induced expression of Sox9 alone (Tew *et al.*, 2005; Cucchiari *et al.*, 2007), or in combination with Sox5 and Sox6 (Ikeda *et al.*, 2004), resulted in the induction of a chondrogenic phenotype, even in OA chondrocytes or in nonchondrogenic cell lines. Further, coexpression of Sox5, Sox6, and Sox9 will suppress the hypertrophic or osteogenic differentiation of transformed cells, offering an enhancement over direct cell therapy for modulation of OA. Similarly, the introduction of β -1,3-glucuronosyltransferase 1, a GAG-synthesizing enzyme, into chondrocytes has been demonstrated to stimulate an abundance of GAG production even in the presence of IL-1 β -induced proteoglycan depletion (Venkatesan *et al.*, 2004).

Stimulation of cellular proliferation by viral overexpression of mitogenic factors, or suppression of cell death via overexpression of antiapoptotic factors, may both inhibit OA disease progression as well as stabilize the articular

tissue. Viral overexpression of FGF-2 in chondrocytes has been demonstrated to stimulate proliferation *in vitro* as well as improved tissue repair when transfected cells are delivered *in vivo* (Cucchiari *et al.*, 2005; Yokoo *et al.*, 2005; Kaul *et al.*, 2006). Further, the combination of FGF-2 and Sox9 overexpression in human OA chondrocytes stimulated mitogenesis, stimulated GAG and collagen type II deposition, and inhibited cellular hypertrophy (Cucchiari *et al.*, 2009). Alternatively, the overexpression of antiapoptotic Bcl-2 (Surendran *et al.*, 2006), or the serine proteinase inhibitor kallistatin, offers a means by which native chondrocytes can be protected from OA-induced cell death (Wang *et al.*, 2005; Hsieh *et al.*, 2009).

As the TGF- β superfamily is paramount for the development of cartilage and supports cartilage-like ECM deposition by chondrocytes, it is only logical to genetically overexpress TGF- β superfamily members in an effort to induce the secretion of a hyaline-associated matrix. Several alternative methodologies to overexpress TGF- β_1 have been investigated including light-activated gene transduction (Ito *et al.*, 2004) or Lipofectamine (Guo *et al.*, 2007). These studies resulted in successful protein overexpression, the enhancement of cartilaginous ECM protein production, and enhanced chondral repair in conjunction with the suppression of the cartilage-degrading matrix metalloproteinases MMP-1 and -3 (Guo *et al.*, 2007). Adeno-associated virus (AAV)-based overexpression of TGF- β_1 in MSCs (Pagnotto *et al.*, 2007) or healthy or OA chondrocytes (Ulrich-Vinther *et al.*, 2005) also resulted in the synthesis of type II collagen and aggrecan, the suppression of MMP-3 (Ulrich-Vinther *et al.*, 2005), and the repair of osteochondral defects (Pagnotto *et al.*, 2007). Similar results were obtained by retroviral (Lee *et al.*, 2005) or adenoviral vector expression, whereby proteoglycan synthesis was enhanced with TGF- β_1 alone (Blaney Davidson *et al.*, 2007) or in combination with insulin-like growth factor (IGF) overexpression (Smith *et al.*, 2000). Similarly, production of bone morphogenetic protein (BMP)-7, a member of the TGF- β superfamily, by adenovirus in bovine chondrocytes (Hidaka *et al.*, 2001) or by retrovirus in MSCs (Mason *et al.*, 1998) has been demonstrated to result in enhanced GAG and type II collagen secretion as well as the suppression of chondrocyte hypertrophy (Smith *et al.*,

2000; Hidaka *et al.*, 2001), resulting in bone and articular cartilage regeneration in osteochondral defects (Mason *et al.*, 1998).

Modification of the inflammatory cascade in OA has been extensively investigated through genetic modification. Suppression of NF- κ B via adenoviral overexpression of small interfering RNA (siRNA) in the synovium and cartilage has been demonstrated to result in the inhibition of synovial inflammation and cartilage degradation by suppressing Cox-2, nitric oxide synthase (NOS)-2, and MMP-9 in IL-1 β - and TNF- α -induced arthritis (Lianxu *et al.*, 2006; L.X. Chen *et al.*, 2008). A similar inhibition of inflammation was observed after IL-4 overexpression in a canine model, resulting in suppression of inflammatory mediators prostaglandins and MMPs (Rachakonda *et al.*, 2008).

Adenoviral overexpression of IGF-1 has been demonstrated to be safe in an equine model when directly administered into the joint (Goodrich *et al.*, 2006), making it an attractive vector for OA human gene therapy. Transfection of articular chondrocytes with AdIGF-1 results in cartilage ECM gene and protein expression, as well as the beneficial inhibition of chondrocyte dedifferentiation (Nixon *et al.*, 2000; Brower-Toland *et al.*, 2001). In addition, AdIGF-1-transfected chondrocytes, when administered intraarticularly, contribute to improved cartilage morphology and the deposition of hyaline-like tissue in cartilaginous lesions (Goodrich *et al.*, 2007). Simple plasmid-based overexpression of IGF-1 in rabbit chondrocytes, when encapsulated in alginate and delivered *in vivo*, results in improved cartilage repair and accelerated subchondral bone formation in osteochondral defects (Madry *et al.*, 2005).

By far, the most extensively investigated mechanism for regulating inflammation is the overexpression of IL-1Ra to inhibit IL-1-mediated OA inflammation. Overexpression of IL-1Ra by retrovirus in equine, canine, or lapine synovial tissues regularly results in reduced lesion severity or cartilage degradation (Pelletier *et al.*, 1997; Frisbie and McIlwraith, 2000; Zhang *et al.*, 2004) in the setting of OA, an effect enhanced by the coadministration of IL-10 (Zhang *et al.*, 2004). Similarly, AAV-based overexpression of IL-1Ra has been demonstrated to reduce OA-associated lameness and to improve cartilage morphology in an equine model (Frisbie *et al.*, 2002). Adenoviral overexpression of IL-1Ra similarly suppresses inflammation (Kay *et al.*, 2009) and protects OA chondrocytes from IL-1-induced GAG degradation (Baragi *et al.*, 1995). Alternatively, delivery of the IL-1Ra gene via chitosan nanoparticles (Zhang *et al.*, 2006) or liposomes (Fernandes *et al.*, 1999) demonstrated a similar suppression of cartilage lesion development without the use of a viral vector. The delivery of IL-1Ra via a gene-based mechanism was shown to be beneficial over treatment with recombinant protein (Gouze *et al.*, 2003), possibly due to sustained protein expression.

The coadministration of IGF-1- and IL-1Ra-containing vectors offers a unique proanabolic response while suppressing catabolic mediators where the suppression of IL-1 α , IL-1 β , and associated MMPs by IL-1Ra, is accompanied by IGF-1-stimulated type II collagen and GAG deposition (Haupt *et al.*, 2005). In the setting of a microfractured full-thickness chondral defect, the coadministration of AdIL-1Ra and AdIGF-1 demonstrated increased type II collagen and proteoglycan deposition (Morisset *et al.*, 2007).

Future Considerations

MSCs offer the potential to open new frontiers in the practice of medicine. However, it is critical to increase our understanding of the mechanisms by which these cells impact the progression of OA, or contribute to the pathogenesis of disease, to enable the development of innovative therapeutic options. Other avenues of research that need to be addressed include (1) establishing the optimal conditions for MSC engraftment in the OA joint, (2) identifying the optimal therapeutic cell by comparing chondrocytes, local synovium-derived or circulating marrow-derived MSC populations, and (3) interrogating the underlying mechanisms of action that contribute to attenuation of osteoarthritic symptoms after joint injury. Current cellular therapies in the joint focus on repair of isolated cartilage lesions and do not as yet address the widespread damage associated with OA. With continued improvements in early diagnosis of the disease, it is conceivable that cellular therapies may be applied to resurfacing of minimally damaged cartilage. Progenitor cells have been isolated from the cartilage surface zone, but this layer is lost with the destruction of the cartilage surface in early OA. Strategies that target regeneration of this layer may ultimately reestablish a functional surface zone and delay progression of the disease. It is also conceivable that targeting strategies using nanomaterials could attract either endogenous stem cells or culture-expanded autologous or allogeneic MSCs applied either locally or systemically. Gene therapies also offer some promise particularly in the modulation of inflammatory mediators associated with the osteoarthritic niche. In summary, the convergence of research into the root causes of OA and the role of stem cells in this etiology should lead to more targeted and efficacious treatment for this debilitating disease in the future.

Acknowledgments

This work was funded by Science Foundation Ireland, Centre for Science Engineering and Technology award (CSET) in Regenerative Medicine. The authors gratefully acknowledge the contribution of Mr. William Curtin, Department of Orthopaedic Surgery, Merlin Park Hospital Galway, to this research.

Author Disclosure Statement

The authors have no conflict of interest to disclose.

References

- Abedin, M., Tintut, Y., and Demer, L.L. (2004). Mesenchymal stem cells and the artery wall. *Circ. Res.* 95, 671–676.
- Alsalameh, S., Amin, R., Gemba, T., and Lotz, M. (2004). Identification of mesenchymal progenitor cells in normal and osteoarthritic human articular cartilage. *Arthritis Rheum.* 50, 1522–1532.
- Ankrum, J., and Karp, J.M. (2010). Mesenchymal stem cell therapy: Two steps forward, one step back. *Trends Mol. Med.* 16, 203–209.
- Attur, M., Samuels, J., Krasnokutsky, S., and Abramson, S.B. (2010). Targeting the synovial tissue for treating osteoarthritis (OA): Where is the evidence? *Best Pract. Res. Clin. Rheumatol.* 24, 71–79.

- Bai, L., Lennon, D.P., Eaton, V., Maier, K., Caplan, A.I., Miller, S.D., and Miller, R.H. (2009). Human bone marrow-derived mesenchymal stem cells induce Th2-polarized immune response and promote endogenous repair in animal models of multiple sclerosis. *Glia* 57, 1192–1203.
- Baragi, V.M., Renkiewicz, R.R., Jordan, H., Bonadio, J., Hartman, J.W., and Roessler, B.J. (1995). Transplantation of transduced chondrocytes protects articular cartilage from interleukin 1-induced extracellular matrix degradation. *J. Clin. Invest.* 96, 2454–2460.
- Barry, F.P., and Murphy, J.M. (2004). Mesenchymal stem cells: Clinical applications and biological characterization. *Int. J. Biochem. Cell Biol.* 36, 568–584.
- Bartlett, W., Skinner, J.A., Gooding, C.R., Carrington, R.W., Flanagan, A.M., Briggs, T.W., and Bentley, G. (2005). Autologous chondrocyte implantation versus matrix-induced autologous chondrocyte implantation for osteochondral defects of the knee: A prospective, randomised study. *J. Bone Joint Surg. Br.* 87, 640–645.
- Bentley, G.W., Cook, P.P., Davis, K., Murphy, M.J., and Berding, C.B. (2003). RN to BSN program: Transition from traditional to online delivery. *Nurse Educ.* 28, 121–126.
- Bianco, P., Robey, P.G., and Simmons, P.J. (2008). Mesenchymal stem cells: Revisiting history, concepts, and assays. *Cell Stem Cell* 2, 313–319.
- Blaney Davidson, E.N., van der Kraan, P.M., and van den Berg, W.B. (2007). TGF- β and osteoarthritis. *Osteoarthritis Cartilage* 15, 597–604.
- Brower-Toland, B.D., Saxer, R.A., Goodrich, L.R., Mi, Z., Robbins, P.D., Evans, C.H., and Nixon, A.J. (2001). Direct adenovirus-mediated insulin-like growth factor I gene transfer enhances transplant chondrocyte function. *Hum. Gene Ther.* 12, 117–129.
- Bruder, S.P., Jaiswal, N., Ricalton, N.S., Mosca, J.D., Kraus, K.H., and Kadiyala, S. (1998). Mesenchymal stem cells in osteobiology and applied bone regeneration. *Clin. Orthop. Relat. Res.* (355 Suppl.), S247–S256.
- Buckwalter, J.A., and Martin, J.A. (2006). Osteoarthritis. *Adv. Drug Deliv. Rev.* 58, 150–167.
- Buckwalter, J.A., Saltzman, C., and Brown, T. (2004). The impact of osteoarthritis: Implications for research. *Clin. Orthop. Relat. Res.* (427 Suppl.), S6–S15.
- Bunnell, B.A., Estes, B.T., Guilak, F., and Gimble, J.M. (2008). Differentiation of adipose stem cells. *Methods Mol. Biol.* 456, 155–171.
- Caplan, A.I. (2008). All MSCs are pericytes? *Cell Stem Cell* 3, 229–230.
- Caplan, A.I. (2009). Why are MSCs therapeutic? New data: New insight. *J. Pathol.* 217, 318–324.
- Chen, F.H., and Tuan, R.S. (2008). Mesenchymal stem cells in arthritic diseases. *Arthritis Res. Ther.* 10, 223.
- Chen, J., Park, H.C., Addabbo, F., Ni, J., Pelger, E., Li, H., Plotkin, M., and Goligorsky, M.S. (2008). Kidney-derived mesenchymal stem cells contribute to vasculogenesis, angiogenesis and endothelial repair. *Kidney Int.* 74, 879–889.
- Chen, L.X., Lin, L., Wang, H.J., Wei, X.L., Fu, X., Zhang, J.Y., and Yu, C.L. (2008). Suppression of early experimental osteoarthritis by *in vivo* delivery of the adenoviral vector-mediated NF- κ Bp65-specific siRNA. *Osteoarthritis Cartilage* 16, 174–184.
- Choi, J.J., Yoo, S.A., Park, S.J., Kang, Y.J., Kim, W.U., Oh, I.H., and Cho, C.S. (2008). Mesenchymal stem cells overexpressing interleukin-10 attenuate collagen-induced arthritis in mice. *Clin. Exp. Immunol.* 153, 269–276.
- Crisan, M., Yap, S., Casteilla, L., Chen, C.W., Corselli, M., Park, T.S., Andriolo, G., Sun, B., Zheng, B., Zhang, L., Norotte, C., Teng, P.N., Traas, J., Schugar, R., Deasy, B.M., Badyrak, S., Buhring, H.J., Giacobino, J.P., Lazzari, L., Huard, J., and Péault, B. (2008). A perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell* 3, 301–313.
- Cucchiarini, M., and Madry, H. (2005). Gene therapy for cartilage defects. *J. Gene Med.* 7, 1495–1509.
- Cucchiarini, M., Schetting, S., Terwilliger, E.F., Kohn, D., and Madry, H. (2007). Restoration of the extracellular matrix in human osteoarthritic articular cartilage by overexpression of the transcription factor SOX9. *Arthritis Rheum.* 56, 158–167.
- Cucchiarini, M., Schetting, S., Terwilliger, E.F., Koh, D., and Madry, H. (2009). rAAV-mediated overexpression of FGF-2 promotes cell proliferation, survival, and α -SMA expression in human meniscal lesions. *Gene Ther.* 16, 1363–1372.
- da Silva Meirelles, L., Chagastelles, P.C., and Nardi, N.B. (2006). Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J. Cell Sci.* 119, 2204–2213.
- De Bari, C., Dell'Accio, F., and Luyten, F.P. (2001a). Human periosteum-derived cells maintain phenotypic stability and chondrogenic potential throughout expansion regardless of donor age. *Arthritis Rheum.* 44, 85–95.
- De Bari, C., Dell'Accio, F., Tylzanowski, P., and Luyten, F.P. (2001b). Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum.* 44, 1928–1942.
- De Bari, C., Dell'Accio, F., Vandenabeele, F., Vermeesch, J.R., Raymackers, J.M., and Luyten, F.P. (2003). Skeletal muscle repair by adult human mesenchymal stem cells from synovial membrane. *J. Cell Biol.* 160, 909–918.
- Delorme, B., and Charbord, P. (2007). Culture and characterization of human bone marrow mesenchymal stem cells. *Methods Mol. Med.* 140, 67–81.
- Dennis, J.E., and Charbord, P. (2002). Origin and differentiation of human and murine stroma. *Stem Cells* 20, 205–214.
- Dennis, J.E., Cohen, N., Goldberg, V.M., and Caplan, A.I. (2004). Targeted delivery of progenitor cells for cartilage repair. *J. Orthop. Res.* 22, 735–741.
- Dowthwaite, G.P., Bishop, J.C., Redman, S.N., Khan, I.M., Rooney, P., Evans, D.J., Houghton, L., Bayram, Z., Boyer, S., Thomson, B., Wolfe, M.S., and Archer, C.W. (2004). The surface of articular cartilage contains a progenitor cell population. *J. Cell Sci.* 117, 889–897.
- Duffy, G.P., Ahsan, T., O'Brien, T., Barry, F., and Nerem, R.M. (2009). Bone marrow-derived mesenchymal stem cells promote angiogenic processes in a time- and dose-dependent manner *in vitro*. *Tissue Eng. Part A* 15, 2459–2470.
- Dufourcq, P., Descamps, B., Tojais, N.F., Leroux, L., Oses, P., Daret, D., Moreau, C., Lamazière, J.M., Couffinhal, T., and Dupl a, C. (2008). Secreted frizzled-related protein-1 enhances mesenchymal stem cell function in angiogenesis and contributes to neovessel maturation. *Stem Cells* 26, 2991–3001.
- Dunlop, D.D., Manheim, L.M., Yelin, E.H., Song, J., and Chang, R.W. (2003). The costs of arthritis. *Arthritis Rheum.* 49, 101–113.
- English, K., Barry, F.P., Field-Corbett, C.P., and Mahon, B.P. (2007). IFN- γ and TNF- α differentially regulate immunomodulation by murine mesenchymal stem cells. *Immunol. Lett.* 110, 91–100.
- English, K., Barry, F.P., and Mahon, B.P. (2008). Murine mesenchymal stem cells suppress dendritic cell migration, maturation and antigen presentation. *Immunol. Lett.* 115, 50–58.

- English, K., Ryan, J.M., Tobin, L., Murphy, M.J., Barry, F.P., and Mahon, B.P. (2009). Cell contact, prostaglandin E₂ and transforming growth factor β_1 play non-redundant roles in human mesenchymal stem cell induction of CD4⁺CD25^{High} forkhead box P3⁺ regulatory T cells. *Clin. Exp. Immunol.* 156, 149–160.
- Estrada, R., Li, N., Sarojini, H., An, J., Lee, M.J., and Wang, E. (2009). Secretome from mesenchymal stem cells induces angiogenesis via Cyr61. *J. Cell. Physiol.* 219, 563–571.
- Feng, Q., Lu, S.J., Klimanskaya, I., Gomes, I., Kim, D., Chung, Y., Honig, G.R., Kim, K.S., and Lanza, R. (2010). Hemangioblastic derivatives from human induced pluripotent stem cells exhibit limited expansion and early senescence. *Stem Cells* 28, 704–712.
- Fernandes, J., Tardif, G., Martel-Pelletier, J., Lascau-Coman, V., Dupuis, M., Moldovan, F., Sheppard, M., Krishnan, B.R., and Pelletier, J.P. (1999). *In vivo* transfer of interleukin-1 receptor antagonist gene in osteoarthritic rabbit knee joints: Prevention of osteoarthritis progression. *Am. J. Pathol.* 154, 1159–1169.
- Fickert, S., Fiedler, J., and Brenner, R.E. (2004). Identification of subpopulations with characteristics of mesenchymal progenitor cells from human osteoarthritic cartilage using triple staining for cell surface markers. *Arthritis Res. Ther.* 6, R422–R432.
- Findlay, D.M. (2010). If good things come from above, do bad things come from below? *Arthritis Res. Ther.* 12, 119.
- Flynn, A., Barry, F., and O'Brien, T. (2007). UC blood-derived mesenchymal stromal cells: An overview. *Cytotherapy* 9, 717–726.
- Friedenstein, A.J. (1995). Marrow stromal fibroblasts. *Calcif. Tissue Int.* 56(Suppl. 1), S17.
- Friedenstein, A.J., Piatetzky-Shapiro, I.I., and Petrakova, K.V. (1966). Osteogenesis in transplants of bone marrow cells. *J. Embryol. Exp. Morphol.* 16, 381–390.
- Friedenstein, A.J., Chailakhjan, R.K., and Lalykina, K.S. (1970). The development of fibroblast colonies in monolayer cultures of guinea-pig bone marrow and spleen cells. *Cell Tissue Kinet.* 3, 393–403.
- Friedenstein, A.J., Chailakhyan, R.K., Latsinik, N.V., Panasyuk, A.F., and Keiliss-Borok, I.V. (1974). Stromal cells responsible for transferring the microenvironment of the hemopoietic tissues: Cloning *in vitro* and retransplantation *in vivo*. *Transplantation* 17, 331–340.
- Frisbie, D.D., and McIlwraith, C.W. (2000). Evaluation of gene therapy as a treatment for equine traumatic arthritis and osteoarthritis. *Clin. Orthop. Relat. Res.* 379(Suppl.), S273–S287.
- Frisbie, D.D., Ghivizzani, S.C., Robbins, P.D., Evans, C.H., and McIlwraith, C.W. (2002). Treatment of experimental equine osteoarthritis by *in vivo* delivery of the equine interleukin-1 receptor antagonist gene. *Gene Ther.* 9, 12–20.
- Gimble, J.M., Katz, A.J., and Bunnell, B.A. (2007). Adipose-derived stem cells for regenerative medicine. *Circ. Res.* 100, 1249–1260.
- Goldring, M.B., and Goldring, S.R. (2007). Osteoarthritis. *J. Cell. Physiol.* 213, 626–634.
- Goldring, M.B., and Goldring, S.R. (2010). Articular cartilage and subchondral bone in the pathogenesis of osteoarthritis. *Ann. N. Y. Acad. Sci.* 1192, 230–237.
- Goodrich, L.R., Brower-Toland, B.D., Warnick, L., Robbins, P.D., Evans, C.H., and Nixon, A.J. (2006). Direct adenovirus-mediated IGF-I gene transduction of synovium induces persisting synovial fluid IGF-I ligand elevations. *Gene Ther.* 13, 1253–1262.
- Goodrich, L.R., Hidaka, C., Robbins, P.D., Evans, C.H., and Nixon, A.J. (2007). Genetic modification of chondrocytes with insulin-like growth factor-1 enhances cartilage healing in an equine model. *J. Bone Joint Surg. Br.* 89, 672–685.
- Gouze, J.N., Gouze, E., Palmer, G.D., Liew, V.S., Pascher, A., Betz, O.B., Thornhill, T.S., Evans, C.H., Grodzinsky, A.J., and Ghivizzani, S.C. (2003). A comparative study of the inhibitory effects of interleukin-1 receptor antagonist following administration as a recombinant protein or by gene transfer. *Arthritis Res. Ther.* 5, R301–R309.
- Grogan, S.P., Miyaki, S., Asahara, H., D'Lima, D.D., and Lotz, M.K. (2009). Mesenchymal progenitor cell markers in human articular cartilage: Normal distribution and changes in osteoarthritis. *Arthritis Res. Ther.* 11, R85.
- Guo, C.A., Liu, X.G., Huo, J.Z., Jiang, C., Wen, X.J., and Chen, Z.R. (2007). Novel gene-modified-tissue engineering of cartilage using stable transforming growth factor- β_1 -transfected mesenchymal stem cells grown on chitosan scaffolds. *J. Biosci. Bioeng.* 103, 547–556.
- Hangody, L., and Fules, P. (2003). Autologous osteochondral mosaicplasty for the treatment of full-thickness defects of weight-bearing joints: Ten years of experimental and clinical experience. *J. Bone Joint Surg. Am.* 85-A(Suppl. 2), 25–32.
- Haupt, J.L., Frisbie, D.D., McIlwraith, C.W., Robbins, P.D., Ghivizzani, S., Evans, C.H., and Nixon, A.J. (2005). Dual transduction of insulin-like growth factor-I and interleukin-1 receptor antagonist protein controls cartilage degradation in an osteoarthritic culture model. *J. Orthop. Res.* 23, 118–126.
- Hidaka, C., Quitoriano, M., Warren, R.F., and Crystal, R.G. (2001). Enhanced matrix synthesis and *in vitro* formation of cartilage-like tissue by genetically modified chondrocytes expressing BMP-7. *J. Orthop. Res.* 19, 751–758.
- Hiraoka, K., Grogan, S., Olee, T., and Lotz, M. (2006). Mesenchymal progenitor cells in adult human articular cartilage. *Biorheology* 43, 447–454.
- Horie, M., Sekiya, I., Muneta, T., Ichinose, S., Matsumoto, K., Saito, H., Murakami, T., and Kobayashi, E. (2009). Intra-articular injected synovial stem cells differentiate into meniscal cells directly and promote meniscal regeneration without mobilization to distant organs in rat massive meniscal defect. *Stem Cells* 27, 878–887.
- Hsieh, J.L., Shen, P.C., Shiau, A.L., Jou, I.M., Lee, C.H., Teo, M.L., Wang, C.R., Chao, J., Chao, L., and Wu, C.L. (2009). Adenovirus-mediated kallistatin gene transfer ameliorates disease progression in a rat model of osteoarthritis induced by anterior cruciate ligament transection. *Hum. Gene Ther.* 20, 147–158.
- Ikeda, T., Kamekura, S., Mabuchi, A., Kou, I., Seki, S., Takato, T., Nakamura, K., Kawaguchi, H., Ikegawa, S., and Chung, U.I. (2004). The combination of SOX5, SOX6, and SOX9 (the SOX trio) provides signals sufficient for induction of permanent cartilage. *Arthritis Rheum.* 50, 3561–3573.
- Im, G.I., Jung, N.H., and Tae, S.K. (2006). Chondrogenic differentiation of mesenchymal stem cells isolated from patients in late adulthood: The optimal conditions of growth factors. *Tissue Eng.* 12, 527–536.
- Ito, H., Goater, J.J., Tiyyapatanaputi, P., Rubery, P.T., O'Keefe, R.J., and Schwarz, E.M. (2004). Light-activated gene transduction of recombinant adeno-associated virus in human mesenchymal stem cells. *Gene Ther.* 11, 34–41.
- Jaenisch, R., and Young, R. (2008). Stem cells, the molecular circuitry of pluripotency and nuclear reprogramming. *Cell* 132, 567–582.
- Jones, E.A., Crawford, A., English, A., Henshaw, K., Mundy, J., Corscadden, D., Chapman, T., Emery, P., Hatton, P., and McGonagle, D. (2008). Synovial fluid mesenchymal stem cells

- in health and early osteoarthritis: Detection and functional evaluation at the single-cell level. *Arthritis Rheum.* 58, 1731–1740.
- Karystinou, A., Dell'Accio, F., Kurth, T.B., Wackerhage, H., Khan, I.M., Archer, C.W., Jones, E.A., Mitsiadis, T.A., and De Bari, C. (2009). Distinct mesenchymal progenitor cell subsets in the adult human synovium. *Rheumatology* 48, 1057–1064.
- Kaul, G., Cucchiari, M., Arntzen, D., Zurakowski, D., Menger, M.D., Kohn, D., Trippel, S.B., and Madry, H. (2006). Local stimulation of articular cartilage repair by transplantation of encapsulated chondrocytes overexpressing human fibroblast growth factor 2 (FGF-2) *in vivo*. *J. Gene Med.* 8, 100–111.
- Kay, J.D., Gouze, E., Oligino, T.J., Gouze, J.N., Watson, R.S., Levings, P.P., Bush, M.L., Dacanay, A., Nickerson, D.M., Robbins, P.D., Evans, C.H., and Ghivizzani, S.C. (2009). Intra-articular gene delivery and expression of interleukin-1Ra mediated by self-complementary adeno-associated virus. *J. Gene Med.* 11, 605–614.
- Khan, I.M., Bishop, J.C., Gilbert, S., and Archer, C.W. (2009). Clonal chondroprogenitors maintain telomerase activity and Sox9 expression during extended monolayer culture and retain chondrogenic potential. *Osteoarthritis Cartilage* 17, 518–528.
- Kinnaird, T., Stabile, E., Burnett, M.S., Lee, C.W., Barr, S., Fuchs, S., and Epstein, S.E. (2004). Marrow-derived stromal cells express genes encoding a broad spectrum of arteriogenic cytokines and promote *in vitro* and *in vivo* arteriogenesis through paracrine mechanisms. *Circ. Res.* 94, 678–685.
- Knutsen, G., Engebretsen, L., Ludvigsen, T.C., Drogset, J.O., Grøntvedt, T., Solheim, E., Strand, T., Roberts, S., Isaksen, V., and Johansen, O. (2004). Autologous chondrocyte implantation compared with microfracture in the knee: A randomized trial. *J. Bone Joint Surg. Am.* 86-A, 455–464.
- Koelling, S., Kruegel, J., Irmer, M., Path, J.R., Sadowski, B., Miro, X., and Miosge, N. (2009). Migratory chondrogenic progenitor cells from repair tissue during the later stages of human osteoarthritis. *Cell Stem Cell* 4, 324–335.
- Koyama, E., Shibukawa, Y., Nagayama, M., Sugito, H., Young, B., Yuasa, T., Okabe, T., Ochiai, T., Kamiya, N., Rountree, R.B., Kingsley, D.M., Iwamoto, M., Enomoto-Iwamoto, M., and Pacifici, M. (2008). A distinct cohort of progenitor cells participates in synovial joint and articular cartilage formation during mouse limb skeletogenesis. *Dev. Biol.* 316, 62–73.
- Ladage, D., Brixius, K., Steingen, C., Mehlhorn, U., Schwinger, R.H., Bloch, W., and Schmidt, A. (2007). Mesenchymal stem cells induce endothelial activation via paracrine mechanisms. *Endothelium* 14, 53–63.
- Lee, D.K., Choi, K.B., Oh, I.S., Song, S.U., Hwang, S., Lim, C.L., Hyun, J.P., Lee, H.Y., Chi, G.F., Yi, Y., Yip, V., Kim, J., Lee, E.B., Noh, M.J., and Lee, K.H. (2005). Continuous transforming growth factor β_1 secretion by cell-mediated gene therapy maintains chondrocyte redifferentiation. *Tissue Eng.* 11, 310–318.
- Lee, K.B., Hui, J.H., Song, I.C., Ardany, L., and Lee, E.H. (2007). Injectable mesenchymal stem cell therapy for large cartilage defects: A porcine model. *Stem Cells* 25, 2964–2971.
- Lee, S.Y., Nakagawa, T., and Reddi, A.H. (2010). Mesenchymal progenitor cells derived from synovium and infrapatellar fat pad as a source for superficial zone cartilage tissue engineering: Analysis of superficial zone protein/lubricin expression. *Tissue Eng. Part A* 16, 317–325.
- Li, X., and Makarov, S.S. (2006). An essential role of NF- κ B in the tumor-like phenotype of arthritic synoviocytes. *Proc. Natl. Acad. Sci. U.S.A.* 103, 17432–17437.
- Lianxu, C., Hongti, J., and Changlong, Y. (2006). NF- κ Bp65-specific siRNA inhibits expression of genes of COX-2, NOS-2 and MMP-9 in rat IL-1 β -induced and TNF- α -induced chondrocytes. *Osteoarthritis Cartilage* 14, 367–376.
- Liu, N., Chen, R., Du, H., Wang, J., Zhang, Y., and Wen, J. (2009). Expression of IL-10 and TNF- α in rats with cerebral infarction after transplantation with mesenchymal stem cells. *Cell Mol. Immunol.* 6, 207–213.
- Madry, H., Kaul, G., Cucchiari, M., Stein, U., Zurakowski, D., Remberger, K., Menger, M.D., Kohn, D., and Trippel, S.B. (2005). Enhanced repair of articular cartilage defects *in vivo* by transplanted chondrocytes overexpressing insulin-like growth factor I (IGF-I). *Gene Ther.* 12, 1171–1179.
- Marcacci, M., Kon, E., Moukhachev, V., Lavroukov, A., Kute-pov, S., Quarto, R., Mastrogiacomo, M., and Cancedda, R. (2007). Stem cells associated with macroporous bioceramics for long bone repair: 6- to 7-year outcome of a pilot clinical study. *Tissue Eng.* 13, 947–955.
- Mason, J.M., Grande, D.A., Barcia, M., Grant, R., Pergolizzi, R.G., and Breitbart, A.S. (1998). Expression of human bone morphogenic protein 7 in primary rabbit periosteal cells: Potential utility in gene therapy for osteochondral repair. *Gene Ther.* 5, 1098–1104.
- Meliga, E., Strem, B.M., Duckers, H.J., and Serruys, P.W. (2007). Adipose-derived cells. *Cell Transplant.* 16, 963–970.
- Meng, J., Adkin, C.F., Arechavala-Gomez, V., Boldrin, L., Muntoni, F., and Morgan, J.E. (2010). The contribution of human synovial stem cells to skeletal muscle regeneration. *Neuromuscul. Disord.* 20, 6–15.
- Michaud, C.M., McKenna, M.T., Begg, S., Tomijima, N., Majmudar, M., Bulzacchelli, M.T., Ebrahim, S., Ezzati, M., Salomon, J.A., Kreiser, J.G., Hogan, M., and Murray, C.J. (2006). The burden of disease and injury in the United States 1996. *Popul. Health Metr.* 4, 11.
- Mizuno, K., Muneta, T., Morito, T., Ichinose, S., Koga, H., Nimura, A., Mochizuki, T., and Sekiya, I. (2008). Exogenous synovial stem cells adhere to defect of meniscus and differentiate into cartilage cells. *J. Med. Dent. Sci.* 55, 101–111.
- Mody, N., Tintut, Y., Radcliff, K., and Demer, L.L. (2003). Vascular calcification and its relation to bone calcification: Possible underlying mechanisms. *J. Nucl. Cardiol.* 10, 177–183.
- Morisset, S., Frisbie, D.D., Robbins, P.D., Nixon, A.J., and McIlwraith, C.W. (2007). IL-1ra/IGF-1 gene therapy modulates repair of microfractured chondral defects. *Clin. Orthop. Relat. Res.* 462, 221–228.
- Murakami, S., Lefebvre, V., and de Crombrughe, B. (2000). Potent inhibition of the master chondrogenic factor Sox9 gene by interleukin-1 and tumor necrosis factor- α . *J. Biol. Chem.* 275, 3687–3692.
- Murphy, J.M., Dixon, K., Beck, S., Fabian, D., Feldman, A., and Barry, F. (2002). Reduced chondrogenic and adipogenic activity of mesenchymal stem cells from patients with advanced osteoarthritis. *Arthritis Rheum.* 46, 704–713.
- Murphy, J.M., Fink, D.J., Hunziker, E.B., and Barry, F.P. (2003). Stem cell therapy in a caprine model of osteoarthritis. *Arthritis Rheum.* 48, 3464–3474.
- Németh, K., Leelahavanichkul, A., Yuen, P.S., Mayer, B., Parmelee, A., Doi, K., Robey, P.G., Leelahavanichkul, K., Koller, B.H., Brown, J.M., Hu, X., Jelinek, I., Star, R.A., and Mezey, E. (2009). Bone marrow stromal cells attenuate sepsis via prostaglandin E₂-dependent reprogramming of host macrophages to increase their interleukin-10 production. *Nat. Med.* 15, 42–49.

- Nishikawa, S., Goldstein, R.A., and Nierras, C.R. (2008). The promise of human induced pluripotent stem cells for research and therapy. *Nat. Rev. Mol. Cell Biol.* 9, 725–729.
- Nixon, A.J., Brower-Toland, B.D., Bent, S.J., Saxer, R.A., Wilke, M.J., Robbins, P.D., and Evans, C.H. (2000). Insulinlike growth factor-I gene therapy applications for cartilage repair. *Clin. Orthop. Relat. Res.* 379(Suppl.), S201–S213.
- Oh, J.S., Ha, Y., An, S.S., Khan, M., Pennant, W.A., Kim, H.J., Yoon do, H., Lee, M., and Kim, K.N. (2010). Hypoxia-preconditioned adipose tissue-derived mesenchymal stem cell increase the survival and gene expression of engineered neural stem cells in a spinal cord injury model. *Neurosci. Lett.* 472, 215–219.
- Ortiz, L.A., Gambelli, F., McBride, C., Gaupp, D., Baddoo, M., Kaminski, N., and Phinney, D.G. (2003). Mesenchymal stem cell engraftment in lung is enhanced in response to bleomycin exposure and ameliorates its fibrotic effects. *Proc. Natl. Acad. Sci. U.S.A.* 100, 8407–8411.
- Ortiz, L.A., Dutreil, M., Fattman, C., Pandey, A.C., Torres, G., Go, K., and Phinney, D.G. (2007). Interleukin 1 receptor antagonist mediates the antiinflammatory and antifibrotic effect of mesenchymal stem cells during lung injury. *Proc. Natl. Acad. Sci. U.S.A.* 104, 11002–11007.
- Owen, M., and Friedenstein, A.J. (1988). Stromal stem cells: Marrow-derived osteogenic precursors. *Ciba Found. Symp.* 136, 42–60.
- Pagnotto, M.R., Wang, Z., Karpie, J.C., Ferretti, M., Xiao, X., and Chu, C.R. (2007). Adeno-associated viral gene transfer of transforming growth factor- β_1 to human mesenchymal stem cells improves cartilage repair. *Gene Ther.* 14, 804–813.
- Pelletier, J.P., Caron, J.P., Evans, C., Robbins, P.D., Georgescu, H.I., Jovanovic, D., Fernandes, J.C., and Martel-Pelletier, J. (1997). *In vivo* suppression of early experimental osteoarthritis by interleukin-1 receptor antagonist using gene therapy. *Arthritis Rheum.* 40, 1012–1019.
- Pittenger, M. (2009). Sleuthing the source of regeneration by MSCs. *Cell Stem Cell* 5, 8–10.
- Pittenger, M.F., Mackay, A.M., Beck, S.C., Jaiswal, R.K., Douglas, R., Mosca, J.D., Moorman, M.A., Simonetti, D.W., Craig, S., and Marshak, D.R. (1999). Multilineage potential of adult human mesenchymal stem cells. *Science* 284, 143–147.
- Polchert, D., Sobinsky, J., Douglas, G., Kidd, M., Moadsiri, A., Reina, E., Genrich, K., Mehrotra, S., Setty, S., Smith, B., and Bartholomew, A. (2008). IFN- γ activation of mesenchymal stem cells for treatment and prevention of graft versus host disease. *Eur. J. Immunol.* 38, 1745–1755.
- Pulavendran, S., Vignesh, J., and Rose, C. (2010). Differential anti-inflammatory and anti-fibrotic activity of transplanted mesenchymal vs. hematopoietic stem cells in carbon tetrachloride-induced liver injury in mice. *Int. Immunopharmacol.* 10, 513–519.
- Quarto, R., Mastrogiacomo, M., Cancedda, R., Kutepov, S.M., Mukhachev, V., Lavroukov, A., Kon, E., and Marcacci, M. (2001). Repair of large bone defects with the use of autologous bone marrow stromal cells. *N. Engl. J. Med.* 344, 385–386.
- Quevedo, H.C., Hatzistergos, K.E., Oskouei, B.N., Feigenbaum, G.S., Rodriguez, J.E., Valdes, D., Pattany, P.M., Zambrano, J.P., Hu, Q., McNiece, I., Heldman, A.W., and Hare, J.M. (2009). Allogeneic mesenchymal stem cells restore cardiac function in chronic ischemic cardiomyopathy via trilineage differentiating capacity. *Proc. Natl. Acad. Sci. U.S.A.* 106, 14022–14027.
- Rachakonda, P.S., Rai, M.F., and Schmidt, M.F. (2008). Application of inflammation-responsive promoter for an *in vitro* arthritis model. *Arthritis Rheum.* 58, 2088–2097.
- Redman, S.N., Oldfield, S.F., and Archer, C.W. (2005). Current strategies for articular cartilage repair. *Eur. Cell Mater.* 9, 23–32; discussion 23–32.
- Sacchetti, B., Funari, A., Michienzi, S., Di Cesare, S., Piersanti, S., Saggio, I., Tagliafico, E., Ferrari, S., Robey, P.G., Riminucci, M., and Bianco, P. (2007). Self-renewing osteoprogenitors in bone marrow sinusoids can organize a hematopoietic micro-environment. *Cell* 131, 324–336.
- Sackstein, R., Merzaban, J.S., Cain, D.W., Dagia, N.M., Spencer, J.A., Lin, C.P., and Wohlgemuth, R. (2008). *Ex vivo* glycan engineering of CD44 programs human multipotent mesenchymal stromal cell trafficking to bone. *Nat. Med.* 14, 181–187.
- Sakaguchi, Y., Sekiya, I., Yagishita, K., and Muneta, T. (2005). Comparison of human stem cells derived from various mesenchymal tissues: Superiority of synovium as a cell source. *Arthritis Rheum.* 52, 2521–2529.
- Scanzello, C.R., Plaas, A., and Crow, M.K. (2008). Innate immune system activation in osteoarthritis: Is osteoarthritis a chronic wound? *Curr. Opin. Rheumatol.* 20, 565–572.
- Scharstuhl, A., Schewe, B., Benz, K., Gaissmaier, C., Bühring, H.J., and Stoop, R. (2007). Chondrogenic potential of human adult mesenchymal stem cells is independent of age or osteoarthritis etiology. *Stem Cells* 25, 3244–3251.
- Shi, S., and Gronthos, S. (2003). Perivascular niche of postnatal mesenchymal stem cells in human bone marrow and dental pulp. *J. Bone Miner. Res.* 18, 696–704.
- Smith, P., Shuler, F.D., Georgescu, H.I., Ghivizzani, S.C., Johnstone, B., Niyibizi, C., Robbins, P.D., and Evans, C.H. (2000). Genetic enhancement of matrix synthesis by articular chondrocytes: Comparison of different growth factor genes in the presence and absence of interleukin-1. *Arthritis Rheum.* 43, 1156–1164.
- Steinert, A.F., Ghivizzani, S.C., Rethwilm, A., Tuan, R.S., Evans, C.H., and Nöth, U. (2007). Major biological obstacles for persistent cell-based regeneration of articular cartilage. *Arthritis Res. Ther.* 9, 213.
- Surendran, S., Kim, S.H., Jee, B.K., Ahn, S.H., Gopinathan, P., and Han, C.W. (2006). Anti-apoptotic Bcl-2 gene transfection of human articular chondrocytes protects against nitric oxide-induced apoptosis. *J. Bone Joint Surg. Br.* 88, 1660–1665.
- Tasso, R., Augello, A., Boccardo, S., Salvi, S., Caridà, M., Postiglione, F., Fais, F., Truini, M., Cancedda, R., and Pennesi, G. (2009). Recruitment of a host's osteoprogenitor cells using exogenous mesenchymal stem cells seeded on porous ceramic. *Tissue Eng. Part A* 15, 2203–2212.
- Tavian, M., and Peault, B. (2005). The changing cellular environments of hematopoiesis in human development *in utero*. *Exp. Hematol.* 33, 1062–1069.
- Tavian, M., Zheng, B., Oberlin, E., Crisan, M., Sun, B., Huard, J., and Peault, B. (2005). The vascular wall as a source of stem cells. *Ann. N. Y. Acad. Sci.* 1044, 41–50.
- Tew, S.R., Li, Y., Pothacharoen, P., Tweats, L.M., Hawkins, R.E., and Hardingham, T.E. (2005). Retroviral transduction with SOX9 enhances re-expression of the chondrocyte phenotype in passaged osteoarthritic human articular chondrocytes. *Osteoarthritis Cartilage* 13, 80–89.
- Tintut, Y., Alfonso, Z., Saini, T., Radcliff, K., Watson, K., Bostrom, K., and Demer, L.L. (2003). Multilineage potential of cells from the artery wall. *Circulation* 108, 2505–2510.
- Troyer, D.L., and Weiss, M.L. (2008). Wharton's jelly-derived cells are a primitive stromal cell population. *Stem Cells* 26, 591–599.
- Tweedell, K.S. (2008). New paths to pluripotent stem cells. *Curr. Stem Cell Res. Ther.* 3, 151–162.

- Ulrich-Vinther, M., Stengaard, C., Schwarz, E.M., Goldring, M.B., and Soballe, K. (2005). Adeno-associated vector mediated gene transfer of transforming growth factor- β_1 to normal and osteoarthritic human chondrocytes stimulates cartilage anabolism. *Eur. Cell Mater.* 10, 40–50.
- Urbach, A., Bar-Nur, O., Daley, G.Q., and Benvenisty, N. (2010). Differential modeling of fragile X syndrome by human embryonic stem cells and induced pluripotent stem cells. *Cell Stem Cell* 6, 407–411.
- van der Kraan, P.M., and van den Berg, W.B. (2007). Osteophytes: Relevance and biology. *Osteoarthritis Cartilage* 15, 237–244.
- Venkatesan, N., Barré, L., Benani, A., Netter, P., Magdalou, J., Fournel-Gigleux, S., and Ouzzine, M. (2004). Stimulation of proteoglycan synthesis by glucuronosyltransferase-I gene delivery: A strategy to promote cartilage repair. *Proc. Natl. Acad. Sci. U.S.A.* 101, 18087–18092.
- Wakitani, S., Goto, T., Pineda, S.J., Young, R.G., Mansour, J.M., Caplan, A.I., and Goldberg, V.M. (1994). Mesenchymal cell-based repair of large, full-thickness defects of articular cartilage. *J. Bone Joint Surg. Am.* 76, 579–592.
- Wakitani, S., Imoto, K., Yamamoto, T., Saito, M., Murata, N., and Yoneda, M. (2002). Human autologous culture expanded bone marrow mesenchymal cell transplantation for repair of cartilage defects in osteoarthritic knees. *Osteoarthritis Cartilage* 10, 199–206.
- Wang, C.R., Chen, S.Y., Wu, C.L., Liu, M.F., Jin, Y.T., Chao, L., and Chao, J. (2005). Prophylactic adenovirus-mediated human kallistatin gene therapy suppresses rat arthritis by inhibiting angiogenesis and inflammation. *Arthritis Rheum.* 52, 1319–1324.
- Wehling, N., Palmer, G.D., Pilapil, C., Liu, F., Wells, J.W., Müller, P.E., Evans, C.H., and Porter, R.M. (2009). Interleukin-1 β and tumor necrosis factor α inhibit chondrogenesis by human mesenchymal stem cells through NF- κ B-dependent pathways. *Arthritis Rheum.* 60, 801–812.
- Wei, X., and Messner, K. (1999). Maturation-dependent durability of spontaneous cartilage repair in rabbit knee joint. *J. Biomed. Mater. Res.* 46, 539–548.
- Weiss, M.L., and Troyer, D.L. (2006). Stem cells in the umbilical cord. *Stem Cell Rev.* 2, 155–162.
- Weiss, M.L., Medicetty, S., Bledsoe, A.R., Rachakatla, R.S., Choi, M., Merchav, S., Luo, Y., Rao, M.S., Velagaleti, G., and Troyer, D. (2006). Human umbilical cord matrix stem cells: Preliminary characterization and effect of transplantation in a rodent model of Parkinson's disease. *Stem Cells* 24, 781–792.
- Yokoo, N., Saito, T., Uesugi, M., Kobayashi, N., Xin, K.Q., Okuda, K., Mizukami, H., Ozawa, K., and Koshino, T. (2005). Repair of articular cartilage defect by autologous transplantation of basic fibroblast growth factor gene-transduced chondrocytes with adeno-associated virus vector. *Arthritis Rheum.* 52, 164–170.
- Zannettino, A.C., Paton, S., Arthur, A., Khor, F., Itescu, S., Gimble, J.M., and Gronthos, S. (2008). Multipotential human adipose-derived stromal stem cells exhibit a perivascular phenotype *in vitro* and *in vivo*. *J. Cell. Physiol.* 214, 413–421.
- Zhang, X., Mao, Z., and Yu, C. (2004). Suppression of early experimental osteoarthritis by gene transfer of interleukin-1 receptor antagonist and interleukin-10. *J. Orthop. Res.* 22, 742–750.
- Zhang, X., Yu, C., Xushi, Zhang, C., Tang, T., and Dai, K. (2006). Direct chitosan-mediated gene delivery to the rabbit knee joints *in vitro* and *in vivo*. *Biochem. Biophys. Res. Commun.* 341, 202–208.
- Zhang, Y.W., Su, Y., Lanning, N., Swiatek, P.J., Bronson, R.T., Sigler, R., Martin, R.W., and Vande Woude, G.F. (2005). Targeted disruption of Mig-6 in the mouse genome leads to early onset degenerative joint disease. *Proc. Natl. Acad. Sci. U.S.A.* 102, 11740–11745.

Address correspondence to:

Dr. J. Mary Murphy

Regenerative Medicine Institute

National Centre for Biomedical Engineering Science

National University of Ireland

Galway, Ireland

E-mail: mary.murphy@nuigalway.ie

Received for publication July 7, 2010;
accepted July 12, 2010.

Published online: August 26, 2010.