

University of Groningen

## Meta-Analysis of Genome-Wide Association Studies in > 80 000 Subjects Identifies Multiple Loci for C-Reactive Protein Levels

Dehghan, Abbas; Dupuis, Josee; Barbalic, Maja; Bis, Joshua C.; Eiriksdottir, Gudny; Lu, Chen; Pellikka, Niina; Wallaschofski, Henri; Kettunen, Johannes; Henneman, Peter

*Published in:*  
Circulation

*DOI:*  
[10.1161/CIRCULATIONAHA.110.948570](https://doi.org/10.1161/CIRCULATIONAHA.110.948570)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2011

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Dehghan, A., Dupuis, J., Barbalic, M., Bis, J. C., Eiriksdottir, G., Lu, C., Pellikka, N., Wallaschofski, H., Kettunen, J., Henneman, P., Baumert, J., Strachan, D. P., Fuchsberger, C., Vitart, V., Wilson, J. F., Pare, G., Naitza, S., Rudock, M. E., Surakka, I., ... Chasman, D. I. (2011). Meta-Analysis of Genome-Wide Association Studies in > 80 000 Subjects Identifies Multiple Loci for C-Reactive Protein Levels. *Circulation*, 123(7), 731-738. <https://doi.org/10.1161/CIRCULATIONAHA.110.948570>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Meta-Analysis of Genome-Wide Association Studies in >80 000 Subjects Identifies Multiple Loci for C-Reactive Protein Levels

Abbas Dehghan, MD, PhD\*; Josée Dupuis, PhD\*; Maja Barbalic, PhD\*; Joshua C. Bis, PhD\*; Gudny Eiriksdottir, MSc;\* Chen Lu, MA; Niina Pellikka, BEng; Henri Wallaschofski, MD; Johannes Kettunen, MSc; Peter Henneman, MSc; Jens Baumert, PhD; David P. Strachan, MD; Christian Fuchsberger, PhD; Veronique Vitart, PhD; James F. Wilson, BSc, DPhil; Guillaume Paré, MD, MSc; Silvia Naitza, PhD; Megan E. Rudock, PhD; Ida Surakka, BSc; Eco J.C. de Geus, PhD; Behrooz Z. Alizadeh, PhD; Jack Guralnik, MD, PhD; Alan Shuldiner, MD; Toshiko Tanaka, PhD; Robert Y.L. Zee, PhD; Renate B. Schnabel, MD, MSc; Vijay Nambi, MD; Maryam Kavousi, MD, MSc; Samuli Ripatti, PhD; Matthias Nauck, MD; Nicholas L. Smith, PhD; Albert V. Smith, PhD; Jouko Sundvall, PhD; Paul Scheet, PhD; Yongmei Liu, MD, PhD; Aimo Ruokonen, MD, PhD; Lynda M. Rose, MSc; Martin G. Larson, ScD; Ron C. Hoogeveen, PhD; Nelson B. Freimer, MD; Alexander Teumer, Dipl-Math; Russell P. Tracy, PhD; Lenore J. Launer, PhD; Julie E. Buring, DSc; Jennifer F. Yamamoto, MA; Aaron R. Folsom, MD, MPH; Eric J.G. Sijbrands, MD, PhD; James Pankow, PhD; Paul Elliott, MBBS, PhD, FMedSci; John F. Keaney, MD; Wei Sun, MD, PhD; Antti-Pekka Sarin, BSc; João D. Fontes, MD; Sunita Badola, MSc; Brad C. Astor, PhD, MPH; Albert Hofman, MD, PhD; Anneli Pouta, MD, PhD; Karl Werdan, MD; Karin H. Greiser, MD; Oliver Kuss, PhD; Henriette E. Meyer zu Schwabedissen, MD; Joachim Thiery, MD; Yalda Jamshidi, PhD; Ilja M. Nolte, PhD; Nicole Soranzo, PhD; Timothy D. Spector, MD, MSc, FRCP; Henry Völzke, MD; Alexander N. Parker, PhD; Thor Aspelund, PhD; David Bates, MD, MSc; Lauren Young; Kim Tsui; David S. Siscovick, MD, MPH; Xiuqing Guo, PhD; Jerome I. Rotter, MD; Manuela Uda, PhD; David Schlessinger, PhD; Igor Rudan, MD; Andrew A. Hicks, PhD; Brenda W. Penninx, PhD; Barbara Thorand, PhD, MPH; Christian Gieger, PhD, MS; Joe Coresh, MD, PhD; Gonke Willemsen, PhD; Tamara B. Harris, MD, MSc; Andre G. Uitterlinden, PhD; Marjo-Riitta Järvelin, MD, MSc, PhD; Kenneth Rice, PhD; Dörte Radke; Veikko Salomaa, MD, PhD; Ko Willems van Dijk, PhD; Eric Boerwinkle, PhD; Ramachandran S. Vasan, MD; Luigi Ferrucci, MD, PhD; Quince D. Gibson, MBA; Stefania Bandinelli, MD; Harold Snieder, PhD; Dorret I. Boomsma, PhD; Xiangjun Xiao; Harry Campbell, MBChB, MD; Caroline Hayward, PhD; Peter P. Pramstaller, MD; Cornelia M. van Duijn, PhD; Leena Peltonen, MD, PhD†; Bruce M. Psaty, MD, PhD; Vilmundur Gudnason, MD, PhD; Paul M. Ridker, MD, MPH; Georg Homuth, PhD\*; Wolfgang Koenig, MD, PhD\*; Christie M. Ballantyne, MD\*; Jacqueline C.M. Witteman, PhD\*; Emelia J. Benjamin, MD, ScM\*; Markus Perola, MD, PhD\*; Daniel I. Chasman, PhD\*

**Background**—C-reactive protein (CRP) is a heritable marker of chronic inflammation that is strongly associated with cardiovascular disease. We sought to identify genetic variants that are associated with CRP levels.

**Methods and Results**—We performed a genome-wide association analysis of CRP in 66 185 participants from 15 population-based studies. We sought replication for the genome-wide significant and suggestive loci in a replication

Received February 24, 2010; accepted November 23, 2010.

A list of the institutions and affiliations for the authors of this report may be found in the Acknowledgments at the end of this article.

†Dr Peltonen passed away in March 2010.

\*These authors contributed equally to this work.

Guest Editor for this article was Kari E. North, PhD.

The online-only Data Supplement is available with this article at <http://circ.ahajournals.org/cgi/content/full/CIRCULATIONAHA.110.948570/DC1>.

Correspondence to Jacqueline C.M. Witteman, PhD, Erasmus Medical Center, Dr Molewaterplein 50, 3015 GE Rotterdam, Netherlands (E-mail [j.witteman@erasmusmc.nl](mailto:j.witteman@erasmusmc.nl)); or Daniel I. Chasman, PhD, Division of Preventive Medicine, Brigham and Women's Hospital, 900 Commonwealth Ave E, Boston, MA 02215 (E-mail [dchasman@rics.bwh.harvard.edu](mailto:dchasman@rics.bwh.harvard.edu)).

© 2011 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.110.948570

panel comprising 16 540 individuals from 10 independent studies. We found 18 genome-wide significant loci, and we provided evidence of replication for 8 of them. Our results confirm 7 previously known loci and introduce 11 novel loci that are implicated in pathways related to the metabolic syndrome (*APOC1*, *HNF1A*, *LEPR*, *GCKR*, *HNF4A*, and *PTPN2*) or the immune system (*CRP*, *IL6R*, *NLRP3*, *IL1F10*, and *IRF1*) or that reside in regions previously not known to play a role in chronic inflammation (*PPP1R3B*, *SALL1*, *PABPC4*, *ASCL1*, *RORA*, and *BCL7B*). We found a significant interaction of body mass index with *LEPR* ( $P < 2.9 \times 10^{-6}$ ). A weighted genetic risk score that was developed to summarize the effect of risk alleles was strongly associated with CRP levels and explained  $\approx 5\%$  of the trait variance; however, there was no evidence for these genetic variants explaining the association of CRP with coronary heart disease.

**Conclusions**—We identified 18 loci that were associated with CRP levels. Our study highlights immune response and metabolic regulatory pathways involved in the regulation of chronic inflammation. (*Circulation*. 2011;123:731-738.)

**Key Words:** genetics ■ inflammation ■ meta-analysis ■ myocardial infarction ■ genome-wide association study

C-reactive protein (CRP) is a general marker of systemic inflammation. High CRP levels are associated with increased risks of mortality<sup>1</sup> and major diseases including diabetes mellitus,<sup>2</sup> hypertension,<sup>3</sup> coronary heart disease (CHD),<sup>4</sup> and stroke.<sup>5</sup> The heritability of CRP levels is estimated to be 25% to 40%,<sup>6–8</sup> suggesting that genetic variation is a major determinant of CRP levels. A genome-wide association (GWA) study in 6345 women found 7 loci associated with CRP levels.<sup>9</sup> These loci were in or close to genes encoding CRP (*CRP*), leptin receptor (*LEPR*), interleukin-6 receptor (*IL6R*), glucokinase regulator (*GCKR*), hepatic nuclear factor 1- $\alpha$  (*HNF1A*), apolipoprotein E (*APOE*), and achaete-scute complex homolog 1 (*ASCL1*). Findings from other GWA studies did not extend the number of loci related to CRP.<sup>10,11</sup>

### Clinical Perspective on p 738

In this study, we sought to discover additional genes related to CRP levels using GWA scans in 66 185 participants from 15 population-based cohort studies and replicate our findings in 16 540 participants from 10 independent studies. To investigate whether the genetic variants identified interact with nongenetic determinants of CRP such as age, sex, smoking, and body mass index (BMI), we examined gene-environment interactions. Finally, the extent to which the genes associated with circulating CRP levels, individually or jointly, affect the risk of cardiovascular diseases is still unknown. To address this question, we examined the association of genetic variants with myocardial infarction (MI) and CHD.

## Methods

### Subjects and Measurements

Participants were of European ancestry. All studies had protocols approved by local institutional review boards. Participants provided written informed consent and gave permission to use their DNA for research purposes. Baseline characteristics for all participating studies are presented in Table I in the online-only Data Supplement. Baseline measures of clinical and demographic characteristics were obtained at the time of cohort entry except for British 1958 Birth Cohort (B58C), the Framingham Heart Study (FHS), Northern Finland Birth Cohort 66 (NFBC66), and the Atherosclerosis Risk in Communities (ARIC) study, in which measures were obtained at the time of phenotype measurement.

### GWA Analysis

Genome-wide scans were performed independently in each cohort with the use of various genotyping technologies (Table VII in the online-only

Data Supplement). Investigators in each study performed association analysis using the genotype-phenotype data within their cohort. Each study imputed single-nucleotide polymorphisms (SNPs) with reference to HapMap release 22 CEU and provided results for a common set of SNPs for meta-analysis. Except for the FHS, all studies conducted a linear regression analysis adjusted for age (except for NFBC66 and B58C), sex (except for the Women's Genome Health Study [WGHS]), and site of recruitment (if necessary) for all SNPs based on an additive genetic model. In the Erasmus Rucphen Family (ERF) study, adjustments for the family structure in the GWA analysis were based on the model residuals in the score test, which accounted for pedigree structure as implemented in GenABEL software<sup>12</sup> function "mmscore."<sup>13</sup> In the FHS, a linear mixed effects model was employed with the use of the lmeKin function of the kinship package in R with a fixed additive effect for the SNP genotype, fixed covariate effects, and random family-specific additive residual polygenic effects.<sup>14</sup> In each study, we estimated the genomic inflation rate, stated as lambda ( $\lambda_{gc}$ ), by comparing each study's median  $\chi^2$  value to 0.4549, the median  $\chi^2$  for the null distribution<sup>15</sup> (Table I in the online-only Data Supplement). *P* values for each cohort were adjusted for underlying population structure with the genomic inflation coefficient.

### Discovery Panel and Replication Panel

The 15-study discovery panel included 5 studies from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium,<sup>16</sup> 4 studies from the European Special Population Network (EUROSPAN), and 6 additional independent studies comprising 66 185 participants. The replication studies included 10 independent studies and 16 540 participants.

### Meta-Analysis

To calculate the combined *P* values and  $\beta$  coefficients, we used an inverse-variance weighted fixed-effects meta-analysis. We used METAL, a software package designed to perform meta-analysis on GWA data sets.<sup>17</sup> We applied an a priori threshold  $5.0 \times 10^{-8}$  for genome-wide significance.<sup>18</sup> When  $>1$  genome-wide significant SNP clustered at a locus, we took the SNP with the smallest *P* value as the lead SNP. To investigate the validity of our findings, we sought replication of the lead SNP in genome-wide significant ( $P < 5 \times 10^{-8}$ ) loci and sought additional evidence for suggestive loci ( $5 \times 10^{-8} < P < 10^{-5}$ ) in our replication panel. We ran a fixed-effect meta-analysis to combine the results of the discovery and replication panels. The first GWA study on serum CRP published by Ridker et al<sup>9</sup> was based on part of the WGHS population. To confirm that our findings were not entirely influenced by these previously published results, we performed a meta-analysis excluding the WGHS population.

### Examination of Heterogeneity

We examined between-study heterogeneity with Cochran's *Q* test. On the basis of Bonferroni adjustment for 18 tests, heterogeneity was

**Table 1. Association of 17 Genome-Wide Significant Loci With CRP Levels in the Discovery Panel**

SNP	Band	Significant SNPs	Coded Allele	Allele Frequency	$\beta^*$ (SE)	<i>P</i>	Gene
rs2794520	1q23.2	121	C	0.66	0.193 (0.007)	$9.5 \times 10^{-189}$	<i>CRP</i>
rs4420638	19q13.32	16	A	0.80	0.240 (0.010)	$2.1 \times 10^{-129}$	<i>APOC1</i>
rs1183910	12q24.31	186	G	0.67	0.152 (0.007)	$3.3 \times 10^{-113}$	<i>HNF1A</i>
rs4420065	1p31.3	291	C	0.61	0.111 (0.007)	$3.2 \times 10^{-64}$	<i>LEPR</i>
rs4129267	1q21.3	90	C	0.60	0.094 (0.007)	$1.1 \times 10^{-47}$	<i>IL6R</i>
rs1260326	2q13	54	T	0.41	0.089 (0.007)	$5.4 \times 10^{-43}$	<i>GCKR</i>
rs12239046	1q44	13	C	0.61	0.048 (0.007)	$1.6 \times 10^{-13}$	<i>NLRP3</i>
rs6734238	2p23.3	92	G	0.42	0.047 (0.007)	$3.4 \times 10^{-13}$	<i>IL1F10</i>
rs9987289	8p23.1	15	G	0.90	0.079 (0.011)	$2.3 \times 10^{-12}$	<i>PPP1R3B</i>
rs10745954	12q23.2	22	A	0.50	0.043 (0.006)	$1.6 \times 10^{-11}$	<i>ASCL1</i>
rs1800961	20q13.12	1	C	0.95	0.120 (0.018)	$2.3 \times 10^{-11}$	<i>HNF4A</i>
rs340029	15q22.2	25	T	0.62	0.044 (0.007)	$2.6 \times 10^{-11}$	<i>RORA</i>
rs10521222	16q12.1	6	C	0.94	0.110 (0.017)	$1.3 \times 10^{-10}$	<i>SALL1</i>
rs12037222	1p32.4	11	A	0.24	0.047 (0.008)	$4.5 \times 10^{-10}$	<i>PABPC4</i>
rs13233571	7q11.23	7	C	0.86	0.054 (0.010)	$2.8 \times 10^{-8}$	<i>BCL7B</i>
rs2836878	21q22.2	2	G	0.72	0.040 (0.007)	$4.0 \times 10^{-8}$	<i>PSMG1</i>
rs4903031	14q24.2	1	G	0.21	0.046 (0.008)	$4.6 \times 10^{-8}$	<i>RGS6</i>

\* $\beta$  coefficient represents 1-unit change in the natural log-transformed CRP (mg/L) per copy increment in the coded allele.

considered significant at a *P* value  $< 2.8 \times 10^{-3}$ . We explored the source of heterogeneity for significant SNPs by fitting a covariate (age, gender, BMI, or smoking) in a meta-regression model.

**Gene-Environment Interaction**

For all genome-wide significant SNPs, we examined gene-by-age, gene-by-sex, gene-by-BMI, and gene-by-smoking interactions in

each study by introducing an interaction term into a linear model with age, sex, and the covariate of interest as the independent variables and natural log-transformed CRP as the outcome. A meta-analysis was performed to combine the reported interaction  $\beta$  and *P* values across studies for each of the top SNPs. On the basis of Bonferroni adjustment for 72 tests (18 SNPs for 4 environmental factors), we used a significance threshold at  $6.9 \times 10^{-4}$ .

**Table 2. Association of 17 Genome-Wide Significant Loci With CRP Levels in the Replication Panel and Combined With the Discovery Results**

SNP	Coded Allele	Replication		Discovery + Replication		<i>R</i> <sup>2</sup> †	<i>P</i> for Heterogeneity	Closest Gene
		$\beta^*$ (SE)	<i>P</i>	Beta*(SE)	<i>P</i>			
rs2794520	C	0.086 (0.010)	$9.9 \times 10^{-19}$	0.160 (0.006)	$2.0 \times 10^{-186}$	1.38	$7.4 \times 10^{-26}$	<i>CRP</i>
rs4420638	A	0.200 (0.032)	$3.0 \times 10^{-10}$	0.236 (0.009)	$8.8 \times 10^{-139}$	0.93	0.03	<i>APOC1</i>
rs1183910	G	0.122 (0.021)	$8.3 \times 10^{-14}$	0.149 (0.006)	$2.1 \times 10^{-124}$	0.76	0.08	<i>HNF1A</i>
rs4420065	C	0.045 (0.009)	$1.5 \times 10^{-6}$	0.090 (0.005)	$3.5 \times 10^{-62}$	0.39	$1.1 \times 10^{-9}$	<i>LEPR</i>
rs4129267	C	0.045 (0.010)	$7.3 \times 10^{-6}$	0.079 (0.005)	$2.1 \times 10^{-48}$	0.31	$2.4 \times 10^{-4}$	<i>IL6R</i>
rs1260326	T	0.031 (0.010)	$1.9 \times 10^{-3}$	0.072 (0.005)	$4.6 \times 10^{-40}$	0.24	$2.6 \times 10^{-6}$	<i>GCKR</i>
rs12239046	C	0.042 (0.018)	$1.8 \times 10^{-3}$	0.047 (0.006)	$1.2 \times 10^{-15}$	0.09	0.77	<i>NLRP3</i>
rs6734238	G	0.072 (0.017)	$4.9 \times 10^{-6}$	0.050 (0.006)	$1.8 \times 10^{-17}$	0.14	0.95	<i>IL1F10</i>
rs9987289	A	0.003 (0.031)	$3.5 \times 10^{-2}$	0.069 (0.011)	$3.4 \times 10^{-13}$	0.08	0.04	<i>PPP1R3B</i>
rs10745954	A	0.018 (0.015)	$1.3 \times 10^{-1}$	0.039 (0.006)	$1.6 \times 10^{-11}$	0.06	$1.1 \times 10^{-3}$	<i>ASCL1</i>
rs1800961	C	0.023 (0.026)	$3.7 \times 10^{-1}$	0.088 (0.015)	$2.2 \times 10^{-9}$	0.06	0.07	<i>HNF4A</i>
rs340029	T	0.004 (0.010)	$5.2 \times 10^{-1}$	0.032 (0.006)	$4.1 \times 10^{-9}$	0.08	0.05	<i>RORA</i>
rs10521222	C	0.089 (0.028)	$1.4 \times 10^{-3}$	0.104 (0.015)	$8.5 \times 10^{-13}$	0.09	0.34	<i>SALL1</i>
rs12037222	A	0.035 (0.017)	$3.9 \times 10^{-2}$	0.045 (0.007)	$6.4 \times 10^{-11}$	0.06	0.40	<i>PABPC4</i>
rs13233571	C	0.049 (0.025)	$4.5 \times 10^{-2}$	0.054 (0.009)	$3.6 \times 10^{-9}$	0.08	0.13	<i>BCL7B</i>
rs2836878	G	0.013 (0.011)	$2.3 \times 10^{-1}$	0.032 (0.006)	$1.7 \times 10^{-7}$	0.05	0.18	<i>PSMG1</i>
rs4903031	G	0.001 (0.012)	$9.1 \times 10^{-1}$	0.032 (0.007)	$5.1 \times 10^{-6}$	0.04	0.21	<i>RGS6</i>

\* $\beta$  coefficient represents 1-unit change in the natural log-transformed CRP (mg/L) per copy increment in the coded allele.

†Median percentage of CRP variance explained by the SNP reported in all participating studies.

**Table 3. Association of 3 Suggestive Loci With CRP Levels That Reached Genome-Wide Significance After Combining Discovery and Replication Panel**

SNP	Coded Allele	Discovery		Replication		Discovery+Replication		$R^2$ †	$P$ for Heterogeneity	Closest Gene
		$\beta^*$ (SE)	$P$	$\beta^*$ (SE)	$P$	$\beta^*$ (SE)	$P$			
rs2847281	A	0.034 (0.007)	$1.7 \times 10^{-7}$	0.018 (0.016)	$4.2 \times 10^{-2}$	0.031 (0.006)	$2.2 \times 10^{-8}$	0.04	0.97	<i>PTPN2</i>
rs6901250	A	0.034 (0.007)	$1.2 \times 10^{-6}$	0.038 (0.015)	$1.2 \times 10^{-2}$	0.035 (0.006)	$4.8 \times 10^{-8}$	0.02	0.89	<i>GPRC6A</i>
rs4705952	G	0.038 (0.008)	$4.1 \times 10^{-6}$	0.065 (0.018)	$3.0 \times 10^{-4}$	0.042 (0.007)	$1.3 \times 10^{-8}$	0.05	0.47	<i>IRF1</i>

\* $\beta$  coefficient represents 1-unit change in the natural log-transformed CRP (mg/L) per copy increment in the coded allele.

†Median percentage of CRP variance explained by the SNP reported in all participating studies.

### Genetic Risk Score

To model the cumulative effect of the identified loci, we created a genetic risk score comprising information from the genome-wide significant SNPs. The risk score was computed for each subject by multiplying the number of alleles associated with higher CRP by the  $\beta$  coefficient from the combined meta-analysis and taking the sum over the SNPs. To make the genetic risk score easier to interpret, we rescaled to range from zero (low CRP level) to 100 (high CRP level).

### Association With MI and CHD

The association of the genome-wide significant SNPs and the genetic risk score with clinical events was tested in the ARIC study, the Age, Gene/Environment Susceptibility-Reykjavik (AGES) study, the Cardiovascular Health Study (CHS), the FHS, the Rotterdam Study (RS), and the WGHS with the use of incident cases of MI and CHD (ie, occurring after CRP concentrations were measured). Incident MI included fatal and nonfatal MI. Incident CHD included incident fatal and nonfatal MI, fatal CHD, and sudden death. Each study examined the associations with the use of a Cox proportional hazards model adjusted for age and sex. We subsequently combined these results by performing a meta-analysis.

### Results


The basic characteristics of the participating studies are shown in Table I in the online-only Data Supplement. Figure I in the online-only Data Supplement shows the QQ plot ( $\lambda=1.09$ ), and Figure II in the online-only Data Supplement presents the  $P$  values for >2.5 million SNPs across 22 autosomal chromosomes. A total of 953 SNPs in 17 loci exceeded the genome-wide significance threshold ( $P < 5 \times 10^{-8}$ ) (Table 1). Moreover, we found suggestive signals ( $P < 10^{-5}$ ) in 47 loci. Sixty-four lead SNPs including 17 SNPs from the genome-wide significant loci and 47 SNPs from the suggestive loci were chosen for the replication stage (Table II in the online-only Data Supplement). Six SNPs close to *CRP*, *APOC1*, *HNF1A*, *LEPR*, *IL6R*, and *IL1F10* exceeded the Bonferroni significance level ( $0.05/64 = 7.8 \times 10^{-4}$ ) in the replication stage. In a fixed-effects meta-analysis of the discovery and replication panels, 18 loci showed a genome-wide significant association: 15 loci of the 17 genome-wide significant loci (Table 2) and 3 loci of the 47 suggestive loci (Table 3). In addition to confirming 7 previously reported associations, the genome-wide significant signals marked 11 novel associations within or close to the NLR family, pyrin domain containing 3 (*NLRP3*), interleukin-1 family, member 10 (*IL1F10*), protein phosphatase 1, regulatory (inhibitor) subunit 3B (*PPP1R3B*), hepatocyte nuclear factor 4- $\alpha$  (*HNF4A*), RAR-related orphan receptor A (*RORA*), Sal-like 1 (*SALL1*), poly(A) binding protein, cytoplasmic 4 (inducible form) (*PABPC4*), B-cell chronic lymphocytic leukemia/lymphoma 7B (*BCL7B*), proteasome assembly chaperone 1 (*PSMG1*), protein

tyrosine phosphatase, nonreceptor type 2 (*PTPN2*), G protein-coupled receptor, family C, group 6, member A (*GPRC6A*), and interferon regulatory factor 1 (*IRF1*). Furthermore, our meta-analysis excluding the WGHS population (Table III in the online-only Data Supplement) confirmed the association of 7 previously known genes,<sup>9</sup> *CRP*, *APOE* (*APOC1*), *HNF1A*, *LEPR*, *IL6R*, *GCKR*, and *ASCL1*, with CRP levels (Bonferroni significance level:  $0.05/7 = 7.1 \times 10^{-3}$ ).


Figure 1 presents the average CRP levels across the genetic risk score in the whole population. Individuals in the highest gene score group had a mean CRP level (4.12 mg/L; 95% confidence interval, 4.96 to 5.25) that was more than double the level observed for individuals in the lowest gene score group (1.40 mg/L; 95% confidence interval, 1.31 to 1.49). The percentage of overall variance in CRP that was explained by the genetic risk score ranged from 1.2% to 10.3% across studies in the discovery and replication panels and was more than 5% in half of the studies.

After adjustment for number of tests, significant heterogeneity was found for rs2794520, rs4420065, rs4129267, rs1260326, and rs10745954 (Tables 2 and 3). Meta-regression was used to explore the source of heterogeneity. Sex was associated with heterogeneity for rs10745954 ( $P < 2.8 \times 10^{-5}$ ) (Table VI in the online-only Data Supplement).

All 18 SNPs that showed genome-wide significant results in the combined meta-analyses were studied for interactions



**Figure 1.** Mean CRP level (right vertical axis) shown as solid black dots connected by solid lines for categories of the genetic risk score. The shaded bars show the distribution of the genetic risk score in the whole population (left vertical axis). The CARdiovascular disease, Living and Aging in Halle (CARLA) Study was not included because of missing values for some of the selected SNPs.



**Figure 2.** The forest plots show the meta-analysis of the association of the CRP genetic risk score with MI (A) and CHD (B). The horizontal axis indicates the hazard ratio for MI or CHD per unit increase in the rescaled genetic risk score. ARIC indicates Atherosclerosis Risk in Communities; RS, Rotterdam Study; FHS, Framingham Heart Study; WGHS, Women's Genome Health Study; AGES, Age, Gene/Environment Susceptibility-Reykjavik; and CHS, Cardiovascular Health Study.

with age, sex, BMI, and smoking (Table IV in the online-only Data Supplement). After adjustment for the number of tests, we found a significant interaction between BMI and the *LEPR* SNP rs4420065 ( $P < 2.9 \times 10^{-6}$ ).

We examined the association of the SNPs related to CRP with risk of MI and CHD. These studies comprised 1845 cases of MI and 2947 cases of CHD. Neither the individual SNPs nor the combined genetic risk score showed consistent or genome-wide significant associations with risk of clinical events (Figure 2).

## Discussion

Through a meta-analysis of GWA scans from 15 cohort studies comprising 66 185 subjects and a replication sample of 16 540 subjects, we identified 18 loci associated with circulating CRP levels and provided evidence of replication for 8 of them. Our results confirm 7 gene-annotated loci reported by Ridker et al.<sup>9</sup> Furthermore, we introduce 11 novel loci associated with CRP levels, annotating *NLRP3*, *IL1F10*, *PPP1R3B*, *HNF4A*, *RORA*, *SALL1*, *PAPBC4*, *BCL7B*, *PTPN2*, *GPRC6A*, and *IRF1*.

A number of these genes including *APOC1*, *HNF1A*, *LEPR*, *GCKR*, *HNF4A*, and *PTPN2* are directly or indirectly related to

metabolic regulatory pathways involved in diabetes mellitus. Mutations in *HNF1A* are associated with impaired insulin secretion and maturity-onset diabetes mellitus of the young (MODY) type 3.<sup>19</sup> *HNF4A* is part of a complex regulatory network in the liver and pancreas for glucose homeostasis.<sup>20</sup> Mutations in the *HNF4A* gene cause MODY type 1.<sup>21</sup> *HNF4A* is a transcription factor involved in the expression of several liver-specific genes including *HNF1A*.<sup>21</sup> Defects in the expression of *GCKR* result in deficient insulin secretion.<sup>22</sup> *PTPN2*, which modulates interferon gamma signal transduction at the  $\beta$  cell level,<sup>23</sup> was recently identified as a novel susceptibility gene for type 1 diabetes mellitus.<sup>24</sup> *PTPN2* also is linked to the inflammatory pathway. The nuclear isoform of *PTPN2* is a regulator of transcription factor *STAT3* in the downstream of interleukin-6 signaling and may affect CRP expression in Hep3B cells.<sup>25</sup>

*CRP*, *IL6R*, *NLRP3*, *ILF10*, and *IRF1* are associated with CRP levels at least partly through pathways related to innate and adapted immune response. *NLRP3* encodes a member of the NALP3 inflammasome complex.<sup>26</sup> The NALP3 inflammasome triggers an innate immune response and can be activated by endogenous "danger signals," as well as compounds associated with pathogens.<sup>27,28</sup> Activated NALP3 inflammasome functions as an activator of nuclear factor- $\kappa$ B signaling. Nuclear factor- $\kappa$ B is a transcription factor that affects CRP expression in Hep3B cells.<sup>29</sup>

Our genetic risk score explained  $\approx 5\%$  of the variation in CRP levels, showing that genetic factors are of importance in determining CRP levels. In comparison, BMI as the main nongenetic determinant of CRP was reported to explain 5% to 7% of the variation in CRP levels in AGES<sup>30</sup> and up to 15% in FHS.<sup>31</sup> Ridker et al reported that 7 SNPs discovered in their study explained 10.1% of the variation in CRP levels after adjustment for age, smoking, BMI, hormone therapy, and menopausal status.<sup>9</sup> However, without adjustment for these covariates,  $< 5\%$  of the variation in CRP levels was explained (D. Chasman, PhD, personal written communication, May 2009).

Adipose tissue can induce chronic low-grade inflammation by producing proinflammatory cytokines such as interleukin-6.<sup>32</sup> Therefore, we examined whether adiposity modifies the effect of any of the 18 genes on CRP. We found that BMI modifies the strength of the association between *LEPR* and *CRP*. This interaction was initially found in WGHS.<sup>33</sup>

There is ample evidence that chronic inflammation is involved in atherosclerosis and cardiovascular disease. In this study, we found no association between genetically elevated CRP and risk of CHD. In agreement with our results, Elliott et al<sup>10</sup> reported in a recent study that variations in the *CRP* gene are not associated with risk of MI and CHD, but they found associations of *LEPR*, *IL6R*, and *APOC1-C1-C2* with CHD. However, the lack of association with clinical events in our study could also be due to lack of power.

Our study has the benefit of a large and homogeneous sample size of 82 725 subjects of European ancestry. This enabled us to find novel genes with small effect on CRP level. Furthermore, this large sample size enabled us to study gene-environment interaction, which hitherto has been less

feasible. In contrast to most other studies, we used only incident cases of cardiovascular events from well-defined population-based studies to examine the relation between the identified SNPs and clinical disease. The study has several limitations. Although we identified 18 loci associated with CRP levels, other genetic loci associated with CRP concentrations may still be missed by our study. Six of the genome-wide significant loci from the discovery panel were significant after Bonferroni correction in the replication panel. The other identified loci need replication for confirmation in larger samples. We acknowledge that our genetic risk score is based on our own findings and may be less efficient when used in another population. Finally, we did not fine map the identified loci; we therefore acknowledge that the identified SNPs may be in linkage disequilibrium with non-HapMap variants causally related to CRP levels.

In conclusion, we identified 11 novel loci and confirmed 7 known loci to affect CRP levels. The results highlight immune response and metabolic regulatory pathways involved in the regulation of chronic inflammation, as well as several loci previously unknown to be related to inflammation. Furthermore, *LEPR* was found to affect CRP differently in the presence of low or high BMI, which may lead to new insights in the mechanisms underlying inflammation.

### Acknowledgments

The following is a list of the author affiliations:

From the Department of Epidemiology, Erasmus Medical Center, Rotterdam, Netherlands (A.D., M.K., A.H., C.M.v.D., J.C.M.W.); Member of Netherlands Consortium for Healthy Aging sponsored by Netherlands Genomics Initiative, Leiden, Netherlands (A.D., M.K., A.H., A.G.U., C.M.v.D., J.C.M.W.); Department of Biostatistics, School of Public Health, Boston University, Boston, MA (J.D., C.L.); National Heart, Lung, and Blood Institute and Boston University's Framingham Heart Study, Framingham, MA (J.D., M.G.L., J.F.Y., J.F.K., J.D.F., R.S.V., E.J.B.); Human Genetics Center and Institute of Molecular Medicine, University of Texas Health Science Center at Houston (M.B., E.B.); Department of Medicine, University of Washington, Seattle (J.C.B.); Icelandic Heart Association, Kopavogur, Iceland (G.E., A.V.S., T.A., V.G.); Unit of Public Health Genomics, Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland (N.P., M.P.); Institute of Clinical Chemistry and Laboratory Medicine, University of Greifswald, Greifswald, Germany (H.W., M.N.); Department of Human Genetics, Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK (J.K., L.P.); Department of Human Genetics, Leiden University Medical Centre, Leiden, Netherlands (P.H.); Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany (J.B., B.T., C.G.); Division of Community Health Sciences, St George's University of London, London, UK (D.P.S.); Institute of Genetic Medicine, European Academy Bozen/Bolzano (EURAC), Bolzano, Italy (Affiliated Institute of University of Lübeck, Lübeck, Germany) (C.F., A.A.H., P.P.P.); Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, Western General Hospital, Edinburgh, UK (V.V., C.H.); Centre for Population Health Sciences, University of Edinburgh, Edinburgh, UK (J.F.W., I.R., H.C.); Center for Cardiovascular Disease Prevention, Harvard Medical School, Boston, MA (G.P., N.B.F.); Istituto di Neurogenetica e Neurofarmacologia, Consiglio Nazionale delle Ricerche, Cagliari, Italy (S.N., M.U.); Department of Epidemiology and Prevention, Wake Forest University School of Medicine, Wake Forest, NC (M.E.R., Y.L.); Institute for Molecular Medicine Finland FIMM, University of Helsinki, Helsinki, Finland (I.S., S.R., A. Sarin); Department of Biological

Psychology, VU University, Amsterdam, Netherlands (E.J.C.d.G., G.W., D.I.B.); Unit of Genetic Epidemiology and Bioinformatics, Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, Netherlands (B.Z.A., I.M.N., H.S.); Laboratory of Epidemiology, Demography and Biometry, National Institute on Aging, National Institutes of Health, Bethesda, MD (J.G., L.J.L., T.B.H.); Division of Endocrinology, Diabetes, and Nutrition, University of Maryland School of Medicine, Baltimore (A. Shuldiner; Q.D.G.); Clinical Research Branch, National Institute on Aging, Baltimore, MD (T.T., L.F.); Medstar Research Institute, Baltimore, MD (T.T.); Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA (R.Y.L.Z., L.M.R., J.E.B., D.B., P.M.R., D.I.C.); Department of Medicine, Johannes Gutenberg University, Mainz, Germany (R.B.S.); Department of Medicine, Baylor College of Medicine and Center for Cardiovascular Prevention, Methodist DeBakey Heart and Vascular Center, Houston, TX (V.N., R.C.H., B.C.A., J.C., C.M.B.); Department of Epidemiology, University of Washington, Seattle (N.L.S.); Seattle Epidemiologic Research and Information Center of the Department of Veterans Affairs Office of Research and Development, Seattle, WA (N.L.S.); Unit of Disease Risk, Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland (J.S.); Department of Epidemiology, MD Anderson Cancer Center, University of Texas, Houston (P.S., X.X.); Department of Clinical Chemistry, University of Oulu, Oulu, Finland (A.R.); Interfaculty Institute for Genetics and Functional Genomics, Ernst-Moritz-Arndt University Greifswald, Greifswald, Germany (A.T., G.H.); Departments of Pathology and Biochemistry, Colchester Research Facility, Colchester, VT (R.P.T.); Division of Epidemiology and Community Health, University of Minnesota, Minneapolis (A.R.F., J.P.); Department of Internal Medicine, Erasmus Medical Center, Rotterdam, Netherlands (E.J.G.S., A.G.U.); Medical Research Council-Health Protection Agency Centre for Environment and Health, Department of Epidemiology and Biostatistics, School of Public Health, St Mary's Campus, Imperial College London, London, UK (P.E., M.R.J.); Department of Biostatistics, Department of Genetics, University of North Carolina, Chapel Hill (W.S.); Amgen, Inc, Cambridge, MA (S.B., A.N.P., L.Y., K.T.); Department of Life Course and Services, National Institute for Health and Welfare, Helsinki, Finland (A.P., M.R.J.); Department of Medicine III, Martin Luther University, Halle-Wittenberg, Germany (K.W.); Institute for Medical Epidemiology, Biostatistics, and Informatics, Martin Luther University, Halle-Wittenberg, Germany (K.H.G., O.K.); Division of Cancer Epidemiology, German Cancer Research Centre, Heidelberg, Germany (K.H.G.); Department of Pharmacology, Ernst-Moritz-Arndt University of Greifswald, Greifswald, Germany (H.E.M.z.S.); Institute of Laboratory Medicine, Clinical Chemistry and Molecular Diagnostics, University of Leipzig, Leipzig, Germany (J.T.); Division of Clinical Developmental Sciences, St George's University of London, London, UK (Y.J.); Department of Twin Research and Genetic Epidemiology Unit, St Thomas' Campus, King's College London, St Thomas' Hospital, London, UK (Y.J.); Wellcome Trust Sanger Institute, Hinxton, UK (N.S.); Department of Twin Research and Genetic Epidemiology Unit, King's College London, London, UK (T.D.S.); Institute for Community Medicine, Ernst-Moritz-Arndt Universität Greifswald, Greifswald, Germany (H.V., D.R.); University of Iceland, Reykjavik, Iceland (T.A., V.G.); Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington, Seattle (D.S.S., B.M.P.); Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, CA (X.G., J.I.R.); Laboratory of Genetics, National Institute on Aging, Baltimore, MD (D.S.); Croatian Centre for Global Health, University of Split Medical School, Split, Croatia (I.R.); Department of Psychiatry/EMGO Institute/Neuroscience Campus, VU University Medical Centre, Amsterdam, Netherlands (B.W.P.); Institute of Health Sciences and Biocenter Oulu, Faculty of Medicine, University of Oulu, Oulu, Finland (M.R.J.); Department of Biostatistics, University of Washington, Seattle (K.R.); Unit of Chronic Disease Epidemiology and Prevention, Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland (V.S.); Departments of Internal Medicine and Human

Genetics, Leiden University Medical Centre, Leiden, Netherlands (K.W.v.D.); Preventive Medicine and Cardiology Sections, Department of Medicine, Boston University School of Medicine, Boston, MA (R.S.V., E.J.B.); Geriatric Unit, Azienda Sanitaria Firenze, Florence, Italy (S.B.); Department of Neurology, General Central Hospital, Bolzano, Italy (P.P.P.); Department of Neurology, University of Lübeck, Lübeck, Germany (P.P.P.); Group Health Research Institute, Group Health Cooperative, Seattle, WA (B.M.P.); and Department of Internal Medicine II–Cardiology, University of Ulm Medical Center, Ulm, Germany (W.K.).

See the online-only Data Supplement.

## Sources of Funding

See the online-only Data Supplement.

## Disclosures

Dr Ridker has received research grant support from Roche, AstraZeneca, and Amgen and is listed as a coinventor on patents held by the Brigham and Women's Hospital that relate to the use of inflammatory biomarkers in cardiovascular disease and diabetes mellitus.

## References

- Hindorf LA, Rice KM, Lange LA, Diehr P, Halder I, Walston J, Kwok P, Ziv E, Nievergelt C, Cummings SR, Newman AB, Tracy RP, Psaty BM, Reiner AP. Common variants in the CRP gene in relation to longevity and cause-specific mortality in older adults: the Cardiovascular Health Study. *Atherosclerosis*. 2008;197:922–930.
- Dehghan A, Kardys I, de Maat MP, Uitterlinden AG, Sijbrands EJ, Bootsma AH, Stijnen T, Hofman A, Schram MT, Witteman JC. Genetic variation, C-reactive protein levels, and incidence of diabetes. *Diabetes*. 2007;56:872–878.
- Sesso HD, Buring JE, Rifai N, Blake GJ, Gaziano JM, Ridker PM. C-reactive protein and the risk of developing hypertension. *JAMA*. 2003;290:2945–2951.
- Danesh J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, Rumley A, Lowe GD, Pepys MB, Gudnason V. C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med*. 2004;350:1387–1397.
- Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med*. 1997;336:973–979.
- Reitherstol L, Eikvar L, Berg K. A twin study of C-reactive protein compared to other risk factors for coronary heart disease. *Atherosclerosis*. 2003;169:279–282.
- Pankow JS, Folsom AR, Cushman M, Borecki IB, Hopkins PN, Eckfeldt JH, Tracy RP. Familial and genetic determinants of systemic markers of inflammation: the NHLBI Family Heart Study. *Atherosclerosis*. 2001;154:681–689.
- Dupuis J, Larson MG, Vasan RS, Massaro JM, Wilson PW, Lipinska I, Corey D, Vita JA, Keaney JF Jr, Benjamin EJ. Genome scan of systemic biomarkers of vascular inflammation in the Framingham Heart Study: evidence for susceptibility loci on 1q. *Atherosclerosis*. 2005;182:307–314.
- Ridker PM, Pare G, Parker A, Zee RY, Danik JS, Buring JE, Kwiatkowski D, Cook NR, Miletich JP, Chasman DI. Loci related to metabolic-syndrome pathways including LEPR, HNF1A, IL6R, and GCKR associate with plasma C-reactive protein: the Women's Genome Health Study. *Am J Hum Genet*. 2008;82:1185–1192.
- Elliott P, Chambers JC, Zhang W, Clarke R, Hopewell JC, Peden JF, Erdmann J, Braund P, Engert JC, Bennett D, Coin L, Ashby D, Tzoulaki I, Brown IJ, Mt-Isa S, McCarthy MI, Peltonen L, Freimer NB, Farrall M, Ruokonen A, Hamsten A, Lim N, Froguel P, Waterworth DM, Vollenweider P, Waeber G, Jarvelin MR, Mooser V, Scott J, Hall AS, Schunkert H, Anand SS, Collins R, Samani NJ, Watkins H, Kooner JS. Genetic loci associated with C-reactive protein levels and risk of coronary heart disease. *JAMA*. 2009;302:37–48.
- Reiner AP, Barber MJ, Guan Y, Ridker PM, Lange LA, Chasman DI, Walston JD, Cooper GM, Jenny NS, Rieder MJ, Durda JP, Smith JD, Novembre J, Tracy RP, Rotter JI, Stephens M, Nickerson DA, Krauss RM. Polymorphisms of the HNF1A gene encoding hepatocyte nuclear factor-1 alpha are associated with C-reactive protein. *Am J Hum Genet*. 2008;82:1193–1201.
- Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: an R library for genome-wide association analysis. *Bioinformatics*. 2007;23:1294–1296.
- Chen WM, Abecasis GR. Family-based association tests for genomewide association scans. *Am J Hum Genet*. 2007;81:913–926.
- Chen MH, Yang Q. GWAf: an R package for genome-wide association analyses with family data. *Bioinformatics*. 2010;26:580–581.
- Devlin B, Roeder K, Wasserman L. Genomic control, a new approach to genetic-based association studies. *Theor Population Biol*. 2001;60:155–166.
- Psaty BM, O'Donnell CJ, Gudnason V, Lunetta KL, Folsom AR, Rotter JI, Uitterlinden AG, Harris TB, Witteman JCM, Boerwinkle E; on behalf of the CHARGE Consortium. Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium: design of prospective meta-analyses of genome-wide association studies from 5 cohorts. *Circ Cardiovasc Genet*. 2009;2:73–80.
- Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26:2190–2191.
- Pe'er I, Yelensky R, Altshuler D, Daly MJ. Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genet Epidemiol*. 2008;32:381–385.
- Yamagata K, Oda N, Kaisaki PJ, Menzel S, Furuta H, Vaxillaire M, Southam L, Cox RD, Lathrop GM, Boriraj VV, Chen X, Cox NJ, Oda Y, Yano H, Le Beau MM, Yamada S, Nishigori H, Takeda J, Fajans SS, Hattersley AT, Iwasaki N, Hansen T, Pedersen O, Polonsky KS, Bell GI. Mutations in the hepatocyte nuclear factor-1alpha gene in maturity-onset diabetes of the young (MODY3). *Nature*. 1996;384:455–458.
- Urhammer SA, Fridberg M, Hansen T, Rasmussen SK, Moller AM, Clausen JO, Pedersen O. A prevalent amino acid polymorphism at codon 98 in the hepatocyte nuclear factor-1alpha gene is associated with reduced serum C-peptide and insulin responses to an oral glucose challenge. *Diabetes*. 1997;46:912–916.
- Yamagata K, Furuta H, Oda N, Kaisaki PJ, Menzel S, Cox NJ, Fajans SS, Signorini S, Stoffel M, Bell GI. Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature*. 1996;384:458–460.
- Kesavan P, Wang L, Davis E, Cuesta A, Sweet I, Niswender K, Magnuson MA, Matschinsky FM. Structural instability of mutant beta-cell glucokinase: implications for the molecular pathogenesis of maturity-onset diabetes of the young (type-2). *Biochem J*. 1997;322:57–63.
- Moore F, Colli ML, Cnop M, Esteve MI, Cardozo AK, Cunha DA, Bugliani M, Marchetti P, Eizirik DL. PTPN2, a candidate gene for type 1 diabetes, modulates interferon-gamma-induced pancreatic beta-cell apoptosis. *Diabetes*. 2009;58:1283–1291.
- Todd JA, Walker NM, Cooper JD, Smyth DJ, Downes K, Plagnol V, Bailey R, Nejentsev S, Field SF, Payne F, Lowe CE, Szeszek JS, Hafler JP, Zeitels L, Yang JH, Vella A, Nutland S, Stevens HE, Schuilenburg H, Coleman G, Maisuria M, Meadows W, Smink LJ, Healy B, Burren OS, Lam AA, Ovington NR, Allen J, Adlem E, Leung HT, Wallace C, Howson JM, Guja C, Ionescu-Tirgoviste C, Simmonds MJ, Heward JM, Gough SC, Dunger DB, Wicker LS, Clayton DG. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat Genet*. 2007;39:857–864.
- Yamamoto T, Sekine Y, Kashima K, Kubota A, Sato N, Aoki N, Matsuda T. The nuclear isoform of protein-tyrosine phosphatase TC-PTP regulates interleukin-6-mediated signaling pathway through STAT3 dephosphorylation. *Biochem Biophys Res Commun*. 2002;297:811–817.
- Petrilli V, Dostert C, Muruve DA, Tschopp J. The inflammasome: a danger sensing complex triggering innate immunity. *Curr Opin Immunol*. 2007;19:615–622.
- Dostert C, Petrilli V, Van Bruggen R, Steele C, Mossman BT, Tschopp J. Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science*. 2008;320:674–677.
- Muruve DA, Petrilli V, Zaiis AK, White LR, Clark SA, Ross PJ, Parks RJ, Tschopp J. The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature*. 2008;452:103–107.
- Voleti B, Agrawal A. Regulation of basal and induced expression of C-reactive protein through an overlapping element for OCT-1 and NF-kappaB on the proximal promoter. *J Immunol*. 2005;175:3386–3390.



30. Eiriksdottir G, Smith AV, Aspelund T, Hafsteinsdottir SH, Olafsdottir E, Launer LJ, Harris TB, Gudnason V. The interaction of adiposity with the CRP gene affects CRP levels: Age, Gene/Environment Susceptibility-Reykjavik study. *Int J Obes (Lond)*. 2009;33:267–272.
31. Kathiresan S, Larson MG, Vasan RS, Guo CY, Gona P, Keaney JF, Jr., Wilson PW, Newton-Cheh C, Musone SL, Camargo AL, Drake JA, Levy D, O'Donnell CJ, Hirschhorn JN, Benjamin EJ. Contribution of clinical correlates and 13 C-reactive protein gene polymorphisms to interindividual variability in serum C-reactive protein level. *Circulation*. 2006;113:1415–1423.
32. Yudkin JS, Kumari M, Humphries SE, Mohamed-Ali V. Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? *Atherosclerosis*. 2000;148:209–214.
33. Pare G, Cook NR, Ridker PM, Chasman DI. On the use of variance per genotype as a tool to identify quantitative trait interaction effects: a report from the Women's Genome Health Study. *PLoS Genet*. 2010;6:e1000981.

### CLINICAL PERSPECTIVE

C-reactive protein (CRP) is a heritable marker of chronic inflammation that is strongly associated with cardiovascular disease. Although environmental factors such as obesity, smoking, and hormone therapy influence levels of serum CRP, genes play an important role in determining serum CRP levels. The advent of genome-wide association studies has provided an opportunity to identify previously unsuspected genetic loci that influence complex traits. In this study, we collected data on >80 000 subjects from 25 studies and identified 18 genetic loci that are associated with serum CRP levels. These genetic loci provide valuable insights into the pathways that affect serum levels of CRP. Although further investigations are needed to understand the exact mechanisms, our findings highlight immune response and metabolic regulatory pathways involved in the regulation of chronic inflammation, as well as several loci previously unknown to be related to inflammation. However, these single-nucleotide polymorphisms were not associated with incident myocardial infarction or coronary heart disease, either individually or in combination. A better knowledge of the molecular mechanisms that control serum CRP levels may lead to a deeper understanding of the complex interactions underlying the inflammatory response in cardiovascular disease.