

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Metabolic versatility of the nitrite-oxidizing bacterium *Nitrospira marina* and its proteomic response to oxygen-limited conditions.

Permalink

<https://escholarship.org/uc/item/9gs8c2sn>

Journal

The ISME journal, 15(4)

ISSN

1751-7362

Authors

Bayer, Barbara
Saito, Mak A
McIlvin, Matthew R
et al.

Publication Date

2021-04-01

DOI

10.1038/s41396-020-00828-3

Peer reviewed



Metabolic versatility of the nitrite-oxidizing bacterium *Nitrospira marina* and its proteomic response to oxygen-limited conditions

Barbara Bayer¹ · Mak A. Saito² · Matthew R. McIlvin² · Sebastian Lücker³ · Dawn M. Moran² · Thomas S. Lankiewicz¹ · Christopher L. Dupont⁴ · Alyson E. Santoro¹

Received: 2 July 2020 / Revised: 20 October 2020 / Accepted: 30 October 2020 / Published online: 23 November 2020
© The Author(s) 2020. This article is published with open access

Abstract

The genus *Nitrospira* is the most widespread group of nitrite-oxidizing bacteria and thrives in diverse natural and engineered ecosystems. *Nitrospira marina* Nb-295^T was isolated from the ocean over 30 years ago; however, its genome has not yet been analyzed. Here, we investigated the metabolic potential of *N. marina* based on its complete genome sequence and performed physiological experiments to test genome-derived hypotheses. Our data confirm that *N. marina* benefits from additions of undefined organic carbon substrates, has adaptations to resist oxidative, osmotic, and UV light-induced stress and low dissolved $p\text{CO}_2$, and requires exogenous vitamin B₁₂. In addition, *N. marina* is able to grow chemoorganotrophically on formate, and is thus not an obligate chemolithoautotroph. We further investigated the proteomic response of *N. marina* to low ($\sim 5.6 \mu\text{M}$) O₂ concentrations. The abundance of a potentially more efficient CO₂-fixing pyruvate:ferredoxin oxidoreductase (POR) complex and a high-affinity *cbb₃*-type terminal oxidase increased under O₂ limitation, suggesting a role in sustaining nitrite oxidation-driven autotrophy. This putatively more O₂-sensitive POR complex might be protected from oxidative damage by Cu/Zn-binding superoxide dismutase, which also increased in abundance under low O₂ conditions. Furthermore, the upregulation of proteins involved in alternative energy metabolisms, including Group 3b [NiFe] hydrogenase and formate dehydrogenase, indicate a high metabolic versatility to survive conditions unfavorable for aerobic nitrite oxidation. In summary, the genome and proteome of the first marine *Nitrospira* isolate identifies adaptations to life in the oxic ocean and provides insights into the metabolic diversity and niche differentiation of NOB in marine environments.

Supplementary information The online version of this article (<https://doi.org/10.1038/s41396-020-00828-3>) contains supplementary material, which is available to authorized users.

✉ Barbara Bayer
bbayer@ucsb.edu

✉ Alyson E. Santoro
asantoro@ucsb.edu

¹ Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, CA, USA

² Marine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

³ Department of Microbiology, IWRW, Radboud University, Nijmegen, The Netherlands

⁴ J. Craig Venter Institute, La Jolla, CA, USA

Introduction

Aerobic nitrite (NO₂⁻) oxidation is the main biochemical nitrate (NO₃⁻)-forming reaction, carried out during the second step of nitrification [1]. In marine ecosystems, nitrate is the dominant form of biologically available nitrogen, which is rapidly assimilated by phytoplankton in surface waters and accumulates in the deep sea [2].

Nitrite-oxidizing bacteria (NOB) are chemolithoautotrophic microorganisms found within four known bacterial phyla (Proteobacteria, Nitrospirae, Nitrospinae, and Chloroflexi) [3]. The genus *Nitrospira*, within the Nitrospirae phylum, is the most diverse NOB genus and consists of at least six phylogenetic sublineages [3]. *Nitrospira* are ubiquitously present in natural and engineered ecosystems, including oceans [4, 5], freshwater habitats [6], soils [7, 8], saline-alkaline lakes [9], hot springs [10], wastewater treatment plants [11–13], and aquaculture biofilters [14, 15]. In human-made ecosystems, *Nitrospira* is generally

considered to be adapted to low NO_2^- concentrations [16]. In the open ocean, however, where NO_2^- concentrations are exceedingly low, NOB affiliated with the phylum Nitrospinae appear to be the dominant nitrite oxidizers [4], whereas *Nitrospira* are found in relatively high NO_2^- environments including sediments and deep sea hydrothermal vent plumes [17, 18]. *Nitrospira* also dominate over Nitrospinae-affiliated bacteria in some deep-sea trench environments [19, 20]. High NO_2^- concentrations are found coincident with low O_2 concentrations in oxygen minimum zone (OMZ) waters [21, 22] in a feature known as the secondary nitrite maximum with concentrations reaching $\sim 1\text{--}5\ \mu\text{M}\ \text{NO}_2^-$ [23, 24]. Despite the O_2 -dependence of all known NOB, NO_2^- oxidation can still be detected at nanomolar O_2 concentrations [25].

Some metabolic features appear to be common among *Nitrospira*, based on genomic analyses to date. Immunocytochemical analyses of a representative of *Nitrospira* sublineage I and metagenomic analyses of *Nitrospira defluvii* indicated a periplasmic orientation of nitrite oxidoreductase (NXR) [26, 27], the key enzyme for NO_2^- oxidation, and the presence of the O_2 -sensitive reductive tricarboxylic acid (rTCA) cycle for inorganic carbon fixation [26]. These results suggest that *Nitrospira* evolved from microaerophilic or anaerobic ancestors [26]. *N. defluvii* lacks genes for classical oxidative stress defense enzymes present in most aerobic organisms including catalase and superoxide dismutase (SOD), indicative of adaptations to low O_2 environmental niches [26]. However, other *Nitrospira* species encode both types of enzymes [28, 29], suggesting different O_2 tolerances within members of the *Nitrospira* genus. In addition to NO_2^- oxidation, *Nitrospira* exhibit a high metabolic versatility, growing aerobically on hydrogen [30] or anaerobically on organic acids while respiring nitrate [28]. Recently, the capability for the complete oxidation of ammonia to nitrate (comammox) was identified in representatives of sublineage II *Nitrospira* [14, 31], however, comammox *Nitrospira* appear to be absent in marine systems [31].

Nitrospira marina Nb-295^T, the type species of the genus *Nitrospira*, was isolated from a water sample collected at a depth of 206 m from the Gulf of Maine in the Atlantic Ocean over 30 years ago [5]. It is the only *Nitrospira* species isolated from the oceanic water column and the environmental conditions favorable for *Nitrospira* in the ocean remain largely unexplored. Here, we investigated the metabolic potential of *N. marina* Nb-295^T based on its complete genome sequence and compared the proteome signatures of cultures grown under atmospheric O_2 tension and under low O_2 conditions typically encountered in OMZs.

Materials and methods

Cultivation of *N. marina* Nb-295

A cryopreserved stock of *N. marina* Nb-295^T was obtained from the culture collection of John B. Waterbury at the Woods Hole Oceanographic Institution. Strain Nb-295 was grown in 60 mL polycarbonate bottles (Nalgene) in 50 mL autotrophic mineral salts medium at pH 7.8 containing 2 mM NaNO_2 (Table S1), and bottles were incubated at 25 °C in the dark without agitation. Mixotrophic growth was tested through the individual addition of the following organic carbon substrates to the culture medium of duplicate cultures (final concentrations): 150 mg L⁻¹ yeast extract, 150 mg L⁻¹ tryptone, 0.5 mM pyruvate, or 1 g L⁻¹ glycerol. NO_2^- consumption was measured as previously described [32] and growth was monitored by flow cytometry (see Supplementary Methods). To test for chemoorganotrophic growth, the culture was transferred (2% inoculum) into NO_2^- -free medium containing either 1 mM formate or 1 mM pyruvate, and 100 μM ammonium chloride, which served as nitrogen source.

For incubations at different O_2 concentrations, triplicate cultures of *N. marina* Nb-295 were grown at 22 °C in 400 mL of medium containing 70% natural seawater and 2 mM NaNO_2 as described by Watson et al. [5] in 500 mL polycarbonate bottles (Nalgene) with a custom-made sparging rig. Bottles were constantly bubbled (flow rate: 15 mL min⁻¹) with one of two sterile custom gas mixes containing either 0.5% or 20% oxygen, 300 ppm CO_2 , and a balance of high-purity N_2 . NO_2^- concentrations were measured as a proxy for growth as described above and 10 mL aliquot of each culture was fixed at the last time point (2% formaldehyde, 1 h, 4 °C) for cell enumeration on an epifluorescence microscope as previously described [33].

DNA extraction, genome sequencing, and annotation

High molecular weight genomic DNA was extracted from stationary phase cultures using a CTAB extraction protocol [34] and sequenced on the PacBio platform at the US Department of Energy Joint Genome Institute (JGI). 754,554 reads were produced, with 209,987 passing quality control. The assembly was conducted using HGAP (v 2.2.0.p1) with improvement with Quiver [35] resulting in a single contig.

Gene annotation was conducted using JGI's Integrated Microbial Genomes and Microbiomes pipeline [36] and the MicroScope platform [37]. Manual curation included sequence similarity searches using BLASTP (e-value $< 1\text{e}^{-30}$) [38] against the Transporter Classification database [39] and protein domain searches using

InterProScan (release 72.0) [40]. Signal peptides were identified with SignalP 5.0 [41] to determine if proteins were potentially addressed to the membrane and/or released to the periplasmic space. A list of manually curated annotations can be found in Data Set 1.

Phylogenomic analysis was performed using 120 concatenated phylogenetic marker genes from representatives of the phylum Nitrospirae/Nitrospirota as implemented in the Genome Taxonomy Database Toolkit (GTDB-tk) version 1.1.1 [42]. (see Supplementary Methods).

Protein extraction and proteome analyses

Cells for proteomic analysis were harvested when $[\text{NO}_2^-]$ dropped to $\sim 500 \mu\text{M}$, corresponding to exponential growth of strain Nb-295. Each culture was mixed with an equal volume of a house-made fixative [43] similar to the commercially available solution RNeasy Lysis Buffer (Qiagen), and filtered by vacuum filtration onto 25 mm, 0.2 μm pore size Supor filters (Pall). 200 mL of fixed culture (equivalent to 100 mL of growth medium) were filtered for protein extraction and proteomic analyses. Filters were frozen at -80°C until extraction. A detailed description of the procedures used for protein extraction and purification can be found in the Supplementary Methods. Global (untargeted) proteomes were analyzed on a Fusion Orbitrap mass spectrometer using one-dimensional nanospray separation and data-dependent acquisition based on Saito et al. [44] (see Supplementary Methods for detailed protocols). Select targeted quantitative proteomic assays using custom-made isotopically-labeled (^{15}N) peptide standards were designed and samples were analyzed again by parallel reaction monitoring mass spectrometry using mass spectral information from the global proteome analyses as previously described [44] (also see Supplementary Methods). Tryptic peptides from ten proteins of interest, including nitrite oxidoreductase subunit alpha (NxrA), were targeted for absolute quantitation (see Data Set 2). Cellular NXR concentrations were calculated based on NxrA peptide concentrations using the average carbon content of *N. marina* cells (152 fg cell $^{-1}$, Santoro et al., *unpublished*) and an estimated cellular protein:carbon ratio of 50% based on experimentally determined values [45]. NXR complex density on the cellular membrane was calculated using an estimated NXR complex size of 63 nm 2 [27] and an estimated cell surface area of 2.45 μm^2 (assuming a cylindrical shape with a length of 1.75 μm and a radius of 0.2 μm based on previously determined cell dimensions of *N. marina* [5]).

Differential levels of expression between the two treatments (i.e., atmospheric O_2 and low O_2 conditions) were tested with the DESeq2 Bioconductor package (version 1.20.0) [46] in the R software environment (version 3.5.0) [47] using spectral counts as input data as previously described [48]. Proteins with

a mean spectral count below 6 across all treatments were excluded from the analysis. In DESeq2, only proteins that increased in abundance under low O_2 conditions were considered ('altHypothesis=greater'), P values were adjusted using the Benjamini-Hochberg method ('pAdjust-method=BH') and independent filtering was omitted ('independentFiltering=FALSE'). Changes in protein abundance (as determined by spectral counts) were considered statistically significant when adjusted P values were lower than or equal to 0.05 (see Data Set 3). While DESeq2 has a high precision and accuracy [49], it is more conservative than other methods on low-count transcripts/proteins [50]. Proteins of interest were visualized with the pheatmap package (version 1.0.12) [51] in the R software environment [47]. The normalized spectral abundance factor was calculated as proxy for relative protein abundances [52], and values were square-root transformed to improve visualization of low abundant proteins.

Results and discussion

Genome analysis

The genome of *N. marina* Nb-295 is a single element of 4,683,627 bp and contains 4272 coding DNA sequences (CDS) including one rRNA operon. A 5578 bp region of 99.7% identity on each end of the scaffold suggests circularization into a single chromosome. No plasmids or extrachromosomal elements were identified. The G + C content is 50.04%, lower than in other *Nitrospira* species and the marine nitrite-oxidizers *Nitrospira gracilis* and *Nitrococcus mobilis* (Table S2). Phylogenomic analysis of available closed genomes, metagenome-assembled genomes (MAGs) and single-cell amplified genomes affiliated with the phylum Nitrospirae (see Supplementary Methods) placed *N. marina* Nb-295 T within a cluster of genomes derived from marine and saline environments (Fig. 1). The most closely related *Nitrospira* MAG (UBA8639) was obtained from a laboratory-scale nitrification reactor; however, the reactor influent consisted of 33% untreated seawater [53], suggesting a marine origin of this MAG. The 16S rRNA gene sequence of *N. marina* Nb-295 T clustered together with environmental sequences derived from marine sediments and marine aquaculture biofilters (Fig. S1), and shared 99.1% and 97.9% sequence identity with the cultured lineage IV representatives *Nitrospira* sp. Ecomares 2.1 [15] and *Ca. Nitrospira alkalitolerans* [9], respectively.

Nitrogen metabolism and respiratory chain

The genome of *N. marina* encodes orthologs of all known proteins required for NO_2^- oxidation, including the putatively periplasm-oriented NXR complex. Three candidates




Fig. 1 Maximum likelihood phylogenetic tree of representatives of the phylum Nitrospirae. The multiple sequence alignment consisting of 120 concatenated phylogenetic marker genes contained 95 genomes and metagenome-assemble genomes (MAGs) from the Genome Taxonomy Database (GTDB) (Release 04-RS89, 19 June 2019), the genome of *N. marina*, the MAG of *Ca. N. alkalitolerans* and two open

ocean single-cell amplified genomes (SAGs), AC-738-G23 and AC-732-L14. All MAGs and SAGs were estimated to be $\geq 50\%$ complete with $\leq 5\%$ contamination. Nodes with UFBoot support of at least 95% are indicated as black filled circles. Cultured representatives are shown in bold. The scale bar represents 0.1 substitutions per site.

each of genes for NxrA, NxrB, and NxrC, and two additional NxrC-like proteins were identified (Table S2, Data Set 1). The genes *nxrA* and *nxrB*, encoding the alpha and beta subunits of NXR, are colocalized in three clusters, whereas all *nxrC* candidate genes are localized separately from *nxrAB*, as previously described for *Nitrospira moscoviensis* [54]. The NxrA subunits share 87.3–88.9% amino acid identity, the NxrB subunits share 98.8–99.5% amino acid identity, and the putative NxrC subunits are less

conserved, sharing between 33.7% and 86.9% amino acid identity.

Like all other analyzed *Nitrospira* genomes, *N. marina* encodes a putative copper-dependent NO-forming nitrite reductase (NirK), yet its function in *Nitrospira* and other NOB remains unknown. *N. marina* also encodes the ferredoxin-dependent nitrite reductase (NirA) for assimilatory nitrite reduction, which appears to be conserved in *Ca. Nitrospira lenta* and *N. defluvii* but absent in other

Nitrospira [55]. In addition, *N. marina* encodes three high affinity ammonium transporters (Amt) enabling direct uptake of reduced N for assimilation, and cyanate lyase to hydrolyze cyanate to ammonium and CO₂ (Data Set 1). In contrast to some *Nitrospira* species [13, 28, 31], *N. marina* does not encode a urea transporter or urease, which would catabolize urea to ammonia for N assimilation (Table S2).

Previously sequenced genomes of *Nitrospira* contain multiple copies of several complexes of the respiratory chain [26, 28, 29, 54]. *N. marina* encodes two paralogous copies of complex I, one of which contains a duplication of NADH:quinone oxidoreductase subunit M (NuoM) and lacks genes for NuoE, NuoF and NuoH (Data Set 1), which is a characteristic feature of *Nitrospira* genomes [54, 56]. Furthermore, the *N. marina* genome contains two copies of complex III, two cytochrome *bd* oxidases and seven putative cytochrome *bd*-like oxidases (Data Set 1), which show limited partial similarity to canonical cytochrome *bd* oxidase subunit I as described for *N. moscoviensis* [54]. One of these cytochrome *bd*-like oxidases (*bd*-like_6) contains putative heme b and copper binding sites potentially functioning as a novel terminal oxidase as previously proposed [26]. *N. marina* also encodes for a *cbb*₃-type terminal oxidase, which usually exhibit high affinities for O₂ [57]. This feature is shared with the closely related *Ca. N. alkalitolerans* [9] and with the more distantly related marine NOB *Nitrospira gracilis* [58], but absent in all other *Nitrospira* species sequenced thus far. In addition to the canonical H⁺-translocating F₁F₀-ATPase (complex V), *N. marina* also encodes a putative alternative Na⁺-translocating N-ATPase (Data Set 1), which potentially contributes to the maintenance of the membrane potential and the generation of a sodium motive force (SMF) as suggested for *Ca. N. alkalitolerans* [9]. Furthermore, a H⁺-translocating pyrophosphatase (H⁺-PPase) with homology to *Leptospira* protozoan/plant-type enzymes [59] was identified. H⁺-PPases couple the translocation of H⁺ to the hydrolysis of the biosynthetic by-product pyrophosphate (PP_i), which is suggested to be an adaptation to life under energy limitation [60]. The *N. marina* genome also contains an alternative complex III (ACIII) module, which shares similarity with that from sulfur-reducing Acidobacteria [61]. Like canonical complex III, ACIII also functions as a quinol oxidase transferring electrons to cytochrome *c* and contributes to energy conservation (Refojo et al., 2012). With the exception of the comammox bacterium *Ca. Nitrospira nitrificans* [14], no homologs of ACIII modules were identified in any other NOB genome.

N. marina encodes a putative *hyb*-like operon containing four subunits of a cytoplasmic Group 3b [NiFe] hydrogenase and six accessory proteins involved in hydrogenase assembly and maturation (Data Set 1). This type of

hydrogenase appears to be conserved in the marine NOB *Nitrospira gracilis* 3/211 and *Nitrococcus mobilis* Nb-231 [58, 62], *Ca. N. alkalitolerans* [9] and in comammox *Nitrospira* [14, 31] (Table S2). In addition to catalyzing the reversible, NAD⁺-dependent oxidation of hydrogen, these so-called sulfhydrogenases have been reported to reduce elemental sulfur (S⁰) or polysulfides to hydrogen sulfide (H₂S) [63]. Furthermore, the *N. marina* genome encodes a putative periplasmic sulfite:cytochrome *c* oxidoreductase, which might couple sulfite (SO₃²⁻) oxidation to sulfate (SO₄²⁻) with the reduction of cytochrome *c* as previously suggested for *Nitrospira gracilis* [58]. Contrastingly, sulfide/quinone oxidoreductase, which is speculated to mediate sulfide oxidation in *Nitrococcus* [62], is lacking. Whether or not these enzymes are involved in energy conservation using H₂S and SO₃²⁻ as alternative substrates in NOB remains to be experimentally validated.

Central carbon metabolism

In agreement with other *Nitrospira* genomes [13, 26, 54, 55], *N. marina* encodes the complete gene repertoire for the rTCA cycle for carbon dioxide (CO₂) fixation, including the key enzymes ATP-citrate lyase and 2-oxoglutarate/pyruvate:ferredoxin oxidoreductase (Data Set 1). In the ocean, inorganic carbon is predominately available in the form of bicarbonate (HCO₃⁻) and to a much lesser extent as CO₂. Five inorganic anion transporters (SulP family) with homology to BicA HCO₃⁻ uptake systems of the cyanobacterium *Synechococcus* [64] were identified in the *N. marina* genome (Data Set 1). Two of these putative BicA-like transporters are colocalized with genes encoding Na⁺/H⁺ antiporters (NhaB family), which could drive the uptake of HCO₃⁻ via Na⁺ extrusion under alkaline conditions as suggested for the cyanobacterium *Aphanothece halophytica* [65] and *Ca. N. alkalitolerans* [9]. *N. marina* also encodes one putative SulP-related bicarbonate transporter fused to a carbonic anhydrase and four genes encoding putative alpha, beta and gamma carbonic anhydrases (Data Set 1), which can convert the imported HCO₃⁻ to CO₂ for inorganic carbon fixation via the rTCA cycle.

In addition to the rTCA cycle, *N. marina* encodes all required genes for the oxidative TCA cycle for pyruvate oxidation via acetyl-CoA, complete gluconeogenesis and glycolysis pathways, and the oxidative and non-oxidative branches of the pentose phosphate pathway (Data Set 1), which are common features of all sequenced *Nitrospira* genomes [9, 26, 28, 29, 55]. Furthermore, biosynthetic pathways for all amino acids except methionine were identified in the *N. marina* genome. Although *N. marina* encodes a vitamin B₁₂-dependent methionine synthase (MetH) (Data Set 1), it appears to lack additional enzymes of known methionine biosynthesis pathways, a trait shared

by all other sequenced *Nitrospira* species and *Nitrospina gracilis* [9, 26, 28, 29, 55, 58]. However, as *N. marina* can grow in artificial seawater medium without added methionine, we hypothesize that an alternative unknown pathway for the early steps of methionine biosynthesis functions in *Nitrospira* and *Nitrospina*. The *N. marina* genome contains genes for the biosynthesis and degradation of the storage compounds glycogen and polyphosphate (Data Set 1). In contrast to other *Nitrospira* and the marine NOB *N. gracilis* and *N. mobilis* that encode a *glgA*-type glycogen synthase, *N. marina* encodes alpha-maltose-1-phosphate synthase (*glgM*) and alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase (*glgE*) for the synthesis of glycogen via alpha-maltose-1-phosphate.

Use of organic carbon substrates

N. marina has been reported to be an obligate chemolithotroph that grows best in medium supplemented with low concentrations of organic compounds including pyruvate, glycerol, yeast extract and peptone [5]. Thus, we investigated the genomic basis for this observation and conducted additional physiological experiments.

In addition to its complete glycolysis pathway and oxidative TCA cycle, a putative carbohydrate degradation operon was identified, consisting of a sugar ABC transporter module, beta-glucosidase, a putatively secreted glycoside hydrolase (GH15) and a carbohydrate-binding protein (Data Set 1). *N. marina* also encodes two putative carbohydrate-selective porins (OprB), a sugar:sodium symporter (SSS family), a putative galactonate/glucarate transporter (MFS superfamily) and a putative carboxylate transporter (DASS family). The genomic repertoire for the catabolic degradation and assimilation of peptides and amino acids, including transporter proteins for di- and oligopeptides (ABC and POT/PTR families), multiple amino acid:cation symporters (SSS, DAACS and AGCS families), amino acid/polyamine transporters (APC superfamily) and multiple putatively secreted peptidases are present (Data Set 1).

In agreement with Watson et al. [5], NO_2^- oxidation activity was enhanced when undefined organic compound mixtures such as tryptone and yeast extract were added to the culture medium (Fig. 2). Interestingly, growth was greatly stimulated by tryptone (Fig. 2), while amendment with defined organic carbon compounds had no effect on NO_2^- oxidation or growth (Table S3). Parallel incubations with ammonium as an N source did not increase activity or growth (Table S3), suggesting that the stimulating effect of tryptone, and to a lesser extent of yeast extract, most likely can be attributed to direct amino acid assimilation and does not only reflect the eliminated energy demand for assimilatory NO_2^- reduction. No growth on yeast extract and




Fig. 2 The effect of undefined organic carbon substrates on nitrite oxidation and growth of *N. marina* Nb-295^T. Error bars represent the range of measurements from duplicate cultures.




Fig. 3 Growth of *N. marina* Nb-295^T on formate (1 mM), pyruvate (1 mM) and yeast extract and tryptone (150 mg L $^{-1}$ each) in the absence of nitrite. Error bars represent the range of measurements from duplicate cultures.

tryptone was observed in the absence of NO_2^- (Fig. 3), corroborating their use as source of amino acids rather than for energy conservation.

In addition to undefined organic carbon substrates, defined organic compounds such as glycerol and pyruvate have been reported to enhance the growth of *N. marina* and *N. defluvii*, respectively [5, 12]. Formate has also been shown to serve as an electron donor and carbon source for some lineage I and II *Nitrospira* [11, 28]. *N. marina* encodes a putative formate dehydrogenase (FdhA) (Data Set 1), which is divergent from those found in *N. moscoviensis* and *N. defluvii* (~24 and 27% amino acid identity, respectively), but shares a relatively high sequence similarity

(~48% amino acid identity) to the functionally characterized formate dehydrogenase Fdh4 from *Methylobacterium extorquens* [66]. *N. marina* was able to grow chemoorganotrophically on 1 mM formate in the absence of NO_2^- (Fig. 3) and is thus not an obligate chemolithotrophic organism. However, the use of formate instead of NO_2^- as electron donor resulted in slower growth rates, as previously also observed in *N. moscoviensis* [28]. In contrast to earlier reports [5], additions of 1 g L^{-1} glycerol did not stimulate NO_2^- oxidation activity or growth (Table S3). Furthermore, pyruvate could neither be used as alternative energy source (Fig. 3), nor did it stimulate metabolic activity in the presence of NO_2^- (Table S3).

Protection against oxidative, osmotic, and UV light-induced stress

The formation of reactive oxygen species is prevalent in oxic environments and oxidative stress defense is an important component of the stress response in marine organisms [67]. *N. marina* encodes multiple enzymes to reduce oxidative stress, including a cytoplasmic Mn/Fe-binding SOD, a periplasmic Cu/Zn-binding SOD, two heme-containing catalases, and various peroxiredoxins (Data Set 1). In contrast, *Nitrospina gracilis* and marine ammonia-oxidizing archaea lack catalase [48, 58], suggesting that *N. marina* is less susceptible to oxidative stress compared to other marine nitrifiers. In addition to its plethora of oxidative stress defense-related proteins, *N. marina* also encodes two putative photolyases—enzymes known to be involved in the repair of UV light-induced DNA damage [68]—suggesting that it is well adapted to conditions characteristic for euphotic environments.

Marine microorganisms counteract the external osmotic stress from high salt concentrations by accumulating a variety of organic solutes (=osmolytes) in the cytoplasm, which can either be synthesized de-novo or transported into the cell from the surrounding environment [69]. In addition to select amino acids that can serve as compatible solutes (e.g., proline and glutamate) [70], biosynthesis pathways for the osmolytes glycine betaine and trehalose were identified in the *N. marina* genome (see Supplementary Results and Discussion). The production and concomitant release of osmolytes (i.e., via diffusion, excretion, predation or cell lysis) could potentially fuel heterotrophic metabolism in the ocean [71] representing a link between chemolithoautotrophic production and heterotrophic consumption of DOM as recently suggested for ammonia-oxidizing archaea [72].

Vitamin B₁₂ auxotrophy

B vitamins are important biochemical co-factors required for cellular metabolism and their concentrations are

depleted to near zero across large areas of the global ocean [73]. *N. marina* encodes complete biosynthetic pathways for the B vitamins thiamin (B₁), riboflavin (B₂), pantothenate (B₅), pyridoxine (B₆), biotin (B₇) and tetrahydrofolate (B₉). However, an incomplete vitamin B₁₂ biosynthesis pathway was identified (Fig. S2), lacking genes for multiple precorrin conversion reactions that ultimately lead to the biosynthesis of the molecule's corrin ring [74]. Since *N. marina* only encodes the cobalamin-dependent versions of ribonucleotide reductase, methionine synthase and methylmalonyl-CoA mutase, it must rely on the supply of vitamin B₁₂ or its precursors by other members of the microbial community. Indeed, nitrite consumption by *N. marina* ceased after repeated transfers into an artificial seawater mineral medium without the addition of vitamin B₁₂ (Fig. 4), and nitrite oxidation activity was restored after adding vitamin B₁₂ to B₁₂-deplete cultures (data not shown). Interestingly, the *N. marina* genome contains genes for multiple reactions that convert the precursor cobyrinate/hydrogenobyrate to cobalamin, and encodes all genes required for cobalamin salvage from cobinamide (Fig. S2, Data Set 1). Hence, in contrast to many other bacteria that lack the complete vitamin B₁₂ biosynthesis pathway [75], *N. marina* appears to obtain its B₁₂ from salvage of multiple intermediates from the environment. Vitamin B₁₂ auxotrophy has recently also been observed for several marine Nitrospinae strains [76]. In contrast, Park et al. [77] suggested that the two novel marine NOB strains MSP and DJ, one of which is closely related to *N. marina*, might be able to synthesize cobalamin. However, given that these cultures were not axenic and that genes involved in precorrin conversion reactions absent in *N. marina* (Fig. S2) are also missing from their genomes, it is likely that both strains depend on the exogenous supply of vitamin B₁₂ as well.

When *N. marina* was grown in natural seawater, cellular concentrations of ribonucleotide reductase were approximately 39-fold higher than for the cyanobacterium




Fig. 4 Nitrite consumption by *N. marina* Nb-295^T in artificial seawater mineral medium with and without addition of vitamin B₁₂ (Table S1). Cultures of strain Nb-295^T were repeatedly transferred into medium without vitamin B₁₂ using 2% inocula. Error bars represent the range of measurements from duplicate cultures.

Table 1 Targeted quantitative analyses of selected proteins using isotopically-labeled (^{15}N) peptide standards.

Target protein ^a	^{15}N -Peptide sequence	Peptide concentration (copies cell ⁻¹)	
		atm. O ₂	low O ₂
Nitrite oxidoreductase, alpha subunit 1	LVVITPEYNPTAYR	845 (52)	1924 (275)
	DYAFPDFANSYSGK	241 (40)	451 (91)
	HPFWEETNESKPQWTR	673 (39)	1571 (388)
Nitrite oxidoreductase, alpha subunit 2	VVVITPEYNPTAQR	1,154 (27)	851 (187)
Nitrite oxidoreductase, alpha subunit 3	DYQFPDFTSTYSGK	1,761 (159)	1,261 (242)
	SGIDPALTGTHR	4,519 (516)	3,400 (526)
	IAVITPEYNPTAYR	4,860 (496)	3,741 (710)
Nitrite oxidoreductase, alpha subunit (all)	GWKPSDPYK	11,467 (1,957)	9,564 (1,083)
	AIALDTGYQSNFR	13,996 (962)	13,116 (2,934)
Pyruvate:ferredoxin oxidoreductase, alpha subunit	TPSFFTGSEVIK	2,026 (192)	1,550 (248)
	EAIAILLEEGER	1,362 (80)	1,054 (170)
	EVSATVPNNER	2,453 (169)	1,736 (215)
Ribonucleotide reductase	TGESPYQTIPFSHR	300 (46)	198 (30)
	EAAVPEPYIHR	705 (132)	472 (77)
	IINQSLPPALR	750 (88)	546 (68)

Nitrite oxidoreductase peptide targets were designed to match a specific NxrA copy (1, 2, or 3), or to match all three as indicated in the target protein column. Values represent the mean of triplicate measurements and standard deviations are shown in brackets.

^aA complete list of quantified proteins can be found in Data Set 2.

Prochlorococcus [78] (Table 1) despite only having a ~two-fold greater cell volume, consistent with the importance of B₁₂ nutrition to *N. marina*. Moreover, this salvage acquisition mode is interesting in the context of dissolved cobalt speciation, which is complexed by strong organic ligands hypothesized to include B₁₂ precursors or degradation products [79]. In the Northwest Atlantic near where *N. marina* was isolated, organic cobalt complexes are abundant, comprising about half the dissolved cobalt inventory [80].

Metabolic response to low oxygen concentrations

Given the presence of multiple signatures of micro-aerophilic adaptation and metabolic diversity of nitrite oxidizers [26, 28, 58, 62] and their occurrence in low oxygen environments [25, 81, 82], we sought to further investigate potential adaptations of *Nitrospira marina* Nb-295^T to low O₂ conditions. *N. marina* was grown at O₂ concentrations characteristic for the upper ocean (~200 μM) and at O₂-limiting conditions (~5.6 μM O₂) found in environments with elevated NO₂⁻ concentrations such as OMZ or sediments [83, 84].

When grown under atmospheric O₂ concentration, *N. marina* oxidized 1.5 mM NO₂⁻ within 12 days, whereas under O₂-limiting conditions depletion of 1.5 mM substrate took 27 days (Fig. 5). This result is in agreement with the initial description by Watson et al. [5], who reported partial,




Fig. 5 Nitrite consumption by *N. marina* Nb-295^T grown under atmospheric (filled circles) and low O₂ conditions (open circles). Cells for proteome analysis were harvested after 12 and 27 days, respectively. Error bars represent standard deviations from measurements of triplicate cultures.

though unquantified, inhibition at low O₂ partial pressure. Cell abundances at the final timepoint (after 1.5 mM NO₂⁻ was oxidized) were comparable for both treatments (Fig. S3), indicating that the reduced NO₂⁻ oxidation rate during O₂-limiting conditions ultimately resulted in similar cell yields.

General proteomic response and upregulation of gene clusters

Cultures grown under both O₂ treatments were harvested for proteomic analysis when [NO₂⁻] dropped to ~500 μM,

corresponding to exponential growth of strain Nb-295 (see “Material and Methods”). A total of 2031 and 2046 proteins were identified by liquid chromatography-tandem mass spectrometry in the atmospheric and low O₂ treatments, respectively, accounting for 48.1 and 48.5% (49.7% combined from a total of 175,653 peptides) of the predicted protein coding sequences in the *N. marina* genome. As previously reported for *N. marina* and *Nitrococcus mobilis*, proteins exhibiting the highest abundances were associated with NO₂⁻ oxidation [44] (Data Set 1). NXR made up on average 4% of all peptide spectral counts and cellular NXR concentrations were ~13,500 copies cell⁻¹ as determined by targeted quantitative proteomic analyses (Table 1), covering an estimated 35% of the membrane surface (see “Material and Methods”). All three NxrA copies were detected in the proteome as determined by detection of unique peptides of each, and NxrA₃ appeared to be more abundant compared to NxrA₁ and NxrA₂ (Table 1). Under low O₂ conditions, NxrA₁ increased in abundance (Table 1, Fig. 6) indicating different metabolic or regulatory roles of the highly similar subunits.

Proteins involved in CO₂ fixation, DNA replication, electron transport and central carbon metabolism were also highly abundant under both conditions, indicating that *N. marina* retained its central metabolism during O₂ deficiency (Data Set 1). Although proteomic spectral counts remained constant for the majority of proteins during both treatments, spectral counts of 93 proteins significantly increased (adjusted *P* value ≤ 0.05) during growth at low O₂ concentrations (Fig. 6, Data Set 3). These results are supported by the targeted quantitative proteomic analyses (Fig. S4, Data Set 2), suggesting that spectral counts are a good proxy for changes in absolute protein abundance in our dataset.


Multiple universal stress proteins (UspA superfamily) were among the proteins that showed the highest increase in abundance under low O₂ conditions compared to the control treatment (Fig. 6, Data Set 3). UspA proteins have versatile regulatory and protective functions to enable survival under diverse external stresses [85], and are induced during growth inhibition [86] and oxygen starvation [87]. In *N. marina*, all four upregulated UspA proteins are located upstream or downstream of operons containing genes that also increased in abundance under low O₂ conditions (Fig. 6), suggesting a regulatory role of UspA-related proteins upon O₂ limitation. These upregulated gene clusters include proteins involved in electron transport, carbon metabolism, and alternative energy metabolism (Fig. 6).

In addition to putative UspA-regulated gene clusters, a gene cluster containing type VI secretion system (T6SS)-related proteins exhibited higher abundances under low O₂ conditions (Fig. 6). The T6SS is typically involved in the

secretion of effectors required for pathogenesis, bacterial competition, biofilm formation, and cell communication (e.g., quorum sensing) [88, 89]. While quorum sensing has recently been shown for diverse NOB including *N. moscoviensis* [90], no LuxI autoinducer synthases and/or LuxR signal receptor homologs were identified in the *N. marina* genome.

Induction of a putative O₂-sensitive 2-oxoacid:ferredoxin oxidoreductase complex

The key enzymes of the rTCA cycle, pyruvate:ferredoxin oxidoreductase (POR) and 2-oxoglutarate:ferredoxin oxidoreductase (OGOR), are typically highly O₂ sensitive because they contain easily oxidized iron-sulfur clusters [91]. In *Hydrogenobacter thermophilus*, five-subunit O₂-tolerant forms of POR and OGOR mainly support aerobic growth, while a O₂-sensitive two-subunit form is used under anaerobic conditions [92]. *N. marina* encodes three 2-oxoacid:ferredoxin oxidoreductase gene clusters that could exhibit POR or OGOR activity (Data Set 1). Two of these gene clusters consist of five CDS that exhibit a high sequence similarity to the O₂-tolerant five-subunit POR/OGOR of *H. thermophilus* [93, 94], as previously described for *N. defluvii* [26]. Both complexes were highly abundant in *N. marina* proteomes from atmospheric and low O₂ treatments (Data Set 1, Table 1) confirming their important role in central carbon metabolism. The third cluster contains alpha, beta, and gamma subunits of a putative POR with homology to the functionally characterized four-subunit PORs of the anaerobic thermophiles *Pyrococcus* and *Thermotoga* [95] and is absent in all other *Nitrospira* with the exception of *Ca. N. alkalitolerans* [9]. A protein with a 4Fe-4S binding domain was identified in the same operon, potentially representing the missing delta subunit of the POR complex (Data Set 1). This putative four-subunit POR was among the proteins that showed the highest increase in abundance under low O₂ conditions in *N. marina* (Fig. 6). The O₂-tolerant POR/OGOR isoforms were reported to have a >5-times lower specific activity [92] and might therefore constitute a substantial part of the cellular soluble protein content in *H. thermophilus* [96]. Hence, it is tempting to speculate that *N. marina* increases the expression of a more efficient (i.e., higher specific activity), O₂-sensitive four-subunit POR under O₂ limitation. While oxidative stress typically decreases under low O₂ conditions, it might still be high enough to damage O₂-sensitive enzymes. In *N. marina*, the abundance of a periplasmic Cu/Zn-binding SOD and a cytoplasmic catalase (KatG) increased under O₂-limited conditions (Fig. 6), while the abundances of other oxidative stress defense-related proteins remained constant (Data Set 1



and 3). SOD has been shown to be efficient in protecting POR activity from oxidative damage in *Entamoeba histolytica* [97], and was among the proteins with the

highest increase in abundance under low O₂ conditions in *N. marina* (fold-change: 27.9), suggesting a role in POR protection.

◀ **Fig. 6 Heat map of *N. marina* Nb-295^T proteins that were more abundant under low O₂ concentrations compared to atmospheric O₂ concentration.** Relative protein abundance values were square-root transformed and hypothetical proteins were excluded to improve readability. The complete set of untransformed values can be found in Data Set 3. Fold-changes and significance values (adj. *P* value ≤ 0.001, ***, ≤ 0.01, **, ≤ 0.05, *) are shown in white boxes next to the corresponding protein. Select low abundant proteins of interest with high fold-changes were included despite being not statistically significant (see “Material and Methods”). Fold-changes of proteins that were not detected under atmospheric O₂ conditions are omitted to avoid dividing by zero (not available, na). Functional categories of depicted proteins are indicated by different colors. Gene clusters are indicated by black brackets.

Expression of a high O₂-affinity *cbb*₃-type terminal oxidase

The majority of proteins related to electron transport showed similar abundance levels at atmospheric and limiting O₂ concentrations (Data Set 1 and 3). Despite its overall low abundance, a putative high-affinity cytochrome *cbb*₃-type terminal oxidase was 3.6-times more abundant under low O₂ concentrations compared to atmospheric O₂ tension (Fig. 6). The *cbb*₃-type terminal oxidase of *Bradyrhizobium japonicum* was reported to have a *K_m* value of 7 nmol L⁻¹ O₂ [98] and NO₂⁻ oxidation rates have been detected at O₂ concentrations in the low nanomolar range (5–33 nmol L⁻¹ O₂) [25]. This suggests that the *cbb*₃-type terminal oxidase might enable *N. marina* to continue aerobic respiration at low O₂ concentrations, albeit at lower NO₂⁻ oxidation rates (Fig. 5). Low-affinity terminal oxidases are typically more efficient in energy conservation [99], suggesting that *N. marina* benefits from the presence of a terminal oxidase with lower O₂ affinity in well-oxygenated environments. While *N. gracilis* encodes a highly similar high affinity *cbb*₃-type terminal oxidase to *N. marina* [58], this enzyme is lacking in other Nitrospinae, including those identified in OMZs and sediments [76, 82, 100]. Interestingly, these genomes also lack the putative terminal oxidase proposed for *Nitrospira* [26]. Still, in an OMZ where Nitrospinae bacteria were the only detected NOB, NO₂⁻ oxidation rates already approached saturation at ~1 μmol L⁻¹ O₂ [25], indicating that Nitrospinae bacteria might be better adapted to low O₂ concentrations compared to *N. marina*, but the enzyme conferring high O₂ affinity in Nitrospinae remains to be identified.

Increase in abundance of proteins involved in alternative energy metabolism

The abundance of proteins putatively involved in alternative energy metabolisms increased under low O₂ concentrations. These included Group 3b [NiFe] hydrogenase and formate dehydrogenase (Fig. 6). Alternative electron donors such as H₂ and formate are common reaction

products at oxic/anoxic interfaces [101]. *N. moscoviensis* can couple H₂ and formate oxidation to NO₃⁻ reduction to remain active under anoxia [28, 30] (whereas no net growth was observed for the former [30]), however, it is unlikely that H₂ or formate were present at the culture conditions in this study. While abundances of hydrogenase and formate dehydrogenase increased under low O₂ conditions, they were overall still comparably low (Fig. 6), suggesting that their expression might be upregulated when conditions become more unfavorable for aerobic NO₂⁻ oxidation. Curiously, multiple subunits of the Na⁺-translocating N-type ATPase exhibited higher abundances at low O₂ concentration (Fig. 6). Na⁺-translocating ATPases are suggested to be ancient enzymes that were later replaced by energetically more favorable H⁺-translocating ATPases [102]. While only few obligate anaerobes with very tight energy budgets (which cannot cover the losses caused by proton leaks) primarily use Na⁺ energetics, many organisms retained Na⁺ pumps and utilize them under energetically less favorable conditions such as anaerobiosis [102]. Hence, expression of a Na⁺-translocating ATPase suggests an adaptation of *N. marina* to overcome periods of starvation when energetically favorable electron donors or acceptors are short in supply.

Conclusions

Although the vast majority of the ocean is well oxygenated, oxygen-depleted zones exist within the oceanic water column and in marine sediments [83, 84] with consequences for microbial adaptation and evolution. Our results show that, in contrast to Nitrospinae-dominated NOB populations in low oxygen waters [25], NO₂⁻ oxidation activity of *Nitrospira marina* was reduced when grown at ~5.6 μM O₂, suggesting different O₂ adaptations among different marine NOB. We confirm that *N. marina* benefits from the addition of undefined organic carbon substrates, which were shown to be inhibitory for *Nitrospina gracilis* [103], potentially further contributing to ecological niche partitioning within marine NOB. Our results indicate that *N. marina* is highly metabolically versatile, which might enable it to survive under unfavorable conditions with fluctuating levels of electron donors and acceptors. Hence, while Nitrospinae bacteria are the dominant nitrite oxidizers in oligotrophic oceanic regions and OMZs, *Nitrospira* might be better adapted to well-oxygenated high productivity regions including coastal systems, deep-sea trenches and hydrothermal vents. Finally, our results also indicate several ways that NOB may interact with other members of the marine microbial community—through the supply of organic carbon-containing osmolytes and their requirement for exogenous vitamin B₁₂.

Data availability

The genome of *Nitrospira marina* Nb-295^T is available in the JGI IMG/M repository under genome ID number 2596583682. Manually curated protein annotations are available in Data Set 1, targeted quantitative proteomic analyses results are available in Data Set 2, global proteomic spectral counts and differential expression analysis results are available in Data Set 3. Raw mass spectra are available in PRIDE as project number PXD021606. Data is archived at Biological and Chemical Oceanography Data Management office (BCO-DMO) under project 806565.

Acknowledgements We thank John B. Waterbury and Frederica Valois for providing the culture of *Nitrospira marina* Nb-295^T and for continued advice about cultivation. The *N. marina* genome was sequenced as part of US Department of Energy Joint Genome Institute Community Sequencing Project 1337 to CLD, AES, and MAS in collaboration with the user community. We thank Claus Pelikan for bioinformatic assistance. This research was supported by a Simons Foundation Early Career Investigator in Marine Microbiology and Evolution Award (345889) and US National Science Foundation (NSF) award OCE-1924512 to AES. Proteomics analysis was supported by NSF awards OCE-1924554 and OCE-1850719, and NIH award R01GM135709 to MAS. BB was supported by the Austrian Science Fund (FWF) Project Number: J4426-B (“The influence of nitrifiers on the oceanic carbon cycle”), SL by the Netherlands Organization for Scientific Research (NWO) grant 016.Vidi.189.050, and CLD by NSF award OCE-125999.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Kuypers MMM, Marchant HK, Kartal B. The microbial nitrogen-cycling network. *Nat Rev Microbiol.* 2018; 16:263–76.
- Gruber N. The marine nitrogen cycle: Overview and challenges. In: G Capone, DA Bronk, MR Mulholland, EJ Carpenter (eds). Nitrogen in the marine environment, 2nd ed. Academic Press: New York, NY, 2008, pp. 1–50.
- Daims H, Lückner S, Wagner M. A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol.* 2016;24:699–712.
- Pachiadaki MG, Sintez E, Bergauer K, Brown JM, Record NR, Swan BK, et al. Major role of nitrite-oxidizing bacteria in dark ocean carbon fixation. *Science.* 2017;358:1046–51.
- Watson SW, Bock E, Valois FW, Waterbury JB, Schlosser U. *Nitrospira marina* gen. nov. sp. nov.: a chemolithotrophic nitrite-oxidizing bacterium. *Arch Microbiol.* 1986;144:1–7.
- Altmann D, Stief P, Amann R, De Beer D, Schramm A. In situ distribution and activity of nitrifying bacteria in freshwater sediment. *Environ Microbiol.* 2003;5:798–803.
- Freitag TE, Chang L, Clegg CD, Prosser JI. Influence of inorganic nitrogen management regime on the diversity of nitrite-oxidizing bacteria in agricultural grassland soils. *Appl Environ Microbiol.* 2005;71:8323–34.
- Pester M, Maixner F, Berry D, Rattei T, Koch H, Lückner S, et al. *NxrB* encoding the beta subunit of nitrite oxidoreductase as functional and phylogenetic marker for nitrite-oxidizing *Nitrospira*. *Environ Microbiol.* 2014;16:3055–71.
- Daebeler A, Kitzinger K, Koch H, Herbold CW, Steinfeder M, Schwarz J, et al. Exploring the upper pH limits of nitrite oxidation: diversity, ecophysiology, and adaptive traits of haloalkalitolerant *Nitrospira*. *ISME J.* 2020. <https://doi.org/10.1038/s41396-020-0724-1>.
- Lebedeva EV, Alawi M, Fiencke C, Namsaraev B, Bock E, Spieck E. Moderately thermophilic nitrifying bacteria from a hot spring of the Baikal rift zone. *FEMS Microbiol Ecol.* 2005;54: 297–306.
- Gruber-Dorninger C, Pester M, Kitzinger K, Savio DF, Loy A, Rattei T, et al. Functionally relevant diversity of closely related *Nitrospira* in activated sludge. *ISME J.* 2015;9:643–55.
- Spieck E, Hartwig C, McCormack I, Maixner F, Wagner M, Lipski A, et al. Selective enrichment and molecular characterization of a previously uncultured *Nitrospira*-like bacterium from activated sludge. *Environ Microbiol.* 2006;8:405–15.
- Ushiki N, Fujitani H, Aoi Y, Tsuneda S. Isolation of *Nitrospira* belonging to sublineage II from a wastewater treatment plant. *Microbes Environ.* 2013;28:346–53.
- Van Kessel MAHJ, Speth DR, Albertsen M, Nielsen PH, Op Den Camp HJM, Kartal B, et al. Complete nitrification by a single microorganism. *Nature.* 2015;528:555–9.
- Keuter S, Kruse M, Lipski A, Spieck E. Relevance of *Nitrospira* for nitrite oxidation in a marine recirculation aquaculture system and physiological features of a *Nitrospira marina*-like isolate. *Environ Microbiol.* 2011;13:2536–47.
- Nowka B, Off S, Daims H, Spieck E. Improved isolation strategies allowed the phenotypic differentiation of two *Nitrospira* strains from widespread phylogenetic lineages. *FEMS Microbiol Ecol.* 2015;91:fiu031.
- Hongxiang X, Min W, Xiaogu W, Junyi Y, Chunsheng W. Bacterial diversity in deep-sea sediment from northeastern Pacific Ocean. *Acta Ecol Sin.* 2008;28:479–85.
- Baker BJ, Sheik CS, Taylor CA, Jain S, Bhasi A, Cavalcoli JD, et al. Community transcriptomic assembly reveals microbes that contribute to deep-sea carbon and nitrogen cycling. *ISME J.* 2013;7:1962–73.
- Nunoura T, Takaki Y, Hirai M, Shimamura S, Makabe A, Koide O, et al. Hadal biosphere: Insight into the microbial ecosystem in the deepest ocean on Earth. *Proc Natl Acad Sci USA.* 2015;112:E1230–6.
- Hiraoka S, Hirai M, Matsui Y, Makabe A, Minegishi H, Tsuda M, et al. Microbial community and geochemical analyses

- of trans-trench sediments for understanding the roles of hadal environments. *ISME J.* 2020;14:740–56.
21. Lam P, Kuypers MMM. Microbial nitrogen cycling processes in oxygen minimum zones. *Ann Rev Mar Sci.* 2011; 3:317–45.
 22. Codispoti LA, Friederich GE, Packard TT, Glover HE, Kelly PJ, Spinrad RW, et al. High nitrite levels off northern Peru: a signal of instability in the marine denitrification rate. *Science.* 1986;233:1200–2.
 23. Buchwald C, Santoro AE, Stanley RHR, Casciotti KL. Nitrogen cycling in the secondary nitrite maximum of the eastern tropical North Pacific off Costa Rica. *Glob Biogeochem Cycles.* 2015;29: 2061–81.
 24. Lam P, Jensen MM, Kock A, Lettmann KA, Plancherel Y, Lavik G, et al. Origin and fate of the secondary nitrite maximum in the Arabian Sea. *Biogeosciences.* 2011;8:1565–77.
 25. Bristow LA, Dalsgaard T, Tian L, Mills DB, Bertagnolli AD, Wright JJ, et al. Ammonium and nitrite oxidation at nanomolar oxygen concentrations in oxygen minimum zone waters. *Proc Natl Acad Sci USA.* 2016;113:10601–6.
 26. Lückner S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B, et al. A *Nitrospira* metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria. *Proc Natl Acad Sci USA.* 2010;107:13479–84.
 27. Spieck E, Ehrlich S, Aamand J, Bock E. Isolation and immunocytochemical location of the nitrite-oxidizing system in *Nitrospira moscoviensis*. *Arch Microbiol.* 1998;169:225–30.
 28. Koch H, Lückner S, Albertsen M, Kitzinger K, Herbold C, Spieck E, et al. Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *Proc Natl Acad Sci USA.* 2015;112:11371–6.
 29. Ushiki N, Fujitani H, Shimada Y, Morohoshi T, Sekiguchi Y, Tsuneda S. Genomic analysis of two phylogenetically distinct *Nitrospira* species reveals their genomic plasticity and functional diversity. *Front Microbiol.* 2018;8:2637.
 30. Koch H, Galushko A, Albertsen M, Schintlmeister A, Gruber-Dorninger C, Lückner S, et al. Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science.* 2014;345: 1052–4.
 31. Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, et al. Complete nitrification by *Nitrospira* bacteria. *Nature.* 2015;528:504–9.
 32. Strickland JDH, Parsons TR. A practical handbook of seawater analysis, 2nd ed. 1972. *Fish. Res. Bd. Can., Bull. No. 167*
 33. Porter KG, Feig YS. The use of DAPI for identifying aquatic microflora. *Limnol Oceanogr.* 1980;25:943–8.
 34. Wilson K. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol.* 2001;56:2.4.1–5.
 35. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods.* 2013;10:563–9.
 36. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Grechkin Y, et al. IMG: the Integrated Microbial Genomes database and comparative analysis system. *Nucleic Acids Res.* 2012;40:D115–22.
 37. Vallenet D, Calteau A, Dubois M, Amours P, Bazin A, Beuvin M, et al. MicroScope: an integrated platform for the annotation and exploration of microbial gene functions through genomic, pangenomic and metabolic comparative analysis. *Nucleic Acids Res.* 2020;48:D579–89.
 38. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215:403–10.
 39. Saier MH. TCDB: the transporter classification database for membrane transport protein analyses and information. *Nucleic Acids Res.* 2006;34:D181–6.
 40. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, et al. InterProScan: protein domains identifier. *Nucleic Acids Res.* 2005;33:W116–20.
 41. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et al. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat Biotechnol* 2019. <https://doi.org/10.1038/s41587-019-0036-z>.
 42. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics.* 2019;36:1925–7.
 43. Malmstrom R. RNAlater Recipe. protocols.io 2015. <https://doi.org/10.17504/protocols.io.c56y9d>.
 44. Saito MA, McIlvin MR, Moran DM, Santoro AE, Dupont CL, Rafter PA, et al. Abundant nitrite-oxidizing metalloenzymes in the mesopelagic zone of the tropical Pacific Ocean. *Nat Geosci.* 2020;13:355–62.
 45. Simon M, Azam F. Protein content and protein synthesis rates of planktonic marine bacteria. *Mar Ecol Prog Ser.* 1989;51:201–13.
 46. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15:550.
 47. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2013. <http://www.R-project.org>.
 48. Bayer B, Pelikan C, Bittner MJ, Reinthaler T, Könneke M, Herndl GJ, et al. Proteomic response of three marine ammonia-oxidizing archaea to hydrogen peroxide and their metabolic interactions with a heterotrophic alphaproteobacterium. *mSystems.* 2019;4: e00181–19.
 49. Langley SR, Mayr M. Comparative analysis of statistical methods used for detecting differential expression in label-free mass spectrometry proteomics. *J Proteom.* 2015;129:83–92.
 50. Raithel S, Johnson L, Galliard M, Brown S, Shelton J, Herndon N, et al. Inferential considerations for low-count RNA-seq transcripts: a case study on the dominant prairie grass *Andropogon gerardii*. *BMC Genomics.* 2016;17:140.
 51. Kolde R. pheatmap: Pretty Heatmaps. R package version 1.0.8. 2015. <http://CRAN.R-project.org/package=pheatmap>.
 52. Zybailov B, Mosley AL, Sardiou ME, Coleman MK, Florens L, Washburn MP. Statistical analysis of membrane proteome expression changes in *Saccharomyces cerevisiae*. *J Proteome Res.* 2006;5:2339–47.
 53. Ye L, Shao MF, Zhang T, Tong AHY, Lok S. Analysis of the bacterial community in a laboratory-scale nitrification reactor and a wastewater treatment plant by 454-pyrosequencing. *Water Res.* 2011;45:4390–8.
 54. Mündinger AB, Lawson CE, Jetten MSM, Koch H, Lückner S. Cultivation and transcriptional analysis of a canonical *Nitrospira* under stable growth conditions. *Front Microbiol.* 2019;10:1325.
 55. Sakoula D, Nowka B, Spieck E, Daims H, Lückner S. The draft genome sequence of “*Nitrospira lenta*” strain BS10, a nitrite oxidizing bacterium isolated from activated sludge. *Stand Genom Sci.* 2018;13:32.
 56. Chadwick GL, Hemp J, Fischer WW, Orphan VJ. Convergent evolution of unusual complex I homologs with increased proton pumping capacity: energetic and ecological implications. *ISME J.* 2018;12:2668–80.
 57. Cosseau C, Batut J. Genomics of the *ccoNOQP*-encoded *ccb₃* oxidase complex in bacteria. *Arch Microbiol.* 2004;181:89–96.
 58. Lückner S, Nowka B, Rattai T, Spieck E, Daims H. The genome of *Nitrospira gracilis* illuminates the metabolism and evolution of the major marine nitrite oxidizer. *Front Microbiol.* 2013;4:27.
 59. Luoto HH, Belogurov GA, Baykov AA, Lahti R, Malinen AM. Na⁺-translocating membrane pyrophosphatases are widespread in the microbial world and evolutionarily precede H⁺-translocating pyrophosphatases. *J Biol Chem.* 2011;286:21633–42.

60. Luoto HH, Baykov AA, Lahti R, Malinen AM. Membrane-integral pyrophosphatase subfamily capable of translocating both Na⁺ and H⁺. *Proc Natl Acad Sci USA*. 2013;110:1255–60.
61. Hausmann B, Pelikan C, Herbold CW, Köstlbacher S, Albertsen M, Eichorst SA, et al. Peatland *Acidobacteria* with a dissimilatory sulfur metabolism. *ISME J*. 2018;12:1729–42.
62. Füssel J, Lückner S, Yilmaz P, Nowka B, van Kessel MAHJ, Bourceau P, et al. Adaptability as the key to success for the ubiquitous marine nitrite oxidizer *Nitrococcus*. *Sci Adv*. 2017;3:2–10.
63. Ma K, Weiss R, Adams MWW. Characterization of hydrogenase II from the hyperthermophilic archaeon *Pyrococcus furiosus* and assessment of its role in sulfur reduction. *J Bacteriol*. 2000;182:1864–71.
64. Price GD, Woodger FJ, Badger MR, Howitt SM, Tucker L. Identification of a SulP-type bicarbonate transporter in marine cyanobacteria. *Proc Natl Acad Sci USA*. 2004;101:18228–33.
65. Fukaya F, Promden W, Hibino T, Tanaka Y, Nakamura T, Takabe T. An Mrp-like cluster in the halotolerant cyanobacterium *Aphanothece halophytica* functions as a Na⁺/H⁺ antiporter. *Appl Environ Microbiol*. 2009;75:6626–9.
66. Chistoserdova L, Crowther GJ, Vorholt JA, Skovran E, Portais JC, Lidstrom ME. Identification of a fourth formate dehydrogenase in *Methylobacterium extorquens* AM1 and confirmation of the essential role of formate oxidation in methylophony. *J Bacteriol*. 2007;189:9076–81.
67. Lesser MP. Oxidative stress in marine environments: biochemistry and physiological ecology. *Annu Rev Physiol*. 2006;68:253–78.
68. Heelis PF, Kim ST, Okamura T, Sancar A. The photo-repair of pyrimidine dimers by DNA photolyase and model systems. *J Photochem Photobiol B*. 1993;17:219–28.
69. Roberts M. Organic compatible solutes of halotolerant and halophilic microorganisms. *Saline Syst*. 2005;1:5.
70. Csonka L. Prokaryotic osmoregulation: genetics and physiology. *Annu Rev Microbiol*. 1991;45:569–606.
71. Boysen AK, Carlson LT, Durham BP, Groussman RD, Aylward FO, Heal KR, et al. Diel Oscillations of Particulate Metabolites Reflect Synchronized Microbial Activity in the North Pacific Subtropical Gyre. *bioRxiv* 2020. <https://doi.org/10.1101/2020.05.09.086173>.
72. Bayer B, Hansman RL, Bittner MJ, Noriega-Ortega BE, Niggemann J, Dittmar T, et al. Ammonia-oxidizing archaea release a suite of organic compounds potentially fueling prokaryotic heterotrophy in the ocean. *Environ Microbiol*. 2019;21:4062–75.
73. Sañudo-Wilhelmy SA, Gómez-Consarnau L, Suffridge C, Webb EA. The role of B vitamins in marine biogeochemistry. *Ann Rev Mar Sci*. 2014;6:339–67.
74. Graham RM, Deery E, Warren MJ. Vitamin B₁₂: Biosynthesis of the Corrin Ring. *Tetrapyrroles: Birth, Life and Death*. New York, NY: Springer; 2009. pp. 286–99.
75. Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, et al. Uneven distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative genomics. *ISME J*. 2019;13:789–804.
76. Mueller AJ, Jung M-Y, Strachan CR, Herbold CW, Kirkegaard RH, Wagner M, et al. Genomic and kinetic analysis of novel Nitrospinae enriched by cell sorting. *ISME J*. 2020. <https://doi.org/10.1038/s41396-020-00809-6>.
77. Park S-J, Andrei A-Ş, Bulzu P-A, Kavagutti VS, Ghai R, Mosier AC. Expanded diversity and metabolic versatility of marine nitrite-oxidizing bacteria revealed by cultivation- and genomics-based approaches. *Appl Environ Microbiol* 2020. <https://doi.org/10.1128/AEM.016667-20>.
78. Hawco NJ, McIlvin MM, Bundy RM, Tagliabue A, Goepfert TJ, Moran DM, et al. Minimal cobalt metabolism in the marine cyanobacterium *Prochlorococcus*. *Proc Natl Acad Sci USA*. 2020;12. <https://doi.org/10.1073/pnas.2001393117>.
79. Saito MA, Rocap G, Moffett JW. Production of cobalt binding ligands in a *Synechococcus* feature at the Costa Rica upwelling dome. *Limnol Oceanogr*. 2005;50:279–90.
80. Noble AE, Ohnemus DC, Hawco NJ, Lam PJ, Saito MA. Coastal sources, sinks and strong organic complexation of dissolved cobalt within the US North Atlantic GEOTRACES transect GA03. *Biogeosciences*. 2017;14:2715–39.
81. Füssel J, Lam P, Lavik G, Jensen MM, Holtappels M, Günter M, et al. Nitrite oxidation in the Namibian oxygen minimum zone. *ISME J*. 2012;6:1200–9.
82. Sun X, Kop LFM, Lau MCY, Frank J, Jayakumar A, Lückner S, et al. Uncultured *Nitrospina*-like species are major nitrite oxidizing bacteria in oxygen minimum zones. *ISME J*. 2019;13:2391–402.
83. D'Hondt S, Inagaki F, Zarikian CA, Abrams LJ, Dubois N, Engelhardt T, et al. Presence of oxygen and aerobic communities from sea floor to basement in deep-sea sediments. *Nat Geosci*. 2015;8:299–304.
84. Karstensen J, Stramma L, Visbeck M. Oxygen minimum zones in the eastern tropical Atlantic and Pacific oceans. *Prog Oceanogr*. 2008;77:331–50.
85. Vollmer AC, Bark SJ. Twenty-five years of investigating the universal stress protein: function, structure, and applications. *advances in applied microbiology*, 1st ed. 2018. Elsevier Inc., pp 1–36.
86. Nyström T, Neidhardt FC. Expression and role of the universal stress protein, UspA, of *Escherichia coli* during growth arrest. *Mol Microbiol*. 1994;11:537–44.
87. O'Toole R, Smeulders MJ, Blokpoel MC, Kay EJ, Loughheed K, Williams HD. A two-component regulator of universal stress protein expression and adaptation to oxygen starvation in *Mycobacterium smegmatis*. *J Bacteriol*. 2003;185:1543–54.
88. Coulthurst S. The Type VI secretion system: a versatile bacterial weapon. *Microbiology*. 2019;165:503–15.
89. Gallique M, Bouteiller M, Merieau A. The type VI secretion system: a dynamic system for bacterial communication? *Front Microbiol*. 2017;8:1454.
90. Mellbye BL, Spieck E, Bottomley PJ, Sayavedra-Soto LA. Acyl-homoserine lactone production in nitrifying bacteria of the genera *Nitrosospira*, *Nitrobacter*, and *Nitrospira* identified via a survey of putative quorum-sensing genes. *Appl Environ Microbiol*. 2017;83:e01540–17.
91. Erb TJ. Carboxylases in natural and synthetic microbial pathways. *Appl Environ Microbiol*. 2011;77:8466–77.
92. Yamamoto M, Arai H, Ishii M, Igarashi Y. Role of two 2-oxoglutarate:ferredoxin oxidoreductases in *Hydrogenobacter thermophilus* under aerobic and anaerobic conditions. *FEMS Microbiol Lett*. 2006;263:189–93.
93. Ikeda T, Ochiai T, Morita S, Nishiyama A, Yamada E, Arai H, et al. Anabolic five subunit-type pyruvate:ferredoxin oxidoreductase from *Hydrogenobacter thermophilus* TK-6. *Biochem Biophys Res Commun*. 2006;340:76–82.
94. Yun NR, Yamamoto M, Arai H, Ishii M, Igarashi Y. A novel five-subunit-type 2-oxoglutarate:ferredoxin oxidoreductases from *Hydrogenobacter thermophilus* TK-6. *Biochem Biophys Res Commun*. 2002;292:280–6.
95. Kletzin A, Adams MWW. Molecular and phylogenetic characterization of pyruvate and 2-ketoisovalerate ferredoxin oxidoreductases from *Pyrococcus furiosus* and pyruvate ferredoxin oxidoreductase from *Thermotoga maritima*. *J Bacteriol*. 1996;178:248–57.
96. Berg IA. Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl Environ Microbiol*. 2011;77:1925–36.
97. Pineda E, Encalada R, Rodríguez-Zavala JS, Olivos-García A, Moreno-Sánchez R, Saavedra E. Pyruvate:ferredoxin

- oxidoreductase and bifunctional aldehyde-alcohol dehydrogenase are essential for energy metabolism under oxidative stress in *Entamoeba histolytica*. FEBS J. 2010;277:3382–95.
98. Preisig O, Zufferey R, Thöny-Meyer L, Appleby CA, Hennecke H. A high-affinity *cbb₃*-type cytochrome oxidase terminates the symbiosis-specific respiratory chain of *Bradyrhizobium japonicum*. J Bacteriol. 1996;178:1532–8.
99. Han H, Hemp J, Pace LA, Ouyang H, Ganesan K, Roh JH, et al. Adaptation of aerobic respiration to low O₂ environments. Proc Natl Acad Sci USA. 2011;108:14109–14.
100. Ngugi DK, Blom J, Stepanauskas R, Stingl U. Diversification and niche adaptations of *Nitrospina*-like bacteria in the polyextreme interfaces of Red Sea brines. ISME J. 2016;10:1383–99.
101. Shao MF, Zhang T, Fang HHP. Sulfur-driven autotrophic denitrification: diversity, biochemistry, and engineering applications. Appl Microbiol Biotechnol. 2010;88:1027–42.
102. Mulkidjanian AY, Dibrov P, Galperin MY. The past and present of the sodium energetics: May the sodium-motive force be with you. Biochim Biophys Acta. 2008;1777:985–92.
103. Watson SW, Waterbury JB. Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. Arch Mikrobiol. 1971;77:203–30.