

Published in final edited form as:

Angew Chem Int Ed Engl. 2007 ; 46(27): 5195–5197. doi:10.1002/anie.200701128.

Metal-Catalyzed [1,2]-Alkyl Shift in Allenyl Ketones: Synthesis of Multisubstituted Furans**


Alexander S. Dudnik and Vladimir Gevorgyan

 Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Room 4500, Chicago, IL 60607 (USA), Fax: (+1)312-355-0836, vlad@uic.edu Homepage: <http://www.chem.uic.edu/vggroup>

Keywords

allenyl ketones; furans; gold; Lewis acids; rearrangement

Cycloisomerization of allenyl ketones is an efficient approach for the assembly of the furan ring, an important heterocyclic unit.^[1] This transformation in the presence of transition-metal catalysts was first reported by Marshall et al.^[2] and later by Hashmi et al.^[3] for the synthesis of furans [G = H, Eq. (1)]. Recently, we have developed a set of transition-metal-catalyzed cascade transformations of allenyl ketones involving 1,2-migration of various groups (G = SR,^[4] Hal,^[5] OP(O)(OR)₂, OC(O)R, OSO₂R^[6]) to produce up to tetrasubstituted furans [Eq. (1)]. Herein, we wish to report a novel metal-catalyzed [1,2]-alkyl shift in allenyl ketones as a key step in the formation of up to fully carbon-substituted furans [Eq. (1)].



Marshall (G = H, cat. = Ag)

Hashmi (G = H, cat. = Pd, Au)

 Gevorgyan (G = SR, Hal, OP(O)(OR)₂, OC(O)R, OSO₂R',


cat. = Cu, Ag, Au)

this work (G = Alk, Ar)

(1)

Recently, we reported the Au-catalyzed regiodivergent synthesis of halofurans.^[5] It was found that in the presence of Au^I catalysts clean hydrogen migration from **1** occurs to form **2** [Eq. (2)]. The absence of H/D-scrambling, in contrast to that observed in the Cu/base-assisted synthesis of pyrroles,^[7] supported the clean [1,2]-hydrogen shift to the carbenoid center in intermediate **i**.^[5]

**The support of the National Institutes of Health (GM-64444) is gratefully acknowledged.




(2)

It occurred to us that 1,2-migration of an alkyl/aryl group by this mechanism is also feasible,^[8–11] which may allow for the assembly of fully carbon-substituted furans. To this end, we have tested the possible cycloisomerization of allene **3** to give furan **4** in the presence of different catalysts (Table 1). We have found that employment of Au^I and Au^{III} halides gave low yields of furan **4** (Table 1, entries 1 and 2). Gratifyingly, switching to cationic Au^I complexes led to formation of **4k** in nearly quantitative yield (Table 1, entries 3 and 4). In analogy to gold halides, Pt^{II}, Pt^{IV}, and Pd^{II} salts were inefficient in this reaction (Table 1, entries 5–7). Use of Cu^I halides resulted in no reaction (Table 1, entry 8), while employment of cationic Ag^I, Cu^I, and Cu^{II} salts produced **4** in moderate to high yields (Table 1, entries 9–13). Encouraged by these results, we also tested main-group metals in this reaction. Surprisingly, Al, Si, Sn, and In triflates provided moderate to excellent yields of desired furan **4** (Table 1, entries 14, 16–19). Although [Au(PPh₃)]OTf, AgOTf, In(OTf)₃, Sn(OTf)₂, and TIPSOTf were nearly equally efficient in the cascade cycloisomerization of **3** to give **4**, In(OTf)₃ appeared to be a more general catalyst with respect to the substrate scope.^[12]

Next, cycloisomerization of differently substituted allenyl ketones **3a–m** was examined under the optimized conditions (Table 2). Thus, cycloisomerization of 4,4-diphenyl-substituted allenyl ketones **3b–d** proceeded smoothly to provide good to high yields of furans **4b–d** (Table 2, entries 2–4). Selective migration of the phenyl over the methyl group occurred in allenyl ketone **3e** to give **4e** in 72% yield (Table 2, entry 5). Not surprisingly, cycloisomerization of allenyl ketone **3i**, possessing two methyl groups, provided the corresponding furan **4i** in low yield only (Table 2, entry 8). In contrast to the disfavored methyl-group migration in Table 2, entry 5, migration of the ethyl group competed with the phenyl group in **3f**, which resulted in formation of a 2.3:1 mixture of regioisomeric furans **4f** and **4g**, respectively (Table 2, entry 6). Cyclopentylidene allenyl ketone **3h** underwent smooth cyclization with ring expansion^[13] to give fused furan **4h** in 75% yield (Table 2, entry 7). It was also demonstrated that a variety of functional groups such as methoxy (Table 2, entry 9), bromo (Table 2, entry 10), nitro (Table 2, entry 11), and cyano (Table 2, entry 12) were perfectly tolerated under these reaction conditions.

In addition, we have shown that trisubstituted furan **4b** can be obtained directly from alkynyl ketone **5b** [Eq. (3)]. However, the yield for this one-pot transformation was somewhat lower than that for cycloisomerization of allene **3b** (Table 2, entry 2).



(3)

We propose the following mechanism for the cascade transformation of allenyl ketone **3** into furan **4** (Scheme 1). Cycloisomerization in the presence of oxophilic Lewis acids, such as In, Sn, and Si triflates, follows path A, according to which, the Lewis acid activates the enone moiety (see **6**) to form vinyl cation **7**.^[14] [1,2]-Alkyl shift in **7** produces the regioisomeric vinyl cation **8**,^[15] which, upon cyclization, transforms into furan **4** and regenerates the Lewis acid catalyst. Alternatively, π -philic catalysts, such as Ag^I, Cu^I, and Au^I salts, activate the carbon-carbon double bond of allene (see **9**) and trigger nucleophilic attack of a carbonyl oxygen lone pair at the terminal carbon of the allene moiety to form cyclic oxonium intermediate **10**.^[2c,5] [1,5]-Alkyl shift^[16] (Scheme 1, path B) to form **11** with subsequent elimination of metal gives **4**. The involvement of an electrophilic mechanism (Scheme 1, paths A and B) is supported by the data presented in Table 2. Thus, the migratory aptitude of a phenyl vs. that of a methyl group (> 100:1) is in good agreement with that reported in the literature for rearrangements of cations.^[17] Although a mechanism involving [1,2]-alkyl shift in the carbenoid intermediate **12**^[5,8] (Scheme 1, path C) cannot be completely ruled out at this point, it is considered to be less likely.^[18,19]

In summary, we have developed a novel metal-catalyzed method for the synthesis of furans, which proceeds by an unprecedented [1,2]-alkyl shift in allenyl ketones. This method allows for efficient synthesis of up to fully carbon-substituted and fused furans.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. For a recent review, see: Kirsch SF. *Org Biomol Chem*. 2006; 4:2076. [PubMed: 16729118]
2. a) Marshall JA, Robinson ED. *J Org Chem*. 1990; 55:3450. b) Marshall JA, Wang XJ. *J Org Chem*. 1991; 56:960. c) Marshall JA, Bartley GS. *J Org Chem*. 1994; 59:7169. d) Marshall JA, Sehon CA. *J Org Chem*. 1995; 60:5966. e) Marshall JA, Wallace EM. *J Org Chem*. 1995; 60:796.
3. a) Hashmi ASK. *Angew Chem*. 1995; 107:1749. *Angew Chem Int Ed Engl*. 1995; 34:1581. b) Hashmi ASK, Schwarz L, Choi JH, Frost TM. *Angew Chem*. 2000; 112:2382. *Angew Chem Int Ed*. 2000; 39:2285.
4. Kim JT, Kel'in AV, Gevorgyan V. *Angew Chem*. 2003; 115:102. *Angew Chem Int Ed*. 2003; 42:98.
5. Sromek AW, Rubina M, Gevorgyan V. *J Am Chem Soc*. 2005; 127:10500. [PubMed: 16045332]
6. Sromek AW, Kel'in AV, Gevorgyan V. *Angew Chem*. 2004; 116:2330. *Angew Chem Int Ed*. 2004; 43:2280.
7. Kel'in AV, Sromek AW, Gevorgyan V. *J Am Chem Soc*. 2001; 123:2074. [PubMed: 11456838]
8. For examples of [1,2]-shifts in carbenoids, see: a) Xiao F, Wang J. *J Org Chem*. 2006; 71:5789. and references therein. [PubMed: 16839167] b) Markham JP, Staben ST, Toste FD. *J Am Chem Soc*. 2005; 127:9708. [PubMed: 15998074] c) Gorin DJ, Davis NR, Toste FD. *J Am Chem Soc*. 2005; 127:11260. [PubMed: 16089452]
9. For general reviews, see: One or more CH and/or CC bond(s) formed by rearrangement: a) Ducrot, PH. *Comprehensive Organic Functional Group Transformations II*. Katritzky, AR.; Taylor, RJK., editors. Vol. 1. Elsevier; Oxford, UK: 2005. p. 375-426. b) Pattenden, G. *Carbon-Carbon σ -Bond Formation: Rearrangement Reactions*. In: Trost, BM.; Fleming, I., editors. *Comprehensive Organic Synthesis: Selectivity, Strategy, and Efficiency in Modern Organic Chemistry*. Vol. 3. Pergamon; New York: 1991. p. 705-1043.
10. While this manuscript was in preparation, two independent works on synthesis of carbocycles involving [1,2]-alkyl shifts to carbenoid center in allenes were reported. See: a) Funami H, Kusama H, Iwasawa N. *Angew Chem Int Ed*. 2007; 46:909. b) Lee JH, Toste FD. *Angew Chem Int Ed*. 2007; 46:912.

11. For a synthesis of dehydrofuranones by [1,2]-alkyl shift, analogous to a formal ketol rearrangement, see: Kirsch SF, Binder JT, Liébert C, Menz H. *Angew Chem.* 2006; 118:6010. *Angew Chem Int Ed.* 2006; 45:5878. and references therein.
12. See the Supporting Information for details.
13. For examples of ring expansions in the synthesis of carbocycles proceeding by a carbenoid mechanism, see Ref. [9].
14. For activation of enone moiety by Lewis acids see, for example: a) Childs RF, Mulholland DL, Nixon A. *Can J Chem.* 1982; 60:801. b) Schwier T, Gevorgyan V. *Org Lett.* 2005; 7:5191. [PubMed: 16268535]
15. For examples of [1,2]-shifts in vinyl cations, see: a) Capozzi G, Lucchini V, Marcuzzi F, Melloni G. *Tetrahedron Lett.* 1976; 17:717. b) Jäckel KP, Hanack M. *Tetrahedron Lett.* 1974; 15:1637.
16. a) Miller B. *J Am Chem Soc.* 1970; 92:432. b) Dolbier WR, Anapolle KE, McCullagh L, Matsui K, Riemann JM, Rolison D. *J Org Chem.* 1979; 44:2845. c) Ode M, Breslow R. *Tetrahedron Lett.* 1973; 14:2537.
17. Saunders WH, Paine RH. *J Am Chem Soc.* 1961; 83:882.
18. The observed migratory aptitude trends (Ph vs. Et, and Ph vs. Me) do not correspond to those reported in literature for [1,2]-alkyl migration to a carbenoid center. See, for example: a) Philip H, Keating J. *Tetrahedron Lett.* 1961; 2:523. b) von der Schulenburg, W Graf; Hopf, H.; Walsh, R. *Angew Chem.* 1999; 111:1200. *Angew Chem Int Ed.* 1999; 38:1128.
19. No cyclopropanation product was observed in the cycloisomerization of dimethylallenyl ketone **3i**; however, this transformation was reported as a major process in the cycloisomerization of a carbocyclic analogue of **12**. See Ref. [9a].

Table 1

Optimization of reaction conditions.^a

Entry	Catalyst ^b	mol%	Solvent	T [°C]	Yield [%] ^c
1	AuBr ₃	5	toluene ^d	100	23
2	AuI	5	toluene ^d	100	traces
3	[Au(PPh ₃)OTf]	1	toluene ^d	100	100 (89)
4	[Au(PPh ₃)OTf]	5	CH ₂ Cl ₂ ^e	RT	99
5	PtCl ₂	5	toluene ^f	100	21
6	PtCl ₄	5	toluene ^f	100	21
7	[PdCl ₂ (PhCN) ₂]	5	toluene ^f	100	35
8	CuX (X = Cl, Br, I)	5	toluene ^f	100	0
9	CuOTf·PhH	5	toluene ^f	100	42
10	Cu(OTf) ₂	5	toluene ^g	100	95
11	AgPF ₆	5	toluene ^g	100	47
12	AgOTf	5	toluene ^g	100	(80)
13	AgOTf	20	CH ₂ Cl ₂ ^e	RT	70 (62)
14	Al(OTf) ₃	5	toluene ^g	100	64
15	Zn(OTf) ₂	5	toluene ^g	100	39
16	TMSOTf	20	CH ₂ Cl ₂ ^e	RT	82 (62)
17	In(OTf) ₃	5	toluene ^g	100	91 (81)
18	Sn(OTf) ₂	5	toluene ^g	100	97 (81)
19	TIPSOTf	5	toluene ^g	100	100 (81)
20	TMSNTf ₂	5	toluene ^g	100	72

^aEntries 1–4: Ar = *p*-Br-C₆H₄; entries 5–20: Ar = Ph.

^bTf = trifluoromethanesulfonyl, TIPS = trisopropylsilyl, TMS = trimethylsilyl.

^cYield determined from NMR spectrum; yield of isolated product in parentheses.

^d0.05 M, solution of **3**.

^e0.02 M solution of **3**.

^f1 M solution of **3**.

^g0.1 M solution of **3**.

Table 2

Lewis acid catalyzed synthesis of furans.



Entry	Allenyl ketone	Furan	Yield [%] ^d
1			81 ^b
2			64 ^c
3			90
4			79 ^d
5			72 (52) ^{e,f}
6			88 ^g
			(76) ^{h,i}
7			75

Entry	Allenyl ketone	Furan	Yield [%] ^a
8			10 ^f
9			62
10			93 (89) ^h
11			85 ^b
12			94 ^b

^aYield of isolated product; 0.25–0.8-mmol scale, In(OTf)₃ was used unless otherwise mentioned.

^b 5 mol% Sn(OTf)₂ was used.

^c 10 mol% In(OTf)₃ was used.

^d 20 mol% AgOTf/*p*-xylene, 140°C, 1 h.

^e 2 mol% [Au(PPh₃)OTf] was used.

^f Yield determined from NMR spectrum.

^g 2:3:1 mixture of **4 f**:**4g** by ¹H NMR spectroscopy.

^h 1 mol% [Au(PPh₃)OTf] was used.

ⁱ 2:2:1 mixture of **4 f**:**4g** by ¹H NMR spectroscopy.