

Toposym 3

David Preiss

Metric spaces in which Prohorov's theorem is not valid

In: Josef Novák (ed.): General Topology and its Relations to Modern Analysis and Algebra, Proceedings of the Third Prague Topological Symposium, 1971. Academia Publishing House of the Czechoslovak Academy of Sciences, Praha, 1972. pp. 359.

Persistent URL: <http://dml.cz/dmlcz/700750>

Terms of use:

© Institute of Mathematics AS CR, 1972

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

METRIC SPACES IN WHICH PROHOROV'S THEOREM IS NOT VALID

D. PREISS

Praha

A well-known Prohorov's theorem says that for every topologically complete metric space X , for every compact set M of measures on X with mass 1 (with the weak topology) and for every $\varepsilon > 0$ there exists a compact set $A \subset X$ such that $\mu(X - A) < \varepsilon$ for each $\mu \in M$. It has been a problem whether this theorem holds in every separable metric space which is a Borel subset of its completion. This problem can be solved by the help of the following theorem.

Theorem 1. *A coanalytic separable metric space is topologically complete if and only if it contains no countable dense-in-itself G_δ subspace.*

If Prohorov's theorem holds in some metric space X then it is easy to prove that it holds also in every G_δ subspace of X . With respect to Theorem 1 and to Sierpiński's result according to which every two countable dense-in-itself metric spaces are homeomorphic it is clear that the negative answer for some countable dense-in-itself metric space implies the negative answer for every separable coanalytic metric space which is not topologically complete. Such countable metric space can be constructed (namely the space of rational numbers), therefore the following theorem holds.

Theorem 2. *If X is a separable coanalytic metric space then Prohorov's theorem holds in X if and only if X is topologically complete.*

CHARLES UNIVERSITY, PRAHA