

## METRICALLY COMPLETE REGULAR RINGS

BY

K. R. GOODEARL<sup>1</sup>

**ABSTRACT.** This paper is concerned with the structure of those (von Neumann) regular rings  $R$  which are complete with respect to the weakest metric derived from the pseudo-rank functions on  $R$ , known as the  $N^*$ -metric. It is proved that this class of regular rings includes all regular rings with bounded index of nilpotence, and all  $\aleph_0$ -continuous regular rings. The major tool of the investigation is the partially ordered Grothendieck group  $K_0(R)$ , which is proved to be an archimedean norm-complete interpolation group. Such a group has a precise representation as affine continuous functions on a Choquet simplex, from earlier work of the author and D. E. Handelman, and additional aspects of its structure are derived here. These results are then translated into ring-theoretic results about the structure of  $R$ . For instance, it is proved that the simple homomorphic images of  $R$  are right and left self-injective rings, and  $R$  is a subdirect product of these simple self-injective rings. Also, the isomorphism classes of the finitely generated projective  $R$ -modules are determined by the isomorphism classes modulo the maximal two-sided ideals of  $R$ . As another example of the results derived, it is proved that if all simple artinian homomorphic images of  $R$  are  $n \times n$  matrix rings (for some fixed positive integer  $n$ ), then  $R$  is an  $n \times n$  matrix ring.

All rings in this paper are associative with 1, and all modules are unital right modules. For the overall theory of regular rings, we refer the reader to [2]; for the general development of  $K_0$  of regular rings as partially ordered abelian groups, and the theory of partially ordered abelian groups via their state spaces, we refer the reader to [2, 4]. In particular, these references should be consulted for more detail on definitions and concepts which are just sketched here.

**I.  $N^*$ -completeness.** Completeness of a regular ring with respect to a rank function, or with respect to a family of pseudo-rank functions, implies that the ring is right and left self-injective [2, Theorems 19.7 and 20.8], hence a considerable amount of structure theory is available for such rings [2, Chapters 9–12]. The purpose of this paper is to investigate a much broader class of regular rings, namely those which are complete with respect to the (pseudo-) metric obtained from the supremum  $N^*$  of all pseudo-rank functions on the ring. In particular, all regular rings complete with respect to a family of pseudo-rank functions are  $N^*$ -complete, but also, as we prove later in this section, all regular rings with bounded index of nilpotence and all  $\aleph_0$ -continuous regular rings are  $N^*$ -complete.

---

Received by the editors January 13, 1981 and, in revised form, April 3, 1981.  
1980 *Mathematics Subject Classification.* Primary 16A30, 16A54, 06F20.

*Key words and phrases.* Regular ring,  $N^*$ -complete,  $K_0$ , interpolation group.

<sup>1</sup> This research was partially supported by a grant from the National Science Foundation.

©1982 American Mathematical Society  
0002-9947/82/0000-1066/\$08.75

In addition to these results, the present section introduces the definition and a few basic properties of  $N^*$ -completeness. §II is devoted to proving that  $K_0$  of any  $N^*$ -complete regular ring is an archimedean norm-complete interpolation group, while the third section develops a number of structural properties of such groups. In §IV, we apply these results to the structure theory of  $N^*$ -complete regular rings.

DEFINITION. Recall that a *pseudo-rank function* on a regular ring  $R$  is a map  $P: R \rightarrow [0, 1]$  such that (a)  $P(1) = 1$ ; (b)  $P(xy) \leq P(x), P(y)$  for all  $x, y \in R$ ; (c)  $P(e + f) = P(e) + P(f)$  for all orthogonal idempotents  $e, f \in R$ . We use  $\mathbf{P}(R)$  to denote the set of all pseudo-rank functions on  $R$ . Considered as a subset of the linear topological space  $\mathbf{R}^R$  (which is given the product topology),  $\mathbf{P}(R)$  is a compact convex set [2, Proposition 16.17]. In fact,  $\mathbf{P}(R)$  is a Choquet simplex [2, Theorem 17.5].

DEFINITION. Let  $R$  be a regular ring. For each  $x \in R$ , we define

$$N^*(x) = \sup\{P(x) \mid P \in \mathbf{P}(R)\},$$

with the proviso that  $N^*(x) = 0$  in case  $\mathbf{P}(R)$  is empty. (This definition is formally different from the definition of  $N^*$  in [2, p. 272], but the two definitions are equivalent, as follows from [2, Proposition 18.10].) Thus  $N^*(x)$  is a real number, and  $0 \leq N^*(x) \leq 1$ . Whenever the ring  $R$  needs to be emphasized, we shall write  $N_R^*(x)$  in place of  $N^*(x)$ . In case  $R$  is unit-regular,  $N^*$  may be computed as in the following proposition. We first recall two pieces of notation.

DEFINITION. Given modules  $A$  and  $B$ , we write  $A \lesssim B$  to mean that  $A$  is isomorphic to a submodule of  $B$ . Given a module  $A$  and a positive integer  $n$ , we write  $nA$  to denote the direct sum of  $n$  copies of  $A$ .

PROPOSITION 1.1. *If  $R$  is a nonzero unit-regular ring, then*

$$N^*(x) = \inf\{k/n \mid k, n \in \mathbf{N} \text{ and } n(xR) \lesssim kR_R\}$$

for all  $x \in R$ .

PROOF. [2, Proposition 18.10].  $\square$

LEMMA 1.2. *Let  $R$  be a regular ring, and let  $x, y, y_1, \dots, y_n \in R$ .*

(a) *If  $t(xR) \lesssim s_1(y_1R) \oplus \dots \oplus s_n(y_nR)$  for some positive integers  $t, s_1, \dots, s_n$ , then*

$$N^*(x) \leq (s_1/t)N^*(y_1) + \dots + (s_n/t)N^*(y_n).$$

(b) *If  $t(xR) \cong s(yR)$  for some positive integers  $s, t$ , then  $N^*(x) = (s/t)N^*(y)$ .*

(c)  *$N^*(xy) \leq N^*(x)$  and  $N^*(xy) \leq N^*(y)$ .*

(d)  *$N^*(x + y) \leq N^*(x) + N^*(y)$ .*

PROOF. These results are clear if  $\mathbf{P}(R)$  is empty, so assume it is nonempty.

(a) For any  $P \in \mathbf{P}(R)$ , we have

$$tP(x) \leq s_1P(y_1) + \dots + s_nP(y_n)$$

by [2, Proposition 16.1], whence

$$\begin{aligned} P(x) &\leq (s_1/t)P(y_1) + \dots + (s_n/t)P(y_n) \\ &\leq (s_1/t)N^*(y_1) + \dots + (s_n/t)N^*(y_n). \end{aligned}$$

Consequently,  $N^*(x) \leq (s_1/t)N^*(y_1) + \dots + (s_n/t)N^*(y_n)$ .

(b) This follows directly from (a).

(c) For all  $P \in \mathbf{P}(R)$ , we have  $P(xy) \leq P(x) \leq N^*(x)$ , hence  $N^*(xy) \leq N^*(x)$ . Likewise,  $N^*(xy) \leq N^*(y)$ .

(d) Since  $(x + y)R \leq xR + yR \lesssim xR \oplus yR$ , we may apply (a).  $\square$

DEFINITION. Let  $R$  be a regular ring. In view of Lemma 1.2, we see that the rule  $\delta(x, y) = N^*(x - y)$  defines a pseudo-metric  $\delta$  on  $R$ . By way of abbreviation, we shall refer to  $\delta$  as *the  $N^*$ -metric on  $R$* , even though  $\delta$  is not always a metric. Note that  $\delta$  is a metric if and only if  $\ker(\mathbf{P}(R)) = 0$ . For all  $x, y, z, w \in R$ , we see that

$$N^*((x + y) - (z + w)) = N^*((x - z) + (y - w)) \leq N^*(x - z) + N^*(y - w),$$

$$N^*(xy - zw) = N^*((x - z)y + z(y - w)) \leq N^*(x - z) + N^*(y - w).$$

Thus addition and multiplication in  $R$  are uniformly continuous with respect to  $N^*$ . For all  $x, y \in R$ , we also have

$$N^*(x) \leq N^*(x - y) + N^*(y),$$

$$N^*(y) \leq N^*(y - x) + N^*(x) = N^*(x - y) + N^*(y),$$

whence  $|N^*(x) - N^*(y)| \leq N^*(x - y)$ . Thus the map  $N^*: R \rightarrow [0, 1]$  is uniformly continuous with respect to the  $N^*$ -metric.

DEFINITION. A regular ring  $R$  is called  *$N^*$ -complete* provided  $\ker(\mathbf{P}(R)) = 0$  (so that the  $N^*$ -metric on  $R$  is actually a metric) and  $R$  is complete in the  $N^*$ -metric. For example, if  $R$  is a simple artinian ring, then there is a unique rank function  $P$  on  $R$ , which takes on only the values  $0, 1/k, 2/k, \dots, 1$ , for some  $k \in \mathbf{N}$  [2, Corollary 16.6]. As a result,  $N^* = P$ , and  $N^*(x) \geq 1/k$  for all nonzero  $x \in R$ , so that the  $N^*$ -metric on  $R$  is discrete. Therefore  $R$  is  $N^*$ -complete. More generally, we have the following result.

THEOREM 1.3. *A regular ring  $R$  has bounded index of nilpotence if and only if  $\ker(\mathbf{P}(R)) = 0$  and the  $N^*$ -metric on  $R$  is discrete, in which case  $R$  is  $N^*$ -complete.*

PROOF. First assume that  $\ker(\mathbf{P}(R)) = 0$  and the  $N^*$ -metric on  $R$  is discrete. Then there exists  $n \in \mathbf{N}$  such that  $N^*(x) \geq 1/n$  for any nonzero  $x \in R$ , and we claim that the index of nilpotence of  $R$  is at most  $n$ . If not,  $R$  must contain a direct sum of  $n + 1$  nonzero pairwise isomorphic right ideals [2, Theorem 7.2]. Consequently, there is a nonzero element  $y \in R$  satisfying  $(n + 1)(yR) \lesssim R_R$ . But then  $N^*(y) > 0$  (because  $\ker(\mathbf{P}(R)) = 0$ ) and  $N^*(y) \leq 1/(n + 1)$  (by Lemma 1.2), contradicting our discreteness assumption. Therefore the index of nilpotence of  $R$  is at most  $n$ , as claimed.

Conversely, assume that  $R$  has bounded index of nilpotence. We first note that any maximal two-sided ideal  $M$  of  $R$  is the kernel of a pseudo-rank function on  $R$ . Namely,  $R/M$  is a simple artinian ring [2, Theorem 7.9], so there exists a unique rank function on  $R/M$ , which pulls back to a pseudo-rank function on  $R$  with kernel  $M$ .

As all primitive factor rings of  $R$  are artinian [2, Corollary 7.10], the intersection of the maximal two-sided ideals of  $R$  is zero. Since each maximal two-sided ideal is the kernel of a pseudo-rank function, we obtain  $\ker(\mathbf{P}(R)) = 0$ .

Let  $n$  be the index of nilpotence of  $R$ . We claim that  $N^*(x) \geq 1/n$  for any nonzero element  $x \in R$ .

Choose a maximal two-sided ideal  $M$  of  $R$  such that  $x \notin M$ . By [2, Theorem 7.9],  $R/M \cong M_k(D)$  for some positive integer  $k \leq n$  and some division ring  $D$ . Then  $R/M$  has a unique rank function  $Q$ , which takes on only the values  $0, 1/k, 2/k, \dots, 1$ , and  $Q(x + M) \geq 1/k \geq 1/n$ . Pulling  $Q$  back to a pseudo-rank function  $P$  on  $R$ , we obtain  $P(x) \geq 1/n$ , and so  $N^*(x) \geq 1/n$ , as claimed.

Therefore the  $N^*$ -metric on  $R$  is discrete, and, consequently,  $R$  is  $N^*$ -complete.

□

For a deeper class of examples, we now proceed to prove that every  $\aleph_0$ -continuous regular ring is  $N^*$ -complete. In particular, it will follow that every regular, right and left self-injective ring is  $N^*$ -complete. Recall that a regular ring  $R$  is defined to be  $\aleph_0$ -continuous provided the lattice of principal right ideals of  $R$  is an  $\aleph_0$ -continuous geometry. Equivalently,  $R$  is  $\aleph_0$ -continuous if and only if every countably generated right (left) ideal of  $R$  is essential in a principal right (left) ideal [2, Corollary 14.4].

We begin with a lemma generalizing [2, Corollary 14.27(a)] to finitely generated projective modules. This lemma is also implicit in [8, Proposition 2.1].

**LEMMA 1.4.** *Let  $R$  be an  $\aleph_0$ -continuous regular ring, let  $A$  and  $B$  be finitely generated projective right  $R$ -modules, and let  $A_1 \leq A_2 \leq \dots$  be an ascending sequence of finitely generated submodules of  $A$ . If  $\bigcup A_n$  is essential in  $A$ , and each  $A_n \lesssim B$ , then  $A \lesssim B$ .*

**PROOF.** Let  $S$  be the maximal right  $\aleph_0$ -quotient ring of  $R$  [2, pp. 177, 178], so that  $S$  is an  $\aleph_0$ -continuous, regular, right and left  $\aleph_0$ -injective overring of  $R$  [2, Theorems 14.12 and 14.17]. Now  $A \otimes_R S$  and  $B \otimes_R S$  are finitely generated projective right  $S$ -modules,  $A \otimes_R S$  has an ascending sequence of finitely generated submodules which may be labelled  $A_1 \otimes_R S \leq A_2 \otimes_R S \leq \dots$ , the submodule  $\bigcup (A_n \otimes_R S)$  is essential in  $A \otimes_R S$ , and each  $A_n \otimes_R S \lesssim B \otimes_R S$ . Moreover, if  $A \otimes_R S \lesssim B \otimes_R S$ , then [2, Proposition 14.28] shows that  $A \lesssim B$ .

Thus there is no loss of generality in assuming that  $R$  is right and left  $\aleph_0$ -injective. Consequently, all matrix rings over  $R$  are  $\aleph_0$ -continuous [2, Proposition 14.19]. Using the standard Morita-equivalences, we may transfer our problem to the category of right modules over any matrix ring  $M_k(R)$ . By choosing  $k$  large enough we may arrange for the new modules corresponding to  $A$  and  $B$  to be cyclic, hence isomorphic to right ideals of  $M_k(R)$ .

Therefore we may now assume, without loss of generality, that  $A$  and  $B$  are principal right ideals of  $R$ . Since  $\bigcup A_n$  is essential in  $A$ , we see that  $A$  is the supremum, in the lattice of principal right ideals of  $R$ , of the family  $\{A_n\}$ . Consequently, [2, Corollary 14.27] shows that  $A \lesssim B$ . □

**LEMMA 1.5.** *Let  $R$  be an  $\aleph_0$ -continuous regular ring, and let  $x, x_1, x_2, \dots$  be elements of  $R$ . If the right ideal  $\sum x_n R$  is essential in  $xR$ , then  $N^*(x) \leq \sum N^*(x_n)$ .*

**PROOF.** Choose elements  $y_1, y_2, \dots \in R$  such that  $y_n R = x_1 R + \dots + x_n R$  for all  $n$ , and note from Lemma 1.2 that  $N^*(y_n) \leq N^*(x_1) + \dots + N^*(x_n)$ . Thus it suffices

to show that  $N^*(x) \leq \sup\{N^*(y_n)\}$ . If not,

$$N^*(x) > s/t > \sup\{N^*(y_n)\}$$

for some  $s, t \in \mathbb{N}$ .

For each  $n$ , we have  $P(y_n) \leq N^*(y_n) < s/t$  and so  $tP(y_n) < sP(1)$ , for all  $P \in \mathbf{P}(R)$ . As a result, [2, Theorem 18.28] implies that  $t(y_nR) \lesssim sR_R$ . Inside the projective module  $t(xR)$ , we have finitely generated submodules  $t(y_1R) \leq t(y_2R) \leq \dots$ , and  $\bigcup(t(y_nR))$  is an essential submodule of  $t(xR)$ . Consequently, Lemma 1.4 says that  $t(xR) \lesssim sR_R$ . But then  $N^*(x) \leq s/t$  (by Lemma 1.2), which is false.

Therefore  $N^*(x) \leq \sup\{N^*(y_n)\}$ .  $\square$

The key to the upcoming completeness argument is the following lemma, which is a modification of a corresponding argument of von Neumann’s [10, Lemma 17.3, p. 228]. Another completeness argument using this method occurs in [2, Lemma 21.6], and we can adapt the proof of that lemma with only minor changes.

LEMMA 1.6. *If  $R$  is an  $\aleph_0$ -continuous regular ring and  $e$  is an idempotent in  $R$ , then  $eR(1 - e)$  is complete in the  $N^*$ -metric.*

PROOF. We shall need the fact that  $R$  is unit-regular [2, Theorem 14.24].

Let  $\{x_1, x_2, \dots\}$  be a sequence in  $eR(1 - e)$  which is Cauchy in the  $N^*$ -metric. By passing to a subsequence, we may assume that  $N^*(x_i - x_j) < 1/2^{k+1}$  whenever  $i, j \geq k$ . Set  $A_n = (1 - e + x_n)R$  for all  $n$ , and note that  $A_n + eR = R$ . For each  $k = 1, 2, \dots$ , there exists a principal right ideal  $B_k$  in  $R$  such that

$$\sum_{n=k}^{\infty} A_n \leq_e B_k,$$

and since  $A_k \leq B_k$ , we see that  $B_k + eR = R$ . Note that  $B_1 \supseteq B_2 \supseteq \dots$ , and set

$$C = \bigcap_{k=1}^{\infty} B_k.$$

Inasmuch as the lattice of principal right ideals of  $R$  is  $\aleph_0$ -continuous, we obtain

$$C + eR = \left( \bigcap_{k=1}^{\infty} B_k \right) + eR = \bigcap_{k=1}^{\infty} (B_k + eR) = R.$$

Consequently, there exists an idempotent  $f \in R$  such that  $fR = eR$  and  $(1 - f)R \leq C$ .

Since  $fR = eR$ , we have  $f = ef$  and  $e = fe$ , hence the element  $x = e - f$  lies in  $eR(1 - e)$ . We shall show that  $x_n \rightarrow x$  in the  $N^*$ -metric.

For each  $k = 1, 2, \dots$ , there exists an element  $d_k \in R$  such that

$$\sum_{j=k+1}^{\infty} (x_j - x_{j-1})R \leq_e d_k R,$$

and Lemma 1.5 shows that

$$N^*(d_k) \leq \sum_{j=k+1}^{\infty} N^*(x_j - x_{j-1}) < \sum_{j=k+1}^{\infty} 1/2^j = 1/2^k.$$

For all  $n \geq k$ , we have

$$A_n = (1 - e + x_n)R = \left(1 - e + x_k + \sum_{j=k+1}^n (x_j - x_{j-1})\right)R$$

$$\leq (1 - e + x_k)R + \sum_{j=k+1}^n (x_j - x_{j-1})R \leq A_k + d_k R.$$

Consequently,  $\sum_{n=k}^\infty A_n \leq A_k + d_k R$ , whence  $B_k \leq A_k + d_k R$ .

Each  $B_k = A_k \oplus u_k R$  for some  $u_k \in R$ . Then

$$A_k \oplus u_k R = B_k \leq A_k + d_k R \lesssim A_k \oplus d_k R$$

and so  $u_k R \lesssim d_k R$  (because  $R$  is unit-regular). As a result,  $N^*(u_k) \leq N^*(d_k) < 1/2^k$  (Lemma 1.2).

The idempotent  $1 - f$  lies in the right ideal

$$C \leq B_k = A_k + u_k R = (1 - e + x_k)R + u_k R,$$

hence  $1 - f = (1 - e + x_k)r + u_k s$  for some  $r, s \in R$ . Since  $x_k \in eR(1 - e)$ , we see that  $1 - e + x_k$  is idempotent, whence

$$(1 - e + x_k)(1 - f) = (1 - e + x_k)r + (1 - e + x_k)u_k s$$

$$= 1 - f + (x_k - e)u_k s.$$

In addition, since  $fR = eR$ , we have  $R(1 - f) = R(1 - e)$ , and so  $1 - e + x_k$  lies in  $R(1 - f)$ . Thus

$$1 - e + x_k = (1 - e + x_k)(1 - f) = 1 - f + (x_k - e)u_k s,$$

and consequently

$$x_k - x = x_k - e + f = (1 - e + x_k) - (1 - f) = (x_k - e)u_k s,$$

hence  $N^*(x_k - x) \leq N^*(u_k) < 1/2^k$ .

Therefore  $x_k \rightarrow x$  in the  $N^*$ -metric.  $\square$

We shall apply Lemma 1.6 in a situation where  $R$  is a matrix ring and  $e$  is a corner idempotent. For this purpose, and for later use, we need the following information concerning  $N^*$  in matrix rings.

**LEMMA 1.7.** *Let  $R$  be a regular ring, let  $n \in \mathbb{N}$ , and set  $T = M_n(R)$ . Let  $\varphi: R \rightarrow T$  be the natural map, and let  $\{e_{ij} \mid i, j = 1, \dots, n\}$  be the standard matrix units in  $T$ .*

- (a)  $N_T^* \varphi = N_R^*$ .
- (b)  $N_T^*(\varphi(x)e_{ij}) = N_R^*(x)/n$  for all  $x \in R$  and all  $i, j$ .
- (c)  $N_R^*(y_{ij}) \leq nN_T^*(y)$  for all  $y \in T$  and all  $i, j$ .
- (d)  $N_T^*(y) \leq \sum_{i,j=1}^n N_R^*(y_{ij})/n$  for all  $y \in T$ .

**PROOF.** (a) According to [2, Corollary 16.10], the rule  $P \mapsto P\varphi$  defines a bijection of  $\mathbf{P}(T)$  onto  $\mathbf{P}(R)$ , hence

$$N_R^*(x) = \sup\{P\varphi(x) \mid P \in \mathbf{P}(T)\} = N_T^*\varphi(x)$$

for any  $x \in R$ .

(b) For each  $k = 1, \dots, n$ , we note that left multiplication by  $e_{ki}$  defines an isomorphism of  $\varphi(x)e_{ij}T$  onto  $\varphi(x)e_{kk}T$ . Consequently,

$$\varphi(x)T = \bigoplus_{k=1}^n \varphi(x)e_{kk}T \cong n(\varphi(x)e_{ij}T),$$

hence  $N_T^*\varphi(x) = nN_T^*(\varphi(x)e_{ij})$ , by Lemma 1.2.

(c) Since  $\varphi(y_{ij})e_{ij} = e_{ii}ye_{jj}$ , it follows from (b) that

$$N_R^*(y_{ij}) = nN_T^*(\varphi(y_{ij})e_{ij}) = nN_T^*(e_{ii}ye_{jj}) \leq nN_T^*(y).$$

(d) Using (b) again, we conclude that

$$\begin{aligned} N_T^*(y) &= N_T^*\left(\sum_{i,j=1}^n \varphi(y_{ij})e_{ij}\right) \leq \sum_{i,j=1}^n N_T^*(\varphi(y_{ij})e_{ij}) \\ &= \sum_{i,j=1}^n N_R^*(y_{ij})/n. \quad \square \end{aligned}$$

**THEOREM 1.8.** *Every  $\aleph_0$ -continuous regular ring is  $N^*$ -complete.*

**PROOF.** For any  $\aleph_0$ -continuous regular ring  $R$ , [4, Proposition II.11.4] shows that  $\ker(\mathbf{P}(R)) = 0$ .

Assume for the moment that  $R$  is right and left  $\aleph_0$ -injective. Then the ring  $T = M_2(R)$  is  $\aleph_0$ -continuous, by [2, Proposition 14.19]. Setting  $e = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ , we obtain from Lemma 1.6 that  $eT(1 - e)$  is complete in the  $N_T^*$ -metric. There is a group isomorphism  $x \mapsto \begin{pmatrix} 0 & 0 \\ x & 0 \end{pmatrix}$  of  $R$  onto  $eT(1 - e)$ , and Lemma 1.7 shows that

$$N_R^*(x) = 2N_T^*\left(\begin{pmatrix} 0 & 0 \\ x & 0 \end{pmatrix}\right)$$

for all  $x \in R$ . Therefore in this case  $R$  is  $N^*$ -complete.

In general, let  $S$  denote the maximal right  $\aleph_0$ -quotient ring of  $R$ . Then  $S$  is an  $\aleph_0$ -continuous, regular, right and left  $\aleph_0$ -injective overring of  $R$ , and  $R$  contains all the idempotents of  $S$  [2, Theorems 14.12 and 14.17]. By the case above,  $S$  is  $N^*$ -complete.

Now  $R$  and  $S$  are unit-regular rings [2, Theorem 14.24], and we may assume they are nonzero. Using Proposition 1.1 and [2, Proposition 14.28], we compute that for any  $x \in R$ ,

$$\begin{aligned} N_R^*(x) &= \inf\{k/n \mid k, n \in \mathbb{N} \text{ and } n(xR) \lesssim kR_R\} \\ &= \inf\{k/n \mid k, n \in \mathbb{N} \text{ and } n(xS) \lesssim kS_S\} = N_S^*(x). \end{aligned}$$

Consequently, the  $N^*$ -completeness of  $R$  will follow from the  $N^*$ -completeness of  $S$  provided  $R$  is closed in  $S$  in the  $N_S^*$ -metric.

We claim that  $N_S^*(x) = 1$  for any  $x \in S - R$ . By [2, Proposition 3.15],  $S$  has a two-sided ideal  $N$  such that  $N \subseteq R$  and the ring  $S/N$  is abelian. Then  $x \notin N$ , hence  $S$  has a primitive ideal  $M$  such that  $N \subseteq M$  but  $x \notin M$ . As  $S/N$  is abelian,  $S/M$  is a division ring, hence  $S/M$  has a unique rank function  $Q$  and  $Q(x + M) = 1$ . Pulling  $Q$  back to a pseudo-rank function  $P$  on  $S$ , we obtain  $P(x) = 1$ , so that  $N_S^*(x) \geq 1$ . Thus  $N_S^*(x) = 1$ , as claimed.

As a result,  $N_S^*(x - y) = 1$  for all  $x \in S - R$  and all  $y \in R$ , whence  $R$  is closed in  $S$  in the  $N_S^*$ -metric, as desired. Therefore  $R$  is  $N^*$ -complete.  $\square$

**COROLLARY 1.9.** *Every regular, right and left self-injective ring is  $N^*$ -complete.*  $\square$

We conclude this section by proving that  $N^*$ -completeness carries over to matrix rings and certain factor rings.

**THEOREM 1.10.** *If  $R$  is an  $N^*$ -complete regular ring, then any matrix ring  $M_n(R)$  is  $N^*$ -complete.*

**PROOF.** Set  $T = M_n(R)$ , let  $\varphi: R \rightarrow T$  be the natural map, and let  $\{e_{ij} \mid i, j = 1, \dots, n\}$  be the standard matrix units in  $T$ . Given any  $y \in \ker(\mathbf{P}(T))$ , we have  $N_T^*(y) = 0$ , whence Lemma 1.7 shows that  $N_R^*(y_{ij}) = 0$  for all  $i, j$ . Then all  $y_{ij} = 0$ , so that  $y = 0$ . Thus  $\ker(\mathbf{P}(T)) = 0$ .

Now consider any sequence  $\{y^{(1)}, y^{(2)}, \dots\}$  in  $T$  that is Cauchy with respect to  $N_T^*$ . Fixing  $i$  and  $j$  for a while, we have

$$N_R^*(y_{ij}^{(k)} - y_{ij}^{(l)}) \leq nN_T^*(y^{(k)} - y^{(l)})$$

for all  $k, l$  (Lemma 1.7), hence the sequence  $\{y_{ij}^{(1)}, y_{ij}^{(2)}, \dots\}$  in  $R$  is Cauchy with respect to  $N_R^*$ . Consequently, there exists  $y_{ij} \in R$  such that  $y_{ij}^{(k)} \rightarrow y_{ij}$  in the  $N_R^*$ -metric. Having gotten such elements  $y_{ij}$  for each  $i, j$ , we obtain a matrix  $y \in T$  with entries  $y_{ij}$ . Inasmuch as

$$N_T^*(y^{(k)} - y) \leq \sum_{i,j=1}^n N_R^*(y_{ij}^{(k)} - y_{ij})/n$$

for all  $k$  (Lemma 1.7 again), we conclude that  $y^{(k)} \rightarrow y$  in the  $N_T^*$ -metric.

Therefore  $T$  is  $N^*$ -complete.  $\square$

**LEMMA 1.11.** *Let  $J$  be a two-sided ideal in a regular ring  $R$ , let  $A$  and  $B$  be finitely generated projective right  $R$ -modules, and let  $n \in \mathbf{N}$ . If  $n(A/AJ) \lesssim B/BJ$ , then there exists a decomposition  $A = A' \oplus A''$  such that  $nA' \lesssim B$  and  $A'' = A''J$ .*

**PROOF.** As  $(nA)/(nA)J \lesssim B/BJ$ , we may apply [2, Proposition 2.20] to obtain a decomposition  $nA = C \oplus D$  such that  $C \lesssim B$  and  $D = DJ$ . Then, by [2, Theorem 2.8], there exist decompositions  $C = C_1 \oplus \dots \oplus C_n$  and  $D = D_1 \oplus \dots \oplus D_n$  such that  $C_i \oplus D_i \cong A$  for all  $i$ . Since  $D = DJ$ , each  $D_i = D_iJ$ , hence  $C_i/C_iJ \cong A/AJ$ . Thus the modules  $C_i/C_iJ$  are pairwise isomorphic, so by [2, Proposition 2.19] there exist decompositions  $C_i = E_i \oplus F_i$  for each  $i$  such that the  $E_i$  are pairwise isomorphic and each  $F_i = F_iJ$ .

Now  $A \cong C_1 \oplus D_1 = E_1 \oplus F_1 \oplus D_1$ , so there is a decomposition  $A = A' \oplus A''$  with  $A' \cong E_1$  and  $A'' \cong F_1 \oplus D_1$ . Then

$$nA' \cong nE_1 \cong E_1 \oplus \dots \oplus E_n \leq C_1 \oplus \dots \oplus C_n = C \lesssim B.$$

Since  $D_1 = D_1J$  and  $F_1 = F_1J$ , we also have  $A'' = A''J$ .  $\square$

For the moment, we now restrict to unit-regular rings. This restriction will be removed when we prove that all  $N^*$ -complete regular rings are unit-regular (Theorem 2.3).



LEMMA 1.12. *Let  $J$  be a two-sided ideal in a unit-regular ring  $R$ . Then*

$$N_{R/J}^*(x + J) = \inf\{N_R^*(y) \mid y \in x + J\}$$

for all  $x \in R$ .

PROOF. This is clear if  $J = R$ , so assume  $J \neq R$ .

Given any  $y \in x + J$ , we see that

$$\begin{aligned} N_{R/J}^*(x + J) &= N_{R/J}^*(y + J) = \sup\{Q(y + J) \mid Q \in \mathbf{P}(R/J)\} \\ &= \sup\{P(y) \mid P \in \mathbf{P}(R) \text{ and } J \subseteq \ker(P)\} \\ &\leq \sup\{P(y) \mid P \in \mathbf{P}(R)\} = N_R^*(y). \end{aligned}$$

Thus  $N_{R/J}^*(x + J) \leq \inf\{N_R^*(y) \mid y \in x + J\}$ .

Given any real number  $\alpha > N_{R/J}^*(x + J)$ , Proposition 1.1 shows that there exist  $k, n \in \mathbf{N}$  for which  $k/n < \alpha$  and

$$n((x + J)(R/J)) \lesssim k(R/J),$$

that is,  $n(xR/xJ) \lesssim (kR)/(kR)J$ . According to Lemma 1.11, there is a decomposition  $xR = A' \oplus A''$  such that  $nA' \lesssim kR_R$  and  $A'' = A''J$ . Then  $x = y + z$  for some  $y \in A'$  and  $z \in A''$ . Note that  $z \in J$ , so that  $y \in x + J$ . Since  $n(yR) \leq nA' \lesssim kR_R$ , we conclude from Proposition 1.1 that  $N_R^*(y) \leq k/n < \alpha$ . Therefore

$$\inf\{N_R^*(y) \mid y \in x + J\} \leq N_{R/J}^*(x + J). \quad \square$$

THEOREM 1.13. *Let  $R$  be an  $N^*$ -complete unit-regular ring, and let  $J$  be a two-sided ideal of  $R$ . Then the following conditions are equivalent.*

- (a)  $R/J$  is  $N^*$ -complete.
- (b)  $J$  is  $N^*$ -closed in  $R$ .
- (c)  $J = \ker(X)$  for some  $X \subseteq \mathbf{P}(R)$ .

PROOF. (a)  $\Rightarrow$  (c): By definition,  $\ker(\mathbf{P}(R/J)) = 0$ , hence if

$$X = \{P \in \mathbf{P}(R) \mid J \subseteq \ker(P)\},$$

then  $J = \ker(X)$ .

(c)  $\Rightarrow$  (b): Given  $x \in R - J$ , we must have  $P(x) > 0$  for some  $P \in X$ . Then

$$N^*(y - x) \geq P(y - x) \geq P(x) - P(y) = P(x)$$

for all  $y \in J$ , hence  $x$  is not in the  $N^*$ -closure of  $J$ . Thus  $J$  is  $N^*$ -closed.

(b)  $\Rightarrow$  (a): Given a nonzero coset  $x + J$  in  $R/J$ , we have  $x \notin J$ , hence there must exist a positive real number  $\epsilon$  such that  $N_R^*(y - x) \geq \epsilon$  for all  $y \in J$ . Then  $N_R^*(z) \geq \epsilon$  for all  $z \in x + J$ , whence  $N_{R/J}^*(x + J) \geq \epsilon$ , by Lemma 1.12. Consequently,  $\ker(\mathbf{P}(R/J)) = 0$ .

Now consider a sequence  $\{a_1, a_2, \dots\}$  in  $R/J$  that is Cauchy with respect to  $N_{R/J}^*$ . There is no loss of generality in assuming that  $N_{R/J}^*(a_{n+1} - a_n) < 1/2^n$  for all  $n$ . Choose  $x_1, x_2, \dots$  in  $R$  such that each  $a_n = x_n + J$ .

Set  $y_1 = x_1$ . Then  $N_{R/J}^*((x_2 - y_1) + J) < 1/2$ , so by Lemma 1.12 there exists  $z$  in  $(x_2 - y_1) + J$  satisfying  $N_R^*(z) < 1/2$ . Set  $y_2 = y_1 + z$ , so that  $y_2 + J = x_2 + J = a_2$  and  $N_R^*(y_2 - y_1) < 1/2$ . Continuing in this manner, we obtain elements  $y_1, y_2, \dots$  in  $R$  such that  $y_n + J = a_n$  and  $N_R^*(y_{n+1} - y_n) < 1/2^n$  for all  $n$ .

Thus  $\{y_1, y_2, \dots\}$  is Cauchy with respect to  $N_R^*$ , hence there exists  $y \in R$  such that  $y_n \rightarrow y$  in the  $N_R^*$ -metric. Setting  $a = y + J$ , we conclude from Lemma 1.12 that  $a_n \rightarrow a$  in the  $N_{R/J}^*$ -metric.

Therefore  $R/J$  is  $N^*$ -complete.  $\square$

**COROLLARY 1.14.** *Let  $R$  be an  $N^*$ -complete unit-regular ring, and let  $M$  be a maximal two-sided ideal of  $R$ . Then  $R/M$  is  $N^*$ -complete.*

**PROOF.** As  $R/M$  is a simple unit-regular ring, [2, Corollary 18.5] shows that there is a rank function on  $R/M$ . Then there exists  $P \in \mathbf{P}(R)$  such that  $\ker(P) = M$ , hence  $R/M$  is  $N^*$ -complete by Theorem 1.13.  $\square$

**II.  $K_0$ .** A good deal of information about a regular ring  $R$ , particularly ideal theory and decomposition properties of the principal right ideals, is stored in the Grothendieck group  $K_0(R)$ . We study this group for an  $N^*$ -complete regular ring  $R$  in this section, proving that  $K_0(R)$  is an archimedean, norm-complete, partially ordered abelian group with the interpolation property. In the following section, we develop a structure theory for such groups, which can then be applied, via  $K_0$ , to the structure theory of  $N^*$ -complete regular rings.

**DEFINITION.** Recall that the Grothendieck group  $K_0$  of a ring  $R$  is an abelian group with generators  $[A]$  corresponding to the finitely generated projective right  $R$ -modules  $A$  and with relations  $[A] + [B] = [C]$  whenever  $A \oplus B \cong C$ . All elements of  $K_0(R)$  are of the form  $[A] - [B]$ , for suitable  $A$  and  $B$ . We set

$$K_0(R)^+ = \{[A] \mid A \text{ is a finitely generated projective right } R\text{-module}\},$$

and we define a relation  $\leq$  on  $K_0(R)$  so that  $x \leq y$  if and only if  $y - x$  lies in  $K_0(R)^+$ . This relation is a translation-invariant pre-order on  $K_0(R)$ , so that  $K_0(R)$  becomes a pre-ordered abelian group. The element  $[R]$  is an *order-unit* in  $K_0(R)$ , meaning that for any  $x \in K_0(R)$  there exists  $n \in \mathbf{N}$  such that  $x \leq n[R]$ .

For a unit-regular ring  $R$ , the relations between  $K_0(R)$  and the finitely generated projective right  $R$ -modules are much cleaner than in general. Namely, for any finitely generated projective right  $R$ -modules  $A, B, C, D$  we have

$$\begin{aligned} [A] - [B] &= [C] - [D] && \text{if and only if } A \oplus D \cong B \oplus C, \\ [A] - [B] &\leq [C] - [D] && \text{if and only if } A \oplus D \lesssim B \oplus C \end{aligned}$$

[2, Proposition 15.2]. In addition, the relation  $\leq$  on  $K_0(R)$  is actually a partial order, so that  $K_0(R)$  is a partially ordered abelian group in this case. Thus, in order to deal effectively with  $K_0$  of  $N^*$ -complete regular rings, we first prove that such rings are unit-regular. Two lemmas will be helpful in doing this.

**LEMMA 2.1.** *Let  $R$  be an  $N^*$ -complete regular ring, and let  $A, B, C$  be finitely generated projective right  $R$ -modules. Let  $\{A_1, A_2, \dots\}$  and  $\{B_1, B_2, \dots\}$  be independent sequences of finitely generated submodules of  $A$  and  $B$ , such that  $A_k \cong B_k$  for all  $k$ . For each  $k$ , let  $A_k^*$  be a submodule of  $A$  such that*

$$A = A_1 \oplus \dots \oplus A_k \oplus A_k^*,$$

*and assume that  $2^k t_k A_k^* \lesssim t_k C$  for some  $t_k \in \mathbf{N}$ . Then  $A \lesssim B$ .*

PROOF. As all matrix rings over  $R$  are  $N^*$ -complete (Theorem 1.10), we may use the standard Morita-equivalences to transfer our problem to the category of right modules over a suitable matrix ring  $M_n(R)$ , with  $n$  chosen large enough so that the modules corresponding to  $A, B, C$  are cyclic. Thus there is no loss of generality in assuming that  $A, B, C$  are actually principal right ideals of  $R$ .

Choose idempotents  $e, f \in R$  such that  $eR = A$  and  $fR = B$ . Applying [2, Proposition 2.13] to the ascending sequence

$$A_1 \leq A_1 \oplus A_2 \leq A_1 \oplus A_2 \oplus A_3 \leq \dots$$

of finitely generated submodules of  $A$ , we obtain orthogonal idempotents  $e_1, e_2, \dots$  in  $eRe$  such that

$$e_1R \oplus \dots \oplus e_kR = A_1 \oplus \dots \oplus A_k$$

for all  $k$ . Similarly, there exist orthogonal idempotents  $f_1, f_2, \dots$  in  $fRf$  such that

$$f_1R \oplus \dots \oplus f_kR = B_1 \oplus \dots \oplus B_k$$

for all  $k$ . Note that each

$$\begin{aligned} e_kR &\cong (e_1R \oplus \dots \oplus e_kR) / (e_1R \oplus \dots \oplus e_{k-1}R) \\ &= (A_1 \oplus \dots \oplus A_k) / (A_1 \oplus \dots \oplus A_{k-1}) \cong A_k \end{aligned}$$

and similarly  $f_kR \cong B_k$ , so that  $e_kR \cong f_kR$ . Thus there exist elements  $x_k \in e_kRf_k$  and  $y_k \in f_kRe_k$  such that  $x_k y_k = e_k$  and  $y_k x_k = f_k$ .

For each  $k$ , we have  $(e - e_1 - \dots - e_k)R \cong A_k^*$ , whence

$$2^k t_k((e - e_1 - \dots - e_k)R) \lesssim t_k C \lesssim t_k R,$$

and consequently  $N^*(e - e_1 - \dots - e_k) \leq 1/2^k$ , by Lemma 1.2. Thus  $\sum e_k \rightarrow e$  in the  $N^*$ -metric. As

$$x_{k+1} = e_{k+1} x_{k+1} = (e - e_1 - \dots - e_k) e_{k+1} x_{k+1}$$

for each  $k$ , we also obtain  $N^*(x_{k+1}) \leq 1/2^k$ , and similarly,  $N^*(y_{k+1}) \leq 1/2^k$ .

Now the partial sums of the series  $\sum x_k$  and  $\sum y_k$  are Cauchy with respect to  $N^*$ , hence there exist  $x, y \in R$  such that  $\sum x_k \rightarrow x$  and  $\sum y_k \rightarrow y$  in the  $N^*$ -metric. As each

$$x_k = e_k x_k f_k = e e_k x_k f_k f = e x_k f,$$

we obtain  $x = exf$ , and likewise  $y = fye$ . Since  $x_i y_j = x_i f_i f_j y_j = 0$  whenever  $i \neq j$ , we have

$$(x_1 + \dots + x_k)(y_1 + \dots + y_k) = x_1 y_1 + \dots + x_k y_k = e_1 + \dots + e_k$$

for all  $k$ , and consequently  $xy = e$ . Therefore  $eR \lesssim fR$ , that is,  $A \lesssim B$ .  $\square$

LEMMA 2.2. *Let  $A, B, C$  be finitely generated projective right modules over a regular ring, such that  $A \oplus C \cong B \oplus C$ . Then there exist decompositions*

$$A = A' \oplus A''; \quad B = B' \oplus B''; \quad C = C' \oplus C''$$

*such that  $A' \cong B'$  and  $A'' \cong C'$ , while also  $A'' \oplus C'' \cong B'' \oplus C''$ .*

PROOF. According to [2, Theorem 2.8], there exist decompositions  $A = A' \oplus A''$  and  $C = D \oplus E$  such that  $A' \oplus D \cong B$  and  $A'' \oplus E \cong C$ . Then we obtain decompositions  $B = B' \oplus B''$  and  $C = C' \oplus C''$  such that  $B' \cong A'$  and  $B'' \cong D$ , while also  $C' \cong A''$  and  $C'' \cong E$ . Finally,

$$A'' \oplus C'' \cong A'' \oplus E \cong C = D \oplus E \cong B'' \oplus C''. \quad \square$$

THEOREM 2.3. *Every  $N^*$ -complete regular ring is unit-regular.*

PROOF. Given an  $N^*$ -complete regular ring  $R$ , we have  $\ker(\mathbf{P}(R)) = 0$  by definition, hence all matrix rings over  $R$  are directly finite [2, Proposition 16.11]. To prove that  $R$  is unit-regular, it suffices to show that if  $A, B, C$  are any finitely generated projective right  $R$ -modules satisfying  $A \oplus C \cong B \oplus C$ , then  $A \cong B$ .

Inducting on Lemma 2.2, we obtain submodules

$$A_1, A'_1, A'_2, A'_2, \dots \leq A; \quad B_1, B''_1, B'_2, B''_2, \dots \leq B; \quad C_1, C''_1, C'_2, C''_2, \dots \leq C$$

such that

$$A = A_1 \oplus A'_1; \quad B = B_1 \oplus B''_1; \quad C = C_1 \oplus C''_1$$

while also

$$A'_i = A'_{i+1} \oplus A''_{i+1}; \quad B''_i = B'_{i+1} \oplus B''_{i+1}; \quad C''_i = C'_{i+1} \oplus C''_{i+1}; \\ A'_i \cong B'_i; \quad A'_i \cong C'_i; \quad A'_i \oplus C''_i \cong B''_i \oplus C''_i$$

for all  $i$ . Note that  $A'_1 \geq A'_2 \geq \dots$ , and that  $C_1, C_2, \dots$  are independent submodules of  $C$ . Consequently, we obtain

$$iA'_i \lesssim A'_1 \oplus A'_2 \oplus \dots \oplus A'_i \cong C_1 \oplus C_2 \oplus \dots \oplus C_i \leq C$$

for all  $i$ .

Now set  $A_1 = A'_1 \oplus A'_2$  and  $B_1 = B'_1 \oplus B'_2$ , while  $A_1^* = A'_2$ . Set

$$A_k = \bigoplus_{i=2^{k-1}+1}^{2^k} A'_i; \quad A_k^* = A'_{2^k}; \quad B_k = \bigoplus_{i=2^{k-1}+1}^{2^k} B'_i$$

for all  $k = 2, 3, \dots$ . Thus  $\{A_1, A_2, \dots\}$  and  $\{B_1, B_2, \dots\}$  are independent sequences of finitely generated submodules of  $A$  and  $B$ , with  $A_k \cong B_k$  for all  $k$ . Also, since

$$A = A'_1 \oplus A'_1 = A'_1 \oplus A'_2 \oplus A'_2 = \dots = A'_1 \oplus A'_2 \oplus \dots \oplus A'_i \oplus A'_i$$

for all  $i$ , we have  $A = A_1 \oplus A_2 \oplus \dots \oplus A_k \oplus A_k^*$  for all  $k$ . Inasmuch as  $2^k A_k^* \lesssim C$  for all  $k$ , we conclude from Lemma 2.1 that  $A \lesssim B$ .

By symmetry,  $B \lesssim A$ . As all matrix rings over  $R$  are directly finite, it follows that  $A \cong B$  [2, Proposition 5.4].  $\square$

DEFINITION. Let  $G$  be a partially ordered abelian group. Then  $G$  is said to be an interpolation group if given any  $x_1, x_2, y_1, y_2$  in  $G$  satisfying  $x_i \leq y_j$  for all  $i, j$ , there exists  $z \in G$  such that  $x_i \leq z \leq y_j$  for all  $i, j$ . Equivalently,  $G$  is an interpolation group if and only if either of the following forms of the Riesz decomposition property holds.

(a) If  $x, y_1, \dots, y_n \in G^+$  and  $x \leq y_1 + \dots + y_n$ , then there exist  $x_1, \dots, x_n \in G^+$  such that  $x = x_1 + \dots + x_n$  and each  $x_i \leq y_i$ .

(b) If  $x_1, \dots, x_n, y_1, \dots, y_k \in G^+$  and  $x_1 + \dots + x_n = y_1 + \dots + y_k$ , then there exist  $z_{ij} \in G^+$  (for  $i = 1, \dots, n$  and  $j = 1, \dots, k$ ) such that  $x_i = z_{i1} + \dots + z_{ik}$  for all  $i$  and  $y_j = z_{1j} + \dots + z_{nj}$  for all  $j$ .

For any unit-regular ring  $R$ , the partially ordered abelian group  $K_0(R)$  is an interpolation group [4, Proposition II.10.3]. In particular, this holds for any  $N^*$ -complete regular ring, because of Theorem 2.3; however, we postpone recording this fact until Theorem 2.11.

DEFINITION. Let  $G$  be a partially ordered abelian group, and let  $n$  be a positive integer. We say that  $G$  is  $n$ -unperforated if for all  $x \in G$ , we have  $nx \geq 0$  only when  $x \geq 0$ . If  $G$  is  $n$ -unperforated for all  $n \in \mathbb{N}$ , then  $G$  is said to be unperforated. The group  $G$  is archimedean provided that whenever  $x, y \in G$  and  $nx \leq y$  for all  $n \in \mathbb{N}$ , then  $x \leq 0$ . It is easily checked that all archimedean directed abelian groups are unperforated [4, Lemma I.5.2]. Our next goal is to prove that  $K_0$  of any  $N^*$ -complete regular ring  $R$  is archimedean; however, since our proof requires  $K_0(R)$  to be 2-unperforated, we show that first.

LEMMA 2.4. Let  $A$  and  $B$  be finitely generated projective right modules over a unit-regular ring  $R$ , and let  $n \in \mathbb{N}$ . If  $A \lesssim nB$ , then there exists a decomposition  $A = A' \oplus A''$  such that  $A' \lesssim B$  and  $nA'' \lesssim (n - 1)A$ .

PROOF. Since  $A \lesssim nB$ , there is a decomposition  $A = A_1 \oplus \dots \oplus A_n$  with each  $A_i \lesssim B$  [2, Corollary 2.9]. Each  $A_i$  is isomorphic to a submodule  $B_i \leq B$ , and we define  $B' = B_1 + \dots + B_n$ . Then  $B'$  is a finitely generated submodule of  $B$ , and

$$B' \lesssim B_1 \oplus \dots \oplus B_n \cong A_1 \oplus \dots \oplus A_n = A,$$

hence we obtain a decomposition  $A = A' \oplus A''$  with  $A' \cong B' \leq B$ . For each  $j = 1, \dots, n$ , we have  $A_j \cong B_j \leq B' \cong A'$  and so

$$A' \oplus A'' = A = A_1 \oplus \dots \oplus A_n \lesssim A' \oplus \left( \bigoplus_{i \neq j} A_i \right),$$

whence  $A'' \lesssim \bigoplus_{i \neq j} A_i$ . Therefore

$$nA'' \lesssim \bigoplus_{j=1}^n \bigoplus_{i \neq j} A_i \cong \bigoplus_{i=1}^n (n - 1)A_i \cong (n - 1)A. \quad \square$$

LEMMA 2.5. Let  $A$  and  $B$  be finitely generated projective right modules over a unit-regular ring  $R$ , and let  $n \in \mathbb{N}$ . If  $nA \lesssim nB$ , then there exist decompositions  $A = A' \oplus A''$  and  $B = B' \oplus B''$  such that  $A' \cong B'$  and  $nA'' \lesssim nB''$ , while also  $2tA'' \lesssim tA$  for some  $t \in \mathbb{N}$ .

PROOF. By Lemma 2.4, there exist decompositions  $A = A_1 \oplus A_1^*$  and  $B = B_1 \oplus B_1^*$  such that  $A_1 \cong B_1$  and  $nA_1^* \lesssim (n - 1)A$ . Since

$$nA_1 \oplus nA_1^* \cong nA \lesssim nB = nB_1 \oplus nB_1^* \cong nA_1 \oplus nB_1^*,$$

we also have  $nA_1^* \lesssim nB_1^*$ . Thus we may continue by induction, obtaining submodules

$$A_1, A_1^*, A_2, A_2^*, \dots \leq A; \quad B_1, B_1^*, B_2, B_2^*, \dots \leq B$$

such that

$$A_i^* = A_{i+1} \oplus A_{i+1}^*; \quad B_i^* = B_{i+1} \oplus B_{i+1}^*; \\ A_{i+1} \cong B_{i+1}; \quad nA_{i+1}^* \lesssim (n-1)A_i^*; \quad nA_{i+1}^* \lesssim nB_{i+1}^*$$

for all  $i$ . In addition,

$$n^i A_i^* \lesssim n^{i-1}(n-1)A_{i-1}^* \lesssim \cdots \lesssim n(n-1)^{i-1}A_1^* \lesssim (n-1)^i A$$

for all  $i$ .

Choose  $i \in \mathbb{N}$  such that  $((n-1)/n)^i \leq 1/2$ , and set  $t = (n-1)^i$ , so that  $2t \leq n^i$ . Setting  $A' = A_1 \oplus \cdots \oplus A_i$  and  $A'' = A_i^*$ , we obtain  $A = A' \oplus A''$  and

$$2tA'' \lesssim n^i A_i^* \lesssim (n-1)^i A = tA.$$

Setting  $B' = B_1 \oplus \cdots \oplus B_i$  and  $B'' = B_i^*$ , we obtain  $B = B' \oplus B''$ , while also  $A' \cong B'$  and  $nA'' \lesssim nB''$ .  $\square$

**THEOREM 2.6.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $A$  and  $B$  be finitely generated projective right  $R$ -modules, and let  $n \in \mathbb{N}$ . If  $nA \lesssim nB$ , then  $A \lesssim B$ .*

**PROOF.** Inducting on Lemma 2.5, we obtain submodules

$$A_1, A_1^*, A_2, A_2^*, \dots \leq A; \quad B_1, B_1^*, B_2, B_2^*, \dots \leq B$$

such that  $A = A_1 \oplus A_1^*$  and  $B = B_1 \oplus B_1^*$ , while also

$$A_k^* = A_{k+1} \oplus A_{k+1}^*; \quad B_k^* = B_{k+1} \oplus B_{k+1}^*; \\ A_k \cong B_k; \quad nA_k^* \lesssim nB_k^*; \quad 2^k t_k A_k^* \lesssim t_k A$$

(for some  $t_k \in \mathbb{N}$ ) for all  $k$ . Applying Lemma 2.1, we conclude that  $A \lesssim B$ .  $\square$

As an  $N^*$ -complete regular ring  $R$  is unit-regular, it follows immediately from Theorem 2.6 that  $K_0(R)$  is unperforated. This result will be subsumed by the stronger result that  $K_0(R)$  is archimedean (Theorem 2.11).

**LEMMA 2.7.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $A, B, C$  be finitely generated projective right  $R$ -modules, and let  $n \in \mathbb{N}$ . If  $2^n A \lesssim 2^n B \oplus C$ , then there exists a decomposition  $A = A' \oplus A''$  such that  $A' \lesssim B$  and  $2^n A'' \lesssim C$ .*

**PROOF.** The proof of [2, Lemma 14.31] may be used, substituting Theorem 2.6 for [2, Theorem 14.30].  $\square$

**THEOREM 2.8.** *Let  $R$  be an  $N^*$ -complete regular ring, and let  $A, B, C$  be finitely generated projective right  $R$ -modules. If  $2^n A \lesssim 2^n B \oplus C$  for all  $n \in \mathbb{N}$ , then  $A \lesssim B$ .*

**PROOF.** Since  $2A \lesssim 2B \oplus C$ , Lemma 2.7 provides us with decompositions  $A = A_1 \oplus A_1^*$  and  $B = B_1 \oplus B_1^*$  such that  $A_1 \cong B_1$  and  $2A_1^* \lesssim C$ . For all  $n \in \mathbb{N}$ ,

$$2^n A_1 \oplus 2^n A_1^* \cong 2^n A \lesssim 2^n B \oplus C \cong 2^n B_1 \oplus 2^n B_1^* \oplus C \cong 2^n A_1 \oplus 2^n B_1^* \oplus C,$$

hence  $2^n A_1^* \lesssim 2^n B_1^* \oplus C$ . Thus we may continue by induction, obtaining submodules

$$A_1, A_1^*, A_2, A_2^*, \dots \leq A; \quad B_1, B_1^*, B_2, B_2^*, \dots \leq B$$

such that

$$A_k^* = A_{k+1} \oplus A_{k+1}^*; \quad B_k^* = B_{k+1} \oplus B_{k+1}^*; \\ A_k \cong B_k; \quad 2^k A_k^* \lesssim C; \quad 2^n A_k^* \lesssim 2^n B_k^* \oplus C$$

for all  $k, n$ . Applying Lemma 2.1, we conclude that  $A \lesssim B$ .  $\square$

It follows directly from Theorem 2.8 that  $K_0$  of any  $N^*$ -complete regular ring is archimedean. We postpone recording this result until Theorem 2.11.

**DEFINITION.** Let  $(G, u)$  be a partially ordered abelian group with order-unit. For any  $x \in G$ , we define

$$\|x\|_u = \inf\{k/n \mid k, n \in \mathbf{N} \text{ and } -ku \leq nx \leq ku\},$$

and we note that  $\|x\|_u$  is a nonnegative real number. When there is no danger of confusion as to the order-unit  $u$ , we just write  $\|x\|$  instead of  $\|x\|_u$ . The function  $\|\cdot\|$  behaves like a seminorm on  $G$ , for  $\|mx\| = |m| \cdot \|x\|$  and  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in G$  and  $m \in \mathbf{Z}$  [4, Lemma I.6.1]. In particular, it follows that the rule  $\delta(x, y) = \|x - y\|$  defines a pseudo-metric  $\delta$  on  $G$ . If  $\delta$  is actually a metric, and  $G$  is complete in this metric, then we say that  $(G, u)$  is *norm-complete*.

It is tempting to expect norm-complete partially ordered abelian groups with order-unit, particularly those that are interpolation groups, to be archimedean, but this is not the case in general. For instance, make the group  $G = \mathbf{R}^2$  into a partially ordered abelian group with positive cone

$$G^+ = \{(0, 0)\} \cup \{(a, b) \in G \mid a > 0 \text{ and } b > 0\}.$$

It is clear that  $G$  is an interpolation group, and that the element  $u = (1, 1)$  is an order-unit in  $G$ . Observing that

$$\|(a, b)\|_u = \max\{|a|, |b|\}$$

for all  $(a, b) \in G$ , we see that  $(G, u)$  is norm-complete. However,  $n(-1, 0) \leq u$  for all  $n \in \mathbf{N}$  while  $(-1, 0) \not\leq 0$ , so that  $G$  is not archimedean.

**LEMMA 2.9.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $v \in K_0(R)$ , and let  $n \in \mathbf{N}$ . If  $\|v\| < 1/2^n$ , then there exist  $x, y \in R$  such that  $v = [xR] - [yR]$ , while also  $N^*(x) \leq 1/2^n$  and  $N^*(y) \leq 1/2^n$ .*

**PROOF.** Write  $v = [A] - [B]$  for some finitely generated projective right  $R$ -modules  $A$  and  $B$ . Since  $\|v\| < 1/2^n$ , there exist  $s, t \in \mathbf{N}$  such that  $-s[R] \leq tv \leq s[R]$  and  $s/t < 1/2^n$ . Since  $R$  is unit-regular (Theorem 2.3), it follows that

$$tA \lesssim sR_R \oplus tB \quad \text{and} \quad tB \lesssim sR_R \oplus tA.$$

Now  $2^n tA \lesssim 2^n tB \oplus tR_R$ , because  $2^n s < t$ . Then  $2^n A \lesssim 2^n B \oplus R_R$  by Theorem 2.6, and similarly  $2^n B \lesssim 2^n A \oplus R_R$ .

In view of Lemma 2.7, there exist decompositions  $A = A_1 \oplus A_2$  and  $B = B_1 \oplus B_2$  such that  $A_1 \cong B_1$  and  $2^n A_2 \lesssim R_R$ . Then

$$2^n B_1 \oplus 2^n B_2 \cong 2^n B \lesssim 2^n A \oplus R_R \cong 2^n A_1 \oplus 2^n A_2 \oplus R_R \cong 2^n B_1 \oplus 2^n A_2 \oplus R_R,$$

whence  $2^n B_2 \lesssim 2^n A_2 \oplus R_R$ . Applying Lemma 2.7 a second time, we obtain decompositions  $B_2 = B_3 \oplus B_4$  and  $A_2 = A_3 \oplus A_4$  such that  $B_3 \cong A_3$  and  $2^n B_4 \lesssim R_R$ .

Now  $B_4 \lesssim 2^n B_4 \lesssim R_R$ , so  $B_4 \cong yR$  for some  $y \in R$ . Then as  $2^n(yR) \lesssim R_R$ , we obtain  $N^*(y) \leq 1/2^n$  from Lemma 1.2. Similarly, since  $2^n A_4 \lesssim 2^n A_2 \lesssim R_R$ , we have  $A_4 \cong xR$  for some  $x \in R$  satisfying  $N^*(x) \leq 1/2^n$ . Finally,

$$\begin{aligned} A &= A_1 \oplus A_2 = A_1 \oplus A_3 \oplus A_4 \cong A_1 \oplus A_3 \oplus xR, \\ B &= B_1 \oplus B_2 = B_1 \oplus B_3 \oplus B_4 \cong A_1 \oplus A_3 \oplus yR, \end{aligned}$$

hence we conclude that  $v = [A] - [B] = [xR] - [yR]$ .  $\square$

**LEMMA 2.10.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $x_1, x_2, \dots \in R$ , and assume that  $N^*(x_n) < 1/2^n$  for all  $n$ . Then there exists  $x \in R$  such that*

$$\|[x_1R] + \dots + [x_nR] - [xR]\| \rightarrow 0$$

as  $n \rightarrow \infty$ .

**PROOF.** We may clearly assume that  $R$  is nonzero.

For each  $n$ , Proposition 1.1 provides us with positive integers  $s_n$  and  $t_n$  such that  $s_n/t_n < 1/2^n$  and  $t_n(x_nR) \lesssim s_nR_R$ . Then since  $2^{n s_n} < t_n$ , we obtain  $2^{n s_n}(x_nR) \lesssim s_nR_R$ , and so  $2^n(x_nR) \lesssim R_R$ , using Theorem 2.6. Corollarily,

$$2^n(x_1R \oplus \dots \oplus x_nR) \lesssim 2^{n-1}R_R \oplus 2^{n-2}R_R \oplus \dots \oplus 2R_R \oplus R_R \lesssim 2^nR_R,$$

whence  $x_1R \oplus \dots \oplus x_nR \lesssim R_R$  (Theorem 2.6 again).

As this holds for all  $n$ , we obtain  $\bigoplus x_nR \lesssim R_R$ , by [2, Proposition 4.8]. Thus  $R$  contains an ascending sequence  $A_1 \leq A_2 \leq \dots$  of principal right ideals such that each

$$A_n \cong x_1R \oplus \dots \oplus x_nR.$$

By [2, Proposition 2.13], there exist orthogonal idempotents  $e_1, e_2, \dots$  in  $R$  such that

$$e_1R \oplus \dots \oplus e_nR = A_n$$

for all  $n$ . Note that  $e_nR \cong A_n/A_{n-1} \cong x_nR$ , when  $N^*(e_n) = N^*(x_n) < 1/2^n$  (Lemma 1.2).

Now the partial sums of the series  $\sum e_n$  are Cauchy with respect to  $N^*$ , hence there exists  $e \in R$  such that  $\sum e_n \rightarrow e$  in the  $N^*$ -metric. Note that  $e$  is an idempotent, and  $e_n e = e e_n = e_n$  for all  $n$ . For each  $n$ , we compute that

$$N^*(e - e_1 - \dots - e_n) = N^*\left(\sum_{k=n+1}^{\infty} e_k\right) \leq \sum_{k=n+1}^{\infty} N^*(e_k) < \sum_{k=n+1}^{\infty} 1/2^k = 1/2^n,$$

whence  $2^n((e - e_1 - \dots - e_n)R) \lesssim R_R$ , using Proposition 1.1 and Theorem 2.6 again. Inasmuch as

$$eR = e_1R \oplus \dots \oplus e_nR \oplus (e - e_1 - \dots - e_n)R,$$

it follows that  $2^n(eR) \lesssim 2^n(e_1R \oplus \dots \oplus e_nR) \oplus R_R$ . Consequently,

$$2^n([e_1R] + \dots + [e_nR]) \leq 2^n[eR] \leq 2^n([e_1R] + \dots + [e_nR]) + [R]$$

in  $K_0(R)$ . Thus

$$0 \leq 2^n([eR] - [e_1R] - \dots - [e_nR]) \leq [R],$$



from which we conclude that

$$\|[eR] - [e_1R] - \dots - [e_nR]\| \leq 1/2^n.$$

Since each  $[e_nR] = [x_nR]$ , this proves that

$$\|[x_1R] + \dots + [x_nR] - [eR]\| \rightarrow 0,$$

as desired.  $\square$

**THEOREM 2.11.** *If  $R$  is an  $N^*$ -complete regular ring, then  $(K_0(R), [R])$  is an archimedean norm-complete interpolation group with order-unit.*

**PROOF.** Since  $R$  is unit-regular (Theorem 2.3), it follows that  $K_0(R)$  is a partially ordered (rather than just pre-ordered) abelian group [2, Proposition 15.2], and that  $K_0(R)$  is an interpolation group [4, Proposition II.10.3]. Given  $x, y \in K_0(R)$  such that  $nx \leq y$  for all  $n \in \mathbb{N}$ , choose finitely generated projective right  $R$ -modules  $A, B, C, D$  such that  $x = [A] - [B]$  and  $y = [C] - [D]$ . Then

$$[nA] - [nB] = nx \leq y \leq [C]$$

and so  $nA \lesssim nB \oplus C$ , for all  $n \in \mathbb{N}$ . By Theorem 2.8,  $A \lesssim B$ , whence  $x \leq 0$ . Therefore  $K_0(R)$  is archimedean. In particular, it now follows from [4, Proposition I.6.2] that the pseudo-metric on  $K_0(R)$  induced by  $\|\cdot\|$  is actually a metric.

Finally, consider a Cauchy sequence  $\{v_1, v_2, \dots\}$  in  $K_0(R)$ . By passing to a subsequence, we may assume that  $\|v_{n+1} - v_n\| < 1/2^{n+1}$  for all  $n$ . Using Lemma 2.9, we obtain elements  $x_n, y_n \in R$  such that

$$v_{n+1} - v_n = [x_nR] - [y_nR],$$

while also  $N^*(x_n) \leq 1/2^{n+1}$  and  $N^*(y_n) \leq 1/2^{n+1}$ . According to Lemma 2.10, there exist elements  $x, y \in R$  such that

$$\begin{aligned} \|[x_1R] + \dots + [x_nR] - [xR]\| &\rightarrow 0, \\ \|[y_1R] + \dots + [y_nR] - [yR]\| &\rightarrow 0. \end{aligned}$$

Since  $v_{n+1} - v_1 = [x_1R] + \dots + [x_nR] - [y_1R] - \dots - [y_nR]$  for all  $n$ , we conclude that

$$\|(v_{n+1} - v_1) - ([xR] - [yR])\| \rightarrow 0,$$

and consequently  $v_{n+1} \rightarrow v_1 + [xR] - [yR]$ . Therefore  $(K_0(R), [R])$  is norm-complete.  $\square$

**III. Archimedean norm-complete interpolation groups.** Given an archimedean norm-complete interpolation group  $(G, u)$  with order-unit, we study the state space  $S(G, u)$ , and the relationship between  $G$  and the space of affine continuous real-valued functions on  $S(G, u)$ . In particular, we investigate extreme points and closed faces of  $S(G, u)$ , and relate them to ideals of  $G$ . These results will be applied, via  $K_0$ , to the ideal theory of  $N^*$ -complete regular rings.

**DEFINITION.** Let  $(G, u)$  be a partially ordered abelian group with order-unit. A *state* on  $(G, u)$  is any positive homomorphism  $s: G \rightarrow \mathbb{R}$  such that  $s(u) = 1$ . The *state space* of  $(G, u)$ , denoted  $S(G, u)$ , is the set of all states on  $(G, u)$ . The state space is

regarded as a subset of the linear topological space  $\mathbf{R}^G$  (which is given the product topology), and as such is a compact convex set [2, Proposition 17.11]. If  $G$  is an interpolation group, then  $S(G, u)$  is a Choquet simplex [4, Theorem I.2.5].

DEFINITION. The *extreme boundary* of a convex set  $S$ , denoted  $\partial_e S$ , is the set of all *extreme points* of  $S$ , that is, points  $s \in S$  such that the only convex combinations  $s = \alpha s' + (1 - \alpha)s''$  with  $0 \leq \alpha \leq 1$  and  $s', s'' \in S$  are those for which  $\alpha = 0$ , or  $\alpha = 1$ , or  $s' = s'' = s$ . Now suppose that  $S = S(G, u)$  for some partially ordered abelian group  $(G, u)$  with order-unit. An extreme state  $s$  in  $\partial_e S$  is said to be *discrete* if  $s(G)$  is a cyclic subgroup of  $\mathbf{R}$ . Note that if  $s$  is discrete, then  $s(G) = (1/m)\mathbf{Z}$  for some  $m \in \mathbf{N}$ , because  $1 = s(u) \in s(G)$ . On the other hand, if  $s$  is not discrete, then  $s(G)$  is dense in  $\mathbf{R}$ .

DEFINITION. We use  $\text{Aff}(S)$  to denote the partially ordered real Banach space of all affine continuous real-valued functions on  $S$  (with the pointwise ordering and the supremum norm). Evaluation at elements of  $G$  provides a map  $\varphi: G \rightarrow \text{Aff}(S)$ , so that  $\varphi(x)(s) = s(x)$  for all  $x \in G$  and  $s \in S$ . Note that  $\varphi$  is a positive homomorphism, and that  $\varphi(u)$  is the constant function 1. We refer to  $\varphi$  as *the natural map from  $G$  to  $\text{Aff}(S)$* . The map  $\varphi$  is also norm-preserving; namely,

$$\|\varphi(x)\| = \sup\{|s(x)| : s \in S\} = \|x\|$$

for all  $x \in G$  [4, Lemma I.6.1].

THEOREM 3.1. *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, and set  $S = S(G, u)$ . For all discrete  $s \in \partial_e S$ , set  $A_s = s(G)$ ; for all other  $s \in \partial_e S$ , set  $A_s = \mathbf{R}$ . Set*

$$A = \{p \in \text{Aff}(S) \mid p(s) \in A_s \text{ for all } s \in \partial_e S\}.$$

*Then the natural map from  $G$  to  $\text{Aff}(S)$  provides an isomorphism of  $(G, u)$  onto  $(A, 1)$  (as partially ordered abelian groups with order-unit).*

PROOF. [3, Theorem 5.1].  $\square$

COROLLARY 3.2. *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit. Then  $G$  is lattice-ordered if and only if  $\partial_e S(G, u)$  is compact.*

PROOF. [3, Corollary 5.4].  $\square$

In order to apply Theorem 3.1 effectively, we must be able to identify  $\partial_e S$  easily. Thus we develop criteria for deciding when states are extreme. For topological considerations, we also develop similar results for compact sets of extreme states, and for closed faces of  $S$ . We begin with a result relating the archimedean property to norm properties.

PROPOSITION 3.3. *Let  $(G, u)$  be an interpolation group with order-unit. Then  $G$  is archimedean if and only if  $G$  is 2-unperforated and  $G^+$  is norm-closed in  $G$ .*

PROOF. If  $G$  is archimedean, then  $G$  is unperforated by [4, Lemma I.5.2]. Now consider elements  $x_1, x_2, \dots$  in  $G^+$  and  $x \in G$  such that  $x_n \rightarrow x$  in norm. We may assume that  $\|x_n - x\| < 1/n$  for all  $n$ . According to [4, Proposition I.6.2],  $n(x_n - x) \leq u$ , and consequently  $n(-x) \leq u$ . Since this holds for all  $n \in \mathbf{N}$ , the archimedean property implies that  $-x \leq 0$ , so that  $x \in G^+$ . Thus  $G^+$  is norm-closed in  $G$ .

Conversely, assume that  $G$  is 2-unperforated and that  $G^+$  is norm-closed in  $G$ . Given  $a, b \in G$  such that  $na \leq b$  for all  $n \in \mathbb{N}$ , we must show that  $a \leq 0$ . Write  $a = x - y$  for some  $x, y \in G^+$ , and choose  $z \in G^+$  with  $b \leq z$ . Then  $nx \leq ny + z$  for all  $n \in \mathbb{N}$ , and we must show that  $x \leq y$ .

For all  $n \in \mathbb{N}$ , we have  $2^n x \leq 2^n y + z$ , hence [4, Lemma I.5.7] says that  $x = v_n + w_n$  for some  $v_n, w_n \in G^+$  such that  $v_n \leq y$  and  $2^n w_n \leq z$ . Then  $\|w_n\| \leq \|z\|/2^n$ , so that  $w_n \rightarrow 0$ , and consequently  $v_n \rightarrow x$ . Thus  $y - v_n \rightarrow y - x$ . Since each  $y - v_n$  is in  $G^+$ , we conclude that  $y - x$  is in  $G^+$ , as desired. Therefore  $G$  is archimedean.  $\square$

In particular, if  $(G, u)$  is an archimedean interpolation group with order-unit, and we have norm-convergent sequences  $x_n \rightarrow x$  and  $y_n \rightarrow y$  in  $G$  with  $x_n \leq y_n$  for all  $n$ , then  $x \leq y$ .

**THEOREM 3.4.** *Let  $(G, u)$  be an interpolation group with order-unit, and let  $s \in S(G, u)$ . Then  $s$  is an extreme point of  $S(G, u)$  if and only if*

$$\min\{s(x), s(y)\} = \sup\{s(z) \mid z \in G^+; z \leq x; z \leq y\}$$

for all  $x, y \in G^+$ .

**PROOF.** [3, Theorem 3.1].  $\square$

**COROLLARY 3.5.** *Let  $(G, u)$  be an interpolation group with order-unit, and let  $X$  be a compact subset of  $\partial_e S(G, u)$ . Given  $x, y \in G^+$  and a positive real number  $\epsilon$ , there exists  $z \in G^+$  such that  $z \leq x$  and  $z \leq y$ , while also*

$$s(z) > \min\{s(x), s(y)\} - \epsilon$$

for all  $s \in X$ .

**PROOF.** Set  $W = \{w \in G^+ \mid w \leq x \text{ and } w \leq y\}$ , and note, because of the interpolation property, that  $W$  is upward directed. For each  $w \in W$ , set

$$V(w) = \{s \in S(G, u) \mid s(w) > \min\{s(x), s(y)\} - \epsilon\},$$

which is an open subset of  $S(G, u)$ . In view of Theorem 3.4, we see that these  $V(w)$ 's cover  $\partial_e S(G, u)$ , and so cover  $X$ . As  $X$  is compact, it follows that

$$X \subseteq V(w_1) \cup \dots \cup V(w_n)$$

for some  $w_1, \dots, w_n \in W$ . Since  $W$  is upward directed, there exists  $z \in W$  such that all  $w_i \leq z$ , and  $z$  has the desired properties.  $\square$

**COROLLARY 3.6.** *Let  $(G, u)$  be an interpolation group with order-unit, let  $a, b \in G^+$ , and let  $m \in \mathbb{N}$ . Let  $X$  be a compact subset of  $\partial_e S(G, u)$ , and assume that  $s(ma) \leq s(b)$  for all  $s \in X$ . Given any positive real number  $\epsilon$ , there exists  $c \in G^+$  such that  $c \leq a$  and  $mc \leq b$ , while also  $s(c) > s(a) - \epsilon$  for all  $s \in X$ .*

**PROOF.** By Corollary 3.5, there exists  $x \in G^+$  such that  $x \leq ma$  and  $x \leq b$ , while also  $s(x) > s(ma) - \epsilon$  for all  $s \in X$ . Then, using Riesz decomposition,  $x = x_1 + \dots + x_m$  for some  $x_i \in G^+$  satisfying  $x_i \leq a$ . For each  $i$ , note that  $x - x_i \leq (m - 1)a$ , whence  $x - (m - 1)a \leq x_i$ . Since  $0 \leq x_i$  for each  $i$  as well, interpolation provides an element  $c \in G$  such that

$$x - (m - 1)a \leq c \leq x_i \quad \text{and} \quad 0 \leq c \leq x_i$$

for all  $i$ . Thus  $c \in G^+$  and  $c \leq x_1 \leq a$ , while

$$mc \leq x_1 + \dots + x_m = x \leq b.$$

As  $x - (m - 1)a \leq c$ , we have  $a - c \leq ma - x$ , whence

$$s(a) - s(c) \leq s(ma) - s(x) < \varepsilon$$

for all  $s \in X$ . Therefore  $s(c) > s(a) - \varepsilon$  for all  $s \in X$ .  $\square$

**THEOREM 3.7.** *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, and let  $X$  be a compact subset of  $\partial_e S(G, u)$ . Given any  $x, y \in G$ , there exist  $z, w \in G$  such that  $z \leq x \leq w$  and  $z \leq y \leq w$ , while also*

$$s(z) = \min\{s(x), s(y)\} \quad \text{and} \quad s(w) = \max\{s(x), s(y)\}$$

for all  $s \in X$ . If  $x, y \in G^+$ , then such  $z, w$  can be found in  $G^+$ .

**PROOF.** First assume that  $x, y \in G^+$ . The rule  $p(s) = \min\{s(x), s(y)\}$  defines a continuous map  $p$  of  $S(G, u)$  into  $\mathbf{R}^+$ . We construct elements  $z_1 \leq z_2 \leq \dots$  in  $G^+$  such that each  $z_n \leq x$  and  $z_n \leq y$ , while also

$$s(z_n) > p(s) - (1/2^n) \quad \text{and} \quad \|z_{n+1} - z_n\| \leq 1/2^n$$

for all  $n$  and all  $s \in X$ . To begin, we obtain  $z_1$  directly from Corollary 3.5.

Now assume that  $z_1, \dots, z_n$  have been constructed, for some  $n$ . According to Corollary 3.5, there exists  $a \in G^+$  such that  $a \leq x$  and  $a \leq y$ , while also

$$s(a) > p(s) - (1/2^{n+2})$$

for all  $s \in X$ . Since  $z_n \leq x$  and  $z_n \leq y$  as well, there is some  $b \in G^+$  satisfying  $a \leq b \leq x$  and  $z_n \leq b \leq y$ . Note that

$$s(b) \geq s(a) > p(s) - (1/2^{n+2})$$

for all  $s \in X$ .

The element  $b - z_n$  lies in  $G^+$ , and for all  $s \in X$  we have

$$s(z_n) > \min\{s(x), s(y)\} - (1/2^n) \geq s(b) - (1/2^n),$$

whence  $s(2^n(b - z_n)) < 1 = s(u)$ . By Corollary 3.6, there exists  $c \in G^+$  such that  $c \leq b - z_n$  and  $2^n c \leq u$ , while also

$$s(c) > s(b - z_n) - (1/2^{n+2})$$

for all  $s \in X$ . Set  $z_{n+1} = z_n + c$ , noting that  $z_n \leq z_{n+1} \leq b \leq x$  and  $z_{n+1} \leq b \leq y$ . For all  $s \in X$ , we have

$$s(z_{n+1}) = s(z_n) + s(c) > s(b) - (1/2^{n+2}) > p(s) - (1/2^{n+1}).$$

Since  $0 \leq 2^n(z_{n+1} - z_n) = 2^n c \leq u$ , we also have  $\|z_{n+1} - z_n\| \leq 1/2^n$ , which completes the induction step.

Having constructed a Cauchy sequence  $\{z_1, z_2, \dots\}$  in  $G$ , we must have  $z_n \rightarrow z$  for some  $z \in G$ . Since  $0 \leq z_n \leq x$  and  $0 \leq z_n \leq y$  for all  $n$ , we obtain  $0 \leq z \leq x$  and  $0 \leq z \leq y$ . On the other hand,  $z_k \geq z_n$  whenever  $k \geq n$ , hence  $z \geq z_n$  for all  $n$ . Consequently, for any  $s \in X$ ,

$$\min\{s(x), s(y)\} \geq s(z) \geq s(z_n) > \min\{s(x), s(y)\} - (1/2^n),$$

and thus  $s(z) = \min\{s(x), s(y)\}$ .

Now consider arbitrary elements  $x, y \in G$ , and choose  $a \in G^+$  such that  $x + a \geq 0$  and  $y + a \geq 0$ . By the above, there exists  $b \in G^+$  such that  $b \leq x + a$  and  $b \leq y + a$ , while also

$$s(b) = \min\{s(x + a), s(y + a)\}$$

for all  $s \in X$ . Setting  $z = b - a$ , we obtain  $z \leq x$  and  $z \leq y$ , while

$$s(z) = \min\{s(x), s(y)\}$$

for all  $s \in X$ .

Finally, use the result above to obtain  $c \in G$  such that  $c \leq -x$  and  $c \leq -y$ , while also

$$s(c) = \min\{s(-x), s(-y)\}$$

for all  $s \in X$ . Set  $w = -c$ .  $\square$

**COROLLARY 3.8.** *If  $(G, u)$  is an archimedean norm-complete interpolation group with order-unit, and  $X$  is a compact subset of  $\partial_e S(G, u)$ , then  $\ker(X)$  is an ideal of  $G$ , and  $G/\ker(X)$  is a lattice-ordered abelian group.*

**PROOF.** Set  $K = \ker(X)$ . Clearly  $K$  is a convex subgroup of  $G$ . Given  $x \in K$ , we have  $s(x) = s(0)$  for all  $s \in X$ . According to Theorem 3.7, there exists  $z \in G$  such that  $z \leq x$  and  $z \leq 0$ , while also  $s(z) = s(x) = s(0)$  for all  $s \in X$ . Thus  $x - z$  and  $-z$  are elements of  $G^+ \cap K$  satisfying  $(x - z) - (-z) = x$ , which proves that  $K$  is a directed subgroup of  $G$ . Therefore  $K$  is an ideal of  $G$ .

Now consider any elements  $x + K$  and  $y + K$  in  $G/K$ . According to Theorem 3.7, there exists  $z \in G$  such that  $z \leq x$  and  $z \leq y$ , while also  $s(z) = \min\{s(x), s(y)\}$  for all  $s \in X$ . Then  $z + K \leq x + K$  and  $z + K \leq y + K$ , and we claim that  $z + K$  is the infimum of  $x + K$  and  $y + K$  in  $G/K$ .

Given  $a + K$  in  $G/K$  satisfying  $a + K \leq x + K$  and  $a + K \leq y + K$ , we have  $a \leq x + k'$  and  $a \leq y + k''$  for some  $k', k'' \in K$ . Since  $K$  is directed, we may choose  $k \in K$  such that  $k' \leq k$  and  $k'' \leq k$ , whence  $a \leq x + k$  and  $a \leq y + k$ . Now

$$a - k \leq x; \quad a - k \leq y; \quad z \leq x; \quad z \leq y.$$

Interpolating, we obtain  $b \in G$  such that  $a - k \leq b \leq x$  and  $z \leq b \leq y$ . In particular,

$$s(z) \leq s(b) \leq \min\{s(x), s(y)\} = s(z)$$

for all  $s \in X$ , whence  $b - z \in K$ . Thus

$$a + K = (a - k) + K \leq b + K = z + K,$$

proving that  $z + K$  is indeed the infimum of  $x + K$  and  $y + K$ .

Therefore  $G/K$  is lattice-ordered.  $\square$

We thank the referee for pointing out that the hypothesis of norm-completeness in Theorem 3.7 and Corollary 3.8 is essential, as the following example shows. (This is a simplified version of the referee's example.)

**EXAMPLE 3.9.** There exists an archimedean interpolation group  $(G, u)$  with order-unit possessing an extreme state  $s$  such that  $\ker(s) = \{0\}$  but  $G$  is not lattice-ordered.

PROOF. Let  $G$  be the  $\mathbf{Q}$ -subspace of  $\mathbf{R}^2$  spanned by the vectors  $u = (1, 1)$  and  $v = (\pi, -\pi)$ . Note that since  $u$  and  $v$  are  $\mathbf{R}$ -linearly independent,  $G$  is dense in  $\mathbf{R}^2$  in the usual Euclidean topology. Give  $\mathbf{R}^2$  the direct product ordering, and give  $G$  the relative ordering inherited from  $\mathbf{R}^2$ , so that  $G^+ = G \cap (\mathbf{R}^2)^+$ . Then  $\mathbf{R}^2$  and  $G$  are archimedean partially ordered abelian groups, and  $u$  is an order-unit in each of them.

To check interpolation, consider elements  $x_1, x_2, y_1, y_2$  in  $G$  satisfying  $x_i \leq y_j$  for all  $i, j$ . Then each

$$x_i = (a_i + b_i\pi, a_i - b_i\pi) \quad \text{and} \quad y_j = (c_j + d_j\pi, c_j - d_j\pi)$$

for suitable  $a_i, b_i, c_j, d_j \in \mathbf{Q}$ . First assume that  $a_i = c_j$  for some  $i, j$ , say  $a_1 = c_1$ . Since

$$a_1 + b_1\pi \leq c_1 + d_1\pi \quad \text{and} \quad a_1 - b_1\pi \leq c_1 - d_1\pi,$$

it follows that  $b_1 = d_1$ , whence  $x_1 = y_1$ . In this case,  $x_i \leq x_1 \leq y_j$  for all  $i, j$ . Now assume that  $a_i \neq c_j$  for all  $i, j$ . Then

$$a_i + b_i\pi \neq c_j + d_j\pi \quad \text{and} \quad a_i - b_i\pi \neq c_j - d_j\pi$$

for all  $i, j$ . Consequently, the set  $W$  consisting of those  $(\alpha, \beta)$  in  $\mathbf{R}^2$  satisfying

$$a_i + b_i\pi < \alpha < c_j + d_j\pi \quad \text{and} \quad a_i - b_i\pi < \beta < c_j - d_j\pi$$

for all  $i, j$  is a nonempty open subset of  $\mathbf{R}^2$ . Since  $G$  is dense in  $\mathbf{R}^2$ , there exists an element  $z$  in  $G \cap W$ , and  $x_i \leq z \leq y_j$  for all  $i, j$ . Therefore  $G$  is an interpolation group.

Define  $t: \mathbf{R}^2 \rightarrow \mathbf{R}$  by the rule  $t(\alpha, \beta) = \alpha$ , and note that  $t$  is an extreme point of  $S(\mathbf{R}^2, u)$ . Let  $s$  denote the restriction of  $t$  to  $G$ . As  $G^+ = G \cap (\mathbf{R}^2)^+$ , all states in  $S(G, u)$  extend to states in  $S(\mathbf{R}^2, u)$ , by [2, Proposition 18.1]. Thus the restriction map  $S(\mathbf{R}^2, u) \rightarrow S(G, u)$  is an affine homeomorphism, from which we see that  $s$  is an extreme point of  $S(G, u)$ . It is clear that  $\ker(s) = \{0\}$ .

Consider any  $x \in G$  satisfying  $x \leq u$  and  $x \leq v$ . Then  $x = (a + b\pi, a - b\pi)$  for some  $a, b \in \mathbf{Q}$ , and from the relations  $x \leq u$  and  $x \leq v$  we obtain  $a \leq \alpha$ , where

$$\alpha = \min\{1 - b\pi, 1 + b\pi, (1 - b)\pi, (b - 1)\pi\}.$$

Note that  $\alpha$  must be irrational. Thus  $a < \alpha$ , and we may choose  $c \in \mathbf{Q}$  such that  $a < c < \alpha$ . Setting  $y = (c + b\pi, c - b\pi)$ , we conclude that  $x < y$  while also  $y \leq u$  and  $y \leq v$ . Therefore the set  $\{u, v\}$  has no infimum in  $G$ , proving that  $G$  is not lattice-ordered.  $\square$

In case the set  $X$  in Corollary 3.8 is a singleton, we can precisely identify the quotient partially ordered abelian group  $G/\ker(X)$ , as follows.

**THEOREM 3.10.** *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, and let  $s \in \partial_e S(G, u)$ . Then  $\ker(s)$  is an ideal of  $G$ , and  $s$  induces an isomorphism of  $G/\ker(s)$  onto  $s(G)$  as partially ordered abelian groups. Moreover, either  $s(G) = \mathbf{R}$  or  $s(G) = (1/m)\mathbf{Z}$  for some  $m \in \mathbf{N}$ .*

PROOF. The subgroup  $K = \ker(s)$  is an ideal of  $G$  by Corollary 3.8. Obviously the induced map  $\bar{s}: G/K \rightarrow s(G)$  is a group isomorphism, and a positive map as well. Given an element  $x + K$  in  $G/K$  with  $\bar{s}(x + K) \geq 0$ , we have  $s(x) \geq 0 = s(0)$ . Then

Theorem 3.7 provides us with an element  $z \in G$  such that  $z \leq x$  and  $z \leq 0$ , while also  $s(z) = 0$ . Now  $z \in K$  and  $x - z \geq 0$ , hence

$$x + K = (x - z) + K \geq 0$$

as well. Thus for all  $a \in G/K$ , we have  $a \geq 0$  if and only if  $\bar{s}(a) \geq 0$ . Therefore  $\bar{s}$  is an isomorphism of partially ordered abelian groups.

If  $s$  is discrete, then  $s(G) = (1/m)\mathbf{Z}$  for some  $m \in \mathbf{N}$ . Now assume that  $s$  is not discrete, so that  $s(G)$  is dense in  $\mathbf{R}$ .

Given  $\alpha \in \mathbf{R}$ , we construct elements  $x_1, x_2, \dots$  in  $G$  such that

$$\alpha - (1/2^n) < s(x_n) < \alpha \quad \text{and} \quad \|x_{n+1} - x_n\| \leq 1/2^n$$

for all  $n$ . To begin, we obtain  $x_1$  from the density of  $s(G)$  in  $\mathbf{R}$ .

Now assume that  $x_1, \dots, x_n$  have been constructed, for some  $n$ . Choose an element  $a \in G$  such that

$$\alpha - (1/2^{n+2}) < s(a) < \alpha.$$

By Theorem 3.7, there exists  $b \in G$  such that  $b \geq x_n$  and  $b \geq a$ , while also  $s(b) = \max\{s(x_n), s(a)\}$ . Note that

$$\alpha - (1/2^{n+2}) < s(b) < \alpha.$$

Since  $s(b) < \alpha$  and  $s(x_n) > \alpha - (1/2^n)$ , we find that

$$s(2^n b) < 2^n \alpha < s(2^n x_n) + 1,$$

whence  $s(2^n(b - x_n)) < 1 = s(u)$ . As  $b - x_n \in G^+$ , Corollary 3.6 provides us with an element  $c \in G^+$  such that  $c \leq b - x_n$  and  $2^n c \leq u$ , while also

$$s(c) > s(b - x_n) - (1/2^{n+2}).$$

Set  $x_{n+1} = x_n + c$ , so that  $x_n \leq x_{n+1} \leq b$ . Thus  $s(x_{n+1}) \leq s(b) < \alpha$ , and

$$s(x_{n+1}) = s(x_n) + s(c) > s(b) - (1/2^{n+2}) > \alpha - (1/2^{n+1}).$$

In addition,  $\|x_{n+1} - x_n\| = \|c\| \leq 1/2^n$ , which completes the induction step.

Now there exists  $x \in G$  such that  $x_n \rightarrow x$  in norm. Since

$$|s(x_n) - s(x)| = |s(x_n - x)| \leq \|x_n - x\|$$

for all  $n$  [4, Lemma I.6.1], it follows that  $s(x_n) \rightarrow s(x)$ , and thus  $s(x) = \alpha$ . Therefore  $s(G) = \mathbf{R}$ , as desired.  $\square$

**DEFINITION.** A *face* of a convex set  $S$  is a convex subset  $F \subseteq S$  (possibly empty) such that whenever  $s = \alpha s' + (1 - \alpha)s''$  is a positive convex combination with  $s \in F$  and  $s', s'' \in S$ , then  $s', s'' \in F$ .

**LEMMA 3.11.** Let  $(G, u)$  be an interpolation group with order-unit, and let  $X$  be either a compact subset of  $\partial_e S(G, u)$  or a closed face of  $S(G, u)$ . Let  $t \in \partial_e S(G, u)$  such that  $t \notin X$ . Then there exists  $x \in G^+$  such that  $t(x) > 1$  but  $s(x) < 1$  for all  $s \in X$ .

**PROOF.** Set  $A = \{a \in G^+ \mid t(a) > 1\}$ , and note that  $2u \in A$ . Also,  $A$  is downward directed, by Theorem 3.4. For all  $a \in A$ , set

$$W(a) = \{s \in S(G, u) \mid s(a) < 1\},$$

which is an open subset of  $S(G, u)$ . We claim that these  $W(a)$ 's cover  $X$ .

Thus consider any  $s \in X$ . Since  $\{t\}$  is a face of  $S(G, u)$ , and either  $\{s\}$  or  $X$  is a face of  $S(G, u)$ , we see that  $s$  and  $t$  lie in disjoint faces of  $S(G, u)$ . By [3, Lemma 2.8],  $2u = a + b$  for some  $a, b \in G^+$  such that  $s(a) + t(b) < 1$ . Then  $t(a) = 2 - t(b) > 1$ , whence  $a \in A$  and  $s \in W(a)$ . Therefore the  $W(a)$ 's do cover  $X$ , as claimed.

By compactness,  $X \subseteq W(a_1) \cup \dots \cup W(a_n)$  for some elements  $a_i \in A$ . As  $A$  is downward directed, there exists  $x \in A$  such that each  $a_i \geq x$ , and  $x$  has the desired properties.  $\square$

**THEOREM 3.12.** *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, and let  $X$  be a compact subset of  $\partial_e S(G, u)$ . Then*

$$X = \{s \in \partial_e S(G, u) \mid G^+ \cap \ker(X) \subseteq \ker(s)\}.$$

**PROOF.** Consider  $t \in \partial_e S(G, u)$  such that  $t \notin X$ . By Lemma 3.11, there exists  $x \in G^+$  such that  $t(x) > 1$  but  $s(x) < 1$  for all  $s \in X$ . Applying Theorem 3.7 to the elements  $x, u \in G^+$  and the compact subset  $X \cup \{t\}$  of  $\partial_e S(G, u)$ , we obtain  $y \in G^+$  such that  $y \geq x$  and  $y \geq u$ , while  $t(y) > 1$  and  $s(y) = 1$  for all  $s \in X$ . Consequently,  $y - u$  is an element of  $G^+ \cap \ker(X)$  which does not lie in  $\ker(t)$ .  $\square$

We now turn to closed faces of state spaces. The results above, concerning compact sets of extreme states, carry over fairly directly, with similar proofs.

**THEOREM 3.13.** *Let  $(G, u)$  be an interpolation group with order-unit, let  $x, y \in G^+$ , and let  $F$  be a closed face of  $S(G, u)$ . Assume that  $s(x) \leq s(y)$  for all  $s \in F$ . Given any positive real number  $\epsilon$ , there exists  $z \in G^+$  such that  $z \leq x$  and  $z \leq y$ , while also  $s(z) > s(x) - \epsilon$  for all  $s \in F$ .*

**PROOF.** [3, Theorem 3.4].  $\square$

**COROLLARY 3.14.** *Let  $(G, u)$  be an interpolation group with order-unit, and let  $F$  be a closed face of  $S(G, u)$ . Let  $a, b \in G^+$  and  $m \in \mathbb{N}$ , and assume that  $s(ma) \leq s(b)$  for all  $s \in F$ . Given any positive real number  $\epsilon$ , there exists  $c \in G^+$  such that  $c \leq a$  and  $mc \leq b$ , while also  $s(c) > s(a) - \epsilon$  for all  $s \in F$ .*

**PROOF.** As Corollary 3.6, using Theorem 3.13 in place of Corollary 3.5.  $\square$

**THEOREM 3.15.** *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, let  $x, y \in G$ , and let  $F$  be a closed face of  $S(G, u)$ . If  $s(x) \leq s(y)$  for all  $s \in F$ , then there exist  $z, w \in G$  such that  $z \leq x \leq w$  and  $z \leq y \leq w$ , while also  $s(z) = s(x)$  and  $s(w) = s(y)$  for all  $s \in F$ . If  $x, y \in G^+$ , then such  $z, w$  can be found in  $G^+$ .*

**PROOF.** As Theorem 3.7, using Theorem 3.13 and Corollary 3.14 in place of Corollaries 3.5 and 3.6.  $\square$

**COROLLARY 3.16.** *If  $(G, u)$  is an archimedean norm-complete interpolation group with order-unit, and  $F$  is a closed face of  $S(G, u)$ , then  $\ker(F)$  is an ideal of  $G$ .*

**PROOF.** As Corollary 3.8, using Theorem 3.15 in place of Theorem 3.7.  $\square$



**THEOREM 3.17.** *Let  $(G, u)$  be an archimedean norm-complete interpolation group with order-unit, and let  $F$  be a closed face of  $S(G, u)$ . Then*

$$F = \{s \in S(G, u) \mid G^+ \cap \ker(F) \subseteq \ker(s)\}.$$

**PROOF.** Set  $F' = \{s \in S(G, u) \mid G^+ \cap \ker(F) \subseteq \ker(s)\}$ . Clearly  $F'$  is a closed convex subset of  $S(G, u)$ , and we claim that  $F'$  is a face of  $S(G, u)$  as well. Thus consider any positive convex combination  $s = \alpha s' + (1 - \alpha)s''$  in  $S(G, u)$  with  $s \in F'$ . For any  $x \in G^+ \cap \ker(F)$ , we have

$$\alpha s'(x) + (1 - \alpha)s''(x) = 0; \quad s'(x) \geq 0; \quad s''(x) \geq 0$$

and so  $s'(x) = s''(x) = 0$ . Therefore  $s', s'' \in F'$ , proving that  $F'$  is indeed a face of  $S(G, u)$ .

Being a compact convex set,  $F'$  equals the closure of the convex hull of its extreme boundary  $\partial_e F'$ , by the Krein-Milman Theorem. Thus if  $F' \not\subseteq F$ , there must be some  $t \in \partial_e F'$  which does not lie in  $F$ . As  $F'$  is a face of  $S(G, u)$ , we see that actually  $t$  is an extreme point of  $S(G, u)$ .

By Lemma 3.11, there exists  $x \in G^+$  such that  $t(x) > 1$  but  $s(x) < 1$  for all  $s \in F$ . According to Theorem 3.15, there exists  $w \in G^+$  such that  $x \leq w$  and  $u \leq w$ , while also  $s(w) = 1$  for all  $s \in F$ . Thus  $w - u$  is an element of  $G^+ \cap \ker(F)$ . On the other hand,  $t(w) \geq t(x) > 1$  and so  $w - u$  is not in  $\ker(t)$ , which contradicts the fact that  $t \in F'$ .

Therefore  $F' \subseteq F$ . The reverse inclusion is automatic, hence  $F = F'$ , as desired.

□

**IV.  $N^*$ -complete regular rings.** We apply the results of §III, via  $K_0$ , to the structure of  $N^*$ -complete regular rings  $R$ . The thrust of most of these results is that  $R$  behaves much like a ring of sections of a sheaf of simple self-injective rings. Namely, for every maximal two-sided ideal  $M$  of  $R$ , the factor ring  $R/M$  is right and left self-injective, and many properties of  $R$  and its projective modules are determined by what happens modulo these maximal two-sided ideals. For instance,  $R$  is an  $n \times n$  matrix ring (for some fixed  $n \in \mathbf{N}$ ) if and only if each  $R/M$  is an  $n \times n$  matrix ring. For another example, given finitely generated projective right  $R$ -modules  $A$  and  $B$ , we have  $A \cong B$  if and only if  $A/AM \cong B/BM$  for all  $M$ . Many of our results generalize parallel results for  $\aleph_0$ -continuous regular rings in [4, 7, 8]. At the end of the section, we indicate how our results relate to these papers.

All the results of this section depend on Theorems 2.3 and 2.11: That any  $N^*$ -complete regular ring  $R$  is unit-regular, and that for such rings  $(K_0(R), [R])$  is an archimedean norm-complete interpolation group with order-unit. We shall use these results repeatedly without further reference to them. One other basic result is needed, to identify the state space of  $(K_0(R), [R])$ .

**PROPOSITION 4.1.** *For any regular ring  $R$ , there is a natural affine homeomorphism  $\theta: S(K_0(R), [R]) \rightarrow \mathbf{P}(R)$  such that  $\theta(s)(x) = s([xR])$  for all  $s$  in  $S(K_0(R), [R])$  and all  $x \in R$ .*

**PROOF.** [2, Proposition 17.12]. □

**THEOREM 4.2.** *Let  $R$  be an  $N^*$ -complete regular ring.*

(a) *If  $X$  is a compact subset of  $\partial_e \mathbf{P}(R)$ , then*

$$X = \{P \in \partial_e \mathbf{P}(R) \mid \ker(X) \subseteq \ker(P)\}.$$

(b) *If  $F$  is a closed face of  $\mathbf{P}(R)$ , then*

$$F = \{P \in \mathbf{P}(R) \mid \ker(F) \subseteq \ker(P)\}.$$

**PROOF.** Set  $S = S(K_0(R), [R])$ , and let  $\theta: S \rightarrow \mathbf{P}(R)$  be the affine homeomorphism given in Proposition 4.1.

(a) Set  $Y = \theta^{-1}(X)$ , which is a compact subset of  $\partial_e S$ . Let  $Q \in \partial_e \mathbf{P}(R)$  such that  $\ker(X) \subseteq \ker(Q)$ , and set  $t = \theta^{-1}(Q)$ . We shall show that  $K_0(R)^+ \cap \ker(Y)$  is contained in  $\ker(t)$ .

Given  $a \in K_0(R)^+ \cap \ker(Y)$ , we have  $a = [A]$  for some finitely generated projective right  $R$ -module  $A$ . Then

$$A \cong x_1 R \oplus \cdots \oplus x_n R \quad \text{and} \quad [A] = [x_1 R] + \cdots + [x_n R]$$

for some elements  $x_i \in R$ . For each  $i$ , we have  $0 \leq [x_i R] \leq [A]$ , whence  $s([x_i R]) = 0$  for all  $s \in Y$ , and so  $P(x_i) = 0$  for all  $P \in X$ . Then each  $x_i$  lies in  $\ker(X)$ , hence  $x_i \in \ker(Q)$ , and so  $t([x_i R]) = 0$ . Consequently,

$$t(a) = t([x_1 R]) + \cdots + t([x_n R]) = 0.$$

Therefore  $K_0(R)^+ \cap \ker(Y) \subseteq \ker(t)$ , as claimed.

Now Theorem 3.12 shows that  $t \in Y$ , and therefore  $Q \in X$ .

(b) As (a), using Theorem 3.17 in place of Theorem 3.12.  $\square$

**COROLLARY 4.3.** *If  $M$  is a maximal two-sided ideal in an  $N^*$ -complete regular ring  $R$ , then  $R/M$  is a simple, unit-regular, right and left self-injective ring. There is a unique rank function on  $R/M$ , and  $R/M$  is complete in the rank-metric.*

**PROOF.** According to Corollary 1.14,  $R/M$  is  $N^*$ -complete, hence there is no loss of generality in assuming that  $M = 0$ .

Since  $R$  is a nonzero unit-regular ring, [2, Corollary 18.5] shows that  $\mathbf{P}(R)$  is nonempty. By the Krein-Milman Theorem, there exists at least one  $P$  in  $\partial_e \mathbf{P}(R)$ . Note that  $\ker(P) = 0$ , because  $R$  is simple. Inasmuch as  $\{P\}$  is a closed face of  $\mathbf{P}(R)$ , we now conclude from Theorem 4.2(b) that  $\mathbf{P}(R) = \{P\}$ . Thus  $P$  is the unique rank function on  $R$ .

Now  $N^* = P$ , hence the  $N^*$ -metric on  $R$  coincides with the  $P$ -metric. Therefore  $R$  is complete in the  $P$ -metric. According to [2, Theorem 19.7],  $R$  is thus right and left self-injective.  $\square$

**COROLLARY 4.4.** *Let  $R$  be an  $N^*$ -complete regular ring, and let  $P \in \mathbf{P}(R)$ . Then the following conditions are equivalent.*

- (a)  *$P$  is an extreme point of  $\mathbf{P}(R)$ .*
- (b)  *$\ker(P)$  is a maximal two-sided ideal of  $R$ .*
- (c) *There is a unique pseudo-rank function on  $R/\ker(P)$ .*

PROOF. Let  $\bar{P}$  denote the rank function on  $R/\ker(P)$  induced by  $P$ .

(a)  $\Rightarrow$  (c): Since  $\{P\}$  is a closed face of  $\mathbf{P}(R)$ , Theorem 4.2(b) says that  $P$  is the only pseudo-rank function on  $R$  whose kernel contains  $\ker(P)$ . Thus  $\bar{P}$  is the only pseudo-rank function on  $R/\ker(P)$ .

(c)  $\Rightarrow$  (b): Since  $R/\ker(P)$  is a unit-regular ring possessing a unique rank function, [2, Corollary 18.6] shows that  $R/\ker(P)$  is a simple ring.

(b)  $\Rightarrow$  (a): By Corollary 4.3,  $\bar{P}$  is the only rank function on  $R/\ker(P)$ , hence  $P$  is the only pseudo-rank function on  $R$  whose kernel contains  $\ker(P)$ . Given a positive convex combination  $P = \alpha P_1 + (1 - \alpha)P_2$  in  $\mathbf{P}(R)$ , we see that  $\ker(P) \subseteq \ker(P_1)$  because  $\alpha > 0$ , whence  $P_1 = P$ , and similarly  $P_2 = P$ . Therefore  $P$  is an extreme point of  $\mathbf{P}(R)$ .  $\square$

With the help of Corollaries 4.3 and 4.4, we can show that in any  $N^*$ -complete regular ring  $R$ , there is a natural bijection between  $\partial_e \mathbf{P}(R)$  and the set of maximal two-sided ideals of  $R$ . In fact, this bijection is continuous with respect to the usual topology on the maximal ideal space, as follows.

DEFINITION. For any ring  $R$ , we use  $\text{MaxSpec}(R)$  to denote the family of all maximal two-sided ideals of  $R$ , equipped with the usual hull-kernel topology.

THEOREM 4.5. *Let  $R$  be an  $N^*$ -complete regular ring.*

(a) *There is a continuous bijection  $\theta: \partial_e \mathbf{P}(R) \rightarrow \text{MaxSpec}(R)$  given by the rule  $\theta(P) = \ker(P)$ .*

(b)  *$\theta$  maps compact subsets of  $\partial_e \mathbf{P}(R)$  onto closed subsets of  $\text{MaxSpec}(R)$ .*

(c)  *$\theta$  is a homeomorphism if and only if  $\partial_e \mathbf{P}(R)$  is compact, if and only if  $\text{MaxSpec}(R)$  is Hausdorff.*

PROOF. (a) It is clear from Corollaries 4.3 and 4.4 that  $\theta$  defines a bijection of  $\partial_e \mathbf{P}(R)$  onto  $\text{MaxSpec}(R)$ . If  $X$  is a closed subset of  $\text{MaxSpec}(R)$ , then

$$X = \{M \in \text{MaxSpec}(R) \mid Y \subseteq M\}$$

for some  $Y \subseteq R$ . Consequently,

$$\theta^{-1}(X) = \{P \in \partial_e \mathbf{P}(R) \mid P(y) = 0 \text{ for all } y \in Y\},$$

which is closed in  $\partial_e \mathbf{P}(R)$ . Therefore  $\theta$  is continuous.

(b) If  $X$  is a compact subset of  $\partial_e \mathbf{P}(R)$ , then

$$X = \{P \in \partial_e \mathbf{P}(R) \mid \ker(X) \subseteq \theta(P)\},$$

by Theorem 4.2(a). As a result,

$$\theta(X) = \{M \in \text{MaxSpec}(R) \mid \ker(X) \subseteq M\},$$

which is closed in  $\text{MaxSpec}(R)$ .

(c) Note that  $\partial_e \mathbf{P}(R)$  is Hausdorff while  $\text{MaxSpec}(R)$  is compact. Thus if  $\theta$  is a homeomorphism, then  $\partial_e \mathbf{P}(R)$  must be compact and  $\text{MaxSpec}(R)$  must be Hausdorff. On the other hand, if  $\partial_e \mathbf{P}(R)$  is compact, then we see from (b) that  $\theta$  is a closed map, whence  $\theta$  is a homeomorphism.

Now assume that  $\text{MaxSpec}(R)$  is Hausdorff. We shall show that  $\partial_e \mathbf{P}(R)$  is compact, by showing that  $\partial_e \mathbf{P}(R)$  is closed in  $\mathbf{P}(R)$ .

If not, then some  $P \in \mathbf{P}(R)$  lies in the closure of  $\partial_e \mathbf{P}(R)$  but not in  $\partial_e \mathbf{P}(R)$ . Set  $K = \ker(P)$ , and note from Corollary 4.4 that there exist more than one pseudo-rank functions on  $R/K$ . Because of the Krein-Milman Theorem, there must exist at least two extreme points in  $\mathbf{P}(R/K)$ . As  $R/K$  is  $N^*$ -complete (Theorem 1.13), it follows from (a) that  $R/K$  has at least two maximal two-sided ideals. Thus there exist distinct  $M_1$  and  $M_2$  in  $\text{MaxSpec}(R)$  which contain  $K$ .

Since  $\text{MaxSpec}(R)$  is Hausdorff, there exist disjoint open sets  $V_1$  and  $V_2$  in  $\text{MaxSpec}(R)$  such that each  $M_i \in V_i$ . There are subsets  $X_i \subseteq R$  such that each

$$V_i = \{M \in \text{MaxSpec}(R) \mid X_i \not\subseteq M\}.$$

For each  $i$ , choose  $x_i \in X_i$  such that  $x_i \notin M_i$ . Note that if  $M$  is in  $\text{MaxSpec}(R)$  and  $x_i \notin M$ , then  $M \in V_i$ . Thus for any  $M$  in  $\text{MaxSpec}(R)$ , we must have either  $x_1 \in M$  or  $x_2 \in M$ .

Inasmuch as  $x_i \notin M_i$ , we have  $x_i \notin K$ . Set

$$W = \{Q \in \mathbf{P}(R) \mid Q(x_1) > 0 \text{ and } Q(x_2) > 0\},$$

which is an open subset of  $\mathbf{P}(R)$ . Note that  $P \in W$ , because each  $x_i \notin \ker(P)$ . Since  $P$  lies in the closure of  $\partial_e \mathbf{P}(R)$ , there exists  $Q$  in  $W \cap \partial_e \mathbf{P}(R)$ . But then  $\ker(Q)$  is a maximal two-sided ideal of  $R$  which contains neither  $x_i$ , a contradiction.

Thus  $\partial_e \mathbf{P}(R)$  is indeed closed in  $\mathbf{P}(R)$ , and so is compact. Therefore  $\theta$  is a homeomorphism in this case also.  $\square$

**COROLLARY 4.6.** *If  $R$  is an  $N^*$ -complete regular ring, then the intersection of the maximal two-sided ideals of  $R$  is zero. Consequently,  $R$  is a subdirect product of simple, unit-regular, right and left self-injective rings.*

**PROOF.** If an element  $x \in R$  lies in all maximal two-sided ideals, then by Theorem 4.5,  $P(x) = 0$  for all  $P$  in  $\partial_e \mathbf{P}(R)$ . As the convex hull of  $\partial_e \mathbf{P}(R)$  is dense in  $\mathbf{P}(R)$  (the Krein-Milman Theorem), it follows that  $P(x) = 0$  for all  $P$  in  $\mathbf{P}(R)$ . Then  $N^*(x) = 0$ , whence  $x = 0$ .

The subdirect product statement now follows from Corollary 4.3.  $\square$

**COROLLARY 4.7.** *Let  $R$  be an  $N^*$ -complete regular ring, and let  $J$  be a two-sided ideal of  $R$ . Then the following conditions are equivalent.*

- (a)  $R/J$  is  $N^*$ -complete.
- (b)  $J$  is  $N^*$ -closed in  $R$ .
- (c)  $J = \ker(X)$  for some  $X \subseteq \mathbf{P}(R)$ .
- (d)  $J = \bigcap Y$  for some  $Y \subseteq \text{MaxSpec}(R)$ .

**PROOF.** As  $R$  is unit-regular, properties (a), (b), and (c) are equivalent by Theorem 1.13. We obtain (a)  $\Rightarrow$  (d) from Corollary 4.6, while it follows from Theorem 4.5(a) that (d)  $\Rightarrow$  (c).  $\square$

**COROLLARY 4.8.** *Let  $R$  be an  $N^*$ -complete regular ring. Then  $K_0(R)$  is a lattice-ordered abelian group if and only if  $\partial_e \mathbf{P}(R)$  is compact, if and only if  $\text{MaxSpec}(R)$  is Hausdorff.*

PROOF. By Corollary 3.2 and Proposition 4.1,  $K_0(R)$  is lattice-ordered if and only if  $\partial_e \mathbf{P}(R)$  is compact. Theorem 4.5 shows that  $\partial_e \mathbf{P}(R)$  is compact if and only if  $\text{MaxSpec}(R)$  is Hausdorff.  $\square$

An alternate view of Theorem 4.5(a) is that for any  $N^*$ -complete regular ring  $R$ , there is another topology on  $\partial_e \mathbf{P}(R)$ , coarser than the usual topology, with respect to which the bijection  $P \mapsto \ker(P)$  provides a homeomorphism of  $\partial_e \mathbf{P}(R)$  onto  $\text{MaxSpec}(R)$ . This topology coincides with a known topology defined on extreme boundaries of compact convex sets, as follows.

DEFINITION. Let  $K$  be any compact convex set, and let  $\mathcal{F}$  denote the family of subsets of  $\partial_e K$  of the form  $F \cap \partial_e K$ , where  $F$  is a closed split-face of  $K$ . (See [1, p. 133] for the definition of a split-face.) According to [1, Proposition II.6.20],  $\mathcal{F}$  is closed under finite unions and arbitrary intersections. Thus  $\mathcal{F}$  is the family of closed sets for a topology on  $\partial_e K$ , known as the *facial topology* [1, p. 143]. When  $K$  is a Choquet simplex, every closed face of  $K$  is a split-face [1, Theorem II.6.22], hence in this case  $\mathcal{F}$  consists of all sets of the form  $F \cap \partial_e K$  where  $F$  is any closed face of  $K$ . This simplification will apply in our considerations because  $\mathbf{P}(R)$ , for any regular ring  $R$ , is a Choquet simplex [2, Theorem 17.5].

THEOREM 4.9. *Let  $R$  be an  $N^*$ -complete regular ring. If  $\partial_e \mathbf{P}(R)$  is given the facial topology, then the rule  $P \mapsto \ker(P)$  defines a homeomorphism of  $\partial_e \mathbf{P}(R)$  onto  $\text{MaxSpec}(R)$ .*

PROOF. By Theorem 4.5(a), the rule  $\theta(P) = \ker(P)$  defines a bijection  $\theta$  of  $\partial_e \mathbf{P}(R)$  onto  $\text{MaxSpec}(R)$ . If  $X$  is a closed subset of  $\text{MaxSpec}(R)$ , then

$$X = \{M \in \text{MaxSpec}(R) \mid Y \subseteq M\},$$

for some  $Y \subseteq R$ . Set  $F = \{P \in \mathbf{P}(R) \mid Y \subseteq \ker(P)\}$ , and recall that  $F$  is a closed face of  $\mathbf{P}(R)$  [2, Lemma 16.18]. Observing that

$$\theta^{-1}(X) = F \cap \partial_e \mathbf{P}(R),$$

we see that  $\theta^{-1}(X)$  is closed in  $\partial_e \mathbf{P}(R)$  in the facial topology. Thus  $\theta$  is continuous with respect to the facial topology.

If  $F$  is any closed face of  $\mathbf{P}(R)$ , then Theorem 4.2(b) says that

$$F = \{P \in \mathbf{P}(R) \mid \ker(F) \subseteq \ker(P)\}.$$

As a result,

$$\theta(F \cap \partial_e \mathbf{P}(R)) = \{M \in \text{MaxSpec}(R) \mid \ker(F) \subseteq M\},$$

which is closed in  $\text{MaxSpec}(R)$ . Thus  $\theta$  is a closed map with respect to the facial topology.  $\square$

Part of Theorem 4.5(c) may be proved using Theorem 4.9, because of the general result that in a Choquet simplex  $K$ , the facial topology on  $\partial_e K$  is Hausdorff if and only if the usual topology on  $\partial_e K$  is compact [1, Theorem II.7.8].

We now turn to the study of finitely generated projective modules over an  $N^*$ -complete regular ring  $R$ . In particular, we show that their isomorphism classes are determined both by the values of pseudo-rank functions on  $R$ , and by their isomorphism classes modulo maximal two-sided ideals of  $R$ .

**THEOREM 4.10.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $A$  and  $B$  be finitely generated projective right  $R$ -modules, and get*

$$A \cong x_1R \oplus \cdots \oplus x_nR; \quad B \cong y_1R \oplus \cdots \oplus y_kR$$

for some elements  $x_1, \dots, x_n, y_1, \dots, y_k \in R$ .

(a)  $A \lesssim B$  if and only if  $A/AM \lesssim B/BM$  for all  $M \in \text{MaxSpec}(R)$ , if and only if

$$P(x_1) + \cdots + P(x_n) \leq P(y_1) + \cdots + P(y_k)$$

for all  $P \in \partial_e \mathbf{P}(R)$ .

(b)  $A \cong B$  if and only if  $A/AM \cong B/BM$  for all  $M \in \text{MaxSpec}(R)$ , if and only if

$$P(x_1) + \cdots + P(x_n) = P(y_1) + \cdots + P(y_k)$$

for all  $P \in \partial_e \mathbf{P}(R)$ .

**PROOF.** (a) If  $A \lesssim B$ , then obviously  $A/AM \lesssim B/BM$  for all  $M$  in  $\text{MaxSpec}(R)$ .

Now assume that  $A/AM \lesssim B/BM$  for all  $M \in \text{MaxSpec}(R)$ . Consider any  $P$  in  $\partial_e \mathbf{P}(R)$ , and set  $M = \ker(P)$ , which is in  $\text{MaxSpec}(R)$  by Corollary 4.4. Let  $\bar{P}$  denote the rank function induced on  $R/M$  by  $P$ . Using  $x \mapsto \bar{x}$  for the natural map  $R \rightarrow R/M$ , we have

$$\bar{x}_1(R/M) \oplus \cdots \oplus \bar{x}_n(R/M) \cong A/AM \lesssim B/BM \cong \bar{y}_1(R/M) \oplus \cdots \oplus \bar{y}_k(R/M).$$

Applying [2, Proposition 16.1], we obtain

$$\bar{P}(\bar{x}_1) + \cdots + \bar{P}(\bar{x}_n) \leq \bar{P}(\bar{y}_1) + \cdots + \bar{P}(\bar{y}_k),$$

and consequently  $\sum P(x_i) \leq \sum P(y_j)$ .

Finally, assume that  $\sum P(x_i) \leq \sum P(y_j)$  for all  $P \in \partial_e \mathbf{P}(R)$ . In view of Proposition 4.1, it follows that

$$s([A]) = s([x_1R] + \cdots + [x_nR]) \leq s([y_1R] + \cdots + [y_kR]) = s([B])$$

for all extreme states  $s$  on  $(K_0(R), [R])$ . Since  $K_0(R)$  is archimedean, we conclude from [4, Proposition I.5.3] that  $[A] \leq [B]$ . Therefore  $A \lesssim B$ .

(b) This follows directly from (a) and the unit-regularity of  $R$ .  $\square$

**DEFINITION.** Let  $R$  be any regular ring, and set  $S = S(K_0(R), [R])$ . By Proposition 4.1, there is a natural affine homeomorphism  $\theta: S \rightarrow \mathbf{P}(R)$  such that  $\theta(s)(x) = s([xR])$  for all  $s \in S$  and all  $x \in R$ . Then  $\theta$  induces an isomorphism  $\theta^*$  of  $\text{Aff}(\mathbf{P}(R))$  onto  $\text{Aff}(S)$ , as partially ordered Banach spaces. We also have the natural evaluation map  $\varphi$  from  $K_0(R)$  to  $\text{Aff}(S)$ . Composing  $\varphi$  with  $(\theta^*)^{-1}$ , we obtain a positive homomorphism

$$\psi = (\theta^*)^{-1} \varphi: K_0(R) \rightarrow \text{Aff}(\mathbf{P}(R))$$

such that  $\psi(x)(P) = \theta^{-1}(P)(x)$  for all  $x \in K_0(R)$  and all  $P \in \mathbf{P}(R)$ . In particular,  $\psi([xR])(P) = P(x)$  for all  $x \in R$  and all  $P \in \mathbf{P}(R)$ . We refer to  $\psi$  as the natural map from  $K_0(R)$  to  $\text{Aff}(\mathbf{P}(R))$ . Note that  $\psi([R])$  is the constant function 1.

**THEOREM 4.11.** *Let  $R$  be an  $N^*$ -complete regular ring. Whenever  $P \in \partial_e \mathbf{P}(R)$  and  $R/\ker(P)$  is isomorphic to an  $m \times m$  matrix ring over a division ring, set  $A_P = (1/m)\mathbf{Z}$ ; for all other  $P \in \partial_e \mathbf{P}(R)$ , set  $A_P = \mathbf{R}$ . Set*

$$A = \{q \in \text{Aff}(\mathbf{P}(R)) \mid q(P) \in A_P \text{ for all } P \in \partial_e \mathbf{P}(R)\}.$$

*Then the natural map  $\psi: K_0(R) \rightarrow \text{Aff}(\mathbf{P}(R))$  provides an isomorphism of  $(K_0(R), [R])$  onto  $(A, 1)$  (as partially ordered abelian groups with order-unit).*

**PROOF.** Define  $S, \theta, \varphi$  as in the definition above, so that  $\psi = (\theta^*)^{-1}\varphi$ . For all discrete  $s \in \partial_e S$ , set  $B_s = s(K_0(R))$ ; for all other  $s \in \partial_e S$ , set  $B_s = \mathbf{R}$ . Set

$$B = \{p \in \text{Aff}(S) \mid p(S) \in B_s \text{ for all } s \in \partial_e S\}.$$

According to Theorem 3.1,  $\varphi$  provides an isomorphism of  $(K_0(R), [R])$  onto  $(B, 1)$ . Thus we need only show that  $(\theta^*)^{-1}$  restricts to an isomorphism of  $(B, 1)$  onto  $(A, 1)$ . To prove this, it suffices to show that  $A_{\theta(s)} = B_s$  for all  $s \in \partial_e S$ .

Thus let  $s \in \partial_e S$ , and set  $P = \theta(s)$  and  $M = \ker(P)$ , so that  $P \in \partial_e \mathbf{P}(R)$  and  $M$  is a maximal two-sided ideal of  $R$ . Let  $\bar{P}$  be the rank function induced by  $P$  on  $R/M$ .

If  $s$  is discrete, then  $s$  induces an isomorphism of  $K_0(R)/\ker(s)$  onto  $(1/m)\mathbf{Z}$  for some  $m \in \mathbf{N}$  (Theorem 3.10), whence

$$\bar{P}(R/M) = P(R) = \{0, 1/m, 2/m, \dots, 1\}.$$

Choosing  $x \in R/M$  such that  $\bar{P}(x) = 1/m$ , we infer that  $x(R/M)$  is a minimal right ideal of  $R/M$ , whence  $R/M$  is a simple artinian ring. Thus  $R/M \cong M_k(D)$  for some  $k \in \mathbf{N}$  and some division ring  $D$ . There is a unique rank function on  $M_k(D)$ , and its range of values is  $\{0, 1/k, 2/k, \dots, 1\}$  [2, Corollary 16.6]. Consequently, we must have  $k = m$ , and so

$$A_{\theta(s)} = A_P = (1/m)\mathbf{Z} = s(K_0(R)) = B_s.$$

If  $s$  is not discrete, then  $s$  induces an isomorphism of  $K_0(R)/\ker(s)$  onto  $\mathbf{R}$ , by Theorem 3.10, whence

$$\bar{P}(R/M) = P(R) = [0, 1].$$

In this case,  $R/M$  cannot be a simple artinian ring, hence

$$A_{\theta(s)} = A_P = \mathbf{R} = B_s,$$

as desired.  $\square$

**COROLLARY 4.12.** *Let  $R$  be an  $N^*$ -complete regular ring. If  $R$  has no simple artinian homomorphic images, then the natural map from  $K_0(R)$  to  $\text{Aff}(\mathbf{P}(R))$  is an isomorphism of partially ordered abelian groups.  $\square$*

**THEOREM 4.13.** *Let  $R$  be an  $N^*$ -complete regular ring, let  $B$  be a finitely generated projective right  $R$ -module, and let  $n \in \mathbf{N}$ . Assume, for each maximal two-sided ideal  $M$  of  $R$  such that  $R/M$  is artinian, that  $B/BM$  is a direct sum of  $n$  pairwise isomorphic submodules. Then  $B$  is a direct sum of  $n$  pairwise isomorphic submodules. In particular, if  $R$  has no simple artinian homomorphic images, then this holds for all  $n \in \mathbf{N}$ .*

PROOF. In the notation of Theorem 4.11, we wish to show that the function  $\psi([B])/n$  lies in  $A$ . Thus consider any  $P \in \partial_e \mathbf{P}(R)$  such that  $R/\ker(P) \cong M_m(D)$  for some  $m \in \mathbf{N}$  and some division ring  $D$ . The ideal  $M = \ker(P)$  is a maximal two-sided ideal of  $R$ , and  $R/M$  has a unique simple right module  $S$ . Then  $B/BM \cong kS$  for some  $k \in \mathbf{Z}^+$ , and  $\psi([B])(P) = k/m$ . Since  $B/BM$  is assumed to be a direct sum of  $n$  pairwise isomorphic submodules,  $n$  must divide  $k$ , hence

$$\psi([B])(P)/n = (k/n)/m \in (1/m)\mathbf{Z} = A_P.$$

As this holds for all  $P \in \partial_e \mathbf{P}(R)$  such that  $R/\ker(P)$  is simple artinian, we find that  $\psi([B])/n$  does lie in  $A$ , as desired.

In fact,  $\psi([B])/n$  must lie in  $A^+$ . Because of the isomorphism given in Theorem 4.11, it follows that there exists  $x \in K_0(R)^+$  such that  $nx = [B]$ . Then  $x = [C]$  for some finitely generated projective right  $R$ -module  $C$ , and  $[nC] = [B]$ . Therefore  $B \cong nC$ .  $\square$

**COROLLARY 4.14.** *Let  $R$  be an  $N^*$ -complete regular ring, and let  $n \in \mathbf{N}$ . If every simple artinian homomorphic image of  $R$  is an  $n \times n$  matrix ring, then  $R$  is an  $n \times n$  matrix ring. In particular, if  $R$  has no simple artinian homomorphic images, then  $R$  is an  $n \times n$  matrix ring for every  $n \in \mathbf{N}$ .  $\square$*

As another application of the affine representation of  $K_0$  of an  $N^*$ -complete regular ring (Theorem 4.11 and Corollary 4.12), we derive criteria for the following properties.

**DEFINITION.** Let  $R$  be a unit-regular ring. We say that  $R$  satisfies *countable interpolation* provided that given any elements  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$  in  $R$  satisfying  $x_i R \lesssim y_j R$  for all  $i, j$ , there exists  $z \in R$  such that  $x_i R \lesssim zR \lesssim y_j R$  for all  $i, j$ . According to [4, Proposition II.12.1], this property is left-right symmetric, and is equivalent to the countable interpolation property in  $K_0(R)$ . The ring  $R$  is said to satisfy *general comparability* provided that given any  $x, y \in R$ , there exists a central idempotent  $e \in R$  such that  $exR \lesssim eyR$  and  $(1 - e)yR \lesssim (1 - e)xR$ .

**DEFINITION.** Let  $X$  be a compact Hausdorff space. The space  $X$  is said to be an *F-space* if disjoint open  $F_e$  subsets of  $X$  always have disjoint closures. The space  $X$  is said to be *basically disconnected* if the closure of every open  $F_e$  subset of  $X$  is open.

**THEOREM 4.15.** *Let  $R$  be an  $N^*$ -complete regular ring with no simple artinian homomorphic images, and assume that  $\partial_e \mathbf{P}(R)$  is compact. Then  $R$  has countable interpolation if and only if  $\partial_e \mathbf{P}(R)$  is an  $F$ -space, if and only if  $\text{MaxSpec}(R)$  is an  $F$ -space.*

PROOF. Set  $X = \partial_e \mathbf{P}(R)$ , and note from Theorem 4.5 that  $X$  is homeomorphic to  $\text{MaxSpec}(R)$ . Combining Corollary 4.12 with [1, Proposition II.3.13], we see that

$$K_0(R) \cong \text{Aff}(\mathbf{P}(R)) \cong C(X, \mathbf{R})$$

as partially ordered abelian groups. Thus  $R$  has countable interpolation if and only if  $C(X, \mathbf{R})$  satisfies the countable interpolation property (as a partially ordered set). According to [9, Theorem 1.1], this happens if and only if  $X$  is an  $F$ -space.  $\square$



Theorem 4.15 may fail if  $R$  is allowed to have simple artinian homomorphic images, as the following example shows.

EXAMPLE 4.16. There exists an  $N^*$ -complete regular ring  $R$  such that  $\partial_e \mathbf{P}(R)$  is a compact  $F$ -space, but  $R$  does not have countable interpolation.

PROOF. Choose a field  $K$ , set  $R_n = M_2(K)$  for all  $n \in \mathbf{N}$ , and set

$$R = \{x \in \prod R_n \mid x_n \in K \text{ for all but finitely many } n \in \mathbf{N}\}.$$

Clearly  $R$  is a regular ring whose index of nilpotence is 2. By Theorem 1.3,  $R$  is  $N^*$ -complete.

The Boolean algebra  $B(R)$  of central idempotents in  $R$  is a direct product of copies of  $\{0, 1\}$  and so is complete. Consequently, its maximal ideal space  $BS(R)$  is compact, Hausdorff, and extremally disconnected. In particular,  $BS(R)$  is a compact  $F$ -space. Observing that  $R$  satisfies general comparability, we see by [2, Theorem 16.28] that  $\partial_e \mathbf{P}(R)$  is homeomorphic to  $BS(R)$ . Thus  $\partial_e \mathbf{P}(R)$  is a compact  $F$ -space.

Choose  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$  in  $R$  so that  $\text{rank}(x_{kn}) = \text{rank}(y_{kn}) = 1$  for all  $n = 1, \dots, k$ , whereas  $x_{kn} = 0$  and  $y_{kn} = 1$  for all  $n > k$ . Clearly  $x_i R \lesssim y_j R$  for all  $i, j$ . If there exists  $z \in R$  such that  $x_i R \lesssim zR \lesssim y_j R$  for all  $i, j$ , then  $x_{nn} R \lesssim z_n R \lesssim y_{nn} R$  for all  $n \in \mathbf{N}$ , whence  $\text{rank}(z_n) = 1$  for all  $n$ . But then  $z_n \notin K$  for all  $n$ , which is impossible for an element  $z \in R$ . Therefore  $R$  does not satisfy countable interpolation.  $\square$

THEOREM 4.17. Let  $R$  be an  $N^*$ -complete regular ring. If  $\partial_e \mathbf{P}(R)$  is a compact totally disconnected  $F$ -space, then  $R$  satisfies general comparability.

PROOF. Set  $\Delta = \partial_e \mathbf{P}(R)$ . Given  $x, y \in R$ , set

$$X = \{P \in \Delta \mid P(x) < P(y)\}; \quad Y = \{P \in \Delta \mid P(x) > P(y)\}.$$

Inasmuch as the rule  $P \mapsto P(x) - P(y)$  defines a continuous real-valued map on  $\Delta$ , we see that  $X$  and  $Y$  are disjoint open  $F_\sigma$  subsets of  $\Delta$ . Since  $\Delta$  is assumed to be an  $F$ -space, the closures of  $X$  and  $Y$  must be disjoint. Consequently, it follows from the total disconnectedness of  $\Delta$  that there is a clopen set  $V \subseteq \Delta$  such that  $X \subseteq V$  and  $Y \subseteq \Delta - V$ .

Now let  $\theta: \Delta \rightarrow \text{MaxSpec}(R)$  be the homeomorphism given by Theorem 4.5, so that  $\theta(V)$  is a clopen subset of  $\text{MaxSpec}(R)$ . As the intersection of the maximal two-sided ideals of  $R$  is zero (Corollary 4.6), there must exist a central idempotent  $e \in R$  such that

$$\theta(V) = \{M \in \text{MaxSpec}(R) \mid e \notin M\}.$$

Given any  $P \in V$ , we have  $\ker(P) \in \theta(V)$  and so  $e \notin \ker(P)$ . Then  $1 - e$  lies in  $\ker(P)$ , hence  $P(x) = P(ex)$  and  $P(y) = P(ey)$ , by [2, Lemma 16.2]. Consequently,  $P(ex) < P(ey)$  for all  $P \in X$ , and  $P(ex) = P(ey)$  for all  $P \in V - X$ . On the other hand, for  $P \in \Delta - V$ , we have  $\ker(P) \notin \theta(V)$  and so  $e \in \ker(P)$ , whence  $P(ex) = 0 = P(ey)$ . Thus  $P(ex) \leq P(ey)$  for all  $P \in \Delta$ , hence  $exR \lesssim eyR$ , by Theorem 4.10. Similarly,  $(1 - e)yR \lesssim (1 - e)xR$ . Therefore  $R$  satisfies general comparability.  $\square$

The proof of Theorem 4.17 can be considerably shortened if  $R$  has no simple artinian homomorphic images. For in this case  $R$  has countable interpolation by Theorem 4.15, and then [4, Theorem II.14.7] shows that  $R$  satisfies general comparability.

**COROLLARY 4.18.** *Let  $R$  be an  $N^*$ -complete regular ring with no simple artinian homomorphic images. Then  $R$  satisfies general comparability if and only if  $\partial_e \mathbf{P}(R)$  is a compact totally disconnected  $F$ -space, if and only if  $\text{MaxSpec}(R)$  is a Hausdorff totally disconnected  $F$ -space.*

**PROOF.** It is clear from Theorem 4.5 that  $\partial_e \mathbf{P}(R)$  is a compact totally disconnected  $F$ -space if and only if  $\text{MaxSpec}(R)$  is a Hausdorff totally disconnected  $F$ -space. These conditions imply general comparability in  $R$  by Theorem 4.17.

Conversely, assume that  $R$  has general comparability. It follows from [2, Theorem 16.28] that  $\partial_e \mathbf{P}(R)$  is compact and totally disconnected.

Now consider any disjoint open  $F_e$  subsets  $X$  and  $Y$  in  $\partial_e \mathbf{P}(R)$ . Then there exists a continuous real-valued function  $f$  on  $\partial_e \mathbf{P}(R)$  such that  $f > 0$  on  $X$  and  $f < 0$  on  $Y$ . (This is an exercise in applying Urysohn’s Lemma, which we leave to the reader.) As  $\partial_e \mathbf{P}(R)$  is compact, we can modify  $f$  to obtain a continuous function

$$g: \partial_e \mathbf{P}(R) \rightarrow [0, 1]$$

such that  $g > 1/2$  on  $X$  and  $g < 1/2$  on  $Y$ . By [1, Proposition II.3.13],  $g$  extends to an affine continuous function  $g^*: \mathbf{P}(R) \rightarrow [0, 1]$ , to which we apply Corollary 4.12. Since  $0 \leq g^* \leq 1$ , we obtain an element  $b \in K_0(R)$  such that  $0 \leq b \leq [R]$  and the map induced by  $b$  in  $\text{Aff}(\mathbf{P}(R))$  coincides with  $g^*$ . Thus  $b = [xR]$  for some  $x \in R$ , and  $g^*(P) = P(x)$  for all  $P \in \mathbf{P}(R)$ .

We use general comparability to compare the projective modules  $2(xR)$  and  $R_R$ , via [2, Proposition 8.8]. Thus we obtain a central idempotent  $e \in R$  such that

$$2(exR) \lesssim eR \quad \text{and} \quad (1 - e)R \lesssim 2((1 - e)xR).$$

Consider any  $P \in X$ , so that  $P(x) = g(P) > 1/2$ . If  $e \notin \ker(P)$ , then  $1 - e$  is in  $\ker(P)$ , because  $\ker(P)$  is a maximal two-sided ideal of  $R$  (Corollary 4.4). But then

$$2P(x) = 2P(ex) \leq P(e) = 1$$

(since  $2(exR) \lesssim eR$ ), which contradicts the fact that  $P(x) > 1/2$ . Thus  $e \in \ker(P)$  and so  $P(e) = 0$ . Similarly, for any  $P \in Y$  we have  $1 - e \in \ker(P)$  and so  $P(e) = 1$ . As the map  $P \mapsto P(e)$  is a continuous map from  $\mathbf{P}(R)$  to  $\mathbf{R}$ , we conclude that  $X$  and  $Y$  must have disjoint closures.

Therefore  $\partial_e \mathbf{P}(R)$  is an  $F$ -space.  $\square$

Corollary 4.18 may fail if  $R$  is allowed to have simple artinian homomorphic images, as the following example shows.

**EXAMPLE 4.19.** There exists an  $N^*$ -complete regular ring  $R$  such that  $R$  satisfies general comparability, but  $\partial_e \mathbf{P}(R)$  is not an  $F$ -space.

**PROOF.** Choose a field  $K$ , and let  $R$  be the ring of all eventually constant sequences

$$(\alpha_1, \alpha_2, \dots, \alpha_n, \alpha, \alpha, \alpha, \dots)$$

of elements of  $K$ . Clearly  $R$  is a regular ring whose index of nilpotence is 1. By Theorem 1.3,  $R$  is  $N^*$ -complete. It is also clear that  $R$  satisfies general comparability.

The maximal ideals of  $R$  are easily identified, from which one sees that  $\text{MaxSpec}(R)$  is homeomorphic to the one-point compactification of  $\mathbb{N}$ . In particular,  $\text{MaxSpec}(R)$  is Hausdorff but is not an  $F$ -space. By Theorem 4.5,  $\partial_e \mathbf{P}(R)$  is homeomorphic to  $\text{MaxSpec}(R)$ , hence  $\partial_e \mathbf{P}(R)$  is not an  $F$ -space.  $\square$

As we mentioned in the introduction to this section, many of our results—particularly Corollaries 4.3 and 4.6, Theorems 4.10 and 4.13, and Corollary 4.14—show that an  $N^*$ -complete regular ring  $R$  behaves much like a ring of sections of a sheaf of simple self-injective rings. The obvious candidate for a topological space on which such a sheaf should live is  $\text{MaxSpec}(R)$ . However,  $\text{MaxSpec}(R)$  is usually not Hausdorff, and even when it is Hausdorff it need not be disconnected, which would seem to be required for sheaf-theoretic proofs of results such as Corollary 4.14. The lack of Hausdorffness, at least, of  $\text{MaxSpec}(R)$  may be remedied by using the space  $\partial_e \mathbf{P}(R)$  instead; the fact that we have to pay attention to how  $\partial_e \mathbf{P}(R)$  sits inside  $\mathbf{P}(R)$  is in some sense the price for overcoming the lack of compactness of  $\partial_e \mathbf{P}(R)$ . This feeling may be made more precise at the level of  $K_0(R)$ : if  $R$  were to resemble the sections of a sheaf based on  $\partial_e \mathbf{P}(R)$ , then  $K_0(R)$  should resemble the sections of a sheaf of functions based on  $\partial_e \mathbf{P}(R)$ . When  $\partial_e \mathbf{P}(R)$  is compact, this does happen: Combining Theorem 4.11 with [1, Proposition II.3.13] yields

$$K_0(R) \cong \{q \in C(\partial_e \mathbf{P}(R), \mathbf{R}) \mid q(P) \in A_P \text{ for all } P \in \partial_e \mathbf{P}(R)\}.$$

On the other hand, when  $\partial_e \mathbf{P}(R)$  is not compact, continuous real-valued functions on  $\partial_e \mathbf{P}(R)$  do not correspond to elements of  $K_0(R)$  unless they can be made to respect the affine relations present in  $\mathbf{P}(R)$ .

There is one situation in which  $R$  is isomorphic to the ring of sections of a sheaf-like object, namely when  $R$  is a continuous regular ring. This idea is developed by Handelman in [5] under the additional assumption that  $R$  is  $N^*$ -complete; however, we now know that  $N^*$ -completeness holds for all continuous regular rings, by Theorem 1.8. Handelman's construction is based on the space  $\partial_e \mathbf{P}(R)$ , which in this case is a compact Hausdorff extremally disconnected space. (Of course, we could equally well use  $\text{MaxSpec}(R)$ , in view of Theorem 4.5.) The stalk at a point  $P \in \partial_e \mathbf{P}(R)$  is the simple self-injective ring  $R/\ker(P)$ . What prevents Handelman's object from being an actual sheaf is that the rank-metric topologies on the rings  $R/\ker(P)$  must be respected, and these topologies are not discrete unless the rings  $R/\ker(P)$  are artinian.

Another aspect of Handelman's development of this construction provides a general means for constructing  $N^*$ -complete regular rings. Namely, given any regular ring  $R$ , one can complete  $R$  with respect to the  $N^*$ -metric. Since the ring operations on  $R$  are uniformly continuous with respect to  $N^*$  (as we observed in §I), the  $N^*$ -completion  $\bar{R}$  is again a ring; moreover,  $\bar{R}$  is actually a regular ring [6, Proposition 1.4]. In addition, the restriction map  $\mathbf{P}(\bar{R}) \rightarrow \mathbf{P}(R)$  is an affine homeomorphism, as stated in [5, Proposition 15]. (The proof of this proposition is incomplete, but Handelman has informed me that he and Walter Burgess have developed a complete proof.) In particular, this result provides a convenient means

for constructing examples. Combining it with [2, Theorem 17.23], we find that any metrizable Choquet simplex  $S$  is affinely homeomorphic to  $\mathbf{P}(R)$  for a suitable  $N^*$ -complete regular ring  $R$ . For instance, there exists an  $N^*$ -complete regular ring  $R$  such that  $\mathbf{P}(R)$  is affinely homeomorphic to the Choquet simplex of all probability measures on the unit interval  $[0, 1]$ , whence  $\partial_e \mathbf{P}(R)$  is homeomorphic to  $[0, 1]$ .

A number of the results proved here for  $N^*$ -complete regular rings were first proved for  $\aleph_0$ -continuous regular rings [7, 8], or, somewhat more generally, for unit-regular rings satisfying countable interpolation [4]. (All  $\aleph_0$ -continuous regular rings, and their factor rings, satisfy countable interpolation by [4, Theorem II.12.3].) In the first case, our results are generalizations, since all  $\aleph_0$ -continuous regular rings are  $N^*$ -complete (Theorem 1.8). However, in the second case our results are not strict generalizations except at the level of  $K_0$ : Namely, a unit-regular ring  $R$  satisfying countable interpolation need not be  $N^*$ -complete, but as long as  $\ker(\mathbf{P}(R)) = 0$ , then  $K_0(R)$  is an archimedean norm-complete interpolation group [4, Theorems II.12.7 and I.6.6]. Thus the appropriate results for unit-regular rings with countable interpolation follow from the  $K_0$  results in §III, in the same manner as the derivations in §IV for  $N^*$ -complete regular rings. The relationship between our results and these earlier results is as follows.

Theorem 4.2 corresponds to [4, Corollary II.13.6 and Theorem II.13.7], while Corollary 4.3 corresponds to [7, Corollary 3.2]. Theorems 4.5 and 4.9 correspond to [4, Proposition II.14.5 and Theorem II.14.6], while Corollary 4.6 and Theorem 4.10 correspond to [8, Theorem 2.3]. Theorems 4.11 and 4.13, along with Corollaries 4.12 and 4.14, correspond to [4, Theorems II.15.1, II.15.3, and II.15.4, and Corollary II.15.2].

NOTE ADDED IN PROOF. The  $N^*$ -completion results mentioned above are included in a paper by Burgess and Handelmann, *The  $N^*$ -metric completion of regular rings*, submitted for publication.

#### REFERENCES

1. E. M. Alfsen, *Compact convex sets and boundary integrals*, Springer-Verlag, Berlin, 1971.
2. K. R. Goodearl, *Von Neumann regular rings*, Pitman, London, 1979.
3. K. R. Goodearl and D. E. Handelman, *Metric completions of partially ordered abelian groups*, Indiana Univ. Math. J. **29** (1980), 861–895.
4. K. R. Goodearl, D. E. Handelman and J. W. Lawrence, *Affine representations of Grothendieck groups and applications to Rickart  $C^*$ -algebras and  $\aleph_0$ -continuous regular rings*, Mem. Amer. Math. Soc. No. 234 (1980).
5. D. Handelman, *Representing rank complete continuous rings*, Canad. J. Math. **28** (1976), 1320–1331.
6. ———, *Simple regular rings with a unique rank function*, J. Algebra **42** (1976), 60–80.
7. ———, *Finite Rickart  $C^*$ -algebras and their properties*, Studies in Analysis, Advances in Math. Suppl. Studies, Vol. 4, 1979, pp. 171–196.
8. D. Handelman, D. Higgs and J. Lawrence, *Directed abelian groups, countably continuous rings, and Rickart  $C^*$ -algebras*, J. London Math. Soc. (2) **21** (1980), 193–202.
9. G. L. Seever, *Measures on  $F$ -spaces*, Trans. Amer. Math. Soc. **133** (1968), 267–280.
10. J. von Neumann, *Continuous geometry*, Princeton Univ. Press, Princeton, N. J., 1960.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 84112