

Microalbuminuria and Its Risk Factors in Type 2 Diabetic Patients

Sanjay Kumar Thakur,¹ Sukesh Purush Dhakal,¹ Shreeya Parajuli,² Anil Kumar Sah,³ Suresh Prasad Nepal,¹ Bishnu Dutta Paudel¹

¹Bir Hospital, Kathmandu, Nepal, ²Annapurna Neurological Institute and Allied Sciences, Kathmandu Nepal, ³Annapurna Research Center, Kathmandu, Nepal.

ABSTRACT

Background: Diabetic nephropathy is one of the dreaded complications of diabetes leading to chronic kidney disease and end stage renal failure globally. Microalbuminuria is the most sensitive marker of early recognition of the diabetic nephropathy. This study was carried out to find out the prevalence and potential risk factors of microalbuminuria which is the marker of diabetic nephropathy among diabetes patients in Nepal.

Methods: A cross-sectional study was conducted on a random sample of 227 in T2DM patients in private diabetic clinics and Bir hospital in Kathmandu. Data were collected using standard questionnaire format to collect demography, anthropometry, and laboratory assessment of, HbA1c, post prandial, fasting glucose and micro albumin in urine. Micro albuminuria was measured using early morning urine specimen. Micro albuminuria was considered positive when urinary albumin to creatinine ratio was found to be 30–300 mg/g creatinine in preferably an early morning or a spot urine sample. The entire lab test was done by applying the internationally accepted standards of tools and techniques. Those that were reported >30mg/l of micro albumin were considered as positive.

Results: Out of total 217 diabetic patients, 56.2% (122/217) were male and 43.8% (95/217) were female. Among all age groups, maximum patients enrolled were between the age group 41 to 80 (95%). Of the total, 20% (44/217) patients were MA positive. A statistical significant association was seen between MA and BMI ($p=0.029$), duration of DM ($p<0.001$), hypertension ($p<0.001$), smoking ($p<0.001$) and physical activity ($p<0.001$).

Conclusions: Diabetic patients in Nepal have prevalence of 20.3% microalbuminurea. Hypertension, obesity, sedentary lifestyles, duration more than 5 years of illness are found the most important risk factors for the development of microalbuminurea in diabetes.

Keywords: Microalbuminurea; type 2 diabetes Mellitus

INTRODUCTION

Diabetes is managed with glycemic control and risk reduction.¹ worldwide it is a major public health problem.² Around 10-20% of T2DM has end stage renal failure and reduction in life expectancy.³

WHO defines diabetes as symptoms suggesting diabetes and Fasting blood sugar >7.0 mmol/l (>126mg/dl) and/or 2 hrs after glucose load >11.1mmol/l (>200mg/dl) or a random plasma glucose of ≥ 200 (mg/dl).⁴

Random spot urine sample for urine albumin to-creatinine ratio (UACR) in patients with 5-yr diabetes duration should be checked.⁵

Diabetic nephropathy (DN) leads to chronic kidney disease requiring dialysis life long.⁶ Halting the progression is important in management.⁷ Microalbuminuria (MA) is the most sensitive marker of the diabetic nephropathy.⁸ Ahmad et al in Pakistan found 31.56% prevalence of microalbuminuria in T2DM.⁹ T2DM patients, 20-40% with MA progress to overt nephropathy and 20 years later, approximately 20% develop end stage renal disease (ESRD).⁸

Intensive control of blood pressure, hyperglycemia and lipid levels needed.¹⁰ Monitoring microalbuminuria and associated factors are important to prevent overt nephropathy.¹¹ Persoon et al, states that early intervention with ACEIs or ARBs reduces

Correspondence: Sanjay Kumar Thakur, Bir Hospital, Kathmandu, Nepal, Email: sanjay617@hotmail.com, Phone: +9779841270366.

microalbuminuria.¹² Urine albumin level is the main marker of DN.¹³ Microalbuminuria should be monitored once or twice a year.¹⁴ Screening of microalbuminuria to modify risk factors are essential to reduce end-stage renal disease.¹⁵

Study was to done to find out prevalence of microalbuminuria and associated risk factors in T2DM.

METHODS

This study was approved by the ethical review committee of the Bir hospital, Kathmandu. A cross-sectional study was conducted on Diabetes type 2 patients aged 20 years above who attended Annapurna Neurological Institutes and Allied Sciences, Global Hospital, Lagankhel Poly Clinic and Bir hospital, Kathmandu, Nepal from September 2017 to May 2018. A total of 217 patients were enrolled in this study after taking informed consent based on the set inclusion and exclusion criteria. Inclusion criteria were; diabetes type 2 patients aged 20 years and above who attend diabetic clinics in Kathmandu.

Exclusion criteria were presence of overt proteinuria, urinary tract infection, hematuria, ketonuria, pregnancy, heart failure, and use of systemic steroids in the past four weeks.

The following variables were obtained using a questionnaire filled through direct interview with the study participants: age, sex, level of education, marital status, smoking habit, family history of hypertension and diabetes mellitus, comorbid illnesses, and type of diabetes therapy. Body mass index (BMI) and blood pressure were measured using standard methods.

Body mass index (BMI) was divided into underweight, Normal weight, over weight and obese with BMI of <17.50, 17.50-22.9, 23.00-27.99, and >28 for Asian population. Patients were categorized as hypertensive if the systolic/diastolic blood pressure was >140/90 mmHg. Micro-albumin (MA) was defined as micro-albumin \geq 30-300 mg/L on urine specimen. While on diabetes treatment, Glycated hemoglobin (HbA1c) <7%; post prandial and fasting blood glucose below 180 mg/dl and 120 respectively were considered a good control of sugar.

Distributions, percentage and frequency of diabetes patients having micro albuminuria, internal group comparison and Pearson's chi-square test was used to assess intergroup significance at 95% confidence interval by using standard formula. BMI calculation-calculated applying the formula, i.e., BMI = (Weight in Kilograms/ (Height in Meters x Height in Meters)

RESULTS

A total of 217 diabetic patients were analyzed for the micro albuminuria; 56.2 %(122/217) weremale and 43.8% (95/217) were female. Among all age groups maximum patients were between ages 41 - 60 years 49.8% (108/217). 61-80, 21-40 and 81-100 as 49.8% (108/217), 46.1% (100/217), 2.8% (6/217) and 1.4% (3/217) respectively. There were 66.6% (145/217), 21.7% (47/217) and 11.5% (25/217) over weight, obese and normal weight respectively. The majority of the patients (84.3%) were <5 years duration of diabetes illness since diagnosed and physically active were found (64.1%). Whereas 30% (65/217), 30% (65/217), and 12% (26/217) had hypertension, good sugar control and smoking respectively (Table 1).

Table 1. Diabetics' sociodemographic features and disease related variable characteristics.

Variables	Frequency (%)
Gender	
Male	122 (56.2)
Female	95 (43.8)
Age	
21-40	6 (2.8)
41-60	108 (49.8)
61-80	100 (46.1)
81-100	3 (1.4)
BMI	
Normal weight	25 (11.5)
Over weight	145 (66.6)
Obese	47 (21.7)
Sugar Control	
Yes	65 (30)
No	152 (70)
Duration of DM	
<5	183 (84.3)
>5	34 (15.7)
Hypertension	
Yes	65 (30)
No	152 (70)
Smoking	
Yes	26 (12)
No	191 (88)
Physical Activity	
Active	139 (64.1)
Sedentary	78 (35.9)

The overall prevalence of MA was found to be 20.3%, the prevalence of MA was higher in male 52.3% than female 47.7% and there was no statistically different % (p=0.554). Similarly, higher prevalence was observed in age group 61-80 (52.3%) and lower at 81-100 (2.3%) years and there were no any significant difference among others age groups (p=0.517). MA was statistically significantly associated (p=0.029) with the BMI of overweight (61.4%) and obese (15.9%) than the normal weight (22.7%). A significantly difference association (p=0.056) was found between the prevalence of MA and status of sugar control (18.2%) and uncontrolled (81.8%) (Table 2).

Table 2. Relation of microalbuminuria with sociodemographic features and disease related variables.

Variables	Microalbumin		P Value
	Positive	Negative	
Sex			
Male	23 (52.3%)	99 (57.2%)	0.554
Female	21 (47.7%)	74 (42.8%)	
Age			
21-40	2 (4.5%)	4 (2.3%)	0.517
41-60	18 (40.9%)	90 (52.0%)	
61-80	23 (52.3%)	77 (44.5%)	
81-100	1 (2.3%)	2 (1.2%)	
BMI			
Normal weight	10 (22.7%)	15 (8.7%)	0.029
Over weight	27 (61.4%)	118 (68.2%)	
Obese	7 (15.9%)	40 (23.1%)	
Sugar Control			
Yes	8 (18.2%)	57 (32.9%)	0.056
No	36 (81.8%)	116 (67.1%)	
Duration of DM			
<5 Years	10 (22.7%)	173 (100.0%)	0.00
>5 Years	34 (77.3%)	0 (0.0%)	
Hypertension			
Yes	24 (54.5%)	41 (23.7%)	0.00
No	20 (45.5%)	132 (76.3%)	
Smoking			
Yes	12 (27.3%)	14 (8.1%)	0.00
No	32 (72.7%)	159 (91.9%)	
Physical Activity			

Active	10 (22.7%)	129 (74.6%)	0.00
Sedentary	34 (77.3%)	44 (25.4%)	

Higher prevalence of MA was observed in >5 years (77.3%) and lower in <5 years (22.7%) of diabetics period, which was significantly associated with each other (p=<0.001). Similarly, there was significant association of MA with hypertension (p=<0.001); yes (54.5%) and no (45.5%), smoking (p=<0.001); yes (27.3%) and no (72.7%), and physical activity (p=<0.001); active (22.7%) and sedentary (77.3%)(Table 2).

DISCUSSION

The risk factors found associated for microalbuminuria were raised blood pressure and poor glycemicontrol. Some studies have revealed duration of diabetes, male sex, and pre-existing retinopathyas major risk factors for microalbuminuria.⁴ According to this model, blood pressure, duration of diabetes, BMI, smoking status and physical activity directly correlates with micro albuminuria.

In this study the prevalence rate of urinary microalbuminuria among diabetes patients was found to be 20.3%. which is quite higher than various epidemiological and cross-sectional studies of Afkhami-Ardekani et al (2008), Huraib et al (1995) and Vijay et al (1994) 14%, 16.8%, 18.7%), respectively.¹⁶⁻¹⁸ However in some studies conducted in India and Nepal higher prevalence of microalbuminuria also reported as 36.3%, 45.5% respectively.¹⁹⁻²⁰ There was not any significant association with microalbuminurea and age (p=0.517) and sex (p=0.554). Similarly in the other studies also showed not statistically significant.^{17,18,20} In contrast Varghese et al., and Chowta et al., reported a statistically significant correlation with age.^{19,21}

In the present study, a good statistical significant correlation was found between appearance of microalbuminuria and body mass index of the patient with the p= 0.029. Similar findings were observed in a study conducted by Gall et al (1991) who reports higher BMI had higher albumin excretion.²² A study conducted by Ko GT et al (1999) in China had also shown a correlation between obesity and appearance of MA in diabetics.⁶ However, in many other studies showed no correlation with MA explaining poorly controlled diabetes induces weight loss and these patients with low body mass index are at higher risk for diabetic complications and microalbuminuria.¹⁸ However, our findings can be supported with the fact that obesity is also associated with an increase in insulin resistance that further raises

the blood glucose level which in turn hastens glomerular arteriolar sclerosis promoting early onset MA. In the study similar to this study of Gupta et al (1991) reported high levels of HbA1c was associated with microalbuminuria.²³ Duration of diabetes has significant contribution for the development microalbuminuria by prolonged exposure to hyperglycemia-induced advanced glycosylation end products accumulations.²¹ Present study has shown positive correlation of microalbuminuria with duration (>5 years) of diabetes mellitus ($p < 0.001$) which is in accordance with many previous reports.¹⁷⁻¹⁸ Similar findings have been observed with Idowu et al., and Varghese et al., who reported that the duration of diabetes and retinopathy were the major predictors of microalbuminuria.^{19,24}

Blood pressure is one of the most important risk factor of micro albuminuria. High blood pressure may cause microalbuminurea by increasing glomerular filtration pressure and subsequent renal damage.²⁵ In this study, High blood pressure was statistically significant with microalbuminurea ($p = 0.00$). Similarly, many studies have shown a positive correlation between these two variables.^{6,16-19,23}

In established diabetic nephropathy, the rate of progression to renal failure amongst smokers is twice in both Type I and Type II diabetes.²⁶

This study has also shown a positive correlation of MA with physical inactivity. Physical activity improves endothelial function and as we know endothelial dysfunction in the renal vasculature is associated with albuminuria.²⁷ This association held for both walking and strenuous activity. The patients who were physically active were negative for MA than those with a sedentary lifestyle which can in turn be supported by the positive correlation between BMI and MA in this study itself. Similar findings were reported by study conducted by where negative MA was associated with patients who were involved in aerobic exercise for 6 months without any change in variable.²⁸

CONCLUSIONS

There is a high prevalence rate 20.3% of urinary microalbuminuria among diabetes patients of Nepal. There are several risk factors/correlates associated with it. In this study, the most important risk factors found are obesity, 5 or more year duration of disease, smoking, hypertension and sedentary lifestyles. Early recognition and institution of risk control strategy along with early start of ARBs or ACEIs in these patients would halt the progression of diabetic nephropathy.

ACKNOWLEDGEMENTS

We would like to thank all the staff member of Bir Hospital, Annapurna Neurological Institutes and allied Sciences, Global hospital, Kathmandu Nepal and we would also like to thank the patients and their family for cooperation in this study.

REFERENCE

1. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes care*. 1998;21(9):1414-31. [\[PubMed\]](#)
2. Zolotov S, Ben Yosef D, Rishie ND, Yesha Y, Karnieli E. Metabolic profiling in personalized medicine: bridging the gap between knowledge and clinical practice in Type 2 diabetes. *Personalized Medicine*. 2011;8(4):445-56. [\[Full Text\]](#)
3. Franco OH, Steyerberg EW, Hu FB, Mackenbach J, Nusselder W. Associations of diabetes mellitus with total life expectancy and life expectancy with and without cardiovascular disease. *Arch Intern Med*. 2007;167(11):1145-51. [\[PubMed\]](#)
4. Organization WH. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia: report of a WHO/IDF consultation. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia: report of a WHO/IDF consultation 2006. Available At. www.who.int/diabetes/publications/diagnosis_diabetes2006/en
5. Association AD. Standards of medical care in diabetes—2015 abridged for primary care providers. *Clin Diabetes*. 2015;33(2):97. [\[PubMed\]](#)
6. Ko GT, Chan JC, Lau M, Cockram CS. Diabetic microangiopathic complications in young Chinese diabetic patients: a clinic-based cross-sectional study. *Journal of diabetes and its complications*. 1999;13(5-6):300-6. [\[PubMed\]](#)
7. Battisti WP, Palmisano J, Keane WE. Dyslipidemia in patients with type 2 diabetes. relationships between lipids, kidney disease and cardiovascular disease. *Clin Chem Lab Med*. 2003;41(9):1174-81. [\[PubMed\]](#)
8. Lutale JJK, Thordarson H, Abbas ZG, Vetvik K. Microalbuminuria among type 1 and type 2 diabetic patients of African origin in Dar Es Salaam, Tanzania. *BMC Nephrology*. 2007;8(1):2. [\[PubMed\]](#)
9. Ahmad T, Ullhaq I, Mawani M, Islam N. Microalbuminuria in Type-2 Diabetes Mellitus; the tip of iceberg of diabetic complications. *Pakistan Journal of Medical Sciences*.

- 2017;33(3):519.[\[PubMed\]](#)
10. Lau AC-W, Lo MK-W, Leung GT-C, Choi FP-T, Yam LY-C, Wasserman K. Altered exercise gas exchange as related to microalbuminuria in type 2 diabetic patients. *Chest*. 2004;125(4):1292-8.[\[PubMed\]](#)
 11. de Zeeuw D, Remuzzi G, Parving HH, Keane WF, Zhang Z, Shahinfar S, et al. Proteinuria, a target for renoprotection in patients with type 2 diabetic nephropathy: lessons from RENAAL. *Kidney Int*. 2004;65(6):2309-20.[\[PubMed\]](#)
 12. Persson F, Lindhardt M, Rossing P, Parving HH. Prevention of microalbuminuria using early intervention with renin-angiotensin system inhibitors in patients with type 2 diabetes: A systematic review. *J Renin Angiotensin Aldosterone Syst*. 2016;17(3).[\[PubMed\]](#)
 13. Roshan B, Stanton RC. A story of microalbuminuria and diabetic nephropathy. *J Nephropathol*. 2013;2(4):234.[\[PubMed\]](#)
 14. Bakris GL, Molitch M. Microalbuminuria as a risk predictor in diabetes: the continuing saga. *Diabetes Care*. 2014;37(3):867-75.[\[PubMed\]](#)
 15. AlFehaid AA. Prevalence of microalbuminuria and its correlates among diabetic patients attending diabetic clinic at National Guard Hospital in Alhasa. *J Fam Community Med*. 2017;24(1):1-5.
 16. Vijay V, Snehalatha C, Ramachandran A, Viswanathan M. Prevalence of proteinuria in non-insulin dependent diabetes. *The Journal of the Association of Physicians of India*. 1994;42(10):792-4.[europepmc.org/abstract/MED/7876049](#)
 17. Huraib S, Abu-Aisha H, Sulimani RA, Famuyiwa FO, Al-Wakeel J, Askar A, et al. The pattern of diabetic nephropathy among Saudi patients with noninsulin-dependent diabetes mellitus. *Ann Saudi Med*. 1995;15(2):120-4.[\[PubMed\]](#)
 18. Afkhami-Ardekani M, Modarresi M, Amirchaghmaghi E. Prevalence of microalbuminuria and its risk factors in type 2 diabetic patients. *Indian J Nephrol*. 2008;18(3):112.[\[PubMed\]](#)
 19. Varghese A, Deepa R, Rema M, Mohan V. Prevalence of microalbuminuria in type 2 diabetes mellitus at a diabetes centre in southern India. *Postgrad Med J*. 2001;77(908):399-402.[\[PubMed\]](#)
 20. Sigdel M, Rajbhandari N, Basnet S, Nagila A, Basnet P, Tamrakar B. Microalbuminuria among type-2 diabetes mellitus patients in Pokhara, Nepal. *Nepal Med Coll J*. 2008;10(4):242-5.[\[Full Text Link\]](#)
 21. Chowta NK, Pant P, Chowta MN. Microalbuminuria in diabetes mellitus: Association with age, sex, weight, and creatinine clearance. *Indian J Nephrol*. 2009;19(2):53-6.[\[PubMed\]](#)
 22. Gall MA, Rossing P, Skott P, Damsbo P, Vaag A, Bech K, et al. Prevalence of micro- and macroalbuminuria, arterial hypertension, retinopathy and large vessel disease in European type 2 (non-insulin-dependent) diabetic patients. *Diabetologia*. 1991;34(9):655-61.[\[PubMed\]](#)
 23. Gupta D, Verma L, Khosla P, Dash S. The prevalence of microalbuminuria in diabetes: a study from north India. *Diabetes Res Clin Pract*. 1991;12(2):125-8.[\[PubMed\]](#)
 24. Idowu AA, Ajose AO, Adedeji AT, Adegoke AO, Jimoh KA. Microalbuminuria, Other Markers of Nephropathy and Biochemical Derangements in Type 2 Diabetes Mellitus: Relationships and Determinants. *Ghana Medical Journal*. 2017;51(2):56-63.[\[PubMed\]](#)
 25. Koroshi A. Microalbuminuria, is it so important? *Hippokratia*. 2007;11(3):105-7.[\[PubMed\]](#)
 26. Biesenbach G, Janko O, Zazgornik J. Similar rate of progression in the predialysis phase in type I and type II diabetes mellitus. *Nephrol Dial Transplant*. 1994;9(8):1097-102.[\[PubMed\]](#)
 27. Robinson ES, Fisher ND, Forman JP, Curhan GC. Physical Activity and Albuminuria. *Am J Epidemiol*. 2010;171(5):515-21.[\[PubMed\]](#)
 28. Lazarevic G, Antic S, Vlahovic P, Djordjevic V, Zvezdanovic L, Stefanovic V. Effects of aerobic exercise on microalbuminuria and enzymuria in type 2 diabetic patients. *Renal Failure*. 2007;29(2):199-205.[\[PubMed\]](#)