

## Microbial diversity and methanogenic potential in a high temperature natural gas field in Japan

Hanako Mochimaru · Hideyoshi Yoshioka · Hideyuki Tamaki ·  
Kohei Nakamura · Nobuyuki Kaneko · Susumu Sakata · Hiroyuki Imachi ·  
Yuji Sekiguchi · Hiroo Uchiyama · Yoichi Kamagata

Published online: 24 May 2008  
© Springer 2008

**Erratum to: Extremophiles (2007) 11:453–461**  
**DOI 10.1007/s00792-006-0056-8**

The unit of acetate concentration in Table 1 and the reference in the text was shown incorrectly. The unit should be  $\text{mmol l}^{-1}$ . This does not affect any of our conclusions.

---

The online version of the original article can be found under doi:[10.1007/s00792-006-0056-8](https://doi.org/10.1007/s00792-006-0056-8).

---

H. Mochimaru · H. Uchiyama  
Graduate School of Life and Environmental Sciences,  
University of Tsukuba, Tsukuba,  
Ibaraki 305-8572, Japan

H. Mochimaru · H. Tamaki · K. Nakamura ·  
Y. Sekiguchi · Y. Kamagata (✉)  
Institute for Biological Resources and Functions,  
National Institute of Advanced Industrial Science  
and Technology (AIST), 1-1-1 Higashi, Tsukuba,  
Ibaraki 305-8566, Japan  
e-mail: y.kamagata@aist.go.jp

Y. Kamagata  
Research Institute of Genome-Based Biofactory,  
National Institute of Advanced Industrial Science  
and Technology (AIST), Sapporo, Hokkaido 062-8517, Japan

H. Yoshioka · N. Kaneko · S. Sakata  
Institute for Geo-resources and Environment,  
National Institute of Advanced Industrial Science  
and Technology (AIST), Tsukuba, Ibaraki 305-8566, Japan

H. Imachi  
Subground Animalcule Retrieval (SUGAR) Program,  
Extremobiosphere Research Center,  
Japan Agency for Marine-Earth Science & Technology  
(JAMSTEC), Yokosuka, Kanagawa 237-0061, Japan