

MICROBIAL ECOLOGY OF METHANE EMISSION IN RICE AGROECOSYSTEM: A REVIEW

S. K. DUBEY

e-mail: dskbot@yahoo.com

Centre of Advanced Study, Department of Botany, Banaras Hindu University, Varanasi, India

(Received 9th April 2005, accepted 28th June 2005)

Abstract. Methane has profound impact on the physico-chemical properties in atmosphere leading to global climate change. Out of the various sources of CH₄, rice fields are the most significant contributors. The processes involved in the emission of CH₄ from rice fields to the atmosphere include CH₄ production (methanogenesis) in the soil by methanogens, methane oxidation (methanotrophy) by methanotrophs and vertical transfer of CH₄ via plant transport and diffusion or ebullition. In the overall methane dynamics rice plants act as : a) source of methanogenic substrate, b) conduit for CH₄ through well developed system of inter cellular air space (aerenchyma), and c) potential methane oxidizing micro-habitat in the rhizosphere by diffusing oxygen which favour the growth and multiplication of methanotrophs. Apart from mechanistic uncertainties, there are several other uncertainties in the estimation of CH₄ flux. Methane dynamics in the paddy field is controlled by a complex set of parameters linking the biological and physical characteristics of soil environment like temperature, carbon source, Eh, pH, soil microbes and properties of rice plants, etc. It has now become possible to isolate, detect and characterize the methanogens and methanotrophs by using molecular biological tools like PCR, FISH, etc. techniques. The apparent half saturation constant (K_m) and maximum oxidation rate (V_{max}) are distinctive parameters which determine the ability of bacteria to survive on atmospheric methane.

Keywords. *methane; methanotrophs; methanotrophy; methanogens; molecular tools*

Introduction

Atmospheric methane (CH₄) is a potent greenhouse gas with high absorption potential for infrared radiation. Methane is present at about 1.8 ppmV in the atmosphere [135]. During the last 20 years, its concentration has been increasing, on an average at the rate of 0.8% y⁻¹ [125]. Due to this, CH₄ is of great concern as a greenhouse gas. Although the tropospheric CH₄ concentration is very low as compared to CO₂ (357 ppmV), methane accounts for 15 to 20% of global warming [71]. The total annual source strength of all methane emissions from anthropogenic origin is estimated to be 550 Tg [133]. Major sources of this input include natural wetlands, rice fields, enteric fermentation in animals, termites and land fills. The contribution from rice cultivation is estimated to range from 20-100 Tg CH₄ y⁻¹ with an average of 60 Tg CH₄ y⁻¹ [71]. The biogenic methane is mostly produced by methanogenic archaea (methanogens) in anaerobic environments i.e. sediments and flooded rice fields [59]. Each year methanogens produce about 400 million metric tons of CH₄ [49]. Recent studies have shown that methane is not only produced in anoxic rice fields soil but also directly from the roots of rice plants which are inhabited by a methanogenic community different from that in the rice field soil [92].

According to a current estimate, rice production will need to expand by around 70% over the next 25 years to support the growing human population [39]. For this, intensified global fertilizer application will be essential, and this will exacerbate the methane problem. It is projected that the methane emission from rice cultivation may increase from the 1990 level of 97 Tg y^{-1} to 145 Tg y^{-1} by 2025 [5]. India is an important rice producing country, comprising 28.6% of world rice cultivated area [65]. During recent years, studies on methane emission from Indian rice fields have focused on the influence of soil type, season, water regime, organic inputs, fertilizers, rice cultivars and agrochemicals [111]. Using the baseline scenario, annual methane emissions for China, India, Indonesia, Philippines and Thailand were calculated to be 3.73, 2.14, 1.65, 0.14 and 0.18 Tg $CH_4 y^{-1}$, respectively [106].

Chemical and biological processes consume methane in the global methane cycle. The only known biological sink for atmospheric methane is its oxidation in aerobic soils by methanotrophic bacteria, this may contribute up to 10-20% to the total methane destruction [128], or between 15 and 45 Tg $CH_4 y^{-1}$ [71]. Methanotrophs oxidize CH_4 with the help of methane monooxygenase (MMO) enzyme. These bacteria are classified into three groups (Type-I, Type-II and Type -X) based on the pathways used for assimilation of formaldehyde and other physiological and morphological features [58]. An enormous effort is being made worldwide by microbial ecologists to isolate, detect and characterize methanotrophs and methanogens in different rice ecosystems by using molecular biological tools and techniques [18, 46, 57]. Methanotrophy is an aerobic process [52], but in marine sediments and in some saline inland waters it could be anaerobic [36]. The apparent half saturation constant (K_m), and maximum rate (V_{max}) of CH_4 oxidation are characteristic parameters, which determine the ability of methanotrophs to grow on atmospheric CH_4 [31]. Several workers have reported that methane oxidation occurs in rice microcosm [56], wetland rice [40, 158] and dryland rice fields [44, 45]. Methane oxidation in rhizospheric soil is considered as an important sink for CH_4 [44, 45, 56]. Therefore, the knowledge of several environmental factors (e.g. temperature, fertilizer inputs, crop phenology and soil moisture) that can provide feedback on the capacity of soil to oxidize atmospheric CH_4 , may have significant consequences on the global atmospheric CH_4 budget.

This review presents an overview of the underlying microbial basis for production, oxidation and emission of methane in paddy fields under the influence of several environmental factors. Molecular ecological approaches for the isolation, detection and characterization of methanogens and methanotrophs are also described.

Production of methane

Methanogens

Methanogens are strictly anaerobic unicellular organisms originally thought to be bacteria but now recognized as belonging to a separate phylogenetic domain, the *archae* [53]. Phenotypic characteristics of methanogenic bacteria are listed in Table 1. 16S rRNA analysis suggested that methanogens can be categorized under three groups. Group I comprises *Methanobacterium* and *Methanobrevibacter*, Group II contains *Methanococcus*, and Group III comprises the genera including *Methanospirillum* and *Methanosarcina* [53]. They proliferate in anaerobic fresh water environments, such as sediments and the digestive tract of animals [147].

Table 1. Characteristics of methanogenic and methanotrophic bacteria (Source: [53, 58, 74, 101]).

Characteristics	Methanogens	Methanotrophs
Cell form	rods, cocci, spirilla, filamentous, sarcina	rods, cocci, vibrios
Gram stain reaction	Gram +/-	Gram -
Classification	Archaeobacteria	Eubacteria
Cell wall	pseudomureine, protein, heteropolysaccharide	peptidoglycon
Metabolism	anaerobic	aerobic
Energy and carbon source	H ₂ +CO ₂ ; H ₂ +methanol; formate; methylamines; methanol, acetate	methane; methanol; dimethyl-ether, methyl formate, dimethyl carbonate
Catabolic products	CH ₄ or CH ₄ +CO ₂	CO ₂
TCA cycle	Incomplete	Incomplete (Type-I) or complete (Type-II)
Carbon assimilation pathways	TCA cycle, gluconeogenesis	ribulose monophosphate pathways (Type-I) or serine pathways (Type-II)
Resting cells	-	cysts (Type-I) or exospore (Type-II)
GC content mol%	26-60	50-62.5
Typical species	<i>Methanobacterium bryanthii</i> <i>Methanobrevibacter smithii</i> <i>Methanomicrobium mobile</i> <i>Methanogenium cariaci</i>	<i>Methylosinus trichosporium</i> <i>Methylomonas methanica</i> <i>Methylocystis minimus</i> <i>Methylobacter albus</i>

In these habitats, methanogens play an important role in the degradation of complex organic compounds. Most methanogens are mesophilic, able to function in temperature ranging from 20 to 40⁰C [147]. They are also found in extreme environments like hydrothermal vents where they thrive at temperatures above 100⁰C. Methanogens mainly use acetate (contributes about 80% to CH₄ production) as a carbon substrate but other substrate like H₂/CO₂ and formats also contribute 10-30% to CH₄ production [27]. All methanogens use NH₄⁺ as a nitrogen source, although the ability to fix molecular nitrogen and the *nif* gene is present in all the three orders (*Methanobacteriales*, *Methanococcales* and *Methanomicrobiales*) of methanogens [120].

Methanogenesis

Methane is produced in the anaerobic layers of paddy soil by bacterial decomposition of organic matter [39]. The organic matter converted to CH₄ is derived mainly from plant-borne material, and organic manure [35], if applied. The anaerobic degradation of organic matter involves four main steps: a) hydrolysis of polymers by hydrolytic organisms, b) acid formation from simple organic compound by fermentative bacteria, c) acetate formation from metabolites of fermentations by homoacetogenic or syntrophic bacteria, and d) CH₄ formation from H₂/CO₂, acetate, simple methylated

compounds or alcohols and CO₂ [163]. CH₄ is produced in rice fields after the sequential reduction of O₂, nitrate, manganese, iron and sulphate, which serve as electron acceptors for oxidation of organic matter to CO₂ [164]. Methanogenesis from all substrates requires a number of unique coenzymes, some of which are exclusively found in methanogens [98]. At least nine methanogen-specific enzymes are involved in the pathway of methane formation from H₂ and CO₂ [140]. In paddy soil, acetate and H₂ are the two main intermediate precursors for CH₄ formation [162].

Methanogenesis from H₂/CO₂

The first step of the pathway comprises the binding of CO₂ to methanofuran (MFR) and its H₂ dependent reduction to formyl-MF (first stable intermediate compound of the pathway). Formation of this complex is catalyzed by *formylmethanofuran dehydrogenase*. Further, the formyl moiety of the formyl-MF is subsequently transferred to tetrahydromethanopterin (H₄MPT), which with H₄MPT *formyl transferase*, forms methenyl-H₄MPT, while reduces to methylene-H₄MPT and then to methyl-H₄MPT. In both reactions, reduced coenzyme F₄₂₀ serves as the reductant. The F₄₂₀-dependent reduction of methenyl-H₄MPT is reversible and is catalyzed by *methylene H₄MPT dehydrogenase*. The reversible F₄₂₀ H₂-dependent reduction of methylene - H₄MPT to methyl-H₄MPT is catalyzed by *methylene-H₄MPT reductase*. In the next step of CO₂ reduction pathway, the methyl group is transferred from N-5-methyl H₄MPT to coenzyme M (2-mercaptoethane sulfonate) giving rise to methyl coenzyme M(methyl-CoM). The reduction of methyl CoM is catalysed by the methyl-CoM-reductase. This reaction involves two unique coenzymes. The first one is N-7-mercaptoheptanoylthreonine phosphate (HS-HTP) which acts as electron donor in the reduction of methyl CoM giving rise to CH₄ and mixed disulfide of HS-CoM and HS-HTP (CoM-S-S-HTP). The second coenzyme involved in this reaction is factor F₄₃₀ (a nickel porphyrinoid) which is the characteristic prosthetic group of the methyl reductase [16, 74, 140].

Methanogenesis from acetate

Methanogenesis from acetate starts with its activation to acetyl-CoA. *Methanosarcina* and *Methanothrix* use different ways of acetate activation. The former organism takes advantage of *acetate kinase* and *phosphotrans acetylase* whereas the later makes use of *acetyl-CoA synthetase*. All three enzymes are soluble and oxygen insensitive. Further breakdown of *acetyl-CoA* catalyzes the cleavage of *acetyl-CoA*, giving rise to a methyl group, a carbonyl group and CoA, all of which are transiently bound to the enzyme. In a further step, the *Co-dehydrogenase* complex catalyzes the oxidation of the carbonyl group. The CO₂ is thereby formed and CoA is released from the enzyme, where the methyl group is transferred to a corrinoid-Fe-S protein. This complex catalyzes not only the cleavage of *acetyl-CoA* and oxidation of the carbonyl group but in addition, the transfer of the methyl moiety to H₄MPT. Further pathway from methyl H₄MPT to CH₄ takes advantage of the pattern similar to that discussed for the utilization of the CO₂/H₂ [16, 98].

Thermodynamics of CH₄ production

The process of methane production in paddy fields is thermodynamically exergonic [163]. The Gibbs free energy (ΔG) for the process is mainly a function of the acetate concentration and H₂-partial pressure. A pre-requisite for early methane production

seems to be a sufficiently high H_2 -partial pressure that corresponds to ΔG of H_2 dependent methanogenesis (hydrogenotrophic) of less than about $-23 \text{ kJ mol}^{-1} \text{ CH}_4$ [162]. The time until the onset of CH_4 production and the magnitude of production is a function of the quantity of easily degradable organic matter, reducible Fe(III) and sulfate [162]. Methanogens are energetically limited by availability of their substrates H_2 and acetate as long as iron or sulfate reducers are able to compete for them [1]. The methanogens have to compete for available substrates with other anaerobic bacteria, namely the nitrate, manganese, ferric iron and sulfate reducers. The competition for carbon substrates in general follows thermodynamic rules: nitrate reducers outcompete the other anaerobic bacteria for the substrates [149]. Several studies have reported different ΔG values for methanogenesis in various paddy fields. For Italian paddy fields the values of ΔG for methane production were found to be -31.6 to $34.8 \text{ kJ mol}^{-1} \text{ CH}_4$ [1] and -24 to $-38 \text{ kJ mol}^{-1} \text{ CH}_4$ [27]. Peters and Conrad [123] found that ΔG ranged from -25 to $-50 \text{ kJ mol}^{-1} \text{ CH}_4$ for German rice fields. Cultures of *Methanobacterium bryantii* required ΔG values of less than $-30 \text{ kJ mol}^{-1} \text{ CH}_4$ for CH_4 production [31]. Kral et al. [86], reported that for cultures of other methanogens ΔG values varied between -32 and $-60 \text{ kJ mol}^{-1} \text{ CH}_4$ for hydrogenotrophic methanogenesis. Yao and Conrad [162] calculated the ΔG for acetate dependent methanogenesis (acetoclastic) and concluded that unlike H_2 -dependent methanogenesis acetate based process was apparently not under thermodynamic control. They found that CH_4 production was less at ΔG equal to $-26 \text{ kJ mol}^{-1} \text{ CH}_4$ and was the average at ΔG equal to $-29 \text{ kJ mol}^{-1} \text{ CH}_4$. Roy et al. [130] have suggested that early CH_4 production is due to H_2 -dependent methanogenesis and that acetate-dependent methanogenesis only starts later when sulfate and Fe(III) have been reduced in paddy soils. Further, methane production from H_2/CO_2 is not started before fermentation has increased the H_2 partial pressure to an amply high value to allow exergonic production of CH_4 at a ΔG of less than about $-26 \text{ kJ mol}^{-1} \text{ CH}_4$ [130, 162].

Methanogenic population and CH_4 emission

The mechanism of methanogenesis in paddy fields worldwide has been investigated in detail. However, information regarding methanogenic population size in rice fields is limited. Rajagopal et al. [126] were the first to carry out isolation and characterization of methanogens from Louisiana paddy fields and reported the presence of two *Methanobacterium*-like strains and two *Methanosarcina*-like strains. Joulain et al. [76] recorded the methanogenic populations from the paddy fields of France, the Philippines, and USA. Their results of the classic counts of methanogenes, and strains isolated and identified by 16S rRNA gene sequencing, suggested the dominance of *Methanobacterium spp.* and *Methano-sarcina spp.* among the culturable organisms. Reichardt et al. [129] revealed that methanogens were abundant in root extracts of mature rice plants. Methanogens are also exist on the rhizoplane of the rice plants [92]. Fetzer et al. [50] isolated four genera (*Methanobacterium*, *Methano-sarcin*, *Methanobrevibacter* and *Methanoculleus*) from Italian rice fields. According to Asakawa et al. [6] there are only two strains (*Methanobrevibacter arboriphilus* and *Methanosarcina mazeii*) of methanogens in rice fields which have been identified to the species level. Adachi [2], isolated *Methanobacterium* and *Methanobrevibacter spp.* from subtropical Japanese rice fields. Kudo et al. [87] reported the presence of *Methanosarcina*, *Methanogenium*, *Methanosaeta* and *Methanoculleus*-like organisms in rice paddy fields of Japan by using a molecular retrieval approach with archaeal small

subunit (SSU) rRNA encoding gene (rDNA) sequences. Results of investigations on methanogenic population size in various types of rice fields are summarized in Table 2.

Table 2. Population size of methanogens and methanotrophs in various rice fields.

Ecozones/location	Methanogens	References	Methanotrophs	References
Rice fields (Australia, France, Philippines, USA, Trinidad)	$<10^1$ – 2.3×10^6 cells g^{-1} dw	[75]	1.5×10^3 – 3.5×10^7 cells g^{-1} dw	[75]
Rice fields (Zhejiang, China)	4.6×10^3 – 1.3×10^7 CFU g^{-1} dw	[108]	3.0×10^6 – 2.3×10^8 CFU g^{-1} dw	[108]
Rice fields (Beijing, China)	1.4×10^5 – 2.3×10^5 g^{-1} dw	[64]	-	-
Rice fields (Germany)	-	-	3.7×10^4 – 3.8×10^7 bacteria g^{-1} dw	[13]
Rice fields (Cuttack, India)	0.2 – 2.8×10^5 MPN g^{-1} soil	[90]	4.2×10^4 – 5.2×10^6 CFU g^{-1} dw	[89]
Rice fields (Cuttack, India)	2.4 – 6.1×10^6 MPN g^{-1} dw	[88]	-	-
Rice fields (Cuttack, India)	5.4 – 7.5×10^3 cells g^{-1} dw	[15]	-	-
Dryland Rice fields (Varanasi, India)	-	-	5.3×10^6 – 7.4×10^7 cells g^{-1} soil	[42]
Wetland Rice fields (Varanasi, India)	-	-	2.6×10^6 – 5.0×10^7 cells g^{-1} soil	[43]
Rice fields (Northern Italy)	-	-	4.2×10^6 – 2.3×10^7 cells g^{-1} dw	[10]
Rice fields (Northern Italy)	-	-	4.0×10^5 – 2.0×10^8 cells g^{-1} dw	[56]
Rice fields (Japan)	5.2×10^4 – 1.1×10^6 cells g^{-1} dw	[151]	1×10^4 – 1×10^7 CFU g^{-1} dw	[157]

Table 3. Variation of estimates of global methane emission ($Tg CH_4 y^{-1}$) from rice fields during 1963-1998.

CH_4 emission $Tg CH_4 y^{-1}$	References
190	[85]
280	[48]
59	[28]
95	[78]
35-59	[138]
120-200	[32]
70-170	[62]
142-190	[14]
47-145	[136]
25-60	[116]

Table 3. continued from page 6.

CH₄ emission Tg CH₄ y⁻¹	References
20-100	[68]
66	[79]
50	[114]
20-150	[124]
60-105	[70]
20-100	[71]
25-54	[132]
30-50	[117]
50-80	[93]

The first measurements of CH₄ emission from paddy fields were conducted in California by Cicerone and Shetter [28]. This was followed by extensive studies in Spain [138], China [26], USA [133], Japan [157], Philippines [115], and India [110, 139], but the estimates of methane emissions vary widely, e.g., from 18.0 to 27.1 mg CH₄ m⁻² h⁻¹ in Indonesia [119], 19.5 to 32.2 mg CH₄ m⁻² h⁻¹ in Thailand [160] and 19 to 79 mg CH₄ m⁻² h⁻¹ in Philippines; CH₄ emission from temperate rice fields is relatively lower (6.67 and 18.25 mg CH₄ m⁻² h⁻¹) [136]. These field experiments varied widely in the methodologies (closed chamber with automatic or manual sampling devices), sampling frequencies (continuous or sporadic sampling), observation periods (one season or several consecutive years) and sampling field designs (randomized plots or single fields). Recent estimates indicated that methane release (m²y⁻¹) from different rice ecosystems follows the order deepwater > irrigated > rainfed rice [115]. The distinction among irrigated, rainfed, and deepwater rice fields is a common feature of the available statistics of rice cultivated area. A specific assessment of these ecosystems will, therefore, directly improve the accuracy of regional and global estimates of the methane source strength [67]. The estimates on global CH₄ emission from paddy fields, however, have varied over time period, mainly depending on approach, technique and database used for extrapolation. Table 3 shows reported estimates of global methane emission rates from various rice fields during 1963-1998. These have varied from as low as 12 Tg CH₄ y⁻¹ to as high as 200 Tg CH₄ y⁻¹. This is largely due to the lack of uniformity in the methods used for collecting field data, location of experimental sites and seasonal variations. This problem was addressed by Khalil and Shearer [79] who developed an inventory of direct flux measurements from a number of studies and modified the information from Matthews et al. [105] on the duration of growing seasons to estimate global and regional annual methane emission rates. They arrived at a figure for the global CH₄ emission rate of 66 Tg CH₄ y⁻¹.

Pathways of methane emission

The net amount of CH₄ emitted from soil to the atmosphere is the balance of two opposite processes - production and oxidation. Methane, the product of methanogenesis, escapes to the atmosphere from soil via aerobic interfaces where CH₄ oxidation takes place. There are three pathways of CH₄-transport into the atmosphere – molecular diffusion, ebullition and plant transport (Fig. 1).

In the temperate rice fields more than 90% of the CH₄ is emitted through plant transport [136] while in the tropical rice fields, significant amounts of CH₄ may evolve

by ebullition (gas transport via gas bubbles) in particular during the early period of the season and in the case of high organic input [38].

Ebullition is also the common and significant mechanism of CH_4 flux in natural wetlands [155]. According to Sass et al. [131], ebullition can play significant role in CH_4 transport under high organic fertilization. If soil is unvegetated or plant aerenchyma is not yet well-developed, ebullition plays a major role in CH_4 emission [22] but it occurs only at surface layer and its rate is regulated by CH_4 concentration, temperature, soil porosity and plant aerenchyma [94].




Figure 1. Conceptual schematic diagram of methane production, oxidation and emission from paddy field.

Diffusion of CH_4 across the flooded soil and overlying water of the rice field to the atmosphere is a function of surface-water concentration of CH_4 , wind speed and CH_4 supply to the surface water [137]. CH_4 diffusion through the soil is a very slow process because the 'diffusion rate' of gaseous CH_4 is very low in liquid phase (about 104 times

slower than diffusion through the gas phase), therefore, it hardly contributes to the total CH₄ flux [8].

Plant mediated transport is the primary mechanism for the CH₄ emission from paddy fields, and contributes 60-90% to the total CH₄ flux [154]. Methane in the soil-water surrounding the roots dissolves into the surface-water of the roots, diffuses into the cell-wall water of root epidermis cells, and then diffuses through the cell-wall water of the root-cortex, depending upon the concentration gradient between the soil-water surrounding the roots and the lysigenous inter-cellular spaces in the roots [118]. Methane is then gasified in the root cortex and transported to the shoots via lysigenous intercellular spaces and aerenchyma. Eventually, CH₄ is released primarily through the micropores in the leaf sheath of the lower leaf position and also through the stomata in the leaf blade [118].

Factors affecting methane emission

Methane emission from paddy fields is controlled by a complex set of parameters linking the physical and biological characteristics of soil environments with specific agricultural practices. Methane production depends on the soil organic carbon content and quality, texture, Eh, pH, Fe content, sulfate content and salinity and application of fertilizers, etc.

Soil pH. Eh and texture

Methane production in flooded rice soils is very sensitive to pH with an optimum range between 6.7 and 7.1 [152]. Effect of soil pH on CH₄ production varied by about two orders of magnitude in four different Indian soils but was found to be maximum at pH around 8.2 [121].

Yagi and Minami [161] reported that values of redox potential (Eh) varied from -100 to -200 mV for the initiation of CH₄ production in paddy soils. Masscheleyn et al. [104] incubated rice soil under controlled redox levels ranging between -250 and +500 mV. They found the threshold for methane production to be -150 mV. Some suggested that soils containing greater amounts of readily decomposable organic substrates (acetate, formate, methanol, methylated amines, etc.) and low amounts of electron acceptors (Fe³⁺, Mn⁴⁺, NO₃⁻, SO₄²⁻) are likely to show high production of CH₄. According to sequential oxidation-reduction order, molecular O₂ is the first to be reduced at an Eh of about +30 mV followed by NO₃⁻ and Mn⁴⁺ at 250 mV, Fe³⁺ at +125 mV and SO₄²⁻ at -150 mV (Patrick, 1981). Subsequent to SO₄²⁻ reduction, methanogens will start producing methane [8].

As texture determines various physico-chemical properties of soil, it could influence CH₄ production indirectly. Jackel et al. [72] found that rates of CH₄ production increased when the aggregate size of the soil increased. A negative co-relationship between CH₄ emission and clay content was reported by Sass and Fisher [133]. Seasonal CH₄ emissions indicated a negative relationship to clay content for Texan paddy soils [133].

Temperature

Methane emission is much more responsive to temperature. Temperature not only has an effect on methane production itself but also has an effect on the decomposition of

organic materials from which the methanogenic substrates are produced [27]. The influence of temperature on CH₄ production rates has been reported for several rice ecosystems [121, 145]. Wassman et al. [156] observed a faster development of CH₄ production rate and higher maximum value with increasing temperatures between 25 and 35°C. Hattori et al. [59] recorded optimum temperature of 40°C for CH₄ production in Japanese paddy fields due to dominance of methanogenic population at this temperature.

Growth period and crop phenology

Wassmann et al. [154] recorded lower CH₄ fluxes in the early growth period of rice plant, which increased gradually during mid to late season and dropped to very low level before or after harvest. Jermawatdipong et al. [73] found that more than 50% of CH₄ was emitted in the first half of the growth period in Thailand rice fields, while CH₄ emissions in Japanese rice fields occurred mainly in the second-half of the growth period [81]. Jermawatdipong et al. [73] argued that the high temperatures from the beginning of rice growth in the tropics caused the main decomposition stage of soil and applied organic materials to shift to early growth stage which resulted in active CH₄ production from the very beginning of rice growth. Seiler [138] observed maximum CH₄ emission at the end of heading and flowering stage of rice plants in Spain.

Flowering period is generally considered as the peak period for methane emission. The peak emission value remains for a period of 10-15 days in the crop duration of 90-100 days. According to Holzapfel-Pschorn et al. [62] this period emits 90% of the total methane during the whole crop season, because the biomass of rice crop increases gradually, reaching the maximum weight by flowering. Up to 50 % of the total methane emission from rice fields can be due to root exudation [35]. Methane emission decreases after flowering because the rate of photosynthesis declines after the commencement of grain development and hence the supply of available assimilates for methane production decreases [142].

Diurnal and seasonal variations

Emission rates of CH₄ generally increase rapidly after sunrise, reach a peak in the early afternoon then decline rapidly and level off at night. Methane emission rates during the early and late phase of plant growth varied with a distinct maximum in the early afternoon, while this variation pattern is less pronounced in the middle stage of plant growth [8]. Buendia et al. [21] reported that diurnal patterns of CH₄ fluxes are relatively similar across study sites in same climates and depend on crop phenology. Three seasonal maxima were found in Italy, the first shortly after flooding, the second during the vegetative growth stage and third during the grain filling and maturity stage of rice plants [136].

Rice cultivars, organic manures and crop residues

Rice cultivars have received high research priority because high yielding rice cultivars with low CH₄ emission rates can be easily extended to farmer's fields without any additional input and management practices [151]. Wang et al. [153] argued that, cultivars influence the CH₄ emission by providing the soil with root exudates, decaying root tissues and leaf litter while Aulakh et al. [7] found significant variations in methane

transport capacity of different rice cultivars. In Korean rice cultivars, the CH₄ flux among the rice varieties ranged from 36.9 g CH₄ m⁻² to 76.0 g CH₄ m⁻² [141]. The amendment of organic matter (cattle manure, pig manure, chicken manure, etc.) to a flooded rice field, increases CH₄ production. It reduces the soil Eh and provides carbon to methanogens. Organic materials influence the CH₄ formation through change in qualitative and quantitative properties of soil.

Fertilizers

Numerous studies have revealed the impact of chemical fertilizers on CH₄ emissions [3, 136, 139]. The effect of fertilizers on CH₄ emission depends on rate, type and mode of applications. Urea application enhances CH₄ fluxes over the growth season possibly by increasing soil pH following urea hydrolysis and the drop in redox potential, which stimulates methanogenic activities [152].

Lindau [95] reported decrease in CH₄ emission rate with ammonium nitrate application due to competitive inhibition of nitrate reduction in favour of methane production. Under field conditions, the application of sulphate based fertilizers such as (NH₄)₂ SO₄ and CaSO₄ have reduced CH₄ emission [24] and application of K₂HPO₄ enhances the CH₄ emission [4].

Methane oxidation

Methanotrophs


Methanotrophs (gram negative, aerobic bacteria belonging to the subset of a physiological group of bacteria known as methylotrophs) oxidize CH₄ via methanomonooxygenase (MMO) enzyme (Table 1). These bacteria are classified into three groups: Type-I, Type-II and Type-X. According to Conrad [29], all the methanotrophs that have so far been isolated and described belong to the Proteobacteria, of the γ sub-class (Type I) or α sub-class (Type II). The Type I group is represented by the *Methylomonas*, *Methylocaldum*, *Methylosphaera*, *Methylomicrobium* and *Methylobacter*. The Type-II comprises *Methylosystis* and *Methylosinus*. The members of the genus *Methylococcus* occupy an intermediate position and have been kept in to a separate group Type-X [58]. By using molecular ecology techniques, it has become clear that methanotrophs are ubiquitous in nature and well adapted to high or low temperature, pH and salinity [148]. Henckel et al. [61] found that both Type-I and Type-II methanotrophs were stimulated in rice fields with unsaturated water content. Bodelier et al. [18] reported that Type II methanotrophs dominated in unplanted, unfertilized soils and the presence of rice plant was an essential factor for Type-I methanotrophs to proliferate. Methanotrophic bacteria are present in the aerobic soil layer, rhizosphere [42, 56, 75] and on the roots and stem bases of flooded rice plants [158]. The physiology, biochemistry and ecology of methanotrophic bacteria have been recently reviewed [29, 41].

Methanotrophy

The oxidation of methane by methanotrophs is initiated by methane monooxygenase (MMO) enzyme. The MMO occurs in two forms: as a membrane bound particulate form (sMMO) in all types of methanotrophs, and as a soluble form in Type-II and Type-

X methanotrophs [102]. Results of comprehensive studies on the structure, function and regulation of the MMO have been recently presented by Murrell et al. [113].

According to Conrad [29], the biggest problem for the energy metabolism is the activation of the relatively inert CH₄ molecule. The activation is achieved in the initial step by the MMO which converts CH₄, O₂ and reducing equivalents to methanol and H₂O, i.e.:



The reducing equivalents are supplied by the subsequent dehydrogenation steps, e.g. the conversion of methanol to formaldehyde to formate to CO₂ in which a total of 6 electrons are liberated [29].

Two main CH₄ oxidation pathways, catabolic and anabolic are present in methanotrophs [34]. Catabolic pathway is purely enzymatic through which methanotrophs oxidize methane to CO₂ via methanol, formaldehyde, and formate catalyzed by the enzymes: methane monooxygenase, methanoldehydrogenase, formaldehyde dehydrogenase and formate dehydrogenase, respectively. In this pathway energy is released but carbon is not incorporated into cellular biomass. The anabolic pathway may be further divided into two sub-pathways, ribulose monophosphate pathway and serine pathway. In both these pathways carbon of methane is incorporated in cellular biomass at the level of formaldehyde. Carboxylic acids and amino acids are the intermediary products of the serine pathway. Serine transhydroxy-methylase is the initiator enzyme of this pathway. On the other hand, phosphoglycerated sugars are intermediary products in ribulose monophosphate pathway. RuMP pathway is a cyclic pathway in which fixation is followed by cleavage and rearrangement reactions. Type-I and Type-X methanotrophs follow RuMP pathway whereas serine pathway is followed by Type-II and Type-X methanotrophs [58].

Methanotrophs population and methane oxidation in paddy soils

In rice fields all type of methanotrophs have been detected [52]. From Italian paddy soils two strains of Type-II methanotrophs were isolated [56]. There is a large variation in number of methanotrophs on rice roots, e.g. < 0.1 MOB mm⁻² to > 120 MOB mm⁻² of root surface and in the soil (Table 2). It has been reported that the population size of methanotrophs depends upon location of experimental site, concentration of CH₄ in the soil [10] and concentration of NH₄⁺-N [42, 75]. Rhizosphere has several orders of magnitude higher MOB population than bulk soil [42, 43, 56]. Population size of MOB in rice planted microcosm, rhizosphere, dryland and flooded rice soils as well as endorhizospheric population in rice roots increase with time [20].

Most quantitative data upon methanotrophic population size rely on MPN methods. The limitations of this method are well known. The medium may be selective for certain strains; cells may be in unculturable state; resting and active cells can not be differentiated and microcolonies may be counted instead of single cells [52]. Considering these limitations, the current state of knowledge about population size of methanotrophs from various paddy fields is given in Table 2.

Uptake of atmospheric CH₄ through biological oxidation has been reported in a variety of rice-agroecosystems (Table 4). First evidence of plant associated CH₄ oxidation came from studies with microcosms [52]. Several scientists estimated the amount of CH₄ that is oxidized in association with rice plants and compared with overall CH₄ oxidation [20, 56]. From the review of the available data of CH₄ uptake,

Minami et al. [109] estimated the total terrestrial CH₄ consumption to be between 7 and 78 Tg y⁻¹. Although, rice field is an important source of CH₄, the data of CH₄ oxidation by unflooded paddy soil after harvest could be important for the CH₄ global budget [145].

Table 4. Methane oxidation rates in a variety of rice fields.

Site	CH ₄ oxidation	Sources
Rice field (Panama)	0.1-1.5 mg CH ₄ m ⁻² d ⁻¹	[77]
Paddy field (Italy)	0.1-7.6 nmol CH ₄ h ⁻¹ cm ⁻²	[30]
Paddy field (USA)	170-460 mg CH ₄ m ⁻² d ⁻¹	[54]
Paddy field (Italy)	0.3-162.5 nmol CH ₄ h ⁻¹ g ⁻¹ dw	[12]
Paddy field (USA)	126.2-348.5 ng CH ₄ h ⁻¹ g ⁻¹	[153]
Paddy field (China)	0.7-1.4 μmol CH ₄ h ⁻¹ g ⁻¹ dw	[17]
Paddy field (Japan)	9.5-18.8 μg CH ₄ g ⁻¹ root d ⁻¹	[151]
Paddy field (China)	54.0-892.0 ng CH ₄ g ⁻¹ h ⁻¹	[23]
Paddy field (Italy)	284-810 nmol CH ₄ h ⁻¹ gfw ⁻¹	[72]
Paddy field (Italy)	0.1-0.2 μmol CH ₄ g ⁻¹ dwh ⁻¹	[51]

Gilbert and Frenzel [56] found that the greater part of the CH₄ produced in paddy soil is probably oxidized either in the surface layer of the paddy soil or in the rhizosphere of rice plants. Thurlow et al. [145] showed that unflooded paddy soils after drainage practices, are able to act as sink of CH₄ and vary in their ability to oxidize it depending on the soil temperature and atmospheric CH₄ concentrations. We have found that rhizospheric soil oxidized greater amount of CH₄ (dryland 64-86% ; flooded rice soil 46 to 64%) as compared to bulk and bare soils [40, 45]. Denier and Neue [37] reported that CH₄ emission from rice plants one week before panicle initiation increased by 40% if CH₄ oxidation in the rhizosphere was blocked.

Kinetics of methane oxidation

The apparent half saturation constant (K_m), and maximum oxidation rate (V_{max}) of CH₄ oxidation are characteristic parameters which determine the ability of methanotrophs to grow on atmospheric methane. The CH₄ concentration is a key determinant of $K_{m(app)}$ but this could be mediated through the MMO enzyme, the methanotrophs or the bacterial community as a whole [47].

A model recently proposed by Koch [84] suggest that $K_{m(app)}$ may change in response to the dynamics of substrate utilization as determined by coupling between transport, growth and internal substrate pools. The general model may be useful when applied to methanotrophs, because the first product of methane oxidation, methanol, is sometime excreted and affects the kinetics of MMO [84]. Affinity of methanotrophs for CH₄ varies with growth conditions [45, 47]. According to King [83] the V_{max} of root associated CH₄ oxidation varied largely with season, indicating quantitative and/or qualitative changes in methanotrophic communities.

Recent studies revealed that there are two types of CH₄ oxidizers present in the soil. One population, having a high affinity for CH₄, typically has K_m in the range 1000 nM CH₄, and the other population, having a high affinity for CH₄, has K_m in the range of 30 to 60 nM CH₄ [13]. These methanotrophs typically occur in upland soils that consume atmospheric methane [11], but can also be activated in the paddy soils [10]. However,

the atmospheric CH₄ oxidation has hardly been studied in irrigated rice fields soil. Henckel and Conrad [61] found that moisturised air dried paddy soil does not oxidize CH₄ at atmospheric concentration unless it has been pre-incubated under elevated CH₄ concentration. A decreasing trend of K_m and V_{max} with decreasing CH₄ uptake rate along the soil depth was reported by Wang et al. [153].

Dubey et al. [44, 45] have found that K_m and V_{max} values for CH₄ oxidation in dryland/flooded rice fields decreased from rhizosphere to bulk to bare soil in confirmity with the decreasing CH₄ oxidation activity. Variations in kinetic parameters (K_m and V_{max}) for different rice fields are shown in Table 5. Bender and Conrad [11] have stated that different K_m values may indicate the existence of different types of methanotrophs in soils. According to Conrad [29] type II methanotrophs, which are frequently found in soils, are able to adopt to CH₄ concentration by changing their K_m. This difference could be due to differences in species composition and/or due to conditioning of methanotrophs under different soil microhabitats. According to King [82] all the methanotrophs that have been isolated from soil thus far do not possess the required kinetic properties. These methanotrophs have an ecological niche that is characterized not by atmospheric CH₄ oxidation but by oxidation of relatively high CH₄ concentration that emerge in the proximity of CH₄ production sites i.e. wetlands [128].

Factors affecting methane oxidation

Temperature

Although microbial community is expected to respond to changes in temperature, there are contradictory reports regarding its effect on methane oxidation. Whalen et al. [159] showed that CH₄ consumption increased with increasing temperature (5-20°C) under high CH₄ (103ppm) amended atmosphere. Bender and Conrad [13] reported a linear response in the temperature range of 20-35°C, but 13-38% of the maximum activity remain even at 0°C. A weak relationship between CH₄ uptake and soil temperature suggests that an abiotic process, such as diffusion of CH₄ or O₂ can be a controlling factor for CH₄ uptake.

Methane concentration and soil moisture

Methanotrophs are highly sensitive to variation in CH₄ concentration in atmosphere [11]. CH₄ concentration affects the rate of consumption both directly or indirectly. Enhanced concentration of CH₄ increases the number of methanotrophs) which have a significant role in methanotrophy [103]. Bender and Conrad [13] reported that increase of microbial methane oxidation activity and number of methanotrophs at CH₄ mixing ratios exceeding about 100-1000 μl CH₄l⁻¹. The threshold value below which no CH₄ uptake occurs is much lower for soils than for sediments. For example, 2-3 ppm and < 0.1 to 0.4 ppm threshold values have been reported for sediments and soils, respectively [19]. The maximal CH₄ oxidation rates are probably determined by the magnitude of the supply of CH₄ to the zone of oxidation [82].

Early studies have shown that methane oxidation was sensitive to desiccation and dramatically decreased at soil moisture below at 20 % WHC [13, 72]. The low solubility of CH₄ in water enhances this effect, mainly at low limiting CH₄ concentrations.

Oxygen availability

Oxygen availability depends upon soil porosity. As the porosity increases, a decreased volume of water is distributed in pore volume, decreasing the water film thickness. This increases the rate of substrate (CH₄) delivery to the methanotrophs for oxidation [102]. Methanotrophs in the rice rhizosphere do not have to compete for methane with microbial or chemical compounds, although there is a strong sink of methane by methane transport. However, intensive competition for oxygen occurs. The available values for K_(app) for O₂ and CH₄ indicated that uptake of both substrates is saturated at concentrations of ≥ 10 μM [82].

Table 5. Observed kinetic parameters of CH₄ oxidation.

Ecozone/location	K _m	V _{max}	Reference
Rice field (Italy)	4.0μM	0.1μ mol g ⁻¹ dw h ⁻¹	[20]
Rice field (Italy)	4.1-165μg	1.2-12.5μ g h ⁻¹ g ⁻¹ dw	[153]
Rice field (Italy)	56.0-186.0nM	-	[47]
Rice field (Italy)	16.8nM	839 n M g dw ⁻¹ h ⁻¹	[72]
Dryland rice field (India)	4.8-81.6μ g g ⁻¹ dw	0.05-0.61μ g h ⁻¹ g ⁻¹ dw	[45]
Flooded rice field (India)	6.2-81.1μ g g ⁻¹ dw	0.03-0.41μ g h ⁻¹ g ⁻¹ dw	[40]

Nitrogenous compounds and soil pH

Inorganic N influences CH₄ oxidation due to shifts in the population structure and the kinetics of methanotrophs [45]. This may affect the threshold value for CH₄ oxidation [82]. NO₃⁻-N fertilization did not affect the CH₄ consumption but NH₄⁺-N fertilization completely ceased CH₄ oxidation [66]. Nitrite was found to inhibit CH₄ oxidation in the cultures of *Methylobacterium albus* BG8 and *M. trichosporium* OB3b [83]. Recently it was shown that methane oxidation is stimulated by increased nitrogen availability due to unquantified nitrogen limitation of methanotrophs [18]. However, the most solidly substantiated explanation for ammonium inhibition of methane oxidation is competitive inhibition at the enzyme level. This occurs because, at the molecular scale, methane and ammonium are similar in size and structure [135]. As a result, the enzyme MMO can bind to ammonium ion and react with it. Because the possibility of competitive inhibition is fundamental to the biochemistry of methane oxidation, it was generally thought that inhibition should occur in paddy fields as well as in upland systems [135].

In a pasture soil, CH₄ oxidation at pH 6.3 was greater than at pH 5.6 and was completely inhibited at pH 4.8-5.1 [66]. In certain other soils, oxidation has been reported at pH as low as 3.2 [143]. In general, low pH has an inhibitory effect on methane consumption although the mechanism responsible for this effect is not fully known [66].

Inhibitors for CH₄ production and oxidation

An inhibitor specific to either methanogens or methanotrophs would be useful for distinguishing which of these organisms is responsible for CH₄ production and oxidation in environments in which such activities occur. A variety of chemicals used in agriculture such as pesticides and herbicides and nitrification inhibitors, are known to affect microbial processes. It is well established that CH₄ production is inhibited by acetylene, aminopurine, ammoniumthiosulphate, carbofurane, calcium carbide (capsulated), DDT, dicyandiamide, methyle chloride, methyle fluoride, nitrapyrine, pyridine, organochlorine and sodium azide and CH₄-acetylene, bromoxynil, dicyandiamide, DDT, 2,4,-D, ethylene, hexachlorocyclohexane, hydrazine, methomyle, nitrapyrine, phenylalanine, sodium thiosulphate, threonine and thiourea [9, 25, 147]. Nitrification inhibitors such as acetylene and nitrapyrin can inhibit the growth of nitrifiers, methanogens and methanotrophs [107]. Lindau et al. [96] found that CH₄ emissions from rice fields decreased by 35% and 14% following the application of encapsulated calcium carbide and dicyandiamide, respectively. Topp [146], found that the pesticides, bromoxynil, methomyl and nitrapyrin were inhibitory to CH₄ oxidation at 50 g l⁻¹. Sathpathy et al. [134] reported that the application of HCH (organochlorine insecticide) to flooded rice soils reduced the production and emission of CH₄. Kumarswamy et al. [89] have revealed that carbofuran inhibited net CH₄ production when applied at low rates (5-10 mg g⁻¹ soil), but stimulated it when applied at a rate of 100 mg g⁻¹ soil. Chan and Parkin [25] have recommended the use of acetylene and ethylene for inhibition of CH₄ oxidation and methyl chloride for inhibition of methanogenesis.

Approach for Methanogens/Methanotrophs Detection at Molecular Ecological Level

Effective representation of the diversity towards desired functional gene is one of the major concerns of microbial ecology of any ecosystem. A preliminary attempt has been reported by Dubey et al. [46] where, by cluster analysis of ARDRA pattern, it has been shown that different types of methanotrophs dominate in rhizosphere and non-rhizosphere soils of tropical rice ecosystems. The growing demand of paddy with increasing population is an emerging issue and the scenario results in the over exposure of soil microflora with pesticide residues. As discussed earlier pesticides play a major role at the physiological level for the expression of key enzymes in methane production and its further utilization as a substrate. The question, what are the emerging criteria that decide the course of evolution for these bacteria, which play key role in maintaining the greenhouse gas methane in the ecosystem and for habitats exposed to such residues, is likely to trigger a lot of interest. Studies are required to compare the performance of diverse strains from such habitats followed by molecular analysis of key genes to understand which is the most susceptible genotype for such stresses. These studies are

likely to generate the tracking probes for the same target genes but having substituted primary coding sequences that will help evaluating the impact on soil methane cycle and its relation to pesticide residues. Similarly, other parameters such as application of fertilizers and accumulation of inorganic compounds in soil will lead to modified soil chemistry which in turn will be changing the overall microbial community structure vis-à-vis their relation in density and survival of methane oxidizing bacteria. Therefore, parallel tracking tools would have to be designed to assess the existing microbial population using selected markers which could act as biomarker for associated biogeochemical cycles.

Several approaches have been adopted in molecular ecological studies for the detection and characterization of methanogens and methanotrophs in the various ecosystems [46, 112]. These are schematically shown in Figure 2. The first approach is indirect and relies on enrichment and/or isolation with subsequent characterization of methanogens and methanotrophs. The second approach is direct and relies on polymerase chain reaction (PCR) technique and use of phylogenetic and functional gene probes (Table 6) for the molecular analysis of these microbes in the environmental samples without the prerequisite of their cultivation. At each stage, in both approaches, molecular biological analysis of the key genes (*pmoA*, *mmoX*, *mxoF* for methanotrophs, *mcr*, *mtd*, *mth*, *mrt*, *frh* for methanogens and 16S r RNA) can be carried out and comparison can be made to see if the same organism that grows in culture derived from the sample is representative of what is actually present in the environmental sample (as revealed by DNA analysis).

Several other approaches have also been used to study methanogens and methanotrophs. Phospholipid analysis has been used successfully for the detection of these microbes in the wetland ecosystem [144].

Table 6. Sequences of oligonucleotides commonly used as PCR primers and probes for molecular analysis of methanogens and methanotrophs.

Primers /Probes	Targets	Sequences (5' → 3')	Tm (°C)	References
PRIMERS				
A 189 f	<i>pmoA</i> ; pMMO/AMO	GGN GAC TGG GAC TTC TGG	56	[63]
A 682 r	<i>pmoA</i> ; pMMO/AMO	GAA SGC NGA GAA GAA SGC	56	[63]
882 f	<i>mmoX</i> ; sMMO ⁺ Methylotrophs	GGC TCC AAG TTC AAG GTC AG	55	[99]
1403 r	<i>mmoX</i> ; sMMO ⁺ Methylotrophs	TGG CAC TCG TAG CGC TCCGGCTCG	55	[99]
1003 f	<i>mxoF</i> ; All Methylotrophs	GCG GCA CCA ACT GGG GCT GGT	55	[100]
1561 r	<i>mxoF</i> ; All Methylotrophs	GGG CAG CAT GAA GGG CTC CC	55	[100]
ME1 f	<i>mcrA</i> ; methanogens	GCM ATG CAR ATH GGW ATG TC	50	[60]
ME2 r	<i>mcrA</i> ; methanogens	TCA TKG CRT AGT TDG GRT AGT	50	[60]
Probes				
MSMX 860	Methanosarcinaceae	GGC TCG CTT CAC GGC TTC CCT	54	[127]
MS821	Methanosarcina spp.	CGC CAT GCC TGA CAC CTA GCG AGC	54	[127]
MC1109	Methanococcaceae	GCA ACA TAG GGC ACC GG TCT	47	[127]
MB310	Methanobacteriaceae	CTT GTC TCA GGT TCC ATC TCC G	52	[127]
MG1200	Methanomicrobiales	CGG ATA ATT CGG GGC ATG CTG	45	[127]

This technique relies on the fact that methanotrophs contain unusual fatty acids (16:1 and 18:1 derivatives). Phospholipid analysis is, however, limited due to small database

on fatty acid profiles from methanotrophs [113] and the relatively high cost of equipment used in this analysis.




Figure 2. Generalized scheme of molecular analysis for detection and phylogenetic study of methanogens and methanotrophs

The detection and characterization of these microbes can also be attempted using antibodies either to whole cells of culturable methanogens/methanotrophs or to key enzymes such as MMO, MDH (methanotrophs) and MCR, MTD (methanogens). One of the most powerful tools in the molecular ecology is FISH (fluorescence in situ hybridization). FISH allows the specific detection and enumeration of methanogens/methanotrophs directly in the natural habitats without cultivation. Grobkopf et al. [57] detected most of the archaeal group members (e.g. *Methanosarcinaceae*,

Methanosaetaceae, *Methanomicrobiaceae*, *Methanobacteriaceae*, RC-I to RC-IV and RC-VI) on rice plant roots by using FISH.

Conclusions

Methane is an important greenhouse gas and it affects the chemistry of the atmosphere. The ecological role of methanogens and methanotrophs in the methane dynamics in rice fields is still unclear. Current information is insufficient for the development of technology and strategy for reduction in methane emission from rice field at regional and global levels. Knowledge of comparative genomics and proteomics of methanogens and methanotrophs will contribute to the deciphering their population structure and existing mechanisms of methane emission in paddy fields.

It has now become possible to isolate, detect and characterize these microbes by using molecular biological tools like PCR, FISH, etc. techniques. Knowledge of structure and function of methanogens and methanotrophs communities will be beneficial for understanding the microbial ecology of methane to control the CH₄ turnover in rice soils.

Acknowledgements. The author acknowledges the kind financial support provided by department of science and Technology, Government of India, New Delhi.

REFERENCES:

- [1] Achnich, C., Bak, F., Conrad, R., (1995): Competition for electron donors among nitrate reducers, ferric iron reducers, sulfate reducers, and methanogens in anoxic paddy soil. – *Biology and Fertility of Soils* 19, 65-72.
- [2] Adachi, K., (1999): Isolation of hydrogenotrophic methanogenic archaea from a subtropical paddy field. *FEMS. – Microbiology Ecology* 30, 77-85.
- [3] Adhya, T.K., Bharati, K., Mohanty, S.R., Ramakrishnan, B., Rao, V.R., Sethunathan, N., Wassmann, R., (2000): Methane emission from rice fields at Cuttack, India. – *Nutrient Cycling in Agroecosystems* 58, 95-105.
- [4] Adhya, T.K., Pattnaik, P., Satpathy, S.N., Kumaraswamy S., Sethunathan, N, (1997): Influence of phosphorus application on methane emission and production in flooded paddy soils. – *Soil Biology and Biochemistry* 30, 177-181.
- [5] Anastasi, C., Dowding, M., Simpson, V.J, (1992): Future CH₄ emissions from rice production. – *Journal of Geophysical Research* 97, 7521-7525.
- [6] Asakawa, S., Agakawa-Matsushita, M., Morii, H., Yago, Y., Hayano, K., (1995): Characterization of *Methanosacina mazeii* TMA isolated from a paddy field soil. – *Current Microbiology* 31, 34-38.
- [7] Aulakh, M.S., Bodenbender, J., Wassmann, R., Reenberg, H., (2000): Methane transport capacity of rice plants. II. Variations among different rice cultivars and relationship with morphological characteristics. – *Nutrient Cycling in Agroecosystems*. 58, 367-375.
- [8] Aulakh, M.S., Wassmann, R., Reenberg, H., (2001): Methane emissions from rice fields quantification, mechanisms, role of management and mitigation options. – *Advances in Agronomy* 70, 193-260.
- [9] Bedard, C., Knowles, R., (1989): Physiology, biochemistry and specific inhibitors of CH₄, NH₄⁺ and CO oxidation by methanotrophs and nitrifiers. – *Microbiological Review* 53,68-84.
- [10] Bender, M., Conrad, R., (1992): Kinetics of CH₄ oxidation in oxic soils exposed to ambient air or high CH₄ mixing ratios. – *FEMS Microbiology Ecology* 101, 261-270.

- [11] Bender, M., Conrad, R., (1993): Kinetics of methane oxidation in oxic soils. – *Chemosphere* 26, 687-696.
- [12] Bender, M., Conrad, R., (1994): Microbial oxidation of methane, ammonium and carbon monoxide, and turnover of nitrous oxide and nitric oxide in soils. – *Biogeochemistry* 27, 97-112.
- [13] Bender, M., Conrad, R., (1995). Effect of CH₄ concentrations and soil conditions on the induction of CH₄ oxidation activity. – *Soil Biology and Biochemistry* 27, 1517-1527.
- [14] Blake, D.R. and Rowland, F.S. (1988): Continuing world wide increase in tropospheric methane (1978-1987). – *Science* 239, 1129-1131.
- [15] Bharati K., Mohanty, S.R., Padmavathi, P.V.L, Rao, V.R., Adhya, T.K., (2000): Influence of six nitrification inhibitors on methane production in a flooded alluvial soil. – *Nutrient Cycling in Agroecosystems* 58, 389-394.
- [16] Blaut, M., (1994): Metabolism of Methanogenes. – *Antonie Van Leeuwenhoek* 66, 187-208.
- [17] Bodelier, P.LE., Frenzel, P., (1999) Contribution of methanotrophic and nitrifying bacteria to CH₄ and NH₄⁺ oxidation in the rhizosphere of rice plants as determined by new methods of discrimination. – *Applied and Environmental Microbiology* 65,1826-1833.
- [18] Bodelier, P.LE., Roslev, P., Henckel, T., Frenzel, P., (2000): Stimulation by ammonium based fertilizers of methane oxidation in soil around rice roots. – *Nature* 403, 421-424.
- [19] Born, M., Dorr, H., Levin, J., (1990): Methane consumption in aerated soils of the temperate zone. – *Tellus B* 42, 2-8.
- [20] Bosse, U., Frenzel, P., (1997): Activity and distribution of methane oxidizing bacteria in flooded rice soil microcosms and in rice plants (*Oryza sativa*). – *Applied and Environmental Microbiology* 63,1199-1207.
- [21] Buendia, L.V., Neue, H.D., Wassmann, R.B., Lantin, R.S., Javellana, A.M., Yuchang, X., Markarim, A.K., Corton, T.M., Charoensilp, N., (1997): Understanding the nature of methane emission from rice ecosystems as basis of mitigation strategies. – *Applied Energy* 56, 433-444.
- [22] Byrenes, B.H., Austin, E.R., Tays, B.K., (1995): Methane emission from flooded rice soils and plants under controlled conditions. – *Soil Biology and Biochemistry* 27, 331-339.
- [23] Cai, Z.C., Mosier, A.R., (2000): Effect of NH₄ an addition on methane oxidation by paddy soils. – *Soil Biology and Biochemistry* 32,1537-1545.
- [24] Cai, Z.C., Xing, H., Yan, X., Xu, H., Tsuruta, H., K. Yagi, K., Minami, K., (1997): Methane and nitrous oxide emissions from rice paddy fields as affected by nitrogen fertilizers and water management. – *Plant and Soil*. 196, 7-14.
- [25] Chan, A.S.K., Parkin, T.B., (2000): Evaluation of potential inhibitors of methanogenesis and methane oxidation in a landfill cover soil. – *Soil Biology and Biochemistry* 32, 1581-1590.
- [26] Chen, Z., Li, D., Shao, K., Wang, B., (1993): Features of CH₄ emission from rice paddy field in Beijing and Nanjing. – *Chemosphere* 26, 239-245.
- [27] Chin, K. J., Conrad, R., (1995): Intermediary metabolism in methanogenic paddy soils and the influence of temperature. – *FEMS Microbiology and Ecology* 18, 85-102.
- [28] Cicerone, R.J., Shetter, J. D., (1981): Source of atmospheric methane: measurement in rice paddies and a discussion. – *Journal of Geophysical Research* 86, 7203-7209.
- [29] Conrad, R., (1999): Soil microorganisms oxidizing atmospheric trace gases (CH₄, CO, H₂, NO). – *Indian Journal of Microbiology* 39, 193-203.
- [30] Conrad, R., Rathfuss, F., (1991): Methane oxidation in soil surface layer of a nooded rice field and effect of ammonium. – *Biology and Fertility of Soils* 12,28-32.
- [31] Conrad, R., Schink, B., Phelps, T.J., (1986): Thermodynamics of H₂ - consuming and H₂ producing metabolic reactions in diverse methanogenic environments under in situ conditions. – *FEMS Microbiology Ecology* 38, 353-360.

- [32] Crutzen, P.J., (1985): The role of the tropics in atmospheric chemistry .In: Dickinson, R., (ed.), *Geophysiology of Amazon*. – Wiley, Chichester, UK. pp. 107-132.
- [33] Crutzen, P.I, (1991): Methane: sources and sinks. *Nature* 350, 380-381.
- [34] Daltan, H., Leakck, D.J., (1985): Microbial gas metabolism. In: Poole, R.K., Dow, C.S., (eds.), *Methane Oxidation by Microorganisms*. – National Academic Press, Washington. pp. 173-295.
- [35] Dannenberg, S. Conrad, R., (1999): Effect of rice plant on methane production and rhizospheric metabolism in paddy soil. – *Biogeochemistry* 45, 53-71.
- [36] De Long, E.F., (2000): Resolving a methane mystery. – *Nature* 407, 577-579.
- [37] Denier van der Gon, H.A.C., Neue, H.U., (1996): Oxidation of methane in the rhizosphere of rice plants. – *Biology and Fertility of Soils* 22, 359-366.
- [38] Denier Van Der Gon, H.A.C., Neue, H.U., (1995): Influence of organic matter incorporation on the methane emission from a wetland rice field. – *Global Biogeochemical Cycles* 11, 11-22.
- [39] Dubey, S.K., (2001): Methane emission and rice agriculture. – *Current Science* 81, 345-346.
- [40] Dubey, S.K., (2003): Spatio-kinetic variation of methane oxidizing bacteria in paddy soil at mid tillering: Effect of N fertilizer. – *Nutrient Cycling in Agroecosystems* 65, 53-59.
- [41] Dubey, S.K., Kashyap, A.K., Singh, J.S., (1996): Methanotrophic bacteria, methanotrophy and methane oxidation in soil and rhizosphere. – *Tropical Ecology* 37, 167-182.
- [42] Dubey, S.K., Singh, J.S., (2000): Spatio-temporal variation and effect of urea fertilization on methanotrophs in a tropical dryland rice field. – *Soil Biology and Biochemistry* 32, 521-525.
- [43] Dubey, S.K., Singh, J.S., (2001): Plant induced spatial variation in the size of methanotrophic population in dryland and flooded rice agroecosystems. – *Nutrient Cycling in Agroecosystems* 59, 161-167.
- [44] Dubey, S.K., Sinha, A.S.K., Singh, J.S., (2000): Spatial variation in the capacity of soil for CH₄ uptake and population size of methane oxidizing bacteria in dryland rice agriculture. – *Current Science* 78,617-620.
- [45] Dubey, S.K., Sinha, A.S.K., Singh, J.S., (2002): Differential inhibition of CH₄ oxidation in bare, bulk and rhizosphere soils of dryland rice field by nitrogen fertilizers. – *Basic and Applied Ecology* 3, 347-355.
- [46] Dubey, S.K., P. Padamabhan, H.J. Purohit and S.N. Upadhyay (2003): Tracking of methanotrophs and their diversity in paddy soil: A molecular approach. – *Current Science* 85, 92-95.
- [47] Dunfield, P.F., Liesack, W., Henckel, T., Knowles, R., Conrad, R., (1999): High affinity methane oxidation by a soil enrichment culture containing a type II methanotrophs. – *Applied and Environmental Microbiology* 65, 1009-1014.
- [48] Ehhalt, D.H., Schmidt, U., (1978): Sources and sinks of atmospheric methane. – *Pageophysics* 116, 452-464.
- [49] Ferry, J.G., (1997): Methane: small molecule, big impact. – *Science* 278, 1413-1414.
- [50] Fetzer, S., Bak, F., Conrad, R., (1993): Sensitivity of methanogenic bacteria from paddy soils to oxygen and dessication. – *FEMS Microbiology Ecology* 12, 107-115.
- [51] Filler, G., Frenzel, P., (2001): Changes in activity and community structure of methane oxidizing bacteria over the growth period of rice. – *Applied and Environmental Microbiology* 67, 2395-2403.
- [52] Frenzel, P., (2000): Plant-associated methane oxidation in rice fields and wetlands. – *Advances in Microbial Ecology* 16, 85-114.
- [53] Garcia, I.L., (1990): Taxonomy and ecology of methanogens. – *FEMS Microbiological Review* 87, 297-308.
- [54] Gerard, G., Chanton, J, (1993): Quantification of methane oxidation in the rhizosphere of emergent aquatic macrophytes: Defining upper limits. – *Biogeochemistry* 23, 79-97.

- [55] Gilbert, B., Frenzel, P., (1995): Methanotrophic bacteria in the rhizosphere of rice microsoms and their effect on pore-water methane concentration and methane emission. – *Biology and Fertility of Soils* 20, 93-100.
- [56] Gilbert, B., Frenzel, P., (1998): Rice roots and CH₄ oxidation: the activity of bacteria, their distribution and the microenvironment. – *Soil Biology and Biochemistry* 30, 1903-1916.
- [57] Grobcof, R., Stubner, S., Liesack, W., (1998): Novel euryarchaeotal lineages detected on rice roots and in the anoxic bulk soil of flooded rice microcosms. – *Applied and Environmental Microbiology* 64, 4983-4989.
- [58] Hanson, R.S., Hanson, T.E., (1996): Methanotrophic bacteria. – *Microbiology Review* 62, 439-471.
- [59] Hattori, C., Ueki, A., Seto, T., Ueki, K., (2001): Seasonal variations in temperature dependence of methane production in paddy soil. – *Microbes and Environments* 16, 227-233.
- [60] Hales B. A., Edward C., Ritchie D.A., Hall G., Pickup R.W., Saunders J.R. (1996): Isolation and identification of methanogen specific DNA from blanket bog peat using PCR amplification and sequence analysis. – *Applied and Environmental Microbiology* 62, 668-675.
- [61] Henckel, T., Friedrich, M., Conrad, R., (1999): Molecular analysis of the methane-oxidizing microbial community in rice field soil by targeting the genes of the 16S rRNA, particulate methane monooxygenase, and methanol dehydrogenase. – *Applied and Environmental Microbiology* 65, 1980-1990.
- [62] Holzapfel-Pschorn, A., Conrad, R., Seiler, W., (1986): Production, oxidation and emission of methane in rice paddies. – *FEMS Microbiology Ecology* 31, 149-158.
- [63] Holmes, A.J., Costello, A., Lidstrom, M.E. and Murrell, J.C. (1995): Evidence that particulate methane monooxygenase and ammoniamonooxygenase may be evolutionarily related. – *FEMS Microbiology Letter* 132, 203-208.
- [64] Hou, A.X., Wang, Z.P., Chen, G.X., Patrick Jr, W.H., (2000): Effect of organic and N fertilizer on methane production potential in a Chinese rice soil and its microbiological aspect. – *Nutrient Cycling in Agroecosystems* 58, 333-338.
- [65] Huke, R.E., Huke, E.H., (1997): *Rice Area by Type of Culture: South, Southeast, and East Asia*. International Rice Research Institute, Los Banos, Laguna, Philippines, p. 15.
- [66] Hutsch, B.W., Webster, C.P., Powlson, D.S., (1994): Methane oxidation in soil as affected by land use, pH, and N-fertilization. – *Soil Biology and Biochemistry* 26, 1613-1622.
- [67] I.R.R.I., (1997): *Rice Almanac*, 2nd ed. IRRI Manila, Philippines. 18. p.
- [68] IPCC, (1992): Climate change. In: Houghton, I.T., Callander, B.A., Barney, S.K., (eds.) *The Supplementary Report to the IPCC Scientific Assessment*. – Cambridge University Press, Cambridge, U.K. p. 200.
- [69] IPCC, (1994): The Science of climate change. In: (Houghton, I.T., Meira Filho, L.G., Bruce, J., Lee, H., Callander, B.A, Harris, N., Haites, E., Maskell, K., (Eds.), *Radioactive Forcing of Climate Change and Evaluation of the IPCC Emission Scenarios*. – Cambridge University Press, Cambridge. U.K.
- [70] IPCC, (1995): The science of climate change. In: (Houghton, I.T., Meira, F., Callander, L.G., Harris, B.A, Kattenberg, A, Maskell, K., (Eds.), *Contribution of Working Group I. To the Second Assessment Report of the Intergovernmental Panel on Climate Change*. – Cambridge University Press, UK.
- [71] IPCC, (1996): XII summary for policy makers. In: (Houghton, I.T., Meira, F., Callander, L.G., Harris, B.A., Kattenberg, A., Maskell, K., (Eds.), *Climate Change 1995: The Scientific Basis of Climate Change*. – Cambridge University Press, Cambridge, UK. p. 572.

- [72] Jackel, U., Schnell, S., Conrad, R., (2001): Effects of moisture, texture and aggregate size of paddy soil on production and consumption of CH₄. – *Soil Biology and Biochemistry* 33, 965-971.
- [73] Jermasawatdipong, P., Murase, I., Prabuddham, P., Hasathon, Y., Khomthong, N., Naklang, K., Watanabe, A., Haaraguchi, H., Kimura, K., (1994): Methane emission from plots with different fertilizer application in Thai paddy fields. – *Soil Science and Plant Nutrition* 40, 63-71.
- [74] Jones, W.J., (1991): Diversity and physiology of methanogens p. 39-56. In: Rogers, J.R., Whitman, W.B., (eds.), *Microbial Production and Consumption of Greenhouse Gas: CH₄, nitrogen oxides, and halo methanes*. – American Society for Microbiology, Washington, DC.
- [75] Joulain C., Escoffier, S., Lemer, J, Neue, H.U., Roger, P.A. (1997): Population and potential activities of methanogens and methanotrophs in rice fields: Relation with soil properties. – *European Journal of Soil Biology* 33, 105-166.
- [76] Joulain C., Ollivier, B., Patel, B.K.C., Roger, P.A., (1998): Phenotypic and phylogenetic characterization of dominant culturable methanogens isolated from rice fields soils. – *FEMS Microbiology Ecology* 25, 135-145.
- [77] Keller, M, Mitre, M.E., Stallard, R.F., (1990): Consumption of atmospheric methane in soils of Central Panama: Effects of agricultural development. – *Global Biogeochemical Cycles* 4, 21-27.
- [78] Khalil, M.A.K., Rasmussen, R.A., 1983. Sources, sinks cycles of atmospheric methane. – *Journal of Geophysical Research* 88, 5131-5144.
- [79] Khalil, M.A.K., Shearer, MJ., (1993): Atmospheric methane: Sources sinks and role in global change. – *Chemosphere* 26, 201-217.
- [80] Kheshti, H.S., Jain, A.K., Kotamarthi, V.R., Wuebbles, D.I, (1999): Future atmospheric methane concentrations in the context of the stabilization of greenhouse gas concentrations. – *Journal of Geophysical Research* 104, 19182-19190.
- [81] Kimura, M., Miura, Y., Watanabe, A, Katoh, T., Haraguehi, H., (1991): Methane emission from paddy field. (I) effect of fertilization, growth stage and midsummer drainage: pot experiment. – *Environmental Science* 14, 265-271.
- [82] King, G.M., (1992): Ecological aspect of methane oxidation, a key determinant of global methane dynamics In: (Mashall, K.c., ed.), *Advances in Microbial Ecology*. Plenum, New York. pp. 431-468.
- [83] King, G.M., Schnell S., (1994): Effect of increasing atmospheric methane concentration on ammonium inhibition of soil methane consumption. – *Nature* 370, 282-284.
- [84] Koch, A.L., (1997): Microbial physiology and ecology of slow growth. – *Microbial and Molecular Biology Review*, 61, 305-318.
- [85] Koyama, T., (1963): Gaseous metabolism in lake sediments and paddy soils and the production of hydrogen and methane. *Journal of Geophysical Research*, 68, 3971-3973.
- [86] Kral, T.A, Brink, K.M, Miller, S.L, Mc Kay, C.P., (1998): Hydrogen consumption by methanogens on the early earth. – *Origins of the Life and Evolution of the Biosphere* 28, 311-319.
- [87] Kudo, Y., Nakojima, T., Miyaki, T, Oyaizu., H., (1997): Methanogen flora of paddy soils in Japan. – *FEMS Microbiology Ecology* 22, 39-48.
- [88] Kumaraswamy, S, Ramakrishnan, B, Sethunathan, N., (2001): Methane production and oxidation in an anoxic rice soil as influenced by inorganic redox species. – *Environmental Quality* 30, 2195-2201.
- [89] Kumaraswamy, S, Ramakrishnan, B., Satpathy, S.N., Rath, A.K., Mishra, S., Rao, V.R., Sethunathan, N., (1997): Spatial distribution of methane oxidizing activity in a flooded rice soil. – *Plant and Soil* 191, 241-248.
- [90] Kumaraswamy, S., Rath, AK., Satpathy, S.N., Ramakrishnan, B., Adhya, TK., Sethunathan, N., (1998): Influence of an insecticide carbofuran on production and oxidation of CH₄ in flooded rice soils. – *Biology and Fertility of Soils* 26, 362-366.

- [91] Le Mer, J., Escoffier, S., Chessel, C., Roger, A.A. (1996): Microbial aspects of methane emission in rice field soil from Camargue (France): 2. Methanotrophy and related microflora. – *European Journal of Soil Biology* 32: 71-80.
- [92] Lehmann-Richter, S., Grosskopf, R., Liesack, W., Frenzel, P., Conrad, R., (1999): Methanogenic archaea and CO₂ dependent methanogenesis on washed rice roots. – *Environmental Microbiology* 1, 159-166.
- [93] Lelieveld, J., Crutzen, P.J., Dentener, F.J., (1998): Changing concentration, life time and climate forcing of atmospheric methane. – *Tellus* 50B, 128-150.
- [94] Li, C.S., (2000): Modelling trace gas emission from agricultural ecosystem. – *Nutrient Cycling in Agroecosystems* 58, 259-267.
- [95] Lindau, C.W., (1994): Methane emission from Louisiana rice fields amended with nitrogen fertilizers. – *Soil Biology and Biochemistry* 26, 353-359.
- [96] Lindau, C.W., Bollich, P.K., Delaune, R.D., Mosier, A.R., Bronson, K.F., (1993): Methane mitigation in flooded Louisiana rice fields. – *Biology and Fertility of Soils* 15, 174-178.
- [97] Lindau, C.W., Bollich, P.K., Delaune, R.D., Patrick Jr., W.H., Law, V.J., (1991): Effect of urea fertilizer and environmental factors on CH₄ emissions from Louisiana USA rice fields. – *Plant and Soil*. 136, 195-203.
- [98] Ludmila, C., Julia A., Rudoyk R., Toms M., Lidstrom E., (1998): G Transfer enzymes and coenzymes linking methylotrophic bacteria and methanogenic archaea. – *Science* 281, 99-101.
- [99] Mc Donald, I.R., Kenna, E.M. and Murrell, J.C. (1995): Detection of methanotrophic bacteria in environmental samples with the PCR. – *Applied and Environmental Microbiology* 61, 116-121.
- [100] Mc Donald, I.R. and Murrell, J.C. (1997): The particulate methane monooxygenase gene *pmoA* and its use as a functional gene probe for methanotrophs. – *FEMS Microbiology Letter* 156, 205-210.
- [101] Mah, R.A., Smith, M.R., (1981): The methanogenic bacteria. In: Starr, M.P., Stolp, H., Truper, H.G., Baws, A., Schlegel, H.G., (eds.), *The Prokaryotes and Handbook on Habitats Isolation, and Identification of Bacteria*. Springer, Berlin, Heidelberg, New York. pp. 948-977.
- [102] Mancinelli, R.L., (1995): The regulation of methane oxidation in soil. – *Annual Review of Microbiology* 49, 581-605.
- [103] Mancinelli, R.L., White, M.R., Bogher, I., (1991): Microbial methane oxidation. In: P.L. Wilkey (ed.) *Rio vista gas leak study*, Argonne National Laboratory Topical Report, April 1989 to January 1991. Argonne National Laboratory IL.
- [104] Masscheleyn P.H., Delaune, R.D., Patrick, W.H., (1993): Methane and nitrous oxide emission from laboratory measurements of rice soil suspension. Effect of soil oxidation - reduction status. – *Chemosphere* 26, 251-260.
- [105] Matthews, E., Fung, I., Lerner, J., (1991): Methane emission from rice cultivation: Geographic and seasonal distribution of cultivated areas and emission. – *Global Biogeochemical Cycles* 5, 3-24.
- [106] Matthews, R.B., Wassmann, R., Knox, J.K., Buendia, L.V., (2000): Using a crop/soil simulation model and GIS techniques to assess methane emissions from rice fields in Asia. IV. Upscaling to national levels. – *Nutrient Cycling in Agroecosystems* 58, 201-217.
- [107] McCarty, G.W., (1999): Modes of action of nitrification inhibitors. – *Biology and Fertility of Soils* 29, 1-9.
- [108] Min, H., Zhao, Y. and X.W. Wu, (2002): Microbial aerobic oxidation in paddy soil. – *Nutrient Cycling in Agroecosystems* 64, 79-85.
- [109] Minami, K., Goudrian, J., Lantinga, E.A., Kimura, T., (1993): Significance of grasslands in emission and absorption of greenhouse gases. – In: *Proceedings of 17th International Grassland Congress*. pp. 1231-1238.

- [110] Mishra, S., Rath, A.K., Adhya, T.K., Rao, V.R., Sethunathan, N., (1997): Effect of continuous and alternate water regims on methane efflux from rice under greenhouse conditions. – *Biology and Fertility of Soils* 24, 399-405.
- [111] Mitra, S., Jain, M.C., Kumar, S., Bandopadhyay, S.K., Kalra, N., (1999): Effect of rice cultivars on methane emission. – *Agriculture Ecosystems and Environment* 73, 177-183.
- [112] Morris, S.A., S., Radajewski, T.W., Willison and J.C., Murrell (2002): Identification of the functionally active methanotroph population in a peat soil microcosm by stable-isotope probing. – *Applied and Environmental Microbiology* 68, 1446-1453.
- [113] Murrell, J.C., McDonald, I.R. and Bourne, D.G. (1998): Molecular methods for the study of methanotroph ecology. – *FEMS Microbiology* 27, 103-114.
- [114] Neue, H.D., (1993): Methane emission from rice fields. – *Bioscience* 43, 466-474.
- [115] Neue, H.D., (1997): Fluxes of methane from rice fields and potential for mitigation. – *Soil Use Management* 13, 258-267.
- [116] Neue, H.D., Becker-Heidmann, P., Scharpenseel, H.W., (1990): Organic matter dynamics, soil properties and cultural practices in rice lands and their relationship to methane production – In: (Bouwman, A.F., ed.), *Soils and Greenhouse Effect*. Wiley, Chichester. pp. 457-466.
- [117] Neue, H.D., Sass, R.L., (1998): The budget of methane from rice fields. – *IGAC Activities* 12, 3-11.
- [118] Nouchi, I, Mariko, S., Aoki, K., (1990): Mechanism of methane transport from the rhizosphere to atmosphere through rice plants. – *Plant Physiology* 94, 59-66.
- [119] Nugraho, S., Lumbanjara, G., Suprpto, I, Sunyoto, H., Ardjasa, W.S., Haraguchi, H. Kimura, M. (1994). Methane emission from an Indonesian paddy fields subjected to several fertilizers treatments. – *Soil Science and Plant Nutrition* 40, 275-281.
- [120] Palmer, R.R., Reeve, I.N., (1993): Methanogene genes and the molecular biology of methanogenesis. – In: Sebald, M.,(ed.), *Genetics and Molecular Biology of Anaerobic Bacteria*. Springer-Verlag, Berlin. pp. 13-35.
- [121] Parashar, D.C., Rai, J, Gupta, P.K., Singh, N., (1991): Parameter affecting methane emission from paddy fields. – *Indian Journal of Radio and Space Physics* 20, 12-17.
- [122] Patrick, W.H. Jr., (1981): The role of inorganic redox systems in controlling reduction in paddy soils. – In: *Proceedings of Symposium on Paddy Soil*. Institute of Soil Science, Academia Sinica. Beijing and Springer Verlag, Berlin. pp. 107-117.
- [123] Peters, V., Conrad, R., (1996): Sequential reduction processes and initiation of CH₄ production upon flooding of oxic upland soils. – *Soil Biology and Biochemistry* 28, 371-382.
- [124] Prinn, R.G., (1994): Global atmospheric-biospheric chemistry. In: Prinn, R.G., (eds) *Global atmospheric-biospheric chemistry*, pp. 1-18. Plenum, New York.
- [125] Prinn, R.G., (1995): Global change: Problems and uncertainties. In: Peng, S., Ingram, K.T, Neue, H.D., Ziska, L.H. (eds.), *Climate Change and Rice*, Springer, Berlin. pp. 3-7.
- [126] Rajgopal, B.S., Belay, N., Daniel, L, (1988): Isolation and characterization of methanogenic bacteria from rice paddies. – *FEMS Microbiology Ecology* 53, 153-158.
- [127] Raskin L, Stromley J M, Rittmann BE , Stahl DA, (1994): Group specific 16 S rRNA hybridization probes to describe natural communities of methanogens. – *Applied and Environmental Microbiology* 60, 1232-1240.
- [128] Reeburgh, W.S., Whjalen, S.C., Alperin, M.J., (1993): The role of methylotrophy in the global methane budget In: Murrell, J.C., Kelly, D.P., (eds.), *Microbial growth on C1 compound*. Intercept Ltd., Andover, D.K. pp. 1-14.
- [129] Reichardt, W., Mascarina, G., Padre, B., Doll, J., (1997): Microbial communities of continuously cropped, irrigated rice fields. – *Applied and Environmental Microbiology* 63, 233-238.
- [130] Roy, R, Kluber, H.D., Curd, R, (1997): Early initiation of methane production in anoxic rice soil despite the presence of oxidants. – *FEMS Microbiology Ecology* 24, 311-320.

- [131] Sass, R.L., Fischer Jr., F.M., Huang, Y., (2000): A process-based model for methane emission from irrigated rice fields: experimental basis and assumption. – *Nutrient Cycling in Agroecosystems* 58, 249-258.
- [132] Sass, R.L., Fisher Jr., F.M., (1997): Methane emission from rice paddies: A process studies summary. – *Nutrient Cycling in Agroecosystems* 49, 119-127.
- [133] Sass, R.L., Fisher Jr., F.M., (1994): CH₄ emission from paddy fields in the United States gulf coast area. In: Minami, C.K., Mosier, A., Sass, R.L., (eds.) CH₄ and N₂O: Global Emissions and Controls from Rice Fields and other Agricultural and Industrial Sources. NIAES Series 2, Tsukuba, Japan. pp. 65- 77.
- [134] Sathpathy, S.N., Rath A.K, Mishra S., Kumarswamy S., Ramkrishnan B., Adhya T.K., Sethunathan N., (1997): Effect of hexachlorocyclohexane on methane production and emission from flooded rice soils. – *Chemosphere* 34, 2663-2671.
- [135] Schimel, J., (2000): Rice, microbes and methane. – *Nature* 403, 375-377.
- [136] Schutz, H., Holzappel-Pschorn, A., Conrad, R., Rennenberg, H., Seiler, W., (1989): A three-year continuous record on the influence of day time season and fertilizer treatment on methane emission rates from an Italian rice paddy. – *Journal of Geophysics Research* 94, 16405-16416.
- [137] Sebacher, D.T., Hams, R.C., Bartlett, K.B., (1983): Methane flux across the air water interface air velocity effects. – *Tellus* 35, 1-10.
- [138] Seiler, W., Holzappel-Pschorn, A, Conrad, R., Scharffe, D., (1984): Methane emission from rice paddies. – *Journal of Atmospheric Chemistry* 1, 241-268.
- [139] Sethunathan, N, Kumaraswamy, S., Rath, AK, Ramakrishnan, B., Satpathy, S.N., Adhya T.K, Rao, V.R, (2000): Methane production, oxidation and emission from Indian rice soils. – *Nutrient Cycle in Agroecosystems* 58, 377-388.
- [140] Shima, S.L., (1998): Mechanism of biological methane formation: Structure and function of methyl-coenzyme M. reductase. *Protein, Nucleic Acid and Enzyme* 43, 1461-1467.
- [141] Shin, Y.K, Yun, S.H., (2000): Varietal differences in methane emission from Korean rice cultivars. – *Nutrient Cycling in Agroecosystems*. 58, 315-319.
- [142] Sinha, S.K, (1995): Global methane emission from rice paddies: Excellent methodology but poor extrapolation. – *Current Science* 68, 643-646.
- [143] Steudler, A.P., Bowden, R.D., Mellilo, J.M., and Aber, J.D., (1989): Influence of nitrogen fertilization on methane uptake in temperate forest. – *Nature* 314-316.
- [144] Sundh, I., Borga, P., Nilsson, M and Svensson, B.H. (1995): Estimation of cell number of methanotrophic bacteria in boreal peatlands based on analysis of specific phospholipid fatty acids. – *FEMS Microbiology Ecology* 18, 103-112.
- [145] Thurlow, M., Karda, K.I., Tsurula, H., Minami, K, (1995): Methane uptake by unflooded paddy soils: The influence of soil temperature and atmospheric methane concentration. – *Soil Science and Plant Nutrition* 41, 371-375.
- [146] Topp, E., (1993): Effects of selected agrochemicals on methane oxidation by an organic agricultural soil. – *Canadian Journal of Soil Science* 73, 287-291.
- [147] Topp, E., Pattey, E., (1997): Soil as a source and sinks for atmospheric methane. – *Canadian Journal of Soil Science* 77, 167-178.
- [148] Trotsenko, Y.A., Khmelenina, V.N., (2002): Biology of extremophilic and extremotolerant methanotrophs. – *Archives of Microbiology* 177, 123-131.
- [149] Van Bodegom, P.M., Stams, A.J.M., (1999): Effect of alternative electron acceptors and temperature on methanogenesis in rice paddy soils. – *Chemosphere* 39, 167-182.
- [150] Van Bodegom, P.M., Stams, F., Mollema, L, Boeke, S., Leffelaar, P., (2001): Methane oxidation and the competition for oxygen in the rice rhizosphere. – *Applied and Environmental Microbiology* 67, 3586-3597.
- [151] Wang, B., Adachi, K., (2000): Differences among rice cultivars in root exudation, methane oxidation and population of methanogenic and methanotrophic bacteria in relation to methane emission. – *Nutrient Cycling in Agroecosystems* 58, 349-356.

- [152] Wang, Z.P., Delaune, R.D., Masscheleyn, P.B., Patrick Jr., W.H., (1993): Soil redox and pH effects on methane production in a flooded rice soils. – *Soil Science Society of American Journal* 57, 382-385.
- [153] Wang, Z.P., Zeng, D. Patrick Jr. W.H., (1997): Characteristics of CH₄ oxidation in a flooded rice profile. – *Nutrient Cycling in Agroecosystems*, 49, 97-103.
- [154] Wassmann, R., Lantin, R.S., Neu, H.D., Buendia, L.V., Corton, T.M., Lu, Y., (2000): Characterization of methane emissions from rice fields in Asia. III Mitigation options and future research needs. – *Nutrient Cycling in Agroecosystems* 58, 23-36.
- [155] Wassmann, R., Martius, C.S., (1997): Methane emission from the Amazon flood plain. In: Junk, W.J., (Ed.), *The Central Amazon floodplain: Ecological Studies* 126.. Springer-Verlag, Berlin. pp. 137-143.
- [156] Wassmann, R., Neue, H.D., Bueno, C., Latin, R.S., Alberto, MCR, Buendia, L.V., Bronson, K., Papen, H., Rennenberg, H., (1998): Methane production capacities of different rice soils derived from inherent and exogenous substrates. – *Plant and Soil* 203, 227-237.
- [157] Watanabe, A., Kajiwara, M., Tashiro, T., Kimura, M., (1995): Influence of rice cultivar on methane emission from paddy fields. – *Plant and Soil* 17, 51-56.
- [158] Watanabe, D., Hashimoto, T., Shimoyama, A., (1997): Methane oxidizing activities and methanotrophic population associated with wetland rice plants. – *Biology and Fertility of Soils* 24, 261-265.
- [159] Whalen, S.L., Reelourgh, W.S., Sandbes, K.A., (1990): Rapid methane oxidation in a landfill cover soil. – *Applied and Environmental Microbiology* 6, 3405-3411.
- [160] Yagi, K., Chairaj, P., Tsurata, H., Cholitkul, W., Minami, K., (1994): Methane emission from rice paddy fields in the central plain of Thailand. – *Soil Science and Plant Nutrition* 40, 29-37.
- [161] Yagi, K., Minami, K., (1990): Effects of organic matter application on methane emission from some Japanese paddy fields. – *Soil Science and Plant Nutrition* 36, 599-610.
- [162] Yao, H., Conrad, R., (1999): Thermodynamics of methane production in different rice paddy soils from China, the Philippines and Italy. – *Soil Biology and Biochemistry* 31, 463-473.
- [163] Yao, H., Conrad, R., (2001): Thermodynamics of propionate degradation in anoxic paddy soil from different rice-growing regions. – *Soil Biology and Biochemistry*. 33, 359-364.
- [164] Yao, H., Conrad, R., Wassmann, R., Neue, H.D., (1999): Effect of soil characteristic on sequential reduction and methane production in sixteen rice paddy soils from China, the Philippines, Italy. – *Biogeochemistry* 47, 269-295.