

## Microbial tannases: advances and perspectives

Cristóbal N. Aguilar · Raúl Rodríguez ·  
Gerardo Gutiérrez-Sánchez · Christopher Augur ·  
Ernesto Favela-Torres · Lilia A. Prado-Barragan ·  
Ascensión Ramírez-Coronel ·  
Juan C. Contreras-Esquivel

Received: 30 January 2007 / Revised: 14 April 2007 / Accepted: 15 April 2007 / Published online: 26 May 2007  
© Springer-Verlag 2007

**Abstract** In the last years, tannase has been the subject of a lot of studies due to its commercial importance and complexity as catalytic molecule. Tannases are capable of hydrolyzing complex tannins, which represent the main chemical group of natural anti-microbials occurring in the plants. The general outline of this work includes information of the substrates, the enzyme, and the applications. This review considers in its introduction the concepts and history of tannase and explores scientific and technological aspects. The “advances” trace the route from the general, molecular, catalytic, and functional information obtained under close to optimal conditions for microbial production through purification, description of the enzyme properties, and the commercial applications to the “perspectives” including expression studies, regulation, and potential uses;

aspects related to the progress in our understanding of tannin biodegradation are also included.

**Keywords** Tannase · Tannins biodegradation · Microbial sources · Advances · Perspectives

### Introduction

Tannin acyl hydrolase (E.C. 3.1.1.20) is commonly referred as tannase, an enzyme accidentally discovered by Tieghem (1867) in an experiment of formation of gallic acid into an aqueous solution of tannins, where grew two fungal species later identified as *Penicillium glaucum* and *Aspergillus niger* (Lekha and Lonsane 1997). Tannase catalyses the hydrolysis reaction of the ester bonds present in the gallotannins, complex tannins, and gallic acid esters. The enzyme is used in food and beverage processing; however, the practical use of this enzyme is at present limited due to insufficient knowledge about its properties, optimal expression, and large-scale application.

The major commercial applications of the tannases reside in the elaboration of instantaneous tea or of acorn liquor and in the production of gallic acid (Coggon et al. 1975; Chae and Yu 1983; Pourrat et al. 1985; Lekha and Lonsane 1997; García-Nájera et al. 2002; Belmares et al. 2004), the latter being an important intermediary compound in the synthesis of the antibacterial drug, trimethoprim, used in the pharmaceutical industry (Sittig 1988) and also in the food industry; gallic acid is a substrate for the chemical or enzymatic synthesis of propyl gallate, a potent antioxidant. Moreover, tannase is used as a clarifying agent in some wines, beers, fruit juices, and in refreshing drinks with coffee flavour (Lekha et al. 1993; Lekha and Lonsane 1994; Belmares et al. 2004). Its commercial production is

---

C. N. Aguilar (✉) · R. Rodríguez · J. C. Contreras-Esquivel  
Food Research Department, School of Chemistry,  
Universidad Autónoma de Coahuila,  
Blvd. Venustiano Carranza and J. Cardenas s/n,  
Col. Republica Oriente,  
25280 Saltillo, Mexico  
e-mail: cag13761@mail.uadec.mx

G. Gutiérrez-Sánchez  
Complex Carbohydrate Research Center, University of Georgia,  
Athens, GA 30602, USA

C. Augur  
IRD–Unité Biotrans IMEP Case 441 Faculté des Sciences de  
Saint Jérôme, Université Paul Cézanne,  
Av. Escadrille Normandie-Niemen,  
13397 Marseille Cedex 20, France

E. Favela-Torres · L. A. Prado-Barragan · A. Ramírez-Coronel  
Department of Biotechnology,  
Universidad Autónoma Metropolitana,  
Iztapalapa 09340 DF, Mexico

through microorganisms using submerged culture (SmC), where it is intracellularly expressed, implying additional costs during its production (Lekha and Lonsane 1994; Belmares et al. 2004). Tannase is commercialized by Biocon (India), Kikkoman (Japan), ASA special enzyme GmbH (Germany), and JFC GmbH (Germany) with different catalytic units depending on the product presentation. Tannases from Kikkoman and Biocon are produced by solid state culture (SSC).


An excellent review about the production and applications of tannase was reported by Lekha and Lonsane (1997), summarizing a lot of useful information generated in almost 130 years of research. During the last 10 years, novel and interesting information has been published. This review pretends to present and discuss these contributions to know the advances and perspectives in the knowledge and application of the microbial tannases.

### Substrates

Tannins are naturally occurring polyphenolic compounds with varying molecular weights that occur naturally in the plant kingdom. These phenolic compounds differ from others by having the ability to precipitate proteins from solutions. In the plant kingdom, these tannins are found in leaves, bark, and wood. Tannins are considered to be plants' secondary metabolic products because they play no direct role in plant metabolism. After lignin, tannins are the second most abundant group of plant phenolics. One of the major characteristics of tannins is their ability to form strong complexes with protein and other macromolecules such as starch, cellulose, and minerals (Lekha and Lonsane 1997; Aguilar and Gutiérrez-Sánchez 2001). It is widely

accepted that tannins are divided into four major groups: gallotannins, ellagitannins, condensed tannins, and complex tannins (Fig. 1). Gallotannins are characterized by the presence of several molecules of organic acids, such as gallic, digallic, and chebulic acids, esterified to a molecule of glucose. On the other hand, ellagitannins have building blocks of ellagic acid units linked to glucosides. Molecules with a core of quinic acid instead of glucose have also been considered as ellagitannins. To maintain its binding capacity, gallotannins, and ellagitannins must have more than two acidic unit constituents esterified to the glucose core. Gallotannins can easily be hydrolysed under mild acid or alkaline conditions, either in hot water or enzymatically (López-Ríos 1984). Ellagitannins are more stable than gallotannins. Condensed tannins or proanthocyanidins are complex compounds made of flavonoid building blocks (from 2 over 50) that are not considered to be easily hydrolyzable (Ramirez-Coronel et al. 2004). Among the major constituents are catechin derivatives such as cyanidin and delphinidin, which are responsible for the astringent taste of fruit and wines (Sanchez 2001). The negative effect of tannins relates not only to taste but also directly to animal nutrition with the known capacity of tannins to bind macromolecules, rendering them indigestible (Mendez 1984; Goel et al. 2005). This results in the formation of stable complexes with enzymes and minerals otherwise required by ruminal microorganisms (Goel et al. 2007). The bitter taste also is the reason why the feed intake is reduced. However, low tannin concentrations in feed have been shown to result in an increase in nitrogen assimilation in ruminants, rendering higher growth rates and milk production (Nip and Burns 1969). Complex tannins can be generated through reactions between gallic or ellagic acids with catechins and glucosides.

**Fig. 1** Main chemical structures of the tannins




Tannins have a range of effects on various organisms—from toxic effects on animals to growth inhibition of microorganisms. Some microbes are, however, resistant to tannins and have developed various mechanisms and pathways for tannin degradation in their natural habitat. Microbial degradation of condensed tannins is, however, less documented than that of gallotannins in both aerobic and anaerobic environments (Bhat et al. 1998). Condensed tannins are not hydrolysed by “classical tannases” (Contreras-Dominguez et al. 2006), with initial degradation steps carried out by mono- or di-oxygenases. However, further studies are needed to further characterize the degradation of condensed tannins. Tannases generally act on gallotannins. However, in the particular case of ellagitannins, the information is scarce and confuse (Scalbert 1991; Vivas et al. 2004), mainly due to their chemical complexity and diversity of kinds of ellagitannins. Saavedra et al. (2005) reported that the production of ellagic acid has not been explored due to its high production cost and the great amount of subproducts generated as a result of ellagitannin biodegradation. This results in serious problems regarding the recovery and purification of ellagic acid. To date, there is no published data on ellagitannin degradation using biological methods (microbial or enzymatic). However, it is known that the selective hydrolysis of galloyl groups of the ellagitannin phyllanthin is catalyzed by tannase (Zhang et al. 2001). Yoshida et al. (1999) reported the production of nobotannin K from complex ellagitannins through catalysis of tannases. However, the biochemical mechanism was not clearly explained.

Vattem and Shetty (2002, 2003) reported on ellagic acid production from cranberry pomace fermented by a SSC using *Lentinus edodes*, attributing the catalysis to the enzyme  $\beta$ -glucosidase. Huang et al. (2005) described a new valonea tannin hydrolase as responsible for the biodegradation of valonea tannins. This enzyme is itself a tannin acyl hydrolase.

Mingshu et al. (2006) reported in their review information generated by Vaquero et al. (2004) on tannase production by species of *Lactobacillus*, *Leuconostoc*, *Oenococcus*, and *Pediococcus*, using as carbon sources gallotannins, ellagitannins, and condensed tannins present in muscadine grapes. However, Vaquero et al. (2004) only considered the tannase production by lactobacilli without evaluation of ellagic acid released from ellagitannins.

Ramirez-Coronel et al. (2004) purified a thermostable bi-functional tannases from *Aspergillus*. The enzyme also had  $b$ -glucosidase activity. Unpublished results from the same group indicated that, in the presence of both tannic acid and cellobiose as substrates and of the purified enzyme, only the glucosidase activity was present. While studying the degradation of ellagitannins, Vattem and Shetty (2003) also observed glucosidase activity. In addition, Lee and Talcott (2005) incubated ellagitannins from muscadine grape seeds with commercial tannase and  $\alpha$ -glucosidase, and they found that tannase activity was not expressed, while  $\alpha$ -glucosidase showed high reactivity to ellagitannins. However, further studies on the subject are required to understand the microbial degradation process of such molecules.

**Fig. 2** Mechanism of action of tannase



## Enzyme

Tannase catalyses the breakdown of hydrolyzable tannins such as tannic acid, methyl gallate, ethyl gallate, *n*-propylgallate, and isoamyl gallate (Fig. 2). It is well known that tannase hydrolyses the ester bonds of tannic acid, although tannic acid is known to denature proteins. Tannase hydrolyses tannic acid completely to gallic acid and glucose through 2,3,4,6,-tetragalloyl glucose and two kinds of monogalloyl glucose. This is supported by the facts that the same products can be detected in the hydrolysate of 1,2,3,4,6,-pentagalloyl glucose and that gallic acid of methyl.*m*.digallate is liberated first.

Albertse (2002) reported that for a true enzyme-substrate complex to form, the criteria to consider are the following: (a) There should be no restriction on the structure of an alcohol composing a substrate ester, although the acid should be gallic acid; (b) any phenolic hydroxyl might react with the binding site of the enzyme and prevent the enzyme from forming a true enzyme-substrate (ES) complex; and (c) an ester bond or carboxyl does not link to the enzyme by itself because an ester or carboxylic compound is not hydrolyzed by or inhibits the enzyme unless it has phenolic hydroxyls.

Although tannase is present in plants, animals, and microorganisms, it is mainly produced by the latter. Tannase is produced by bacteria, yeasts, and fungi (Table 1).

The production and applications of tannase have been extensively studied; researches related to strain isolation and improvement, process development, and application of tannases have resulted in a great number of scientific publications and patents. Table 2 presents some of the published patents regarding tannase production and application.

Filamentous fungi of the *Aspergillus* genus and bacteria of the *Bacillus* genus have been widely used for tannase production (Mondal et al. 2001a, b; Pinto et al. 2001). Although tannase production by *Aspergillus* can occur in the absence of tannic acid, this fungus (mainly *A. niger*) tolerates tannic acid concentrations as high as 20% without having a deleterious effect on both growth and enzyme production (Van Diepeningen et al. 2004; Cruz-Hernández et al. 2006). Studies on tannase production by *Aspergillus* have been carried out in submerged and solid-state cultures. Depending on the strain and the culture conditions, the enzyme is induced and expressed with different levels of activity, showing different production patterns. Phenolic compounds such as gallic acid, pyrogallol, methyl gallate, and tannic acid induces tannase synthesis (Bajpai and Patil 1997). However, the induction mechanism has not been clearly demonstrated, and there is some controversy about the role of some of the hydrolyzable tannin constituents as related to the synthesis of tannase (Deschamps et al. 1983; Aguilar et al. 2001a). For instance, gallic acid, one of the

**Table 1** Microbial sources of tannase

Microorganism	Reference
<b>Bacteria</b>	
<i>Achromobacter</i> sp.	Lewis and Starkey (1969)
<i>Bacillus pumilus</i>	Deschamps et al. (1983)
<i>Bacillus polymyxa</i>	Deschamps et al. (1983)
<i>Corynebacterium</i> sp.	Deschamps et al. (1983)
<i>Bacillus cereus</i>	Mondal et al. (2001b)
<i>Klebisella planticola</i>	Deschamps et al. (1983)
<i>Klebisella pneumoniae</i>	Deschamps et al. (1983)
<i>Pseudomonas solanaceanum</i>	Deschamps et al. (1983)
<i>Streptococcus bovis</i>	Belmares et al. (2004)
<i>Streptococcus gallolyticus</i>	Sasaki et al. (2005)
<i>Lactobacillus plantarum</i>	Ayed and Hamdi (2002); Kostinek et al. (2007)
<i>Lactobacillus paraplantarum</i>	Nishitani and Osawa (2003); Nishitani et al. (2004)
<i>Lactobacillus pentosus</i>	Nishitani et al. (2004); Kostinek et al. (2007)
<i>Lactobacillus acidophilus</i>	Nishitani et al. (2004); Sabu et al. (2006)
<i>Lactobacillus animalis</i>	Nishitani et al. (2004)
<i>Lactobacillus murinus</i>	Nishitani et al. (2004)
<i>Enterococcus faecalis</i>	Goel et al. (2005)
<i>Weissella paramesenteroides</i>	Kostinek et al. (2007)
<i>Leuconostoc fallax</i>	Kostinek et al. (2007)
<i>Leuconostoc mesenteroides</i>	Kostinek et al. (2007)
<i>Pediococcus acidilactici</i>	Nishitani et al. (2004)
<i>Pediococcus pentosaceus</i>	Nishitani et al. (2004)
<i>Citrobacter freundii</i>	Belmares et al. (2004)
<i>Selenomonas ruminantium</i>	Belmares et al. (2004)
<b>Yeasts</b>	
<i>Candida</i> sp.	Aoki et al. (1976)
<i>Saccharomyces cerevisiae</i>	Zhong et al. (2004)
<i>Mycotorula japonica</i>	Belmares et al. (2004)
<i>Pichia</i> spp.	Deschamps et al. (1983)
<i>Debaryomyces hansenii</i>	Deschamps et al. (1983)
<b>Fungi</b>	
<i>Aspergillus niger</i>	Bradoo et al. (1996); Rana and Bhat (2005); Cruz-Hernandez et al. (2006); Treviño-Cueto et al. (2007); Murugan et al. (2007)
<i>Aspergillus japonicus</i>	Bradoo et al. (1997)
<i>Aspergillus gallonyces</i>	Belmares et al. (2004)
<i>Aspergillus awamori</i>	Bradoo et al. (1996); Mahapatra et al. (2005)
<i>Aspergillus fumigatus</i>	Batra and Saxena (2005)
<i>Aspergillus versicolor</i>	Batra and Saxena (2005)
<i>Aspergillus flavus</i>	Yamada et al. (1968); Batra and Saxena (2005)
<i>Aspergillus caespitosum</i>	Batra and Saxena (2005)
<i>Aspergillus oryzae</i>	Bradoo et al. (1996)
<i>Aspergillus aculeatus</i>	Banerjee et al. (2001)
<i>Aspergillus Aureus</i>	Bajpai and Patil (1997)
<i>Aspergillus fischeri</i>	Bajpai and Patil (1997)

**Table 1** (continued)

Microorganism	Reference
<i>Aspergillus rugulosus</i>	Bradoo et al. (1996)
<i>Aspergillus terreus</i>	Bajpai and Patil (1997)
<i>Aspergillus foetidus</i>	Banerjee et al. (2005)
<i>Penicillium notatum</i>	Ganga et al. (1977)
<i>Penicillium islandicum</i>	Ganga et al. (1977)
<i>Penicillium chrysogenum</i>	Bradoo et al. (1996)
<i>Penicillium digitatum</i>	Bradoo et al. (1996)
<i>Penicillium acrellanum</i>	Bradoo et al. (1996)
<i>Penicillium caryophilum</i>	Bradoo et al. (1996)
<i>Penicillium citrinum</i>	Bradoo et al. (1996)
<i>Penicillium charlessii</i>	Bradoo et al. (1996); Batra and Saxena (2005)
<i>Penicillium variable</i>	Batra and Saxena (2005)
<i>Penicillium glaucum</i>	Lekha and Lonsane (1997)
<i>Penicillium crustosum</i>	Batra and Saxena (2005)
<i>Penicillium restrictum</i>	Batra and Saxena (2005)
<i>Penicillium glabrum</i>	Van de Lagemaat and Pyle (2005)
<i>Trichoderma viride</i>	Bradoo et al. (1996)
<i>Trichoderma hamatum</i>	Bradoo et al. (1996)
<i>Trichoderma harzianum</i>	Bradoo et al. (1996)
<i>Fusarium solani</i>	Bradoo et al. (1996)
<i>Fusarium oxysporium</i>	Bradoo et al. (1996)
<i>Mucor</i> sp.	Belmares et al. (2004)
<i>Paecilomyces variotii</i>	Mahendran et al. (2005); Battestin and Alves-Macedo (2007)
<i>Rhizopus oryzae</i>	Hadi et al. (1994); Purohit et al. (2006)
<i>Cryphonectria parasitica</i>	Farias et al. (1994)
<i>Heliocostylum</i> sp.	Bradoo et al. (1996)
<i>Cunninghamella</i> sp.	Bradoo et al. (1996)
<i>Syncephalastrum racemosum</i>	Bradoo et al. (1996)
<i>Neurospora crassa</i>	Bradoo et al. (1996)

structural constituents of some hydrolyzable tannins, such as tannic acid, has been reported as an inducer of tannase synthesis under submerged fermentation, while it represses tannase synthesis under solid-state fermentation. Nevertheless, independent of the involved mechanism, it has been well accepted that, due to the complex composition of the hydrolyzable tannins, some of their hydrolysis products induce tannase synthesis (Aguilar et al. 2002).

The addition of carbon sources such as glucose, fructose, sucrose, maltose, and arabinose to the culture medium at initial concentrations from 10 to 30 g/l improves tannase production by *A. niger* (Bradoo et al. 1997). Nitrogen requirements can be supplied by different organic and inorganic sources. Inorganic nitrogen can be supplemented as ammonium salts (sulphate, carbonate, chloride, nitrate, monohydrated phosphate) or nitrate salts (sodium, potassium, or ammonium). Other nutritional requirements such as potassium, magnesium, zinc, phosphorous, and sulfur are supplied as salts. Although *A. niger* does not require

**Table 2** Published patents regarding tannase production and application

Year	Title	Patent no.
1974	Conversion of green tea and natural tea leaves using tannase	USP3812266
1975	Production of tannase by <i>Aspergillus</i>	JP7225786
1975	Tea soluble in cold water	UKP1280135
1976	Extraction of tea in coldwater	GP2610533
1976	Enzymatic solubilization of tea cream	USP3959497
1985	Gallic acid ester(s) preparation	EP-137601
1985	Preparation of gallic acid esters, e.g., propylgallate	EP-137601
1985	Enzymatic treatment of black tea leak	EP135222
1987	Preparation of tannase	JP62272973
1987	Manufacturing of tannase with <i>Aspergillus</i>	JP62272973
1988	Production of tannase by <i>Aspergillus oryzae</i>	JP63304981
1988	Elaboration of tannase by fermentation	JP63304981
1989	Preparation of spray-concrete coating in mining shaft	SUP1514947
1989	Antioxidant catechin and gallic acid preparation	JP01268683
1989	Tannase production by culture of <i>Aspergillus tamaritii</i>	EP-339011
1989	New <i>Aspergillus niger</i> B1 strain	EP307071
1989	Tannase production process by <i>Aspergillus</i> and its application to obtain gallic acid	EP339011
1992	Tannase preparation method	JP4360684
1995	Enzymatic clarification of tea extracts	USP5445836
1997	DNA fragment containing a tannases gene, a recombinant plasmid, a process for producing tannases, and a promoter	USP5665584
2000	Tea concentrate prepared by enzymatic extraction and containing xanthan gum that is stable at ambient temperature	USP6024991
2000	Producing theaflavin	USP6113965
2004	Compositions based on vanilloid–catechin synergies for	USP6759064
2006	Diagnostic agent and test method for colon cancer using tannases as index	USP7090997
2006	Isolation of a dimmer di-gallate a potent endothelium-dependent vasorelaxing compound	USP7132446

cofactor supplementation, folic and pantothenic acid are eventually added to the cultured medium (Belmares et al. 2004).

Tannase production in submerged culture by *Aspergillus* sp. is improved at high aeration rates. It is favored at 30–33°C and initial pH values from 3.5 to 6.5. The maximal enzymatic activity is attained after 1 to 3 days of cultivation. Tannase production has been mostly studied in submerged fermentation. Strain improvement for tannase production from co-culture of *A. foetidus* and *Rhizopus oryzae* was reported by Purohit et al. (2006) using ultraviolet light, heat, and 3-nitro,5-methylguanidine mutagenesis.

However, several studies have reported interesting advantages between the tannase produced by SSC in relation with that produced by SmC. On this topic, several reports have been published (Barthomeuf et al. 1994; García-Peña 1996; Chaterjee et al. 1996; Lekha and Lonsane 1997; García-Peña et al. 1999; Ramírez-Coronel et al. 1999; Aguilar et al. 1999, 2001a and b, 2002; Kar et al. 2002; Viniegra-González et al. 2003; Van de Lagemaat and Pyle 2001, 2005; Rana and Bhat 2005; Pinto et al. 2006). In those, attractive advantages were indicated: the high-production titles (up to 5.5 times more than in SmC), the extracellular nature of the enzymes, and the stability towards wide pH and temperature ranges (Lekha and Lonsane 1994). Aguilar et al. (1999) reported productivities of 6.667 and 1.275 UE/Lh for SSC and SmC, respectively. Maximum tannase activity expressed intracellularly is also 18 times more in SSC than in SmC, while the extracellular activity is 2.5 times higher in SSC than in SmC.

It is important to note that such higher tannase activity levels in SSC than SmC have been clearly associated with the concomitant production of proteolytic activities in the latter culture system (Aguilar et al. 2002; Viniegra-González et al. 2003). Also, in SSC, the tannase produced exhibits a higher tolerance to wide range of pH and temperature (Lekha and Lonsane 1994; Rana and Bhat 2005).

Table 3 shows the materials used as supports of SSC for tannase production. Also, it presents the tannin-rich materials employed in both culture systems (SmC and SSC) for the production of the microbial enzyme.

For optimization of tannase production, Pinto et al. (2003) evaluated the tannic acid/wheat bran ratio, different moisture levels, addition of supplementary nitrogen sources, addition of supplementary phosphate, and concentration of supplementary nitrogen and phosphate added to the medium. Their results showed that the best medium was with 15% of tannic acid, 37.5% of initial moisture, 1.7 ammonium sulphate, and 2.0% of sodium phosphate. The presence of phosphate was of great importance for optimization because it promoted the increase in the synthesis level and a very expressive decrease in the maximum production time, from 72 to 24 h of fermentation. The optimized process promoted an increase of 861% in yield and 2783% in productivity.


Hatamoto et al. (1996, 1997) cloned and sequenced the gene-encoding tannase, and a structural study of the enzyme subunit from *A. oryzae* gave the possibility to manipulate the producer systems to increase and to improve the levels of tannase activity. In that study, authors reported that gene sequence did not have any introns. They found that the gene code for a 588-amino acid sequence with an 18-amino acid signal sequence and a molecular weight of approximately 64,000 Da. They hypothesized that their tannase consisted out of two subunits with molecular

**Table 3** Materials used as supports of SSC for tannase production and tannin-rich materials used as enzyme inducer in SSC and SmC

Traditional supports	Reference
Sugarcane bagasse	Lekha and Lonsane (1994); García-Peña et al. (1999)
Wheat bran	Chaterjee et al. (1996); Sabu et al. (2005)
Polyurethane foam	Ramírez-Coronel et al. (1999); Aguilar et al. (2001b); Van de Lagemaat and Pyle (2001, 2005)
Tamarind seed powder	Sabu et al. (2005)
Palm kernel cake	Sabu et al. (2006)
Tannin rich materials	
Chestnut bark	Deschamps et al. (1983)
Tara ( <i>Caesalpinia spinosa</i> ) tannins	Pourrat et al. (1985)
Gall nuts ( <i>Quercus infectoria</i> )	Barthomeuf et al. (1994)
<i>Rhus coriaria</i> leaves	Barthomeuf et al. (1994)
Fruits of <i>Terminalia chebula</i>	Banerjee et al. (2005)
Pod cover of <i>Caesalpinia digyna</i>	Banerjee et al. (2005)
<i>Quercus aegylops</i> tannins	Shi et al. (2005)
Jawar leaves ( <i>Sorghum vulgare</i> )	Kumar et al. (2006)
Amla leaves ( <i>Phyllanthus emblica</i> )	Kumar et al. (2006)
Jamun leaves ( <i>Syzygium cumini</i> )	Kumar et al. (2006)
Ber leaves ( <i>Zyzyphus mauritiana</i> )	Kumar et al. (2006)
Creosote bush leaves	Treviño-Cueto et al. (2007)

weights of 30,000 and 33,000 Da linked by a disulfide bond. The tannase gene was transcribed as a single polypeptide chain, after which, the 18 amino acid signal sequence was cleaved off, and the polypeptide chain was cleaved into two subunits. It was reported that the single polypeptide chain was cleaved by a KEX-II-like protease in two separate polypeptide chains. They concluded that native tannase consisted of four pairs of the two subunits, forming a hetero-octamer with a molecular weight of about 300,000 Da (Fig. 3).

Albertse (2002) sequenced the genes of positive isolates of *Aspergillus* and expressed a tannase gene in *Saccharomyces cerevisiae*, evaluating the characterization and comparison of the recombinant tannase with published properties of other tannases. However, the recombinant tannase was expressed in very low amounts, probably due to the phenomenon exhibited in yeasts to hyper-glycosylate glycoproteins, therefore, expressing proteins that are often catalytically inactive or different than the native enzymes.



**Fig. 3** Descriptive scheme of the proposed post-translational modification of tannase precursor (Hatamoto et al. 1996)

Recently, Cerda-Gomez et al. (2006) reported the use of conserved tannase gene sequences from five different *Aspergillus* species (Fig. 4) to design a set of primers (Tan1, Tan2), which later were used to amplify by polymerase chain reaction (PCR) a DNA segment of 435 bp from four different *Aspergillus* strains (Fig. 5). Zhong et al. (2004) reported the heterologous expression of *A. oryzae* tannase, suggesting that the *Pichia pastoris* system represents an attractive means of generating large quantities of tannases (7,000 IU/l) for both research and industrial purpose.

On the other hand, tannase extraction strongly depends on the fermentation system used. As tannase is mostly extracellular when produced by SSC, it can be easily extracted with water or a buffer. Two to three volumes of the agent extraction is well mixed with the fermented mass and pressed to obtain the enzymatic extract. Tannase location during its production by SmC depends on the cultivation time (Rajkumar and Nandy 1983). It is mainly intracellular at the beginning of the culture, and it is further secreted to the culture medium. However, up to 80% of tannase remained bound to the mycelium when the maximum overall tannase titer is attained. Bound tannase can be extracted after cell-wall hydrolysis with digestive enzymes such as chitinase.

The cells can also be mechanically disrupted to recover the bound tannase. An extracellular tannase produced by solid-state cultures of *A. niger* was purified to homogeneity


from the cell-free culture broth by preparative isoelectric focusing and by fast protein liquid chromatography (FPLC) using anion-exchange and gel-filtration chromatography (Ramirez-Coronel et al. 2003). SDS-PAGE analysis, as well as gel localization studies of purified tannase, indicated the presence of two enzyme forms.

Tannase produced by *A. awamori nakazawa* was purified and characterized by Mahapatra et al (2005). Tannase from this new isolate exhibited optimum activity at 35°C and at a pH of 5.0. Urea concentrations higher than 3 M were inhibitory. Increasing concentrations of sodium lauryl sulfate (SLS) also led to decrease in activity. Two percent SLS was inhibitory. Increasing concentrations of ethylenediaminetetraacetic acid (EDTA) had an inhibitory effect on tannase. Tannase was found to be a glycoprotein. Circular dichroism analysis of purified fractions of tannase indicates that the β-sheet structure in tannase was predominant, indicating its globular nature.

Because a fraction of the produced tannase remains bound to the cell under submerged fermentation conditions, the produced biomass may be recycled as a biocatalyst. Several strategies can be used for tannase concentration or purification and immobilization after extraction from the biomass (submerged fermentation) or from the culture medium (submerged solid state fermentation). To increase the specific activity of the enzymatic preparation, tannase should be concentrated. For that, classical methods such as salt or solvent precipitation, ultrafiltration followed by ion

**Fig. 4** Tannase gene sequence from five different *Aspergillus* species used for primer design (Cerda-Gomez et al. 2006)

```
ATTACGAGTGC GGGTCTTTGGGCCACCTCCCATCCGTGTCTATTATACCCTGTT
GCTTCGGCGGGCCCGCCGGRTAGAACTGGTACCACTTGTTCGGCCGCCGGGGGG
GCGCCTTTGCCCGCCGGCCCGTGCCTCGCGGAGACCCCAACACGAACACGAG
TGCGGGTCTTTGGGCCAACCTCCCATCCGTGTCTATTATACCCTGTTGCTTCGG
CGGGCCCGCCGCTTGTTCGGCCGCCGGGGGGGCGCCTTTGCCCGCCGGCCCGT
GCCCGCCGAGACCCCAACACGAACACTGTCTGAAAGCGTGCAGTCTGAGTTGAT
TGAATGCAATCAGTTAAACTTTCAACAATGGATCTCTTGGTTCCGGCATCGATGAA
GAACGCAGCGAAATGCGATAACTAATGTGAATTGCAGAATTCAGTGAATCATCGAG
TCTTTGAACGCACATTGCGCCCCCTGGTATTCCGGGGGGCATGCCTGTTTTACCT
CTCCTGGCAGTYGCAGGTTGACCTCGGATCAGGTAGGGATACCCGCTGAACTTAA
GCATATCCAATAAGCGGAAA
```



**Fig. 5** Amplification of a tannase gene segment (435 bp) from *Aspergillus* strains. Lanes 1 and 6 Molecular marker, lane 2 GH1 strain, lane 3 ESH strain, lane 4 PSH strain, and lane 5 GS strain (Cerdeira-Gomez et al. 2006)

exchange or size exclusion chromatography, as well as solvent extraction can be used (Lekha and Lonsane 1994).

Tannases of *Aspergillus* strains have a molecular weight around 150–350 kDa. Their activity and stability pH are 5–6.0 and 3.5–8.0, respectively, while optima temperatures from 35 to 40°C have been reported (Farias et al. 1994). Tannase is stable for several months at 30°C. Tannase produced by *Penicillium* strains present similar characteristics in terms of pH and temperature activity and stability. When tannic acid is used as substrate, Km values of 11.25 and 0.048 mM were obtained with *Aspergillus* and *Penicillium*, respectively (Yamada et al. 1968; Adachi et al. 1971; Ibuchi et al. 1972; Aoki et al. 1976; Chae and Yu 1983).

Ramirez-Coronel et al. (2003) produced by SSC, a single tannase present in monomeric and dimeric forms with molecular masses of 90 and 180 kDa. The tannase had an isoelectric point of 3.8, a temperature optimum of 60–70°C, and a pH optimum of 6.0. The substrate specificity of the tannase was determined by high performance liquid chromatography (HPLC) analysis of tannin substrates and products. The enzyme was able to remove gallic acid from both condensed and hydrolyzable tannins. Internal sequences were obtained from each of the gel-purified and trypsin-digested tannase forms. The peptide sequences obtained from both forms were identical to sequences within a  $\beta$ -glucosidase from *A. kawachii* (Fig. 6). The purified tannase was tested for  $\beta$ -glucosidase activity and was shown to hydrolyze cellobiose efficiently. However, no  $\beta$ -glucosidase activity was detected when the enzyme was assayed in the presence of tannic acid.

Tannase requires the presence of metal ions to express its full catalytic activity; in this sense, it is important to know the kind of ions and their concentrations in achieving maximal reaction efficiency. The effect of metal ions on tannase activity was studied by Kar et al. (2003). One mM  $Mg^{+2}$  or  $Hg^{+2}$  activated tannase activity.  $Ba^{+2}$ ,  $Ca^{+2}$ ,

$Zn^{+2}$ ,  $Hg^{+2}$ , and  $Ag^{+}$  inhibited tannase activity at 1.0 mM concentration, and  $Fe^{+3}$  and  $Fe^{+2}$  completely inhibited tannase activity.  $Ag^{+}$ ,  $Ba^{+2}$ , and  $Hg^{+2}$  competitively inhibited tannase activity (Mukherjee and Banerjee 2005). Among the anions studied, 1 mM bromide or  $S_2O_3$  enhanced tannase activity. Other chemical substances reported to play a role in the tannase activity are the surfactants and chelators. Kar et al. (2003) reported that palmitic acid and oleic acid enhanced tannase activity, whereas stearic acid inhibited tannase activity; also, SLS and Triton X-100 inhibited tannase activity. Urea stimulated tannase activity at a concentration of 1.5 M. Among the chelators, 1 mM EDTA or 1,10-*o*-phenanthroline inhibited tannase activity. Dimethyl sulfoxide and *b*-mercapto ethanol inhibited tannase activity at 1 mM concentration, whereas soybean extract inhibited tannase activity at concentrations varying from 0.05 to 1.0% (w/v). Among the nitrogen sources selected, ammonium ferrous sulfate, ammonium sulfate, ammonium nitrate, and ammonium chloride enhanced tannase activity at 0.1% (w/v) concentration.

## Applications

With respect to the uses of tannase, one of the major applications of tannase is in the manufacturing of instantaneous tea. Tannase applications in food and beverage industrial products contribute to remove the undesirable effects of tannins (Boadi and Neufeld 2001).

Enzymatic treatment of fruit juices to reduce the bitterness has got advantages such as the higher quality of juice due to the lower haze and non-deterioration of juice quality. New fruit juices (pomegranate, cranberry, raspberry, cold tea, etc.) have recently been acclaimed for their health benefits, in particular, for its disease-fighting antioxidant potential. The presence of high tannin content in those fruits is responsible for haze and sediment formation, as well as for color, bitterness, and astringency of the juice upon storage. Due to the inability of conventional fruit juice debittering processes to remove the bitterness effectively, enzymatic debittering should be preferred. Preliminary results reported by Rout and Banerjee (2006) for pomegranate juice demonstrated that tannase treatment resulted in 25% degradation of tannin, while a combination of tannase and gelatin (1:1) resulted in 49% of tannin degradation. However, further studies in this topic are needed.

Other important application of tannase is the production of gallic acid and propylgallate, fine-chemicals employed as antioxidants in foods, cosmetics, hair products, adhesives, and lubricant industry (Kar et al. 2002; Yu et al. 2004; Yu and Li 2006). Both antioxidants (gallic acid and propylgallate) have been biosynthesized in fungal SmC (Gaathon et



**Fig. 6** Alignment of the amino acid sequences obtained from the 90- and 180-kDa forms of the *A. niger* tannase with the deduced amino acid sequence of *A. kawachii*  $\beta$ -D-glucosidase (Glc; accession no. BAA19913, gi|2077896|). Thirteen sequences were obtained from the 180-kDa form (D180) and two sequences from the 90-kDa form (M90). Identical amino acids are shaded (Ramirez-Coronel et al. 2003)

Glc D180 M90	MRFTLIEAVA	LTAVSLASAD	ELAYSPPYYP	SPWANGQGDW	AQAYQRAVDI RAVDI	VSQMTLAEKV VSQMTLAEKV	60
Glc D180 M90	NLTTGTGWEL	ELCVGQTGGV	PRLGVPGMCL	QDSPLGVRDS	DYNSAFFPSGM	NVAATWDKNL NL	120
Glc D180 M90	AYLRGKAMGQ AYLRGKAMGQ GKAMGQ	EFSDKGADIQ EFSDK EFSDK	LGPAAGPLGR	SPDGGRNWEG NWEG	FSPDLAPSGV FSPDLAPSGV	LFAETIKGIQ LFAETIKGIQ	180
Glc D180 M90	DAGVVATKH DAGVVATK	YIAYEQEHR	QAPEAQCYGF	NISESGSANL	DDKTMHELYL TMHELYL	WPFADAIRAG WPFADAIR	240
Glc D180 M90	GDVDYDSGTS	YWGNTLTVSV	LNGTVPQWRV V	DDMAVRIMAA DDMAVRIMAA	YYKVRDRLN YYK LN	TPPNFSSWTR TPPNFSSWTR	360
Glc D180 M90	DEYGYKYYV DEYGYK	SEGPYEKVNH	YVNVQRNHSE	LIRRIGADST RIGADST	VLLKNDGALP VLLKNDGALP	LTGKERLVAL LTGK	420
Glc D180 M90	IGEDAGSNPY	GANGCSDRGC	DNGTLAMGWG	SGTANFPYLV	TPEQAISNEV	LKNKNGVFTA NGVFTA	480
Glc D180 M90	TDNWAIDQIE TDNWAIDQIE	ALAKTASVSL ALAKTASVSL	VFVNADSGEG VFVNADSGEG	YINVDGNLGD YINVDGNLGD	RKNLTLWRNG R	DNVIAAASN	540
Glc D180 M90	CNNTVIHIS	VGPVLVNEWY	DNPNTAILW	GGLPGQESGN	SLADVLYGRV	NPGAKSPFTW SPFTW	600
Glc D180 M90	GKTREAYQDY GK	LVTEPNNGNG	APQEDFVEGV	FIDYRGFDKR	NETPIYEFY	GLSYTTFNYS	660
Glc D180 M90	GDASYGQDSS	DYLPEGATDG	SAQPILPAGG	GPGGNPRLYD	ELIRVSVTIK	NTGKVAGDEV VAGDEV	780
Glc D180 M90	PQLYVSLGGP PQLYVSLGGP	NEPKIVLRQF NEPK	REITLQPSEE ITLQPSEE	TKWSTTLTRR TKWSTTLTR	DLANWNVEKQ	DWEITSYPKM	840
Glc D180 M90	VFGSSSRKP VFGSSSR	PLRASLPTVH ASLPTVH					

al. 1989; Kar and Banerjee 2000; García-Nájera et al. 2002; Sharma and Gupta 2003), in SSC (Kar and Banerjee 2000), and in modified SSC (Kar et al. 2002) using micro-encapsulated tannase in chitosan and as free enzyme (Yu et al. 2004), and by mycelium-bound tannase in organic solvent like 1-propanol (Yu and Li 2006). This last is an important contribution to discover the new reactions of tannase in non-conventional media because the industrial biocatalysis has grown considerably in recent years for its potential applications in fine-chemical, agrochemical, perfumery, flavor, pharmaceutical, and drug industries. Advantages of the tannase action of non-aqueous system lie in dramatically higher substrate solubility, the ability to use tannase synthetically rather than hydrolytically, and the capability to modify native selectivity by simply tailoring the reaction medium rather than the tannase itself.

Tannase has been also immobilized to reuse its catalytic action in agarose, chitosan, alginate, ceolite, and different derivatives of silicious materials can be used for tannase

immobilization (Sharma et al. 2002; Sharma and Gupta 2003; Mahendran et al. 2005). Abdel-Naby et al. (1999) immobilized tannase from *A. oryzae* on various carriers. However, the enzyme immobilized on chitosan glutaraldehyde showed the highest activity. The bound enzyme retained 20.3% of the original specific activity. On the other hand, Sharma et al. (2002) immobilized tannase from *A. niger* on concavalin A-sepharose via bioaffinity interaction. The immobilized preparation was quite stable to reuse; there was no loss of enzyme activity after three cycles, and it retained 81% activity even after the sixth cycle. Ester hydrolysis using the immobilized enzyme led to a 40% conversion into gallic acid as compared with 30% obtained with the free enzyme.

A stopped-flow manifold was developed to assay and characterize immobilized tannase. The immobilized enzyme reactor was inserted within the tube-type electrode pair (cell constant=103.2 cm<sup>-1</sup>) for a real-time conductometric measurement. Tris buffer (2 mM, pH=7.0) was used as

the carrier for sensitivity improvement. The activities and kinetic parameters ( $K_m$  values) for propyl gallate, methyl gallate, and tannic acid were reported by Chang et al. (2006). Raab et al. (2007) immobilized fungal tannase on Eupergit C to produce complex tannins based on catechins esterified with gallic acid.

The gallic acid is used in the pharmaceutical industry for the synthesis of antibacterial drugs and in the food industry as substrate for the chemical synthesis of food preservatives such as pyrogallol and gallates (García-Nájera et al. 2002).

Tannase may play an important role in plant cell-wall degradation by cleaving some of the cross-links existing between cell-wall polymers (García-Conesa et al. 2001). Tannase can be potentially used for the degradation of tannins present in the effluents of tanneries, which represent serious environmental problems (Van de Lagemaat and Pyle 2001; Gammoun et al. 2006). Also, it can be used in the preparation of animal feeding using as culture support the mycelial wastes from penicillin manufacture (Nuero and Reyes 2002). Bioactive polyphenolic compounds can be produced by tannases applied to lentil flours (Dueñas et al. 2007).

### Concluding remarks

Tannins are complex and diverse molecules in structure. It is, therefore, clear that “tannase” is a generic term that groups a number of enzymes from different organisms that are all able to hydrolyze tannins. The use of tannase from different microbial sources may have benefits for different areas such as food, beverage, cosmetic, and pharmaceutical industries, as well as environmental depollution. A tannase may be efficient on one substrate and not on another. For that, more effort is needed to develop specific tannases to answer specific needs. In this sense, solid-state fermentation presents more advantages than the submerged type of culture. However, new bioreactors open the possibility to improve the tannase productivity in both culture systems. Study of biodegradation of ellagitannins is a field that remains unclear. In addition, improvements on tannase immobilization are needed for the development of cheaper processes. For that, enzymatic preparations with improved catalytic characteristics must be developed, and new conditions of reaction should be tested. Expression studies have been conducted but more efforts are necessary to improve and optimize the production of tannase under new culture conditions. To fully elucidate the kinetics of the recombinant expressed tannase, future expression will be required in an established microbial expression host. Finally, studies on the structure and protein engineering should be conducted to enhance the levels of activity of tannase.

**Acknowledgment** C. N. Aguilar thanks CONACYT–SEP (project no. 42244) and COAH–CONACYT (COAH-2002-CO1.2565 and 4652) for financial support. The present work was conducted within the framework of the ECOS program (M02A02, project between Mexico–France).

### References

- Abdel-Naby MA, Sherif AA, El-Tanash AB, Mankarios AT (1999) Immobilization of *Aspergillus oryzae* tannase and properties of the immobilized enzyme. *J Appl Microbiol* 87:108–114
- Adachi O, Watanabe M, Yamada H (1971) Physicochemical properties of the tannase from *Aspergillus flavus*. *Agric Biol Chem* 32:1079–1085
- Aguilar CN, Augur C, Favela-Torres E, Viniegra-González G (2001a) Induction and repression patterns of fungal tannase in solid-state and submerged cultures. *Proc Biochem* 36:565–570
- Aguilar CN, Augur C, Favela-Torres E, Viniegra-González G (2001b) Production of tannase by *Aspergillus niger* Aa-20 in submerged and solid state fermentation: influence of glucose and tannic acid. *J Ind Microbiol Biotechnol* 26:296–302
- Aguilar CN, Augur C, Viniegra-González G, Favela-Torres E (1999) A comparison of methods to determine tannin acyl hydrolase activity. *Braz Arch Biol Technol* 42:355–361
- Aguilar CN, Favela-Torres E, Viniegra-González G, Augur C (2002) Culture conditions dictate protease and tannase production in submerged and solid-state cultures by *Aspergillus niger* Aa-20. *Appl Biochem Biotechnol* 102–103:407–414
- Aguilar CN, Gutiérrez-Sánchez G (2001) Review sources, properties, applications and potential uses of tannin acyl hydrolase. *Food Sci Technol Int* 7:373–382
- Albertse EK (2002) Cloning, expression and characterization of tannase from *Aspergillus* species. M.Sc. thesis, Faculty of Natural and Agricultural Sciences, Department of Microbiology and Biotechnology, University of the Free State Bloemfontein, South Africa
- Aoki K, Shinke R, Nishira H (1976) Purification and some properties of yeast tannase. *Agric Biol Chem* 40:79–85
- Ayed L, Hamdi M (2002) Culture conditions of tannase production by *Lactobacillus plantarum*. *Biotechnol Lett* 24:1763–1765
- Bajpai B, Patil S (1997) Introduction of tannin acyl hydrolase (EC 3.1.1.20) activity in some members of *fungi imperfecti*. *Enzyme Microb Technol* 20:612–614
- Banerjee D, Mondal K, Bikas R (2001) Production and characterization of extracellular and intracellular tannase from newly isolated *Aspergillus aculeatus* DBF9. *J Basic Microbiol* 6:313–318
- Banerjee R, Mukherjee G, Patra KC (2005) Microbial transformation of tannin-rich substrate to gallic acid through co-culture method. *Bioresour Technol* 96:949–953
- Barthomeuf C, Regerat F, Pourrat H (1994) Production, purification and characterization of tannase from *Aspergillus niger* LCF8. *J Ferment Technol* 77:137–142
- Batra A, Saxena RK (2005) Potential tannase producers from the genera *Aspergillus* and *Penicillium*. *Proc Biochem* 40:1553–1557
- Battestin V, Alves-Macedo G (2007) Tannase production by *Paecilomyces variotii*. *Bioresour Technol* 98:1832–1837
- Belmares R, Contreras-Esquivel JC, Rodríguez-Herrera R, Ramírez Coronel A, Aguilar CN (2004) Microbial production of tannase: an enzyme with potential use in food industry. *Lebensm Wiss Technol* 37:857–864
- Bhat TK, Singh B, Sharma OP (1998) Microbial degradation of tannins. A current perspective. *Biodegradation* 9:343–357
- Boadi DK, Neufeld RJ (2001) Encapsulation of tannase for the hydrolysis of tea tannins. *Enzyme Microb Technol* 28:590–595

- Bradoo S, Gupta R, Saxena RK (1997) Parametric optimization and biochemical regulation of extracellular tannase from *Aspergillus japonicus*. Proc Biochem 32:135–139
- Bradoo S, Gupta R, Saxena R (1996) Screening of extracellular tannase producing fungi: development of a rapid simple plate assay. J Gen Appl Microbiol 42:325–329
- Cerda-Gomez A, Contreras-Esquivel JC, Reyes-Valdes H, Rodríguez R, Aguilar CN (2006) Molecular characterization of *Aspergillus* strains producers of tannase. Proceedings of the second international congress on food science and technology in developing countries, Saltillo, Coahuila, Mexico (FEMB-18)
- Chae S, Yu T (1983) Experimental manufacture of a com wine by fungal tannase. Hanguk Sipkum Kwahakoechi 15:326–332
- Chang FS, Chen PC, Chen RC, Lu FM, Cheng TJ (2006) Real-time assay of immobilized tannase with a stopped-flow conductometric device. Bioelectrochemistry 69:113–116
- Chatterjee R, Dutta A, Banerjee R, Bhattacharyya BC (1996) Production of tannase by solid state fermentation. Bioprocess Eng 14:159–162
- Coggon P, Graham NH, Sanderson GW (1975) UK Patent 1,380,135
- Contreras-Dominguez M, Guyot S, Marnet N, Le Petit J, Perraud-Gaime I, Roussos S, Augur C (2006) Degradation of procyanidins by *Aspergillus fumigatus*: identification of a novel aromatic ring cleavage product. Biochimie 88:1899–1908
- Cruz-Hernández M, Augur C, Rodríguez R, Contreras-Esquivel J, Aguilar CN (2006) Evaluation of culture conditions for tannase production by *Aspergillus niger* GH1. Food Technol Biotechnol 44:541–544
- Deschamps AM, Otuk G, Lebeault JM (1983) Production of tannase and degradation of chestnut tannin by bacteria. J Ferment Technol 61:55–59
- Dueñas M, Hernández T, Estrella I (2007) Changes in the content of bioactive polyphenolic compounds of lentils by the action of exogenous enzymes. Effect on their antioxidant activity. Food Chem 101:90–97
- Farias GM, Gorbea C, Elkins JR, Griffin GJ (1994) Purification, characterization, and substrate relationships of the tannase from *Cryphonectria parasitica*. Physiol Mol Plant Pathol 44:51–63
- Gaathon A, Gross Z, Rozhanski M (1989) Propyl gallate: enzymatic synthesis in a reverse micelle system. Enzyme Microb Technol 11:604–609
- Gammoun A, Moros J, Tahiri S, Garriques S, Guardia M (2006) Partial least-squares near-infrared determination of hydrocarbons removed from polluted waters by tanned solid wastes. Anal Bioanal Chem 385:766–770
- Ganga PS, Nandy SC, Santappa M (1977) Effect of environmental factors on the production of fungal tannase. Leather Sci 24:8–16
- García-Conesa MT, Ostergaard P, Kauppinen S, Williamson G (2001) Hydrolysis of diethyl diferulates by a tannase from *Aspergillus oryzae*. Carbohydr Polym 44:319–324
- García-Nájera JA, Contreras-Esquivel JC, Rodríguez-Herrera R, Prado-Barragan LA, Aguilar CN (2002) Fungal production of 3,4,5-trihydroxybenzoic acid in submerged culture (in Spanish). Proceedings of the third international symposium on bioprocess engineering, IBT-UNAM, Cuernavaca, Morelos, Mexico
- García-Peña I (1996) Production and characterization of a tannase produced by *Aspergillus niger* in solid state culture (in Spanish). M.Sc. thesis, Universidad Autónoma Metropolitana, Iztapalapa, Mexico
- García-Peña I, Augur C, Favela-Torres E (1999) Partial purification of *Aspergillus niger* tannase produced by solid state culture (in Spanish). In: Prado A, Huerta S, Rodríguez G, Sancedo G (eds) Advances in enzyme purification and applied biotechnology. Ediciones UAM-Iztapalapa, Mexico, DF, pp 247–261, 2207–2212
- Goel G, Puniya AK, Aguilar CN, Singh K. (2005) Interaction of gut microflora with tannins in feeds. Naturwissenschaften 92:497–503
- Goel G, Puniya AK, Singh K (2007) Phenotypic characterization of tannin–protein complex degrading bacteria from faeces of goat. Small Rumin Res 69:217–220
- Hadi TA, Banerjee R, Bhattacharyya BC (1994) Optimization of tannase biosynthesis by a newly isolated *Rhizopus oryzae*. Bioprocess Eng 11:239–243
- Hatamoto O, Watarai T, Kikuchi M, Mizusawa K, Sekine H (1996) Cloning and sequencing of the gene encoding tannase and a structural study of the tannase subunit from *Aspergillus oryzae*. Gene 175:215–221
- Hatamoto O, Watarai T, Kikuchi M, Mizusawa K, Sekine H (1997) Erratum to: 'cloning and sequencing of the gene encoding tannase and a structural study of the tannase subunit from *Aspergillus oryzae*' [Gene 175 (1996) 215–221]. Gene 186:1–153
- Huang W, Ni J, Borthwick AGL (2005) Biosynthesis of valonia tannin hydrolase and hydrolysis of valonia tannin to ellagic acid by *Aspergillus* SHL 6. Process Biochem 40:1245–1249
- Ibuchi S, Minoda Y, Yamada K (1972) Hydrolyzing pathway, substrate specificity and inhibition of tannin acyl hydrolase. Agric Biol Chem 32:803–809
- Kar B, Banerjee R (2000) Biosynthesis of tannin acyl hydrolase from tannin rich forest residue under different fermentation conditions. J Ind Microbiol Biotechnol 25:29–38
- Kar B, Banerjee R, Bhattacharyya BC (2002) Optimization of physicochemical parameters for gallic acid production by evolutionary operation-factorial design technique. Process Biochem 37:1395–1401
- Kar B, Banerjee R, Bhattacharyya BC (2003) Effect of additives on the behavioural properties of tannin acyl hydrolase. Process Biochem 38:1285–1293
- Kostinek M, Specht I, Edward VA, Pinto C, Egonlety M, Sossa C, Mbugua S, Dortu C, Thonart P, Taljaard L (2007) Characterisation and biochemical properties of predominant lactic acid bacteria from fermenting cassava for selection as starter cultures. Int J Food Microbiol 114:342–351
- Kumar R, Sharma J, Singh R (2006) Production of tannase from *Aspergillus ruber* under solid state fermentation using jamun (*Syzygium cumini*) leaves. Microbiol Res (in press). DOI 10.1016/J.micres.2006.06.012
- Lee J, Talcott ST (2005) Enzyme hydrolysis of ellagic acid derivatives in muscadine grapes (*Vitis rotundifolia*). Session 36E, fruit and vegetable products: general. IFT annual meeting, July 15–20, New Orleans, La, USA
- Lekha PK, Lonsane BK (1997) Production and application of tannin acyl hydrolase: state of the art. Adv Appl Microbiol 44:215–260
- Lekha P, Lonsane B (1994) Comparative titres, location and properties of tannin acyl hydrolase produced by *Aspergillus niger* PKL 104 in solid-state, liquid surface and submerged fermentations. Process Biochem 29:497–503
- Lekha P, Ramakrishna M, Lonsane B (1993) Strategies for isolation of potent culture capable of producing tannin acyl hydrolase in higher titres. Chem Mikrobiol Technol Lebensm 15:5–10
- Lewis JA, Starkey RL (1969) Decomposition of plant tannins by some soil microorganisms. Soil Sci 107:235–241
- López-Ríos GF (1984) Fitoquímica, 1st edn. Universidad Autónoma de Chapingo, Estado de México, p 13
- Mahapatra K, Nanda RK, Bag SS, Banerjee R, Pandey A, Szakacs G (2005) Purification, characterization and some studies on secondary structure of tannase from *Aspergillus awamori nakazawa*. Process Biochem 40:3251–3254
- Mahendran B, Raman N, Kim D (2005) Purification and characterization of tannase from *Paecilomyces variotii*: hydrolysis of tannic acid using immobilized tannase. Appl Microbiol Biotechnol 70:445–451
- Mendez A (1984) Energía metabolizable del sorgo y efecto de la adición de aceite a dietas con sorgo dulce o amargo. Avances de investigación (resúmenes), Colegio de Postgraduados
- Mingshu L, Kai Y, Qiang H, Dongying J (2006) Biodegradation of gallotannins and ellagitannins. J Basic Microbiol 46:68–84

- Mondal K, Banerjee D, Banerjee R, Pati B (2001b) Production and characterization of tannase from *Bacillus cereus* KBR9. *J Gen Appl Microbiol* 47:263–267
- Mondal K, Samanta S, Giri S, Pati B (2001a) Distribution of tannic acid degrading microorganisms in the soil and comparative study of tannase from two fungal strains. *Acta Microbiol Pol* 50:75–82
- Mukherjee G, Banerjee R (2005) Effects of temperature, pH and additives on the activity of tannase produced by a co-culture of *Rhizopus oryzae* and *Aspergillus foetidus*. *World J Microbiol Biotechnol* 22:207–211
- Murugan K, Saravanababu S, Arunachalam M (2007) Screening of tannin acyl hydrolase (E.C.3.1.1.20) producing tannery effluent fungal isolates using simple agar plate and SmF process. *Bioresour Technol* 98:946–949
- Nip WK, Burns EE (1969) Pigment characterization in grain sorghum, I. Red varieties. *Cereal Chem* 46:490–495
- Nishitani Y, Osawa R (2003) A novel colorimetric method to quantify tannase activity of viable bacteria. *J Microbiol Methods* 54:281–284
- Nishitani Y, Sasaki E, Fujisawa T, Osawa R (2004) Genotypic Analyses of *Lactobacilli* with a range of tannase activities isolated from human feces and fermented foods. *Syst Appl Microbiol* 27:109–117
- Nuero OM, Reyes F (2002) Enzymes for animal feeding from *Penicillium chrysogenum* mycelial wastes from penicillin manufacture. *Lett App Microbiol* 34:413–416
- Pinto G, Bruno L, Hamacher M, Tarzi S, Couri S (2003) Increase of tannase production in solid state fermentation by *Aspergillus niger* 3T5B8. 25th Symposium on biotechnology for fuels and chemicals, poster presentation, Breckenridge, CO, USA, pp 3–68
- Pinto G, Couri S, Goncalves E (2006) Replacement of methanol by ethanol on gallic acid determination by rhodanine and its impacts on the tannase assay. *EJEAFChE* 5:5. [http://ejeafche.uvigo.es/5\(5\)2006/009552006F.htm](http://ejeafche.uvigo.es/5(5)2006/009552006F.htm)
- Pinto G, Leite S, Tarzi S, Couri S (2001) Selection of tannase-producing *Aspergillus niger* strains. *Braz J Microbiol* 32:24–26
- Pourrat H, Regeat F, Pourrat A, Jean D (1985) Production of gallic acid from tara tannin by a strain of *A. niger*. *J Ferment Technol* 63:401–403
- Purohit JS, Dutta JR, Nanda RK, Banerjee R (2006) Strain improvement for tannase production from coculture of *Aspergillus foetidus* and *Rhizopus oryzae*. *Bioresour Technol* 97:795–801
- Raab T, Bel-Rhliid R, Williamson G, Hansen CE, Chaillot D (2007) Enzymatic galloylation of catechins in room temperature ionic liquids. *J Mol Catal B Enzym* 44:60–65
- Rajkumar GS, Nandy SC (1983) Isolation purification and some properties of *Penicillium chrysogenum* tannase. *Appl Environ Microbiol* 46:525–527
- Ramirez-Coronel A, Marnet N, Kumar V, Rousses S, Guyot S, Augur C (2004) Characterization and estimation of proanthocyanidins and other phenolics in coffee pulp (*Coffea arabica*) by thiolysis-high-performance liquid chromatography. *J Agric Food Chem* 52:1344–1349
- Ramirez-Coronel A, Viniestra-González G, Augur C (1999) Purification of a tannase produced by *Aspergillus niger* Aa-20, in solid state fermentation. Proceedings of the VIII Mexican congress and IV Latin American congress of biotechnology and bioengineering, Huatulco, Oaxaca, Mexico
- Ramirez-Coronel MA, Viniestra-Gonzalez G, Darvill A, Augur C (2003) A novel tannase from *Aspergillus niger* with  $\beta$ -glucosidase activity. *Microbiology* 149:2941–2946
- Rana N, Bhat T (2005) Effect of fermentation system on the production and properties of tannase of *Aspergillus niger* van Tieghem MTCC 2425. *J Gen Appl Microbiol* 51:203–212
- Rout S, Banerjee R (2006) Production of tannase under mSSF and its application in fruit juice debittering. *Ind J Biotechnol* 5:351–356
- Saavedra G, Couri S, Ferreira S, Sousa de Brito E (2005) Tannase: conceitos, productos e aplicacao (in Portuguese). *B.CEPPA Curitiba* 23:435–462
- Sabu A, Pandey A, Jaafar Daud M, Szakacs G (2005) Tamarind seed powder and palm kernel cake: two novel agro residues for the production of tannase under solid state fermentation by *Aspergillus niger* ATCC 16620. *Bioresour Technol* 96:1223–1228
- Sabu A, Augur C, Swati C, Pandey A (2006) Tannase production by *Lactobacillus* sp. ASR-S1 under solid-state fermentation. *Process Biochem* 41:575–580
- Sanchez EE (2001) Applications and potential uses of tannase and tannins (in Spanish). B.Sc. thesis, Universidad Autónoma de Coahuila, Saltillo, Coah, p 25
- Sasaki E, Shimada T, Osawa R, Nishitani Y, Spring S, Lang E (2005) Isolation of tannin-degrading bacteria isolated from feces of the Japanese large wood mouse, *Apodemus speciosus*, feeding on tannin-rich acorns. *Syst Appl Microbiol* 28:358–365
- Scalbert A (1991) Antimicrobial properties of tannins. *Phytochemistry* 30:3875–3883
- Sharma S, Gupta MN (2003) Synthesis of antioxidant propyl gallate using tannase from *Aspergillus niger* van Tieghem in nonaqueous media. *Bioorg Med Chem Lett* 13:395–397
- Sharma S, Bhat TK, Gupta MN (2002) Bioaffinity immobilization of tannase from *Aspergillus niger* on concavalin A-sepharose CL-4B. *Biotechnol Appl Biochem* 35:165–169
- Shi B, He Q, Yao K, Huang W, Li Q (2005) Production of ellagic acid from degradation of valonea tannins by *Aspergillus niger* and *Candida utilis*. *J Chem Technol Biotechnol* 80:1154–1159
- Sittig M (1988) Trimethoprim. In: Sittig M (ed) *Pharmaceutical manufacturing encyclopedia*. William Andrew/Noyes, New Jersey, pp 282–284
- Tieghem, P (1867) Sur la fermentation gallyque. *CR Acad Sci (Paris)* 65:1091–1094
- Treviño-Cueto B, Luis M, Contreras-Esquivel JC, Rodríguez R, Aguilar A, Aguilar CN (2007) Gallic acid and tannase accumulation during fungal solid state culture of a tannin-rich desert plant (*Larrea tridentate* Cov). *Bioresour Technol* 98:721–724
- Van de Lagemaat J, Pyle DL (2001) Solid-state fermentation and bioremediation: development of a continuous process for the production of fungal tannases. *Chem Eng J* 84:115–123
- Van de Lagemaat J, Pyle DL (2005) Modelling the uptake and growth kinetics of *Penicillium glabrum* in a tannic acid-containing solid-state fermentation for tannase production. *Process Biochem* 40:1773–1782
- Van Diepeningen AD, Debet A, Varga J, Van der Gaag M, Swart K, Hoekstra R (2004) Efficient degradation of tannic acid by black *Aspergillus* species. *Mycol Res* 108:919–925
- Vaquero I, Marcobal A, Muñoz R (2004) Tannase activity by lactic acid bacteria isolated from grape must and wine. *Int J Food Microbiol* 96:199–204
- Vattem DA, Shetty K (2002) Solid-state production of phenolic antioxidants from cranberry pomace by *Rhizopus oligosporum*. *Food Biotechnol* 16:189–210
- Vattem DA, Shetty K (2003) Ellagic acid production and phenolic antioxidants activity in cranberry pomace (*Vaccinium macrocarpo*) mediated by *Lentinus edodes* using a solid-state system. *Process Biochem* 39:367–379
- Viniestra-González G, Favela-Torres E, Aguilar CN, Romero-Gomez SJ, Díaz-Godínez G, Augur C (2003) Advantages of fungal enzyme production in solid state over liquid fermentation systems. *Biochem Eng J* 13:157–167

- Vivas N, Laguierre M, Pianet de Boissel I, Vivas de Gaulejac N, Nonier MF (2004) Conformational interpretation of vascalagin and castalagin physicochemical properties. *J Agric Food Chem* 52:2073–2078
- Yamada K, Iibuchi S, Minoda Y (1968) Studies on tannin acyl hydrolase of microorganisms. Isolation and identification of producing molds and studies on the conditions of cultivation. *Agric Biol Chem* 45:233–240
- Yoshida T, Amakura Y, Koyura N, Ito H, Isaza JH, Ramírez S, Peláez DP, Renner SS (1999) Oligomeric hydrolysable tannins from *Tibouchina multiflora*. *Phytochemistry* 52:1661–1666
- Yu X, Li Y, Wu D (2004) Enzymatic synthesis of gallic acid esters using microencapsulated tannase: effect of organic solvents and enzyme specificity. *J Mol Catal B Enzym* 30(2):69–73
- Yu XW, Li YQ (2006) Kinetics and thermodynamics of synthesis of propyl gallate by mycelium-bound tannase from *Aspergillus niger* in organic solvent. *J Mol Catal B Enzym* 40:44–50
- Zhang YJ, Abe T, Tanaka T, Yang CR, Kouna I (2001) Phyllanem-blinins A–F, new ellagitannins from *Phyllanthus emblica*. *J Nat Prod* 64:1527–1532
- Zhong X, Peng L, Zheng S, Sun Z, Ren Y, Dong M, Xu A (2004) Secretion, purification, and characterization of a recombinant *Aspergillus oryzae* tannase in *Pichia pastoris*. *Protein Expr Purif* 36:165–169