

Cite this: *Food Funct.*, 2015, **6**, 1035

Microencapsulation of bioactives for food applications†

 Maria Inês Dias,^{a,b} Isabel C. F. R. Ferreira^{*a} and Maria Filomena Barreiro^{*b}

Health issues are an emerging concern to the world population, and therefore the food industry is searching for novel food products containing health-promoting bioactive compounds, with little or no synthetic ingredients. However, there are some challenges in the development of functional foods, particularly in which the direct use of some bioactives is involved. They can show problems of instability, react with other food matrix ingredients or present strong odour and/or flavours. In this context, microencapsulation emerges as a potential approach to overcome these problems and, additionally, to provide controlled or targeted delivery or release. This work intends to contribute to the field of functional food development by performing a comprehensive review on the microencapsulation methods and materials, the bioactives used (extracts and isolated compounds) and the final application development. Although several studies dealing with microencapsulation of bioactives exist, they are mainly focused on the process development and the majority lack proof of concept for final applications. These factors, together with the lack of regulation, in Europe and in the United States, delay the development of new functional foods and, consequently, their market entry. In conclusion, the potential of microencapsulation to protect bioactive compounds ensuring their bioavailability is shown, but further studies are required, considering both its applicability and incentives by regulatory agencies.

Received 19th December 2014,

Accepted 7th February 2015

DOI: 10.1039/c4fo01175a

www.rsc.org/foodfunction

1. Introduction

1.1. The increasing interest in functional foods

Nowadays, food not only serves to satisfy the primal urge of hunger, but also is a means to promote consumer's health. In this context, the food industry has focused on avoiding the potential harmfulness of synthetic food additives and on developing novel food products containing health-promoting ingredients. Therefore, bioactive natural products are considered as viable and safer substitutes to satisfy the world market demand for new products.¹

"Functional foods" arise as the frontier between nutrition and health, providing a long-term beneficial physiological/health effect beyond their nutritional properties.¹ The concept of functional food appeared 40 years ago, however the growing interest in this type of product, either from industry (through patents) or academia (through scientific research articles and

reviews), was only observed from the second half of the 1990s, indicating an increasing tendency (Fig. 1). The exponential growth of patents and scientific research articles/reviews observed since 2005 was accompanied by the regulation (EC) no. 1924/2006 publication by the European Parliament on nutrition and health claims in foods, which was completed and finalized in 2011 by the European Food Safety Authority (EFSA) regarding beneficial health claims in certain food ingredients.^{2,3} In the United States (US) the regulation of functional foods is facilitated, as the food industry itself provides the product definition that will be placed on the market; food companies are only obliged to follow labelling and safety rules implemented by the Food and Drug Administration (FDA).⁴

Nowadays, consumers' awareness of health issues is growing together with the increasing incidence of chronic age-related diseases, such as neurodegenerative diseases, diabetes and cancer, usually correlated with the lifestyle and dietary habits of our societies.⁵ Moreover, as the life expectancy is rising, with the consequent increase of health care costs, pharmaceutical and food industries have started to consider functional foods as a new market with huge growth potential. Nowadays, Japan, the United States (US) and the European Union (EU) are the leading markets for functional foods, representing in total 90% of the world market supply for this type of product.⁶ In 2006, US and EU markets were valued at 33 billion US\$ and at 15 billion US\$, respectively, with a

^aMountain Research Centre (CIMO), ESA, Polytechnic Institute of Bragança, Campus Santa Apolónia Ap. 1172, 5301-855 Bragança, Portugal. E-mail: iferreira@ipb.pt; Fax: +351-273-325405; Tel: +351-273-303219

^bLaboratory of Separation and Reaction Engineering (LSRE), Associate Laboratory LSRE/LCM, Polytechnic Institute of Bragança, Campus Santa Apolónia Ap. 1134, 5301-857 Bragança, Portugal. E-mail: barreiro@ipb.pt; Fax: +351-273-325405; Tel: +351-273-303089

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c4fo01175a




Fig. 1 Number of research articles and reviews, and patents published in the period from 1970 to 2014 regarding functional foods (obtained on web of science, October 2014; keyword: functional food).

tendency to grow. Germany, France, the United Kingdom and the Netherlands are considered the most important countries within the European functional food market.⁷

1.2. The problems related to the use of free bioactives

Despite the known beneficial health effects of natural bioactive matrices and isolated individual compounds, as will be discussed in this section, they show some fragility that has to be considered regarding their direct use or incorporation into foods.

The main factors limiting the use of bioactives in food applications are shown in Fig. 2. Bioactive ingredients are generally prone to degradation, both during storage and food processing, as many of them are physically, chemically and/or enzymatically unstable leading to their degradation or transformation with the consequent loss of bioactivity. In many cases the mechanism involved in the degradation of these bioactive molecules is very complex and still unknown.^{5,8} Wu *et al.*⁹ reported the reduction of the anthocyanin content in blackberry fruits after six months of canned and jam storage and also after drying treatment. Various types of cereals (wheat, barley and oat) were also tested for the content of biologically active compounds, such as tocopherols, phenolic compounds and microelements, and after hydrothermal processing, the concentration of these molecules severely decreased.¹⁰ Rawson *et al.*¹¹ described major losses of bioactive compounds after processing exotic fruits such as mangoes, *açaí*, pineapple and *pitanga*, subjecting them to heat treatments, pasteurization and drying, canning and even to storage processing steps. All these processes affect, to a lesser or greater extent, the stability, chemical characteristics, concentration, and even antioxidant activity of a number of compounds such as vitamins and phenolic compounds. Another study that describes the modifications occurring in fruits and vegetables during the processing steps was published by Nicoli



Maria Inês Dias

Maria Inês Dias has been a PhD student at Mountain Research Centre (CIMO) of the Polytechnic Institute of Bragança, REQUIMTE and Laboratory of Separation Reaction Engineering (LSRE) of the University of Porto since 2013. She obtained her degree in Biotechnological Engineering in 2009 and her Master's degree in Biotechnology in 2011, both at Polytechnic Institute of Bragança. Her main research interests are improvement of secondary metabolite production through in vitro culture techniques, chemical characterization and bioactive properties of edible plants, and microencapsulation of phenolic fractions.



Isabel C.F.R. Ferreira

Isabel C.F.R. Ferreira is Coordinator Professor at Polytechnic Institute of Bragança (Portugal). She is a member of the Board of Directors of the Mountain Research Centre (CIMO) and the principal investigator of the Food Safety and Technology Group. She is an associate editor of Food & Function (a journal of the Royal Society of Chemistry). She obtained her degree in Biochemistry (1996) at the University of Porto, Master's in Sciences (1999), PhD in Sciences—Chemistry (2003), and habilitation in Sciences—Chemistry (2011) at the University of Minho (Portugal). She is the principal investigator of several financed research projects, and an evaluator of international and national research projects, post-doc and PhD grants. She had supervised several post-doc, PhD and master's students in the BioChemCore group. She has received awards from several different organizations such as Calouste Gulbenkian Foundation. She has published over 260 papers in refereed journals and is a highly cited scientist (top 1%) in Agricultural Sciences (Researcher ID: E-8500-2013; ORCID ID: 0000-0003-4910-4882; SCOPUS ID: 7102135224). Her main research interests are nutraceuticals and functional foods, chemistry of natural matrices/products, and emerging technologies for conservation of food matrices.




Fig. 2 Limiting factors for the use of free bioactives in food applications.

et al.,¹² focusing on the antioxidant activity decrease of the food matrix due to the loss and transformation of the antioxidant compounds, and also due to their interaction with other molecules. The processing steps of a food matrix involve the action of endogenous enzymes, water activity, oxygen pressure and also thermal/mechanical energy, and all of these factors can influence the degradation/transformation of the bioactive molecules leading to the loss of their intended characteristics.



Maria Filomena Barreiro

Filomena Barreiro graduated in Chemical Engineering in 1990 at the Faculty of Engineering, University of Porto (FEUP). In 2000, she obtained her Ph.D. also at FEUP. She joined the Polytechnic Institute of Bragança (IPB, <http://www.ipb.pt>) in 1995 where she is Coordinator Professor at the Chemical and Biological Department (DTQB). Since 2000 she has been a member of the research staff of the Laboratory of Separation and Reaction

Engineering (LSRE, <http://lsre.fe.up.pt/>) (Researcher ID: L-9802-2014; ORCID ID: 0000-0002-6844-333X; SCOPUS ID: 23093315700). Her main research interests are synthesis and characterization of polymeric materials with special emphasis on materials from renewable resources and polyurethane chemistry, functional coatings and microencapsulation applied to different industrial fields.

Nevertheless, not all the compounds are equally affected; phenolic compounds and vitamins (*e.g.* vitamins C and E) are more sensitive to blanching and long-term freezing treatments than minerals or dietary fibres.¹³ Despite the processing steps, the perishability of food is also a limitation in their intake in the free form. This is because the shelf life determines whether a particular food maintains its characteristics and bioactive properties. For instance, edible mushrooms have a very short shelf life and the postharvest changes, such as browning, cap transformation, texture and weight loss changes, occur immediately, which decrease their bioactive components.¹⁴

The ingested amount of the bioactive compound, its structure and chemical form, its interaction with other molecules, and also the organism itself (mucosal mass, intestinal and gastric behaviour, metabolism and protein bonding) will influence the stability and functionality within the human body, and consequently its bioavailability.^{15,16} For instance, phenolic compounds present very low bioavailability due to their poor solubility and stability, especially those with high molecular weight. Furthermore, there are no reports on specific receptors on the small intestinal epithelial cell surface, and thus the transport mechanism involves active diffusion and active efflux, lowering the permeability of such compounds.¹⁷ In the case of anthocyanins, they are very sensitive to pH and temperature changes in the medium.¹⁸ Concerning carotenoid compounds, the nature of the food matrix, the particle size and the processing method, and also their interaction with other food constituents, will affect their bioavailability; moreover, fibre constituents decrease the absorption of carotenoids. The nutritional state of the organism itself will influence the absorption of these molecules (*e.g.*, protein deficiency affects the bioavailability).^{19,20} As an example, the interaction of mineral elements with other molecules can decrease their bioavailability, as is the case of calcium where compounds such as oxalates, tannins and dietary fibres decrease its absorption due to precipitation.²¹ Also, the gastrointestinal environment and epithelial transport can decrease the bioavailability of natural extracts, as described by Vermaak *et al.*²² who investigated the biological activity of green tea and sage extracts under simulated gastrointestinal conditions; the authors observed an accentuated decrease in the antimicrobial activity. Lipophilic compounds have also low solubility, which restricts their incorporation into many food matrices, especially in water-based carriers. The molecular weight, functionality and polarity seriously influence their solubility, physical state, chemical stability and bioavailability.^{8,23} It is very difficult to evaluate the bioavailability of these types of compounds, since once metabolized they reach the systemic circulatory system where they can be stored, utilized or excreted. Depending on the concentration and time of these molecules in a particular tissue, or their use in some biological function, the bioavailability can be estimated.²⁴ For instance, the bioavailability of lycopene, a highly lipophilic carotenoid compound, is influenced dramatically by the intestinal lymphatic uptake. Faisal *et al.*²⁵ applied an *in vivo* model to increase its solubility using digestible lipid excipients. A similar study was

performed by Balakrishnan *et al.*²⁶ in order to increase the solubility and bioavailability of the Coenzyme Q₁₀, practically insoluble in aqueous medium, by using oil and surfactant compounds for its oral delivery.

Another factor that drives researchers to invest their knowledge into the design of novel food delivery systems is the organoleptic behaviour of some bioactive extracts/compounds. They can present unpleasant taste, odour and/or textures. This is a crucial point for the food industry when developing a new product because the consumer gives importance not only to the price, but also to the taste, smell and appearance. Accordingly, consumers will choose the non-functional counterpart of a similar product, even if it has lower bioactive properties.^{16,27} It is known that many people avoid eating fruits and vegetables because most of their compounds such as polyphenols, terpenes and glucosinolates have bitter or astringent taste, making them unappealing to the consumer.²⁸

To overcome the problems related to the direct use of bioactive extracts/compounds, microencapsulation techniques arise as a potential approach in the food industry to deal with their incorporation, either to impart additional functional properties or to protect the bioactive component itself.

The main goal of the present review is to highlight the use of microencapsulation techniques for food applications, as well as to discuss the advantages of microencapsulating bioactive extracts/compounds. Various extracts and compounds that have been encapsulated using different techniques and formulations will be enumerated focusing on the potential for functional food development. A particular emphasis will be given to examples where the final application (incorporation into food matrices) is explored.

2. Overview of microencapsulation techniques and materials

2.1. The advantages of using microencapsulated bioactives

Microencapsulation can provide a tool to protect natural extracts and compounds from the action of biotic, abiotic, and biological factors. It emerges as a reliable methodology not only for the food industry, but also for the fields of nutrition and health, where the stability, efficacy and bioavailability of these extracts and compounds are needed. As described previously, there are several factors affecting the bioactives' stability in their free form (Fig. 2), however with microencapsulation technology protection from factors such as light, moisture, heat and oxygen is provided. Also, the organoleptic characteristics of many food products can be masked, but most importantly functional/biological characteristics can be maintained after ingestion together with controlled release in a specific target. The success of a delivery system based on microencapsulation can be measured by the bioactives' behaviour during food processing and storage, and after ingestion.⁸

From a practical point of view, microencapsulation techniques protect the core material from the outside environment; it increases the product shelf life by reducing the

transfer between the core and the surrounding medium, and by protecting the molecules from reaction with other food constituents, which can decrease their bioavailability.²⁹ It also increases the solubility, dispersibility and flow of the bioactives.³⁰

Depending on the applied technology and encapsulated bioactive, the response of the produced delivery system will be different; each compound has specific characteristics that should be considered in the design of a novel microencapsulation process. For instance, phenolic compounds are very powerful antioxidant molecules; however they present problems in their bioavailability because they are transformed, after ingestion, into methylated, glucuronated and sulphated metabolites.³¹ Nano- and microparticle based delivery systems appear as a response to overcome these problems, increasing the phytochemical absorption of phenolic compounds in epithelial cells.^{17,32} In particular, Davidov-Pardo & McClements³³ showed that the microencapsulation of resveratrol increased its bioavailability.

Essential oils have also some organoleptic related problems, most of them presenting an unpleasant taste and odour, with very poor water solubility and high volatility. All these limitations can be overcome by using microencapsulation techniques that increase the effectiveness of their biological functions and decrease the sensory impact on food products.³⁴

2.2. Microencapsulation techniques

The microencapsulation concept was primarily developed by the pharmaceutical industrial sector, whose goal was to control and/or modify the release of drug substances. Nowadays, it still represents the major field using microencapsulation (68%) while the food sector accounts for only 13%.³⁵ The amount of scientific reports and patents regarding microencapsulation for food purposes (Fig. 3) is indicative of the growing interest in this technique regarding the incorporation of bioactive extracts and compounds. Nevertheless, the absence of regulation for novel food ingredients, including the ones derived from using nano- and micro-technologies in their preparation, is still remaining. In the USA the FDA is currently developing a recognition program for nanomaterials to overcome the existing scarcity of information, and also to assess food safety of these new ingredients.³⁶ The introduction of microencapsulation technologies into the food industry allows the incorporation of flavouring agents in certain types of foods, and also the improvement of their functional and health properties.^{30,37} Regarding food science and biotechnology, the incorporation of natural ingredients intends to stabilize, protect and preserve the bioactives into a core, surrounded by a wall, or dispersed in a matrix, made of a material chosen to be suitable for the target delivery system.³⁴ There have already been reviews on microencapsulation of bioactive compounds and extracts for food applications,^{29,30,34,37–40} nevertheless they mainly explored the available techniques for microencapsulation, lacking specificity in the existing examples of microencapsulated bioactive extracts and com-




Fig. 3 Number of research articles and reviews, and patents published in the period from 1970 to 2014 regarding microencapsulation for food purposes (obtained on web of science, October 2014; keywords: microencapsulation and food).

pounds together with the applicability of the performed studies. Fig. 4 shows the logical chain, from the choice of bioactives, materials and microencapsulation process to final applications evidencing the crucial points involved in each step.

Microcapsules are particles with diameters ranging from 1 to 1000 micrometers. The most common morphology is of two types: (1) shell type, where the core, the bioactive component itself or a carrier containing it (compounds that facilitate the release) is protected by a membrane; (2) matrix type, where the bioactive component is dispersed in a material's matrix. The encapsulation materials, production process, final morphology and ultimate application are the most important factors to be taken into account when designing a novel delivery system based product. Also, the stability and functional properties of the bioactive component must be taken into account when selecting the microencapsulation technique. Furthermore, to achieve high encapsulation yields it is necessary to ensure process reproducibility, release profile and overcome limiting drawbacks such as microspheres aggregation and adherence.³⁰

The encapsulation methods and materials most commonly used in food applications are described in Tables 1 and 2,

respectively (as also in the ESI†). The definition of categories presented in Table 1 was somehow difficult because the microencapsulation processes can be categorized according to the formation mechanism, the consolidation method, and even according to the specific equipment used. A clear distinction among the described possibilities is not always clear in the published work. Therefore, in this work, efforts were made to define categories according to the microcapsule formation process and a set of general categories are proposed: coacervation, extrusion-based processes, spray-based processes, emulsion-based processes, liposomes, supercritical fluid based processes, ultrasound-based processes and others.

2.2.1. Spray-based process. Spray-based processes are by far the most common methods being divided into spray-drying, electrospray, spray-coagulation (according to internal or external gelation) and spray-freeze drying methods. Spray-drying, the oldest microencapsulation process used by the food industry, is a very straightforward technique. It can be described as flexible, allowing continuous production, making it a cost effective process and consequently the most economical among several encapsulation methods. It can be easily industrialized in terms of equipment and materials, which have a low cost compared with other available techniques.⁴¹ The most commonly used shell materials in this technique are carbohydrates which may limit the encapsulation of some bioactives.³⁹ It produces high quality microcapsules, with a size less than 40 μm , by atomizing a liquid solution or emulsion through a nozzle to a hot gas chamber giving rise to the prompt formation of a powder. The method's speed and effectiveness ensure the production of microbiologically stable products, with lower costs and specific properties.^{37,41} There are several applications dealing with the encapsulation of bioactive compounds and extracts by spray-drying. Examples in the published literature are crude extracts,^{42–52} carotenoids,^{53,54} enzymes,^{55,56} essential oils,^{57–62} fatty acids,^{63–66} phenolic compounds (including anthocyanins)^{67–87} and vitamins.⁸⁸ It is also noticeable (ESI†) that the vast majority of the used shell materials, as was previously reported, are carbo-




Fig. 4 Schematic procedures for the development of microencapsulation protocols (GRAS—generally recognized as safe).

Table 1 The encapsulation methodologies mostly used for food applications, and the corresponding examples

Method category	Examples	Ref.
Coacervation	Complex coacervation Simple coacervation	90, 93, 102, 104–107, 125, 126, 159 108–111, 120, 128, 139
Extrusion-based processes	Electrostatic extrusion Co-extrusion	105, 122 123, 154
Spray-based processes	Spray drying Electrospray Spray-coagulation ^a Spray-freeze drying	42–88, 108, 150, 152, 157 103 89–94, 107 96–102
Emulsion based processes		106, 109–121, 160
Liposomes	Liposomes and niosomes	124–129, 156
Supercritical fluid based processes	Supercritical antisolvent process Rapid extraction of supercritical solution Supercritical fluid impregnation	87, 130 94 62
Ultrasound based processes	Sonication Ultrasound	131, 133 132
Others	Co-crystallization Core-shell printing Nanoprecipitation Fluidized bed Inclusion Lyophilization Microwave Molecular inclusion Phase separation method Response surface methodology Solvent evaporation Spinning disc reactor	136, 137 138 111, 139 134 153, 158 140, 141 142 155 143 144 145, 146 135

^a Coagulation mostly achieved with internal or external gelation.

Table 2 Main materials used for encapsulating bioactive extracts and compounds for food applications (based on Kuang *et al.*³⁰)

Material category	Encapsulation material	Ref.
Water soluble polymers	Carbohydrate and carbohydrate derivatives (<i>e.g.</i> alginate, gums, chitosan, amylose, k-carrageenan and pectin), protein and protein derivatives (<i>e.g.</i> whey, milk and soybean proteins), synthetic polymers (<i>e.g.</i> polyethyleneglycol) and others (<i>e.g.</i> ethyl cellulose and mucilage extract of <i>Opuntia ficus indica</i>)	43–60, 63–66, 70–79, 81–85, 87–94, 99, 102, 100–108, 110, 111, 113–115, 117–119, 123, 126, 131, 134, 140–144, 146, 153, 154, 157, 159
Water soluble non-polymers	Carbohydrate and carbohydrate derivatives (<i>e.g.</i> cyclodextrin, maltodextrin, inulin and lactose), synthetic polymers (<i>e.g.</i> PEG2000-DSPE, polyvinyl alcohol and high and low HLP lipophilic polymeric emulsifiers) and others (Tween, buffer and alcoholic solutions and ascorbic acid)	42, 44–46, 48, 49, 51, 53, 56, 57, 59, 63–65, 67–71, 76–80, 82, 83, 94, 96–99, 101, 104, 111, 112, 114, 120, 121, 128, 131–135, 139, 140, 143, 155, 156, 158
Non-water soluble polymers	Carbohydrate and carbohydrate derivatives (<i>e.g.</i> starch), protein and protein derivatives (<i>e.g.</i> casein), synthetic polymers (<i>e.g.</i> low-density polyethylene, poly(ϵ -caprolactone) and poly-D,L-lactide (PLA)) and others (liquid vaseline)	51, 56, 59, 62, 71, 73, 100, 101, 103, 109, 110, 112, 115, 122, 129, 139, 142, 145, 150, 152, 153, 160
Non-water soluble non-polymers	Carbohydrate and carbohydrate derivatives (<i>e.g.</i> sucrose) and others (lecithin, supercritical CO ₂ , stearic acid and wax)	61, 62, 66, 114, 116, 124–129, 136–138, 144

hydrates and derivatives. However, Medina-Torres *et al.*⁷² encapsulated gallic acid in mucilage obtained directly from *Opuntia ficus indica*, while Cortés-Rojas *et al.*⁶¹ encapsulated eugenol in lipid formulations, both affording good results and high encapsulation yields. These results show the constant evolution of this method, and the possibility to overcome constraints related to the limited number of available shell materials, as stated by Gouin *et al.*³⁹

Coagulation processes are also commonly used to encapsulate bioactive extracts and compounds for food applications, the most common being those based on alginate beads.^{89–94} Alginate beads are formed from the polyanionic copolymer derived from the brown marine algae, alginate, which is frequently used as a stabilizer and thickener of many food products. Its coagulation can be promoted by external gelation (*e.g.* using calcium chloride as the calcium source added to the

coagulation solution) or internal gelation (*e.g.* using calcium carbonate as the calcium source added to the alginate solution). In the first case, gelation occurs mainly at the particle surface and in the second case gelation occurs mainly inside the formed particles. The formed materials, due to their degree of ionic reticulation and functionality, allow the control of water intake and thus the release of the bioactive component.⁹⁵ The preparation of such alginate beads is easily performed at the lab-scale, and they have been used to encapsulate a wide variety of compounds (hydrophilic, lipophilic, oils among others), and the controlled release is achieved by pH changes.^{39,95}

Freeze-drying technology allows the encapsulation of many food constituents, being used on a daily basis to stabilize compounds and increase controlled release.³⁹ It is mostly used to encapsulate bioactive extracts,⁹⁶ phenolic compounds,^{97–99} vitamin C^{100,101} and even essential oils.¹⁰² To the best of our knowledge the use of electrospray technology for food applications is not very common and only one work was found in the reviewed literature.¹⁰³ This work refers to the encapsulation of folic acid (vitamin B₉), and according to the provided description, it is a very appealing technology since the use of organic solvents and high temperatures is not required.

2.2.2. Coacervation. Coacervation is the second most commonly used encapsulation technique for food applications, not only because it provides high encapsulation efficiency, but also due to the triggered controlled release that can be based on the temperature, mechanical or biological mechanisms, providing the needed versatility to support the development of a wide range of food products.³⁹ It can be divided into complex and simple coacervation; the first is based on the complexation of two oppositely charged polymers that form a strong polymeric shell or matrix.¹⁰⁴ For the complex coacervation, chitosan is the preferable wall material, and alginate is the most commonly used polymer in all the mentioned studies.^{92,93,105–107} Chitosan has low toxicity, antimicrobial activity, and biocompatibility, but it is mainly muco-adherence that allows transmucosal absorption and better release of the bioactives.¹⁰⁷ In simple coacervation the initially soluble polymer is precipitated by changing the pH or temperature.³⁴ Milk proteins^{108,109} and pectins with PGPR (polyglycerol polyricinoleate)¹¹⁰ are some examples of wall materials used in simple coacervation.

2.2.3. Emulsion based process. Emulsion based processes are also commonly used for food encapsulation applications. It allows the encapsulation of both water and oil soluble food ingredients.^{34,37} Emulsion based techniques have been successfully used to encapsulate bioactive compounds including fatty acids,^{111,112} vitamins,¹¹³ phenolic compounds,^{109,114–117} anthocyanins,^{110–118} oils^{119,120} and bioactive extracts.^{106,121} This technique is sometimes coupled with a second one, in most cases a spray-drying based process, which gives rise to a dry powder that can be promptly introduced into a food matrix.³⁷ In fact, several of the commonly used encapsulation processes start with the first step comprising the preparation of an emulsion. This is the reason why a straightforward divi-

sion of the encapsulation techniques is not easy to achieve and some superimposition exists. In this work, and given the importance of spray-based processes, the cases dealing with emulsion coupled with spray techniques were included in the spray-based process category.

2.2.4. Extrusion based process. Extrusion methodologies, unlike the above described methods, are not so usual. They can be divided into electrostatic extrusion and co-extrusion. The extrusion method comprises the passage of the polymer melt with the solubilized bioactive through a nozzle, or the polymer melt and bioactive through concentric nozzles, leading to the formation of particles with high density and encapsulation efficiency.^{30,37} This technique is primarily used for the encapsulation of volatiles and unstable flavours.³⁹ Belščak-Cvitanović *et al.*¹⁰⁵ and Barbosa-Pereira *et al.*¹²² demonstrated the efficiency of this method for the encapsulation of phenolic compounds. Co-extrusion is used to prepare spherical microbeads with a hydrophobic core,³⁷ nevertheless it can also be used for the encapsulation of hydrophilic compounds in alginate beads as was done by Piazza & Roversi.¹²³

2.2.5. Liposomes. Liposome technology has been mostly used in pharmaceutical and cosmetic fields, for targeted delivery of therapeutic agents and inclusion of stabilizers in creams and lotions, respectively. For food applicability they represent a highly valuable resource due to their high encapsulation efficiency, stability and easy production.³⁹ Foremost, liposomes have been used to stabilize and increase the bioavailability of bioactive molecules.^{124–127} Moreover it is widely used to encapsulate compounds that are poorly soluble in certain solvents. Coimbra *et al.*¹²⁸ demonstrated the efficacy of liposomes for the encapsulation of resveratrol, caffeic acid, carvacrol, among others (compounds poorly soluble in water), while Rasti *et al.*¹²⁹ increased the oxidative stability of polyunsaturated fatty acids by means of their encapsulation in liposomes.

2.2.6. Supercritical fluid based process. Supercritical fluid based processes have many advantages for the encapsulation of sensitive substances such as essential oils or enzymes, always being coupled with other encapsulation techniques. Almeida *et al.*⁶² used a supercritical fluid impregnation technique to encapsulate oregano essential oil into a starch matrix, achieving a homogeneous product in a faster way due to the low viscosity and higher diffusion of supercritical CO₂. On the other hand, Santos *et al.*⁹⁴ by using rapid extraction of a supercritical solution and Sosa *et al.*¹³⁰ and Visentin *et al.*⁸⁷ by using a supercritical antisolvent process applied this technique to encapsulate bioactive extracts with high encapsulation efficiencies. The main advantages of supercritical fluids are related to their physical properties such as viscosity, density, solvating power, diffusion and mass transfer. The solubilisation of the core and shell materials is therefore faster as microcapsule formation is facilitated, *i.e.* they are formed by using lower temperatures and in the absence of water.³⁹

2.2.7. Ultrasound based process. Ultrasound based processes, such as sonication and ultrasound, are also reliable techniques for food applications, mostly being used with the double functions of extracting the bioactives and forming the

microcapsules.^{131,132} However, Kalogeropoulos *et al.*¹³³ used sonication to aggregate the inclusion complex of propolis extract and β -cyclodextrins to form microcapsules.

2.2.8. Others. Despite all those described above, there are other methods not so common for food applications. An example is the fluidized bed, a microencapsulation technique for powder compounds. It needs the preparation of a suspension with the coating material (polysaccharides, proteins, emulsifiers and fats) and subsequent spraying, offering a more effective controlled release of the core material than with other existing technologies.^{30,37,39} Li *et al.*¹³⁴ used this technology achieving good integrity and stability of the core compound after the drying process. Molecular inclusion is another process that is not so commonly used, generally referred to as a supramolecular method in the sense that the bonding between the encapsulated compound and the shell material occurs in a cavity-bearing substrate by hydrogen bonds, van der Waals forces or entropy-driven hydrophobic effects. Cyclodextrins and hydrophobic vitamins are the most commonly used shell materials in molecular inclusion methods.³⁹ Spinning-disk and centrifugal co-extrusion are new atomisation methods, possibly used in modified spray encapsulation methods; the difference lies in the formation of the capsule, involving the creation of a film with much smaller dimensions than those obtained in common atomisers.³⁹ Akhtar *et al.*¹³⁵ showed the reduction of the particle size using a spinning-disk reactor to encapsulate flavonoids by means of a double emulsion technique, achieving better stabilization of the prepared emulsions by this technique. Other microencapsulation methods that are not commonly used in the food sector are co-crystallization,^{136,137} core-shell printing,¹³⁸ nanoprecipitation,^{111,139} lyophilisation,^{140,141} microwave,¹⁴² phase separation methods,¹⁴³ response surface methodology¹⁴⁴ and solvent evaporation methods.^{145,146}

2.3. Encapsulation materials

When designing an experiment protocol for the development of encapsulated products (Fig. 4), the shell material choice is one of the most important steps, firstly because it has to be non-toxic to the organism, its preparation should consider environmental issues and use clean solvents (water soluble materials are therefore preferable) and, finally, because it plays a crucial role in the bioactive release behaviour. Conditions such as pH, temperature, salts and ion concentration also have to be taken into account and defined in accordance with the ultimate objective of the developed microcapsules. In this work the materials were divided into four categories (Table 2), according to Kuang *et al.*³⁰ which discriminate them as water and non-water soluble materials, and as polymer and non-polymer materials. Within each category it was also possible to sub-divide them into carbohydrate and its derivatives, protein and its derivatives, synthetic polymers and other types of materials.

The coating material and its physical structure strongly influence the product development; nevertheless there are some constraints since law does not allow the application of

some materials in food. They must be considered “generally recognized as safe” (GRAS), biodegradable and efficient as the protective barrier between the nucleus and the surrounding medium. Both EU through the EFSA and the US through FDA have many strict rules about material usage for food applications.^{37,147} The most commonly used materials are carbohydrate polymers (starch and cellulose and their derivatives), plant exudates and extracts (gum, galactomannans, pectins and soybean polysaccharide), marine extracts (carrageenan and alginate), microbial and animal derived polysaccharides (xanthan, gellan, dextran and chitosan), and also proteins, lipids and others (paraffin and some inorganic materials).¹⁴⁸ This is in accordance with our survey, where it can be observed that water soluble materials, both polymer (*e.g.* alginate and chitosan) and non-polymer (*e.g.* cyclodextrins) types, are the most commonly used, followed by non-water soluble polymers (*e.g.* starch and casein) and, finally, non-water soluble non-polymers (*e.g.* sucrose and lecithin).

Concerning the EU, no access is provided to a list of authorized materials for food product development by EFSA. There is a lack of information, as the existing list is under construction. They include only food additives and nutrient sources, listing only those who are not considered food additives (*e.g.* starch), but without any reference to whether they are authorized or not.¹⁴⁹ Regarding the US, the FDA has a list of approved food ingredients that allows the companies and academia to design microencapsulation protocols more suitable to serve food industry purposes. Despite the above listed compounds, identified as the most commonly used, not all have been approved by the FDA (or they were not considered for review or the assessment is pending). From Table 2, and following the guidelines of FDA, it can be observed that the approved materials are stearic acid, sucrose, amylopectin, maize starch, calcium caseinate, casein, FHCO (fully hydrogenized canola oil), PGPR, β -cyclodextrin, ethanol, lactose, PEG (polyethylene glycol), alginate, chitosan, whey protein, cellulose, xanthan, ethyl cellulose, soy protein, inulin, pectin and lysozyme. The materials with pending requests for assessment are lecithin, caffeine, arabic gum, milk proteins and poloxamers. For the remaining materials no information is available. It is also necessary to understand that some investigations are conducted to find new encapsulation materials, meaning that although they are not currently present in the FDA list, they could be added in the future. Many of them are of natural origin such as starch from *Araucaria angustifolia* (Bertol.) Kuntze seeds,^{100,101} mucilage extract from *Opuntia ficus Indica*⁷² and gelatinized sweet potato starch,¹⁵⁰ and therefore further studies are needed to establish the safety of these materials.

3. Incorporation of microencapsulated bioactives in food matrices

3.1. Bioactive extracts

The main reason to consider a bioactive extract is related to synergistic effects occurring among their components that

often result in increased bioactive characteristics. The information regarding microencapsulated bioactive extracts obtained from different plant materials and other natural matrices after extraction with various solvents is summarized in Table 3. Crude extracts represent a significant part of the microencapsulation studies, followed by polyphenols (as also anthocyanins), essential oils, vitamins, proteins and fat extracts.

The majority of the microencapsulation studies for food purposes have focused on the technique development itself which includes the definition of the best suitable materials and the achievement of microcapsules with the adequate morphology, encapsulation efficiency, stability and release behaviour. The studies calling up the development of final applications, *i.e.* the test of the microencapsulated materials with real food matrices, are much scarcer. Chiou & Langrish⁴⁷ used the crude extract (water) of *Hibiscus sabdariffa* L. for encapsulation with the fibres extracted from the same fruit as the wall material, aiming at developing a novel nutraceutical product using a by-product usually not consumed. A similar study conducted by Berg *et al.*⁷⁰ in which pectin (natural polysaccharide) was used as the encapsulation wall material to protect anthocyanins extracted from *Vaccinium* genus fruits showed that the addition of gelling substances provided a higher encapsulation efficiency. The optimization of encapsulation methodologies is constantly evolving, as is the case of supercritical fluid-based processes, which were used to encapsulate green tea extract from *Camellia sinensis* L. leaves with polycaprolactone (PCL), by high pressure antisolvent coprecipitation demonstrating high retention of catechins in the coprecipitates, and also to encapsulate ethanolic extracts from *Rosmarinus officinalis* L. leaves with poloxamer polymers, with similar results.^{87,130} With a different goal, but intending to improve encapsulation and delivery of bioactive extracts, Averina & Allémann¹¹¹ developed pH sensitive micro- and nanoparticle containing natural sources of polyunsaturated fatty acids, namely oils extracted from *Thymallus baikalensis* Dybowski muscle and *Pinus sibirica* Du Tour seeds, and commercial fish oil, by using the emulsification–diffusion and nanoprecipitation techniques with promising results. Barras *et al.*¹²⁴ developed lipid nanoparticles loaded with polyphenol extracts to enhance their solubility and stability. Many of the studies with phenolic compounds are performed with the main objective to optimize the encapsulation process,^{80,118,125,131} using different types of extracts (*e.g.* alcoholic, aqueous, hydroalcoholic *etc.*). In fact, there is no specific standard protocols for the extraction of each class of phenolic compounds, depending on the nature of the sample and the objective of the work (structure elucidation and quantification).¹⁵¹ In terms of proteins,^{138,152} vitamins,⁸⁸ phyto-sterols¹⁵³ and essential oils,^{57,59,60} the majority of the studies was also conducted with the aim of developing new encapsulating methodologies and materials, and optimizing the process.

After optimization of the encapsulation process, it is necessary to establish whether the extracts maintain, reduce or increase their bioactive characteristics. Therefore, several

bioactivity assays can be conducted to evaluate the antioxidant and antimicrobial activities, and quantify total phenolic compounds. To assess the antioxidant activity, the DPPH (2,2-diphenyl-1-picrylhydrazyl) scavenging activity is the most commonly used assay not only to characterize a given sample, but also to evaluate the bioactivity maintenance. The studies performed by López-Córdoba *et al.*¹³⁶ and Chan *et al.*¹⁵⁴ with crude extracts of *Ilex paraguariensis* A. St. Hil. aerial parts and *Piper sarmentosum* Roxb., respectively, showed that encapsulation did not affect, positively or negatively, the antioxidant activity of the extracts. On the other hand, in the studies conducted by Igual *et al.*⁴⁹ and Parthasarathi *et al.*⁴³ with *Solanum quitoense* L. pulp and *Garcinia cowa* Roxb. fruit, respectively, the encapsulation proved to be very effective, since an increase in the antioxidant activity of the extracts was observed, which can be explained by protection of the bioactives from degradation. Anthocyanin extracts obtained from *Garcinia indica* Choisy fruit pulp,⁶⁸ *Euterpe oleracea* Mart. fruit pulp⁷¹ and *Daucus carota* L. roots⁶⁷ were encapsulated with maltodextrins, which proved to be efficient in protecting these extracts whose stability and antioxidant activity increased after microencapsulation. With another goal Deladino *et al.*⁹⁰ used the DPPH assay to assess the diffusion and kinetic behaviour of the produced microencapsulated system. The oxygen radical absorbance capacity (ORAC), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid assay (ABTS) and trolox equivalent antioxidant capacity (TEAC) assays have also been used to evaluate the antioxidant activity of microencapsulated extracts.^{50,62,76,82,105,115,117} As previously mentioned, the quantification of phenolic compounds is also a very common methodology to assess the effectiveness of the encapsulation process.^{42,44,46,48,64,78,79,85,92,98,141,155} Some studies also describe the use of carotenoids to infer the efficacy of the microencapsulation process.^{94,141}

Antibacterial and antifungal properties are among the most studied and important bioactivities not only due to the increasing resistance of the microorganisms to commercially available synthetic antibiotics, but also because natural matrices present great potential for the discovery of new drugs. There are several studies focusing on the microencapsulation of natural extracts presenting antibacterial and antifungal activities. Sansone *et al.*⁵² and Fernandes *et al.*⁴⁵ reported the antifungal activity of *Paeonia rockii* (S.G. Haw & Lauener) roots and *Lippia sidoides* Cham. leaves, respectively, showing the advantage of their microencapsulation and enhancement of the antifungal activity was obtained compared with the extracts in the free form. The antibacterial activity of the essential oil extracted from *Citrus hydrix* D.C. fruit skins was assessed by Adamiec *et al.*,⁵⁸ also reporting the enhancement of the activity in the microencapsulated extract. Souza *et al.*⁸⁶ studied the antimicrobial effect of *Vitis labrusca* L. ethanol–water (67.6%) encapsulated extract, showing a very good growth inhibiting capacity of *Staphylococcus aureus* and *Listeria monocytogenes*.

Studies considering the improvement of bone quality in rats¹²¹ and *in vitro* cytotoxicity¹⁰⁷ were performed with micro-

Table 3 Microencapsulated bioactive extracts

Bioactive extract	Source	Extraction solvent	Ref.
Anthocyanin extracts	<i>Bactris guineensis</i> L. fruits	Methanol–acetic acid (19 : 1, v/v)	69
	<i>Daucus carota</i> L. roots	Ethanol	67
	<i>Euterpe oleracea</i> Mart. fruit pulp	Juice	71
	<i>Garcinia indica</i> Choisy fruit pulp	Acidified water	68
	<i>Myrciaria cauliflora</i> (Mart.) fruit peels	Acidified ethanol	82, 94
	<i>Vaccinium</i> genus fruits	n.a.	70, 110, 118
Crude extracts	<i>Bidens pilosa</i> L. aerial parts	Ethanol	48
	<i>Camellia sinensis</i> L. leaves	Acetone; ethanol	121, 130
	<i>Eugenia uniflora</i> L. fruits	Juice	141
	<i>Fadogia ancyllantha</i> Schweinf. aerial parts	Ethanol–water (70 : 30, v/v)	44
	<i>Garcinia cowa</i> Roxb fruits	Water	43
	<i>Hibiscus sabdariffa</i> L fruits	Water	47, 50
	<i>Ilex paraguariensis</i> A. St. Hil. aerial parts	Water	136
	<i>Ipomoea batatas</i> L. Lam variety, Sinjami tuber	n.a.	42
	<i>Lippia sidoides</i> Cham. Leaves	Ethanol–water (50 : 50, v/v)	45
	<i>Melissa officinalis</i> L. aerial parts	Ethanol–water (70 : 30, v/v)	44
	<i>Morinda citrifolia</i> L. fruits	Ethyl acetate	46
	<i>Paeonia rockii</i> (S.G. Haw & Lauener) roots	Polar	52
	Five herbs: <i>Paeonia suffruticosa</i> Andrews, <i>Phellodendron chinense</i> Schneid, <i>Lonicera japonica</i> Thunb, <i>Mentha Spicata</i> L. and <i>Atractylodes lancea</i> Thunb.	Water	106
	<i>Piper sarmentosum</i> Roxb.	Water	154
	Propolis	Ethanol	133
	<i>Quercus resinosa</i> Liebm. leaves	Water	51
	<i>Solanum quitoense</i> L. pulp	n.a.	49
	<i>Tussilago farfara</i> L.	n.a.	44
	Crude and fatty acid extracts	Fish oil	Hydrolysis
<i>Pinus sibirica</i> Du Tour seeds		n.a.	111
<i>Thymallus baikalensis</i> Dybowski muscle		Ethanol	111
Essential oil extracts	<i>Citrus hydrix</i> D.C. fruit skins	Water	58
	<i>Cymbopogon nardus</i> G. aerial parts	n.a.	57
	<i>Majorana hortensis</i> L. aerial parts	n.a.	57
	<i>Origanum vulgare</i> L. aerial parts	n.a.	57
	<i>Origanum vulgare</i> L. flowers and leaves	Water	59, 60, 62
Fatty acid extracts	Commercial	n.a.	63, 65, 119
	<i>Hibiscus cannabinus</i> L. seeds	Hexane	64
Phytosterol ester extracts	Commercial	n.a.	153
Polyphenol extracts	<i>Achillea millefolium</i> L. aerial parts	Water	105
	<i>Cabernet Sauvignon</i> fruits	Juice (wine)	98
	<i>Camellia sinensis</i> L. leaves	Ethanol	107
	Commercial	n.a.	122, 124, 156
	<i>Crategus laevigata</i> (Poir.) Dc. aerial parts	Water	105
	<i>Glechoma hederacea</i> L. aerial parts	Water	105
	<i>Hypericum perforatum</i> L. leaves and flowers	Methanol	155
	<i>Ilex paraguariensis</i> A. St. Hil. aerial parts	Water	90
	<i>Myrica</i> genus fruits	Ethanol	143
	<i>Olea europea</i> L. leaves	Water	105
	<i>Orthosiphon stamineus</i> Benth leaves	Methanol–water (50 : 50, v/v)	79
	<i>Prunus cerasus</i> L. pomace	Ethanol–water (50 : 50, v/v)	131
	<i>Punica granatum</i> L. fruits	Ethanol and juice	85
	<i>Punica granatum</i> L. peels	Water	77
	<i>Quercus resinosa</i> Liebm. leaves	Water	78
	<i>Ribes nigrum</i> L. pomace	Ethanol–water–citric acid (80 : 20 v/v; 5%)	76
	<i>Rosmarinus officinalis</i> L. leaves	Ethanol	87
	<i>Rubus chamaemorus</i> L. fruits	Water–acetone (70 : 30, v/v)	97

Table 3 (Contd.)

Bioactive extract	Source	Extraction solvent	Ref.
	<i>Rubus idaeus</i> L. leaves	Water	105
	<i>Rubus ulmifolius</i> Schott flowers	Methanol–water (80 : 20, v/v)	92
	<i>Urtica dioica</i> L. leaves	Water	105
	<i>Vaccinium myrtillus</i> L. fruits	n.a.	117
	<i>Vitis labrusca</i> L. seeds and fruits	Water–ethanol (67.6 : 32.4, v/v)	86
	<i>Vitis vinifera</i> L. seeds	Buffer acetate	125
	<i>Aristolelia chilensis</i> [Molina] Stuntz leaves	Ethanol–water (40 : 60, v/v)	115
Polyphenol and betalain extracts	<i>Opuntia ficus-indica</i> fruits	Juice and ethanol	80
Polyphenol and oil extracts	Commercial	n.a.	128
Protein extracts	Commercial	n.a.	138
	<i>Pisum sativum</i> L. grain	n.a.	152
Vitamin extracts	<i>Capsicum annuum</i> L. variety Piquillo seeds, skins and stems	CO ₂	88
Vitamin and enzyme extracts	Commercial	n.a.	160
Oil extracts	Commercial	n.a.	120, 150

n.a. – information not available.

encapsulated *C. sinensis* tea. The antioxidant α -glucosidase inhibitory activity of microencapsulated aqueous extracts from *Punica granatum* L. peel and the anti-inflammatory effect of commercial polyphenols and oil extracts were also studied.^{77,128}

As can be observed in Fig. 4, *in vitro* release studies are among the most important steps to consider when developing and validating a microencapsulated product. A successful microencapsulated system not only has to protect the bioactive compound ensuring its bioavailability, but also needs to guarantee the intended release behaviour (temporal and target oriented). *In vitro* release studies can be performed by simulating the gastrointestinal environment using pH buffers mimicking the conditions of digestion,^{106,156} or using *in vitro* gastrointestinal models comprising enzymes and pH buffers.^{110,133,143,150} Tavano *et al.*¹⁵⁶ showed, by *in vitro* release studies, that curcumin and quercetin when microencapsulated in niosomes improved the solubility after gastrointestinal digestion. Frank *et al.*¹¹⁰ and Park *et al.*¹⁵⁰ reported that after *in vitro* gastrointestinal digestion, microencapsulated anthocyanin extracts of *V. myrtillus* and a commercial oil extract, respectively, presented good resistance to pH change during digestion, being released only under intestinal conditions. This corroborates the efficacy of microencapsulation in designing adequate delivery systems for water and non-water soluble compounds to be incorporated in innovative food products.

3.2. Bioactive compounds

The importance of studying individual bioactive compounds lies in their powerful bioactivities, with different applications, such as in pharmaceutical and food industries. In this context their isolation from the original matrix is an interesting topic of study and provides an added value to the developed pro-

ducts. A set of microencapsulated individual bioactive compounds used for food application purposes is described in Table 4. The number of articles concerning the encapsulation of individual compounds is markedly lower than that for bioactive extracts. However, phenolic compounds are once again the individual molecules most commonly used in microencapsulation experiments. Most of these studies are focused on the development and optimization of microencapsulation techniques,^{74,82,132,140,144,145,157} including new encapsulation materials. An example is the work performed by Medina-Torres *et al.*,⁷² in which commercial gallic acid was encapsulated using mucilage extracted from *O. ficus Indica*. Robert *et al.*⁷³ also encapsulated gallic acid using acetylated starch and inulin, obtaining higher encapsulation efficiency with the first material. On the other hand, for quercetin and vanillin phenolic compounds, inulin gave the best results.⁸¹ Despite the beneficial health effects of phenolic compounds, their stability and bioavailability are severely compromised during food processing, storage and digestion, as mentioned in the previous sections. So, microencapsulation of individual phenolic compounds could provide a way to maintain or increase their antioxidant activity,^{114,139} stability^{75,97} and bioavailability.^{96,127} The antimicrobial activity was also tested in microcapsules containing chlorogenic acid isolated from *Nicotiana tabacum* L. leaves, indicating that its activity was not affected by microencapsulation, being an alternative in the development of food products with antimicrobial properties.¹⁵⁸

Polyunsaturated fatty acids are also the target of microencapsulation studies. Their known beneficial health effects make them very appealing to enrich food matrices. However, their lipophilic nature and rancidity tendency are obstacles in the development of efficient delivery systems. Naik *et al.*¹⁰² developed an encapsulation technique for the delivery of

Table 4 Microencapsulated individual bioactive compounds

Class	Individual bioactive compounds	Source	Ref.
Carotenoids	Curcumin	Commercial	114, 127, 159
	Lutein	Commercial	104
	β -Carotene	Commercial	48, 100,101
	β -Carotene	<i>Capsicum annuum</i> L. fruits	53
Carotenoids and vitamins	Curcumin and retinol	Commercial	109
Enzymes	Cellulases and xylanases	Commercial	55
	Coenzyme Q10	Commercial	56
Essential oil	Cardamom oleoresin	Commercial	137
	Engenol and eugenyl acetate	<i>Syzygium aromaticum</i> L. buds	61
Fatty acid	α -Linolenic acid	<i>Lepidium sativum</i> Linn. seeds	102
	ω -3 Fatty acids	Commercial	66, 129
Phenolic compounds	Caffeine	Commercial	157
	Catechins	<i>Camellia sinensis</i> L. leaves	96
	Chlorogenic acid	<i>Nicotiana tabacum</i> L. leaves	158
	Ellagic acid	Commercial	126
	Gallic acid	Commercial	72, 73, 140
	Isoflavone	Commercial	116
	Mangiferin	<i>Mangifera indica</i> L. bark	74
	Naringenin and quercetin	Commercial	75
	Quercetin	Commercial	139
	Quercetin and vanillin	Commercial	81
	Quercitrin	<i>Albizia chinensis</i> L. flowers (90 : 10, v/v)	145
	Resveratrol	<i>Arachis hypogaea</i> L. sprout	144
	Resveratrol	<i>Polygonum cuspidatum</i> Siebold & Zucc roots	132
	Rutin and anthocyanins	<i>Hibiscus sabdariffa</i> L. dried calyx	135
Proteins	Albumin and hirudin	Commercial	93
	Papain	Commercial	91
Organic acids	Citric acid	Commercial	123
	(-)-Hydroxycitric acid	<i>Garcinia cowa</i> Roxb fruit	142
Organosulfur compound	Allicin	<i>Allium sativum</i> L. bulb cloves	83, 84, 99
Vitamins	Folic acid	Commercial	134
	Riboflavin (vitamin B ₂)	Commercial	123
Mixtures of bioactives	Fish oil, resveratrol, tributyrin	Commercial	123
	Glucose, vitamin B12, olive oil	Commercial	103, 146
	Fish oil, phytosterols (5 α -cholestane, β -sitosterol, campesterol and stigmasterol) and limonene	Commercial	89, 113

α -linoleic acid isolated from the seeds of *Lepidium sativum* Linn. using freeze drying to achieve a stable and bioavailable compound. On the other hand, Shaw *et al.*⁶⁶ and Rasti *et al.*¹²⁹ developed different lipophilic delivery systems for commercial ω -3-fatty acids. Shaw *et al.*⁶⁶ used the spray-drying technique with lecithin and chitosan as the wall material, to prevent lipid oxidation and to study the reconstruction of the enriched microcapsules in aqueous medium, showing that this multi-layer system is very promising. Rasti *et al.*¹²⁹ used liposome based delivery systems to microencapsulate ω -3-fatty acids, using soybean phospholipids as the wall material. The authors demonstrated that the formation of liposomes in aqueous medium, combined with the antioxidant protection of the phospholipids, increased the stability and prevented fatty acid peroxidation. Other compounds, also very unstable and there-

fore benefiting from microencapsulation, are essential oils or their constituents. In addition to the lipophilic character they are also very volatile, needing the protection by microencapsulation. Lipid carriers involve the formulation of a lipidic solution containing solid lipids, surfactants and drying carriers (*e.g.* polysaccharides) and have provided high encapsulation efficiencies for eugenol and eugenyl acetate isolated from *Syzygium aromaticum* L. buds.⁶¹ Microencapsulation by co-crystallization of cardamom oleoresin also protected their major components, 1,8-cineole and α -terpinyl acetate; nevertheless, some degradation occurred during packaging and storage.¹³⁷

Carotenoids are a family of compounds largely used for food coloration in place of synthetic dyes, presenting additionally antioxidant and antiangiogenic effects. Nevertheless, their tendency to oxidation and isomerization is high. Qv *et al.*¹⁰⁴

and Xu *et al.*¹⁵⁹ studied the stability of lutein and curcumin, respectively, after microencapsulation by complex coacervation with Ca-alginate/k-carrageenan, and Ca-alginate/lysozyme, respectively. Both achieved good encapsulation efficiencies and demonstrated the efficacy of the used method. Spada *et al.*^{100,101} microencapsulated commercial β -carotene in starch obtained from *Araucaria angustifolia* (Bertol.) Kuntze seeds, and concluded that a modified gelation form of this starch led to higher carotenoid encapsulation efficiency ensuring protection against adverse conditions. Aissa *et al.*⁵⁴ tested microcapsules enriched with β -carotene for its genotoxic and antiangiogenic effects, using arabic gum as the wall material. The authors observed preservation of the genotoxic effects, but a decrease in antiangiogenic activity, maybe due to the loss of bioavailability during microencapsulation.

Organic acids,^{83,84,99,142} enzymes^{55,56} and proteins^{91,93} are examples of other individual compounds that have been subjected to microencapsulation.

Vitamin B₂ (riboflavin) and vitamin B₉ (folic acid) have also been microencapsulated for food purposes. Due to their known beneficial health effects, coupled with a high tendency to degradation and loss of bioavailability, *in vitro* release tests were used to evaluate new delivery systems. Chen & Subirade¹¹³ tested the release of riboflavin using simulated gastric, intestinal and pancreatic fluids, concluding that riboflavin microcapsules made of alginate/whey protein are semi-destroyed by the intestinal fluid and completely released with the pancreatic fluid. To estimate the product shelf life, Wichchukit *et al.*⁸⁹ studied the release of riboflavin incorporated into a food product, a model beverage. Prasertmanakit *et al.*¹⁴⁶ studied the *in vitro* release of folic acid from ethyl cellulose microcapsules, a material that had good encapsulation efficiency. The addition of a water soluble polymer, sucrose, caused swelling of the polymer matrix, which allowed better controlled release of folic acid.

An improvement in delivery system development is the encapsulation of a mixture of bioactive compounds within the same microcapsule, thereby obtaining several beneficial effects. Augustin *et al.*¹¹² developed an oil-in-water emulsion to stabilize commercial fish oil, resveratrol and tributyrin using caseinate, glucose and starch, to study their behaviour in the gastrointestinal tract, obtaining increased bioavailability for all the compounds. Pan *et al.*¹⁰⁹ studied the oxidative stability of curcumin (carotenoid) and retinol (essential oil) in oil-in-water emulsions, with very satisfactory results.

3.3. Incorporation in food matrices

Some examples of applicability studies with microencapsulated bioactive extracts or individual compounds are described in Table 5. After an exhaustive search in the literature, it was confirmed that the vast majority of the studies do not include the validation of the developed microencapsulated bioactives through their incorporation into food matrices. Only twelve studies were found where this final step, so important for the food industry, was included. In general, milk and dairy products such as cheese and yoghurt, and ice creams are the pre-

ferable food matrices under study. The sector of cereals, bread and pasta, is also significant in applicability studies. Tea, soup and meat are also food matrices that have been tested for incorporation of bioactive microcapsules. Phenolic extracts of *Punica granatum* L. peels were studied by Çam *et al.*⁷⁷ and were added to ice cream to enhance antioxidant and α -glucosidase inhibitory activities. Martins *et al.*⁹² and Robert *et al.*⁸⁵ also incorporated phenolic extracts in yogurt using *Rubus ulmifolius* Schott flowers and *Punica granatum* L. fruits, respectively. Martins *et al.*⁹² obtained a higher antioxidant activity in yogurt with microencapsulated extracts, compared with the use of extracts in the free form and with the control (yogurts without extracts); on the other hand, Robert *et al.*⁸⁵ also reported a higher content of phenolic compounds and anthocyanins in yogurt with microencapsulated extracts. The incorporation technique developed by Barbosa-Pereira *et al.*¹²² to add phenolic extracts in active packaging to extend the shelf-life of meat products gave promising results retarding lipid oxidation and microbial growth. In terms of individual phenolic compounds, a water soluble isoflavone was microencapsulated in a polyglycerol monostearate emulsion and further incorporated in milk to study its stability during storage and after *in vitro* digestion. It was demonstrated that the microencapsulated isoflavone did not affect milk taste and that its absorption in the intestine increased.¹¹⁶

Citric acid and its derivative, (–)-hydroxycitric acid, were also used in incorporation studies; in particular, the derivative extracted from the fruits of *Garcinia cowa* Roxb. was incorporated into bread^{83,99} and pasta;⁸⁴ in both cases, bread and pasta enriched with microencapsulated bioactives showed good sensory and quality attributes, which proves the viability of using such strategies in food product development. Citric acid was also incorporated in chewing gum at a micronized scale, using a technique based on casein and inulin to form bioactive microcapsules, to develop chewing gums with health promoting properties.¹⁴² Soups, among the most highly consumed food products worldwide, also served as the matrix for the incorporation study developed by Rubilar *et al.*⁶⁵ Microcapsules containing fatty acids (linseed oil) were added to an instant soup in powder form in order to develop a new functional product; moreover, since the linseed oil was microencapsulated in a polymeric matrix consisting of arabic gum and maltodextrin, a higher controlled release of the lipophilic core was successfully achieved. Sardar *et al.*¹³⁷ also encapsulated a lipophilic compound, cardamom oleoresin. Since the stability of this compound for spray-drying was very poor, a sucrose wall matrix was used with a co-crystallization method giving rise to small flavouring sugar cubes for tea beverages. The produced cubes were stable to storage when packed in a three-layer metalized laminate. Cheese, although appreciated by many consumers, is rich in fat and, therefore, there have been efforts for the addition of vegetable oils to this matrix. However, oils degrade very quickly, benefiting from the addition of antioxidants such as vitamins A and E and coenzymes. In this context, the work of Stratulat *et al.*¹⁶⁰ was intended to inhibit lipid peroxidation (rancidity), by formulat-

Table 5 Examples of studies with microencapsulated bioactive extracts or individual compounds incorporated in food matrices

Food matrix	Bioactive	Source	Encapsulation method	Encapsulation material	Ref.
Bread	(-)-Hydroxycitric acid	<i>Garcinia cowa</i> Roxb fruit rinds	Spray-drying	Whey protein and maltodextrin	83
	(-)-Hydroxycitric acid	<i>Garcinia cowa</i> Roxb fruit rinds	Freeze-drying	Whey protein and maltodextrin	99
Cheese	Vitamins E and A; Coenzyme 10	Commercial	Emulsion	Calcium caseinate	160
Chewing gum	Citric acid	Commercial	Microwave	Casein and inulin	142
Ice cream	Phenolic extracts	<i>Punica granatum</i> L. peels	Spray drying	Maltodextrin	77
Meat	Phenolic extracts	Residues from brewing industry	Extrusion	Ethylene vinyl acetate and LDPE	122
Milk	Isoflavone	Commercial	Emulsion	Polyglycerol monostearate	116
Pasta	(-)-Hydroxycitric acid	<i>Garcinia cowa</i> Roxb. fruits	Spray-drying	Whey protein	84
Soup	Fatty acid (linseed oil)	Commercial	Spray-drying	Gum arabic and maltodextrin	65
Tea	Cardamom oleoresin	Commercial	Co-crystallization	Sucrose	137
Yogurt	Phenolic extract	<i>Rubus ulmifolius</i> Schott flowers	Atomization/coagulation	Alginate	92
	Phenolic extract	<i>Punica granatum</i> L. fruits	Spray-drying	Maltodextrin or soybean protein	85

ing emulsions, stabilized with calcium caseinate, containing vitamins A and E, and Coenzyme Q10. The results showed that the vegetables oils did not affect the cheese stability, due to the presence of antioxidants.

4. Conclusion

Nowadays, food not only serves to satisfy the primal urge of hunger but is also intended to overcome dietary flaws and/or impart health benefits. Bioactives are sources of functional molecules with recognized health effects in populations that otherwise would not be able to benefit from them. Nevertheless, they show organoleptic constraints and instability to food processes, storage and ingestion, which has led to research in the field of bioactive protection and controlled release. Among the proposed technologies, microencapsulation emerged as a viable route to valorise natural bioactives in functional foods, thus extending their benefits to a wider population.

According to the present review, there are several examples available of microencapsulation of bioactives using a wide range of processes and encapsulating materials. Among the various possibilities, the spray-based processes, *e.g.* spray-drying, are the most commonly used techniques. The advantages are its easy implementation, namely at the industrial level, and the fact of being inexpensive. Nevertheless, green techniques, such as supercritical and ultrasound based processes, are nowadays attracting much attention.

Water soluble materials, both polymer and non-polymer ones, are the most commonly used encapsulation materials. They include carbohydrate polymers (starch and cellulose and their derivatives), plant exudates and extracts (gum, galactomannans and pectins), marine extracts (carrageenan and alginate), and microbial and animal derived polysaccharides (xanthan, gellan, dextran and chitosan). In most of the cases,

the industrial applicability in the field of food production is prevented by current regulations.

Crude and phenolic extracts, together with individual phenolic compounds, are the most studied bioactives for food purposes. Nevertheless, studies dealing with final food applications are scarce, demanding investment from academia, industry and regulatory agencies. Finally, the consumers also have a crucial role in the acceptance of new products in the market.

Acknowledgements

The authors are grateful to Fundação para a Ciência e a Tecnologia (FCT, Portugal) for financial support to CIMO (strategic project PEst-OE/AGR/UI0690/2011) and LSRE (PEst-C/EQB/LA0020/2011), and M. I. Dias grant (SFRH/BD/84485/2012). Financial support from QREN, ON2 and FEDER (NORTE-07-0124-FEDER-000014) is also acknowledged.

Notes and references

- 1 J. A. Milner, *Am. J. Clin. Nutr.*, 2010, 7, 1654–1659.
- 2 European Parliament, Regulation (EC) No. 1924/2006, 2006.
- 3 EFSA, Regulation (EC) No. 1924/2006, 2006.
- 4 FDA, Guide to nutrition labeling and education act (NLEA) requirements, 1994.
- 5 J. C. Espín, M. T. Carcía-Conesa and F. A. Tomás-Barberán, *Phytochemistry*, 2007, 68, 2986–3008.
- 6 B. Bigliardia and F. Galati, *Trends Food Sci. Technol.*, 2013, 31, 118–129.
- 7 I. Siró, E. Kápolna, B. Kápolna and A. Lugasi, *Appetite*, 2008, 51, 456–467.

- 8 I. J. Joye, G. Davidov-Pardo and D. J. McClements, *Trends Food Sci. Technol.*, 2014, **40**, 1–15.
- 9 R. Wu, B. Frei, J. A. Kennedy and Y. Zhao, *LWT-Food Sci. Technol.*, 2010, **43**, 1253–1264.
- 10 H. Zielinski, H. Kozłowska and B. Lewczuk, *Innovative Food Sci. Emerging Technol.*, 2001, **2**, 159–169.
- 11 A. Rawson, A. Patras, B. K. Tiwari, F. Noci, T. Koutchma and N. Brunton, *Food Res. Int.*, 2011, **44**, 1875–1887.
- 12 M. C. Nicoli, M. Anese and M. Parpinel, *Trends Food Sci. Technol.*, 1999, **10**, 94–100.
- 13 R. Puupponen-Pimiä, S. T. Häkkinen, M. Aarni, T. Suortti, A. Lampi, M. Eurola, V. Piironen, A. M. Nuutila and K. Oksman-Caldentey, *J. Sci. Food Agric.*, 2003, **83**, 1389–1402.
- 14 A. Fernandes, A. L. Antonio, M. B. P. P. Oliveira, A. Martins and I. C. F. R. Ferreira, *Food Chem.*, 2012, **135**, 641–650.
- 15 B. Holst and G. Williamson, *Curr. Opin. Biotechnol.*, 2008, **19**, 73–82.
- 16 S. Y. Leong and I. Oey, *Food Chem.*, 2012, **133**, 1577–1578.
- 17 Z. Li, H. Jiang, C. Xu and L. Gu, *Food Hydrocolloids*, 2015, **43**, 153–164.
- 18 I. Fernandes, A. Faria, C. Calhau, V. Freitas and N. Mateus, *J. Funct. Foods*, 2014, **7**, 54–66.
- 19 D. B. Rodriguez-Amaya, *J. Food Compos. Anal.*, 2010, **23**, 726–740.
- 20 E. Fernández-García, I. Carvajal-Lérida, M. Jarén-Galán, J. Garrido-Fernández, A. Pérez-Gálvez and D. Hornero-Méndez, *Food Res. Int.*, 2012, **46**, 438–450.
- 21 A. Amalraj and A. Pius, *Food Chem.*, 2015, **170**, 430–436.
- 22 I. Vermaak, A. M. Viljoen, J. H. Hamman and S. F. V. Vuuren, *Phytochem. Lett.*, 2010, **2**, 166–170.
- 23 D. J. McClements, E. A. Decker and J. Weiss, *J. Food Sci.*, 2007, **72**, 109–124.
- 24 D. J. McClements and Y. Li, *Adv. Colloid Interface Sci.*, 2010, **159**, 213–228.
- 25 W. Faisal, T. Ruane-O’Hora, C. M. O’Driscoll and B. T. Griffin, *Int. J. Pharm.*, 2013, **453**, 307–314.
- 26 P. Balakrishnan, B. Lee, D. H. Oh, J. O. Kim, Y. Lee, D. Kim, J. Jee, Y. Lee, J. S. Woo, C. S. Yong and H. Choi, *Int. J. Pharm.*, 2010, **374**, 66–72.
- 27 T. Bech-Larsen and J. Scholderer, *Trends Food Sci. Technol.*, 2007, **18**, 231–234.
- 28 A. Drewnowski and C. Gomez-Carneros, *Am. J. Clin. Nutr.*, 2000, **72**, 1424–1435.
- 29 Z. Fang and B. Bhandari, *Trends Food Sci. Technol.*, 2010, **21**, 510–523.
- 30 S. S. Kuang, J. C. Oliveira and A. M. Crean, *Crit. Rev. Food Sci. Nutr.*, 2010, **50**, 951–968.
- 31 S. Heleno, A. Martins, M. J. R. P. Queiroz and I. C. F. R. Ferreira, *Food Chem.*, 2015, **173**, 501–513.
- 32 S. Wang, R. Su, S. Nie, M. Sun, J. Zhang, D. Wu and N. Moustaid-Moussa, *J. Nutr. Biochem.*, 2014, **25**, 363–376.
- 33 G. Davidov-Pardo and D. J. McClements, *Trends Food Sci. Technol.*, 2014, **38**, 88–103.
- 34 F. Nazzaro, P. Orlando, F. Fratianni and R. Coppola, *Curr. Opin. Biotechnol.*, 2012, **23**, 182–186.
- 35 I. M. Martins, M. F. Barreiro, M. Coelho and A. E. Rodrigues, *Chem. Eng. J.*, 2014, **245**, 191–200.
- 36 H.-S. Kwak, *Nano- and Microencapsulation for Foods*, John Wiley & Sons, United Kingdom, 2014.
- 37 V. Nedovic, A. Kalusevic, V. Manojlovic, S. Levic and B. Bugarski, *Procedia Food Sci.*, 2011, **1**, 1806–1815.
- 38 P. M. M. Schrooyen, R. van der Meer and C. G. De Kruif, *Proc. Nutr. Soc.*, 2001, **60**, 475–479.
- 39 S. Gouin, *Trends Food Sci. Technol.*, 2004, **15**, 330–347.
- 40 C. P. Champagne and P. Fustier, *Curr. Opin. Biotechnol.*, 2007, **18**, 184–190.
- 41 A. Gharsallaoui, G. Roudaut, O. Chambin, A. Voilley and R. Saurel, *Food Res. Int.*, 2007, **40**, 1107–1121.
- 42 M. Ahmed, M. S. Akter, J. Lee and J. Eun, *LWT-Food Sci. Technol.*, 2010, **43**, 1307–1312.
- 43 S. Parthasarathi, P. N. Ezhilarasi, B. S. Jena and C. Anandharamakrishnan, *Food Bioprod. Process.*, 2013, **9**, 103–110.
- 44 F. Sansone, T. Mencherini, P. Picerno, M. d’Amore, R. P. Aquino and M. R. Lauro, *J. Food Eng.*, 2011, **105**, 468–476.
- 45 L. P. Fernandes, R. C. Candido and W. P. Oliveira, *Food Bioprod. Process.*, 2012, **90**, 425–432.
- 46 D. Krishnaiah, R. Sarbatly and R. Nithyanandam, *Chem. Eng. Res. Des.*, 2012, **90**, 622–632.
- 47 D. Chiou and T. A. G. Langrish, *J. Food Eng.*, 2007, **82**, 84–91.
- 48 D. F. Cortés-Rojas, C. R. F. Souza and W. P. Oliveira, *Chem. Eng. Res. Des.*, 2015, **93**, 366–376.
- 49 M. Igual, S. Ramires, L. H. Mosquera and N. Martínez-Navarrete, *Powder Technol.*, 2014, **256**, 233–238.
- 50 T. A. G. Langrish and R. Premarajah, *Adv. Powder Technol.*, 2013, **24**, 771–779.
- 51 N. E. Rocha-Guzmán, J. A. Gallegos-Infante, R. A. González-Laredo, F. Harte, L. Medina-Torres, L. A. Ochoa-Martínez and M. Soto-García, *J. Food Sci.*, 2010, **75**, 57–61.
- 52 F. Sansone, P. Picerno, T. Mencherini, A. Porta, M. R. Lauro, P. Russo and R. P. Aquino, *J. Food Eng.*, 2014, **120**, 260–267.
- 53 A. Y. Guadarrama-Lezama, L. Dorantes-Alvarez, M. E. Jaramillo-Flores, C. Pérez-Alonso, K. Niranjan, G. F. Gutiérrez-López and L. Alamilla-Beltrán, *J. Food Eng.*, 2012, **112**, 29–37.
- 54 A. F. Aissa, M. L.P. Bianchi, J. C. Ribeiro, L. C. Hernandez, A. F. Faria, A. Z. Mercadante and L. M. G. Antunes, *Food Chem. Toxicol.*, 2012, **50**, 1418–1424.
- 55 M. Santa-Maria, H. Scher and T. Jeoh, *J. Microencapsulation*, 2012, **29**, 286–295.
- 56 M. V. Bule, R. S. Singhal and J. F. Kennedy, *Carbohydr. Polym.*, 2010, **82**, 1290–1296.
- 57 R. Baranauskienė, R. R. Venskutonis, K. Dewettinck and R. Verhé, *Food Res. Int.*, 2006, **39**, 413–425.

- 58 J. Adamiec, C. Borompichaichartkul, G. Srzednicki, W. Panket, S. Piriypunsakul and J. Zhao, *Dry. Technol.*, 2012, **30**, 914–920.
- 59 J. M. G. Costa, S. V. Borges, A. A. C. T. Hijo, E. K. Silva, G. R. Marques, M. A. Cirillo and V. M. Azevedo, *J. Microencapsulation*, 2013, **30**, 717–727.
- 60 L. C. Garcia, R. V. Tonon and M. D. Hubinger, *Dry. Technol.*, 2012, **30**, 1413–1421.
- 61 D. F. Cortés-Rojas, C. R. F. Souza and W. P. Oliveira, *J. Food Eng.*, 2014, **127**, 34–42.
- 62 A. P. Almeida, S. Rodríguez-Rojo, A. T. Serra, H. Vila-Real, A. L. Simplicio, I. Delgado, S. B. Costa, L. B. Costa, I. D. Nogueira and C. M. M. Duarte, *Innovative Food. Sci. Emerging Technol.*, 2013, **20**, 140–145.
- 63 G. Gallardo, L. Guida, V. Martinez, M. C. López, D. Bernhardt, R. Blasco, R. Pedroza-Islas and L. G. Hermida, *Food Res. Int.*, 2013, **52**, 473–482.
- 64 S. Ng, P. Wong, C. Tan, K. Long and K. Nyam, *Eur. J. Lipid Sci. Technol.*, 2013, **115**, 1309–1318.
- 65 M. Rubilar, E. Morales, K. Contreras, C. Ceballos, F. Acevedo, M. Villarreal and C. Shene, *Eur. J. Lipid Sci. Technol.*, 2012, **114**, 423–433.
- 66 L. A. Shaw, D. J. McClements and E. A. Decker, *J. Agric. Food Chem.*, 2007, **55**, 3112–3119.
- 67 S. Ersus and U. Yurdagel, *J. Food Eng.*, 2007, **80**, 805–812.
- 68 C. A. Nayak and N. K. Rastogi, *Dry. Technol.*, 2010, **28**, 1396–1404.
- 69 C. Osorio, B. Acevedo, S. Hillebrand, J. Carriazo, P. Winterhalter and A. L. Morales, *J. Agric. Food Chem.*, 2012, **58**, 6977–6985.
- 70 S. Berg, M. Bretz, E. A. Hubbermann and K. Schwarz, *J. Food Eng.*, 2012, **108**, 158–165.
- 71 R. V. Tonon, C. Brabet and M. D. Hubinger, *Food Res. Int.*, 2010, **43**, 907–914.
- 72 L. Medina-Torres, E. E. García-Cruz, F. Calderas, R. F. G. Laredo, G. Sánchez-Olivares, J. A. Gallegos-Infante, N. E. Rocha-Guzmán and J. Rodríguez-Ramírez, *LWT-Food Sci. Technol.*, 2013, **50**, 642–650.
- 73 P. Robert, P. García, N. Reyes, J. Chávez and J. Santos, *Food Chem.*, 2012, **134**, 1–8.
- 74 J. R. R. Souza, J. P. A. Feitosa, N. P. S. Ricardo, M. T. S. Trevisan, H. C. B. Paula, C. M. Ulrich and R. W. Owen, *Food Hydrocolloids*, 2013, **33**, 10–18.
- 75 F. Sansone, P. Picerno, T. Mencherini, F. Villecco, A. M. D'Ursi, R. P. Aquino and M. R. Lauro, *J. Food Eng.*, 2011, **103**, 188–196.
- 76 A. M. Bakowska-Barczak and P. P. Kolodziejczyk, *Ind. Crops Prod.*, 2011, **34**, 1301–1309.
- 77 M. Çam, N. C. İcyer and F. Erdoğan, *LWT-Food Sci. Technol.*, 2014, **55**, 117–123.
- 78 J. A. Gallegos-Infante, N. E. Rocha-Guzmán, R. B. González-Laredo, L. Medina-Torres, C. A. Gomez-Aldap, L. A. Ochoa-Martínez, C. A. Martínez-Sánchez, B. A. Hernández-Santos and J. Rodríguez-Ramírez, *Food Biosci.*, 2013, **2**, 31–38.
- 79 S. F. Pang, M. M. Yusoff and J. Gim bun, *Food Hydrocolloids*, 2014, **37**, 159–165.
- 80 C. Saénz, S. Tapia, J. Chávez and P. Robert, *Food Chem.*, 2009, **114**, 616–622.
- 81 D. Sun-Waterhouse, S. S. Wadhwa and G. I. N. Waterhouse, *Food Bioprocess Technol.*, 2013, **6**, 2376–2388.
- 82 P. I. Silva, P. C. Stringheta, R. F. Teófilo and I. R. N. Oliveira, *J. Food Eng.*, 2013, **117**, 538–544.
- 83 P. N. Ezhilarasi, D. Indrani, B. S. Jena and C. Anandharamakrishnana, *J. Sci. Food Agric.*, 2013, **94**, 1116–1123.
- 84 D. S. Pillai, B. Prabhasankar, B. S. Jena and C. Anandharamakrishnan, *Int. J. Food Prop.*, 2012, **15**, 590–604.
- 85 P. Robert, T. Gorena, N. Romero, E. Sepulveda, J. Chavez and C. Saenz, *Int. J. Food Sci. Technol.*, 2010, **45**, 1386–1394.
- 86 V. B. Souza, A. Fujita, M. Thomazini, E. R. Silva, J. F. Lucon Jr., M. I. Genovese and C. S. Favaro-Trindade, *Food Chem.*, 2014, **164**, 380–386.
- 87 A. Visentin, S. Rodríguez-Rojo, A. Navarrete, D. Maestri and M. J. Cocero, *J. Food Eng.*, 2012, **109**, 9–15.
- 88 A. Romo-Hualde, A. I. Yetano-Cunchillos, C. González-Ferrero, M. J. Sáiz-Abajo and C. J. González-Navarro, *Food Chem.*, 2012, **133**, 1045–1049.
- 89 S. Wichchukit, M. H. Oztop, M. J. McCarthy and K. L. McCarthy, *Food Hydrocolloids*, 2013, **33**, 66–73.
- 90 L. Deladino, P. S. Anbinder, A. S. Navarro and M. N. Martino, *Carbohydr. Polym.*, 2008, **71**, 126–134.
- 91 D. Betancur-Ancona, J. Pacheco-Aguirre, A. Castellanos-Ruelas and L. Chel-Guerrero, *Innovative Food. Sci. Emerging Technol.*, 2011, **12**, 67–72.
- 92 A. Martins, L. Barros, A. M. Carvalho, C. Santos-Buelga, I. P. Fernandes, F. Barreiro and I. C. F. R. Ferreira, *Food Funct.*, 2014, **5**, 1091–1100.
- 93 T. Chandy, D. L. Mooradian and G. H. R. Rao, *J. Appl. Polym. Sci.*, 1998, **70**, 2143–2153.
- 94 D. T. Santos, J. Q. Albarelli, M. M. Beppu and M. Angela, *Food Res. Int.*, 2013, **50**, 617–624.
- 95 C. H. Goh, P. W. S. Heng and L. W. Chan, *Carbohydr. Polym.*, 2012, **88**, 1–12.
- 96 M. H. Jung, P. N. Seong, M. H. Kim, N. Myong and M. Chang, *Nutr. Res. Pract.*, 2013, **7**, 366–372.
- 97 P. Laine, P. Kylli, M. Heinonen and K. Jouppila, *J. Agric. Food Chem.*, 2008, **56**, 11251–11261.
- 98 V. Sanchez, R. Baeza, M. V. Galmarini, M. C. Zamora and J. Chirife, *Food Bioprocess Technol.*, 2011, **6**, 1350–1354.
- 99 P. N. Ezhilarasi, D. Indrani, B. S. Jena and C. Anandharamakrishnana, *J. Food Eng.*, 2013, **117**, 513–520.
- 100 J. C. Spada, C. P. Z. Noreña, L. D. F. Marczak and I. C. Tessaro, *Carbohydr. Polym.*, 2012, **89**, 1166–1173.
- 101 J. C. Spada, L. D. F. Marczak, I. C. Tessaro and C. P. Z. Noreña, *Int. J. Food Sci. Technol.*, 2012, **47**, 186–194.

- 102 A. Naik, V. Meda and S. S. Lele, *Eur. J. Lipid Sci. Technol.*, 2014, **116**, 837–846.
- 103 R. Pérez-Masiá, R. López-Nicolás, M. J. Periago, G. Ros, J. M. Lagaron and A. López-Rubio, *Food Chem.*, 2015, **168**, 124–133.
- 104 X. Qv, Z. Zeng and J. Jiang, *Food Hydrocolloids*, 2011, **25**, 1596–1603.
- 105 A. Belščak-Cvitanović, R. Stojanović, V. Manojlović, D. Komes, I. J. Cindrić, V. Nedović and B. Bugarski, *Food Res. Int.*, 2011, **44**, 1094–1101.
- 106 P. C. Hui, W. Wang, C. Kan, F. S. Ng, E. Wat, V. X. Zhang, C. Chan, C. B. Lau and P. Leung, *Colloids Surf., B*, 2013, **111**, 156–161.
- 107 J. Liang, F. Li, Y. Fang, W. Yang, X. An, L. Zhao, Z. Xin, L. Cao and Q. Hu, *Colloids Surf., B*, 2011, **82**, 297–301.
- 108 Q. Chen, D. McGillivray, J. Wen, F. Zhong and S. Y. Quek, *J. Food Eng.*, 2013, **117**, 505–512.
- 109 Y. Pan, R. V. Tikekar, M. S. Wang, R. J. Avena-Bustillos and N. Nitin, *Food Hydrocolloids*, 2014, **43**, 1–9.
- 110 K. Frank, E. Walz, V. Gräf, R. Greiner, R. Köhler and H. P. Schuchmann, *J. Food Sci.*, 2012, **77**, 50–57.
- 111 E. Averina and E. Allémann, *LWT–Food Sci. Technol.*, 2013, **53**, 271–277.
- 112 M. A. Augustin, M. Y. Abeywardena, G. Patten, R. Head, T. Lockett, A. D. Luca and L. Sanguansri, *J. Funct. Food*, 2011, **3**, 25–37.
- 113 L. Chen and M. Subirade, *Biomaterials*, 2006, **27**, 4646–4654.
- 114 P. Malik, R. K. Ameta and M. Singh, *Chem.-Biol. Interact.*, 2014, **222**, 77–86.
- 115 J. L. Vidal, L. M. Avello, C. Loyola, P. J. Campos, M. P. Aqueveque, S. R. Dungan, L. M. Galotto and M. A. Guarda, *Chil. J. Agric. Res.*, 2012, **73**, 17–23.
- 116 J. S. Seok, J. S. Kim and H. S. Kwak, *Arch. Pharmacol. Res.*, 2003, **26**, 426–431.
- 117 M. Betz, B. Steiner, M. Schantz, J. Oidtmann, K. Mäder, E. Richling and U. Kulozik, *Food Res. Int.*, 2012, **47**, 51–57.
- 118 M. Betz and U. Kulozik, *Procedia Food Sci.*, 2011, **1**, 2047–2056.
- 119 S. S. Gupta and M. Ghosh, *J. Food Eng.*, 2014, **121**, 64–72.
- 120 F. Ostertag, J. Weiss and D. J. McClements, *J. Colloid Interface Sci.*, 2012, **388**, 95–102.
- 121 L. Haidong, Y. Fang, T. Zhihong and R. Changle, *Int. J. Biol. Macromol.*, 2011, **49**, 561–566.
- 122 L. Barbosa-Pereira, I. Angulo, J. M. Lagarón, P. Paseiro-Losada and J. M. Cruz, 2014, **26**, 310–318.
- 123 L. Piazza and T. Roversi, *Procedia Food Sci.*, 2011, **1**, 1374–1380.
- 124 A. Barras, A. Mezzetti, A. Richard, S. Lazzaroni, S. Roux, P. Melnyk, D. Betbeder and N. Monfilliette-Dupont, *Int. J. Pharm.*, 2009, **379**, 270–277.
- 125 M. Gibis, K. Thellmann, C. Thongkaew and J. Weiss, *Food Hydrocolloids*, 2014, **41**, 119–131.
- 126 S. Madrigal-Carballo, S. Lim, G. Rodriguez, A. O. Vila, C. G. Krueger, S. Gunasekaran and J. D. Reed, *J. Funct. Foods*, 2010, **2**, 99–106.
- 127 M. Hasan, N. Belhaj, H. Benachour, M. Barberi-Heyob, C. J. F. Kahn, E. Jabbari, M. Lindera and E. Arab-Tehrany, *Int. J. Pharm.*, 2014, **461**, 519–528.
- 128 M. Coimbra, B. Isacchi, L. V. Bloois, J. S. Torano, A. Keta, X. Wu, F. Broere, J. M. Metselaar, C. J. F. Zijcken, G. Storm, R. Bilia and R. M. Schiffelers, *Int. J. Pharm.*, 2011, **416**, 433–442.
- 129 B. Rasti, S. Jinap, M. R. Mozafari and A. M. Yazid, *Food Chem.*, 2012, **135**, 2761–2770.
- 130 M. V. Sosa, S. Rodríguez-Rojo, F. Mattea, M. Cismondi and M. J. Cocero, *J. Supercrit. Fluids*, 2011, **56**, 304–311.
- 131 B. Cilek, A. Luca, V. Hasirci, S. Sahin and G. Sumnu, *Eur. Food Res. Technol.*, 2012, **235**, 587–596.
- 132 S. Mantegna, A. Binello, L. Boffa, M. Giorgis, C. Cena and G. Cravotto, *Food Chem.*, 2012, **130**, 746–750.
- 133 N. Kalogeropoulos, S. Konteles, I. Mourtzinos, E. Troullidou, A. Chiou and V. T. Karathanos, *J. Microencapsulation*, 2009, **26**, 603–613.
- 134 Y. Li, S. Xu and D. Sun, *J. Food Eng.*, 2007, **83**, 76–83.
- 135 M. Akhtar, B. S. Murray, E. I. Afeisume and S. H. Khew, *Food Hydrocolloids*, 2014, **34**, 62–67.
- 136 A. López-Córdoba, L. Deladino, L. Agudelo-Mesa and M. Martino, *J. Food Eng.*, 2014, **124**, 158–165.
- 137 B. R. Sardar, K. M. Tarade and R. S. Singhal, *J. Food Eng.*, 2013, **117**, 530–537.
- 138 N. Blanco-Pascual, R. B. J. Koldeweij, R. S. A. Stevens, M. P. Montero, M. C. Gómez-Guillén and A. T. T. Cate, *Food Bioprocess Technol.*, 2014, **7**, 2472–2483.
- 139 T. Wu, F. Yen, L. Lin, T. Tsai, C. Lin and T. Cham, *Int. J. Pharm.*, 2008, **346**, 160–168.
- 140 C. G. Rosa, C. D. Borges, R. C. Zambiazzi, M. R. Nunes, E. V. Benvenutti, S. R. Luza, R. F. D'Avila and J. K. Rutz, *Ind. Crops Prod.*, 2013, **46**, 138–146.
- 141 J. K. Rutz, R. C. Zambiazzi, C. D. Borges, F. D. Krumreich, S. R. Luz, N. Hartwig and C. G. Rosa, *Carbohydr. Polym.*, 2013, **98**, 1256–1265.
- 142 S. Abbasi, S. Rahimi and M. Azizi, *J. Microencapsulation*, 2009, **26**, 90–96.
- 143 L. Zheng, Z. Ding, M. Zhang and J. Sun, *J. Food Eng.*, 2011, **104**, 89–95.
- 144 Y. Lee, S. Ahn and H. Kwak, *Food Hydrocolloids*, 2013, **30**, 307–314.
- 145 A. Kumari, S. K. Yadav, Y. B. Pakade, V. Kumar, B. Singh, A. Chaudhary and S. C. Yadav, *Colloids Surf., B*, 2010, **82**, 224–232.
- 146 S. Prasertmanakit, N. Praphairaksit, W. Chiangthong and N. Muangsin, *AAPS PharmSciTech*, 2009, **10**, 1104–1112.
- 147 P. Vos, M. Bučko, P. Gemeiner, M. Navrátil, J. Švitel, M. Faas, B. L. Strand, G. Skjak-Braek, Y. A. Morch, A. Vikartovská, I. Lacík, G. Kolláriková, G. Orive, D. Poncelet, J. L. Pedraz and M. B. Ansorge-Schumacher, *Biomaterials*, 2009, **30**, 2559–2570.

- 148 N. J. Zuidam and V. A. Nedović, in *Encapsulation Technologies for Active Food Ingredients and Food Processing*, Springer Science Business Media, 2010.
- 149 European Parliament, Regulation (EC) No. 1333/2008, 2008.
- 150 K. M. Park, H. Sung, S. J. Choi, Y. J. Choi and P. Chang, *Food Chem.*, 2014, **161**, 53–59.
- 151 C. Santos-Buelga, S. Gonzalez-Manzano, M. Dueñas and A. M. Gonzalez-Paramas, *Natural Products Isolation, Methods in Molecular Biology, Chapter 17- Extraction and isolation of phenolic compounds*, Springer Science+Business Media, 2012.
- 152 A. Gharsallaoui, R. Saurel, O. Chambin and A. Voilley, *Food Bioprocess Technol.*, 2012, **5**, 2211–2221.
- 153 U. V. L. Ma, J. D. Floros and G. R. Ziegler, *Carbohydr. Polym.*, 2011, **83**, 1869–1878.
- 154 E. Chan, Z. Yim, S. Phan, R. F. Mansa and P. Ravindra, *Food Bioprod. Process.*, 2010, **88**, 195–201.
- 155 N. Kalogeropoulos, K. Yannakopoulou, A. Gioxari, A. Chiou and D. P. Makris, *LWT–Food Sci. Technol.*, 2010, **43**, 882–889.
- 156 L. Tavano, R. Muzzalupo, N. Picci and B. Cindio, *Colloids Surf., B*, 2014, **114**, 82–88.
- 157 L. Bagheri, A. Madadlou, M. Yarmand and M. E. Mousavi, *Food Res. Int.*, 2014, **62**, 1113–1119.
- 158 M. Zhao, H. Wang, B. Yang and H. Tao, *Food Chem.*, 2010, **120**, 1138–1142.
- 159 W. Xu, W. Jin, C. Zhang, Z. Li, L. Lin, Q. Huang, S. Ye and B. Li, *Food Res. Int.*, 2014, **59**, 61–66.
- 160 I. Stratulat, M. Britten, S. Salmieri, P. Fustier, D. St-Gelais, C. P. Champagne and M. Lacroix, *J. Funct. Food*, 2014, **6**, 48–59.