

2017

Microplastics are not important for the cycling and bioaccumulation of organic pollutants in the oceans – but should microplastics be considered POPs themselves?

Rainer Lohmann
University of Rhode Island, rlohmann@uri.edu

Follow this and additional works at: <https://digitalcommons.uri.edu/gsofacpubs>

The University of Rhode Island Faculty have made this article openly available.
Please let us know how Open Access to this research benefits you.

This is a pre-publication author manuscript of the final, published article.

Terms of Use

This article is made available under the terms and conditions applicable towards Open Access Policy Articles, as set forth in our [Terms of Use](#).

Citation/Publisher Attribution

Lohmann, Rainer. Microplastics are not important for the cycling and bioaccumulation of organic pollutants in the oceans – but should microplastics be considered POPs themselves? *Integr Env Assess Monit* 2017, 13, 460-465.

Available at: <http://dx.doi.org/10.1002/ieam.1914>

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

1 **Microplastics are not important for the cycling and bioaccumulation of organic pollutants**
2 **in the oceans – but should microplastics be considered POPs themselves?**

3 by **Rainer Lohmann**

4

5 Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island
6 02882-1197, United States.

7 Email: rlohmann@uri.edu

8

9 **Background**

10 This commentary discusses the current evidence about current prevailing themes on the
11 relationship between marine microplastics and organic pollutants. In this context, microplastics
12 can be defined as particles < 5 mm in size [Thompson *et al.*, 2004]. This plastic debris, mostly
13 from anthropogenic land-based sources, fragments into smaller pieces over time [Jambeck *et al.*,
14 2015].

15

16 **Do microplastics accumulate high concentrations of organic pollutants.**

17 The current body of evidence suggests that this is true, as microplastics act – as do polymers in
18 general - as passive samplers of organic pollutants. The specific affinity of a given organic
19 pollutant for a polymer dictates its overall enrichment factor (partitioning constant) in the
20 microplastic [Rusina *et al.*, 2007]. The specific affinity for various hydrophobic organic
21 contaminants (HOCs) for polymers has been determined in numerous laboratory and field
22 calibrations in which the partitioning constants between passive samplers and water or air were
23 measured [Adams *et al.*, 2007; Rusina *et al.*, 2007; Smedes *et al.*, 2009; Lohmann, 2012;

24 *Pintado-Herrera et al.*, 2016; *Ziccardi et al.*, 2016]. The polymers most commonly used as
25 passive samplers in field experiments include polyethylene (PE), silicone rubber (SR) and
26 polyoxymethylene (POM) sheets.

27 There is also plenty of field evidence showing that generic plastic debris accumulates organic
28 pollutants [*Karapanagioti et al.*, 2011; *Rochman et al.*, 2012; *Endo et al.*, 2013]. A prominent
29 example is the so-called ‘pellet watch’ global monitoring program, which relies on plastic pellets
30 collected by volunteers from across the globe [*Hirai et al.*, 2011]. In these studies, strong
31 enrichment of HOCs in the polymers, often exceeding 10^6 times relative to their dissolved
32 concentrations was found.

33

34 **How significantly will microplastics contribute to the dispersion and global cycling of** 35 **POPs?**

36 It has become common knowledge that microplastics are present around the globe, and have
37 been found in all ocean gyres, coastal seas and beaches [*Jambeck et al.*, 2015; *Seville et al.*,
38 2015]. It should therefore be no surprise that the concept of microplastics as being important for
39 the global dispersion of organic pollutants, in particular persistent organic pollutants (POPs), a
40 subgroup of persistent HOCs, has been suggested. Yet, numerous studies have refuted that idea
41 [*Zarfl and Matthies*, 2010; *Gouin et al.*, 2011; *Koelmans et al.*, 2016; *Ziccardi et al.*, 2016].

42 There is simply not enough microplastic and plastic debris present in the oceans to outcompete
43 the partitioning of POPs to water and natural organic matter (such as phytoplankton). In
44 *Koelman et al.* [2016]’s analysis of a strongly HOC (concentrated 10^7 times from water), ocean
45 water nonetheless contained 99% of the HOC, followed by DOC and colloids (0.4% each);
46 microplastics captured $\sim 10^{-4}$ % of the total mass present in the oceans.

47 In addition, diffusion of HOCs in and out of microplastics is slow. The time for various dissolved
48 polychlorinated biphenyls (PCBs) to reach equilibrium with a 50 μm or 500 μm PE sheet ranges
49 from days to decades [Lohmann and Muir, 2010; Endo *et al.*, 2013]. Thus the release of these
50 contaminants from the microplastic present in the remote ocean will be strongly retarded [Endo
51 *et al.*, 2013; Bakir *et al.*, 2014b] and only add a small contribution relative to already present
52 POPs at any given place and time. Results by Zarfl and Matthies [2010] also implied that
53 microplastics are not an efficient transport vector of HOCs in comparison to long range transport
54 by ocean or atmosphere, except for very high log K_{ow} chemicals, which have otherwise limited
55 transport potential in air and water.

56 Lastly, a comparison of microplastic particle density in the Pacific Ocean as detected by the Sea
57 Education Association (SEA; Lavender-Law, personal communication) and measured
58 concentrations of PCBs in surface seawater in the region [Zhang and Lohmann, 2010] found
59 little correlation between both.

60

61 **To what extent to microplastics contribute to the bioaccumulation and foodweb transfer of** 62 **POPs?**

63 There has been a long-standing assumption in many articles and studies that microplastics are
64 efficient carriers of organic pollutants into biota and the foodweb [Teuten *et al.*, 2009; Rochman
65 *et al.*, 2013; Chua *et al.*, 2014; Batel *et al.*, 2016; Wardrop *et al.*, 2016]. Such arguments have
66 been based on the notion that microplastics enrich various POPs (correct, see above), coupled
67 with the assumption that inside an animal, these pollutants are stripped off or leach out of the
68 microplastic and are taken up by the organism (e.g., [Teuten *et al.*, 2007; Bakir *et al.*, 2014a]). It
69 is worth recalling that chemicals diffuse to achieve the same chemical activity in the

70 environment, be that water, microplastic or biota [*Schwarzenbach et al.*, 2003]. Just because
71 microplastics display greater concentrations of POPs than present in water does not mean that
72 there is a greater tendency for these pollutants to diffuse out of the plastic particles. The potential
73 importance of microplastics as carriers of POPs into animals remains a strong theme in
74 discussion on microplastics, though, seemingly corroborated by empirical evidence [*Teuten et*
75 *al.*, 2009], it requires deeper examination. This will be addressed in more detail below using the
76 three scenarios outlined in Figure 1, in which (a) a naturally contaminated fish ingests a naturally
77 contaminated microplastic (e.g., both collected in the wild); (b) a clean fish consumes a
78 contaminated microplastic particle (e.g., in laboratory experiments), and (c) a reverse set-up,
79 where a contaminated fish consumes a pollutant-free microplastic particle. Examples from the
80 literature supporting these different scenarios are listed in Table 1. At a very basic level,
81 equilibrium partitioning thinking can be used to define simple expectations in which way organic
82 pollutants will move in a bioaccumulation thought experiment.

83

84 Scenario A

85 As outlined in scenario a) (Figure 1) a fish and piece of microplastic both contain POPs
86 already, simply from occurring in the environment. The presence of POPs in the fish, and a
87 microplastic residing in the same environment, are driven by the contaminants' chemical activity
88 and ought to be the same in both fish and microplastic particles. The ingestion of the
89 microplastic by the fish does not change the contaminant burden by the fish or the microplastic,
90 as they are both already in equilibrium [*Gouin et al.*, 2011]. This should be the most prevalent
91 interaction of biota, microplastics, and POPs in the natural environment, as animals are
92 constantly taking up POPs from the environment via their diet and respiration. There are neither

93 clean (i.e., POPs-free) oceans nor animals present, which means that they will bioaccumulate
94 POPs regardless of whether they ingest microplastics or not.

95 In the South Atlantic Ocean, there was generally no correlation between HOCs in microplastics
96 and amphipods [*Rochman et al.*, 2014]. Of the targeted HOCs (bisphenol A (BPA), alkylphenols,
97 alkylphenol ethoxylates, PCBs, and polybrominated diphenyl ethers (PBDEs)), only PBDEs
98 displayed increased body burdens in regions where more microplastics were present. The
99 presence of PBDEs in tissues could be due to the presence of small microplastic particles during
100 the extraction (see below). Another good example of scenario of (a) is the recent study in which
101 birds and the microplastics in their gut were analyzed for PCBs [*Herzke et al.*, 2016]. The
102 authors concluded that the presence of PCBs in the fulmars due to the ingestion of plastics was
103 negligible relative to the uptake of PCBs via their prey.

104 Similar conclusions were reached earlier by Gouin et al. (2011) based on theoretical
105 considerations based on a bioaccumulation food web model. Lastly, Koelmans et al. (2016) also
106 concluded that there is no experimental or theoretical evidence for an important role of
107 microplastics in the transfer of POPs into animals.

108

109 Scenario B

110 This scenario consists of exposing clean animals from a reference site to microplastics
111 containing a high concentration of POPs, either from laboratory dosing or from microplastics
112 exposed to contaminants at urban/industrialized sites. Most reported bioaccumulation studies
113 with microplastics are based on this scenario, such as the ingestion of PBDE-spiked particles by
114 amphipods in the laboratory [*Chua et al.*, 2014]. Interestingly, the experiment actually resulted in
115 decreased PBDE bioaccumulation uptake relative to control animals. Other experiments used

116 field-contaminated microplastic particles. For example, Teuten et al. (2009) described the
117 feeding of microplastics naturally contaminated by PCBs from Tokyo Bay to shearwater chicks
118 hidden in a fish diet. Initially, some uptake of lower chlorinated PCBs was observed, but the
119 PCBs ingested from the bird's prey fish outweighed the birds' body burden over time [*Teuten et*
120 *al.*, 2009].

121 In general, an efficient transfer of POPs from the microplastic to the animals is observed. This is
122 due to the experimental design, and shows that microplastic can be used as a vector for POPs
123 into animals. ***It does not demonstrate, however, that this pathway is relevant in the field.*** As
124 noted above, animals in the wild are typically as 'contaminated' with respect to POPs as the
125 microplastic particles they might consume. The conclusion that microplastics is not an important
126 transfer process was also reached in a study exposing lugworms to sediments enriched with field-
127 contaminated polystyrene particles [*Besseling et al.*, 2013]. The observed increase in PCB
128 bioaccumulation (1.1 – 1.5 times relative to controls) was only observed at low concentrations of
129 polystyrene particles.

130

131 Scenario C

132 This scenario is the reverse of scenario b) in which a POP-contaminated fish is fed clean
133 microplastic to determine if this will lower its body burden with respect to the POP. The idea is
134 based on research that olestra, a non-digestible fat, can be used to remove POPs from
135 contaminated animals [*Moser and McLachlan*, 1999]. Gouin et al. (2011) picked up this idea in
136 their bioaccumulation model, suggesting that the ingestion of clean microplastic could indeed
137 cause a decreased body burden in animals. Recently, Rummel et al. (2016), did not observe a

138 significant decrease in bioaccumulation of PCBs in rainbow trout allowed to ingest clean
139 microplastics.

140

141 **Do microplastics transfer other organic contaminants into biota?**

142 As discussed above, there is little evidence that microplastics play a major role in the
143 bioaccumulation of POPs, when compared to the role of diet in nature. As already discussed by
144 [Teuten *et al.*, 2009] and [Gouin *et al.*, 2011], microplastics could become an important pathway
145 for polymer additives that otherwise would not be easily transferred into the marine environment.
146 In particular, [Teuten *et al.*, 2009] suggested research should focus on the release of phenolic
147 additive-derived chemicals (i.e., alkylphenols and BPA) from microplastics in the food web. Yet
148 neither a modeling study by [Koelmans *et al.*, 2014] nor the field study by [Rochman *et al.*,
149 2014] found evidence that the ingestion of microplastics is relevant for the uptake of these
150 compounds by biota.

151 Several recent studies highlighted that certain chemicals, likely originating from plastic particles,
152 can indeed be transferred into animals. The presence of highly brominated BDEs 183 and 209 in
153 seabirds was linked to their ingestion of marine plastics [Tanaka *et al.*, 2013]. The birds' prey
154 items had no detectable BDE 183 and 109 concentrations but these contaminants were observed
155 in both the birds and ingested plastic debris particles. Similarly, the presence of
156 hexabromocyclododecanes (HBCDs) in Styrofoam and blue mussels from coastal South Korea
157 were linked [Jang *et al.*, 2016]. Elevated concentrations, and a α/γ HBCD ratio closer to that of
158 Styrofoam were detected in mussels colonizing Styrofoam buoys, when compared to mussels
159 collected from other substrates and regions along the coast. Overall, strong evidence was

160 presented for a direct pathway of HBCD from the Styrofoam buoy into the mussel, including the
161 detection of Styrofoam particles in the mussels themselves.

162 As there is good evidence that brominated compounds can be metabolized in animals [*Stapleton*
163 *et al.*, 2004], the presence of several low solubility brominated compounds (highly brominated
164 BDEs, HBCD) in biota, linked to the ingestion of microplastics, seems surprising at first. Yet it
165 might actually indicate that these compounds are not properly dissolved in the animals, but rather
166 part of nanoplastic particles dispersed within the animals' tissue and organs.

167

168 **Should microplastics be considered POPs?**

169 While the preceding discussion highlighted that microplastics in the oceans do little to affect the
170 presence and transfer of most organic pollutants at this point, there is still plenty of evidence that
171 microplastics are harmful and their impact should be minimized, as far as possible. This be could
172 be seen as a contribution towards a sustainable use of resources.

173 One approach would be to consider classifying microplastics as potential pollutants under the
174 Stockholm Convention on POPs [*UNEP*, 2001]. Four criteria, namely persistence,
175 bioaccumulation, long-range transport and adverse effects (Table 2), must be met for a
176 compound to be listed as a POP. There is strong evidence that microplastics are persistent, as a
177 result of their industrial polymer properties and additives [*Gewert et al.*, 2015], and that they
178 undergo long-range transport, as documented by their widespread presence in remote oceans
179 [*Lavender Law et al.*, 2010; *Sebille et al.*, 2015]. Several ecotoxicological studies highlight
180 adverse effects, though these experiments are often performed at unrealistically high doses of
181 microplastic exposure. The classical concept of bioaccumulation and biomagnification on a

182 molecular level is not met, but there is evidence that microplastics are present in top predators
183 and are transferred up the food chain.

184

185 In summary, there is little evidence that marine microplastics affect the global transport or
186 bioaccumulation of POPs in the oceans. In terms of bioaccumulation, experimental designs can
187 be manipulated to show that microplastics are a vector of POPs into organisms in the laboratory.
188 Yet, there is scant evidence from field studies that the ingestion of microplastics affects the
189 bioaccumulation of POPs. While there are some studies that show several low-solubility
190 compounds increase in animals that have ingested more microplastics, this might in fact be from
191 the presence of micro- and nanoplastic particles in those animals. Just because microplastics are
192 not relevant for the transport of POPs does not take away from their potential for detrimental
193 impacts on the environment. A possibility to address these concerns could be to consider marine
194 (micro)plastics as POPs, and rely on the Stockholm Convention to reduce their sources.

195

196 **References**

- 197 Adams, R. G., R. Lohmann, L. A. Fernandez, J. K. MacFarlane, and P. M. Gschwend (2007),
198 Polyethylene devices: Passive samplers for measuring dissolved hydrophobic organic
199 compounds in aquatic environments, *Environ. Sci. Technol.*, *41*(4), 1317–1323,
200 doi:10.1021/es0621593.
- 201 Bakir, A., S. J. Rowland, and R. C. Thompson (2014a), Enhanced desorption of persistent
202 organic pollutants from microplastics under simulated physiological conditions, *Environ.*
203 *Pollut.*, *185*, 16–23, doi:10.1016/j.envpol.2013.10.007.
- 204 Bakir, A., S. J. Rowland, and R. C. Thompson (2014b), Estuarine , Coastal and Shelf Science
205 Transport of persistent organic pollutants by microplastics in estuarine conditions, *Estuar.*
206 *Coast. Shelf Sci.*, *140*, 14–21, doi:10.1016/j.ecss.2014.01.004.
- 207 Batel, A., F. Linti, M. Scherer, L. Erdinger, and T. Braunbeck (2016), Transfer of
208 benzo[a]pyrene from microplastics to *Artemia Nauplii* and further to zebrafish via a trophic
209 food web experiment: CYP1A induction and visual tracking of persistent organic pollutants,
210 *Environ. Toxicol. Chem.*, *35*(7), 1656–1666, doi:10.1002/etc.3361.
- 211 Besseling, E., A. Wegner, E. M. Foekema, M. J. Van Den Heuvel-Greve, and A. A. Koelmans
212 (2013), Effects of Microplastic on Fitness and PCB Bioaccumulation by the Lugworm

213 Arenicola marina (L.), *Environ. Sci. Technol.*, *47*, 593–600.

214 Chua, E. M., J. Shimeta, D. Nugegoda, P. D. Morrison, and B. O. Clarke (2014), Assimilation of
 215 Polybrominated Diphenyl Ethers from Microplastics by the Marine Amphipod, *Allorchestes*
 216 *Compressa*, *Environ. Sci. Technol.*, *48*, 8127–8134.

217 Endo, S., M. Yuyama, and H. Takada (2013), Desorption kinetics of hydrophobic organic
 218 contaminants from marine plastic pellets, *Mar. Pollut. Bull.*, *74*(1), 125–131,
 219 doi:10.1016/j.marpolbul.2013.07.018.

220 Gewert, B., M. M. Plassmann, and M. Macleod (2015), Pathways for degradation of plastic
 221 polymers floating in the marine environment, *Environ. Sci. Process. Impacts*, *17*, 1513–
 222 1521, doi:10.1039/C5EM00207A.

223 Gouin, T., N. Roche, R. Lohmann, and G. Hodges (2011), A thermodynamic approach for
 224 assessing the environmental exposure of chemicals absorbed to microplastic., *Environ. Sci.*
 225 *Technol.*, *45*, 1466–72, doi:10.1021/es1032025.

226 Herzke, D., T. Anker-nilssen, T. H. Nøst, A. Go, S. Christensen-dalsgaard, M. Langset, K.
 227 Fangel, and A. A. Koelmans (2016), Negligible Impact of Ingested Microplastics on Tissue
 228 Concentrations of Persistent Organic Pollutants in Northern Fulmars off Coastal Norway, ,
 229 doi:10.1021/acs.est.5b04663.

230 Hirai, H. et al. (2011), Organic micropollutants in marine plastics debris from the open ocean
 231 and remote and urban beaches, *Mar. Pollut. Bull.*, *62*(8), 1683–1692,
 232 doi:10.1016/j.marpolbul.2011.06.004.

233 Jambeck, J. R., R. Geyer, C. Wilcox, T. R. Siegler, M. Perryman, A. Andrady, R. Narayan, and
 234 K. L. Law (2015), Plastic waste inputs from land into the ocean., *Science (80-.)*,
 235 doi:10.1126/science.1260352.

236 Jang, M., W. J. Shim, G. M. Han, M. Rani, Y. K. Song, and S. H. HONG (2016), Styrofoam
 237 debris as a source of hazardous additives for marine organisms, *Environ. Sci. Technol.*,
 238 acs.est.5b05485, doi:10.1021/acs.est.5b05485.

239 Karapanagioti, H. K., S. Endo, Y. Ogata, and H. Takada (2011), Diffuse pollution by persistent
 240 organic pollutants as measured in plastic pellets sampled from various beaches in Greece.,
 241 *Mar. Pollut. Bull.*, *62*, 312–317.

242 Koelmans, A. A., E. Besseling, and E. M. Foekema (2014), Leaching of plastic additives to
 243 marine organisms, *Environ. Pollut.*, *187*, 49–54, doi:10.1016/j.envpol.2013.12.013.

244 Koelmans, A. A., A. Bakir, G. A. Burton, and C. R. Janssen (2016), Microplastic as a Vector for
 245 Chemicals in the Aquatic Environment: Critical Review and Model-Supported
 246 Reinterpretation of Empirical Studies, *Env. Sci Technol*, *50*, 3315–3326,
 247 doi:10.1021/acs.est.5b06069.

248 Lavender Law, K., S. Morét-Ferguson, N. A. Maximenko, G. Proskurowski, E. E. Peacock, J.
 249 Hafner, and C. M. Reddy (2010), Plastic Accumulation in the North Atlantic Subtropical
 250 Gyre, *Science (80-.)*, *329*, 1185–1188, doi:10.1126/science.1192321.

251 Lohmann, R. (2012), Critical Review of Low-Density Polyethylene’s Partitioning and Diffusion
 252 Coefficients for Trace Organic Contaminants and Implications for Its Use As a Passive
 253 Sampler, *Environ. Sci. Technol.*, *46*, 606–618.

254 Lohmann, R., and D. Muir (2010), Global aquatic passive sampling (AQUA-GAPS): Using
 255 passive samplers to monitor POPs in the waters of the world, *Environ. Sci. Technol.*, *44*(3),
 256 860–864, doi:10.1021/es902379g.

257 Moser, G. A., and M. S. McLachlan (1999), A non-absorbable dietary fat substitute enhances
 258 elimination of persistent lipophilic contaminants in humans., *Chemosphere*, *39*(9), 1513–

259 1521.

260 Pintado-Herrera, M. G., P. A. Lara-Martín, E. González-Mazo, and I. J. Allan (2016),
 261 Determination of silicone rubber and low-density polyethylene diffusion and polymer/water
 262 partition coefficients for emerging contaminants, *Environ. Toxicol. Chem.*, *35*(9), 2162–
 263 2172, doi:10.1002/etc.3390.

264 Rochman, C. M., E. Hoh, B. T. Hentschel, and S. Kaye (2012), Long-Term Field Measurement
 265 of Sorption of Organic Contaminants to Five Types of Plastic Pellets: Implications for
 266 Plastic Marine Debris,

267 Rochman, C. M., E. Hoh, T. Kurobe, and S. J. Teh (2013), Ingested plastic transfers hazardous
 268 chemicals to fish and induces hepatic stress, *Sci. Rep.*, *3*, 3263–3269,
 269 doi:10.1038/srep03263.

270 Rochman, C. M., R. L. Lewison, M. Eriksen, H. Allen, A. Cook, and S. J. Teh (2014),
 271 Polybrominated diphenyl ethers (PBDEs) in fish tissue may be an indicator of plastic
 272 contamination in marine habitats, *Sci. Total Environ.*, *476–477*, 622–633,
 273 doi:10.1016/j.scitotenv.2014.01.058.

274 Rusina, T. P., F. Smedes, J. Klanova, K. Booij, and I. Holoubek (2007), Polymer selection for
 275 passive sampling : A comparison of critical properties, *Chemosphere*, *68*, 1344–1351,
 276 doi:10.1016/j.chemosphere.2007.01.025.

277 Schwarzenbach, R. P., P. M. Gschwend, and D. M. Imboden (2003), *Environmental Organic*
 278 *Chemistry*, 2nd ed., Wiley-Interscience, New York.

279 Sebille, E. Van, C. Wilcox, L. Lebreton, N. Maximenko, B. D. Hardesty, J. Franeker, M.
 280 Eriksen, D. Siegel, F. Galgani, and K. L. Law (2015), A global inventory of small floating
 281 plastic debris, *Environ. Res. Lett.*, *10*(12), 124006, doi:10.1088/1748-9326/10/12/124006.

282 Smedes, F., R. W. Geertsma, T. Von der Zande, and K. Booij (2009), Polymer - Water Partition
 283 Coefficients of Hydrophobic Compounds for Passive Sampling : Application of Cosolvent
 284 Models for Validation, *Environ. Sci. Technol.*, *43*(18), 7047–7054.

285 Stapleton, H. M., R. J. Letcher, and Bak (2004), Debromination of Polybrominated Diphenyl
 286 Ether Congeners BDE 99 and BDE 183 in the Intestinal Tract of the Common Carp (
 287 *yprius carpio*), *Env. Sci Technol*, *38*(4), 1054–1061.

288 Tanaka, K., H. Takada, R. Yamashita, K. Mizukawa, and M. Fukuwaka (2013), Accumulation of
 289 plastic-derived chemicals in tissues of seabirds ingesting marine plastics, *Mar. Pollut. Bull.*,
 290 *69*(1–2), 219–222, doi:10.1016/j.marpolbul.2012.12.010.

291 Teuten, E. L., S. J. Rowland, T. S. Galloway, and R. C. Thompson (2007), Potential for Plastics
 292 to Transport Hydrophobic Contaminants, *Environ. Sci. Technol.*, *41*, 7759–7764.

293 Teuten, E. L. et al. (2009), Transport and release of chemicals from plastics to the environment
 294 and to wildlife, *Philos. Trans. R. Soc. B Biol. Sci.*, *364*, 2027–2045,
 295 doi:10.1098/rstb.2008.0284.

296 Thompson, R. C., Y. Olsen, R. P. Mitchell, A. Davis, S. J. Rowland, A. W. G. John, D.
 297 McGonigle, and A. E. Russell (2004), Lost at Sea: Where Is All the Plastic?, *Science* (80-
 298), *304*(5672), 838.

299 UNEP (2001), *Final act of the plenipotentiaries on the Stockholm Convention on persistent*
 300 *organic pollutants.*, Geneva, Switzerland.

301 Wardrop, P., J. Shimeta, D. Nugegoda, P. D. Morrison, A. Miranda, M. Tang, and B. O. Clarke
 302 (2016), Chemical Pollutants Sorbed to Ingested Microbeads from Personal Care Products
 303 Accumulate in Fish, *Env. Sci Technol*, *50*, 4037–4044, doi:10.1021/acs.est.5b06280.

304 Zarfl, C., and M. Matthies (2010), Are marine plastic particles transport vectors for organic

305 pollutants to the Arctic ?, *Mar. Pollut. Bull.*, doi:10.1016/j.marpolbul.2010.05.026.
306 Zhang, L., and R. Lohmann (2010), Cycling of PCBs and HCB in the surface ocean-lower
307 atmosphere of the open pacific, *Environ. Sci. Technol.*, 44(10), 3832–3838,
308 doi:10.1021/es9039852.
309 Ziccardi, L. M., A. Edgington, K. Hentz, K. J. Kulacki, and S. Kane Driscoll (2016),
310 Microplastics as vectors for bioaccumulation of hydrophobic organic contaminants in the
311 marine environment: A state-of-the-science review, *Environ. Toxicol. Chem.*, 35(7), 1667–
312 1676, doi:10.1002/etc.3461.
313

314 **List of Tables and Figures**

- 315 - **Figure 1 with bioaccumulation set-up**
- 316 - **Table 1: Bioaccumulation studies involving microplastics and POPs**
317
- 318 - **Table 2: Criteria for inclusion of compounds to the Stockholm Convention on POPs**
- 319
- 320