



# MicroRNA dysregulation in spinal cord injury: causes, consequences, and therapeutics

Manuel Nieto-Díaz<sup>1</sup>, Francisco J. Esteban<sup>2</sup>, David Reigada<sup>1</sup>, Teresa Muñoz-Galdeano<sup>1</sup>, Mónica Yunta<sup>1,3</sup>, Marcos Caballero-López<sup>1</sup>, Rosa Navarro-Ruiz<sup>1</sup>, Ángela del Águila<sup>1</sup> and Rodrigo M. Maza<sup>1\*</sup>

<sup>1</sup> Molecular Neuroprotection Group, Experimental Neurology Unit, Hospital Nacional de Paraplégicos (Servicio de Salud de Castilla-La Mancha), Toledo, Spain

<sup>2</sup> Departamento de Biología Experimental, Facultad de Ciencias Experimentales y de la Salud, Universidad de Jaén, Jaén, Spain

<sup>3</sup> Unidad de Patología Mitocondrial, Unidad Funcional de Investigación en Enfermedades Crónicas, Instituto de Salud Carlos III, Madrid, Spain

## Edited by:

Tommaso Pizzorusso, Università degli Studi di Firenze, Italy

## Reviewed by:

Simone Di Giovanni, University of Tübingen, Germany

Ulkan Kilic, Bezmialem Vakif University, Turkey

## \*Correspondence:

Rodrigo M. Maza, Molecular Neuroprotection Group, Experimental Neurology Unit, Hospital Nacional de Paraplégicos (Servicio de Salud de Castilla-La Mancha), Finca de la Peraleda, s/n, Toledo 45071, Spain  
e-mail: rodrigom@sescam.jccm.es

Trauma to the spinal cord causes permanent disability to more than 180,000 people every year worldwide. The initial mechanical damage triggers a complex set of secondary events involving the neural, vascular, and immune systems that largely determine the functional outcome of the spinal cord injury (SCI). Cellular and biochemical mechanisms responsible for this secondary injury largely depend on activation and inactivation of specific gene programs. Recent studies indicate that microRNAs function as gene expression switches in key processes of the SCI. Microarray data from rodent contusion models reveal that SCI induces changes in the global microRNA expression patterns. Variations in microRNA abundance largely result from alterations in the expression of the cells at the damaged spinal cord. However, microRNA expression levels after SCI are also influenced by the infiltration of immune cells to the injury site and the death and migration of specific neural cells after injury. Evidences on the role of microRNAs in the SCI pathophysiology have come from different sources. Bioinformatic analysis of microarray data has been used to identify specific variations in microRNA expression underlying transcriptional changes in target genes, which are involved in key processes in the SCI. Direct evidences on the role of microRNAs in SCI are scarcer, although recent studies have identified several microRNAs (miR-21, miR-486, miR-20) involved in key mechanisms of the SCI such as cell death or astrogliosis, among others. From a clinical perspective, different evidences make clear that microRNAs can be potent therapeutic tools to manipulate cell state and molecular processes in order to enhance functional recovery. The present article reviews the actual knowledge on how injury affects microRNA expression and the meaning of these changes in the SCI pathophysiology, to finally explore the clinical potential of microRNAs in the SCI.

**Keywords:** spinal cord injury, microRNA, nervous system, cell death, inflammation, astrogliosis, therapeutics

## INTRODUCTION

Spinal cord injury (SCI) is a discapacitating pathology with a global-incident rate estimated at 23 new traumatic SCI cases per million in 2007 (180,000 new cases/year; Lee et al., 2013). Recovery, even partial, from SCI was considered unachievable until the early 1980s when David and Aguayo (1981) demonstrated that spinal axons could regenerate under appropriate conditions. This seminal work initiated a burst of research aimed to understand the pathophysiology of the SCI and the development of therapeutic tools. The capacity of microRNAs to regulate cell state and function through post-transcriptionally silencing hundreds of genes are being acknowledged as an important actor in the pathophysiology of SCI. In this article, we review the known and potential roles of microRNAs in SCI and their possible therapeutic application. The article begins with an overview of the pathophysiology of SCI and the gene programs involved, followed by a general picture of microRNA biogenesis, function, and regulation, particularly in the central nervous system (CNS). Afterward, we discuss the observed global changes in microRNA expression following SCI and the role of mechanisms regulating microRNA biogenesis on

SCI. We next explore the contribution of microRNA in the regulation of key processes of SCI such as inflammation, cell death, regeneration, or gliosis. The review ends with a brief perspective on the feasibility of microRNA-based therapeutics in the treatment of SCI.

## AN OVERVIEW OF THE PATHOPHYSIOLOGY OF THE SPINAL CORD INJURY

The term spinal cord injury refers to the damage to the spinal cord caused by trauma or disease but also to the physiological response that develops after injury and that largely determines the functional deficits that will face the injured person. Although most SCI are traumatic, inflicted by mechanical causes due to accidents or violence, there are also a number of non-traumatic SCIs, mainly due to discopathies and tumors (Biering-Sorensen et al., 1990). The pathophysiology of the SCI is a complex intermingled set of events, responses, mechanisms, and processes, affecting the nervous, vascular, and immune systems, that develop during the months following the initial damage. Most participating cells reside in the spinal cord, but others are summoned to the

injury site from the circulatory system. A brief description of the pathophysiology is provided in the following paragraphs. Detailed descriptions can be obtained in different reviews (Mautés et al., 2000; Dumont et al., 2001; Profyris et al., 2004; Liverman et al., 2005; Rowland et al., 2008; Oyinbo, 2011).

In the traumatic SCI, the mechanical trauma causes the compression, stretching, laceration, or transection of the spinal cord. The most common form of SCI is a compressive–contusive-type injury in which displaced components of the vertebral column, exert force on the cord causing both immediate traumatic injury and often sustained compression (Rowland et al., 2008). Once compression and contusion surpass structural thresholds, physical and biochemical alterations of the cells induce a cascade of systemic and local events that constitute the primary damage (Oyinbo, 2011). Local events include axon severing, membrane rupture and death of neurons, glia and endothelial cells. Surviving neurons at the injury site respond firing action potentials that shift the local levels of ions together with the ions released due to membrane shear. The resulting ion concentration reaches toxic levels that kill the nearby neurons. The barrage of action potentials also causes the release and accumulation of neurotransmitters that will cause further neuron and glial cell death by excitotoxicity. Mechanical trauma causes intraparenchymal hemorrhage (mainly in the small vessels of the gray substance) and, consequently, the disruption of the blood–spinal cord barrier together with edema and swelling at the spinal cord (Mautés et al., 2000). Vasospasm and thrombosis in the superficial vessels accompany hemorrhage causing hypoxia, ischemia, and increasing neural cell death. At a systemic level, primary damage causes a transient increase in systemic blood pressure that is followed by a prolonged hypotension (either hemorrhagic or neurogenic) causing further oxygen deprivation to the spinal cord. Hypoxia – together with ion shifts inside and outside the neuron – seems to cause a temporal switch off of the spinal cord function at and below the injury site known as spinal shock.

In the minutes to months that follow the initial damage, the secondary phase of the SCI takes place. This secondary phase comprises several interrelated damage processes including vascular alterations, biochemical disturbances and cellular responses that lead to an inflammatory response and cell death that significantly expand the area of damage. Vascular alterations resulting from hemorrhage and ischemia are central constituents of the secondary injury cascade. Reduced perfusion of the spinal cord due to vasospasm and hypotension is followed by a period of reperfusion, which increases the production of oxygen- and nitrogen-derived free radicals [superoxide, hydroxyl radicals, nitric oxide (NO), peroxynitrite] already being produced during the period of ischemia (Dumont et al., 2001). All these species contribute to oxidative stress and exacerbates damage and cell death. Alterations in the vascular system also include the disruption of the blood–spinal cord barrier that extends far beyond the injury site for days and even weeks after injury. Release of cytokines [interleukin (IL)-1 $\beta$ , tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )], matrix metalloproteinases, reactive oxygen species (ROS), etc., contribute to enhance vascular permeability (Mautés et al., 2000; Donnelly and Popovich, 2008) and – together with upregulation of cell adhesion molecules (CAMs and selectins) by endothelial cells (Mautés et al., 2000;

Profyris et al., 2004; Donnelly and Popovich, 2008) – participate in the recruitment and infiltration of immune cells to the injured spinal cord.

Immune cells develop a key role in the pathophysiology of SCI. Neutrophils arrive to the injury site in the first hours after injury [peaking at 1 day post-injury (dpi) in rats and 1–3 dpi in humans, see Fleming et al., 2006] and disappear during the first week, although some evidences indicate a continued presence for long times (180 dpi in rats, according to Beck et al., 2010). Neutrophils remove debris, but mainly release assortments of proteins, including proteolytic and oxidative enzymes that “sterilize” the area but also contribute to extend tissue damage (Taoka et al., 1997). Neutrophils also release signaling proteins that attract macrophages. Macrophages resulting from the activation of spinal cord microglia or from blood monocytes infiltrate the injury in the first days after the injury, presenting a peak during the first week and persisting for months (Fleming et al., 2006). Microglial activation is triggered early after injury and induces a morphological and functional change in the phenotype of this cell, from a resting, ramified phenotype to a phagocytosis-capable, “macrophage-like” phenotype (Byrnes et al., 2006). Macrophages remove debris and dead cells, present antigens, and release pro-inflammatory and protective cytokines, ROS, NO, and proteases (Fleming et al., 2006). T lymphocytes enter the injured spinal cord mainly 1 week after injury. T cells are responsible for cell-mediated adaptive immunity, although their role in SCI remains controversial (Fleming et al., 2006). In rat models, it seems that immune cells tend to maintain or reduce their presence after this first burst of immune response following SCI. However, a recent study in rats demonstrates that immune cells present a time-dependent multiphasic response, with a late phase that mainly involves a peak of macrophages at 60 dpi (Beck et al., 2010). Contrary to the mixed beneficial and detrimental effects on the immune response in the initial phase, this late phase seems to be mainly beneficial and its blocking causes further functional deficits (Beck et al., 2010).

All previous events have strong effects on neural cells. Necrotic cell death initiated by the mechanical trauma spreads during the secondary phase due to excitotoxicity and the accumulation of free radicals (ROS and RNS) released by immune cells or during reperfusion. Free radicals cause lipid peroxidation as well as oxidative and nitrate damage to lipids, proteins, and nucleic acids, inducing the lysis of the cell membrane, altering the cytoskeleton and the organelles, and ultimately causing the death of neural cells (Oyinbo, 2011). Apoptosis and other forms of programmed cell death are also important actors in secondary damage after SCI. Programmed cell death seems to occur in at least two phases: an initial phase, in which apoptosis accompanies necrosis and a later phase, which is predominantly confined to white matter and that affects oligodendrocytes and microglia (Profyris et al., 2004). Calcium influx and possibly signaling through Fas/CD95 pathway are among the triggers proposed for programmed cell death although other mechanisms may be also acting, including loss of trophic support (Liverman et al., 2005; Rowland et al., 2008).

Apoptosis of oligodendrocytes results in extended demyelination, the loss of the oligodendrocyte myelin sheath that insulate nerve axons and permit effective nervous signal conduction. As a

consequence, axons crossing the injured segment/s but deprived from myelin sheath and experiencing alterations in the ion channels become unable to transmit signals to the brain and the body, even though they remain intact. Axotomy (axon sectioning) is also a major factor in SCI. Depending on aspects such as distance of axotomy to cell body, trophic support or neuronal type, the fate of axotomized neurons will differ strongly. A significant proportion of neurons, being severed close to the cell body or lacking enough trophic support, die after axotomy. Others retract the severed axon and develop a terminal retraction bulb. Conversely, some surviving neurons develop a regenerating response. These neurons experience hypertrophy of the cell body due to the increased synthesis of proteins that will be used in an attempt to regenerate the severed axon. Within this pro-regenerative response, intact neurons can also develop sprouts (collateral axonal branches) that may rewire unconnected targets (sometimes with undesired side effects). However, axon growth is strongly if not completely inhibited due to several inhibitors present at the injured spinal cord. Much of this inhibition is due to molecules associated to the oligodendrocyte myelin, such as NOGO, and other are consequence of the changes experienced by the astrocytes following injury. Astrocyte reactivity after SCI involves cellular changes characterized by an initial hypertrophic and a late hyperplastic response (Fawcett and Asher, 1999; Sahni et al., 2010). During the hypertrophic response, reactive astrocytes present a characteristic enlarged somata with thickened processes and upregulation of intermediate filament proteins, such as glial fibrillary acidic protein (GFAP) and vimentin (Pekny and Pekna, 2004). This initial hypertrophic response protects neural cells by up-taking the potentially excitotoxic glutamate, producing oxidative stress scavengers and repairing the blood-spinal cord barrier (Faulkner et al., 2004; Sofroniew, 2005). In the following hyperplastic phase during the subacute and chronic phases of the SCI, newly proliferated astrocytes with thinner processes forms a dense scar in the damaged area that inhibits of axon regeneration (Fawcett and Asher, 1999; Silver and Miller, 2004).

In the months and years that follow the damage, the SCI becomes chronic. Cell death, scarring, gliosis, and other local alterations of the tissue lead to the formation of cavities filled of fluid and surrounded by glial scar that can extend for several segments above and below the injury site (Liverman et al., 2005). Myelin loss and alterations in the functioning of the ion channels can lead to changes in the surviving neurons and dependent networks leading to chronic neuropathic pain and/or spasticity.

### GENE EXPRESSION CHANGES AFTER SPINAL CORD INJURY

Most cell functions, responses, and phenotypic changes depend on the activation and/or suppression of a large number of transcriptional pathways (Di Giovanni et al., 2003). Microarray analyses in SCI models have identified gene expression changes that, to some extent, can be observed across different studies and even across different strains or species (Di Giovanni et al., 2003; Velardo et al., 2004). Prominent expression changes in the spinal cord following injury comprise clustered expression changes in genes associated to:

### TRANSCRIPTION AND STRESS RESPONSE

Clusters including factors such as *NF- $\kappa$ B* (nuclear factor kappa-light-chain-enhancer of activated B cells), *c-fos*, *HSP-70* (70 kD heat shock protein) become upregulated early after injury and continue for at least 24 h (Song et al., 2001). Activation of transcription factors such as *NF- $\kappa$ B* or interferon regulatory factor (IRF)-1 mediates the increased expression of pro-inflammatory genes and modulates pivotal processes such as apoptosis or regeneration (Song et al., 2001). On the other hand, there is an early increase in chaperone expression (*HSP-70*) together with metallothioneins 1 and 2 (Aimone et al., 2004) that may protect cells from protein damage or oxidative stress. Further anti-oxidative stress genes become upregulated at much later times, including superoxide dismutases (SODs) 1 and 2, *catalase*, and *glutathione peroxidase* (GPX; Aimone et al., 2004).

### INFLAMMATORY RESPONSE

Upregulation of pro-inflammatory genes, including cyclooxygenase (COX)-2, interleukins *IL-1 $\beta$*  and *IL-6*, *TNF- $\alpha$* , is observed early after injury and persists during the first weeks to go back to normal levels at 14 days (Carmel et al., 2001; Song et al., 2001; Di Giovanni et al., 2003; Zhang et al., 2004). Chemokines involved in immune cell attraction [monocyte chemoattractant protein (MCP)-1, macrophage inflammatory protein (MIP)-1 $\beta$ ] and adhesion molecules participating in cell infiltration (integrins, intercellular and vascular CAMs, cadherins, and selectins) are also upregulated during the first hours after injury (Aimone et al., 2004). Changes in the expression of inflammatory genes take place in different spinal cord cells but has been particularly studied in microglia. According to a profiling study developed by Byrnes et al. (2006), activated microglia first express (peak at 4–24 h) pro-inflammatory molecules, including *IL-1 $\beta$* , *IL-6*, *CCL2* [chemokine (C–C motif) ligand 2]/MCP-1, *CXCL2* [chemokine (C–X–C motif) ligand 2]/MIP2 $\alpha$ , involved in the recruitment of immune cells to the damaged area. A second pulse of change in microglial gene expression occurs later (3–7 dpi in rats) and involves genes coding for cytochrome b-245 light chain protein (*CYBa*), cathepsin Y, Galectin-3, microglial response factor (MRF)-1, P38, cyclin D1, caspase 1, and leukocyte surface antigen CD53/OX44, that participate in the regulation of the immune response, phagocytosis, production of ROS, proliferation, and cell death. Other immune response genes related to phagocytosis including the classical complement pathway and the FC receptors show a persistent upregulation after injury (Aimone et al., 2004).

### NEURON-ASSOCIATED GENES

A large cluster of genes coding for proteins involved in the potassium, calcium, and sodium pumps and channels as well as in synapsis, cell excitability, and neurotransmission show a significant decrease during the first week (Carmel et al., 2001; Velardo et al., 2004; De Biase et al., 2005; Wu et al., 2005). This decrease can reflect changes in the gene profile of the neurons but it may also reflect the advance of neuronal cell death that takes place after injury (Carmel et al., 2001; De Biase et al., 2005). Attempts of axonal regeneration in the weeks following injury are also accompanied by expression changes in a large group of genes that includes the overexpression of several plasticity and

regeneration-associated proteins (*Ninjurin*, *Coronin 1b*, *Rab13*, *Growth Associated Protein-43*, *Neuritin*, *Ankyrin*, *Myelin oligodendrocyte glycoprotein*, and *cAMP-related genes*; Carmel et al., 2001; Song et al., 2001; Di Giovanni et al., 2003, 2005b).

### CELL CYCLE AND CELL DEATH

Changes in the expression of cell cycle genes has been detected 24 h after injury, including upregulation of *c-myc*, *pcdna*, *gadd45a*, and cyclins (Di Giovanni et al., 2003). Activation of cell cycle genes may induce apoptosis in post-mitotic cells and, thus underlie post-SCI apoptosis of neurons. They may also be involved in astrocytic proliferation during the glial scar formation (Di Giovanni et al., 2003). Pro-apoptotic and anti-apoptotic genes also show significant expression changes, including upregulation of caspase-3, *Bax*, *Bak-1* in the first week after SCI and the later upregulation of protective *PI3K* and *Stat3* and downregulation of pro-apoptotic *GSK-3* (Carmel et al., 2001; Aimone et al., 2004). In addition, it has been observed the upregulation at 24 h of genes coding for different growth factors [transforming growth factor- $\beta$  (TGF- $\beta$ ), platelet-derived growth factor, vascular endothelial growth factor] and anti-apoptotic proteins (survival of motor neurons proteins), which may contribute to prevent neural cell death (Song et al., 2001).

### VASCULAR SYSTEM REGULATION AND ANGIOGENESIS

Hemorrhage and other early vascular events are reflected in the expression profile. Blood coagulation genes – platelet factor 4 (CXCL4), coagulation factors VIII, protein C, etc., – appear over-expressed in the first 24 h after injury and some remain for several weeks (Chamankhah et al., 2013). Genes such as angiopoietin are increased in the injury area in the week that follows the damage (Aimone et al., 2004). Other dysregulated genes related to vascular events include HIG (hypoxia-induced gene), which is downregulated late after injury (Aimone et al., 2004).

### CHANGES IN GLIAL CELLS

Although less explored in global profile studies, there are important changes in gene expression in the glial cells related to astrocyte reactivity and the formation of the glial scar, as well as to the proliferation and remyelination attempts of oligodendrocytes. Among the best-characterized genes related to astrocyte reactivity and glial scar formation are those coding for GFAP, vimentin, and nestin, intermediate filaments highly overexpressed in reactive astrocytes. These genes are markedly upregulated during the first week after injury (Carmel et al., 2001; Wu et al., 2005). On the other hand, oligodendrocyte profiles are characterized by a decrease in cell specific genes due to the advance of oligodendrocyte death during the first weeks followed by an increase in proliferation and myelination genes (Wu et al., 2005).

### MicroRNA BIOGENESIS, FUNCTION, AND REGULATION

As we have shown in the previous sections, SCI causes profound cellular changes that result from dysregulation of signaling pathways and structural proteins. Alteration of gene expression following SCI is likely accompanied by the post-transcriptional regulation of these modified gene networks. Among the known post-transcriptional regulators, microRNAs

have recently attracted much attention due to their ability to inhibit mRNA translation.

MicroRNAs were first identified in *C. elegans* in 1993 (Lee et al., 1993). A small non-coding RNA (*lin-4*) was shown to regulate translation of *lin-14* through RNA–RNA interaction. MicroRNAs constitute an abundant class of highly conserved small non-coding RNA molecules composed of 20–24 nucleotides in length, that post-transcriptionally regulate gene expression. More than 2500 mature forms of microRNA sequences have been identified in humans (miRBase; Kozomara and Griffiths-Jones, 2011). MicroRNAs are transcribed from genomic DNA by RNA polymerase II or III in the form of large primary transcripts (pri-miRNAs) with functional secondary structures of stem-loop hairpins. The stem-loops structures are recognized and cleaved by the complex formed by the RNase III Droscha and the RNA-binding protein DGCR8, which leads to liberation of a precursor microRNA (pre-miRNA). Once processed, exportin 5 transports the pre-miRNAs from the nucleus to the cytosol, where an enzymatic complex containing Dicer process them to yield the 20- to 25-nucleotide duplex mature miRNAs. One strand (passenger) of the duplex is degraded and the other strand (guide) is integrated into a RNA-induced silencing complex (RISC), which facilitates the binding of the microRNAs to their target mRNA.

The predominant role of microRNAs in RISCs is to regulate post-transcriptionally the expression of their target genes by translational repression, mRNA cleavage, or mRNA decay. The employed mechanism depends on the degree of complementarity between the microRNAs and their target mRNAs. When microRNAs perfectly or near-perfectly pairs the targeted mRNAs, as occurs in plants, cleavage of the mRNA takes place. On the contrary, when microRNAs imperfectly pair to their target mRNAs, as usually occurs in animals, translational repression or decay (due to microRNA-driven deadenylation) become the mechanisms mediating gene regulation. In animals, microRNAs have been proposed to downregulate gene expression mostly through the translational repression (Krichevsky, 2007), although a recent study showed that destabilization (and degradation) of target mRNAs is the major mechanism of miRNA gene repression in mammals (Guo et al., 2010).

Since microRNAs do not require perfect complementarity for target recognition in animals, a single microRNA can regulate multiple, even hundreds mRNAs (Lim et al., 2005; Baek et al., 2008; Selbach et al., 2008). At the same time, each mRNA can be regulated by many microRNAs (Krek et al., 2005; Krichevsky, 2007) and a given microRNA may have multiple binding sites in the same mRNA, thereby enhancing its overall effect (Bhalala et al., 2013). Computational predictions suggest that the more than 2000 known human microRNAs are able to repress the expression of at least a 20–30% of all protein-coding genes (Krichevsky, 2007) and up to the 60% according to Friedman et al. (2009) and Sayed and Abdellatif (2011). Besides a single miRNA can tune protein synthesis from thousands of genes, quantitative proteomics studies have found that the magnitude of the change is small under physiological conditions (less than fourfold; Guo et al., 2010). In fact, blocking microRNA biogenesis through DICER inhibition has modest effects on cell differentiation and organism patterning, although opposite results have also been reported (Kosik, 2010).

MicroRNAs are common components of regulatory pathways, and in many cases constitute molecular on–off switches in establishing cell fate and identity both under physiological and pathological conditions. The power of this regulatory mechanism lies in the unique ability of microRNAs to guide processes and cellular functions through precise titration of gene dosage, and the ability of a single microRNA to control the levels of a large cohort of gene products. Because microRNAs target multiple mRNAs, they can exert distributed control over broad target fields of functionally related mRNAs as opposed to focusing their control on a small number of genes in a “final common pathway.” The action of an individual microRNA can lead to a cumulative reduction in expression of multiple components of one specific functional network, and several microRNAs may cooperatively target various mRNAs whose protein products are part of the same molecular pathway (Krek et al., 2005; Krol et al., 2010; Sluijter, 2013). MicroRNAs are often physically clustered in the genome, and these sets of microRNAs may target mRNAs with related biological functions at short distances in their protein–protein interaction map (Kim et al., 2009). Coordinated microRNA targeting of closely connected genes seems to be prevalent across pathways (Tsang et al., 2010). Thus, microRNAs provide broad and robust transcriptional regulation that can be governed either by individual microRNAs or by the combined action of multiple microRNAs.

MicroRNA networks are often specialized for specific cell types and there is a strong correlation between cell identity and patterns of microRNA expression (Kosik, 2010). The anticorrelated expression of microRNAs and their target mRNAs in developmental transitions and the mutually exclusive expression of target genes and microRNAs in neighboring tissues argues that microRNAs confer accuracy to developmental gene expression programs, thus ensuring tissue identity and supporting cell-lineage decisions, and reflect the basic role of miRNAs in establishing cell identity during development (Ebert and Sharp, 2012). MicroRNAs also serve as a buffer to assist cells in coping with environmental contingencies (Kosik, 2010). As markers of cell identity, miRNAs encode a representation of multiple cell states that all correspond to a single identity. That is, many different states comprise a single identity because cells must retain their identities in the face of both environmental changes and internal noise that can result in large variations in molecular composition. MicroRNAs are good candidates for setting boundary conditions upon coding transcripts to restrict protein levels within a range of values that maintain cell identity in the face of homeostatic compensatory changes. The RISC allows both the constitutive maintenance of cell identity by silencing mRNAs that are not part of the specialized cell’s repertoire as well as the holding of mRNAs of an alternative identity in reserve (Lim et al., 2005). The environment that cells face is many times more complex than the biological adaptations available within the genome. Among the adaptive responses of cells to an environmental contingency is the up- or downregulation of proteins. The properties of miRNAs to adjust protein levels, their dispensability under basal conditions, their conservation, as well as the ease with which new miRNAs appear over evolutionary time all suggest that they are suited for environmental contingencies (Kosik, 2010). Wu et al. (2009)

have proposed that miRNAs keep the system close to the mean and set expression boundaries of transcription factors, which are otherwise noisy.

### MicroRNA IN THE NORMAL AND PATHOLOGICAL CENTRAL NERVOUS SYSTEM

MicroRNAs are highly expressed in the mammalian CNS, including the spinal cord (Miska et al., 2004; Kosik, 2006; Krichevsky, 2007; Bak et al., 2008). Their expression in the spinal cord seems to be specific and preserved through vertebrate evolution (Yunta et al., 2012). Moreover, experimental data reveal that some miRNAs are cell-type specific, such as miR-124 and miR-128, which are preferentially expressed in neurons, or miR-23 or miR-219, which are restricted to astrocytes and oligodendrocytes, respectively (Sempere et al., 2004; Smirnova et al., 2005; Lau et al., 2008). MicroRNAs serve essential roles in virtually every aspect of CNS function, including neurogenesis, neural development, and cellular responses leading to changes in synaptic plasticity (Krichevsky et al., 2003; Miska et al., 2004; Sempere et al., 2004; Stefani and Slack, 2008; Gangaraju and Lin, 2009; Li and Jin, 2010; Smith et al., 2010; Cochella and Hobert, 2012; Goldie and Cairns, 2012). For example, experimental overexpression or inhibition of miR-124 have demonstrated its key role in neuronal differentiation (Krichevsky et al., 2006; Makeyev et al., 2007; Visvanathan et al., 2007), whereas let-7b regulates neural stem cell proliferation and differentiation by targeting the stem cell regulator TLX and the cell cycle regulator cyclin D1 (Zhao et al., 2010a). MicroRNAs are also involved in the specification of glia. They have been shown to be critical regulators of oligodendrocyte differentiation and myelination in the vertebrate CNS, in particular miR-219 and miR-338 (Zhao et al., 2010b).

In addition to their role in the development and the functioning in the normal CNS, numerous evidences indicate that microRNA dysregulation is implicated in a wide range of neurological diseases. Several studies indicate that microRNA dysregulation can be associated to neurodegeneration, as in Alzheimer’s disease where the downregulation of the miR-29 cluster has been proposed to contribute to increase BACE1 and amyloid-beta levels and thus to the development of the disease (see Saugstad, 2010). The role of microRNAs in neurodegeneration extends to other pathologies, such as Huntington and Parkinson diseases, or even to prion diseases (Saugstad, 2010). The evidences also suggest that microRNAs participate in psychiatric disorders, such as Schizophrenia, Tourette’s syndrome, and bipolar disorder, as well as in developmental disorders, such as the Rett and fragile X syndromes (Kosik, 2006; Meza-Sosa et al., 2012). MicroRNAs also participate in the profound cellular changes that occur in the damaged CNS, where they play an active role in the regulation of typical features of CNS injuries, such as inflammation, apoptosis, cell proliferation, and differentiation (see this review for SCI or Bhalala et al., 2013 for a more general approach).

### GLOBAL CHANGES IN microRNA EXPRESSION AND REGULATION OF microRNA BIOGENESIS IN SPINAL CORD INJURY

Several studies have analyzed microRNA expression and function in SCI (34 published paper in August, 2013 according to PubMed).

Seven of these studies include global analyses of microRNA expression based on microarrays. As shown in **Table 1**, the results from these studies strongly vary in the resulting overall patterns, both in the number (from as few as 10 significant expression changes to more than 250) and profile of changing microRNAs. Much of this variability can be ascribed to differences in SCI model, from ischemia-reperfusion to transection, compression, or contusion, to differences in the sampling time after injury, from 4 h to 14 days after injury, or to differences in the animal model (rats and mice of different strains). Besides, some results from comparable injury models and times still show strong differences. Technical differences among platforms, laboratories, sampling and analytical procedures, etc., may account for variability in microarray data (Git et al., 2010; Callari et al., 2012; Saugstad, 2013). In agreement, we have observed that the number of significant changes

identified at 7 dpi vary from 257 to 2 depending on whether parametric (Student's *t*-test) or non-parametric (rank product test) tests are used (Yunta et al., 2012). However, discrepancies may also reveal variability in microRNA expression associated to differences in injury features. Injury severity is one of such factors. It is well known that severity determines several aspects of the SCI pathophysiology such as the inflammatory response or the nerve fiber preservation/destruction (Fehlings and Tator, 1995; Yang et al., 2005). In agreement, the expression of specific microRNAs (miR-129-2 and miR-146a) has been observed to correlate with functional score after injury, an accepted proximate to SCI severity (Strickland et al., 2011). Additional support for a severity effect on transcription comes from mRNA data. De Biase et al. (2005) observed strong differences in gene expression in contusive injuries of different severities, with a dramatic increase in

**Table 1 | Global analyses of microRNA expression after SCI.**

Study	SCI model	miRNA expression changes
Liu et al. (2009)	Animal: Sprague Dawley rats Contusion, NYU impactor (10 g, 12.5 mm) Level: T10 Sampling: 4 h, 1 and 7 days	4 hpo: 18 down, 23 up 1 dpo: 30 down, 27 up 7 dpo: 30 down, 30 up
Nakanishi et al. (2010)	Animal: C57BL/6 mice Compression, forceps (10 s). Level: T11–T12 Sampling: 12 h	12 hpo: 5 down, 5 up
Strickland et al. (2011)	Animal: Sprague Dawley rats Contusion, MASCIS impactor (10 g, 12.5 mm) Level: T12–T13 Sampling: 4 and 14 days	4 and 14 dpo: 32 down, 4 up
Jee et al. (2012a,b), Im et al. (2012)	Animal: ICR mice Transection Level: T9–T10 Sampling: ?	265 exp. changes
Yunta et al. (2012)	Animal: Wistar rats IH spinal cord impactor (200 kdynes) Level: T8 Sampling: 1, 3, and 7 days	1 dpo: 0 down, 0 up 3 dpo: 46 down, 5 up 7 dpo: 192 down, 11 up
Hu et al. (2013b)	Animal: Sprague Dawley rats Contusion, mod. Allen Weight Drop (8 g, 40 mm) Level: T10 Sampling: 1 and 3 days	1 dpo: 5 down, 9 up 3 dpo: 5 down, 3 up
Hu et al. (2013a)	Animals: Rats Ischemic-Reperfusion Level: lumbo-sacral segments Sampling: 2 days	2 dpo: 10 down, 38 up

*Summary of models and results employed in published microarray analyses of miRNA expression in spinal cord injury. In all cases, data from injured animals was compared to data from Sham injured animals. Information from several studies was very limited and we would encourage authors to submit microarray data to public repositories as part of the process of publication, as suggested by the Microarray Gene Expression Data Society (<http://www.mged.org>). T, thoracic level; hpo, hours post-operation; dpo, days post-operation; up and down indicate up- and down-regulated microRNAs.*

the number of genes with altered expression in moderate injuries relative to mild and severe ones. Thus, injury severity could be an important and interesting source of variation of microRNA expression that would merit further studies. Other injury features contribute to condition the microRNA profile of the spinal cord. Recently, Ziu et al. (2013) have compared the expression of six microRNAs after SCI with variable compression times showing that the duration of compression significantly alter the expression of different microRNAs.

Besides general variability in the microRNA expression patterns, Strickland et al. (2011) and our group (Yunta et al., 2012) observed a global downregulation of miRNA expression after a moderate contusive injury. Interestingly, studies by De Biase et al. (2005) and Byrnes et al. (2006) showed that moderate contusive injuries present a significant bias toward gene upregulation with a maximum at 7 days after injury, that contrast with the more balanced numbers of up- and downregulated genes observed in severe and mild injuries. Thus, the increase in the number of upregulated mRNA correlates with the general decrease of miRNA expression, which becomes particularly evident 1 week after injury (Yunta et al., 2012). Bioinformatic analyses based on miRNA target prediction data allowed us to show that almost one-third of the upregulated mRNAs in De Biase et al.'s (2005) study where targets of the downregulated microRNAs (Yunta et al., 2012). It is tempting to hypothesize that a generalized decrease in miRNA abundance reduces post-transcriptional regulation and thus causes an increase in mRNA levels after moderate SCI. Two questions immediately arise: why in moderate injuries? and, how do injuries regulate microRNA expression or biogenesis? Although we have no information to approach the first question, some is available to explore the effects of injuries on microRNA biogenesis. MicroRNA biogenesis can be regulated at different stages. The first layer governing miRNA abundance is the regulation of pri-miRNA transcription by binding of transcription factors (Treiber et al., 2012). C-Myc is one of such transcription factors, which is known to directly upregulate miR-17-92 cluster and, at the same time, to cause a widespread repression of microRNA expression (Chang et al., 2008). Interestingly, this transcription factor is significantly overexpressed at 4 and 24 h following a SCI (Di Giovanni et al., 2003). In agreement, the levels of several microRNAs repressed by c-Myc in Chang et al. (2008) experiments – miR-26a, miR-26b, miR-29a-c, miR-34a, miR-146a, miR-30a-e, let-7a, let-7d, let-7g, miR-99b, and miR-125 – become significantly decreased after SCI in our analysis whereas two members of the miR-17-92 cluster (miR-20a and miR-17) appear upregulated (Yunta et al., 2012). In addition to transcriptional regulation, processing of the pri- and pre-miRNA transcripts can be also regulated (Treiber et al., 2012). Blockage or downregulation of key proteins in the biogenesis pathway such Dicer, Drosha, DGCR8, or Exportin 5 leads to a reduction in the abundance of mature microRNAs and an accumulation of pri- and pre-microRNAs (Lee et al., 2008). Examples of this regulatory pathway are common in cancer (Lu et al., 2005) but also in other processes, such as liver regeneration after damage (Shu et al., 2011). Little is known about the variation in the expression and function of the microRNA biogenic machinery after SCI, however, Jee et al. (2012a) have recently described a downregulation of Dicer 7 days after contusive spinal cord. Dicer

downregulation is consistent with the observed general depletion in miRNA abundance observed by Strickland et al. (2011) and Yunta et al. (2012). Additionally, growth factors such as bone morphogenetic proteins (BMPs) and TGF- $\beta$  – overexpressed after SCI (McTigue et al., 2000; Xiao et al., 2010) – can contribute to regulate the microRNA biogenesis through activation of Smad proteins, which bind to pri-miRNA and enhance their Drosha-mediated processing to pre-miRNA (Davis et al., 2010). To what extent all these mechanisms contribute to the different patterns of microRNA expression following SCI remains to be elucidated.

Microarray data allows identifying consistent changes in microRNA abundance following SCI. Considering those changes observed in at least two different studies, 36 microRNAs show consistent patterns of change (see **Table 2**). These changes illustrate the complexity of interpreting the microRNA expression changes from microarray data in the SCI due to the cellular heterogeneity of the spinal cord and the multiple changes that take place after injury. Expression changes in heterogeneous samples such as the spinal cord actually correspond to the weighted mean of the transcription programs of all cell types present in the sample (Lu et al., 2003). Thus, the observed expression changes may result either from changes in gene expression within a given cell type or to changes in the relative abundance of the expressing cell types, which severely constraints the conclusion that can be derived from the expression data (Wang et al., 2006; Gosink et al., 2007). In fact, heterogeneity could be a major reason why many gene expression analyses fail a rigorous validation (Clarke et al., 2010). After injury, the spinal cord experiences changes in the relative proportions of different cell types due to the necrotic and apoptotic death of neurons and oligodendrocytes (Grossman et al., 2001; Rowland et al., 2008) and the infiltration of immune cells (Grossman et al., 2001; Profyris et al., 2004). The death of specific cell types explains the downregulation of microRNAs associated with neurons, such as miR-124 and oligodendrocytes, miR-219 (Smirnova et al., 2005; Lau et al., 2008). Both microRNA show a sustained decrease in their levels that follows the progression of cell death in these neural cells. Interestingly, Liu et al. (2009) observed an increase in miR-124 abundance during the first 4 h. A similar upregulation was previously described in brain ischemia-reperfusion by Jeyaseelan et al. (2008), which proposed that such increase “indicates that injured brain cells could be actively involved in regeneration during the first 24 h of reperfusion.” In parallel, the infiltration of immune and vascular cells explains the overexpression of specific microRNAs. The best characterized is miR-223, a neutrophil microRNA (Lindsay, 2008; Sonkoly and Pivarcsi, 2009; Izumi et al., 2011), whose upregulation reflects the infiltration of these immune cells during the acute phase of the spinal cord (Izumi et al., 2011; Yunta et al., 2012). Similarly, miR-451 – a red blood cell marker (Merkerova et al., 2008) – appears clearly increased in the first hours after injury to be later repressed or returned to control condition probably reflects the entrance of erythrocytes in the damaged area during the acute phase and their clearance later on.

## MicroRNAs IN THE REGULATION OF INFLAMMATION FOLLOWING SCI

Spinal cord injury activates an inflammatory response that is initiated by the alteration of the blood–spinal cord barrier, followed by

**Table 2 | Prominent changes in microRNA expression in murine models of spinal cord injury.**


MicroRNA	Time after injury						
	4 h	12 h	1 days	3 days	4 days	7 days	14 days
miR-1		Up <sup>2</sup>	Up <sup>1</sup>	Down <sup>4</sup>	Down <sup>3</sup>	Up <sup>1</sup> /Down <sup>4</sup>	Down <sup>3</sup>
miR-100	Up <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-103	Up <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-107	Up <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-124	Up <sup>1</sup>	Down <sup>2</sup>	Down <sup>1</sup>	Down <sup>4</sup>	Down <sup>3</sup>	Down <sup>1,4</sup>	Down <sup>3</sup>
miR-125b-3p			Down <sup>5</sup>			Down <sup>4</sup>	
miR-126			Down <sup>5</sup>	Down <sup>5</sup>		Down <sup>4</sup>	
miR-127	Up <sup>1</sup>		Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-128	Up <sup>1</sup>		Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-129-1					Down <sup>3</sup>		Down <sup>3</sup>
miR-129-2					Down <sup>3</sup>		Down <sup>3</sup>
miR-129-3p		Down <sup>2</sup>				Down <sup>4</sup>	
miR-129*	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-133a	Up <sup>1</sup>	Up <sup>2</sup>	Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-133b	Up <sup>1</sup>	Up <sup>2</sup>	Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-138	Down <sup>1</sup>		Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-146a			Up <sup>1</sup>			Up <sup>1,4</sup>	
miR-17			Up <sup>1</sup>	Up <sup>4</sup>		Up <sup>1</sup>	
miR-181a	Up <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-21			Up <sup>1</sup>	Up <sup>4,5</sup>	Up <sup>3</sup>	Up <sup>1,4</sup>	Down <sup>3</sup>
miR-219-2-3p	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-219-5p	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-223	Up <sup>1</sup>	Up <sup>2</sup>	Up <sup>1,4</sup>	Up <sup>4</sup>	Up <sup>3</sup>	Up <sup>1,4</sup>	Up <sup>3</sup>
miR-30b-5p	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-30c	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-30d	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-338*	Down <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-342		Down <sup>2</sup>				Down <sup>4</sup>	
miR-34a	Down <sup>1</sup>		Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-379*	Down <sup>1</sup>		Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-451	Up <sup>1</sup>	Up <sup>2</sup>	Down <sup>1</sup> /Up vs 7 dpo <sup>4</sup>			Down <sup>1</sup>	
miR-487b	Up <sup>1</sup>		Down <sup>1</sup>			Down <sup>1,4</sup>	
miR-495	Down <sup>1</sup>	Down <sup>2</sup>	Down <sup>1</sup>	Down <sup>4</sup>		Down <sup>1,4</sup>	
miR-708			Down <sup>5</sup>	Down <sup>4</sup>		Down <sup>4</sup>	
miR-99a	Up <sup>1</sup>		Down <sup>1</sup>	Down <sup>5</sup>		Down <sup>1,4</sup>	
miR-let-7b			Down <sup>5</sup>	Down <sup>5</sup>		Down <sup>4</sup>	

The table details the changes in microRNA expression following spinal cord injury of dysregulated microRNAs according to, at least, two different studies. Data derived from the following microarray analyses: <sup>1</sup>Liu et al. (2009); <sup>2</sup>Nakanishi et al. (2010); <sup>3</sup>Strickland et al. (2011); <sup>4</sup>Yunta et al. (2012); <sup>5</sup>Hu et al. (2013a,b). Unless stated, Up and Down correspond to upregulated and downregulated respect to Sham control individuals respectively.

the sequential infiltration of peripheral immune cells, the activation of the microglia and the induction of inflammatory signaling pathways. MicroRNAs play important roles in controlling signaling pathways and the dynamics of the immune response during pathogenic immunological conditions (Lindsay, 2008; Sonkoly

and Pivarcsi, 2009; Chen et al., 2013). Following SCI, several microRNAs undergo expression changes that can be related to the immune response, either associated with the invading immune cells or participating in the modulation of inflammatory pathways (Figure 1).





Immediately after the injury, the blood–spinal cord barrier becomes disturbed and the blood immune cells infiltrate the damaged area (Profyris et al., 2004; Jones et al., 2005). Early increases in the expression of CAMs play a key role in the process of immune cells recruitment and extravasation into the nervous tissue. Some of these expression changes can be controlled by microRNAs. In particular, the upregulation of *VCAM1* mRNA (Aimone et al., 2004) occurs in parallel to the downregulation of its regulator miR-126 (Harris et al., 2008) during the first week after injury (Yunta et al., 2012; Hu et al., 2013b). The subsequent immune cell infiltration is responsible for several changes in the microRNA

profile of the spinal cord, as previously commented. Neutrophil infiltration explains the upregulation of miR-223 (Lindsay, 2008; Sonkoly and Pivarcsi, 2009; Izumi et al., 2011), whereas increased expression of the lymphocyte specific miR-142 (Wu et al., 2007) correlates with the access of these immune cells to the injury site during the first week (Yunta et al., 2012). MicroRNAs are also involved in the activation of microglia and macrophages. Particularly, the downregulation of miR-124 contributes to resting phenotype of microglia by targeting CEBPα, a master transcription factor important for the development of myeloid cells (Ponomarev et al., 2011; Guedes et al., 2013). miR-124 shows a

sustained downregulation after injury (Deo et al., 2006; Liu et al., 2009; Nakanishi et al., 2010; Yunta et al., 2012) that may underlie microglial activation. However, miR-124 is a well-characterized neuronal microRNA and its downregulation likely reflects the extension of neuronal death that characterizes the secondary damage of SCI.

The inflammatory response is modulated by a number of key molecular immune mediators, such as cytokines, chemokines (TNF- $\alpha$ , IL-6, IL-1 $\beta$ ) or the complement cascade (C1qb; Hausmann, 2003; Profyris et al., 2004; Jones et al., 2005), which are known targets of microRNAs. Particularly, overexpression of three key pro-inflammatory cytokines that modulate the inflammatory response in the SCI is likely regulated by microRNA experiencing expression changes after injury. Different SCI studies (Liu et al., 2009; Yunta et al., 2012) have suggested that increasing levels of the pro-inflammatory and pro-apoptotic factor TNF- $\alpha$  after injury (Tyor et al., 2002) may result from downregulation of its regulators miR-181 (Hutchison et al., 2013) and miR-125b (Tili et al., 2007) after SCI (Liu et al., 2009; Yunta et al., 2012; Hu et al., 2013b). Other microRNAs, such as miR-411, miR-99a, that appear downregulated after SCI (Liu et al., 2009; Yunta et al., 2012; Hu et al., 2013b) have been also predicted to target TNF- $\alpha$  by bioinformatics analysis. On the other hand, the increased levels of cytokine IL-6 during the first days after injury correlate with a reduced expression of its regulators let-7a (Iliopoulos et al., 2009) or miR-181a (Tili et al., 2007; Iliopoulos et al., 2009). Finally, the observed downregulation of miR-30b-5p and miR-30c during the first week after injury (Liu et al., 2009; Yunta et al., 2012) could be related to the overexpression of their target IL-1 $\beta$ . Additionally, it has been recently described that upregulation of miR-20a induces a cytotoxic environment with increased IL-6, TNF- $\alpha$ , IL-1- $\beta$ , and COX-2 expression in the spinal cord (Jee et al., 2012b) due to miR-20a-mediated inhibition of neurogenin 1 (*NGN-1*). Inhibition of this microRNA – which is upregulated after injury (Liu et al., 2009; Jee et al., 2012b) – lead to a significant improvement in functional recovery associated to a reduced inflammation and the increased survival of motor neurons (Jee et al., 2012b).

Interestingly, pro-inflammatory cytokines lead to the activation of the NF- $\kappa$ B signaling pathway, which is also under microRNAs regulation (Ma et al., 2010). In particular, downregulation of miR-9 and miR-199 (Yunta et al., 2012) may induce the overexpression of the NF- $\kappa$ B pathway genes *p50Nfkb* and *ikkb* (Chen et al., 2008; Bazzoni et al., 2009; Wang et al., 2011). The increased expression of miR-21 may also contribute to the regulation of this pathway but its role is less clear, as it exhibits both pro- and anti-inflammatory effects. miR-21 targets PTEN, a negative regulator of NF- $\kappa$ B (Iliopoulos et al., 2010), but also PDCD4, which promotes NF- $\kappa$ B activation and inhibits the expression of IL-10 (Frankel et al., 2008; Sheedy et al., 2010; Young et al., 2010). On the opposite side, expression changes in several microRNAs that have been observed after SCI may attenuate the activation of NF- $\kappa$ B pathway, contributing to the attempts of the damaged spinal cord to recover homeostasis (Bareyre and Schwab, 2003). These changes include the increased expression of miR-146a at 7 days after injury (Liu et al., 2009; Yunta et al., 2012), which negatively regulates NF- $\kappa$ B expression (Taganov et al., 2006; Ma et al., 2010). Interestingly, miR-146a expression is induced by NF- $\kappa$ B and, thus, its

overexpression at 7 days after injury may be consequence of the increased levels of NF- $\kappa$ B in the previous days (Bethea et al., 1998), forming a negative feedback that may cause the inactivation of NF- $\kappa$ B pathway.

A second group of factors with a prominent role in inflammation after SCI are the complement proteins (Brennan et al., 2012). Complement activation is involved in the removal of cellular debris, but it may also promote clearance of mildly damaged cells contributing to secondary cell death and demyelination. Complement protein C1qb increases its expression in the first day after injury and persist upregulated at least 5 weeks later (Aimone et al., 2004). C1q knockout mice show improved locomotor recovery and reduced secondary tissue damage after contusive SCI (Galvan et al., 2008). Interestingly, C1qb is a predicted target of miR-103 (Perri et al., 2012), which appears downregulated in the first week after injury (Liu et al., 2009; Yunta et al., 2012). Thus, miR-103 downregulation could be responsible for the overexpression of the complement protein C1qb and its associated deleterious effects.

Inflammation is also stimulated through the inhibition of anti-inflammatory pathways, such as the downregulation of pSMAD2, SMAD4, and TGFBR2 by observed upregulation of members of the miR-17-92 microRNA cluster (Mestdagh et al., 2010) or the silencing of the anti-inflammatory neuroprotective cytokine IL-10 by miR-98, miR-106a (Sharma et al., 2009; Liu et al., 2011). Many other microRNAs have been related to inflammation in SCI based on *in silico* predictions. Bioinformatics analyses predict that anti-inflammatory mRNAs annexin A1, annexin A2 and annexin A7 mRNAs are potential targets of the SCI, upregulated microRNAs miR-221, miR-1, and miR-323, respectively (Liu et al., 2009; Hu et al., 2013a). The list also includes miR-127, miR-411, and miR-34a, significantly downregulated after SCI in adult rats and which according to Liu et al. (2009) should lead to increased inflammation. Other pro-inflammatory microRNAs, such as miR-152, miR-214, miR-206, and miR-221, show significant changes in isolated studies or even show opposite expression trend (i.e., miR-1) in different studies (Liu et al., 2009; Strickland et al., 2011; Yunta et al., 2012). All these results will require further assays to elucidate the role in the inflammatory process of the SCI.

### MicroRNA REGULATION OF CELL DEATH REGULATION IN SCI

Cell death is a hallmark of the pathophysiology of SCI (Crowe et al., 1997; Liu et al., 1997). The programmed cell death (apoptosis) that characterizes the SCI secondary damage is a gene-controlled process that is stimulated or inhibited by a variety of regulatory factors including several microRNAs (Wang, 2010). Previous studies have shown that SCI alters the transcription levels of a substantial number of genes associated with the regulation of apoptosis (Aimone et al., 2004). The resulting scenario is complex, combining both temporal and spatial changes in the expression of both pro and anti-apoptotic signals. In agreement, many microRNAs have been proposed to promote or inhibit cell death during the course of SCI, with discrepancies among microRNA expression profiling studies. These discrepancies highlight the intricate roles that miRNAs play in the regulation of cellular processes and the difficulty to identify the precise activity of dysregulated microRNAs. However, two recent studies have provided direct evidence of

microRNAs involvement in cell death modulation following SCI (**Figure 1**).

The first study deals with miR-20a, which shows an increased expression early after injury (24 h) that persists at least for 1 week (Liu et al., 2009; Jee et al., 2012b). miR-20a inhibits the expression of *Ngn1*, a protein with a key role in maintenance of cell survival, self-renewal, and neurogenesis in normal and injured spinal cords. Silencing *Ngn1* by siNgn1 or infusion of miR-20a into uninjured mouse spinal cord reproduced SCI-like symptoms including apoptotic death of neural cells, whereas the administration of anti-miR-20a decreased apoptosis, reduced tissue damage and functional deficits were significantly ameliorated (Liu et al., 2009; Jee et al., 2012b). In a similar way, miR-486 is upregulated at 7 days after injury (Jee et al., 2012a). miR-486 represses neurogenic differentiation 6 (*NeuroD6*), a protein that promotes neuronal survival by increased expression of the ROS scavenger proteins (Jee et al., 2012a). Infusion of miR-486 into the normal spinal cord of mice reproduces SCI symptoms including increased neuronal death, whereas silencing miR-486 after SCI produced a decrease in the magnitude of neuronal death and led to a significant improvement in motor recovery. Therefore upregulation of miR-486 following SCI promotes neurodegeneration by suppressing *NeuroD6*, pointing to this microRNA as a potential target for therapeutic interventions (Jee et al., 2012a).

Further evidences of the roles of microRNAs on secondary cell death come from microRNA expression changes that are accompanied by changes in the expression of their apoptotic gene targets. Several pro and anti-apoptotic microRNAs act on key apoptosis molecules, such as caspases, *Fas/CD95*, *c-Myc*, *TNF- $\alpha$* , or members of the *BCL-2* family. For example, the decreased expression of the *let-7/miR-98* family members miR-96 and miR-146a (Liu et al., 2009; Strickland et al., 2011; Yunta et al., 2012) would promote apoptosis by increasing the expression of their targets, the pro-apoptotic proteins caspase 3 (Citron et al., 2000; Aimone et al., 2004) and *Fas/CD95* (Casha et al., 2001). On the contrary, overexpression of miR-21 would protect neural cells from death by repressing the expression of the pro-apoptotic molecules *Fas ligand* (Buller et al., 2010), *TPM1* and *PTEN* (Hafez et al., 2012; Han et al., 2012), and *PDCD4* (Frankel et al., 2008). *BCL-2* modulation is highly representative of the complexity of microRNA regulation of cell death in SCI. Upregulation of miR-15b (Liu et al., 2009) would decrease *BCL-2* (Cimmino et al., 2005; Saito et al., 2006) and induce apoptosis. However, upregulation of miR-15b is counteracted by the decreased expression during the first week of miR-138 and miR-148b, which also target *BCL-2* (Liu et al., 2009; Yunta et al., 2012). Downregulation of these microRNAs is broadly consistent with the increase in the number of *BCL-2*-positive cells present 3 days after injury (Saito et al., 2000; however, see Qiu et al., 2001), although microRNA downregulation extends throughout the 7-day period after injury, which is the time-point when the number of *BCL-2*-positive cells is progressively reduced. Other microRNAs targeting *BCL-2* appear dysregulated after SCI. Regulation of *BCL-2* by miR-107 was discussed in the profiling study by Liu et al. (2009). These authors observed a miR-107 upregulation 4 h after injury, which they proposed should decrease *BCL-2* levels and induce apoptosis, to be later downregulated at 7 dpi (also observed in Yunta et al.,

2012) promoting cell survival. miR-1 represents a puzzling case that appears upregulated in Liu et al. (2009) and downregulated in the analyses by Strickland et al. (2011) and Yunta et al. (2012).

In addition to modulation of genes that regulate apoptosis, microRNAs also participate in the disruption of the calcium signaling or the oxidative stress events triggered after SCI that contribute to secondary cell death. Expression of the gene coding for the  $\text{Ca}^{2+}$ -related genes such as  $\text{Ca}^{2+}$  pump, voltage-gated (L-type)  $\text{Ca}^{2+}$  channels or  $\text{Ca}^{2+}$ -permeable ionotropic glutamate (AMPA) channels, is decreased with injury and post-transcriptionally regulated by microRNAs. Several studies have shown that upregulated miR-223 reduces the expression of the NR2B and GluR2 subunits of the NMDA and AMPA receptors, respectively (Kaur et al., 2012). Similarly, decreased expression of voltage-gated (L-type)  $\text{Ca}^{2+}$  channels may be result of upregulated miR-21 (Carrillo et al., 2011). These could cause an increment of intracellular  $\text{Ca}^{2+}$  concentration level that accompanies traumatic SCI, and could trigger mechanisms of secondary cell death, such as calpain activation. MicroRNAs also play an important role in the regulation of oxidative stress, a hallmark of the secondary damage of SCI that has received much attention in the attempts to develop effective therapies (Jia et al., 2012). Recent reports have demonstrated that miR-486 repress the expression of *NeuroD6*, a neuroprotective protein that promotes the expression of ROS scavenger proteins, such as *GPX3*, *selenoprotein-N*, and *thioredoxin* (Jee et al., 2012a). Upregulation of miR-486 – observed in motor neurons at 7 days after injury in murine models of SCI – leads to the repression of *NeuroD6* expression, and consequently to a decrease in the expression of ROS scavenger proteins and increased neurodegeneration mediated by oxidative stress (Jee et al., 2012a).

Microarray analyses revealed increased expression of genes associated with anti-oxidant actions, such as *SOD1*, *SOD2*, *catalase*, and *GPX* (Di Giovanni et al., 2003; Aimone et al., 2004). This overexpression of the mitochondrial *SOD2* gene (*sod2*) 7 days after injury (Santoscoy et al., 2002; Sugawara et al., 2002) is consistent with the downregulation of its modulator miR-145 (Dharap et al., 2009) described in Yunta et al. (2012). However, the bioinformatics analysis performed by Liu et al. (2009) revealed that some anti-oxidant genes such as *SOD1* and *catalase* gene are potential targets of the upregulated miR-206, miR-152, and miR-214. Moreover, it has been proposed that downregulation of miR-1 and miR-129, which regulate transcription, differentiation or prevent post-mitotic cells from re-entering the cell cycle, could cause neural cells to become aberrantly mitotic, increasing the number of apoptotic cells observed at the injury site after SCI (Bhalala et al., 2013).

### MicroRNA MODULATION OF ASTROCYTE REACTIVITY AND GLIAL SCAR

Astrogliosis is another hallmarks of the cellular response to SCI. It consists in an early hypertrophic neuroprotective phase followed by a hyperplastic phase characterized by the formation of a dense glial scar that inhibits CNS regeneration during the subacute and chronic phases of the SCI (Sofroniew, 2009). Recent genomic analyses have shown reactive astrogliosis is associated to a rapid, but quickly attenuated, induction of gene expression

(Zamanian et al., 2012). Increasing evidence supports the involvement of several microRNAs in the regulation of the astrocyte response to injury, including four microRNAs that appear dysregulated in studies of SCI. The best characterized is miR-21. Its expression increases in a time-dependent manner following SCI (Liu et al., 2009; Bhalala et al., 2012; Yunta et al., 2012; Hu et al., 2013b) and is highly expressed in astrocytes during the chronic stage (Bhalala et al., 2012). miR-21 expression after SCI shows a marked spatial pattern, with highest expression in the astrocytes adjacent to the lesion area (Bhalala et al., 2012). The role of miR-21 in astrogliosis has been studied in detail using transgenic mice that overexpress in astrocytes either miR-21 or a miRNA sponge designed to inhibit miR-21 function (Bhalala et al., 2012). The results from these studies demonstrate that miR-21 overexpression in astrocytes abrogates the hypertrophic astrocytic response after severe SCI, which is consistent with previous studies *in vitro* (Sahni et al., 2010; Sayed and Abdellatif, 2011). On the contrary, miR-21 inhibition enhances the hypertrophic response in early and chronic stages after SCI (Bhalala et al., 2012). BMP signaling following SCI mediates the miR-21 and astrocytic response through the opposing effects of the BMP receptors BMPRIa and BMPRIb (Sahni et al., 2010). BMPRIa signaling decreases levels of miR-21 and induces reactive astrocytic hypertrophy, whereas BMPRIb signaling increases miR-21 levels and negatively regulates astrogliosis. These findings point to the BMP–BMPRI–miR-21 axis as a key regulator of astrocytic hypertrophy and glial scar progression after SCI, modulating the pro-reactive effects of the inflammatory signaling.

A second microRNA that has been related to astrogliosis is miR-125b. Overexpression of miR-125b correlates with the overexpression of the astrogliosis markers GFAP and vimentin in several neurological disorders (Pogue et al., 2010). *In vitro* studies show that miR-125b downregulation in IL-6 stimulated reactive astrocytes increases the expression of its target cyclin-dependent kinase inhibitor 2A (CDKN2A), a negative regulator cell growth, and attenuates cell proliferation. Thus, evidences indicate that miRNA-125b upregulation contributes to astrogliosis. However, contrary to expectations, miR-125b appears downregulated during the first week after injury (Yunta et al., 2012), which would contribute to inhibit astrocyte proliferation and astrogliosis.

The miR-181 family of miRNAs is another candidate for post-transcriptional regulation of neuroinflammation and reactive gliosis (Hutchison et al., 2013). miR-181s are constitutively expressed in astrocytes but inflammation causes its downregulation, in agreement with *in vivo* observations following SCI (Liu et al., 2009; Yunta et al., 2012). Genetic studies demonstrated that miR-181 inhibits the production of multiple pro-inflammatory cytokines (TNF- $\alpha$ , IL-6, IL-1 $\beta$ , IL-8, LIF, and HMGB1) and increases the levels of the anti-inflammatory cytokine IL-10 (Hutchison et al., 2013) in cultured astrocytes under LPS inflammatory exposure. Thus, miR-181s act as negative regulators of astrogliosis, reducing the expression of reactivity promoters such as pro-inflammatory cytokines (Balasingam et al., 1994; Sofroniew, 2009) and FGF2 (Goddard et al., 2002), and increasing the expression of reactivity inhibitors, such as IL-10 (Balasingam and Yong, 1996).

Similarly to miR-181s, miR-146a is a negative regulator of the astrocyte response to inflammation and, consequently, a

negative regulator of astrogliosis (Iyer et al., 2012). miR-146a is expressed in reactive astrocytes in the areas of prominent gliosis (Iyer et al., 2012), and appears upregulated 7 dpi after SCI (Liu et al., 2009; Strickland et al., 2011; Yunta et al., 2012) as well as in several neurodegenerative pathologies (Junker et al., 2009; Cui et al., 2010; Iyer et al., 2012). Studies with astroglial cell lines and primary astrocytes have shown that IL-1 $\beta$  stimulation induces a prominent upregulation of miR-146a expression. Overexpression of miR-146a has anti-inflammatory effects and significantly reduces the expression of signaling molecules downstream IL-1 $\beta$ , such as IRAK-1, IRAK-2, and TRAF-6 and inhibits the release of pro-reactive and pro-inflammatory factors, including IL-6 and COX-2 (Iyer et al., 2012). Thus, miR-146a forms a negative feedback of the IL-1 $\beta$  signaling, being induced by IL-1 $\beta$  but blocking the expression of its downstream response. Overexpression of miR-146a following SCI adds to the overexpression of miR-21 and miR-125b to limit astrocyte reactivity. On the contrary, downregulation of miR-181 family members would promote astrocyte reactivity, indicating a fairly complex regulation of the astrocyte response. It is interesting to note that according to recent studies, the disruption of microRNA biogenesis by deletion of Dicer leads to an altered mature astroglial transcriptome signature that resembles to a reactive state (Tao et al., 2011), which adds a further layer of complexity to the whole picture.

#### THE ROLES OF microRNAs IN AXONAL REGENERATION, MYELINATION, AND OTHER PROCESSES

Although less studied, microRNAs also seem to contribute to the regulation of other pivotal processes in the SCI pathophysiology, such as axonal regeneration, remyelination, or pain. As we mentioned before, failure to produce a sustained regenerative response is one of the critical features of the CNS. Local environmental clues are largely responsible for the lack of regeneration but intracellular specific features associated to neural cell maturation are also involved (see, for example, Goldberg, 2003). Different evidences indicate that microRNAs can contribute to these changes. In fact, microRNAs have a critical role in neurite outgrowth in post-mitotic neurons as demonstrated in Dicer conditional knockout mice. In this animals, silencing of Dicer and the subsequent inhibition of the microRNA biogenesis causes defects in neurite outgrowth and decreased soma size but has not influence in neurogenesis, cortical patterning or cell survival (Hong et al., 2013). Additional evidences have been obtained from the zebrafish, a model of spontaneous axonal regeneration in the damaged spinal cord (Becker et al., 1997). In this fish, miR-133b is overexpressed in regenerating neurons following SCI and its inhibition using antisense morpholinos result in reduced axonal regeneration (Yu et al., 2011). Functionally, miR-133b contributes to spinal cord regeneration through the downregulation of its target RhoA, a small GTPase that inhibits axonal growth. Contrary to zebrafish, miR-133b shows a significant downregulation at 1 and 7 days after contusive SCI in mammals (Liu et al., 2009; Yunta et al., 2012), which may contribute to their reduced neuroregenerative capacity. miRNA-124 presents a somehow similar behavior that may also contribute to reduce axonal regeneration after SCI. Previous studies have

shown that miR-124 overexpression in differentiating mouse P19 cells promotes neurite outgrowth, while miR-124 inhibition reduces it (Yu et al., 2008). Thus the observed miR-124 downregulation after SCI may also contribute to hinder the axonal regenerative capacities of spinal cord neurons. However, microRNAs may also contribute to the activation of pro-regenerative gene programs after injury. Di Giovanni et al. (2005a) described the overexpression at 7 and 28 days after SCI of a gene cluster that comprise known promoters of the neural plasticity and the neurite outgrowth, including synaptotagmin-1. Interestingly, synaptotagmin-1 is a target of miR-34a (Agostini et al., 2011), and its upregulation after SCI is consistent with the observed downregulation of miR-34a at 3 and 7 after injury (Liu et al., 2009; Yunta et al., 2012).

The progressive loss of myelin in the areas surrounding the injury is another critical feature of the SCI that results from the combined effects of damage to oligodendrocytes and remyelination failure. Evidences have confirmed that microRNA loss of function due to Dicer1 ablation in mature oligodendrocytes causes demyelination, gliosis, and neuronal degeneration (Shin et al., 2009; Dugas et al., 2010). More precisely, Shin et al. (2009) identified miR-219 as a central actor in myelin maintenance and remyelination. miR-219 is highly expressed in mature oligodendrocytes and when is lost due to Dicer1 ablation, miR-219 target ELOVL7 increases its expression resulting in lipid accumulation in myelin-rich areas and disrupting the stability of the membranes (Shin et al., 2009). Strikingly, miR-219 abundance is markedly reduced after SCI (Liu et al., 2009; Yunta et al., 2012) although this decrease may also reflect the loss of spinal cord oligodendrocytes that takes place after injury. Further studies are needed to determine the contribution of microRNAs in demyelination and remyelination and to evaluate their use as therapeutic tools in the SCI and other CNS pathologies.

In addition to their direct roles in most processes implicated in the pathophysiology of the SCI, microRNAs are also involved in the functional consequences of SCI, including the neuropathic pain. Neuropathic pain is the manifestation of maladaptive plasticity in the nervous system characterized by pain in the absence of a stimulus and reduced nociceptive thresholds (Scholz and Woolf, 2007). It is a debilitating accompaniment of SCI that affects up to 50% SCI patients and limit their ability to achieve an optimal level of activity (Mann et al., 2013). Plastic changes in sensory neuron excitability are considered the cellular basis of neuropathic pain, although a growing body of evidence also implicates activated microglia and astrocytes as key players in the development of pain (Scholz and Woolf, 2007). Although information on the roles of miRNAs in neuropathic pain following SCI is very restricted, available evidences indicate that microRNAs expression at the spinal cord respond to pain induction (Kusuda et al., 2011; Genda et al., 2013; Li et al., 2013), although evidences are sometimes contradictory (Brandenburger et al., 2012). There are also evidences of the contribution of microRNAs in the development of central neuropathic pain. Recently, Im et al. (2012) demonstrated that SCI reduced the expression of miR-23b in GABAergic neurons from the spinal cord while increasing the levels of NADPH oxidase 4 (NOX4), a target of miR-23b and a key factor in the production of ROS

and pain induction. Reduction of NOX4 expression and neuropathic pain was observed after infusion of miR-23b to the spinal cord confirming the involvement of this microRNA in pain regulation. Similarly, intrathecal administration of miR-124 inhibits the activation of spinal cord microglia, reducing inflammation and preventing neuropathic pain (Willemsen et al., 2012). Many other microRNAs have been described to affect neuropathic pain at different levels, particularly at the dorsal root ganglia. However, a review of this complex system lies beyond the scope of the present article.

## CLINICAL APPLICATIONS: microRNA-BASED THERAPEUTIC AND DIAGNOSTIC TOOLS

The capability of individual microRNAs to reduce the expression of numerous components of cellular networks supposes an opportunity to modulate the cell phenotypes by manipulating the expression or function of microRNAs. The possibility of a therapeutic use of the miRNAs is highly appealing due to their capability to modulate entire gene programs through tuning, not blunting, the expression of their targets (Hydbring and Badalian-Very, 2013). Moreover, microRNA deregulation has a critical role in a wide range of pathologies, and, indeed, several studies have identified specific deregulated microRNAs that can constitute potential targets of therapeutic approaches in a wide range of pathologies. As a consequence, in the short time from its discovery in humans, a microRNA-based therapeutic for suppression of hepatitis C virus has already entered phase II clinical trial and several others are on their way (see review in Hydbring and Badalian-Very, 2013).

Therapeutic approaches are based on local or systemic administration of either antagonists (anti-miRs) of endogenous microRNAs that show a gain-of-function in diseased tissues or mimics that replace downregulated microRNAs. Both modulators incorporate chemical modifications (phosphorothionate backbones, locked nucleic acids or LNAs, etc.) to confer resistance to nucleases, increase stability during delivery, and facilitate cellular uptake. Anti-miRs are siRNAs designed to inhibit miRNAs through complementary base pairing, in a similar way to siRNA. Since binding is irreversible, the resulting miRNA duplex cannot be processed by RISC and/or degraded. Anti-miRs are synthesized as short single-stranded oligonucleotides of small size that make delivery possible without vehicle-systems. On the contrary, administration of microRNA mimics, also known as “miRNA replacement therapy,” aims to re-introduce miRNAs into diseased cells that are normally expressed in healthy cells. miRNA mimics is expected to re-activate pathways required for normal cellular welfare and block those driving disease. To be processed correctly by the cellular RNAi-machinery, miRNA mimics need to be double-stranded, which confer them greater chemical complexity and larger size. Thus, miRNA mimics require delivery techniques similar to those employed in siRNA therapeutics, including microvesicles, exosomes or adeno-associated viruses, which increase the complexity of replacement therapies.

The critical contribution of the phenotypic changes in the neural, immune, and vascular cells in the pathophysiology of the SCI and the capabilities of miRNAs to modulate these changes (see previous sections) make miRNA therapeutics a highly promising approach to be explored. Although no microRNA-based therapy

has entered clinical trials for SCI up to now, preclinical assays provide the necessary proof of concept. Experimental treatments with anti-miRs have demonstrated that inhibition of specific dysregulated microRNAs back to pre-injury levels can effectively reduce cell death. For example, Jee et al. (2012b) showed that local infusion of miR-20a inhibitor in the injured spinal cord reduced the expression of pro-apoptotic genes, and promoted neuron survival and functional recovery. Similarly, Hu et al. (2013b) also employed microRNA inhibition to explore the neuroprotective role of miR-21. Intrathecal infusion of miR-21 antagonist resulted in over-expression of pro-apoptotic genes, increased cell death and reduced recovery of the hindlimb motor function. On the other hand, two articles have proven that miRNA replacement can be also a viable therapeutic approach. Willemsen et al. (2012) showed that intrathecal administration of miR-124 prevents persistent pain in rats, probably due to its modulation of microglial activation (Ponomarev et al., 2011). Im et al. (2012) reported similar results infusing miR-23b intrathecally to a murine model of neuropathic pain. According to these authors, restoring the normal levels of miR-23b reduced the expression of inflammatory proteins, particularly NOX4 in GABAergic neurons, protecting them from cell death, and ameliorating the neuropathic pain derived from SCI. These preclinical data strongly supports the feasibility of microRNA-based therapeutics in SCI treatment, although key aspects –including timing and side effects – remain to be elucidated. Detailed analysis characterizing the beneficial effects and determining the underlying mechanisms are strongly needed before microRNAs can reach the clinic in the treatment of the SCI.

Another clinically relevant but yet unexplored topic concerns how SCI alters the profiles of circulating microRNAs, and how information on these alterations can be used for diagnostic and prognostic purposes. Circulating or cell-free microRNAs are released to the body fluids either actively by secretion in exosomes or microvesicles or in association with RNA-binding proteins such as AGO2 and HDL; or passively, within apoptotic bodies liberated from dying cells (Chen et al., 2012; Zampetaki and Mayr, 2012). Encapsulation within lipid vesicles or association to binding proteins confers high stability to circulating microRNAs, despite the presence of large amounts of RNase in the body fluids (Chen et al., 2012; Li et al., 2012). Stability of circulating microRNAs together with the changes in microRNA expression in pathological states make circulating microRNAs promising biomarkers. In fact, since their discovery in all body fluids in 2008, nearly 500 articles have proposed several circulating microRNAs as biomarkers for different pathologies, including cancer, cardiovascular pathologies, and CNS injuries, among others (Laterza et al., 2009; Kosaka et al., 2010; Zhang et al., 2011; Pritchard et al., 2012). For example, analysis of the expression of circulating microRNAs after a traumatic brain injury has identified several potential biomarkers in humans (Redell et al., 2010) and rats (Balakathiresan et al., 2012). Also, it has been shown that the plasma concentration of neuron marker miR-124 becomes significantly increased after acute stroke (Laterza et al., 2009). However, quantification of circulating microRNAs can be challenging due to (i) their low concentrations, (ii) the effects of cell contaminants, and (iii) the absence of endogenous

controls for normalization. Low concentrations of circulating microRNAs suppose a technical challenge for microRNA extraction and quantification, and also increase the risk that microRNAs from contaminant cells, which are at much higher concentrations, would mask or confound the circulating microRNA profile (Kirschner et al., 2011).

Circulating microRNAs are also highly interesting due to their possible role as paracrine regulators. Circulating microRNAs can be transferred to neighboring or distant cells, altering the expression of target genes and regulating various functions, including proliferation, death or even tumor cell invasion (see Zhu and Fan, 2011, and references therein). Interestingly, microRNA transfer occurs when microRNAs are wrapped in exosomes, microvesicles, or apoptotic bodies, as well as when packaged with proteins (Zhu and Fan, 2011). Although most available evidence deals with circulating microRNA transfer in immune and vascular cells, a recent article has also demonstrated that miRNA transfer occurs between neural cells. According to Morel et al. (2013), neurons are able to secrete exosomes containing miR-124a, which are internalized by astrocytes causing an increase in the glutamate transporter GLT1.

## CONCLUDING REMARKS

Spinal cord injury is a complex pathology that induces strong cellular and molecular changes in the nervous, immune, and vascular systems. These changes alter the expression of the microRNAs - small non-coding RNAs that post-transcriptionally regulate the expression of thousands of genes- to different degrees up to a general downregulation of the microRNA expression. Bioinformatic analyses of the microRNA and mRNA expression profiles in the injured spinal cord have predicted that microRNA dysregulation strongly affects processes developing after the SCI. However, much more research analyzing the expression of specific cell populations and evaluating the effects of microRNA dysregulation is still needed if we want to validate the bioinformatic predictions, and to precisely characterize the changes in microRNA expression after SCI as well as their causes and their functional consequences. The pioneering studies developed up to now have been able to demonstrate the active role of individual microRNAs in the regulation of key processes of the SCI, such as cell death, inflammation, and astrogliosis. These results strongly suggest that microRNAs can be highly valuable therapeutic targets to modulate the deleterious events that follow SCI and to promote regenerative responses that will contribute to reduce the functional deficits associated to the SCI.

## AUTHOR CONTRIBUTIONS

All authors contributed in the conception and design of the present review, as well as in drafting and revising the manuscript. All authors have given full approval to the present version for its publication.

## ACKNOWLEDGMENTS

We would like to thank Dr. Tommaso Pizzorusso for the invitation to contribute to the present volume, and to two anonymous reviewers for their useful comments. Present contribution has been funded by Instituto de Salud Carlos III (MINECO), PI

12/28282. Rosa Navarro-Ruiz is funded by FISCAM (grant PI 2010/19) and Ángela del Águila by La Marato de TV3 foundation (grant exp.112131).

## REFERENCES

- Agostini, M., Tucci, P., Steinert, J. R., Shalom-Feuerstein, R., Rouleau, M., Aberdam, D., et al. (2011). microRNA-34a regulates neurite outgrowth, spinal morphology, and function. *Proc. Natl. Acad. Sci. U.S.A.* 108, 21099–21104. doi: 10.1073/pnas.1112063108
- Aimone, J. B., Leasure, J. L., Perreau, V. M., and Thallmair, M. (2004). Spatial and temporal gene expression profiling of the contused rat spinal cord. *Exp. Neurol.* 189, 204–221. doi: 10.1016/j.expneurol.2004.05.042
- Baek, D., Villen, J., Shin, C., Camargo, F. D., Gygi, S. P., and Bartel, D. P. (2008). The impact of microRNAs on protein output. *Nature* 455, 64–71. doi: 10.1038/nature07242
- Bak, M., Silaharoglu, A., Moller, M., Christensen, M., Rath, M. F., Skryabin, B., et al. (2008). MicroRNA expression in the adult mouse central nervous system. *RNA* 14, 432–444. doi: 10.1261/rna.783108
- Balakathiresan, N., Bhomia, M., Chandran, R., Chavko, M., McCarron, R. M., and Maheshwari, R. K. (2012). MicroRNA let-7i is a promising serum biomarker for blast-induced traumatic brain injury. *J. Neurotrauma* 29, 1379–1387. doi: 10.1089/neu.2011.2146
- Balasingam, V., Tejada-Berges, T., Wright, E., Bouckova, R., and Yong, V. W. (1994). Reactive astrogliosis in the neonatal mouse brain and its modulation by cytokines. *J. Neurosci.* 14, 846–856.
- Balasingam, V., and Yong, V. W. (1996). Attenuation of astroglial reactivity by interleukin-10. *J. Neurosci.* 16, 2945–2955.
- Bareyre, F. M., and Schwab, M. E. (2003). Inflammation, degeneration and regeneration in the injured spinal cord: insights from DNA microarrays. *Trends Neurosci.* 26, 555–563. doi: 10.1016/j.tins.2003.08.004
- Bazzoni, F., Rossato, M., Fabbri, M., Gaudiosi, D., Mirolo, M., Mori, L., et al. (2009). Induction and regulatory function of miR-9 in human monocytes and neutrophils exposed to proinflammatory signals. *Proc. Natl. Acad. Sci. U.S.A.* 106, 5282–5287. doi: 10.1073/pnas.0810909106
- Beck, K. D., Nguyen, H. X., Galvan, M. D., Salazar, D. L., Woodruff, T. M., and Anderson, A. J. (2010). Quantitative analysis of cellular inflammation after traumatic spinal cord injury: evidence for a multiphasic inflammatory response in the acute to chronic environment. *Brain* 133, 433–447. doi: 10.1093/brain/awp322
- Becker, T., Wullimann, M. F., Becker, C. G., Bernhardt, R. R., and Schachner, M. (1997). Axonal regrowth after spinal cord transection in adult zebrafish. *J. Comp. Neurol.* 377, 577–595. doi: 10.1002/(SICI)1096-9861(19970127)377:4<577::AID-CNE8>3.0.CO;2-#
- Bethea, J. R., Castro, M., Keane, R. W., Lee, T. T., Dietrich, W. D., and Yezierski, R. P. (1998). Traumatic spinal cord injury induces nuclear factor- $\kappa$ B activation. *J. Neurosci.* 18, 3251–3260.
- Bhalala, O. G., Pan, L., Sahni, V., McGuire, T. L., Gruner, K., Tourtellotte, W. G., et al. (2012). microRNA-21 regulates astrocytic response following spinal cord injury. *J. Neurosci.* 32, 17935–17947. doi: 10.1523/JNEUROSCI.3860-12.2012
- Bhalala, O. G., Srikanth, M., and Kessler, J. A. (2013). The emerging roles of microRNAs in CNS injuries. *Nat. Rev. Neurol.* 9, 328–339. doi: 10.1038/nrneuro.2013.67
- Biering-Sorensen, E., Pedersen, V., and Clausen, S. (1990). Epidemiology of spinal cord lesions in Denmark. *Paraplegia* 28, 105–118. doi: 10.1038/sc.1990.13
- Brandenburger, T., Castoldi, M., Brendel, M., Grievink, H., Schlosser, L., Werdehausen, R., et al. (2012). Expression of spinal cord microRNAs in a rat model of chronic neuropathic pain. *Neurosci. Lett.* 506, 281–286. doi: 10.1016/j.neulet.2011.11.023
- Brennan, F. H., Anderson, A. J., Taylor, S. M., Woodruff, T. M., and Ruitenberg, M. J. (2012). Complement activation in the injured central nervous system: another dual-edged sword? *J. Neuroinflammation* 9, 137. doi: 10.1186/1742-2094-9-137
- Buller, B., Liu, X., Wang, X., Zhang, R. L., Zhang, L., Hozeska-Solgot, A., et al. (2010). MicroRNA-21 protects neurons from ischemic death. *FEBS J.* 277, 4299–4307. doi: 10.1111/j.1742-4658.2010.07818.x
- Byrnes, K. R., Garay, J., Di Giovanni, S., De Biase, A., Knobloch, S. M., Hoffman, E. P., et al. (2006). Expression of two temporally distinct microglia-related gene clusters after spinal cord injury. *Glia* 53, 420–433. doi: 10.1002/glia.20295
- Callari, M., Dugo, M., Musella, V., Marchesi, E., Chiorino, G., Grand, M. M., et al. (2012). Comparison of microarray platforms for measuring differential microRNA expression in paired normal/cancer colon tissues. *PLoS ONE* 7:e45105. doi: 10.1371/journal.pone.0045105
- Carmel, J. B., Galante, A., Soteropoulos, P., Tolia, P., Recce, M., Young, W., et al. (2001). Gene expression profiling of acute spinal cord injury reveals spreading inflammatory signals and neuron loss. *Physiol. Genomics* 7, 201–213.
- Carrillo, E. D., Escobar, Y., Gonzalez, G., Hernandez, A., Galindo, J. M., Garcia, M. C., et al. (2011). Posttranscriptional regulation of the  $\beta$ 2-subunit of cardiac L-type  $\text{Ca}^{2+}$  channels by MicroRNAs during long-term exposure to isoproterenol in rats. *J. Cardiovasc. Pharmacol.* 58, 470–478. doi: 10.1097/FJC.0b013e31822a789b
- Casha, S., Yu, W. R., and Fehlings, M. G. (2001). Oligodendroglial apoptosis occurs along degenerating axons and is associated with FAS and p75 expression following spinal cord injury in the rat. *Neuroscience* 103, 203–218. doi: 10.1016/S0304-4522(00)00538-8
- Chamankhah, M., Eftekharpour, E., Karimi-Abdolrezaee, S., Boutros, P. C., San-Marina, S., and Fehlings, M. G. (2013). Genome-wide gene expression profiling of stress response in a spinal cord clip compression injury model. *BMC Genomics* 14:583. doi: 10.1186/1471-2164-14-583
- Chang, T. C., Yu, D., Lee, Y. S., Wentzel, E. A., Arking, D. E., West, K. M., et al. (2008). Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat. Genet.* 40, 43–50. doi: 10.1038/ng.2007.30
- Chen, C. Z., Schaffert, S., Fragoso, R., and Loh, C. (2013). Regulation of immune responses and tolerance: the microRNA perspective. *Immunol. Rev.* 253, 112–128. doi: 10.1111/immr.12060
- Chen, R., Alvero, A. B., Silasi, D. A., Kelly, M. G., Fest, S., Visintin, I., et al. (2008). Regulation of IKK $\beta$  by miR-199a affects NF- $\kappa$ B activity in ovarian cancer cells. *Oncogene* 27, 4712–4723. doi: 10.1038/onc.2008.112
- Chen, X., Liang, H., Zhang, J., Zen, K., and Zhang, C. Y. (2012). Secreted microRNAs: a new form of intercellular communication. *Trends Cell Biol.* 22, 125–132. doi: 10.1016/j.tcb.2011.12.001
- Cimmino, A., Calin, G. A., Fabbri, M., Iorio, M. V., Ferracin, M., Shimizu, M., et al. (2005). miR-15 and miR-16 induce apoptosis by targeting BCL2. *Proc. Natl. Acad. Sci. U.S.A.* 102, 13944–13999. doi: 10.1073/pnas.0506654102
- Citron, B. A., Arnold, P. M., Sebastian, C., Qin, F., Malladi, S., Ameenuddin, S., et al. (2000). Rapid upregulation of caspase-3 in rat spinal cord after injury: mRNA, protein, and cellular localization correlates with apoptotic cell death. *Exp. Neurol.* 166, 213–226. doi: 10.1006/exnr.2000.7523
- Clarke, J., Seo, P., and Clarke, B. (2010). Statistical expression deconvolution from mixed tissue samples. *Bioinformatics* 26, 1043–1049. doi: 10.1093/bioinformatics/btq097
- Cochella, L., and Hobert, O. (2012). Diverse functions of microRNAs in nervous system development. *Curr. Top. Dev. Biol.* 99, 115–143. doi: 10.1016/B978-0-12-387038-4.00005-7
- Crowe, M. J., Bresnahan, J. C., Shuman, S. L., Masters, J. N., and Beattie, M. S. (1997). Apoptosis and delayed degeneration after spinal cord injury in rats and monkeys. *Nat. Med.* 3, 73–76. doi: 10.1038/nm0197-73
- Cui, J. G., Li, Y. Y., Zhao, Y., Bhattacharjee, S., and Lukiw, W. J. (2010). Differential regulation of interleukin-1 receptor-associated kinase-1 (IRAK-1) and IRAK-2 by microRNA-146a and NF- $\kappa$ B in stressed human astroglial cells and in Alzheimer disease. *J. Biol. Chem.* 285, 38951–38960. doi: 10.1074/jbc.M110.178848
- David, S., and Aguayo, A. J. (1981). Axonal elongation into peripheral nervous system “bridges” after central nervous system injury in adult rats. *Science* 214, 931–933. doi: 10.1126/science.6171034
- Davis, B. N., Hilyard, A. C., Nguyen, P. H., Lagna, G., and Hata, A. (2010). Smad proteins bind a conserved RNA sequence to promote microRNA maturation by Drosha. *Mol. Cell* 39, 373–384. doi: 10.1016/j.molcel.2010.07.011
- De Biase, A., Knobloch, S. M., Di Giovanni, S., Fan, C., Molon, A., Hoffman, E. P., et al. (2005). Gene expression profiling of experimental traumatic spinal cord injury as a function of distance from impact site and injury severity. *Physiol. Genomics* 22, 368–381. doi: 10.1152/physiolgenomics.00081.2005
- Deo, M., Yu, J. Y., Chung, K. H., Tippens, M., and Turner, D. L. (2006). Detection of mammalian microRNA expression by in situ hybridization with RNA oligonucleotides. *Dev. Dyn.* 235, 2538–2548. doi: 10.1002/dvdy.20847
- Dharap, A., Bowen, K., Place, R., Li, L. C., and Vemuganti, R. (2009). Transient focal ischemia induces extensive temporal changes in rat cerebral microRNAome. *J. Cereb. Blood Flow Metab.* 29, 675–687. doi: 10.1038/jcbfm.2008.157
- Di Giovanni, S., De Biase, A., Yakovlev, A., Finn, T., Beers, J., Hoffman, E. P., et al. (2005a). *In vivo* and *in vitro* characterization of novel neuronal plasticity

- factors identified following spinal cord injury. *J. Biol. Chem.* 280, 2084–2091. doi: 10.1074/jbc.M411975200
- Di Giovanni, S., Faden, A. I., Yakovlev, A., Duke-Cohan, J. S., Finn, T., Thouin, M., et al. (2005b). Neuronal plasticity after spinal cord injury: identification of a gene cluster driving neurite outgrowth. *FASEB J.* 19, 153–154.
- Di Giovanni, S., Knobloch, S. M., Brandoli, C., Aden, S. A., Hoffman, E. P., and Faden, A. I. (2003). Gene profiling in spinal cord injury shows role of cell cycle in neuronal death. *Ann. Neurol.* 53, 454–468. doi: 10.1002/ana.10472
- Donnelly, D. J., and Popovich, P. G. (2008). Inflammation and its role in neuroprotection, axonal regeneration and functional recovery after spinal cord injury. *Exp. Neurol.* 209, 378–388. doi: 10.1016/j.expneurol.2007.06.009
- Dugas, J. C., Cuellar, T. L., Scholze, A., Ason, B., Ibrahim, A., Emery, B., et al. (2010). Dicer1 and miR-219 Are required for normal oligodendrocyte differentiation and myelination. *Neuron* 65, 597–611. doi: 10.1016/j.neuron.2010.01.027
- Dumont, R. J., Okonkwo, D. O., Verma, S., Hurlbert, R. J., Boulos, P. T., Ellegala, D. B., et al. (2001). Acute spinal cord injury, part I: pathophysiologic mechanisms. *Clin. Neuropharmacol.* 24, 254–264. doi: 10.1097/00002826-200109000-00002
- Ebert, M. S., and Sharp, P. A. (2012). Roles for microRNAs in conferring robustness to biological processes. *Cell* 149, 515–524. doi: 10.1016/j.cell.2012.04.005
- Faulkner, J. R., Herrmann, J. E., Woo, M. J., Tansey, K. E., Doan, N. B., and Sofroniew, M. V. (2004). Reactive astrocytes protect tissue and preserve function after spinal cord injury. *J. Neurosci.* 24, 2143–2155. doi: 10.1523/JNEUROSCI.3547-03.2004
- Fawcett, J. W., and Asher, R. A. (1999). The glial scar and central nervous system repair. *Brain Res. Bull.* 49, 377–391. doi: 10.1016/S0361-9230(99)00072-6
- Fehlings, M. G., and Tator, C. H. (1995). The relationships among the severity of spinal cord injury, residual neurological function, axon counts, and counts of retrogradely labeled neurons after experimental spinal cord injury. *Exp. Neurol.* 132, 220–228. doi: 10.1016/0014-4886(95)90027-6
- Fleming, J. C., Norenberg, M. D., Ramsay, D. A., Dekaban, G. A., Marcillo, A. E., Saenz, A. D., et al. (2006). The cellular inflammatory response in human spinal cords after injury. *Brain* 129, 3249–3269. doi: 10.1093/brain/awl296
- Frankel, L. B., Christoffersen, N. R., Jacobsen, A., Lindow, M., Krogh, A., and Lund, A. H. (2008). Programmed cell death 4 (PDCD4) is an important functional target of the microRNA miR-21 in breast cancer cells. *J. Biol. Chem.* 283, 1026–1033. doi: 10.1074/jbc.M707224200
- Friedman, R. C., Farh, K. K., Burge, C. B., and Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 19, 92–105. doi: 10.1101/gr.082701.108
- Galvan, M. D., Luchetti, S., Burgos, A. M., Nguyen, H. X., Hooshmand, M. J., Hamers, F. P., et al. (2008). Deficiency in complement C1q improves histological and functional locomotor outcome after spinal cord injury. *J. Neurosci.* 28, 13876–13888. doi: 10.1523/JNEUROSCI.2823-08.2008
- Gangaraju, V. K., and Lin, H. (2009). MicroRNAs: key regulators of stem cells. *Nat. Rev. Mol. Cell Biol.* 10, 116–125. doi: 10.1038/nrm2621
- Genda, Y., Arai, M., Ishikawa, M., Tanaka, S., Okabe, T., and Sakamoto, A. (2013). microRNA changes in the dorsal horn of the spinal cord of rats with chronic constriction injury: a TaqMan(R) Low Density Array study. *Int. J. Mol. Med.* 31, 129–137. doi: 10.3892/ijmm.2012.1163
- Git, A., Dvinge, H., Salmon-Divon, M., Osborne, M., Kutter, C., Hadfield, J., et al. (2010). Systematic comparison of microarray profiling, real-time PCR, and next-generation sequencing technologies for measuring differential microRNA expression. *RNA* 16, 991–1006. doi: 10.1261/rna.1947110
- Goddard, D. R., Berry, M., Kirvell, S. L., and Butt, A. M. (2002). Fibroblast growth factor-2 induces astroglial and microglial reactivity *in vivo*. *J. Anat.* 200, 57–67. doi: 10.1046/j.0021-8782.2001.00002.x
- Goldberg, J. L. (2003). How does an axon grow? *Genes Dev.* 17, 941–958. doi: 10.1101/gad.1062303
- Goldie, B. J., and Cairns, M. J. (2012). Post-transcriptional trafficking and regulation of neuronal gene expression. *Mol. Neurobiol.* 45, 99–108. doi: 10.1007/s12035-011-8222-0
- Gosink, M. M., Petrie, H. T., and Tsinoremas, N. F. (2007). Electronically subtracting expression patterns from a mixed cell population. *Bioinformatics* 23, 3328–3334. doi: 10.1093/bioinformatics/btm508
- Grossman, S. D., Rosenberg, L. J., and Wrathall, J. R. (2001). Temporal-spatial pattern of acute neuronal and glial loss after spinal cord contusion. *Exp. Neurol.* 168, 273–282. doi: 10.1006/exnr.2001.7628
- Guedes, J., Cardoso, A. L., and Pedrosa de Lima, M. C. (2013). Involvement of microRNA in microglia-mediated immune response. *Clin. Dev. Immunol.* 2013, 186872. doi: 10.1155/2013/186872
- Guo, H., Ingolia, N. T., Weissman, J. S., and Bartel, D. P. (2010). Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature* 466, 835–840. doi: 10.1038/nature09267
- Hafez, M. M., Hassan, Z. K., Zekri, A. R., Gaber, A. A., Al Rejaie, S. S., Sayed-Ahmed, M. M., et al. (2012). MicroRNAs and metastasis-related gene expression in Egyptian breast cancer patients. *Asian Pac. J. Cancer Prev.* 13, 591–598. doi: 10.7314/APJCP.2012.13.2.591
- Han, M., Liu, M., Wang, Y., Chen, X., Xu, J., Sun, Y., et al. (2012). Antagonism of miR-21 reverses epithelial–mesenchymal transition and cancer stem cell phenotype through AKT/ERK1/2 inactivation by targeting PTEN. *PLoS ONE* 7:e39520. doi: 10.1371/journal.pone.0039520
- Harris, T. A., Yamakuchi, M., Ferlito, M., Mendell, J. T., and Lowenstein, C. J. (2008). MicroRNA-126 regulates endothelial expression of vascular cell adhesion molecule 1. *Proc. Natl. Acad. Sci. U.S.A.* 105, 1516–1521. doi: 10.1073/pnas.0707493105
- Hausmann, O. N. (2003). Post-traumatic inflammation following spinal cord injury. *Spinal Cord* 41, 369–378. doi: 10.1038/sj.sc.3101483
- Hong, J., Zhang, H., Kawase-Koga, Y., and Sun, T. (2013). MicroRNA function is required for neurite outgrowth of mature neurons in the mouse postnatal cerebral cortex. *Front. Cell. Neurosci.* 7:151. doi: 10.3389/fncel.2013.00151
- Hu, J. R., Lv, G. H., and Yin, B. L. (2013a). Altered microRNA expression in the ischemic-reperfusion spinal cord with atorvastatin therapy. *J. Pharmacol. Sci.* 121, 343–346. doi: 10.1254/jphs.122355C
- Hu, J. Z., Huang, J. H., Zeng, L., Wang, G., Cao, M., and Lu, H. B. (2013b). Anti-apoptotic effect of microRNA-21 after contusion spinal cord injury in rats. *J. Neurotrauma* 30, 1349–1360. doi: 10.1089/neu.2012.2748
- Hutchison, E. R., Kawamoto, E. M., Taub, D. D., Lal, A., Abdelmohsen, K., Zhang, Y., et al. (2013). Evidence for miR-181 involvement in neuroinflammatory responses of astrocytes. *Glia* 61, 1018–1028. doi: 10.1002/glia.22483
- Hydbring, P., and Badalian-Very, G. (2013). Clinical applications of microRNAs. *F1000Research* 2, 136. doi: 10.12688/f1000research.2-136.v3
- Iliopoulos, D., Hirsch, H. A., and Struhl, K. (2009). An epigenetic switch involving NF- $\kappa$ B, lin28, Let-7 microRNA, and IL6 links inflammation to cell transformation. *Cell* 139, 693–706. doi: 10.1016/j.cell.2009.10.014
- Iliopoulos, D., Jaeger, S. A., Hirsch, H. A., Bulyk, M. L., and Struhl, K. (2010). STAT3 activation of miR-21 and miR-181b-1 via PTEN and CYLD are part of the epigenetic switch linking inflammation to cancer. *Mol. Cell* 39, 493–506. doi: 10.1016/j.molcel.2010.07.023
- Im, Y. B., Jee, M. K., Choi, J. I., Cho, H. T., Kwon, O. H., and Kang, S. K. (2012). Molecular targeting of NOX4 for neuropathic pain after traumatic injury of the spinal cord. *Cell Death Dis.* 3, e426. doi: 10.1038/cddis.2012.168
- Iyer, A., Zurolo, E., Prabowo, A., Fluiter, K., Splet, W. G., van Rijen, P. C., et al. (2012). MicroRNA-146a: a key regulator of astrocyte-mediated inflammatory response. *PLoS ONE* 7:e44789. doi: 10.1371/journal.pone.0044789
- Izumi, B., Nakasa, T., Tanaka, N., Nakanishi, K., Kamei, N., Yamamoto, R., et al. (2011). MicroRNA-223 expression in neutrophils in the early phase of secondary damage after spinal cord injury. *Neurosci. Lett.* 492, 114–118. doi: 10.1016/j.neulet.2011.01.068
- Jee, M. K., Jung, J. S., Choi, J. I., Jang, J. A., Kang, K. S., Im, Y. B., et al. (2012a). MicroRNA 486 is a potentially novel target for the treatment of spinal cord injury. *Brain* , 135, 1237–1252. doi: 10.1093/brain/awo047
- Jee, M. K., Jung, J. S., Im, Y. B., Jung, S. J., and Kang, S. K. (2012b). Silencing of miR20a is crucial for Ngn1-mediated neuroprotection in injured spinal cord. *Hum. Gene Ther.* 23, 508–520. doi: 10.1089/hum.2011.121
- Jeyaseelan, K., Lim, K. Y., and Armugam, A. (2008). MicroRNA expression in the blood and brain of rats subjected to transient focal ischemia by middle cerebral artery occlusion. *Stroke* 39, 959–966. doi: 10.1161/STROKEAHA.107.500736
- Jia, Z., Zhu, H., Li, J., Wang, X., Misra, H., and Li, Y. (2012). Oxidative stress in spinal cord injury and antioxidant-based intervention. *Spinal Cord* 50, 264–274. doi: 10.1038/sc.2011.111
- Jones, T. B., McDaniel, E. E., and Popovich, P. G. (2005). Inflammatory-mediated injury and repair in the traumatically injured spinal cord. *Curr. Pharm. Des.* 11, 1223–1236. doi: 10.2174/1381612053507468



- Junker, A., Krumbholz, M., Eisele, S., Mohan, H., Augstein, F., Bittner, R., et al. (2009). MicroRNA profiling of multiple sclerosis lesions identifies modulators of the regulatory protein CD47. *Brain* 132, 3342–3352. doi: 10.1093/brain/awp300
- Kaur, P., Armugam, A., and Jayaseelan, K. (2012). MicroRNAs in neurotoxicity. *J. Toxicol.* 2012, 870150. doi: 10.1155/2012/870150
- Kim, Y. K., Yu, J., Han, T. S., Park, S. Y., Namkoong, B., Kim, D. H., et al. (2009). Functional links between clustered microRNAs: suppression of cell-cycle inhibitors by microRNA clusters in gastric cancer. *Nucleic Acids Res.* 37, 1672–1681. doi: 10.1093/nar/gkp002
- Kirschner, M. B., Kao, S. C., Edelman, J. J., Armstrong, N. J., Vallely, M. P., van Zandwijk, N., et al. (2011). Haemolysis during sample preparation alters microRNA content of plasma. *PLoS ONE* 6:e24145. doi: 10.1371/journal.pone.0024145
- Kosaka, N., Iguchi, H., and Ochiya, T. (2010). Circulating microRNA in body fluid: a new potential biomarker for cancer diagnosis and prognosis. *Cancer Sci.* 101, 2087–2092. doi: 10.1111/j.1349-7006.2010.01650.x
- Kosik, K. S. (2006). The neuronal microRNA system. *Nat. Rev. Neurosci.* 7, 911–920. doi: 10.1038/nrn2037
- Kosik, K. S. (2010). MicroRNAs and cellular phenotypy. *Cell* 143, 21–26. doi: 10.1016/j.cell.2010.09.008
- Kozomara, A., and Griffiths-Jones, S. (2011). miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res.* 39, D152–D157. doi: 10.1093/nar/gkq1027
- Krek, A., Grun, D., Poy, M. N., Wolf, R., Rosenberg, L., Epstein, E. J., et al. (2005). Combinatorial microRNA target predictions. *Nat. Genet.* 37, 495–500. doi: 10.1038/ng1536
- Krichevsky, A. M. (2007). MicroRNA profiling: from dark matter to white matter, or identifying new players in neurobiology. *ScientificWorldJournal* 7, 155–166. doi: 10.1100/tsw.2007.201
- Krichevsky, A. M., King, K. S., Donahue, C. P., Khrapko, K., and Kosik, K. S. (2003). A microRNA array reveals extensive regulation of microRNAs during brain development. *RNA* 9, 1274–1281. doi: 10.1261/rna.5980303
- Krichevsky, A. M., Sonntag, K. C., Isacson, O., and Kosik, K. S. (2006). Specific microRNAs modulate embryonic stem cell-derived neurogenesis. *Stem Cells* 24, 857–864. doi: 10.1634/stemcells.2005-0441
- Krol, J., Loedige, I., and Filipowicz, W. (2010). The widespread regulation of microRNA biogenesis, function and decay. *Nat. Rev. Genet.* 11, 597–610. doi: 10.1038/nrg2843
- Kusuda, R., Cadetti, F., Ravanelli, M. I., Sousa, T. A., Zanon, S., De Lucca, F. L., et al. (2011). Differential expression of microRNAs in mouse pain models. *Mol. Pain* 7, 17. doi: 10.1186/1744-8069-7-17
- Laterza, O. F., Lim, L., Garrett-Engle, P. W., Vlasakova, K., Muniappa, N., Tanaka, W. K., et al. (2009). Plasma MicroRNAs as sensitive and specific biomarkers of tissue injury. *Clin. Chem.* 55, 1977–1983. doi: 10.1373/clinchem.2009.131797
- Lau, P., Verrier, J. D., Nielsen, J. A., Johnson, K. R., Notterpek, L., and Hudson, L. D. (2008). Identification of dynamically regulated microRNA and mRNA networks in developing oligodendrocytes. *J. Neurosci.* 28, 11720–11730. doi: 10.1523/JNEUROSCI.1932-08.2008
- Lee, B. B., Cripps, R. A., Fitzharris, M., and Wing, P. C. (2013). The global map for traumatic spinal cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord* 110–116. doi: 10.1038/sc.2012.158
- Lee, E. J., Baek, M., Gusev, Y., Brackett, D. J., Nuovo, G. J., and Schmittgen, T. D. (2008). Systematic evaluation of microRNA processing patterns in tissues, cell lines, and tumors. *RNA* 14, 35–42. doi: 10.1261/rna.804508
- Lee, R. C., Feinbaum, R. L., and Ambros, V. (1993). The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75, 843–854. doi: 10.1016/0092-8674(93)90529-Y
- Li, H., Huang, S., Guo, C., Guan, H., and Xiong, C. (2012). Cell-free seminal mRNA and microRNA exist in different forms. *PLoS ONE* 7:e34566. doi: 10.1371/journal.pone.0034566
- Li, H., Shen, L., Ma, C., and Huang, Y. (2013). Differential expression of miRNAs in the nervous system of a rat model of bilateral sciatic nerve chronic constriction injury. *Int. J. Mol. Med.* 32, 219–226. doi: 10.3892/ijmm.2013.1381
- Li, X., and Jin, P. (2010). Roles of small regulatory RNAs in determining neuronal identity. *Nat. Rev. Neurosci.* 11, 329–338. doi: 10.1038/nrn2739
- Lim, L. P., Lau, N. C., Garrett-Engle, P., Grimson, A., Schelter, J. M., Castle, J., et al. (2005). Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 433, 769–773. doi: 10.1038/nature03315
- Lindsay, M. A. (2008). microRNAs and the immune response. *Trends Immunol.* 29, 343–351. doi: 10.1016/j.it.2008.04.004
- Liu, N. K., Wang, X. E., Lu, Q. B., and Xu, X. M. (2009). Altered microRNA expression following traumatic spinal cord injury. *Exp. Neurol.* 219, 424–429. doi: 10.1016/j.expneurol.2009.06.015
- Liu, X. Z., Xu, X. M., Hu, R., Du, C., Zhang, S. X., McDonald, J. W., et al. (1997). Neuronal and glial apoptosis after traumatic spinal cord injury. *J. Neurosci.* 17, 5395–5406.
- Liu, Y., Chen, Q., Song, Y., Lai, L., Wang, J., Yu, H., et al. (2011). MicroRNA-98 negatively regulates IL-10 production and endotoxin tolerance in macrophages after LPS stimulation. *FEBS Lett.* 585, 1963–1968. doi: 10.1016/j.febslet.2011.05.029
- Liverman, C. T., Altevogt, B. M., Joy, J. E., and Johnson, R. T. (2005). *Spinal Cord Injury: Progress, Promise, and Priorities*. Washington, DC: National Academies Press.
- Lu, J., Getz, G., Miska, E. A., Alvarez-Saavedra, E., Lamb, J., Peck, D., et al. (2005). MicroRNA expression profiles classify human cancers. *Nature* 435, 834–838. doi: 10.1038/nature03702
- Lu, P., Nakorchevskiy, A., and Marcotte, E. M. (2003). Expression deconvolution: a reinterpretation of DNA microarray data reveals dynamic changes in cell populations. *Proc. Natl. Acad. Sci. U.S.A.* 100, 10370–10375. doi: 10.1073/pnas.1832361100
- Ma, X., Becker Buscaglia, L. E., Barker, J. R., and Li, Y. (2010). MicroRNAs in NF- $\kappa$ B signaling. *J. Mol. Cell Biol.* 3, 159–166. doi: 10.1093/jmcb/mjr007
- Makeyev, E. V., Zhang, J., Carrasco, M. A., and Maniatis, T. (2007). The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol. Cell* 27, 435–448. doi: 10.1016/j.molcel.2007.07.015
- Mann, R., Schaefer, C., Sadosky, A., Bergstrom, F., Baik, R., Parsons, B., et al. (2013). Burden of spinal cord injury-related neuropathic pain in the United States: retrospective chart review and cross-sectional survey. *Spinal Cord* 51, 564–570. doi: 10.1038/sc.2013.34
- Mautes, A. E., Weinzierl, M. R., Donovan, F., and Noble, L. J. (2000). Vascular events after spinal cord injury: contribution to secondary pathogenesis. *Phys. Ther.* 80, 673–687.
- McTigue, D. M., Popovich, P. G., Morgan, T. E., and Stokes, B. T. (2000). Localization of transforming growth factor-beta1 and receptor mRNA after experimental spinal cord injury. *Exp. Neurol.* 163, 220–230. doi: 10.1006/exnr.2000.7372
- Merkerova, M., Belickova, M., and Bruchova, H. (2008). Differential expression of microRNAs in hematopoietic cell lineages. *Eur. J. Haematol.* 81, 304–310. doi: 10.1111/j.1600-0609.2008.01111.x
- Mestdagh, P., Bostrom, A. K., Impens, F., Fredlund, E., Van Peer, G., De Antonellis, P., et al. (2010). The miR-17-92 microRNA cluster regulates multiple components of the TGF- $\beta$  pathway in neuroblastoma. *Mol. Cell* 40, 762–773. doi: 10.1016/j.molcel.2010.11.038
- Meza-Sosa, K. F., Valle-Garcia, D., Pedraza-Alva, G., and Perez-Martinez, L. (2012). Role of microRNAs in central nervous system development and pathology. *J. Neurosci. Res.* 90, 1–12. doi: 10.1002/jnr.22701
- Miska, E. A., Alvarez-Saavedra, E., Townsend, M., Yoshii, A., Sestan, N., Rakic, P., et al. (2004). Microarray analysis of microRNA expression in the developing mammalian brain. *Genome Biol.* 5, R68. doi: 10.1186/gb-2004-5-9-r68
- Morel, L., Regan, M., Higashimori, H., Ng, S. K., Esau, C., Vidensky, S., et al. (2013). Neuronal exosomal miRNA-dependent translational regulation of astroglial glutamate transporter GLT1. *J. Biol. Chem.* 288, 7105–7116. doi: 10.1074/jbc.M112.410944
- Nakanishi, K., Nakasa, T., Tanaka, N., Ishikawa, M., Yamada, K., Yamasaki, K., et al. (2010). Responses of microRNAs 124a and 223 following spinal cord injury in mice. *Spinal Cord* 48, 192–196. doi: 10.1038/sc.2009.89
- Oyinbo, C. A. (2011). Secondary injury mechanisms in traumatic spinal cord injury: a nugget of this multiply cascade. *Acta Neurobiol. Exp. (Wars)* 71, 281–299.
- Pekny, M., and Pekna, M. (2004). Astrocyte intermediate filaments in CNS pathologies and regeneration. *J. Pathol.* 204, 428–437. doi: 10.1002/path.1645
- Perri, R., Nares, S., Zhang, S., Barros, S. P., and Offenbacher, S. (2012). MicroRNA modulation in obesity and periodontitis. *J. Dent. Res.* 91, 33–38. doi: 10.1177/0022034511425045
- Pogue, A. I., Cui, J. G., Li, Y. Y., Zhao, Y., Culicchia, F., and Lukiw, W. J. (2010). MicroRNA-125b (miRNA-125b) function in astrogliosis and glial cell proliferation. *Neurosci. Lett.* 476, 18–22. doi: 10.1016/j.neulet.2010.03.054

- Ponomarev, E. D., Veremeyko, T., Barteneva, N., Krichevsky, A. M., and Weiner, H. L. (2011). MicroRNA-124 promotes microglia quiescence and suppresses EAE by deactivating macrophages via the C/EBP- $\alpha$ -PU.1 pathway. *Nat. Med.* 17, 64–70. doi: 10.1038/nm.2266
- Pritchard, C. C., Cheng, H. H., and Tewari, M. (2012). MicroRNA profiling: approaches and considerations. *Nat. Rev. Genet.* 13, 358–369. doi: 10.1038/nrg3198
- Profyris, C., Cheema, S. S., Zang, D., Azari, M. F., Boyle, K., and Petratos, S. (2004). Degenerative and regenerative mechanisms governing spinal cord injury. *Neurobiol. Dis.* 15, 415–436. doi: 10.1016/j.nbd.2003.11.015
- Qiu, J., Nestic, O., Ye, Z., Rea, H., Westlund, K. N., Xu, G. Y., et al. (2001). Bcl-xL expression after contusion to the rat spinal cord. *J. Neurotrauma* 18, 1267–1278. doi: 10.1089/089771501317095304
- Redell, J. B., Moore, A. N., Ward, N. H. III, Hergenroeder, G. W., and Dash, P. K. (2010). Human traumatic brain injury alters plasma microRNA levels. *J. Neurotrauma* 27, 2147–2156. doi: 10.1089/neu.2010.1481
- Rowland, J. W., Hawryluk, G. W., Kwon, B., and Fehlings, M. G. (2008). Current status of acute spinal cord injury pathophysiology and emerging therapies: promise on the horizon. *Neurosurg. Focus* 25, E2. doi: 10.3171/FOC.2008.25.11.E2
- Sahni, V., Mukhopadhyay, A., Tysseling, V., Hebert, A., Birch, D., McGuire, T. L., et al. (2010). BMPR1a and BMPR1b signaling exert opposing effects on gliosis after spinal cord injury. *J. Neurosci.* 30, 1839–1855. doi: 10.1523/JNEUROSCI.4459-09.2010
- Saito, N., Yamamoto, T., Watanabe, T., Abe, Y., and Kumagai, T. (2000). Implications of p53 protein expression in experimental spinal cord injury. *J. Neurotrauma* 17, 173–182. doi: 10.1089/neu.2000.17.173
- Saito, Y., Liang, G., Egger, G., Friedman, J. M., Chuang, J. C., Coetzee, G. A., et al. (2006). Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells. *Cancer Cell* 9, 435–443. doi: 10.1016/j.ccr.2006.04.020
- Santoscoy, C., Rios, C., Franco-Bourland, R. E., Hong, E., Bravo, G., Rojas, G., et al. (2002). Lipid peroxidation by nitric oxide supplements after spinal cord injury: effect of antioxidants in rats. *Neurosci. Lett.* 330, 94–98. doi: 10.1016/S0304-3940(02)00716-4
- Saugstad, J. A. (2010). MicroRNAs as effectors of brain function with roles in ischemia and injury, neuroprotection, and neurodegeneration. *J. Cereb. Blood Flow Metab.* 30, 1564–1576. doi: 10.1038/jcbfm.2010.101
- Saugstad, J. A. (2013). MicroRNAs as effectors of brain function. *Stroke* 44(Suppl. 1), S17–S19. doi: 10.1161/STROKEAHA.113.000985
- Sayed, D., and Abdellatif, M. (2011). MicroRNAs in development and disease. *Physiol. Rev.* 91, 827–887. doi: 10.1152/physrev.00006.2010
- Scholz, J., and Woolf, C. J. (2007). The neuropathic pain triad: neurons, immune cells and glia. *Nat. Neurosci.* 10, 1361–1368. doi: 10.1038/nn1992
- Selbach, M., Schwanhauser, B., Thierfelder, N., Fang, Z., Khanin, R., and Rajewsky, N. (2008). Widespread changes in protein synthesis induced by microRNAs. *Nature* 455, 58–63. doi: 10.1038/nature07228
- Sempere, L. F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004). Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biol.* 5, R13. doi: 10.1186/gb-2004-5-3-r13
- Sharma, A., Kumar, M., Aich, J., Hariharan, M., Brahmachari, S. K., Agrawal, A., et al. (2009). Posttranscriptional regulation of interleukin-10 expression by hsa-miR-106a. *Proc. Natl. Acad. Sci. U.S.A.* 106, 5761–5766. doi: 10.1073/pnas.0808743106
- Sheedy, F. J., Palsson-McDermott, E., Hennessy, E. J., Martin, C., O'Leary, J. J., Ruan, Q., et al. (2010). Negative regulation of TLR4 via targeting of the proinflammatory tumor suppressor PDCD4 by the microRNA miR-21. *Nat. Immunol.* 11, 141–147. doi: 10.1038/ni.1828
- Shin, D., Shin, J. Y., McManus, M. T., Ptacek, L. J., and Fu, Y. H. (2009). Dicer ablation in oligodendrocytes provokes neuronal impairment in mice. *Ann. Neurol.* 66, 843–857. doi: 10.1002/ana.21927
- Shu, J., Kren, B. T., Xia, Z., Wong, P. Y., Li, L., Hanse, E. A., et al. (2011). Genomewide microRNA down-regulation as a negative feedback mechanism in the early phases of liver regeneration. *Hepatology* 54, 609–619. doi: 10.1002/hep.24421
- Silver, J., and Miller, J. H. (2004). Regeneration beyond the glial scar. *Nat. Rev. Neurosci.* 5, 146–156. doi: 10.1038/nrn1326
- Sluijter, J. P. G. (2013). MicroRNAs in cardiovascular regenerative medicine: directing tissue repair and cellular differentiation. *ISRN Vasc. Med.* 2013, 16.
- Smirnova, L., Grafe, A., Seiler, A., Schumacher, S., Nitsch, R., and Wolczyn, F. G. (2005). Regulation of miRNA expression during neural cell specification. *Eur. J. Neurosci.* 21, 1469–1477. doi: 10.1111/j.1460-9568.2005.03978.x
- Smith, B., Treadwell, J., Zhang, D., Ly, D., McKinnell, I., Walker, P. R., et al. (2010). Large-scale expression analysis reveals distinct microRNA profiles at different stages of human neurodevelopment. *PLoS ONE* 5:e11109. doi: 10.1371/journal.pone.0011109
- Sofroniew, M. V. (2005). Reactive astrocytes in neural repair and protection. *Neuroscientist* 11, 400–407. doi: 10.1177/1073858405278321
- Sofroniew, M. V. (2009). Molecular dissection of reactive astrogliosis and glial scar formation. *Trends Neurosci.* 32, 638–647. doi: 10.1016/j.tins.2009.08.002
- Song, G., Cechvala, C., Resnick, D. K., Dempsey, R. J., and Rao, V. L. (2001). GeneChip analysis after acute spinal cord injury in rat. *J. Neurochem.* 79, 804–815. doi: 10.1046/j.1471-4159.2001.00626.x
- Sonkoly, E., and Pivarcsi, A. (2009). microRNAs in inflammation. *Int. Rev. Immunol.* 28, 535–561. doi: 10.3109/08830180903208303
- Stefani, G., and Slack, F. J. (2008). Small non-coding RNAs in animal development. *Nat. Rev. Mol. Cell Biol.* 9, 219–230. doi: 10.1038/nrm2347
- Strickland, E. R., Hook, M. A., Balamanna, S., Huie, J. R., Grau, J. W., and Miranda, R. C. (2011). MicroRNA dysregulation following spinal cord contusion: implications for neural plasticity and repair. *Neuroscience* 186, 146–160. doi: 10.1016/j.neuroscience.2011.03.063
- Sugawara, T., Lewen, A., Gasche, Y., Yu, F., and Chan, P. H. (2002). Overexpression of SOD1 protects vulnerable motor neurons after spinal cord injury by attenuating mitochondrial cytochrome c release. *FASEB J.* 16, 1997–1999.
- Taganov, K. D., Boldin, M. P., Chang, K. J., and Baltimore, D. (2006). NF- $\kappa$ B-dependent induction of microRNA miR-146, an inhibitor targeted to signaling proteins of innate immune responses. *Proc. Natl. Acad. Sci. U.S.A.* 103, 12481–12486. doi: 10.1073/pnas.0605298103
- Tao, J., Wu, H., Lin, Q., Wei, W., Lu, X. H., Cante, J. P., et al. (2011). Deletion of astroglial dicer causes non-cell-autonomous neuronal dysfunction and degeneration. *J. Neurosci.* 31, 8306–8319. doi: 10.1523/JNEUROSCI.0567-11.2011
- Taoka, Y., Okajima, K., Uchiba, M., Murakami, K., Kushimoto, S., Johnno, M., et al. (1997). Role of neutrophils in spinal cord injury in the rat. *Neuroscience* 79, 1177–1182. doi: 10.1016/S0304-4522(97)00011-0
- Tili, E., Michaille, J. J., Cimino, A., Costinean, S., Dumitru, C. D., Adair, B., et al. (2007). Modulation of miR-155 and miR-125b levels following lipopolysaccharide/TNF- $\alpha$  stimulation and their possible roles in regulating the response to endotoxin shock. *J. Immunol.* 179, 5082–5089.
- Treiber, T., Treiber, N., and Meister, G. (2012). Regulation of microRNA biogenesis and function. *Thromb. Haemost.* 107, 605–610. doi: 10.1160/TH11-12-0836
- Tsang, J. S., Ebert, M. S., and van Oudenaarden, A. (2010). Genome-wide dissection of microRNA functions and cotargeting networks using gene set signatures. *Mol. Cell* 38, 140–153. doi: 10.1016/j.molcel.2010.03.007
- Tyor, W. R., Avgeropoulos, N., Ohlandt, G., and Hogan, E. L. (2002). Treatment of spinal cord impact injury in the rat with transforming growth factor- $\beta$ . *J. Neurosci.* 22, 33–41. doi: 10.1016/S0022-510X(02)00113-2
- Velardo, M. J., Burger, C., Williams, P. R., Baker, H. V., Lopez, M. C., Mareci, T. H., et al. (2004). Patterns of gene expression reveal a temporally orchestrated wound healing response in the injured spinal cord. *J. Neurosci.* 24, 8562–8576. doi: 10.1523/JNEUROSCI.3316-04.2004
- Visvanathan, J., Lee, S., Lee, B., Lee, J. W., and Lee, S. K. (2007). The microRNA miR-124 antagonizes the anti-neural REST/SCP1 pathway during embryonic CNS development. *Genes Dev.* 21, 744–749. doi: 10.1101/gad.1519107
- Wang, J., Gu, Z., Ni, P., Qiao, Y., Chen, C., Liu, X., et al. (2011). NF- $\kappa$ B P50/P65 hetero-dimer mediates differential regulation of CD166/ALCAM expression via interaction with microRNA-9 after serum deprivation, providing evidence for a novel negative auto-regulatory loop. *Nucleic Acids Res.* 39, 6440–6455. doi: 10.1093/nar/gkr302
- Wang, M., Master, S. R., and Chodosh, L. A. (2006). Computational expression deconvolution in a complex mammalian organ. *BMC Bioinformatics* 7:328. doi: 10.1186/1471-2105-7-328
- Wang, Z. (2010). MicroRNA: a matter of life or death. *World J. Biol. Chem.* 1, 41–54. doi: 10.4331/wjbc.v1.i4.41
- Willemsen, H. L., Huo, X. J., Mao-Ying, Q. L., Zijlstra, J., Heijnen, C. J., and Kavelaars, A. (2012). MicroRNA-124 as a novel treatment for persistent hyperalgesia. *J. Neuroinflammation* 9, 143. doi: 10.1186/1742-2094-9-143

- Wu, C. I., Shen, Y., and Tang, T. (2009). Evolution under canalization and the dual roles of microRNAs: a hypothesis. *Genome Res.* 19, 734–743. doi: 10.1101/gr.084640.108
- Wu, H., Neilson, J. R., Kumar, P., Manocha, M., Shankar, P., Sharp, P. A., et al. (2007). miRNA profiling of naive, effector and memory CD8 T cells. *PLoS ONE* 2:e1020. doi: 10.1371/journal.pone.0001020
- Wu, X., Yoo, S., and Wrathall, J. R. (2005). Real-time quantitative PCR analysis of temporal-spatial alterations in gene expression after spinal cord contusion. *J. Neurochem.* 93, 943–952. doi: 10.1111/j.1471-4159.2005.03078.x
- Xiao, Q., Du, Y., Wu, W., and Yip, H. K. (2010). Bone morphogenetic proteins mediate cellular response and, together with Noggin, regulate astrocyte differentiation after spinal cord injury. *Exp. Neurol.* 221, 353–366. doi: 10.1016/j.expneurol.2009.12.003
- Yang, L., Jones, N. R., Blumbergs, P. C., Van Den Heuvel, C., Moore, E. J., Manavis, J., et al. (2005). Severity-dependent expression of pro-inflammatory cytokines in traumatic spinal cord injury in the rat. *J. Clin. Neurosci.* 12, 276–284. doi: 10.1016/j.jocn.2004.06.011
- Young, M. R., Santhanam, A. N., Yoshikawa, N., and Colburn, N. H. (2010). Have tumor suppressor PDCD4 and its counteragent oncogenic miR-21 gone rogue? *Mol. Interv.* 10, 76–79. doi: 10.1124/mi.10.2.5
- Yu, J. Y., Chung, K. H., Deo, M., Thompson, R. C., and Turner, D. L. (2008). MicroRNA miR-124 regulates neurite outgrowth during neuronal differentiation. *Exp. Cell Res.* 314, 2618–2633. doi: 10.1016/j.yexcr.2008.06.002
- Yu, Y. M., Gibbs, K. M., Davila, J., Campbell, N., Sung, S., Todorova, T. I., et al. (2011). MicroRNA miR-133b is essential for functional recovery after spinal cord injury in adult zebrafish. *Eur. J. Neurosci.* 33, 1587–1597. doi: 10.1111/j.1460-9568.2011.07643.x
- Yunta, M., Nieto-Diaz, M., Esteban, F. J., Caballero-Lopez, M., Navarro-Ruiz, R., Reigada, D., et al. (2012). MicroRNA dysregulation in the spinal cord following traumatic injury. *PLoS ONE* 7:e34534. doi: 10.1371/journal.pone.0034534
- Zamanian, J. L., Xu, L., Foo, L. C., Nouri, N., Zhou, L., Giffard, R. G., et al. (2012). Genomic analysis of reactive astrogliosis. *J. Neurosci.* 32, 6391–6410. doi: 10.1523/JNEUROSCI.6221-11.2012
- Zampetaki, A., and Mayr, M. (2012). Analytical challenges and technical limitations in assessing circulating miRNAs. *Thromb. Haemost.* 108, 592–598. doi: 10.1160/TH12-02-0097
- Zhang, K. H., Xiao, H. S., Lu, P. H., Shi, J., Li, G. D., Wang, Y. T., et al. (2004). Differential gene expression after complete spinal cord transection in adult rats: an analysis focused on a subchronic post-injury stage. *Neuroscience* 128, 375–388. doi: 10.1016/j.neuroscience.2004.07.008
- Zhang, Y., Liao, Y., Wang, D., He, Y., Cao, D., Zhang, F., et al. (2011). Altered expression levels of miRNAs in serum as sensitive biomarkers for early diagnosis of traumatic injury. *J. Cell. Biochem.* 112, 2435–2442. doi: 10.1002/jcb.23168
- Zhao, C., Sun, G., Li, S., Lang, M. F., Yang, S., Li, W., et al. (2010a). MicroRNA let-7b regulates neural stem cell proliferation and differentiation by targeting nuclear receptor TLX signaling. *Proc. Natl. Acad. Sci. U.S.A.* 107, 1876–1881. doi: 10.1073/pnas.0908750107
- Zhao, X., He, X., Han, X., Yu, Y., Ye, F., Chen, Y., et al. (2010b). MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* 65, 612–626. doi: 10.1016/j.neuron.2010.02.018
- Zhu, H., and Fan, G. C. (2011). Extracellular/circulating microRNAs and their potential role in cardiovascular disease. *Am. J. Cardiovasc. Dis.* 1, 138–149.
- Ziu, M., Fletcher, L., Savage, J. G., Jimenez, D. F., Digicaylioglu, M., and Bartanusz, V. (2013). Spatial and temporal expression levels of specific microRNAs in a spinal cord injury mouse model and their relationship to the duration of compression. *Spine J.* 14, 353–360. doi: 10.1016/j.spinee.2013.08.015

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 15 December 2013; accepted: 06 February 2014; published online: 25 February 2014.

Citation: Nieto-Diaz M, Esteban FJ, Reigada D, Muñoz-Galdeano T, Yunta M, Caballero-López M, Navarro-Ruiz R, del Águila Á and Maza RM (2014) MicroRNA dysregulation in spinal cord injury: causes, consequences, and therapeutics. *Front. Cell. Neurosci.* 8:53. doi: 10.3389/fncel.2014.00053

This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2014 Nieto-Diaz, Esteban, Reigada, Muñoz-Galdeano, Yunta, Caballero-López, Navarro-Ruiz, del Águila and Maza. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.