

MicroRNA profiling in human diploid fibroblasts uncovers miR-519 role in replicative senescence

Bernard S. Marasa^{1,2}, Subramanya Srikantan¹, Jennifer L. Martindale¹, Mihee M. Kim¹, Eun Kyung Lee¹, Myriam Gorospe¹, and Kotb Abdelmohsen¹

¹ *Laboratory of Cellular and Molecular Biology, NIA-IRP, NIH, Baltimore, MD 21224, USA*

² *Department of Biology, The Catholic University of America, Washington, DC 20064, USA*

Key words: HuR, miR-519, tumor suppression

Abbreviations: CR, coding region; HDF, human diploid fibroblasts; RISC, RNA-induced silencing complex; RBP, RNA-binding protein; SA, senescence-associated; UTR, untranslated region

Received: 05/21/10; **accepted:** 06/17/10; **published on line:** 06/19/10

Correspondence: abdelmohsenk@grc.nia.nih.gov

Copyright: © Marasa et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract: MicroRNAs (miRNAs) are short non-coding RNAs that regulate diverse biological processes by controlling the pattern of expressed proteins. In mammalian cells, miRNAs partially complement their target sequences leading to mRNA degradation and/or decreased mRNA translation. Here, we have analyzed transcriptome-wide changes in miRNAs in senescent relative to early-passage WI-38 human diploid fibroblasts (HDFs). Among the miRNAs downregulated with senescence were members of the let-7 family, while upregulated miRNAs included miR-1204, miR-663 and miR-519. miR-519 was recently found to reduce tumor growth at least in part by lowering the abundance of the RNA-binding protein HuR. Overexpression of miR-519a in either WI-38 or human cervical carcinoma HeLa cells triggered senescence, as measured by monitoring β -galactosidase activity and other senescence markers. These data suggest that miR-519 can suppress tumor growth by triggering senescence and that miR-519 elicits these actions by repressing HuR expression.

INTRODUCTION

MicroRNAs are short (~ 22-nt) RNA molecules that modulate changes in gene expression [1,2]. They are generated from precursor transcripts (primary microRNAs) which are exported to the cytoplasm and are cleaved by Dicer; mature miRNAs then assemble into ribonucleoprotein silencing complexes (RISC) that are recruited to specific mRNAs [3]. MicroRNAs function primarily as repressors of mRNA stability and translation [4]. Through their influence on the patterns of expressed genes, microRNAs have been implicated in numerous physiologic processes, such as development of the muscular, immune, neuronal, epithelial and other systems, and in pathologies including neurodegeneration and cancer [5-8]. The latter studies have revealed a number of miRNAs that can function as tu-

mor suppressors (TS-miRNAs) or tumor promoters (oncomiRs) [9].

Cellular senescence is achieved when cells reach the end of their replicative lifespan [10,11]. It is believed to represent a tumor-suppressive mechanism and a contributing factor in aging [12,13]. MicroRNAs have been implicated in replicative senescence, since loss of miRNA biogenesis through Dicer ablation causes senescence in primary cells [14]. Several specific miRNAs were reported to be differentially expressed in senescent cells compared to young, proliferating cells. For example, miRNA-146a and miR-146b are up-regulated in senescent cells and modulate inflammatory responses by suppressing secretion of IL-6 and IL-8 and by downregulating IRAK1 [15].

Recently, four microRNAs (miR-15b, miR-24, miR-25, and miR-141) that jointly lower expression of the kinase MKK4 were found to decline during replicative senescence and to contribute to the senescence process [16]. miR-24 was also found to regulate translation of the cyclin-dependent kinase inhibitor p16, thereby allowing increased p16 expression in senescent cells [17].

Several miRNAs differentially expressed with aging have also been identified. For example, miR-17, miR-19b, miR-20a, and miR-106a were less abundant in cells from older humans [18]. Reduced expression of miR-103, miR-107, miR-128, miR-130a, miR-155, miR-24, miR-221, miR-496, and miR-1538 in older individuals was also recently reported [19]. Age-regulated changes in the expression of microRNAs were also found in mouse liver and brain [20,21]. MicroRNA changes in Ames dwarf mouse liver led to the identification of microRNAs that might delay aging [22]. Studies in *Caenorhabditis elegans* revealed that the microRNA *lin-4* represses *lin-14* transcripts and *lin-14* protein to extend lifespan by reducing DAF-16; miRNA profiling in *C. elegans* provided evidence that microRNAs may potentially influence the biology of aging [23-25].

Many studies have focused on the role of microRNAs in tumorigenesis and age-related diseases. Here, we have studied changes in expressed microRNAs during repli-

cative senescence of WI-38 human diploid fibroblasts (HDFs). We identified subsets of microRNAs that were differentially expressed in young compared with senescent WI-38 cells. miR-519, a microRNA that suppresses tumorigenesis and lowers expression of RNA-binding protein HuR, was upregulated in senescent cells. Overexpression of miR-519 induced senescence in WI-38 and HeLa cells. Our data support the hypothesis that senescence-associated changes in microRNA expression patterns can affect the susceptibility to age-related diseases such as cancer.

RESULTS

Global changes in microRNAs between early-passage and senescent WI-38 human diploid fibroblasts

Compared with early-passage, 'young' proliferating [Y, at population doubling (pdl) 22] WI-38 cells, the senescent (S, pdl 52) WI-38 cells displayed a flattened morphology and senescence-associated (SA) β -galactosidase (SA- β -gal) activity, a widely used senescence marker [26,27] (Figure 1A). Western blot analysis also revealed that senescent cells expressed lower levels of SIRT1 and HuR, whereas p16 and p53 were upregulated (Figure 1B), in keeping with reported literature [28-30].




Figure 1. Characterization of early-passage and senescent WI-38 cells. (A) Micrographs illustrating β -galactosidase activity in young (Y) early-passage (pdl 22) and senescent (S), late-passage (pdl 52) WI-38 cells. **(B)** Western blot analysis of the proteins indicated in whole-cell lysates prepared from Y and S WI-38 populations; β -actin served as a loading control.

To test how the pattern of expressed microRNAs is affected by replicative senescence, we studied transcriptome-wide changes in microRNAs using miRNome arrays (not shown); we then validated individual microRNAs by reverse transcription (RT) followed by real-time, quantitative (q)PCR amplification (see Materials and Methods). Depicted in Figures 2 and 3 and in supplemental Table S1 are all of the microRNAs

validated using sequence-specific qPCR primers. As shown in Figure 2, several microRNAs were markedly more abundant in senescent cells (e.g., miR-1204, miR-663, miR-548b-3p and miR-431). Other microRNAs were expressed at lower levels in senescent cells [e.g., miR-24, miR-141, and miR-10a (Figure 3, Supplemental Table S1)]. MicroRNAs changing less than twofold with senescence are listed in the supplemental Figure S1.




Figure 2. MicroRNAs upregulated in senescent cells. RNA extracted from Y (pdl 22-25) and S (pdl 50-55) WI-38 cells was used to measure the levels of the microRNAs listed, using RT-qPCR (Materials and Methods). MicroRNA abundance was normalized to U1 snRNA levels. Data are the means and S.D. from three independent experiments.




Figure 3. MicroRNAs downregulated in senescent cells. RNA was extracted and analyzed as explained in the legend of Figure 2. Data show the means and S.D. from three independent experiments.




Figure 4. Influence of miR-519 on WI-38 senescence. (A) Fold differences in miR-519 expression in S relative to Y cells, calculated as explained in the legend of Figure 2. (B) Forty-eight h after transfection of either control (Ctrl) siRNA or miR-519a, the levels of miR-519a were measured by RT-qPCR. (C,D) In cells transfected as explained in panel (B), the levels of HuR protein and loading control GAPDH were assessed by Western blot analysis (C), and the levels of *HuR* mRNA and normalization control 18S rRNA were measured by RT-qPCR (D). (E) WI-38 cell numbers in cells transfected as in (B) were counted every 7 days. (F) SA- β -galactosidase activity in WI-38 cells by week 4 after sequential transfection (every 7 days) of either Ctrl siRNA or miR-519a. (G) Western blot analysis of p27 and loading control GAPDH in Y and S WI-38 cells (*top*) or in WI-38 cells by week 4 after transfection as explained in (E) (*bottom*). The data in B,D,E represent the means and S.D. from three independent experiments.

miR-519-induced senescence in HDFs

We were particularly interested in the miR-519 family. miR-519 was recently found to inhibit translation of the RNA-binding protein HuR through its interaction with the HuR coding region [31]. In a separate study, miR-519 suppressed the growth of tumor xenografts in an HuR-dependent manner [32]. Given that HuR promotes cell proliferation and decreases senescence [33,34], we hypothesized that the elevated miR-519 in senescent cells (Figure 4A) might lower HuR expression in WI-38 HDFs, and hence promote senescence. To test this possibility, we overexpressed miR-519a in young-HDFs (Figure 4B); western blot analysis confirmed that miR-519a overexpression repressed HuR (Figure 4C). In

keeping with earlier results [31], miR-519a did not influence the levels of *HuR* mRNA (Figure 4D), in agreement with the view that miR-519a inhibited *HuR* mRNA translation without affecting *HuR* mRNA stability. Moreover, sustained miR-519a overexpression for 4 weeks caused a marked reduction in cell number as compared to control transfection groups (Figure 4E). miR-519a-overexpressing cells also showed increased SA- β -gal activity (Figure 4F) and elevated expression of the senescence marker p27 [35,36] (Figure 4G, *bottom*). Together, these data indicate that miR-519a induced cellular senescence and inhibited cell proliferation, resulting in accelerated senescence. They further suggest that miR-519a-induced senescence may be mediated in part by repression of HuR.




Figure 5. Influence of miR-519 on the senescent phenotype of HeLa cells. (A) Forty-eight h after transfection of HeLa cells with either control (Ctrl) siRNA or miR-519a, miR-519a levels were measured by RT-qPCR. (B) Number of HeLa cells remaining by 72 h after transfection of Ctrl siRNA or miR-519a as explained in (A). (C) β -galactosidase activity in HeLa cells 5 days after transfection with either Ctrl siRNA or miR-519a. (D) Seventy-two hours after transfection as indicated in (A), the levels of the proteins shown were assessed by Western blot analysis. The data in A,B represent the means and S.D. from three independent experiments.

miR-519-induced senescence in HeLa cells

As indicated above, miR-519 was found to suppress tumor growth [32]. Since cellular senescence is considered to be an anti-tumorigenic process, we examined the effect of miR-519 on the senescent phenotype of cancer cells. Upon miR-519a overexpression (Figure 5A), HeLa cell numbers declined significantly (Figure 5B). Five days after transfection of miR-519a, cells showed a strong increase in SA- β -gal activity compared to the control transfection group (Figure 5C); in addition, miR-519-induced senescence in HeLa cells was accompanied by increased levels of the senescence markers p53 and p27 (Figure 5D). Together, these data indicate that miR-519 reduced HeLa cell proliferation and promoted HeLa cell senescence. Accordingly, we postulate that one of the mechanisms by which miR-519 suppress tumor growth is by inducing senescence, and further propose that miR-519 triggers senescence—at least in part—by reducing HuR levels.

DISCUSSION

Cells become senescent as a result of factors such as the accumulation of reactive oxygen species, DNA damage, erosion of telomeres, and oncogenic activation. Collectively, these triggers cause cells to undergo morphological changes, to become unable to replicate DNA and to display altered gene expression patterns [10-13]. Here, we investigated microRNA levels in WI-38 human diploid fibroblasts by comparing microRNA patterns in senescent relative to young, proliferating cells. Among the microRNAs showing increasing abundance with senescence, miR-519 was of particular interest because it was shown to inhibit translation of HuR and to diminish tumor growth [31,32]. Through its influence on the expression of many genes, HuR plays a key role in cell proliferation, tumorigenesis, and senescence [37,38]. We found that overexpression of miR-519a decreased HuR levels, lowered cell proliferation, and promoted replicative senescence in both WI-38 and HeLa cells.

microRNAs and senescence

We previously used miRNA microarrays to identify changes in a limited number of microRNAs in senescent cells [16]. Here, we have expanded this analysis and have verified many individual microRNAs whose abundance changes with replicative senescence. Many of them target key proteins implicated in senescence and cancer. For example, miR-146b is upregulated in senescent cells (Figure 2), in keeping with earlier findings that miR-146a and miR-146b increased with

senescence and repressed the senescence-associated inflammatory mediators IL-6 and IL-8 [15]. miR-34a regulates SIRT1 expression and induced senescence of cancer cells [39-41]; here, we observed higher miR-34c (not miR-34a) in senescent cells, likely a reflection of the variability and complexity of the senescence process. Several let-7 members were also upregulated in senescent cells (Figure 2 and Supplemental Figure S1); this observation supports the view that the ability of let-7 microRNAs can suppress tumor growth [reviewed in 42], which could contribute to the senescence process. Similarly, upregulation of miR-20 in senescence cells correlates with the ability of miR-20 to inhibit proliferation of K562 human erythromyeloblastoid leukemia cells [43]. In conjunction with the finding that miR-519 reduced tumorigenesis in a xenograft model [32], we propose that the coordinated action of senescence-upregulated microRNAs can suppress tumor growth by reducing the levels of oncogenes or tumor promoters.

Conversely, many microRNAs were downregulated in senescent cells (Figure 3). Among the myriad of senescence-associated proteins that they might regulate, these microRNAs likely repress several tumor suppressors. In this regard, as miR-21 has been shown to lower expression of the tumor suppressor PTEN [44], the downregulated of miR-21 in senescent cells (Figure 3) could allow increased PTEN expression, in turn reducing tumor cell proliferation, migration, and invasion [45].

miR-519a-induced senescence by lowering HuR

We previously reported that miR-519 represses the production of HuR, an RNA-binding protein which is highly abundant in cancer cells and is low in untransformed cells [11,38]. HuR overexpression delays the senescent phenotype while the loss of HuR enhances it [11]. Moreover, while HuR levels are high in tumors and low in normal tissues, miR-519 levels are high in normal tissues and low in cancer tissues [32]. Since HuR potently enhances the expression of cancer-promoting proteins, and reducing HuR levels promotes HDF senescence [11,38], we propose that miR-519 represses tumor growth at least in part, by lowering HuR and thereby promoting senescence (Figs. 4 and 5). Additionally, miR-519 could further repress tumor growth by lowering the expression of other genes, such as ABCG2 or HIF-1 α [46,47].

In summary, we have identified collections of microRNAs displaying altered abundance with replicative senescence. As shown here for miR-519, we postulate that these changes help to meet the needs of

senescent cells in eliciting tumor suppression and growth arrest. Future studies will help to recognize more fully the proteins and processes modulated by senescence-regulated microRNAs.

MATERIAL AND METHODS

Cell culture, transfections, and β -galactosidase staining. Early-passage, proliferating ('young', ~20 to 30 pdl) and late-passage, senescent (~50 to 55 pdl) WI-38 human diploid fibroblasts (HDFs; Coriell Cell Repositories) were cultured in Dulbecco's modified Eagle's medium (DMEM, Invitrogen) supplemented with 10% fetal bovine serum and 0.1 mM nonessential amino acids (Invitrogen). HeLa cells were cultured in DMEM supplemented with 10% FBS and antibiotics. miR-519a (Ambion) or control siRNA (AATTCTCCGAACGTGTCACGT, Qiagen) were transfected at a final concentration of 100 nM using Lipofectamine 2000 (Invitrogen). Where indicated, transfections were performed every 7 days for 4 weeks. WI-38 HDFs and HeLa cells were stained with a senescence-associated β -galactosidase (Cell Signaling Technology) detection kit, according to the manufacturer's protocol.

RNA isolation and miRNA profiling. Total cellular RNA was isolated using Trizol (Invitrogen). Isolated RNA was used to measure miRNA levels in young and senescent cells with a 7900HT real-time PCR instrument (Applied Biosystems). All microRNAs were measured and validated using miRNA-specific forward primers (Supplemental Table S2) and a universal reverse primer (System Biosciences, SBI), according to the manufacturer's protocol. The levels of U1 snRNA, used for normalization, were determined using the specific forward primer CGACTGCATAATTTGTGGTAGTGG.

Protein analysis. Whole-cell lysates were prepared with RIPA buffer [10 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1% NP-40, 1 mM EDTA, 0.1% SDS, and 1 mM dithiothreitol]. Proteins were resolved by SDS-polyacrylamide gel electrophoresis and transferred to polyvinylidene difluoride membranes (Invitrogen). After incubation with primary antibodies recognizing SIRT1, HuR, p16, p53, p27, GAPDH (all from Santa Cruz Biotechnology) or β -actin (Abcam), blots were incubated with the appropriate secondary antibodies and the signals were detected by ECL Plus (GE Healthcare).

ACKNOWLEDGEMENTS

This research was supported in full by the National Institute on Aging-Intramural Research Program, National Institutes of Health.

CONFLICT OF INTERESTS STATEMENT

The authors of this manuscript have no conflict of interests to declare.

REFERENCES

1. Chekulaeva M, Filipowicz W. Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells. *Curr. Opin. Cell Biol.* 2009; 21:452-460.
2. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009; 8:215-233.
3. Winter J, Jung S, Keller S, Gregory RI, Diederichs S. Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat. Cell. Biol.* 2009; 11:228-234.
4. Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R. Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev* 2006; 20:515-524.
5. Ambros V. The functions of animal microRNAs. *Nature* 2004; 431:350-355.
6. Kloosterman WP, Plasterk RH. The diverse functions of microRNAs in animal development and disease. *Dev. Cell* 2006; 11:441-450.
7. Baltimore D, Boldin MP, O'Connell RM, Rao DS, Taganov KD. MicroRNAs: new regulators of immune cell development and function. *Nat. Immunol.* 2008; 9:839-845.
8. Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat. Rev. Mol. Cell. Biol.* 2008; 9:219-230.
9. Croce CM. Causes and consequences of microRNA dysregulation in cancer. *Nat. Rev. Genet.* 2009; 10:704-714.
10. Hayflick L. The Limited In Vitro Lifetime of Human Diploid Cell Strains. *Exp. Cell Res.* 1965; 37: 614-636.
11. Cristofalo VJ, Lorenzini A, Allen RG, Torres C, Tresini M. Replicative senescence: A critical review. *Mech. Ageing Dev.* 2004; 125:827-848.
12. Smith JR, Pereira-Smith OM. Replicative senescence: Implications for in vivo aging and tumor suppression. *Science* 1996; 273:63-67.
13. Campisi J. Senescent cells, tumor suppression, and organismal aging: Good citizens, bad neighbors. *Cell* 2005; 120:513-522.
14. Mudhasani R, Zhu Z, Hutvagner G, Eischen CM, Lyle S, Hall LL, Lawrence JB, Imbalzano AN, Jones SN. Loss of miRNA biogenesis induces p19Arf-p53 signaling and senescence in primary cells. *J. Cell Biol.* 2008; 181:1055-1063.
15. Bhaumik D, Scott GK, Schokrpur S, Patil CK, Orjalo AV, Rodier F, Lithgow GJ, Campisi J. MicroRNAs miR-146a/b negatively modulate the senescence-associated inflammatory mediators IL-6 and IL-8. *Aging (Albany NY)* 2009; 1:402-411.
16. Marasa BS, Srikantan S, Masuda K, Abdelmohsen K, Kuwano Y, Yang X, Martindale JL, Rinker-Schaeffer CW, Gorospe M. Increased MKK4 abundance with replicative senescence is linked to the joint reduction of multiple microRNAs. *Sci Signal* 2009; 2: ra69.
17. Lal A, Kim HH, Abdelmohsen K, Kuwano Y, Pullmann R Jr, Srikantan S, Subrahmanyam R, Martindale JL, Yang X, Ahmed F, Navarro F, Dykxhoorn D, Lieberman J, Gorospe M. p16(INK4a) translation suppressed by miR-24. *PLoS ONE* 2008; 3:p.e1864.
18. Hackl M, Brunner S, Fortschegger K, Schreiner C, Micutkova L, et al., miR-17, miR-19b, miR-20a, and miR-106a are down-regulated in human aging. *Aging Cell* 2010; 9:291-296.

19. Noren Hooten N, Abdelmohsen K, Gorospe M, Ejiogu N, Zonderman AB, and Evans MK. microRNA expression patterns reveal differential expression of target genes with age. *PLoS ONE* 2010, in press.
20. Maes OC, An J, Sarojini H, Wang E. Murine microRNAs implicated in liver functions and aging process. *Mech. Ageing Dev.* 2008; 129:534-541.
21. Li N, Bates DJ, An J, Terry DA, Wang E. Up-regulation of key microRNAs, and inverse down-regulation of their predicted oxidative phosphorylation target genes, during aging in mouse brain. *Neurobiol. Aging* 2009; [epub]
22. Bates DJ, Li N, Liang R, Sarojini H, An J, Masternak MM, Bartke A, Wang E. MicroRNA regulation in Ames dwarf mouse liver may contribute to delayed aging. *Aging Cell* 2009; 2010; 9:1-18.
23. Kato M, Slack FJ. microRNAs: small molecules with big roles - C. elegans to human cancer. *Biol. Cell* 2008; 100:71-81.
24. Boehm M, Slack F. A developmental timing microRNA and its target regulate life span in C. elegans. *Science* 2005; 310:1954-1957.
25. Ibanez-Ventoso C, Yang M, Guo S, Robins H, Padgett RW, Driscoll M. Modulated microRNA expression during adult lifespan in *Caenorhabditis elegans*. *Aging Cell* 2006; 5:235-246.
26. Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I, Pereira-Smith O, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc. Natl. Acad. Sci. USA* 1995;92:9363-9367.
27. Debacq-Chainiaux F, Erusalimsky JD, Campisi J, Toussaint O. Protocols to detect senescence-associated beta-galactosidase (SA-beta-gal) activity, a biomarker of senescent cells in culture and in vivo. *Nat. Protoc.* 2009; 4:1798-806.
28. Abdelmohsen K, Pullmann R Jr, Lal A, Kim HH, Galban S, Yang X, Blethrow JD, Walker M, Shubert J, Gillespie DA, Furneaux H, Gorospe M. Phosphorylation of HuR by Chk2 regulates SIRT1 expression. *Mol. Cell* 2007; 25:543-557.
29. Rodier F, Coppé JP, Patil CK, Hoeijmakers WA, Muñoz DP, Raza SR, Freund A, Campeau E, Davalos AR, Campisi J. Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat. Cell Biol.* 2009; 11:973-979.
30. McConnell BB, Starborg M, Brookes S, Peters G. Inhibitors of cyclin-dependent kinases induce features of replicative senescence in early passage human diploid fibroblasts. *Curr. Biol.* 1998; 8:351-354.
31. Abdelmohsen K, Srikantan S, Kuwano Y, Gorospe M. miR-519 reduces cell proliferation by lowering RNA-binding protein HuR levels. *Proc. Natl. Acad. Sci. USA* 2008; 105: 20297-20302.
32. Abdelmohsen K, Kim MM, Srikantan S, Mercken EM, Brennan SE, Wilson GM, de Cabo R, Gorospe M. miR-519 suppresses tumor growth by reducing HuR levels. *Cell Cycle* 2010; 9 [epub].
33. Yi J, Chang N, Liu X, Guo G, Xue L, Tong T, Gorospe M, Wang W. Reduced nuclear export of HuR mRNA by HuR is linked to the loss of HuR in replicative senescence. *Nucleic Acids Res.* 2010; 38:1547-1558.
34. Wang W, Yang X, Cristofalo VJ, Holbrook NJ, Gorospe M. Loss of HuR is linked to reduced expression of proliferative genes during replicative senescence. *Mol. Cell. Biol.* 2001; 21:5889-5898.
35. Alexander K, Hinds PW. Requirement for p27(KIP1) in retinoblastoma protein-mediated senescence. *Mol. Cell. Biol.* 2001; 21:3616-3631.
36. Zeng J, Wang L, Li Q, Li W, Björkholm M, Jia J, Xu D. FoxM1 is up-regulated in gastric cancer and its inhibition leads to cellular senescence, partially dependent on p27 kip1. *J. Pathol.* 2009; 218:419-427.
37. Hinman MN, Lou H. Diverse molecular functions of Hu proteins. *Cell Mol. Life Sci.* 2008; 65:3168-3181.
38. Abdelmohsen K, Gorospe M. Post-transcriptional regulation of cancer traits by HuR. *WIRES RNA* 2010; in press.
39. Christoffersen NR, Shalgi R, Frankel LB, Leucci E, Lees M, Klausen M, Pilpel Y, Nielsen FC, Oren M, Lund AH. p53-independent upregulation of miR-34a during oncogene-induced senescence represses MYC. *Cell Death Differ.* 2010; 17:236-245.
40. Tazawa H, Tsuchiya N, Izumiya M, Nakagama H. Tumor-suppressive miR-34a induces senescence-like growth arrest through modulation of the E2F pathway in human colon cancer cells. *Proc. Natl. Acad. Sci. USA* 2007; 104:15472-15477.
41. Yamakuchi M, Ferlito M, and Lowenstein CJ. miR-34a repression of SIRT1 regulates apoptosis. *Proc. Natl. Acad. Sci. USA* 2008; 105:13421-13426.
42. Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor suppressors. *Dev Biol.* 2007; 302:1-12.
43. Scherr M, Venturini L, Battmer K, Schaller-Schoenitz M, Schaefer D, Dallmann I, Ganser A, Eder M. Lentivirus-mediated antagomir expression for specific inhibition of miRNA function. *Nucleic Acids Res.* 2007;35; e149.
44. Meng F, Henson R, Wehbe-Janek H, Ghoshal K, Jacob ST, Patel T. MicroRNA-21 regulates expression of the PTEN tumor suppressor gene in human hepatocellular cancer. *Gastroenterology* 2007; 133:647-658.
45. Salmena L, Carracedo A, Pandolfi PP. Tenets of PTEN tumor suppression. *Cell* 2008; 133:403-414.
46. To KK, Robey RW, Knutsen T, Zhan Z, Ried T, Bates SE. Escape from hsa-miR-519c enables drug-resistant cells to maintain high expression of ABCG2. *Mol. Cancer Ther.* 2009; 8:2959-2968.
47. Cha ST, Chen PS, Johansson G, Chu CY, Wang MY, Jeng YM, Yu SL, Chen JS, Chang KJ, Jee SH, Tan CT, Lin MT, Kuo ML. MicroRNA-519c suppresses hypoxia-inducible factor-1alpha expression and tumor angiogenesis. *Cancer Res.* 2010; 70:2675-2685.

SUPPLEMENTAL TABLES

miRNA	FC	SD	miRNA	FC	SD	miRNA	FC	SD
miR-605	1.98	0.61	miR-504	1.25	0.35	miR-129-5p	0.98	0.06
miR-1231	1.97	0.10	miR-508-3p	1.24	0.40	miR-518c	0.95	0.40
miR-1256	1.95	0.55	miR-1262	1.24	0.30	miR-640	0.93	0.20
miR-128	1.79	0.26	miR-509-3-5p	1.24	0.32	miR-1243	0.91	0.31
miR-562	1.79	1.11	miR-509-5p	1.23	0.40	miR-653	0.88	0.43
miR-548l	1.79	0.76	miR-511	1.23	0.30	miR-1263	0.83	0.48
miR-520b	1.75	0.59	miR-512-5p	1.23	0.50	miR-618	0.82	0.60
miR-373	1.72	0.75	miR-515-3p	1.22	0.31	miR-106b	0.82	0.60
miR-483-5p	1.71	1.01	miR-513c	1.22	0.41	miR-135a	0.81	0.51
miR-550	1.71	1.07	miR-516b	1.22	0.22	miR-548d-3p	0.81	0.13
miR-25	1.70	0.75	miR-516a-3p	1.22	0.21	miR-520a-3p	0.80	0.44
miR-518e	1.68	0.53	miR-517b	1.22	0.51	miR-651	0.79	0.35
miR-1224-3p	1.66	0.52	miR-518a-3p	1.21	0.24	miR-133b	0.78	0.17
miR-876-5p	1.66	0.52	miR-518b	1.21	0.36	miR-1202	0.78	0.07
miR-507	1.64	1.04	miR-518d-3p	1.21	0.51	miR-1197	0.78	0.28
miR-1266	1.64	0.57	miR-520a-5p	1.17	0.40	miR-548i	0.77	0.15
miR-577	1.64	0.13	miR-494	1.16	0.20	miR-548c-3p	0.77	0.19
miR-1304	1.62	0.76	miR-520g	1.15	0.31	miR-520d-3p	0.75	0.14
miR-1253	1.59	0.41	miR-523	1.15	0.30	miR-34a	0.75	0.38
miR-760	1.58	1.14	miR-521	1.15	0.29	miR-664	0.74	0.17
miR-519e	1.58	1.02	miR-524-5p	1.15	0.50	miR-1208	0.73	0.13
miR-206	1.57	1.07	miR-525-5p	1.15	0.46	miR-9	0.73	0.13
miR-127-5p	1.54	0.75	miR-526b	1.15	0.44	miR-1268	0.72	0.15
miR-1227	1.50	0.11	miR-532-3p	1.15	0.43	miR-217	0.72	0.02
miR-370	1.50	0.45	miR-539	1.14	0.33	miR-1237	0.70	0.04
miR-371-5p	1.49	0.51	miR-542-3p	1.14	0.63	miR-758	0.70	0.15
miR-1207-3p	1.48	0.11	miR-548b-5p	1.12	0.23	miR-548h	0.70	0.15
miR-510	1.48	0.32	miR-633	1.11	0.40	miR-495	0.69	0.13
miR-888	1.47	1.10	miR-208a	1.10	0.07	miR-1229	0.67	0.10
miR-376a	1.46	0.71	miR-548g	1.10	0.20	miR-1236	0.66	0.13
miR-376c	1.46	0.52	miR-1276	1.10	0.14	miR-320c	0.65	0.10
miR-378	1.46	0.76	miR-1269	1.10	0.52	miR-1179	0.65	0.07
miR-1272	1.46	0.08	miR-508-5p	1.10	0.44	miR-24	0.64	0.21
miR-380	1.46	0.39	miR-549	1.10	0.32	miR-1281	0.63	0.13
miR-553	1.46	0.18	miR-1826	1.09	0.68	miR-548a-5p	0.62	0.13
miR-382	1.45	0.60	miR-1266	1.09	0.20	miR-212	0.61	0.13
miR-1226	1.43	0.10	miR-554	1.09	0.20	miR-450b-5p	0.60	0.17
miR-384	1.42	0.23	miR-552	1.09	0.19	miR-587	0.60	0.08
miR-409-5p	1.41	0.42	miR-622	1.09	0.18	miR-1255a	0.59	0.19
miR-421	1.41	0.51	miR-556-3p	1.09	0.25	miR-1249	0.59	0.15
miR-1278	1.40	0.45	miR-557	1.08	0.42	miR-375	0.59	0.14
miR-423-3p	1.40	0.41	miR-559	1.08	0.34	miR-335	0.59	0.15
miR-551a	1.40	0.52	miR-1230	1.07	0.51	miR-1275	0.58	0.12
miR-424	1.39	0.38	miR-544	1.06	0.20	miR-365	0.57	0.13
miR-429	1.38	0.40	miR-564	1.06	0.27	miR-626	0.57	0.04
miR-330-3p	1.38	0.37	miR-567	1.06	0.37	miR-1296	0.56	0.09
miR-296-5p	1.38	0.13	miR-571	1.06	0.33	miR-1207-5p	0.54	0.15
miR-1184	1.38	0.50	miR-573	1.06	0.43	miR-1264	0.54	0.15
miR-448	1.37	0.41	miR-574-5p	1.05	0.51	miR-132	0.54	0.15
miR-449b	1.37	0.39	miR-577	1.05	0.20	miR-423-5p	0.53	0.24
miR-548j	1.34	0.59	miR-579	1.05	0.24	miR-607	0.53	0.17
miR-450b-3p	1.34	0.50	miR-616	1.05	0.02	miR-320d	0.53	0.21
miR-613	1.33	0.57	miR-135b	1.04	0.48	miR-520c-3p	0.52	0.16
miR-105	1.32	0.60	miR-338-5p	1.04	0.22			
miR-1224-5p	1.32	0.30	miR-125a-3p	1.04	0.03			
miR-1261	1.32	0.45	miR-522	1.03	0.04			
miR-453	1.31	0.50	miR-584	1.03	0.23			
miR-371-3p	1.31	0.65	miR-582-5p	1.03	0.12			
miR-455-3p	1.30	0.51	miR-581	1.03	0.10			
miR-483-3p	1.30	0.31	miR-21	1.03	0.50			
miR-484	1.30	0.40	miR-586	1.03	0.21			
miR-1245	1.28	0.02	miR-1205	1.02	0.19			
miR-486-5p	1.28	0.37	miR-588	1.02	0.16			
miR-485-5p	1.28	0.31	miR-590-3p	1.02	0.21			
miR-487b	1.28	0.41	miR-591	1.01	0.42			
miR-489	1.28	0.51	miR-383	1.01	0.02			
miR-490-5p	1.28	0.29	miR-593	1.01	0.3			
miR-491-5p	1.27	0.41	miR-596	1.00	0.41			
miR-493	1.27	0.50	miR-604	1.00	0.30			
miR-497	1.27	0.27	miR-602	1.00	0.42			
miR-548a-3p	1.26	0.45	miR-598	1.00	0.28			
miR-500	1.26	0.50	miR-1286	0.98	0.32			
miR-501-5p	1.26	0.42	miR-1285	0.98	0.38			
miR-502-5p	1.25	0.39	miR-1267	0.98	0.15			

Supplemental Table S1. MicroRNAs showing less than twofold differences in abundance in senescent relative to early-passage cells. RNA extracted from Y (pdl 22-25) and S (pdl 50-55) WI-38 cells was used to measure the levels of the microRNAs listed, using RT-qPCR (Materials and Methods). microRNA abundance was normalized to U1 snRNA levels. Data are the means and S.D. from three independent experiments.

miR name	miRBase Acc No.	Primer Sequence	miR name	miRBase Acc No.	Primer Sequence
miR-96	MIMAT0000095	TTTGGCACTAGCACATTTTGTGCT	miR-206	MIMAT0000462	TGGAATGTAAGGAAGTGTGTGG
miR-944	MIMAT0004987	AAATTAATTGTACATCGGATGAG	miR-203	MIMAT0000264	GTGAAATGTTTAGGCCACTAG
miR-9	MIMAT0000441	UCUUUGUUUUCUAGCUGUAUGA	miR-194	MIMAT0000460	TGTAACAGCACTCCATGTGGA
miR-890	MIMAT0004912	TACTTGGAAGGCATCAGTTG	miR-18b	MIMAT0001412	TAAGGTGCATCTAGTGCAGTTAG
miR-888	MIMAT0004916	TACTCAAAAAGCTGTCAAGTCA	miR-18a	MIMAT0000072	TAAGGTGCATCTAGTGCAGATAG
miR-876-5p	MIMAT0004924	TGGAATTTCTTTGTGAATCACCA	miR-182b	MIMAT0006766	ATTGATCATCGACACTTCAACGCAAT
miR-874	MIMAT0004911	CTGGCCTGGCCCGAGGGACCGA	miR-15a	MIMAT0000068	TAGCAGCACAATAGTGTGTGTG
miR-760	MIMAT0004957	CGGCTCTGGGCTGTGGGGA	miR-155	MIMAT0000646	TTAATGCTAATCGTGATAGGGGT
miR-758	MIMAT0003879	UUUGUGACCUUGUCCACUAAACC	miR-1537	MIMAT0007399	AAAACCGCTAGTTACAGTTGT
miR-7	MIMAT0000252	TGGAAGACTAGTATTTTGTGTG	miR-146b-3p	MIMAT0004766	TGCCCTGTGGACTCAGTTCTGG
miR-664	MIMAT0005949	UAUUCAUUUUCUCCAGCCUACA	miR-141	MIMAT0000432	TAACACTGTCTGGTAAAGATGG
miR-663	MIMAT0003326	AGGGGGGCGCCGCGGACCGC	miR-140-5p	MIMAT0000431	CAGTGTGTTTACCCTATGGTAG
miR-658	MIMAT0003336	GGCGGAGGGAAGTGGTCCGTTGGT	miR-140-3p	MIMAT0004597	TACCACAGGGTGAACACCGG
miR-653	MIMAT0003328	GTGTTGAAACAATCTCTACTG	miR-135b	MIMAT0000758	TATGGCTTTTCATTCTATGTGA
miR-651	MIMAT0003321	TTTATGATAAGCTTGTCTTTTG	miR-135a	MIMAT0000428	TATGGCTTTTATTCCTATGTGA
miR-649	MIMAT0003319	AAACCTGTGTTGTTCAAGAGTC	miR-133b	MIMAT0000770	TTTGGTCCCTTCAACCAGCTA
miR-641	MIMAT0003311	AAAGACATAGGATAGATCACCTC	miR-1323	MIMAT0005795	TCAAACTGAGGGGCATTTTCT
miR-640	MIMAT0003310	ATGATCCAGGAACCTGCCTCT	miR-132	MIMAT0000426	TAACAGTCTACAGCATGGTCCG
miR-639	MIMAT0003303	CTAATAGTATCTACCAATAAA	miR-1305	MIMAT0005893	TTTTCAACTTAATGGGAGAGA
miR-628-5p	MIMAT0004809	ATGCTGACATATTACTAGAGG	miR-1304	MIMAT0005892	TTTGGGCTACAGTGAATGTG
miR-626	MIMAT0003295	AGCUGUCUGAAAUGUCUU	miR-1303	MIMAT0005891	TTTAGAGACGGGGTCTTGCTCT
miR-625	MIMAT0003294	AGGGGGAAAGTCTATAGTCC	miR-1296	MIMAT0005794	TTAGGGCCCTGGCTCCATCTCC
miR-622	MIMAT0003291	ACAGTCTGCTGAGGTTGGAGC	miR-129-5p	MIMAT0000242	CTTTTGGCGGCTGGGCTTGC
miR-618	MIMAT0003287	AAACTCTACTTGTCTCTGAGT	miR-129-3p	MIMAT0004605	AAGCCCTTACCCTAAAAGCAT
miR-616	MIMAT0004805	AGTCATTGAGGGTTGAGCAG	miR-1291	MIMAT0005881	TGCCCTGACTGAAGACCAGCAGT
miR-613	MIMAT0003281	AGGAATGTTCTCTTTTGGC	miR-1289	MIMAT0005879	TGGAGTCCAGGAATCTGCAATTTT
miR-607	MIMAT0003275	GTTCAAATCCAGATCTATAAC	miR-1288	MIMAT0005942	TGGACTGCCCTGATCTGGAGA
miR-605	MIMAT0003273	TAAATCCCATGGTCCCTCTCCT	miR-1287	MIMAT0005878	TGCTGGATCAGTGGTTCGAGTC
miR-600	MIMAT0003268	ACTTACAGACAAGGCCTTGCTC	miR-1286	MIMAT0005877	TGCAGGACCAAGTAGAGCCCT
miR-587	MIMAT0003253	UUUCCAUJAGGUGAUGAGUCAC	miR-1285	MIMAT0005876	TCTGGGCAACAAGTGAGACCT
miR-584	MIMAT0003249	TTATGTTTGGCCTGGGACTGAG	miR-1284	MIMAT0005941	TCTATACAGCCCTGGCTTTTC
miR-577	MIMAT0003242	TAGATAAAATATTGGTACCTG	miR-1283	MIMAT0005799	TCTACAAAGGAAGCGCTTTCT
miR-576-3p	MIMAT0004796	AAGATGTGAAAAATTTGAAATC	miR-1282	MIMAT0005940	TGCTTTGCTTTTCTGCTGT
miR-572	MIMAT0003237	GTCGCTCGGCGGTGGCCCA	miR-1281	MIMAT0005939	UCGCCUCCUCCUCCUCC
miR-569	MIMAT0003234	AGTTAATGAATCCCTGGAAAAT	miR-1280	MIMAT0005946	TCCCACCGCTGCCACCC
miR-562	MIMAT0003226	AAAGTAGCTGTACCAATTTGC	miR-128	MIMAT0000424	TCACAGTGAACCGGTCTCTTT
miR-561	MIMAT0003225	CAAAAGTTAAGATCCTTGAAT	miR-1279	MIMAT0005937	TCATATTGCTTCTTTCT
miR-553	MIMAT0003216	AAAACGGTGAGATTTTGT	miR-1278	MIMAT0005936	TAGTACTGTGATATCATCTAT
miR-551a	MIMAT0003214	GCGACCCACTCTTGGTTTCCA	miR-1276	MIMAT0005930	TAAAGAGCCCTGTGGAGACA
miR-550	MIMAT0004800	AGTGCCTGAGGAGTAAGAGCC	miR-127-5p	MIMAT0004604	CTGAAAGCTCAGAGGCTCTGAT
miR-548p	MIMAT0005934	TAGCAAAAAGTGCAGTACTTT	miR-1275	MIMAT0005929	GUGGGGAGAGGCUJUC
miR-548l	MIMAT0005935	AAAAGTAATTGCGGATTTTGCC	miR-1274a	MIMAT0005927	GTCCTGTTTCAGGGGCCA
miR-548k	MIMAT0005882	AAAAGTACTTGGCGATTTTGTG	miR-127-3p	MIMAT0000446	TGGATCCGCTGAGCTTGGCT
miR-548j	MIMAT0005875	AAAAGTAATTGCGGCTTTTGTG	miR-1273	MIMAT0005926	GGCGGACAAAGCAAGCTTTTCT
miR-548i	MIMAT0005935	AAAAGUAAUUGCGGAUUUUGCC	miR-1272	MIMAT0005925	GATGATGATGGCAGCAAATCTGAAA
miR-548h	MIMAT0005928	AAAAGUAAUUGCGGUUUUUGUC	miR-1271	MIMAT0005796	CTTGGCACCTAGCAAGCACTCA
miR-548e	MIMAT0005874	AAAAACTGAGACTACTTTGCA	miR-1270	MIMAT0005924	CTGGAGATAGGAGAGCTGTGT
miR-548d-5p	MIMAT0005799	TCTACAAGGAAAGCGCTTTGTC	miR-1269	MIMAT0005923	CTGGACTGAGCCGTGCTACTGG
miR-548d-3p	MIMAT0003323	CAAAACCCACAGTTTCTTTGTC	miR-1268	MIMAT0005922	CGGGCUGUGGUGGGGG
miR-548c-5p	MIMAT0004806	AAAAGTAATTGCGGTTTTTGGC	miR-1267	MIMAT0005921	CCTGTTGAAGTGAATCCCCA
miR-548c-3p	MIMAT0003285	CAAAAUCUCUAAUUCUUUUGC	miR-1266	MIMAT0005920	CCTCAGGGCTGTAGACAGGGCT
miR-548b-3p	MIMAT0003254	CAAGAACCTCAGTGTCTTTGT	miR-1265	MIMAT0005918	CAGGATGTGGTCAAGTGTGTT
miR-548a-5p	MIMAT0004803	AAAAGUAAUUGCGAGUUUJACC	miR-1264	MIMAT0005791	CAAGTCTTATTGAGCACCTGTT
miR-548a-3p	MIMAT0003251	CAAAACTGGCAATTTACTTTTGC	miR-1263	MIMAT0005915	ATGGTACCCTGGCATACTGAGT
miR-545	MIMAT0003165	TCAGCAAACATTTATTGTGTGC	miR-1262	MIMAT0005914	ATGGGTGAATTTGTAAGGAT
miR-544	MIMAT0003164	ATTCTGCATTTTAGCAAGTTC	miR-1261	MIMAT0005913	ATGGATAAGGCTTTGGCTT
miR-543	MIMAT0004954	AAACATTGCGGCTGCACTTCTT	miR-1260	MIMAT0005911	ATCCCACCTCTGCCACCA
miR-522	MIMAT0002868	AAAATGGTTCCCTTTAGAGTGT	miR-125b	MIMAT0000423	TCCTGAGACCCCTAACCTGTGA
miR-520e	MIMAT0002825	AAAGTCTTCTTTTGGAGG	miR-125a-5p	MIMAT0000443	TCCCTGAGACCCCTTAACTGTGA
miR-520d-3p	MIMAT0002856	AAAGUUCUUCUUCUUGGUGGU	miR-125a-3p	MIMAT0004602	ACAGGTGAGGTTCTTTGGAGCC
miR-520c-3p	MIMAT0002846	AAAGTCTTCCCTTTAGAGGGT	miR-1259	MIMAT0005910	ATATATGATGACTTAGCTTTT
miR-520b	MIMAT0002843	AAAGTCTTCCCTTTAGAGGG	miR-1258	MIMAT0005909	AGTTAGGATTAGGTCGTGGAA
miR-520a-3p	MIMAT0002834	AAAGTCTTCCCTTTGGACTGT	miR-1257	MIMAT0005908	AGTAAATGATGGTTCTGACC
miR-519e	MIMAT0002829	AAGTGCTTCCCTTTAGAGTGT	miR-1256	MIMAT0005907	AGGCATTGACTTCTCACTAGCT
miR-519c-3p	MIMAT0002832	AAAGTGCATCTTTTAGAGGAT	miR-1255b	MIMAT0005945	CGGATGAGCAAGAAAGTGGTT
miR-519b-3p	MIMAT0002837	AAAGTGCATCTTTTAGAGGTT	miR-1255a	MIMAT0005906	AGGAGGCAAGAAAGAAUUAU
miR-519a	MIMAT0002869	AAAGTGCATCTTTTAGAGTGT	miR-1254	MIMAT0005905	AGCCTGGAAGCTGGAGCCTGCA
miR-518e	MIMAT0002861	AAAGCGCTTCCCTTCAAGTGT	miR-1253	MIMAT0005904	AGAGAAGAAGTACGCCCTGCA
miR-518d-3p	MIMAT0002864	CAAAAGCGCTTCCCTTTGGAGC	miR-1250	MIMAT0005902	ACGGTCTGGATGTGGCCCTTT
miR-518c	MIMAT0002848	CAAAAGCGCTTCTTTAGAGTGT	miR-1249	MIMAT0005901	ACGCCUJCCUCCUUCUJUCA
miR-518b	MIMAT0002844	CAAAAGCGCTTCCCTTTAGAGTGT	miR-1248	MIMAT0005900	ACCTTCTGTATAAGCACTGTCTAAA
miR-518a-3p	MIMAT0002863	GAAAGCGCTTCCCTTTGCTGGA	miR-1247	MIMAT0005899	ACCCGTCCTGGTTCCGCCGA
miR-517b	MIMAT0002857	TCGTGCATCCCTTTAGAGTGT	miR-1246	MIMAT0005898	AATGGATTTTGGAGCAGG
miR-516b	MIMAT0002859	ATCTGAGGTAAGAAGCACTTT	miR-1245	MIMAT0005897	AAGTACTTAAAGGCCACTAT

Supplemental Table S2. Primers used for RT-qPCR detection of the microRNAs in Figures 2 and 3 and in supplemental Figure S1.