

Research Article

Mild Solutions for Fractional Differential Equations with Nonlocal Conditions

Fang Li

School of Mathematics, Yunnan Normal University, Kunming 650092, China

Correspondence should be addressed to Fang Li, fangli860@gmail.com

Received 8 January 2010; Accepted 21 January 2010

Academic Editor: Gaston Mandata N'Guerekata

Copyright © 2010 Fang Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the existence and uniqueness of mild solution of the fractional differential equations with nonlocal conditions $d^q x(t)/dt^q = -Ax(t) + f(t, x(t), Gx(t))$, $t \in [0, T]$, and $x(0) + g(x) = x_0$, in a Banach space X , where $0 < q < 1$. General existence and uniqueness theorem, which extends many previous results, are given.

1. Introduction

The fractional differential equations can be used to describe many phenomena arising in engineering, physics, economy, and science, so they have been studied extensively (see, e.g., [1–8] and references therein).

In this paper, we discuss the existence and uniqueness of mild solution for

$$\begin{aligned} \frac{d^q x(t)}{dt^q} &= -Ax(t) + f(t, x(t), Gx(t)), \quad t \in [0, T], \\ x(0) + g(x) &= x_0, \end{aligned} \tag{1.1}$$

where $0 < q < 1$, $T > 0$, and $-A$ generates an analytic compact semigroup $\{S(t)\}_{t \geq 0}$ of uniformly bounded linear operators on a Banach space X . The term $Gx(t)$ which may be interpreted as a control on the system is defined by

$$Gx(t) := \int_0^t K(t, s)x(s)ds, \tag{1.2}$$

where $K \in C(D, \mathbb{R}^+)$ (the set of all positive function continuous on $D := \{(t, s) \in \mathbb{R}^2 : 0 \leq s \leq t \leq T\}$) and

$$G^* = \sup_{t \in [0, T]} \int_0^t K(t, s) ds < \infty. \quad (1.3)$$

The functions f and g are continuous.

The nonlocal condition $x(0) + g(x) = x_0$ can be applied in physics with better effect than that of the classical initial condition $x(0) = x_0$. There have been many significant developments in the study of nonlocal Cauchy problems (see, e.g., [6, 7, 9–14] and references cited there).

In this paper, motivated by [1–7, 9–15] (especially the estimating approach given by Xiao and Liang [14]), we study the semilinear fractional differential equations with nonlocal condition (1.1) in a Banach space X , assuming that the nonlinear map f is defined on $[0, T] \times X_\alpha \times X_\alpha$ and g is defined on $C([0, T], X_\alpha)$ where $X_\alpha = D(A^\alpha)$, for $0 < \alpha < 1$, the domain of the fractional power of A . New and general existence and uniqueness theorem, which extends many previous results, are given.

2. Preliminaries

In this paper, we set $I = [0, T]$, a compact interval in \mathbb{R} . We denote by X a Banach space with norm $\|\cdot\|$. Let $-A : D(A) \rightarrow X$ be the infinitesimal generator of a compact analytic semigroup of uniformly bounded linear operators $\{S(t)\}_{t \geq 0}$, that is, there exists $M > 1$ such that $\|S(t)\| \leq M$; and without loss of generality, we assume that $0 \in \rho(A)$. So we can define the fractional power A^α for $0 < \alpha \leq 1$, as a closed linear operator on its domain $D(A^\alpha)$ with inverse $A^{-\alpha}$, and one has the following known result.

Lemma 2.1 (see [15]). (1) $X_\alpha = D(A^\alpha)$ is a Banach space with the norm $\|x\|_\alpha := \|A^\alpha x\|$ for $x \in D(A^\alpha)$.

(2) $S(t) : X \rightarrow X_\alpha$ for each $t > 0$ and $\alpha > 0$.

(3) For every $u \in D(A^\alpha)$ and $t \geq 0$, $S(t)A^\alpha u = A^\alpha S(t)u$.

(4) For every $t > 0$, $A^\alpha S(t)$ is bounded on X and there exists $M_\alpha > 0$ such that

$$\|A^\alpha S(t)\| \leq M_\alpha t^{-\alpha}. \quad (2.1)$$

Definition 2.2. A continuous function $x : I \rightarrow X$ satisfying the equation

$$x(t) = S(t)(x_0 - g(x)) + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} S(t-s) f(s, x(s), Gx(s)) ds \quad (2.2)$$

for $t \in I$ is called a mild solution of (1.1).

In this paper, we use $\|f\|_p$ to denote the L^p norm of f whenever $f \in L^p(0, T)$ for some p with $1 \leq p < \infty$. We denote by C_α the Banach space $C([0, T], X_\alpha)$ endowed with the sup norm given by

$$\|x\|_\infty := \sup_{t \in I} \|x\|_\alpha, \tag{2.3}$$

for $x \in C_\alpha$.

The following well-known theorem will be used later.

Theorem 2.3 (Krasnoselkii, see [16]). *Let Ω be a closed convex and nonempty subset of a Banach space X . Let A, B be two operators such that*

- (i) $Ax + By \in \Omega$ whenever $x, y \in \Omega$.
- (ii) A is compact and continuous,
- (iii) B is a contraction mapping.

Then there exists $z \in \Omega$ such that $z = Az + Bz$.

3. Main Results

We require the following assumptions.

- (H1) The function $f : [0, T] \times X_\alpha \times X_\alpha \rightarrow X$ is continuous, and there exists a positive function $\mu(\cdot) : [0, T] \rightarrow \mathbb{R}^+$ such that

$$\begin{aligned} \|f(t, x, y)\| &\leq \mu(t), \text{ the function } s \mapsto \frac{\mu(s)}{(t-s)^\alpha} \text{ belongs to } L^p([0, t], \mathbb{R}^+), \\ \gamma(t) &:= \left(\int_0^t \left(\frac{\mu(s)}{(t-s)^\alpha} \right)^p ds \right)^{1/p} \leq M_T < \infty, \text{ for } t \in [0, T], \end{aligned} \tag{3.1}$$

where $p > 1/q > 1$.

- (H2) The function $g : C_\alpha \rightarrow X_\alpha$ is continuous and there exists $b > 0$ such that

$$\|g(x) - g(y)\|_\alpha \leq b \|x - y\|_\infty, \tag{3.2}$$

for any $x, y \in C_\alpha$.

Theorem 3.1. *Let $-A$ be the infinitesimal generator of an analytic compact semigroup $\{S(t)\}_{t \geq 0}$ with $\|S(t)\| \leq M$, $t \geq 0$, and $0 \in \rho(A)$. If the maps f and g satisfy (H1), (H2), respectively, and $Mb < 1$, then (1.1) has a mild solution for every $x_0 \in X_\alpha$.*

Proof. Set $\lambda = \sup_{x \in C_\alpha} \|g(x)\|_\alpha$ and choose r such that

$$r \geq M(\|x_0\|_\alpha + \lambda) + \frac{M_\alpha M_T}{\Gamma(q)} M_{p,q} \cdot T^{(q-1)/p}, \tag{3.3}$$

where $M_{p,q} := ((p-1)/(pq-1))^{(p-1)/p}$.

Let $B_r = \{x \in C([0, T], X_\alpha) \mid \|x\|_\infty \leq r\}$.
Define

$$\begin{aligned}(Ax)(t) &:= \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} S(t-s) f(s, x(s), Gx(s)) ds, \\ (Bx)(t) &:= S(t)(x_0 - g(x)).\end{aligned}\tag{3.4}$$

Let $x, y \in B_r$, then for $t \in [0, T]$ we have the estimates

$$\begin{aligned}& \| (Ax)(t) + (By)(t) \|_\alpha \\ & \leq \|S(t)\| (\|x_0\|_\alpha + \lambda) + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} \|A^\alpha S(t-s) f(s, x(s), Gx(s))\| ds \\ & \leq M(\|x_0\|_\alpha + \lambda) + \frac{M_\alpha}{\Gamma(q)} \int_0^t (t-s)^{q-1} \frac{\mu(s)}{(t-s)^\alpha} ds \\ & \leq M(\|x_0\|_\alpha + \lambda) + \frac{M_\alpha}{\Gamma(q)} \left(\int_0^t (t-s)^{(q-1)p/(p-1)} ds \right)^{(p-1)/p} \cdot \left(\int_0^t \left(\frac{\mu(s)}{(t-s)^\alpha} \right)^p ds \right)^{1/p} \\ & \leq M(\|x_0\|_\alpha + \lambda) + \frac{M_\alpha M_T}{\Gamma(q)} M_{p,q} \cdot T^{q-1/p} \\ & \leq r.\end{aligned}\tag{3.5}$$

Hence we obtain $Ax + By \in B_r$.

Now we show that A is continuous. Let $\{x_n\}$ be a sequence of B_r such that $x_n \rightarrow x$ in B_r . Then

$$f(s, x_n(s), Gx_n(s)) \longrightarrow f(s, x(s), Gx(s)), \quad n \longrightarrow \infty,\tag{3.6}$$

since the function f is continuous on $I \times X_\alpha \times X_\alpha$. For $t \in [0, T]$, using (2.1), we have

$$\begin{aligned}& \| (Ax_n)(t) - (Ax)(t) \|_\alpha \\ & = \frac{1}{\Gamma(q)} \left\| \int_0^t (t-s)^{q-1} S(t-s) [f(s, x_n(s), Gx_n(s)) - f(s, x(s), Gx(s))] ds \right\|_\alpha \\ & \leq \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} \|A^\alpha S(t-s) [f(s, x_n(s), Gx_n(s)) - f(s, x(s), Gx(s))]\| ds \\ & \leq \frac{M_\alpha}{\Gamma(q)} \int_0^t (t-s)^{q-1} \|f(s, x_n(s), Gx_n(s)) - f(s, x(s), Gx(s))\| (t-s)^{-\alpha} ds.\end{aligned}\tag{3.7}$$

In view of the fact that

$$\|f(s, x_n(s), Gx_n(s)) - f(s, x(s), Gx(s))\| \leq 2\mu(s), \quad s \in [0, T], \quad (3.8)$$

and the function $s \rightarrow 2\mu(s)(t - s)^{-\alpha}$ is integrable on $[0, t]$, then the Lebesgue Dominated Convergence Theorem ensures that

$$\int_0^t (t - s)^{q-1} \|f(s, x_n(s), Gx_n(s)) - f(s, x(s), Gx(s))\| (t - s)^{-\alpha} ds \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.9)$$

Therefore, we can see that

$$\lim_{n \rightarrow \infty} \|(Ax_n)(t) - (Ax)(t)\|_\infty = 0, \quad (3.10)$$

which means that A is continuous.

Noting that

$$\begin{aligned} \|(Ax)(t)\|_\alpha &= \frac{1}{\Gamma(q)} \left\| \int_0^t (t - s)^{q-1} S(t - s) f(s, x(s), Gx(s)) ds \right\|_\alpha \\ &\leq \frac{1}{\Gamma(q)} \int_0^t (t - s)^{q-1} \|A^\alpha S(t - s) f(s, x(s), Gx(s))\| ds \\ &\leq \frac{M_\alpha}{\Gamma(q)} \int_0^t (t - s)^{q-1} \frac{\mu(s)}{(t - s)^\alpha} ds \\ &\leq \frac{M_\alpha M_\Gamma}{\Gamma(q)} M_{p,q} \cdot T^{q-1/p}, \end{aligned} \quad (3.11)$$

we can see that A is uniformly bounded on B_r .

Next, we prove that $(Ax)(t)$ is equicontinuous. Let $0 < t_2 < t_1 < T$, and let $\varepsilon > 0$ be small enough, then we have

$$\begin{aligned} \|(Ax)(t_1) - (Ax)(t_2)\|_\alpha &\leq \frac{1}{\Gamma(q)} \left\| \int_0^{t_2} [(t_1 - s)^{q-1} - (t_2 - s)^{q-1}] S(t_2 - s) f(s, x(s), Gx(s)) ds \right\|_\alpha \\ &\quad + \frac{1}{\Gamma(q)} \left\| \int_{t_2}^{t_1} (t_1 - s)^{q-1} S(t_1 - s) f(s, x(s), Gx(s)) ds \right\|_\alpha \\ &\quad + \frac{1}{\Gamma(q)} \left\| \int_0^{t_2} (t_1 - s)^{q-1} [S(t_1 - s) - S(t_2 - s)] f(s, x(s), Gx(s)) ds \right\|_\alpha \\ &= I_1 + I_2 + I_3. \end{aligned} \quad (3.12)$$

Using (2.1) and (H1), we have

$$\begin{aligned}
I_1 &= \frac{1}{\Gamma(q)} \left\| \int_0^{t_2} [(t_1 - s)^{q-1} - (t_2 - s)^{q-1}] S(t_2 - s) f(s, x(s), Gx(s)) ds \right\|_{\alpha} \\
&\leq \frac{1}{\Gamma(q)} \int_0^{t_2} |(t_1 - s)^{q-1} - (t_2 - s)^{q-1}| \|A^{\alpha} S(t_2 - s) f(s, x(s), Gx(s))\| ds \\
&\leq \frac{M_{\alpha}}{\Gamma(q)} \int_0^{t_2} |(t_1 - s)^{q-1} - (t_2 - s)^{q-1}| \frac{\mu(s)}{(t_2 - s)^{\alpha}} ds \\
&\leq \frac{M_{\alpha}}{\Gamma(q)} \int_0^{t_2 - \varepsilon} [(t_2 - s)^{q-1} - (t_1 - s)^{q-1}] \frac{\mu(s)}{(t_2 - s)^{\alpha}} ds \\
&\quad + \frac{M_{\alpha}}{\Gamma(q)} \int_{t_2 - \varepsilon}^{t_2} (t_2 - s)^{q-1} \frac{\mu(s)}{(t_2 - s)^{\alpha}} ds \\
&= I_1' + I_1''.
\end{aligned} \tag{3.13}$$

It follows from the assumption of $\mu(s)$ that I_1' tends to 0 as $t_2 \rightarrow t_1$. For I_1'' , using the Hölder inequality, we can see that I_1'' tends to 0 as $t_2 \rightarrow t_1$ and $\varepsilon \rightarrow 0$.

For I_2 , using (2.1), (H1), and the Hölder inequality, we have

$$\begin{aligned}
I_2 &= \frac{1}{\Gamma(q)} \left\| \int_{t_2}^{t_1} (t_1 - s)^{q-1} S(t_1 - s) f(s, x(s), Gx(s)) ds \right\|_{\alpha} \\
&\leq \frac{1}{\Gamma(q)} \int_{t_2}^{t_1} (t_1 - s)^{q-1} \|A^{\alpha} S(t_1 - s) f(s, x(s), Gx(s))\| ds \\
&\leq \frac{M_{\alpha}}{\Gamma(q)} \int_{t_2}^{t_1} (t_1 - s)^{q-1} \frac{\mu(s)}{(t_1 - s)^{\alpha}} ds \rightarrow 0 \quad \text{as } t_2 \rightarrow t_1.
\end{aligned} \tag{3.14}$$

Moreover,

$$\begin{aligned}
I_3 &\leq \frac{1}{\Gamma(q)} \left\| \int_0^{t_2 - \varepsilon} (t_1 - s)^{q-1} [S(t_1 - s) - S(t_2 - s)] f(s, x(s), Gx(s)) ds \right\|_{\alpha} \\
&\quad + \frac{1}{\Gamma(q)} \left\| \int_{t_2 - \varepsilon}^{t_2} (t_1 - s)^{q-1} [S(t_1 - s) - S(t_2 - s)] f(s, x(s), Gx(s)) ds \right\|_{\alpha}
\end{aligned}$$

$$\begin{aligned}
 &\leq \frac{1}{\Gamma(q)} \int_0^{t_2-\varepsilon} (t_1-s)^{q-1} \left\| S\left(\frac{t_1-t_2}{2} + \frac{t_1-s}{2}\right) - S\left(\frac{t_2-s}{2}\right) \right\| \\
 &\quad \cdot \left\| A^\alpha S\left(\frac{t_2-s}{2}\right) f(s, x(s), Gx(s)) \right\| ds \\
 &\quad + \frac{M_\alpha}{\Gamma(q)} \int_{t_2-\varepsilon}^{t_2} (t_1-s)^{q-1} \left[\frac{\mu(s)}{(t_1-s)^\alpha} + \frac{\mu(s)}{(t_2-s)^\alpha} \right] ds \\
 &\leq \frac{2^\alpha M_\alpha}{\Gamma(q)} \int_0^{t_2-\varepsilon} (t_1-s)^{q-1} \left\| S\left(\frac{t_1-t_2}{2} + \frac{t_1-s}{2}\right) - S\left(\frac{t_2-s}{2}\right) \right\| \cdot \frac{\mu(s)}{(t_2-s)^\alpha} ds \\
 &\quad + \frac{M_\alpha}{\Gamma(q)} \int_{t_2-\varepsilon}^{t_2} (t_1-s)^{q-1} \left[\frac{\mu(s)}{(t_1-s)^\alpha} + \frac{\mu(s)}{(t_2-s)^\alpha} \right] ds \\
 &= I'_3 + I''_3. \tag{3.15}
 \end{aligned}$$

Using the compactness of $S(t)$ in X implies the continuity of $t \mapsto \|S(t)\|$ for $t \in [0, T]$; integrating with $s \mapsto \mu(s)/(t_2-s)^\alpha \in L^1_{\text{loc}}([0, t_2], \mathbb{R}^+)$, we see that I'_3 tends to 0, as $t_2 \rightarrow t_1$. For I''_3 , from the assumption of $\mu(s)$ and the Hölder inequality, it is easy to see that I''_3 tends to 0 as $t_2 \rightarrow t_1$ and $\varepsilon \rightarrow 0$.

Thus, $\|(Ax)(t_1) - (Ax)(t_2)\|_\alpha \rightarrow 0$, as $t_2 \rightarrow t_1$, which does not depend on x .

So, $A(B_r)$ is relatively compact. By the Arzela-Ascoli Theorem, A is compact.

Now, let us prove that B is a contraction mapping. For $x, y \in C([0, T], X_\alpha)$ and $t \in [0, T]$, we have

$$\|(Bx)(t) - (By)(t)\|_\alpha \leq \|S(t)\| \|g(x) - g(y)\|_\alpha \leq Mb \|x - y\|_\infty < \|x - y\|_\infty. \tag{3.16}$$

So, we obtain

$$\|(Bx)(t) - (By)(t)\|_\infty < \|x - y\|_\infty. \tag{3.17}$$

We now conclude the result of the theorem by Krasnoselkii's theorem. □

Now we assume the following.

(H3) There exists a positive function $\mu_1(\cdot) : [0, T] \rightarrow \mathbb{R}^+$ such that

$$\|f(t, x(t), Gx(t)) - f(t, y(t), Gy(t))\| \leq \mu_1(t) (\|x - y\|_\alpha + \|Gx - Gy\|_\alpha), \tag{3.18}$$

the function $s \mapsto \mu_1(s)/(t-s)^\alpha$ belongs to $L^1([0, t], \mathbb{R}^+)$ and

$$\gamma'(t) := \left(\int_0^t \left(\frac{\mu_1(s)}{(t-s)^\alpha} \right)^p ds \right)^{1/p} \leq M'_T < \infty, \quad \text{for } t \in [0, T]. \tag{3.19}$$

(H4) The function $L_{\alpha,q} : I \rightarrow \mathbb{R}^+$, $0 < \alpha$, $q < 1$ satisfies

$$L_{\alpha,q}(t) = Mb + \frac{M_\alpha M'_T}{\Gamma(q)} M_{p,q} \cdot t^{q-1/p} (1 + G^*) \leq \omega < 1, \quad t \in [0, T]. \quad (3.20)$$

Theorem 3.2. Let $-A$ be the infinitesimal generator of an analytic semigroup $\{S(t)\}_{t \geq 0}$ with $\|S(t)\| \leq M$, $t \geq 0$ and $0 \in \rho(A)$. If $x_0 \in X_\alpha$ and (H2)–(H4) hold, then (1.1) has a unique mild solution $x \in C_\alpha$.

Proof. Define the mapping $\mathcal{F} : C([0, T], X_\alpha) \rightarrow C([0, T], X_\alpha)$ by

$$(\mathcal{F}x)(t) = S(t)(x_0 - g(x)) + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} S(t-s) f(s, x(s), Gx(s)) ds. \quad (3.21)$$

Obviously, \mathcal{F} is well defined on $C([0, T], X_\alpha)$.

Now take $x, y \in C([0, T], X_\alpha)$, then we have

$$\begin{aligned} & \|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)\|_\alpha \\ & \leq \|S(t)(g(x) - g(y))\|_\alpha \\ & \quad + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} \|S(t-s)[f(s, x(s), Gx(s)) - f(s, y(s), Gy(s))]\|_\alpha ds \\ & \leq M \|g(x) - g(y)\|_\alpha \\ & \quad + \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} \|A^\alpha S(t-s)[f(s, x(s), Gx(s)) - f(s, y(s), Gy(s))]\| ds \quad (3.22) \\ & \leq Mb \|x - y\|_\infty + \frac{M_\alpha}{\Gamma(q)} \int_0^t (t-s)^{q-1} \frac{\mu_1(s)}{(t-s)^\alpha} (\|x - y\|_\alpha + \|Gx - Gy\|_\alpha) ds \\ & \leq Mb \|x - y\|_\infty + \frac{M_\alpha M'_T}{\Gamma(q)} M_{p,q} \cdot t^{q-1/p} (1 + G^*) \|x - y\|_\alpha \\ & \leq L_{\alpha,q}(t) \|x - y\|_\infty. \end{aligned}$$

Therefore, we obtain

$$\|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)\|_\infty \leq \omega \|x - y\|_\infty < \|x - y\|_\infty, \quad (3.23)$$

and the result follows from the contraction mapping principle. \square

Acknowledgment

This work is supported by the NSF of Yunnan Province (2009ZC054M).

References

- [1] R. P. Agarwal, M. Belmekki, and M. Benchohra, "A survey on semilinear differential equations and inclusions involving Riemann-Liouville fractional derivative," *Advances in Difference Equations*, vol. 2009, Article ID 981728, 47 pages, 2009.
- [2] M. M. El-Borai and D. Amar, "On some fractional integro-differential equations with analytic semigroups," *International Journal of Contemporary Mathematical Sciences*, vol. 4, no. 25–28, pp. 1361–1371, 2009.
- [3] V. Lakshmikantham, "Theory of fractional functional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 10, pp. 3337–3343, 2008.
- [4] V. Lakshmikantham and A. S. Vatsala, "Basic theory of fractional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 8, pp. 2677–2682, 2008.
- [5] Z. W. Lv, J. Liang, and T. J. Xiao, "Solutions to fractional differential equations with nonlocal initial condition in Banach spaces," reprint, 2009.
- [6] H. Liu and J.-C. Chang, "Existence for a class of partial differential equations with nonlocal conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 9, pp. 3076–3083, 2009.
- [7] G. M. N'Guérékata, "A Cauchy problem for some fractional abstract differential equation with non local conditions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 5, pp. 1873–1876, 2009.
- [8] X.-X. Zhu, "A Cauchy problem for abstract fractional differential equations with infinite delay," *Communications in Mathematical Analysis*, vol. 6, no. 1, pp. 94–100, 2009.
- [9] J. Liang, J. van Casteren, and T.-J. Xiao, "Nonlocal Cauchy problems for semilinear evolution equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 50, no. 2, pp. 173–189, 2002.
- [10] J. Liang, J. Liu, and T.-J. Xiao, "Nonlocal Cauchy problems governed by compact operator families," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 57, no. 2, pp. 183–189, 2004.
- [11] J. Liang, J. H. Liu, and T.-J. Xiao, "Nonlocal Cauchy problems for nonautonomous evolution equations," *Communications on Pure and Applied Analysis*, vol. 5, no. 3, pp. 529–535, 2006.
- [12] J. Liang, J. H. Liu, and T.-J. Xiao, "Nonlocal impulsive problems for nonlinear differential equations in Banach spaces," *Mathematical and Computer Modelling*, vol. 49, no. 3–4, pp. 798–804, 2009.
- [13] J. Liang and T.-J. Xiao, "Semilinear integrodifferential equations with nonlocal initial conditions," *Computers & Mathematics with Applications*, vol. 47, no. 6–7, pp. 863–875, 2004.
- [14] T.-J. Xiao and J. Liang, "Existence of classical solutions to nonautonomous nonlocal parabolic problems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 63, no. 5–7, pp. e225–e232, 2005.
- [15] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, vol. 44 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1983.
- [16] D. R. Smart, *Fixed Point Theorems*, Cambridge University Press, 1980.