

# Mimivirus: leading the way in the discovery of giant viruses of amoebae

Philippe Colson<sup>1</sup>, Bernard La Scola<sup>1</sup>, Anthony Levasseur<sup>1</sup>, Gustavo Caetano-Anollés<sup>2</sup> and Didier Raoult<sup>1</sup>

**Abstract** | The accidental discovery of the giant virus of amoeba — *Acanthamoeba polyphaga* mimivirus (APMV; more commonly known as mimivirus) — in 2003 changed the field of virology. Viruses were previously defined by their submicroscopic size, which probably prevented the search for giant viruses, which are visible by light microscopy. Extended studies of giant viruses of amoebae revealed that they have genetic, proteomic and structural complexities that were not thought to exist among viruses and that are comparable to those of bacteria, archaea and small eukaryotes. The giant virus particles contain mRNA and more than 100 proteins, they have gene repertoires that are broader than those of other viruses and, notably, some encode translation components. The infection cycles of giant viruses of amoebae involve virus entry by amoebal phagocytosis and replication in viral factories. In addition, mimiviruses are infected by virophages, defend against them through the mimivirus virophage resistance element (MIMIVIRE) system and have a unique mobilome. Overall, giant viruses of amoebae, including mimiviruses, marseilleviruses, pandoraviruses, pithoviruses, faustoviruses and molliviruses, challenge the definition and classification of viruses, and have increasingly been detected in humans.

The discovery of giant viruses of amoebae changed the field of virology<sup>1,2</sup>. Since the nineteenth century, viruses have been defined by their submicroscopic size, and this was a dogmatic obstacle that probably prevented (for decades) researchers from searching for, and discovering, giant protozoan viruses, which can be visualized by light microscopy using the usual dyes<sup>3</sup> (BOX 1; FIG. 1). Indeed, for years the *Acanthamoeba polyphaga* mimivirus (APMV) was considered to be an intracellular bacterium of amoebae<sup>1</sup> (BOX 2). In a similar manner, pandoraviruses were thought to be atypical parasites of amoebae<sup>4</sup>. However, the absence of ribosomal DNA from an isolate that was presumed to be a bacterium eventually led to the discovery of APMV<sup>1,2</sup>, and, interestingly, the absence of a ribosome essentially separates giant viruses from the three defined domains of life. The genomic and proteomic analyses of giant viruses revealed that they are more complex than other viruses. Furthermore, giant viruses have several novel biological features, which include the fact that mimiviruses can be infected by parasitic viruses (termed virophages), they can contain mobile DNA elements, (termed transpovirons), and they have a defence mechanism against virophages termed the mimivirus virophage resistance element (MIMIVIRE)<sup>5–7</sup>. Thus, how should giant viruses, which do not have a ribosome but have

a level of complexity that is approaching that of numerous bacteria and even eukaryotes with an intracellular growth, be defined<sup>8</sup> (BOX 1; [Supplementary information S1](#) (table))? Furthermore, there is disagreement as to whether the large genomes of giant viruses are a result of smaller viruses acquiring genes or of a genome with cellular ancestry adapting to escape the cell nucleus<sup>2,9–13</sup>. In this Review, we reflect on 13 years of research into giant viruses and detail the advances that have been made in characterizing their genomes, structures and mechanisms of replication, as well as the virophages and mobile elements with which they are associated. We place these findings in the context of the ongoing debate on the evolutionary origin of giant viruses and on the extent of similarity between giant viruses and bacteria and eukaryotic cells.

## Structure and genomes of giant viruses

APMV and other giant viruses of amoebae have several structural and genomic features that had not been described in viruses before their discovery.

**Understanding APMV.** APMV has remarkable features compared with other viruses. The APMV capsid is ~500 nm in size and is covered by fibrils that are ~120–140 nm long and 1.4 nm thick<sup>1</sup>. These fibrils are

<sup>1</sup>Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes (URMITE), Aix-Marseille University, UM63, CNRS 7278, IRD 198, INSERM 1095, Institut Hospitalo-Universitaire (IHU) - Méditerranée Infection, AP-HM, 19–21 Boulevard Jean Moulin, 13385 Marseille, France.

<sup>2</sup>Evolutionary Bioinformatics Laboratory, Department of Crop Sciences, University of Illinois, 332 National Soybean Research Center, 1101 West Peabody Drive, Urbana, Illinois 61801, USA.

Correspondence to D.R. [didier.raoult@gmail.com](mailto:didier.raoult@gmail.com)

doi:10.1038/nrmicro.2016.197  
Published online 27 Feb 2017

Box 1 | Defining viruses and microorganisms and their origin and evolution

**The definition of viruses and giant viruses of amoebae**

During the 1950s, Lwoff introduced a definition of viruses that was based mainly on negative criteria<sup>112</sup> (Supplementary information S1 (table)). Viruses were considered as simple entities that relied on a cellular host to replicate, were devoid of machineries for energy production and translation, and consisted of a nucleic acid enclosed in a symmetrical protein shell known as the capsid<sup>112,113</sup>. In 2003, the discovery of *Acanthamoeba polyphaga* mimivirus (APMV) challenged this paradigm and caused debates in regard to the definition and classification of viruses<sup>2,8,55</sup>. Although APMV has many of the phenotypic and genotypic characteristics of viruses, it also shares many features with microorganisms<sup>55</sup>. For example, APMV particles are visible using a light microscope<sup>1</sup>, have larger genomes than small bacteria and contain more diverse genes than other viruses<sup>2</sup>. Moreover, APMV is more complex than other viruses, as its capsid contains two types of nucleic acid — a DNA genome and some mRNAs<sup>2</sup> — as well as 114 proteins<sup>19</sup>. In addition, APMV contains translation factors and aminoacyl-tRNA synthetases (which is unprecedented for a virus), and six tRNAs (tRNA is rarely detected in other double-stranded DNA (dsDNA) viruses)<sup>2</sup>. Furthermore, the mobilome of APMV is more diverse than that of other viruses and includes several self-excising introns and inteins, as well as pro-viropages and transpovirons<sup>5,6</sup>. Finally, unlike many viruses, APMV can enter host cells through phagocytosis that does not require specific viral–host interactions, owing to its large size<sup>76</sup>; this may mean that APMV has a broad host range.

**The origin of giant viruses of amoebae**

The presence of genes that are homologous to bacterial, archaeal and eukaryotic genes in APMV, marseillevirus and other nucleocytoplasmic large DNA viruses (NCLDVs)<sup>2</sup> has enabled phylogenetic reconstructions and comparative analyses of gene repertoires, the results of which have been interpreted as evidence for a ‘fourth domain’ in the tree of life<sup>2,50,54</sup> alongside Bacteria, Archaea and Eukarya, which are domains that were proposed by Woese in 1977. Such a phylogeny would suggest an archaic origin that is contemporary to that of proto-eukaryotes<sup>13,52</sup>. As these analyses did not involve ribosomal genes, it was proposed that APMV and the other members of the proposed order *Megavirales* comprise a fourth ‘TRUC’ (things resisting uncompleted classification) of microorganisms rather than a fourth domain<sup>3</sup>. Phylogenetic studies that were based on entire repertoires of the structural domains of proteins<sup>51</sup> also suggested that a fourth major branch of the tree of life may exist. However, these claims are controversial and have been contested by arguments that the phylogenies were biased by lateral gene transfer from cellular organisms (mostly eukaryotes) to giant viruses and inappropriate methodologies<sup>10,12,13,114,115</sup>. Therefore, the evolutionary origins of giant viruses remain under debate.

**Virophages**

Viruses that depend on the co-infection of their amoebal host by mimiviruses. Virophages replicate within mimivirus factories.

**Transpovirons**

Transposable elements that are ~7 kb long and contain long terminal inverted repeats.

**Mimivirus virophage resistance element**

(MIMIVIRE). A viral defence system that confers a nucleic-acid-based immunity against virophage infection.

**Fourth domain**

A suggested additional domain to the three domains of life (Bacteria, Archaea and Eukarya) that were proposed by Woese. The term was initially coined in 2010.

morphologically unique among viruses and, although their structures have not been fully elucidated<sup>1,14</sup> (Supplementary information S2 (figure)), they form a dense layer, are extensively glycosylated, and enable the attachment of APMV to amoebae, bacteria, arthropods and fungi through glycans<sup>15</sup>. The capsid comprises proteins that have a double jelly-roll fold and is icosahedral except at one vertex, which is covered by a unique five-branch starfish-shaped structure (termed ‘stargate’) that is devoid of fibres<sup>16</sup>. Beneath the capsid, and surrounded by an inner lipid membrane and fibres, is a spherical lipid bilayer compartment that is ~340 nm in size and contains the genome (with an estimated packing density of ~0.06 nm<sup>3</sup> per bp) and proteins<sup>17,18</sup>. This nucleocapsid has a large depression that faces the ‘stargate’, which creates a cavity and indicates that the nucleocapsid has a fixed position relative to the external capsid. APMV particles contain 114 proteins, which is only 12% of the number of predicted genes (see below), among which 12 are involved in transcription, 5 are involved in DNA topology and repair, 2 are involved in RNA modification, 5 are involved in particle structure and 7 are involved in protein or lipid modifications<sup>19</sup>.

APMV has a double-stranded DNA (dsDNA) genome that is 1.2 Mb in length and contains 979 genes that putatively encode proteins with a coding density of 89%<sup>2,20</sup> (FIG. 2; Supplementary information S3 (table)). The genome is also AT rich, comprising 72% AT nucleobases. Several APMV genes are not found in viruses other than giant viruses of amoebae, including those that encode translation factors and aminoacyl tRNA synthetases; some of the genes encoding aminoacyl tRNA synthetases are expressed<sup>21,22</sup> and encode functional proteins<sup>23</sup>. In addition, the APMV genome encodes four different tRNAs<sup>2</sup>. Other genes that are unique to APMV and other giant viruses of amoebae encode proteins that are involved in nucleotide synthesis, amino acid metabolism, protein modification, lipid or polysaccharide metabolism, DNA repair or protein folding<sup>2</sup>. In addition, the APMV genome contains four major groups of genes, including core genes that are shared with poxviruses, ascoviruses, iridoviruses, asfarviruses, phycodnaviruses and other giant viruses of amoebae, as well as large sets of horizontally transferred genes, paralogous genes (in addition to genes that were involved in large genome duplication events) and orphan genes (also known as ORFan genes)<sup>2,9,24–26</sup>. Orphan genes are estimated to represent ~48% of the predicted gene content<sup>2,27</sup>, and the proportion of orphan-encoded proteins is also very high (40%) in APMV particles<sup>19</sup>. The APMV genome also contains early and late gene promoters, and mRNAs are expressed as polyadenylated transcripts that most often end with short palindromic sequences that form hairpin-like structures<sup>2,22,28</sup>. Recoding events, including a frameshift and a readthrough, were described for a gene that encodes a translation termination factor in the APMV genome<sup>29</sup>. In addition, introns and inteins were detected in a few conserved genes, including those that encode the major capsid protein and the DNA polymerase<sup>30,31</sup>, and a ‘mobilome’ that is unique to APMV was identified<sup>6</sup> (see below). Finally, a few mRNAs were detected in mimivirus capsids<sup>2</sup> (FIG. 2).

**Other giant viruses of amoebae.** In 2005, APMV became the founding member of the family *Mimiviridae*<sup>32</sup>. Since then, about 100 new mimivirus strains have been isolated by culturing on amoebae from water, soil, insect and human samples that were collected worldwide (Supplementary information S3,S4 (table, figure)), most recently using high-throughput strategies<sup>33–35</sup> (BOX 3); the second mimivirus to be cultured was named mamavirus<sup>5</sup>. The sizes, morphologies and genomes of the other mimivirus isolates are similar to those of APMV<sup>30,33,36</sup>. They have capsids 370–600 nm in diameter and their genomes are 1.02–1.26 Mb in length, AT rich (72–75% AT nucleobases) and encode 930–1,120 putative proteins. Phylogenomics has enabled mimiviruses that infect amoebae to be divided into three lineages that were named lineage A (in which APMV is the prototype), lineage B (in which moumouvirus is the prototype)<sup>30</sup> and lineage C (in which *Megavirus chiliensis* is the prototype)<sup>36</sup>. In 2010, a distant mimivirus relative named *Cafeteria roenbergensis* virus, the capsid and genome of which are smaller than those of the mimiviruses of *Acanthamoeba*, was isolated from an abundant marine

**TRUC**

(Things resisting uncompleted classifications). A term that was coined in 2013 and proposes an alternative classification of microorganisms to the one that is based on ribosomal genes, which groups giant viruses in a fourth TRUC of microorganisms.

**Double jelly-roll fold**

A protein fold that is found exclusively in double-stranded DNA viruses and is comprised of two connected single jelly-roll folds, which are composed of eight  $\beta$ -strands arranged in two four-stranded sheets.

**Orphan genes**

Genes that lack a homologue in any sequence database.

**Mobilome**

This term represents the mobile genetic elements in the mimivirus genome and corresponds to proviropages and transpovirons, in addition to inteins and introns.

dinoflagellate<sup>37</sup>. Subsequently, a few other viruses that infect marine unicellular eukaryotes, including *Phaeocystis globosa* virus, were linked, albeit distantly, to mimiviruses<sup>38–40</sup>.

In addition to mimiviruses, other giant viruses of amoebae were discovered using amoebal co-culture methods. The first of these was marseillevirus, which was described in 2009 (REF. 41). Since 2013, the number and diversity of giant viruses of amoebae have expanded considerably, and, as of 2016, two virus families, the *Mimiviridae* and *Marseilleviridae*, have been described<sup>32,42</sup>. However, other giant viruses, including pandoraviruses, pithoviruses, faustoviruses and Mollivirus sibericum, represent putative new giant virus families<sup>5,43–46</sup> (Supplementary information S3,S5,S6 (table, table, figure)). Since the APMV genome was described, giant viruses of amoebae have been linked, through phylogenomic analyses, to nucleocytoplasmic large DNA viruses (NCLDV), which is a group of dsDNA viruses that was described in 2001 and comprises poxviruses, ascoviruses, iridoviruses, asfarviruses and phycodnaviruses; these viruses infect a wide range of eukaryotic cells, from algae to insects and mammals<sup>2,47,48</sup>. In 2009, it was noted that giant viruses of amoebae and NCLDVs share a small subset of nine core genes, five of which are found in all of their genomes (these encode a major capsid protein, a D5 helicase, a family B DNA polymerase, an A32-like

packaging ATPase and a very late transcription factor), and a larger subset of ~200 genes are shared by at least two NCLDV families<sup>48,49</sup>. Moreover, giant viruses of amoebae and NCLDVs were described to have a common ancestor, the genome of which is thought to contain ~50 conserved genes that are likely, based on phylogenomic analyses, to have an early origin that is possibly concomitant with eukaryogenesis<sup>48–52</sup>. In 2012, it was proposed that giant viruses of amoebae and NCLDV families should be classified into a new viral order, the *Megavirales*, on the basis of their common origin, which was suggested by the fact that they share a large set of ancestral genes that encode key viral functions, a common virus particle architecture and major biological features, including replication that occurs inside cytoplasmic factories<sup>53</sup>. Nevertheless, the architecture of pandoraviruses, Pithovirus sibericum and *M. sibericum* (see below) differs considerably from that of other giant viruses of amoebae and no capsid-resembling structure, or even gene for pandoraviruses, was detected in these viruses, which, together with poxviruses and ascoviruses that have brick-shaped virus particles and allantoid capsids, respectively, challenges the criteria that are used to classify viruses in the proposed order *Megavirales*<sup>43,44,46,53</sup>.

Members of the proposed order *Megavirales* that were isolated as a result of them being cultured on various amoebae and described during the past 13 years have a




Figure 1 | Particle and genome size of giant viruses of amoebae. Families (namely, the *Mimiviridae* and *Marseilleviridae*; represented by dark blue circles) or putative families of giant viruses of amoebae (namely, those that include pandoraviruses, pithoviruses, faustoviruses and mollivirus; dark blue circles) are shown, along with other families in the proposed order *Megavirales* (namely, *Poxviridae*, *Asfarviridae*, *Phycodnaviridae*, *Iridoviridae* and *Ascoviridae*; light blue circles). Some families or genera of smaller viruses (grey circles) are shown in the inset, which magnifies a section of the larger graph and shows viruses that have a genome size ≤ 400 kb and a particle size ≤ 400 nm. Circle sizes are proportional to virus particle sizes. For each family or genus, the size of the largest member is shown. Viruses are referred to by their family or genus name unless there is no family or genus name.

## Box 2 | The discovery of mimivirus

In the early 1980s, Rowbotham first described the interaction between *Legionella pneumophila* and amoebae in the genus *Acanthamoeba*<sup>119</sup>. Amoebae are host cells for *Legionella* spp., and, during the 1980s and 1990s, Rowbotham assembled a large collection of amoeba-associated bacteria from environmental water samples that were gathered as part of investigations into outbreaks of Legionnaires' disease. These bacteria were mostly strict intracellular *Legionella* spp.<sup>104,120</sup>, but two isolates were unidentified and, by their morphology, seemed to be small Gram-positive coccoid bacteria<sup>121</sup>. In 1995, Birtles brought this collection to the laboratory of D. Raoult with the objective of characterizing the two unidentified isolates by 16S ribosomal DNA (rDNA) amplification and sequencing. All attempts to amplify the isolate that became known as the 'Bradford coccus' were unsuccessful and, as an alternative strategy, the ultrastructure of this isolate was studied by electron microscopy<sup>1</sup>. Unexpectedly, the electron micrographs showed that the coccoid 'bacterium' in infected amoebae consisted of very regular icosahedral forms that resembled a giant iridovirus. Additional observations suggested that, despite its large diameter of ~600 nm, the Bradford coccus was in fact a virus, as the isolate exhibited eclipse-phase replication, which is typical of viruses, the production of virus particles into 'virus factories' and contained a large double-stranded DNA (dsDNA) chromosome that, according to preliminary genome analyses, belonged to the group of viruses that are known as nucleocytoplasmic large DNA viruses (NCLDVs)<sup>47</sup>. These studies revealed that the Bradford coccus was a virus, and it was subsequently renamed 'mimivirus', with an etymology that is based both on the concept of 'mimicking a microorganism' and on a childhood recollection of stories of 'Mimi the amoeba'.

wide range of sizes, shapes, structures, genome lengths, percentage of GC nucleobases, gene repertoires and replicative sites (Supplementary information S3,S5,S6,S7 (table, table, figure, figure)). Nonetheless, they still comprise a monophyletic clade that is based on a limited set of core genes and informational genes<sup>48,49,53–55</sup>. In addition, all of these giant viruses enter amoebae through phagocytosis, after which fusion occurs between the vacuole membrane and the internal membrane of the virus; this leads to the release of the genome into the cytoplasm of the amoeba<sup>56,57</sup>. Finally, the virus factory has a cytoplasmic location, with the exception of *M. sibericum*<sup>46</sup>.

Giant viruses of amoebae were isolated from various environmental samples, ecosystems and geographical locations, and from hosts, including amoebae, invertebrates and mammals<sup>33,57,58</sup>. They were also detected in several metagenomes that were generated from environmental, animal and human samples<sup>59–64</sup>, and in plant genomes<sup>65</sup>. Finally, sequences from new putative giant viruses were detected in marine environmental metagenomes by approaches that revealed that previously overlooked sequences are related to members of the *Megavirales*<sup>54,66,67</sup>. These facts suggest that giant viruses of amoebae are common in our biosphere.

Giant viruses of amoebae have various morphological features (Supplementary information S3,S5,S6 (table, table, figure)). Marseilleviruses and faustoviruses have icosahedral capsids that are 250 nm and 200 nm in size, respectively, and have no 'stargate' structure<sup>41,45</sup>. The surface of the marseillevirus prototype isolate is covered by fibres that are 12 nm long, whereas the faustovirus prototype isolate has no fibres. Contrary to all other DNA viruses, the faustovirus prototype isolate has a double protein shell to encapsidate, with an outer shell that is composed of a double jelly-roll protein and is similar

to that of many double-stranded DNA viruses (including giant viruses of amoebae), and an inner shell that is composed of a repeated hexameric unit, the structure of which differs from that of known capsid proteins. In addition, the major capsid protein of the outer shell is encoded by a 17,000 bp genomic fragment that contains several introns and exons<sup>68</sup>. Pandoraviruses, pithoviruses and *M. sibericum* exhibit different morphologies to other giant viruses of amoebae; virus particles have ovoid or spherical (in the case of *M. sibericum*) shapes<sup>43,44,46</sup>. In addition, pandoraviruses and pithoviruses are larger than mimiviruses and all other giant viruses of amoebae. Pandoravirus and pithovirus particles have a wall that is 60–70 nm thick and an apical pore at one of their extremities, which enables the internal virus components to be delivered into the amoebal cytoplasm and has the appearance of a honeycomb grid in pithoviruses<sup>43,44</sup>. The *M. sibericum* tegument is covered by two layers of fibres that are ~10–14 nm long and has a funnel shaped aperture at its apex<sup>46</sup>.

The genomes of giant viruses of amoebae other than mimiviruses have different lengths, GC content, gene numbers, functions and origins<sup>2,40,42–45</sup> (Supplementary information S6,S7 (figures)). They all include numerous orphan genes, similarly to APMV, which indicates that many of their structural and functional characteristics are novel and remain to be deciphered<sup>2,41,43–46</sup>. In addition, these genomes all have a substantial level of mosaicism; this is well exemplified by marseillevirus<sup>41</sup>. This suggests that many genes were transferred laterally between themselves and other viruses and cellular organisms. Among examples of particular features in the genomes of giant viruses of amoebae, it should be mentioned that the genomes of marseilleviruses encode histone-like proteins<sup>41,69</sup>; the genes in pandoraviruses frequently include spliceosomal introns<sup>43</sup> and the genome of Pandoravirus salinus contains particular transposons that were termed miniature inverted-repeat transposable elements<sup>70</sup>; and the genomes of faustoviruses encode membrane occupation and recognition nexus (MORN)-repeat-containing proteins, similar to marseilleviruses and pandoraviruses<sup>44</sup>. In the faustovirus prototype isolate it was found that the capsid protein is encoded by a ~17,000 bp large genomic region with several exons and introns, which was most unexpected<sup>45,68</sup>. Of note, it has been hypothesized that giant virus genomes may evolve through a complex accordion-like process, with successive steps of genome expansions through duplications and gene transfers followed by genome reduction<sup>71</sup>. Hence, the genomes of giant viruses of amoebae both exhibit features that are shared between some or all giant viruses of amoebae and others that are particular to each family or putative family.

**Giant viruses of amoebae that are living fossils.** *P. sibericum* and *M. sibericum* were isolated from a Siberian permafrost sample that was more than 30,000 years old, which dates back to the Pleistocene epoch, and were therefore considered to be 'living fossils' (REF. 44). *P. sibericum* has an AT-rich (64%) dsDNA genome that is 610,033 bp in length, which is unexpectedly small

### Monophyletic clade

A group of organisms, or taxon, that consists of all of the descendants of an ancestral species.

### Phagocytosis

The engulfment of a solid particle by a cell, which forms an internal compartment named a phagosome.

### Mosaicism

Applied to microbial genomes, this term is used to describe the coexistence in a genome of genes with different origins (viral, bacteria, archaeal or eukaryotic) as the result of lateral gene transfers.

### Permafrost

Rock or soil at a temperature equal to, or below, 0 °C for two or more years. Permafrost is mainly located in and around the Arctic and Antarctic regions.




Figure 2 | **Major genomic and structural features of APMV.** Major structural features of the *Acanthamoeba polyphaga* mimivirus (APMV) particle are shown in the middle of the figure. Black arrows link APMV to the major features of the gene repertoire and nucleic acid content of APMV, including its mobilome and mRNAs, which are shown in grey boxes. The red arrow indicates that virophages can infect mimivirus factories. The minus sign indicates that mimivirus virophage resistance element (MIMIVIRE) can protect against virophages. \*Gene definitions and putative functions are given, as well as their relationship to other viral genes. †Corresponds to mimivirus genes, the closest match to which in the NCBI GenBank protein sequence database is from a virus that does not belong in the family *Mimiviridae* (except distant mimiviruses). <sup>ii</sup>Maximum and minimum proportions of genes that are inferred to be involved in lateral transfer are given, as assessed in three different studies<sup>9,25,26</sup>. NCLDV, nucleocytoplasmic large DNA viruses; VLTF2, viral late transcription factor 2.

considering that it has the largest particle size (~1.5 µm in length and 0.5 µm in diameter) among giant viruses of amoebae<sup>44</sup>. Moreover, its coding density is only 68%, owing to a considerable number of ~150-nm-long regularly interspersed palindromic sequences that are distributed in tandem repeats within arrays that are ~2,000 bp long. Although *P. sibericum* morphologically resembles pandoraviruses, it is most closely related to marseilleviruses and iridoviruses based on phylogenetic analyses. A total of 159 proteins were identified in

purified virus particles, of which two-thirds and the four most abundant have unknown functions<sup>44</sup>. The genome of *M. sibericum* is 651,523 bp in size, with inverted repeats approximately 10,000 bp long at its extremities, and, similar to pandoraviruses, is GC rich (60%)<sup>46</sup>. The majority of the sequences that are most similar to genes in *M. sibericum* correspond to pandoravirus genes, albeit with a low level of homology (about 40% on average), and this result is congruent with phylogenetic analyses. Purified virus particles contain 136 proteins, of

## dN/dS ratio

A ratio of the number, over a given period of time, of non-synonymous substitutions per non-synonymous site (dN) to the number of synonymous substitutions per synonymous site (dS), which quantifies selection pressures.

## Viral factories

Structural and functional elements that are associated with the replication of viral nucleic acids and the substantial production of virus particles.

## Box 3 | Strategies for the discovery of giant viruses

The use of amoebae as a support to isolate intra-amoebal microorganisms (known as the ‘amoebal enrichment method’ (REF. 11 G)) led to the isolation of the bacterium *Legionella pneumophila*. However, as samples that were grown using this technique were often contaminated by extracellular and intracellular bacteria, antibiotics and antifungal compounds were subsequently added to the media<sup>33,43,45,94,109</sup>. The antifungal amphotericin B limited the viability of amoebae, but this drawback was overcome by the adaptation of *Acanthamoeba castellanii* to higher concentrations of amphotericin B<sup>43</sup>, the use of thiabendazole as an antifungal, which is less toxic to amoebae<sup>45</sup>, and then the use of voriconazole, which is not toxic to amoebae, even at high doses. In addition, procedures to pre-enrich samples for amoebae, including the addition of nutrients to water samples followed by their incubation in the dark<sup>36,94</sup>, or the use of Prescott and James medium to resuspend the sample prior to culture, were also established<sup>44</sup>.

More recently, an automated system for the high-throughput isolation of giant viruses, based on the detection of lysed protozoa in liquid media by flow cytometry (virus particles stained by SYBR Green are sorted based on their size and DNA content)<sup>35</sup>, was implemented. Furthermore, whereas the use of *Acanthamoeba polyphaga* as an amoebal host for culture only enabled the isolation of mimiviruses and marseilleviruses, the culture of *A. castellanii* led to the isolation of Pandoravirus spp.<sup>43</sup>, Pithovirus sibericum<sup>44</sup> and Mollivirus sibericum<sup>46</sup>. This shows that *A. polyphaga* is less permissive to infection by several giant viruses of amoebae as a support for culture than *A. castellanii*. Changing the amoebal host was especially fruitful in the isolation of several faustoviruses<sup>45</sup>, which form a new putative family of giant viruses that are specific to the host *Vermamoeba vermiformis*<sup>117</sup>. However, each host must be adapted to a liquid medium that enables minimal survival before the multiplication of viruses. For example, the quick encystment of *V. vermiformis* and its inability to survive in Page’s amoeba saline (PAS) prevented its use as a host for the isolation of giant viruses, although this limitation could be circumvented by using a starvation medium in which this host could survive for culture<sup>45</sup>. Varying the host species has also provided insight into host specificity; as assessed with a large set of giant viruses of amoebae, a sample could be positive for viruses when inoculated on a given *Acanthamoeba* species and negative when inoculated on others<sup>118</sup>.

Finally, a major challenge is to separate viruses when virus mixtures are suspected by current flow cytometric gating strategies, owing to the fact that the end-point dilution method is time consuming and favours the isolation of fast-growing viruses. Fluorescence-activated cell sorting (FACS) for the isolation of giant viruses is under development in our laboratory with encouraging preliminary results.

which more than half and the three most abundant are orphans. In addition, a homologue of the major capsid protein in members of the proposed order *Megavirales* is translated, although it is apparently not involved in the structure of the virus particle<sup>46</sup>.


Another pithovirus, named Pithovirus massiliensis, was recently isolated from sewage in southern France and was compared with its fossil counterpart<sup>72</sup>. This analysis indicated that the genomic content of pithoviruses evolves slowly, as it was conserved after thousands of years, with selective pressure on the conserved genes. This suggests that the mechanisms of evolution are comparable in giant viruses and bacteria, and include selection, gene fixation and then selective constraints. This enabled the first estimation of a molecular clock in giant viruses, and the mutation rate that was estimated based on the dN/dS ratio was  $2 \times 10^{-6}$  substitutions per site per year, which is lower than that of RNA viruses and is very similar to that estimated for poxviruses ( $0.5\text{--}7 \times 10^{-6}$  substitutions per site per year)<sup>73</sup> and in the same order of magnitude as those found in bacteria and archaea<sup>72</sup>. Second, this showed that genes that were thought to be horizontally acquired were selected and highly conserved, which indicates that they are essential genes. The GC content and codon usage of these genes tend to adapt progressively to that of the recipient genome<sup>72</sup>. In light of these results, pithoviruses can select for genes that are acquired by horizontal gene transfer, followed by their long-term fixation and adaptation to viral codon usage. Third, orphans that are highly abundant in pithoviruses<sup>44,72</sup> were also constrained by strong selective pressure, which indicates that their accumulation is not random and is biologically

relevant. These findings do not support the hypothesis that giant viruses are ‘bags’ of genes and pseudogenes that have been randomly taken from cellular organisms and not used, but rather indicate that horizontally acquired genes and orphan genes are functional and biologically active. Similar findings were reported previously for two marseilleviruses that were isolated in south-eastern France and Australia<sup>74</sup>.

## Infection cycle of giant viruses

The replicative cycles of giant viruses in amoebal cells show several similarities, including phagocytic entry, DNA release and replication in ‘viral factories’. However, differences exist between giant viruses in the duration of their replication cycles, the involvement of the amoebal nucleus in virus replication, and the assembly and release of virus particles.

The replicative cycle of APMV, which lasts about 12 h and occurs in the amoebal cytoplasm<sup>1,56,75</sup> (FIG. 3), has several features that have not been observed before in viruses. APMV is internalized through phagocytosis<sup>76</sup> before its genome-containing internal core is released through the ‘stargate’ into the cytoplasm, through the fusion of viral and phagosome membranes; transcription may be initiated in these cores<sup>77</sup>. Subsequently, the genome is released from the core and is replicated at high levels, which generates an early cytoplasmic replicative centre that is thought to be engulfed by a membrane layer of the endoplasmic reticulum. Replicative centres then merge into a single virus factory that contains, from the inside to the outside, zones that are involved in replication, membrane



f	Marseilleviruses	Faustoviruses	Pandoraviruses	Pithovirus	Mollivirus
Amoebal host replication	<i>A. polyphaga</i> , <i>A. castellanii</i>	<i>V. vermiformis</i>	<i>A. castellanii</i>	<i>A. castellanii</i>	<i>A. castellanii</i>
Approximate duration	6–16h	18–20h	10–18h	10–20h	6h
Entry process	Possible entry through giant vesicles and endosome-stimulated pathways	–	–	–	–
Amoebal nucleus	Strong transient modification of its morphology during early replication steps	Some reorganization, including the loss of a spherical appearance and a reduction of size	Disorganized; exhibits numerous membrane invaginations; seems to be replaced by the virus factory	Unaffected	Disorganized; exhibits numerous membrane invaginations
Virion assembly	–	–	The wall and the interior of the viral particles seem to assemble simultaneously		
Virion egress	Through amoebal lysis	Through amoebal lysis	Through amoebal lysis	Through amoebal lysis	No amoebal lysis; new virions emerge at the amoebal nucleus periphery

Figure 3 | **The mimivirus replication cycle and key replicative features of other giant viruses of amoebae.**

**a** | Schematic of the replication cycle of *Acanthamoeba polyphaga* mimivirus (APMV). Virus particles can be seen at the surface of an *Acanthamoeba* sp. amoeba at the first stage of the cycle. Then, after virus entry through phagocytosis, the eclipse phase begins and the release of the APMV genome into the amoebal cytoplasm seeds the virus factory, which appears at a different location to the cell nucleus. Starting from ~8 h post-infection, the assembly of virus particles can be seen at the periphery of the virus factory. Amoebal lysis occurs ~12 h post-infection. **b–e** | Electron microscopy images of APMV particles and *Acanthamoeba* sp. cells that are infected with APMV. **b** | A mimivirus particle is shown. **c** | A mimivirus factory in the cytoplasm of an *Acanthamoeba* sp. amoeba 8 h post-infection. **d** | This image shows the edge and periphery of a virus factory, in which internal membrane biogenesis and assembly, capsid assembly and DNA packaging, and fibre acquisition of the APMV particles occur 8 h post-infection. **e** | This image shows an *Acanthamoeba* sp. amoeba ~12 h post-infection with mimivirus, with the amoebal cytoplasm filled with mimivirus particles. **f** | The key replicative features of giant viruses of amoebae are shown in this table. Dashes indicate where there is no noticeable feature for a virus. *A. castellanii*, *Acanthamoeba castellanii*; *V. vermiformis*, *Vermamoeba vermiformis*. Microscopy images in parts **b–e** courtesy of I. Pagnier, Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, Aix Marseille University, France.

**Exocytosis**

Active transport of molecules, such as proteins, out of a cell through a process that uses energy.

biogenesis, capsid assembly and DNA packaging, and fibre acquisition<sup>56,77</sup>. The virus internal layer seems to originate from multivesicular membrane structures that bud from the endoplasmic reticulum and become open membrane sheets; the major capsid protein is assembled around these sheets while acting as a scaffold<sup>78</sup>. Genome packaging occurs through a transient aperture that is distinct from the ‘stargate’ (REF. 78). Finally, the layer of fibres is assembled. The involvement of the amoebal nucleus in the life cycle of APMV has been debated<sup>56,57,75</sup>. Indeed, the size of the amoebal nucleus decreases approximately twofold during APMV replication, even though this process occurs in a viral factory in the cytoplasm; amoebal nuclear factors may be involved.

In marseilleviruses, the replication cycle lasts about 6–16 h (REFS 41, 69) (FIG. 3). The cell nucleus undergoes transient morphological changes during early stages of the replication cycle, and marseillevirus viral factories tend to have a more extensive distribution in the cytoplasm than APMV viral factories. Marseilleviruses can form giant vesicles, comprised of dozens of virus particles, that are wrapped by membranes derived from the amoebal endoplasmic reticulum<sup>79</sup>. The viruses can be released outside of the amoeba within these giant vesicles; this may cause the amoeba to undergo phagocytosis through an acidification-independent process<sup>79</sup>. In addition, single marseillevirus particles may enter *Acanthamoeba* spp. through an endosome-stimulated pathway or may group together to trigger their own phagocytic uptake<sup>79</sup>.

The replication cycle of faustoviruses occurs in *Vermamoeba vermiformis* (not in *Acanthamoeba* spp.) and lasts 18–20 h (REF. 45). It occurs within the amoebal cytoplasm independently of the cell nucleus, and the viral factory occupies almost the entire amoebal cell, although it should be noted that some virus particles that are produced by the factory may only correspond to empty capsids.

The life cycles of pandoraviruses, *P. sibericum* and *M. sibericum* last ~10–18 h, 10–20 h and 6 h, respectively<sup>43,44,46</sup>. The wall and interior of these virus particles seem to assemble simultaneously. The replication of *P. sibericum* does not seem to affect the amoebal nucleus<sup>44</sup>. By contrast, infection by pandoraviruses and *M. sibericum* results in a disorganized amoebal nucleus that exhibits numerous membrane invaginations<sup>43,46</sup>. The pandoravirus factory seems to replace the cell nucleus, whereas the nucleus persists during infection with *M. sibericum*, with new virus particles emerging at its periphery. These observations are consistent with the fact that transcription-associated proteins are absent from pandoravirus and mollivirus particles, which suggests that they require the amoebal nucleus, but are present in pithovirus particles, which suggests that they can initiate transcription independently from the nucleus<sup>57</sup>.

Overall, giant viruses of amoebae enter cells through a phagocytic-like process and release hundreds of new virus particles through amoebal lysis; the exception is *M. sibericum*, for which the release of virus particles

seems to occur through exocytosis without lysis<sup>46</sup>. *Acanthamoeba* spp. or *V. vermiformis* can protect giant viruses against physical and chemical threats, as they are highly stable organisms that can become encysted<sup>80</sup>. In addition, APMV particles were stable for 9 months in environmental freshwater, saline water and hospital ventilator devices<sup>81</sup>, and resisted antibiotics, 48 h of desiccation, and several chemical biocides, including alcohols<sup>82</sup>. Finally, the phagocytosis-like entry process of giant viruses contrasts with the need of specific interactions for other viruses to enter their host and could enable giant viruses to infect a broader range of hosts. Although amoebae are the primary hosts of giant viruses, mimiviruses and marseilleviruses have been detected in oysters, insects, monkeys, cattle and humans<sup>83–85</sup>. Moreover, APMV can enter various human myeloid cells and replicate in peripheral blood mononuclear cells<sup>76,86</sup>, and marseillevirus can cause a productive infection in human T lymphocyte cells<sup>87</sup>. In addition, the virus factories in which these giant viruses replicate were described for other viruses, including members of the proposed order *Megavirales* that are not discussed here, other DNA viruses (including herpesviruses) and some RNA viruses (such as flaviviruses or coronaviruses)<sup>88</sup>. These factories are replication organelles that recruit viral and cellular components for virus assembly and maturation. Viral factories were considered as the nuclei of ‘virocells’ (that is, cells infected by a virus, the aim of which is to produce virions), which would themselves correspond to the cellular forms of viruses when not strictly assimilated into a virus particle (or virion)<sup>26</sup>. In the case of APMV and other giant viruses of amoebae, if the virocell is the infected amoeba, the giant viral factory can be considered as its nucleus. This view is connected to two evolutionary scenarios: viral eukaryogenesis<sup>13</sup>, in which the eukaryotic nucleus originated from the viral factory of an ancient giant virus<sup>26</sup>, and nuclear virogenesis, in which some giant viruses originated from the nucleus of ancient proto-eukaryotic cells<sup>89</sup>.

**Virophages and other mobile elements**

The discovery and study of mimiviruses led to the identification of a ‘mobilome’ that is specific to these giant viruses and comprises viruses that can infect their viral factories and integrate into their genomes.

**Virophage discovery and diversity.** Virophages, which were discovered in 2008 (REF. 5), are viruses that infect mimivirus factories. They have small virus particles that are 35–74 nm in diameter and have an icosahedral capsid and a dsDNA genome of 17–29 kb. Their genomes are predicted to contain 16–34 genes, most of which are orphans or genes of unknown function, and six core genes<sup>90,91</sup>. Virophages cannot replicate alone in their amoebal host but, instead, replicate only in the presence of mimiviruses<sup>5</sup>. A co-culture procedure that uses a culturable helper virus was established to explore virophage diversity, and it enabled the isolation of new virophages and the analysis of their ability to infect mimivirus lineages<sup>92</sup>. The first virophage, named



Sputnik, was isolated with mamavirus and was shown to impair its replication cycle and morphogenesis<sup>5</sup>. Specifically, the ability of mamavirus to lyse amoebae was decreased and mamavirus particles with abnormal morphologies were observed. So far, three Sputnik-like virophages, namely Sputnik2, Sputnik 3 and Rio Negro virophage, have been isolated, and they all infect lineage A, lineage B and lineage C of the *Mimiviridae*<sup>92–94</sup>. A divergent virophage that is associated with mimiviruses of *Acanthamoeba* spp., named Zamilon, was also isolated<sup>95</sup>. Another virophage, named Mavirus, depends on *Cafeteria roenbergensis* virus for its replication and was the second virophage to be discovered after Sputnik<sup>96</sup>. The study of virophage diversity is still in its infancy, and several studies suggest that virophages are common and especially abundant in the oceans. For example, virophages were identified in the genome of the marine alga *Bigeloviella natans*<sup>97</sup>, and virophage genomes were assembled from metagenomic datasets<sup>98</sup>. Virophages that infect mimiviruses have been recently classified in the family *Lavidaviridae*<sup>91</sup>. This classification highlights that, as suggested by their particle and genome size, virophages are similar in complexity to genuine dsDNA viruses, and they share a set of six core genes that strongly suggests their monophyletic origin. In addition, the promoter and transcription termination signals of mimivirus virophages are similar to those of mimiviruses, which suggests that these virophages rely on their associated giant virus for mRNA synthesis<sup>99</sup>.

Although practical for isolating new virophages, the use of a culture procedure that is based on *A. polyphaga* and mimiviruses as a reporter system<sup>92</sup> was hindered by the unexpected discovery that mimiviruses of lineage A, but not of lineage B and lineage C, were resistant to the Zamilon virophage<sup>95</sup>. The search for a system that would explain this resistance was inspired by an analogy to the CRISPR–Cas system, which is widely present in bacteria and archaea and relies on the integration of short DNA sequences from invaders<sup>7,100</sup>. The resistant lineage A isolates indeed have an insertion of four 15-nucleotide-long repeated Zamilon virophage sequences within an operon, named the MIMIVIRE, the resistance mechanism of which was suggested to rely on the sequence-specific recognition of a nucleic acid sequence<sup>7</sup>. The probability of finding such a sequence in this single mimivirus lineage by chance was very low ( $<1 \times 10^{-9}$ ). MIMIVIRE-associated genes encode a helicase and nuclease that are involved in the degradation of foreign nucleic acids<sup>101</sup> and the functions of which were experimentally validated<sup>7</sup>; MIMIVIRE also includes a gene that contains the repeated insert. Silencing each of these three genes by RNA interference restored the susceptibility of mimivirus to the Zamilon virophage. However, the exact molecular mechanism of MIMIVIRE is unclear. The MIMIVIRE system is thought to differ from CRISPR–Cas systems owing to the absence of a Cas1 homologue, of protospacer-adjacent motifs and of a well-conserved organization that includes bona fide CRISPR-like repeats<sup>100,102</sup>. Furthermore, the helicase and nuclease that are described in this MIMIVIRE system are only

distantly related to proteins that are classified in the same superfamilies as Cas3 and Cas4 proteins, and they do not seem to be related to them<sup>100</sup>. An alternative mechanism to explain the resistance of mimivirus to virophages is one that would rely on interactions between proteins, rather than on nucleic acid-based recognition, and would involve a restriction factor protein of the virophage replication machinery<sup>102</sup>.

**Provirophages, transpovirons and mobilome.** Mobile genetic elements are common features in microorganisms, and mimiviruses have a complex mobilome (FIG. 3). Indeed, the DNA of virophages can integrate into the mimiviral host genome as ‘pro-virophages’ — that is, proviruses of virophages<sup>6</sup>. In addition, a novel category of mobile elements, named transpovirons, was reported. Transpovirons are linear DNA elements of ~7 kb that encompass 6–8 protein-coding genes and have terminal inverted repeats of ~530 bp (REF. 6). Transpovirons are randomly integrated into giant virus genomes and are strictly dependent on giant viruses for their replication and spread; they replicate in the mimivirus factory and accumulate inside mimivirus particles, virophage particles and the amoebal cytoplasm<sup>6</sup>. Distinct transpovirons associate with different mimiviruses, including, for example, mamavirus and courdo7 virus isolates, which are classified in mimivirus lineage A and lineage C, respectively.

In evolution, the giant virus mobilome represents a vehicle and route for genetic novelty and adaptation. Thus, a substantial proportion of genes in the genomes of mimiviruses and other giant viruses are predicted to have been exchanged with bacteria, archaea, eukaryotes (including their amoebal hosts) and other viruses, although the direction of putative transfers is often unclear<sup>26,103</sup>. This high level of genome mosaicism might be related to the co-infection of amoebae by giant viruses and microorganisms, such as bacteria or fungi<sup>41,103</sup>. Indeed, giant viruses can multiply within amoebae that are infected with other microorganisms, which provides opportunities for horizontal gene transfer. *Acanthamoeba* spp. are well-known hosts for several bacteria and some fungi, in addition to viruses<sup>104</sup>. *Acanthamoeba* spp. that were isolated from a contact lens cleaning liquid were found to be co-infected with a mimivirus and two bacteria<sup>93</sup>, and *Acanthamoeba* spp. have been observed experimentally to be co-infected with marseillevirus and two bacteria<sup>41</sup>. In addition, the fact that, in their natural environment, giant viruses are sympatric with other microorganisms that infect *Acanthamoeba* helps to explain the large size of their genomes, which accumulate genes that increase their fitness against other microorganisms that replicate in amoebae<sup>103,105</sup>. Interestingly, a 16% decrease in genome size was observed by co-culturing APMV in amoebae in the absence of other microorganisms<sup>105</sup>; it was determined that the APMV genes that were inactivated were primarily those with the lowest expression, which suggests the loss of obsolete genes<sup>106</sup>. Moreover, these gene losses were associated with the loss of the external fibres that cover the APMV capsid and a decreased rate of replication<sup>105</sup>.

Sympatric  
Inhabiting the same ecological  
niche.

## Seroconversion

The development of antibodies in the blood that are specifically directed against an infectious agent and become detectable by serological tests.

## Box 4 | Do giant viruses of amoebae and virophages infect humans?

The detection of giant viruses of amoebae in humans and the study of their potential pathogenicity are emerging fields<sup>107</sup>. Mimiviruses have been detected in serological, molecular and culture studies of patients with pneumonia<sup>107</sup>; serological studies were almost exclusively conducted in our laboratory, although some of the sera samples were collected by other laboratories<sup>61,107</sup>. Serological evidence notably included seroconversion to *Acanthamoeba polyphaga* mimivirus (APMV) in patients with unexplained pneumonia, including in a laboratory technician who manipulated large quantities of APMV<sup>108</sup>. In addition, two febrile Laotian patients were seropositive for the mimivirus virophage<sup>122</sup>. In culture studies, a mimivirus was isolated from two Tunisian patients who had pneumonia; in one case it was isolated from bronchoalveolar fluid<sup>107,109</sup>. As recently reviewed (see REFS. 61, 107), several studies have failed to detect APMV DNA in respiratory samples from patients with pneumonia and only three cases were positive for APMV DNA when samples were analysed by PCR<sup>123,109,124</sup>. This may indicate that APMV is only rarely associated with pneumonia. Alternatively, this may be the consequence of considerable genetic diversity in mimiviruses, which may have prevented their detection by PCR in these studies<sup>125</sup>. Finally, mice that were inoculated intracardially with APMV developed pneumonia lesions<sup>126</sup>, and APMV can enter, and replicate in, human monocytes, macrophages<sup>76</sup> and peripheral blood mononuclear cells<sup>86</sup>, although the virus was not propagated using these human blood cells.

Marseilleviruses were first serendipitously isolated from the stool of a healthy Senegalese individual<sup>61</sup> and were subsequently detected in human blood by metagenomics, serology, fluorescence *in situ* hybridization (FISH) and PCR<sup>87,110,127,128</sup>. However, the detection of marseillevirus in human blood is controversial, as two other studies failed to detect marseillevirus DNA in blood from polytransfused patients in Paris and Cameroon and in blood from blood donors in southern France, Paris, the United States and Burkina Faso<sup>129,130</sup>. By contrast, positive serologies for Lausannevirus, which is a marseillevirus, were detected in young healthy adults in Switzerland (immunoglobulin G (IgG), 1.7–2.5%)<sup>127</sup>. In 2013, a high antibody titre to marseillevirus was reported in an 11-month-old child who presented with symptoms of lymphadenitis; marseillevirus was also detected by FISH and immunohistochemistry in a lymph node of this individual, and by PCR in the serum<sup>110</sup>. Next, marseillevirus was detected in a lymph node of a 30-year-old woman who was diagnosed with Hodgkin lymphoma by PCR, FISH, immunofluorescence and immunohistochemistry, concurrently, with specific antibodies<sup>111</sup>. Finally, marseillevirus was detected twice, with a one-year interval, by PCR in the pharynx of a patient with neurological disorders and once in the blood<sup>128</sup>. Very few data are available for the infection of humans by giant viruses that have only recently been described, but sequences that might belong to these giant viruses, including pithoviruses, pandoraviruses, faustoviruses, as well as to mimiviruses, marseilleviruses and virophages, have been identified in metagenomes that were generated from human samples<sup>61–64,98</sup>.

## Conclusions and future directions

The replication cycle, structure, genomic make-up and plasticity of giant viruses differ from those of traditional viruses. They have virus particles that are as large as some microorganisms and have a stunning level of complexity. Their genomes are mosaic and contain large repertoires of genes, some of which are hallmarks of cellular organisms, although the majority have unknown functions. These giant viruses enter amoebae through phagocytosis and replicate inside viral factories. In addition, as shown for mimiviruses, they are associated with a specific mobilome and are parasitized by viruses that they can defend against. Giant viruses not only challenge the classification of viruses but also raise intriguing questions about their origin. They extend the definition of viruses into a broader range of biological entities, some of which are very simple and others of which have a complexity comparable to that of other microorganisms.

More giant viruses of amoebae must be identified to gain a better knowledge of their prevalence and diversity, which is largely unexplored but has expanded over the past three years and is likely to expand further now that we have learnt not to neglect them and owing to

high-throughput detection and isolation strategies (BOX 3). The role that giant viruses of amoebae have in evolution also warrants further attention. Several hypotheses on their ancient origin and their evolutionary relationship with cellular organisms have been proposed, and these topics should continue to be debated. In these viruses, it will be important to search for a translation system that does not involve ribosomes, and/or that is anterior to the ribosome and of which we would find remnants. Finally, the detection of giant viruses of amoebae in humans and the study of their potential pathogenicity are emerging fields<sup>107</sup>. To date, giant viruses of amoebae that have been linked to humans were discovered before 2013 (BOX 4). Mimiviruses were associated with pneumonia<sup>108,109</sup>, whereas marseillevirus was detected in the blood and the lymphoid tissue, and was associated with adenitis and lymphoma<sup>110,111</sup>. More systematic investigations of human samples for giant viruses of amoebae must be conducted; such research is more accessible to the next generation of virologists who have entered the field after the description of APMV, and will surely reveal more giant viruses and refine their definition for years to come.

1. La Scola, B. *et al.* A giant virus in amoebae. *Science* **299**, 2033 (2003).

**This study describes the first giant virus of amoebae (APMV) and the circumstances behind its isolation and discovery.**

2. Raoult, D. *et al.* The 1.2-megabase genome sequence of Mimivirus. *Science* **306**, 1344–1350 (2004).

**This study deciphers the APMV genome and gene**

**repertoire, and highlights their unique position in the viral world.**

3. Raoult, D. TRUC or the need for a new microbial classification. *Intervirology* **56**, 349–353 (2013).

4. Scheid, P., Balczun, C. & Schaub, G. A. Some secrets are revealed: parasitic keratitis amoebae as vectors of the scarcely described pandoraviruses to humans. *Parasitol. Res.* **113**, 3759–3764 (2014).

5. La Scola, B. *et al.* The viroplasm as a unique parasite of the giant mimivirus. *Nature* **455**, 100–104 (2008).

**This study describes the discovery of small viruses that replicate within mimivirus factories and impair their growth, which led to the coining of the term viroplasm.**

6. Desnues, C. *et al.* Provirophages and transprovions as the diverse mobilome of giant viruses. *Proc. Natl Acad. Sci. USA* **109**, 18078–18083 (2012).

- This study presents evidence that the genomes of mimiviruses have a specific mobilome that contains virophages and transposons.**
7. Levasseur, A. *et al.* MIMIVIRE is a defence system in mimivirus that confers resistance to virophage. *Nature* **531**, 249–252 (2016).  
**This study shows that mimiviruses can defend against virophages that infect their viral factory.**
  8. Raoult, D. & Forterre, P. Redefining viruses: lessons from Mimivirus. *Nat. Rev. Microbiol.* **6**, 315–319 (2008).  
**This study suggests that, based on the properties of giant viruses of amoebae, viruses should be redefined as capsid-encoding organisms.**
  9. Filee, J., Siguier, P. & Chandler, M. I am what I eat and I eat what I am: acquisition of bacterial genes by giant viruses. *Trends Genet.* **23**, 10–15 (2007).
  10. Moreira, D. & Lopez-Garcia, P. Ten reasons to exclude viruses from the tree of life. *Nat. Rev. Microbiol.* **7**, 306–311 (2009).
  11. Raoult, D. There is no such thing as a tree of life (and of course viruses are out!). *Nat. Rev. Microbiol.* **7**, 615 (2009).
  12. Moreira, D. & Lopez-Garcia, P. Evolution of viruses and cells: do we need a fourth domain of life to explain the origin of eukaryotes? *Phil. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20140327 (2015).
  13. Forterre, P. & Gaia, M. Giant viruses and the origin of modern eukaryotes. *Curr. Opin. Microbiol.* **31**, 44–49 (2016).  
**This study provides a summary of, and discussion on, the findings and theories of the origin of giant viruses of amoebae.**
  14. Klose, T. *et al.* A Mimivirus enzyme that participates in viral entry. *Structure* **23**, 1058–1065 (2015).
  15. Rodrigues, R. A. *et al.* Mimivirus fibrils are important for viral attachment to microbial world by a diverse glycoside interaction repertoire. *J. Virol.* **89**, 11812–11819 (2015).
  16. Zauberman, N. *et al.* Distinct DNA exit and packaging portals in the virus *Acanthamoeba polyphaga* mimivirus. *PLoS Biol.* **6**, e114 (2008).
  17. Kuznetsov, Y. G. *et al.* Atomic force microscopy investigation of the giant mimivirus. *Virology* **404**, 127–137 (2010).
  18. Xiao, C. *et al.* Structural studies of the giant mimivirus. *PLoS Biol.* **7**, e92 (2009).
  19. Renesto, P. *et al.* Mimivirus giant particles incorporate a large fraction of anonymous and unique gene products. *J. Virol.* **80**, 11678–11685 (2006).
  20. Legendre, M., Santini, S., Rico, A., Abergel, C. & Claverie, J. M. Breaking the 1000-gene barrier for Mimivirus using ultra-deep genome and transcriptome sequencing. *Viol. J.* **8**, 99 (2011).
  21. Silva, L. C. *et al.* Modulation of the expression of mimivirus-encoded translation-related genes in response to nutrient availability during *Acanthamoeba castellanii* infection. *Front. Microbiol.* **6**, 539 (2015).
  22. Legendre, M. *et al.* mRNA deep sequencing reveals 75 new genes and a complex transcriptional landscape in Mimivirus. *Genome Res.* **20**, 664–674 (2010).  
**This study describes the first analysis of the transcriptome of a giant virus of amoebae.**
  23. Abergel, C., Rudinger-Thirion, J., Giege, R. & Claverie, J. M. Virus-encoded aminoacyl-tRNA synthetases: structural and functional characterization of mimivirus TyrRS and MetRS. *J. Virol.* **81**, 12406–12417 (2007).
  24. Suhre, K. Gene and genome duplication in *Acanthamoeba polyphaga* Mimivirus. *J. Virol.* **79**, 14095–14101 (2005).
  25. Moreira, D. & Brochier-Armanet, C. Giant viruses, giant chimeras: the multiple evolutionary histories of Mimivirus genes. *BMC Evol. Biol.* **8**, 12 (2008).
  26. Forterre, P. Giant viruses: conflicts in revisiting the virus concept. *Intervirology* **53**, 362–378 (2010).
  27. Boyer, M., Gimenez, G., Suzan-Monti, M. & Raoult, D. Classification and determination of possible origins of ORFans through analysis of nucleocytoplasmic large DNA viruses. *Intervirology* **53**, 310–320 (2010).
  28. Byrne, D. *et al.* The polyadenylation site of Mimivirus transcripts obeys a stringent 'hairpin rule'. *Genome Res.* **19**, 1233–1242 (2009).
  29. Jeudy, S., Abergel, C., Claverie, J. M. & Legendre, M. Translation in giant viruses: a unique mixture of bacterial and eukaryotic termination schemes. *PLoS Genet.* **8**, e1003122 (2012).
  30. Yoosuf, N. *et al.* Related giant viruses in distant locations and different habitats: *Acanthamoeba polyphaga* momovirus represents a third lineage of the *Mimiviridae* that is close to the megavirus lineage. *Genome Biol. Evol.* **4**, 1324–1330 (2012).
  31. Azza, S., Cambillau, C., Raoult, D. & Suzan-Monti, M. Revised Mimivirus major capsid protein sequence reveals intron-containing gene structure and extra domain. *BMC Mol. Biol.* **10**, 39 (2009).
  32. La Scola, B., de Lamballerie, X. N., Claverie, J. M., Drancourt, M. & Raoult, D. *Genus Mimivirus in Virus Taxonomy* (eds Fauquet, M., Mayo, M. A., Maniloff, J., Desselberger, U. & Ball, L. A.) 275–276 (Elsevier Academic Press, 2005).
  33. Pagnier, I. *et al.* A decade of improvements in *Mimiviridae* and *Marseilleviridae* isolation from amoeba. *Intervirology* **56**, 354–363 (2013).
  34. Khalil, J. Y., Andreani, J. & La Scola, B. Updating strategies for isolating and discovering giant viruses. *Curr. Opin. Microbiol.* **31**, 80–87 (2016).
  35. Khalil, J. Y. *et al.* High-throughput isolation of giant viruses in liquid medium using automated flow cytometry and fluorescence staining. *Front. Microbiol.* **7**, 26 (2016).
  36. Arslan, D., Legendre, M., Seltzer, V., Abergel, C. & Claverie, J. M. Distant Mimivirus relative with a larger genome highlights the fundamental features of *Megaviridae*. *Proc. Natl Acad. Sci. USA* **108**, 17486–17491 (2011).
  37. Fischer, M. G., Allen, M. J., Wilson, W. H. & Suttle, C. A. Giant virus with a remarkable complement of genes infects marine zooplankton. *Proc. Natl Acad. Sci. USA* **107**, 19508–19513 (2010).
  38. Yau, S. *et al.* Virophage control of antarctic algal host-virus dynamics. *Proc. Natl Acad. Sci. USA* **108**, 6163–6168 (2011).
  39. Santini, S. *et al.* Genome of *Phaeocystis globosa* virus PpV-16T highlights the common ancestry of the largest known DNA viruses infecting eukaryotes. *Proc. Natl Acad. Sci. USA* **110**, 10800–10805 (2013).
  40. Yutin, N., Colson, P., Raoult, D. & Koonin, E. V. *Mimiviridae*: clusters of orthologous genes, reconstruction of gene repertoire evolution and proposed expansion of the giant virus family. *Viol. J.* **10**, 106–110 (2013).
  41. Boyer, M. *et al.* Giant marseillevirus highlights the role of amoebae as a melting pot in emergence of chimeric microorganisms. *Proc. Natl Acad. Sci. USA* **106**, 21848–21853 (2009).
  42. Colson, P. *et al.* "Marseilleviridae", a new family of giant viruses infecting amoebae. *Arch. Virol.* **158**, 915–920 (2013).
  43. Philippe, N. *et al.* Pandoraviruses: amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic eukaryotes. *Science* **341**, 281–286 (2013).  
**This study shows that giant viruses of amoebae, the particles and genomes of which are larger than those of APMV, exist and have an atypical morphology with no detectable capsid protein.**
  44. Legendre, M. *et al.* Thirty-thousand-year-old distant relative of giant icosahedral DNA viruses with a pandoravirus morphology. *Proc. Natl Acad. Sci. USA* **111**, 4274–4279 (2014).  
**This study describes the most ancient giant virus of amoebae, which was isolated from permafrost.**
  45. Reteno, D. G. *et al.* Faustovirus, an asfarvirus-related new lineage of giant viruses infecting amoebae. *J. Virol.* **89**, 6585–6594 (2015).  
**This study shows that new giant viruses of amoebae can be discovered using amoebae other than *Acanthamoeba* spp. as culture support.**
  46. Legendre, M. *et al.* In-depth study of Mollivirus sibiricum, a new 30,000-y-old giant virus infecting *Acanthamoeba*. *Proc. Natl Acad. Sci. USA* **112**, E5327–E5335 (2015).
  47. Iyer, L. M., Aravind, L. & Koonin, E. V. Common origin of four diverse families of large eukaryotic DNA viruses. *J. Virol.* **75**, 11720–11734 (2001).
  48. Yutin, N., Wolf, Y. I., Raoult, D. & Koonin, E. V. Eukaryotic large nucleocytoplasmic DNA viruses: clusters of orthologous genes and reconstruction of viral genome evolution. *Viol. J.* **17**, 223 (2009).
  49. Yutin, N. & Koonin, E. V. Hidden evolutionary complexity of nucleocytoplasmic large DNA viruses of eukaryotes. *Viol. J.* **9**, 161 (2012).
  50. Boyer, M., Madoui, M. A., Gimenez, G., La Scola, B. & Raoult, D. Phylogenetic and phyletic studies of informational genes in genomes highlight existence of a 4 domain of life including giant viruses. *PLoS ONE* **5**, e15530 (2010).  
**This study presents additional evidence that giant viruses of amoebae may comprise a fourth branch of the tree of life.**
  51. Nasir, A. & Caetano-Anolles, G. A phylogenomic data-driven exploration of viral origins and evolution. *Sci. Adv.* **1**, e1500527 (2015).
  52. Koonin, E. V. & Yutin, N. Origin and evolution of eukaryotic large nucleocytoplasmic DNA viruses. *Intervirology* **53**, 284–292 (2010).
  53. Colson, P. *et al.* "Megavirales", a proposed new order for eukaryotic nucleocytoplasmic large DNA viruses. *Arch. Virol.* **158**, 2517–2521 (2013).
  54. Sharma, V., Colson, P., Giorgi, R., Pontarotti, P. & Raoult, D. DNA-dependent RNA polymerase detects hidden giant viruses in published databanks. *Genome Biol. Evol.* **6**, 1603–1610 (2014).  
**This study describes the use of DNA-dependent RNA polymerase genes as bait to explore the dark matter of metagenomes and detect sequences that are related to giant viruses of amoebae.**
  55. Sharma, V., Colson, P., Pontarotti, P. & Raoult, D. Mimivirus inaugurated in the 21st century the beginning of a reclassification of viruses. *Curr. Opin. Microbiol.* **31**, 16–24 (2016).
  56. Suzan-Monti, M., La Scola, B., Barrassi, L., Espinosa, L. & Raoult, D. Ultrastructural characterization of the giant volcano-like virus factory of *Acanthamoeba polyphaga* Mimivirus. *PLoS ONE* **2**, e328 (2007).
  57. Abergel, C., Legendre, M. & Claverie, J. M. The rapidly expanding universe of giant viruses: Mimivirus, Pandoravirus, Pithovirus and Mollivirus. *FEMS Microbiol. Rev.* **39**, 779–796 (2015).
  58. Aherfi, S., Colson, P., La Scola, B. & Raoult, D. Giant viruses of amoebae: an update. *Front. Microbiol.* **7**, 349 (2016).
  59. Ghedin, E. & Claverie, J. M. Mimivirus relatives in the Sargasso sea. *Viol. J.* **2**, 62 (2005).
  60. Kristensen, D. M., Mushegian, A. R., Dolja, V. V. & Koonin, E. V. New dimensions of the virus world discovered through metagenomics. *Trends Microbiol.* **18**, 11–19 (2010).
  61. Colson, P. *et al.* Evidence of the megavirome in humans. *J. Clin. Virol.* **57**, 191–200 (2013).
  62. Law, J. *et al.* Identification of hepatotropic viruses from plasma using deep sequencing: a next generation diagnostic tool. *PLoS ONE* **8**, e60595 (2013).
  63. Verneau, J., Levasseur, A., Raoult, D., La Scola, B. & Colson, P. MG-Digger: an automated pipeline to search for giant virus-related sequences in metagenomes. *Front. Microbiol.* **7**, 428 (2016).
  64. Rampelli, S. *et al.* ViromeScan: a new tool for metagenomic viral community profiling. *BMC Genomics* **17**, 165–2446 (2016).
  65. Maumus, F., Epert, A., Nogue, F. & Blanc, G. Plant genomes enclose footprints of past infections by giant virus relatives. *Nat. Commun.* **5**, 4268 (2014).
  66. Wu, D. *et al.* Stalking the fourth domain in metagenomic data: searching for, discovering, and interpreting novel, deep branches in marker gene phylogenetic trees. *PLoS ONE* **6**, e18011 (2011).
  67. Mozar, M. & Claverie, J. M. Expanding the *Mimiviridae* family using asparagine synthase as a sequence bait. *Virology* **466–467**, 112–122 (2014).
  68. Klose, T. *et al.* Structure of faustovirus, a large dsDNA virus. *Proc. Natl Acad. Sci. USA* **113**, 6206–6211 (2016).
  69. Thomas, V. *et al.* Lausannevirus, a giant amoebal virus encoding histone doublets. *Environ. Microbiol.* **13**, 1454–1466 (2011).
  70. Sun, C., Feschotte, C., Wu, Z. & Mueller, R. L. DNA transposons have colonized the genome of the giant virus Pandoravirus salinus. *BMC Biol.* **13**, 38–0145 (2015).
  71. Filee, J. Genomic comparison of closely related giant viruses supports an accordion-like model of evolution. *Front. Microbiol.* **6**, 593 (2015).
  72. Levasseur, A. *et al.* Comparison of a modern and fossil pithovirus reveals its genetic conservation and evolution. *Genome Biol. Evol.* **8**, 2333–2339 (2016).
  73. Babkin, I. V. & Babkina, I. N. Molecular dating in the evolution of vertebrate poxviruses. *Intervirology* **54**, 253–260 (2011).
  74. Doutre, G., Philippe, N., Abergel, C. & Claverie, J. M. Genome analysis of the first *Marseilleviridae* representative from Australia indicates that most of its genes contribute to the virus fitness. *J. Virol.* **88**, 14340–14349 (2014).
  75. Mutsafi, Y., Zauberman, N., Sabanay, I. & Minsky, A. Vaccinia-like cytoplasmic replication of the giant Mimivirus. *Proc. Natl Acad. Sci. USA* **107**, 5978–5982 (2010).
  76. Ghigo, E. *et al.* Amoebal pathogen mimivirus infects macrophages through phagocytosis. *PLoS Pathog.* **4**, e1000087 (2008).

77. Mutsafi, Y., Fridmann-Sirkis, Y., Milrot, E., Hevroni, L. & Minsky, A. Infection cycles of large DNA viruses: emerging themes and underlying questions. *Virology* **466–467**, 3–14 (2014).
78. Mutsafi, Y., Shimoni, E., Shimon, A. & Minsky, A. Membrane assembly during the infection cycle of the giant Mimivirus. *PLoS Pathog.* **9**, e1003367 (2013).
79. Arantes, T. S. *et al.* The large marseillevirus explores different entry pathways by forming giant infectious vesicles. *J. Virol.* **90**, 5246–5255 (2016).
80. Boratto, P. V. *et al.* Amoebas as mimivirus bunkers: increased resistance to UV light, heat and chemical biocides when viruses are carried by amoeba hosts. *Arch. Virol.* **159**, 1039–1043 (2013).
81. Dornas, F. P. *et al.* *Acanthamoeba polyphaga* mimivirus stability in environmental and clinical substrates: implications for virus detection and isolation. *PLoS ONE* **9**, e87811 (2014).
82. Silmani, M., Pagnier, I., Boughalmi, M., Raoult, D. & La Scola, B. Alcohol disinfection procedure for isolating giant viruses from contaminated samples. *Intervirology* **56**, 434–440 (2013).
83. Boughalmi, M. *et al.* First isolation of a giant virus from wild *Hirudo medicinalis* leech: *Mimiviridae* isolation in *Hirudo medicinalis*. *Viruses* **5**, 2920–2930 (2013).
84. Dornas, F. P. *et al.* Mimivirus circulation among wild and domestic mammals, Amazon Region, Brazil. *Emerg. Infect. Dis.* **20**, 469–472 (2014).
85. Andrade, K. R. *et al.* Oysters as hot spots for mimivirus isolation. *Arch. Virol.* **160**, 477–482 (2015).
86. Silva, L. C. *et al.* A resourceful giant: APMV is able to interfere with the human type I Interferon system. *Microbes Infect.* **16**, 187–195 (2014).
87. Popgeorgiev, N. *et al.* Marseillevirus-like virus recovered from blood donated by asymptomatic humans. *J. Infect. Dis.* **208**, 1042–1050 (2013). **This study details the first detection of a giant virus of amoebae in human blood.**
88. Netherton, C. L. & Wileman, T. Virus factories, double membrane vesicles and viroplasm generated in animal cells. *Curr. Opin. Virol.* **1**, 381–387 (2011).
89. Claverie, J. M. Viruses take center stage in cellular evolution. *Genome Biol.* **7**, 110 (2006).
90. Yutin, N., Raoult, D. & Koonin, E. V. Virophages, polintons, and transpovirons: a complex evolutionary network of diverse selfish genetic elements with different reproduction strategies. *Virology* **450**, 158 (2013).
91. Krupovic, M., Kuhn, J. H. & Fischer, M. G. A classification system for virophages and satellite viruses. *Arch. Virol.* **161**, 233–247 (2016).
92. Gaia, M. *et al.* Broad spectrum of *Mimiviridae* virophage allows its isolation using a mimivirus reporter. *PLoS ONE* **8**, e61912 (2013).
93. Cohen, G., Hoffart, L., La Scola, B., Raoult, D. & Drancourt, M. Ameba-associated Keratitis, France. *Emerg. Infect. Dis.* **17**, 1306–1308 (2011).
94. Campos, R. K. *et al.* Samba virus: a novel mimivirus from a giant rain forest, the Brazilian Amazon. *Virology* **466–467**, 95 (2014).
95. Gaia, M. *et al.* Zamilon, a novel virophage with *Mimiviridae* host specificity. *PLoS ONE* **9**, e94923 (2014).
96. Fischer, M. G. & Suttle, C. A. A virophage at the origin of large DNA transposons. *Science* **332**, 231–234 (2011).
- This study describes the discovery of a virophage of a distant mimivirus and its evolutionary link to polinton transposons.**
97. Blanc, G., Gallot-Lavallee, L. & Maumus, F. Provirophages in the *Bigelowiella* genome bear testimony to past encounters with giant viruses. *Proc. Natl Acad. Sci. USA* **112**, E5318–E5326 (2015).
98. Zhou, J. *et al.* Diversity of virophages in metagenomic data sets. *J. Virol.* **87**, 4225–4236 (2013).
99. Claverie, J. & Abergel, C. Mimivirus and its virophage. *Annu. Rev. Genet.* **43**, 49–66 (2009).
100. Mohanraju, P. *et al.* Diverse evolutionary roots and mechanistic variations of the CRISPR–Cas systems. *Science* **353**, aad5147 (2016).
101. Marraffini, L. A. CRISPR–Cas immunity in prokaryotes. *Nature* **526**, 55–61 (2015).
102. Claverie, J. M. & Abergel, C. CRISPR–Cas-like system in giant viruses: why MIMIVIRE is not likely to be an adaptive immune system. *Virology* **51**, 193–196 (2016).
103. Raoult, D. & Boyer, M. Amoebae as genitors and reservoirs of giant viruses. *Intervirology* **53**, 321–329 (2010).
104. Greub, G. & Raoult, D. Microorganisms resistant to free-living amoebae. *Clin. Microbiol. Rev.* **17**, 413–433 (2004).
105. Boyer, M. *et al.* Mimivirus shows dramatic genome reduction after intraamoebal culture. *Proc. Natl Acad. Sci. USA* **108**, 10296–10301 (2011).
106. Colson, P. & Raoult, D. Lamarckian evolution of the giant Mimivirus in allopatric laboratory culture on amoebae. *Front. Cell. Infect. Microbiol.* **2**, 91 (2012).
107. Colson, P., Aherfi, S., La Scola, B. & Raoult, D. The role of giant viruses of amoebas in humans. *Curr. Opin. Microbiol.* **31**, 199–208 (2016).
108. Raoult, D., Renesto, P. & Brouqui, P. Laboratory infection of a technician by mimivirus. *Ann. Intern. Med.* **144**, 702–703 (2006).
109. Saadi, H. *et al.* First isolation of Mimivirus in a patient with pneumonia. *Clin. Infect. Dis.* **57**, e127–e134 (2013). **This study describes the first isolation of a mimivirus from a human sample, specifically the bronchoalveolar fluid of a patient that presented with unexplained pneumonia.**
110. Popgeorgiev, N., Michel, G., Lepidi, H., Raoult, D. & Desnues, C. Marseillevirus adenitis in an 11-month-old child. *J. Clin. Microbiol.* **51**, 4102–4105 (2013).
111. Aherfi, S. *et al.* Marseillevirus in lymphoma: a giant in the lymph node. *Lancet Infect. Dis.* **16**, e225–e234 (2016).
112. Lwoff, A. The concept of virus. *J. Gen. Microbiol.* **17**, 239–253 (1957).
113. Lwoff, A. & Tournier, P. The classification of viruses. *Annu. Rev. Microbiol.* **20**, 45–74 (1966).
114. Williams, T. A., Embley, T. M. & Heinz, E. Informational gene phylogenies do not support a fourth domain of life for nucleocytoplasmic large DNA viruses. *PLoS ONE* **6**, e21080 (2011).
115. Yutin, N., Wolf, Y. I. & Koonin, E. V. Origin of giant viruses from smaller DNA viruses not from a fourth domain of cellular life. *Virology* **466–467**, 38–52 (2014).
116. Rowbotham, T. J. Isolation of *Legionella pneumophila* from clinical specimens via amoebae, and the interaction of those and other isolates with amoebae. *J. Clin. Pathol.* **36**, 978–986 (1983).
117. Pagnier, I., Valles, C., Raoult, D. & La Scola, B. Isolation of *Vermamoeba vermiformis* and associated bacteria in hospital water. *Microb. Pathog.* **80**, 14–20 (2015).
118. Dornas, F. P. *et al.* Isolation of new Brazilian giant viruses from environmental samples using a panel of protozoa. *Front. Microbiol.* **6**, 1086 (2015).
119. Rowbotham, T. J. Preliminary report on the pathogenicity of *Legionella pneumophila* for freshwater and soil amoebae. *J. Clin. Pathol.* **33**, 1179–1183 (1980).
120. Birtles, R. J., Rowbotham, T. J., Raoult, D. & Harrison, T. G. Phylogenetic diversity of intra-amoebal legionellae as revealed by 16S rRNA gene sequence comparison. *Microbiology* **142**, 3525–3530 (1996).
121. Birtles, R. J., Rowbotham, T. J., Storey, C., Marrie, T. J. & Raoult, D. Chlamydia-like obligate parasite of free-living amoebae. *Lancet* **349**, 925–926 (1997).
122. Parola, P. *et al.* *Acanthamoeba polyphaga* mimivirus virophage seroconversion in travelers returning from Laos. *Emerg. Infect. Dis.* **18**, 1500–1502 (2012).
123. La Scola, B., Marrie, T. J., Auffray, J. P. & Raoult, D. Mimivirus in pneumonia patients. *Emerg. Infect. Dis.* **11**, 449–452 (2005).
124. Zhang, X. A. *et al.* Lack of Mimivirus detection in patients with respiratory disease, China. *Emerg. Infect. Dis.* **22**, 10 (2016).
125. Ngounga, T. *et al.* Real-time PCR systems targeting giant viruses of amoebae and their virophages. *Intervirology* **56**, 413–423 (2013).
126. Khan, M., La Scola, B., Lepidi, H. & Raoult, D. Pneumonia in mice inoculated experimentally with *Acanthamoeba polyphaga* mimivirus. *Microb. Pathog.* **42**, 56–61 (2007).
127. Mueller, L., Baud, D., Bertelli, C. & Greub, G. Lausannevirus seroprevalence among asymptomatic young adults. *Intervirology* **56**, 430–433 (2013).
128. Aherfi, S., Colson, P. & Raoult, D. Marseillevirus in the pharynx of a patient with neurologic disorders. *Emerg. Infect. Dis.* **22**, 2008–2010 (2016).
129. Sauvage, V. *et al.* No evidence of marseillevirus-like virus presence in blood donors and recipients of multiple blood transfusions. *J. Infect. Dis.* **210**, 2017–2018 (2014).
130. Phan, T. G. *et al.* Absence of giant blood Marseille-like virus DNA detection by polymerase chain reaction in plasma from healthy US blood donors and serum from multiply transfused patients from Cameroon. *Transfusion* **55**, 1256–1262 (2015).

**Acknowledgements**

The authors thank I. Pagnier, J. Bou Khalil and J. Andreani for providing electron microscopy images.

**Competing interests statement**

The authors declare no competing interests.

**SUPPLEMENTARY INFORMATION**

See online article: S1 (table) | S2 (figure) | S3 (table) | S4 (figure) | S5 (table) | S6 (figure) | S7 (figure)

ALL LINKS ARE ACTIVE IN THE ONLINE PDF