



## BIROn - Birkbeck Institutional Research Online

Senju, Atsushi and Southgate, Victoria and White, S. and Frith, U. (2009) Mindblind eyes: an absence of spontaneous theory of mind in Asperger syndrome. *Science* 325 (5942), pp. 883-885. ISSN 0036-8075.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/2566/>

*Usage Guidelines:*

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>  
contact [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk).

or alternatively



## **BIROn** - Birkbeck Institutional Research Online

---

Enabling open access to Birkbeck's published research output

### Mindblind eyes: an absence of spontaneous theory of mind in Asperger syndrome

#### **Journal Article**

<http://eprints.bbk.ac.uk/2566>

Version: Accepted (Refereed)

#### **Citation:**

Senju, A.; Southgate, V.; White, S.; Frith, U. (2009) Mindblind eyes: an absence of spontaneous theory of mind in Asperger syndrome <i>Science</i> 325 (5942), pp.883-885
---

© 2009 AAAS

[Publisher version](#)

---

All articles available through Birkbeck ePrints are protected by intellectual property law, including copyright law. Any use made of the contents should comply with the relevant law.

---

[Deposit Guide](#)

Contact: [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk)

*SCIENCE VOL 325 14 AUGUST 2009 PAGES 883-885*

## **Mindblind eyes: an absence of spontaneous Theory of Mind in Asperger Syndrome**

Atsushi Senju\*<sup>a</sup>, Victoria Southgate<sup>a</sup>, Sarah White<sup>b</sup>, and Uta Frith<sup>b,c</sup>

<sup>a</sup>Centre for Brain and Cognitive Development, Birkbeck, University of London, London, UK

<sup>b</sup>Institute of Cognitive Neuroscience, University College London, London, UK

<sup>c</sup>CFIN Aarhus University

\* Correspondence concerning this paper should be sent to Atsushi Senju, Centre for Brain and Cognitive Development, Birkbeck, University of London, Malet Street, London WC1E 7HX, UK; Tel: +44 207 631 6895; Fax: +44 207 631 6587; E-mail: a.senju@bbk.ac.uk

### **Abstract**

Adults with Asperger syndrome can understand mental states such as desires and beliefs (mentalizing) when explicitly prompted to do so, despite having impairments in social communication. We directly tested the hypothesis that such individuals nevertheless fail to mentalize spontaneously. To this end, we used an eye tracking task that has revealed the spontaneous ability to mentalize in typically-developing infants. We showed that, like infants, neurotypical adults' (N = 17) eye movements anticipated an actor's behaviour on the basis of her false belief. This was not the case for individuals with Asperger syndrome (N = 19). Thus, these individuals do not attribute mental states spontaneously, but may be able to do so in explicit tasks through compensatory learning.


Impairment in reciprocal social interaction and communication is a core feature of Autism Spectrum Disorders, regardless of age and ability. This core feature is manifest in a wide range of social impairments, including characteristic deficits in comprehension and use of pretend play, expressive gestures, deception and irony (1). One influential account that can explain these varied and characteristic impairments proposes that they are a consequence of a failure in the neurologically-based capacity to 'mentalize', that is, the automatic ability to attribute mental states to self and others. The first evidence for this hypothesis, which is also known as a deficit in Theory of Mind or 'mindblindness' (2, 3), comes from the finding that children with autism fail the verbally instructed Sally-Ann False Belief Task (FBT), while four-year-old neurotypical children, and children with Down syndrome of similar verbal mental age, pass (4).

In this task, considered a stringent-test of Theory of Mind (5), one character (Sally), places a marble in a basket and leaves the room. In her absence, another character (Ann) moves the marble to a box. When Sally returns, children are asked where she will look for her marble. If children understand that Sally's actions will be based on what she believes to be true, rather than the actual state of affairs, they should answer that she will look in the basket, rather than the box. This correct answer requires the child to predict Sally's behaviour based on her now false belief.

Despite still exhibiting atypical social features characteristic of autism, individuals of

higher verbal ability, and in particular those with Asperger syndrome, can pass such False Belief attribution tasks (6-9). This competence presents a puzzle for the mindblindness hypothesis (10), and has prompted the proposal that these high ability individuals have acquired the ability to reason explicitly about false beliefs by compensatory learning, while difficulties in spontaneous mental states attribution may nevertheless persist (11). To date, there is only indirect evidence in support of this hypothesis (12-16). Here we seek to provide direct evidence by contrasting the ability to pass the standard FBT with spontaneous looking behaviour during a nonverbal form of this task.

Spontaneous looking behaviour during a FBT scenario was used in a groundbreaking study by Onishi & Baillargeon (17), which exploited the fact that infants look longer at events that they do not expect. The authors showed that 15-month-old infants looked significantly longer when an actor searched in a location where an object was hidden that she could not know about, that is, when her behaviour was incongruent with her belief. Southgate, Senju & Csibra (18) extended this paradigm so that, rather than measuring whether young children look longer at unexpected outcomes, they measured whether children actually anticipate the outcomes before they happen. They designed a task that made it possible to assess directly whether children had an understanding of the content of an actors' belief. Briefly, 25-month-old children were familiarized to an event in which a puppet hid a ball in one of two boxes (Figure 1A) and then an actor reached through one of two windows to retrieve the ball from the box (Figure 1C). Before she reached, a light and simultaneous chime signalled that the actor was about to open a window to retrieve the hidden object (Figure 1B). In the test trial the puppet transferred the ball from one box to another and then removed it altogether, while the actor was looking away (Figure 1D). An eye-tracker was used to assess whether children expected, by making anticipatory eye-movements, the actor to open the door, which would be consistent with her having a false belief about the location of the ball. Indeed the authors found that these typically developing children made eye movements towards the box, which was consistent with the actor's belief about the location of the ball, despite the fact that it no longer contained the ball. These children, who would not be able to perform the traditional verbally instructed FBT, thus correctly anticipated the actor's behaviour in line with her false belief.



**Figure 1.** Selected scenes from stimulus movies (see also *Movies S1* and *S2*). In familiarization trials, participants were familiarized to an event in which (A) the puppet placed a ball in one of two boxes (B) both windows were illuminated and a chime sounded, and (C) an actor reached through the window above the box in which the ball was placed, and retrieved the ball. The participants were familiarized to the contingency between (B) and (C). In (D), the puppet moves the ball while the actor is looking away. This operation induces a false belief in the actor about the location of the ball.

It is this task that we used for the present study (see also *Movies S1* and *S2*). We asked

whether or not adults with Asperger syndrome would, through their anticipatory looking, reveal a similar spontaneous capacity for false belief attribution. At the same time we had to establish that neurotypical adults would show the same anticipatory looking as young children. Prior to the main analyses, we confirmed that all the participants showed anticipatory looking towards the correct location during familiarization trials (19). Written informed consent was obtained from each participant before the study began. The study was approved by the UCL Research Ethics Committee.

**Table 1.** Mean Chronological Age (CA), verbal IQ (VIQ), performance IQ (PIQ), full scale IQ (FIQ) (WAIS-III UK), composite Theory of mind score (ToM), Strange Story test score (SS), scores of Autism Quotient (AQ) and Autism Diagnostic Observation Schedule - Generic (ADOS-G).

GROUP	Asperger Syndrome			Neurotypical		
	Mean	SD	range	Mean	SD	range
CA	36.8	14.3	21-67	39.6	11.7	26-63
VIQ	116.8	14.4	85-144	116.1	13.2	91-138
PIQ	109.6	13.0	80-132	111.5	10.6	97-132
FIQ	115.6	14.9	89-144	115.3	11.0	95-129
ToM <sup>1</sup>	9.7	2.0	4-13.5	10.6	1.3	8.5-12.5
SS <sup>2</sup>	13.2	1.8	10-16	13.6	1.3	12-16
AQ <sup>3</sup>	34.9	7.6	17-48	16.5	7.6	6-37
ADOS-G	7.9	4.7	0-17	-	-	-

<sup>1</sup>The ToM tests consisted of five first-order false belief tests (*Sally-Ann*: (4); *Smarties*: (23); *Interpretational false belief*: (24); *Belief-emotion* and *Real-apparent emotion*: (25)) and two second-order false belief tests (*Ice Cream Van*: (26); *Coat Story*: (6)).

<sup>2</sup>The *Strange Stories* were taken from (27), and required the participant to either interpret another's behaviour or understand another's emotion.


<sup>3</sup>AS and NT groups differed significantly on the Autism-Spectrum Quotient confirming their diagnostic status (AQ: (28),  $t(34) = 7.23$ ,  $p < .001$ , *Cohen's d* = 2.41, t-test). No other variables were significantly different between the two groups.

As shown in Table 1 The Asperger and Neurotypical groups in our study were very similar in age and IQ. As no sex differences were shown in any of our measures, we pooled results over gender in each group. We found that the participants with Asperger syndrome performed as did neurotypical adults on verbally instructed versions of a variety of standard Theory of mind (ToM) tests, including the previously described Sally-Anne Task. There were no group differences in the composite ToM score or in the Strange Stories (both  $t_s < 1.61$ , both  $p_s > .1$ , t-test). Crucially all participants with Asperger syndrome passed the two standard False Belief Tests.

However, the Eye-Tracking version of the FBT revealed a very different picture. We used Differential Looking Times (DLS; (20)) considered a highly reliable measure of looking bias, (e.g. (21, 22)) compiled over a 6 second period (19). The Asperger group showed significantly less looking bias towards the correct window than did the neurotypical group (*Figure 2A*, main effect of group:  $F(1, 32) = 4.93$ ,  $p = .003$ ,  $\eta_p^2 = 0.13$ , ANOVA). Follow-up t-tests revealed that the neurotypical group scored significantly above zero (mean = 0.42,  $t(16) = 2.76$ ,  $p = .014$ , *Cohen's d* = 0.67, t-test), meaning that they showed a significant bias towards the correct target, in line with the actor's false belief. This was not the case for the Asperger group, whose bias did not differ from zero (mean: -0.001,  $t(18) = -.010$ ,  $p = .99$ ,

*Cohen's d* = 0.002, t-test).

We also coded the direction of first saccade (*Figure 2B*). Here 13 out of 17 neurotypical participants made their first eye movement towards the correct location, which was significantly above chance ( $p = .049$ , binominal test). In contrast, only 8 out of 19 individuals in the Asperger group made a correct saccade first, but this did not significantly differ from chance ( $p = .647$ , binominal test). These results are consistent with the Differential Looking Score reported above. However, the first saccade, which represents a broad categorical response, did not differentiate significantly between the groups ( $p = .30$ , Fisher's exact-test, two-tailed).



**Figure 2.** (A) Mean ( $\pm$  SEM) difference looking scores (DLS, (19)) and (B) the ratio of the number of participants who made correct first saccades in each group. AS: participants with Asperger Syndrome ( $N = 19$ ), NT: neurotypical participants ( $N = 17$ ), \*:  $p < .05$ , \*\*:  $p < .01$ , dotted lines: chance level, statistical test used: (A) t-test and (B) binominal test.

Could our results be due to gaze abnormalities in the Asperger group? Note that they could not be due to an avoidance of eye gaze since the actor's eyes were hidden beneath a visor. However, the duration of fixations to the actor's face were significantly shorter in the Asperger group (mean: 1.9s) than in the neurotypical group (mean: 3.1s) ( $t(34) = -2.48$ ,  $p = .018$ , *Cohen's d* = 0.827, t-test). This was not due to overall shorter fixations in the Asperger group, as the duration of fixations toward the windows (correct and incorrect combined) did not differ between groups (mean: 2.5s in Asperger group and 1.9s in neurotypical group,  $t(34) = 1.53$ ,  $p = .14$ , *Cohen's d* = 0.506, t-test). Importantly, the correlation between DLS and the duration of face fixation was not significant in either the Asperger or neurotypical group. Furthermore, total time spent looking at the five regions of interest (face, two windows and two boxes) did not differ between groups (4.8s in Asperger group and 5.1s in neurotypical group, see supporting online text for further details). Thus, the lack of bias towards the 'correct' target in the Asperger group is not explained by gaze abnormalities.

This study demonstrated that adults with Asperger Syndrome do not spontaneously anticipate others' actions in a non-verbal task, closely modeled on the standard FBT which they pass with ease. In particular, the contrast with neurotypical two-year-olds who show spontaneous looking to the correct location on the same task (18) is striking. It is unlikely that differences in motivation are to blame since neurotypical adults showed the same bias as typically-developing children, and the Asperger group exhibited correct anticipatory looking on familiarization trials when no belief reasoning was required. The current results confirm indirect indications (12-15) that individuals with Asperger Syndrome have a persistent impairment in spontaneous mentalizing and are also consistent with a previous finding (16) that children with autism are more likely to give a correct verbal answer than a correct anticipatory look, when asked to infer someone's preference.

Although it is plausible that the documented early emerging spontaneous capacity to mentalize (17, 18) is a prerequisite for the later ability to justify behaviour in terms of mental states in verbal tasks, our results suggest that this need not necessarily be the case. Instead, our data raise the surprising possibility that an early developing form of the cognitive ability to mentalize, evident in spontaneous looking behaviour, is not a necessary precursor of the

later developing form of mental state attribution, which supports explicit reasoning. The former would require spontaneous encoding of socially relevant information and automatic on-line computation of others' mental states, whereas the latter could also be achieved by verbally mediated reasoning prompted by explicit task structure and instructions.

More boldly, we suggest that compensatory learning can circumvent neurophysiological limitations, even without removing the original cause of the limitation. Such compensatory learning might explain the apparent paradox between success on explicit false belief tasks and continued difficulty in everyday social interaction in individuals with Asperger syndrome.

## REFERENCES AND NOTES

1. U. Frith, *Autism: Explaining the Enigma*. (Blackwell, Oxford, ed. 2nd, 2003).
2. S. Baron-Cohen, *Mindblindness : An essay on autism and theory of mind*. . (MIT Press., Cambridge, MA, 1995).
3. U. Frith, *Neuron* **32**, 969 (2001).
4. S. Baron-Cohen, A. M. Leslie, U. Frith, *Cognition* **21**, 37 (1985).
5. D. Dennett, *Behav. Brain Sci.* **4**, 568 (1978).
6. D. M. Bowler, *J. Child Psychol. Psychiatry* **33**, 877 (1992).
7. F. G. Happé, *Child Dev.* **66**, 843 (1995).
8. C. C. Peterson, V. P. Slaughter, J. Paynter, *J. Child Psychol. Psychiatry* **48**, 1243 (2007).
9. S. G. Shamay-Tsoory, *J. Autism Dev. Disord.* **38**, 1451 (2008).
10. A. Klin, W. Jones, R. Schultz, F. Volkmar, D. Cohen, *Am. J. Psychiatry* **159**, 895 (2002).
11. U. Frith, *J. Child Psychol. Psychiatry* **45**, 672 (2004).
12. F. Abell, F. Happé, U. Frith, *Cogn. Dev.* **15**, 1 (2000).
13. A. Klin, *J. Child Psychol. Psychiatry* **41**, 831 (2000).
14. F. Castelli, C. Frith, F. Happé, U. Frith, *Brain* **125**, 1839 (2002).
15. R. K. Kana, T. A. Keller, V. L. Cherkassky, N. J. Minshew, M. A. Just, *Soc. Neurosci.* **4**, 135 (2009).
16. T. Ruffman, W. Garnham, P. Rideout, *J. Child Psychol. Psychiatry* **42**, 1083 (2001).
17. K. H. Onishi, R. Baillargeon, *Science* **308**, 255 (2005).
18. V. Southgate, A. Senju, G. Csibra, *Psychol. Sci.* **18**, 587 (2007).
19. Materials and methods are available as supporting material on *Science Online*.
20. A. Senju, G. Csibra, *Curr. Biol.* **18**, 668 (2008).
21. V. Corkum, C. Moore, *Dev. Psychol.* **34**, 28 (1998).
22. H. M. Wellman, S. Lopez-Duran, J. LaBounty, B. Hamilton, *Dev. Psychol.* **44**, 618 (2008).
23. J. Perner, S. R. Leekam, H. Wimmer, *Br. J. Dev. Psychol.* **5**, 125 (1987).
24. T. Lockett, S. D. Powell, D. J. Messer, M. E. Thornton, J. Schulz, *J. Autism Dev. Disord.* **32**, 127 (2002).
25. H. M. Wellman, D. Liu, *Child Dev.* **75**, 523 (2004).
26. J. Perner, H. Wimmer, *J. Exp. Child Psychol.* **39**, 437 (1985).
27. P. C. Fletcher *et al.*, *Cognition* **57**, 109 (1995).
28. S. Baron-Cohen, S. Wheelwright, R. Skinner, J. Martin, E. Clubley, *J. Autism Dev. Disord.* **31**, 5 (2001).
29. We would like to thank Devorah Coniston for her invaluable help in data collection,

and Gergely Csibra and Chris Frith for their helpful discussions. AS was supported by an ESRC Research Fellowship (RES-063-27-0207); SW by an MRC/ESRC fellowship (PTA 037-27-0107) and UF by a MRC grant (G0701484) and the Aarhus Research Foundation.

**Supporting Online Material**

[www.sciencemag.org](http://www.sciencemag.org)

Materials and Methods

Supporting Text

Movies S1, S2