



Published in final edited form as:

Psychol Addict Behav. 2014 June ; 28(2): 580–585. doi:10.1037/a0036512.

Mindfulness Predicts Lower Affective Volatility among African Americans During Smoking Cessation

Claire E. Adams, Ph.D.¹, Minxing Chen, M.S.², Lin Guo³, Cho Y. Lam, Ph.D.³, Diana W. Stewart, Ph.D.³, Virmarie Correa-Fernández, Ph.D.³, Miguel A. Cano, Ph.D.³, Whitney L. Heppner, Ph.D.⁴, Jennifer Irvin Vidrine, Ph.D.³, Yisheng Li, Ph.D.², Jasjit S. Ahluwalia, M.D., M.P.H.⁵, Paul M. Cinciripini, Ph.D.⁶, and David W. Wetter, Ph.D.³

¹Department of Psychology, The Catholic University of America, Washington, DC

²Department of Biostatistics, UT MD Anderson Cancer Center, Houston, TX

³Department of Health Disparities Research, UT MD Anderson Cancer Center, Houston, TX

⁴Department of Psychological Science, Georgia College & State University, Milledgeville, GA

⁵The Center for Health Equity, University of Minnesota Medical School, Minneapolis, MN

⁶Department of Behavioral Science, UT MD Anderson Cancer Center, Houston, TX

Abstract

Recent research suggests that mindfulness benefits emotion regulation and smoking cessation. However, the mechanisms by which mindfulness affects emotional and behavioral functioning are unclear. One potential mechanism, lower affective volatility, has not been empirically tested during smoking cessation. This study examined longitudinal associations among mindfulness and emotional responding over the course of smoking cessation treatment among predominantly low-socioeconomic status (SES) African American smokers, who are at high risk for relapse to smoking and tobacco-related health disparities. Participants ($N = 399$, 51% female, mean age=42, 48% with annual income <\$10,000) completed a baseline measure of trait mindfulness. Negative affect, positive affect, and depressive symptoms were assessed at 5 time points during smoking cessation treatment (up to 31 days post-quit). Volatility indices were calculated to quantify within-person instability of emotional symptoms over time. Over and above demographic characteristics, nicotine dependence, and abstinence status, greater baseline trait mindfulness predicted lower volatility of negative affect and depressive symptoms surrounding the quit attempt and up to one month post-quit, $ps < 0.05$. Although volatility did not mediate the association between greater mindfulness and smoking cessation, these results are the first to show that mindfulness is linked to lower affective volatility (or greater stability) of negative emotions during the course of smoking cessation. The present study suggests that mindfulness is linked to greater emotional stability and augments the study of mindfulness in diverse populations. Future studies should examine the effects of mindfulness-based interventions on volatility and whether lower volatility explains effects of mindfulness-based treatments on smoking cessation.

Keywords

Mindfulness; Volatility; Smoking; Emotion Regulation; African Americans

Although the majority of current smokers in the U.S. indicate a desire to quit smoking, actual quit rates are low (CDC, 2011). Negative affect is a core symptom of nicotine withdrawal (Hendricks, Ditre, Drobes, & Brandon, 2006) and a consistent predictor of relapse to tobacco use (Baker, Piper, McCarthy, Majeskie, & Fiore, 2004). Studies examining emotional patterns during quit attempts indicate that in addition to the severity of affective symptoms, greater volatility (i.e., lability/scatter over time) predicts lapse and relapse (Cofta-Woerpel et al., 2011; Piasecki, Jorenby, Smith, Fiore, & Baker, 2003a, 2003b; Piasecki et al., 2000). Identifying factors that reduce affective volatility could be useful in smoking cessation treatment, particularly for populations with higher rates of relapse such as those with low socioeconomic status (SES) and African Americans (AAs; CDC, 2011; Fagan, Moolchan, Lawrence, Fernander, & Ponder, 2007; Shavers, Fagan, & McDonald, 2007). Mindfulness is fundamentally linked to affective experience and shows promise for regulating emotion (Szanton, Wenzel, Connolly, & Piferi, 2011) and enhancing smoking cessation (Heppner et al., under review) in low-SES AAs.

Mindfulness has been defined as purposeful, present-focused attention with an accepting, non-judgmental attitude (Baer, Smith, Hopkins, Krietemeyer, & Toney, 2006; Kabat-Zinn, 1990, 1994). Mindfulness involves observing thoughts and emotions as mental events that are not necessarily “true” or reflective of reality. Thus, mindfulness may help people to experience thoughts and feelings without getting “stuck” in their content or reacting to them in impulsive ways. This way of paying attention to thoughts and emotions without overly identifying with them is hypothesized to foster more flexible, adaptive responses (rather than impulsive reactions) to stressors (Arch & Craske, 2006, 2010). Mindfulness is linked to improved mood, anxiety, and stress (Baer et al., 2006; Brown & Ryan, 2003; Smith et al., 2011). Mindfulness may also promote more successful smoking cessation. In a pilot study ($N = 18$), Davis and colleagues (2007) found that 8 weeks of Mindfulness-based Stress Reduction (which included mindfulness-based instructions for coping with cravings) was associated with greater abstinence rates at 6-week follow-up than comparable smoking cessation studies. In a randomized controlled trial ($N = 88$), Brewer et al. (2011) reported that 8 sessions of mindfulness-based smoking cessation treatment produced better abstinence rates at 17-week follow-up than a standard smoking cessation treatment. Furthermore, Heppner et al. (under review) found that among 399 AA smokers, those with higher levels of mindfulness were both more likely to successfully quit and to recover abstinence if they experienced an early lapse.

Although research suggests that mindfulness predicts more positive and less negative affect, a more in-depth study of mechanisms by which mindfulness influences affective experience is needed. Indeed, there appear to be relatively stable inter-individual differences in intra-individual affective variability (Chow, Ram, Boker, Fujita, & Clore, 2005; Eaton & Funder, 2001; Larsen, 1987). Greater affective instability has been linked to poorer psychological health (Peeters, Berkhof, Delespaul, Rottenberg, & Nicolson, 2006; Trull et al., 2008).

Mindfulness is thought to promote a “decentered” perspective in which thoughts and emotions are observed as temporary mental events that do not necessarily represent reality (Teasdale et al., 2002). This mode of relating to experiences (or “metacognitive awareness;” Teasdale et al., 2002) might reduce the tendency for automatic reactions. For example,

Teasdale and colleagues' model of mindfulness-based relapse prevention for depression (2002, 1995) posits that nonjudgmental attention to mild depressive symptoms prevents cognitive and emotional reactivity to these experiences, thus preventing further cycles of more extreme symptoms. Furthermore, research suggests that mindfulness (both trait mindfulness and mindfulness-based training) reduces emotional reactivity to experiences such as distressing images, pain, and social stressors (Arch & Craske, 2006, 2010; Britton, Shahar, Szepsenwol, & Jacobs, 2012; Brown, Goodman, & Inzlicht, 2012). By promoting a decentered approach to experience, mindfulness may attenuate reactivity to day-to-day emotional experiences, thereby reducing affective instability over time.

A "volatility index" (indicating intra-individual variability in emotions over time) provides a quantitative method to study affective instability. In the only known study of mindfulness and affective volatility, Hill and Updegraff (2012) examined patterns of emotions among college students who indicated their emotional experiences six times per day for one week. Mean within-person standard deviations of positive and negative emotions were calculated to indicate degree of emotional instability. Results indicated that greater baseline mindfulness predicted lower volatility with regard to both positive and negative emotion. To the best of our knowledge, associations between mindfulness and affective volatility have not been evaluated during smoking cessation.

Affective volatility may be a critical factor that interferes with smoking cessation in low-SES AAs. Compared to higher-SES and other racial/ethnic groups, individuals with low SES and AAs might be particularly likely to smoke in an attempt to alleviate negative emotions, which are consistent predictors of nicotine dependence and difficulty quitting in this population (Bennett, Wolin, Robinson, Fowler, & Edwards, 2005; Landrine & Klonoff, 2000; Ludman et al., 2002). Although the few studies of mindfulness in low-SES AAs suggest it to be beneficial to emotion regulation (Szanton et al., 2011) and smoking cessation (Heppner et al., under review), more research is needed, particularly regarding mechanisms underlying these effects.

The current study tested the hypothesis that mindfulness predicts lower affective volatility in predominantly low-SES AAs during smoking cessation. Secondary analyses were conducted using data from a larger smoking cessation trial (Cano et al., in preparation). We were specifically interested in volatility of negative affect, which is problematic during cessation (Piasecki et al., 2000). However, we also examined volatility of positive affect, consistent with a previous finding of an association between greater mindfulness and lower volatility of positive emotion in college students (Hill & Updegraff, 2012). Finally, given that trait mindfulness predicts enhanced cessation outcomes in the present sample of low-SES AAs (Heppner et al., under review), we examined whether reduced affective volatility mediates this association.

Method

Participants

Data were collected as part of a randomized clinical trial examining a culturally tailored, palmtop computer-delivered smoking cessation treatment for AA smokers (Kendzor et al.,

2008). Participants were eligible if they self-identified as AA, were between 21–65 years old, had been smoking 5 cigarettes per day for 12 months, had an expired carbon monoxide level of 8 parts per million, planned to quit smoking within 2 weeks, possessed a functioning home telephone number and permanent home address, and were able to understand English at a sixth grade level. Exclusion criteria were regular use of tobacco products other than cigarettes, use of pharmacological cessation treatments other than nicotine patches supplied by the study, medical contraindication of the nicotine patch, or current pregnancy/lactation. Procedures were approved by the Institutional Review Board, and informed consent was obtained from all participants.

Procedure

Participants attended 7 study visits between 2005 and 2007: Pre-Quit Day –19 (baseline), Day –12, Day –5, Post-Quit Day +3, Day +10, Day +31, and Week +26. Participants were provided \$20 gift cards at each visit through Day +10, and \$40 gift cards at Day +31 and Week +26. At baseline, participants were randomly assigned to a standard smoking cessation treatment (ST) that included the nicotine patch, culturally sensitive self-help materials, and individual counseling, or ST in combination with palmtop computer-delivered treatment (CDT). Both treatments were provided through the study. Although analyses revealed no effect of treatment on abstinence (Cano et al., in preparation), treatment group was included as a covariate. This study utilizes data up to Day +31 to capture affective responding during the process of quitting.

Materials

Demographic and smoking characteristics—Demographics (collected at baseline) were age, gender, years of education, total annual family income, and partner status. Two items assessed pre-quit nicotine dependence: “How many cigarettes a day do you smoke on average?” and “How soon after you wake up do you smoke your first cigarette?” (“time to first cigarette;” Heatherton, Kozlowski, Frecker, Rickert, & Robinson, 1989). These items are strong indicators of nicotine dependence (Heatherton et al., 1989) and predictors of relapse (Baker et al., 2007).

Trait mindfulness—The Mindful Attention Awareness Scale (MAAS; Brown & Ryan, 2003) was administered at baseline. Participants responded on a 6-point Likert scale (1=*Almost Always*, 6=*Almost Never*) to 15 statements (e.g., “I could be experiencing some emotion and not be conscious of it until some time later,” “It seems I am ‘running on automatic,’ without much awareness of what I’m doing”). The MAAS showed excellent internal consistency ($\alpha = 0.92$).

Positive and negative affect—The Positive and Negative Affect Schedule (PANAS; Watson, Clark, & Tellegen, 1988) was administered at Day –12, Day –5, Day +3, Day +10, and Day +31. Participants rated the extent to which they experienced each of 20 emotions (e.g., distressed, ashamed, enthusiastic, excited) in the past week (1 =*Very Slightly or Not at All*, 5 =*Extremely*). The PANAS yields two factors: Positive Affect (PA) and Negative Affect (NA; Watson et al., 1988). Both subscales showed excellent internal consistency (α : 0.92 – 0.93).

Depressive symptoms—The Center for Epidemiological Studies–Depression (CES-D; Radloff, 1977) is a 20-item scale of depressive symptoms. Participants rated how often they experienced each symptom during the past week from “rarely or none of the time” to “most or all of the time.” Scores range from 0 to 60. The CES-D was administered at Day –12, Day –5, Day +3, Day +10, and Day +31, with acceptable internal consistency (α : 0.86 – 0.88).

Smoking abstinence—Abstinence (using an intent-to-treat approach) was assessed at Day +3 (abstinence since quit date), Day +10, and Day +31 (both as 7-day point prevalence abstinence). Abstinence was biochemically verified through expired carbon monoxide levels of <10 ppm (Hajek et al., 2001) and/or a cotinine value of <20 ng/ml (McBride et al., 1999).

Statistical Analyses

Several methods have been proposed to calculate volatility (Cofta-Woerpel et al., 2011; Jahng, Wood, & Trull, 2008; Piasecki et al., 2003a). Jahng et al. (2008) suggested that an optimal volatility index include information about general within-person variability as well as temporal instability (taking into account sequencing of scores over time). The exclusive use of within-person variance (or within-person standard deviation) accounts for general variability but not temporal instability. For this study, the mean square successive difference (MSSD), or average of the squared difference between successive observations at times $i + 1$ and i (Jahng et al., 2008), was chosen to capture both affective variability and temporal instability. Two volatility indices were created for each affective variable (PANAS NA, PANAS PA, CES-D) in order to examine emotional processes in the first month of quitting and also isolate the time period immediately surrounding the quit day (given that the vast majority of smokers lapse early in the quit attempt and greater volatility within the first week is linked to increased likelihood of an early lapse; Cofta-Woerpel et al., 2011). The first (indicating volatility surrounding the quit attempt) included Day –12, Day –5, Day +3, and Day +10. The second (indicating volatility surrounding the quit attempt and up to one month post-quit) also included Day +31. Given the unequal time intervals in the latter index, an adjustment in the calculation of the successive difference ($\lambda = 0.25$) was made for all indices including Day +31 (Jahng et al., 2008).

Linear regression models were fit to predict volatility indices from trait mindfulness using Stata/SE 12.1. Analyses controlled for baseline demographic characteristics (age, gender, education, income, partner status; chosen on the basis of past research; e.g., Businelle et al., 2010), dependence, and treatment type (ST vs. CDT). Additional analyses were conducted adjusting for abstinence status at Day +3, Day +10, and Day +31 in order to control for changes in negative affect associated with nicotine withdrawal (Hendricks et al., 2006). To test the hypothesis that lower affective volatility mediates the association between mindfulness and smoking cessation, the PROCESS macro (Hayes, 2013) was used with SPSS 21.0. Models tested the indirect effects of mindfulness on Day +31 abstinence through volatility indices (without Day +31; PANAS NA, PA, and CES-D tested in separate models). For each indirect effect, a 95% percentile bootstrap confidence interval was computed based on 1,000 bootstrap samples.

Results

Three hundred and ninety-nine AA smokers participated in the study. Approximately half (50.9%) were female. Average age was 42.44 years ($SD = 9.74$), 21.7% were married or living with a partner, 51.6% had less than or equal to a high school education, and 48.3% reported total family annual income of $< \$10,000$. Participants smoked 20.56 ($SD = 12.16$) cigarettes per day on average, and 58.6% reported smoking within 5 minutes of waking. Because of attrition, 308 participants had complete data up to Day +10, and 284 had complete data up to Day +31. Participants with complete data were older ($p = 0.01$) and had slightly higher MAAS scores ($p = 0.03$) than those with incomplete data; no other baseline differences emerged.

Greater baseline mindfulness was associated with lower volatility of negative affect and depressive symptoms, both surrounding the quit attempt and up to one-month post-quit ($ps < .05$). These associations remained significant after controlling for covariates, $ps < 0.05$ (Table 1). Baseline mindfulness was not associated with volatility of positive affect ($ps > 0.12$); this pattern remained once covariates were controlled ($ps > 0.24$; Table 1). Tests of indirect effects revealed that the volatility indices were not related to abstinence and were not significant mediators of associations between mindfulness and abstinence.

Discussion

Among predominantly low-SES AA smokers, those with greater baseline mindfulness exhibited lower volatility of negative emotions during the course of a smoking quit attempt. Although greater baseline mindfulness predicted enhanced smoking cessation in this sample (see Heppner et al., under review), affective volatility did not mediate this association. Findings support the hypothesis that greater mindfulness is linked to lower volatility of negative emotions when faced with major life stressors such as quitting smoking. This is the first known study to show a connection between greater mindfulness and lower volatility of negative emotion during the course of smoking cessation. Volatility of negative emotion is a critical determinant of relapse to smoking (Cofta-Woerpel et al., 2011) and is linked to poor psychological functioning more generally (Peeters et al., 2006; Trull et al., 2008). Regardless of the lack of support for volatility as a mediator in this particular study, findings suggest that mindfulness is linked to greater emotional stability, an important aspect of emotional functioning during the course of smoking cessation.

Research consistently shows mindfulness to be associated with greater positive and lower negative emotion (Baer et al., 2006; Brown & Ryan, 2003; Smith et al., 2011). Perhaps the benefits of mindfulness are not only due to more positive and less negative emotion, but also to less volatility of negative emotion. The tendency to observe experiences without impulsively reacting may lessen the likelihood of extreme shifts in negative emotion, thereby fostering a sense of equanimity. Researchers are striving to understand the construct of “mindful emotion regulation” (Chambers, Gullone, & Allen, 2009), and lower volatility of negative emotion might be one critical mechanism to consider.

In comparison to the only other known study of mindfulness and emotional volatility, Hill and Updegraff (2012) found that greater baseline mindfulness (particularly “nonreactivity”) predicted lower volatility of both negative and positive emotion. The current study found a relationship between mindfulness and volatility of negative, but not positive affect. This discrepancy could be related to differences in the study samples and methodologies. Hill and Updegraff utilized a sample of mostly non-Latino white college students, collected data over a typical week of college, utilized a different measure of mindfulness (Five Facet Mindfulness Questionnaire; Baer et al., 2006), and calculated volatility using within-person standard deviations (capturing general variability but not temporal instability; Jahng et al., 2008). The current study examined volatility using the MSSD approach during smoking cessation among predominantly low-SES AA smokers. Perhaps mindfulness is more relevant to processes regarding negative affect (a core symptom of nicotine withdrawal) than positive affect during smoking cessation.

Moreover, optimal emotion regulation might be better reflected by low reactivity to negative events and higher reactivity to positive events. For example, mindful attention might lessen reactivity in the context of stressful events (such as smoking cessation) but increase the emotional benefits of pleasant experiences. Mindful attention to pleasant experiences is akin to the process of “savoring,” through which individuals purposefully pay attention to positive life events to experience greater emotional benefit (Bryant & Veroff, 2007). Indeed, mindfulness is associated with greater reactivity to pleasant experiences (as evidenced by “boosts” in positive emotions; Catalino & Fredrickson, 2011). Further research is needed to elucidate associations between mindfulness and volatility with regard to positive emotions.

The current study is limited by self-reported questionnaire data, which may suffer from retrospective recall bias. There is a need for mindfulness research to examine moment-to-moment experiences in natural environments (e.g., using ecological momentary assessment [EMA]; Shiffman, Paty, Gny, Kassel, & Hickcox, 1996). Only one known study has used EMA to study associations between mindfulness and volatility (Hill & Updegraff, 2012), and future research should use EMA to examine mechanisms of mindfulness in more diverse populations. We encourage researchers to revisit the question of whether volatility mediates the association between mindfulness and smoking using EMA data, which would provide a more fine-grained analysis than possible with the current study. Researchers should also examine other potential mechanisms (e.g., levels of negative and positive affect, self-efficacy, perceived social support) that might explain associations between mindfulness and abstinence on a moment-to-moment basis.

This study is also limited by a unidimensional measure of mindfulness. Although the MAAS appears reliable and valid (Brown & Ryan, 2003), recent research has emphasized the multidimensionality of mindfulness (Baer et al., 2006). Furthermore, given that our study examined correlational associations between mindfulness and affective volatility (which does not necessarily imply causality), research that examines the direct effects of mindfulness training on affective volatility is needed. Finally, given the specific nature of our sample, we do not know whether results would generalize to other populations. Although it is entirely possible that these findings might be relevant in other contexts, the

current data cannot speak to affective processes in non-smokers or in life circumstances other than quitting smoking.

This study is strengthened by its assessment of multiple affective variables, use of longitudinal data, control for socio-demographics, dependence, and abstinence status, use of a volatility indicator of both within-person variability and temporal instability (Jahng et al., 2008), and utilization of a sample of predominantly low-SES AAs (an underserved population at high risk for affectively-triggered smoking and tobacco-related disparities). Results revealed that greater mindfulness is associated with lower volatility of negative affect and depressive symptoms over the course of smoking cessation. Results provide information on how mindfulness might benefit emotion regulation (empirically elucidating mechanisms of mindfulness) and suggest benefits of mindfulness for low-SES AAs. Lower volatility of negative emotion may be one mechanism by which mindfulness enhances emotion regulation in underserved populations.

Acknowledgments

Funding Statement: This research was supported by the National Cancer Institute through grants R01CA94826, R25-TCA57730, the University of Texas MD Anderson Cancer Center's Support Grant CA016672, and the Latinos Contra el Cancer Community Networks Program Center Grant U54CA153505. This work was also supported by faculty fellowships from the University of Texas MD Anderson Cancer Center Duncan Family Institute for Cancer Prevention and Risk Assessment.

References

- Arch JJ, Craske MG. Mechanisms of mindfulness: emotion regulation following a focused breathing induction. *Behaviour Research and Therapy*. 2006; 44(12):1849–1858. S0005-7967(05)00274-3 [pii]. 10.1016/j.brat.2005.12.007 [PubMed: 16460668]
- Arch JJ, Craske MG. Laboratory stressors in clinically anxious and non-anxious individuals: the moderating role of mindfulness. *Behaviour Research and Therapy*. 2010; 48(6):495–505. S0005-7967(10)00023-9 [pii]. 10.1016/j.brat.2010.02.005 [PubMed: 20303471]
- Baer RA, Smith GT, Hopkins J, Krietemeyer J, Toney L. Using self-report assessment methods to explore facets of mindfulness. *Assessment*. 2006; 13(1):27–45. 13/1/27 [pii]. 10.1177/1073191105283504 [PubMed: 16443717]
- Baker TB, Piper ME, McCarthy DE, Bolt DM, Smith SS, Kim SY, Toll BA. Time to first cigarette in the morning as an index of ability to quit smoking: implications for nicotine dependence. *Nicotine & Tobacco Research*. 2007; 9(Suppl 4):S555–570. 788258006 [pii]. 10.1080/14622200701673480 [PubMed: 18067032]
- Baker TB, Piper ME, McCarthy DE, Majeskie MR, Fiore MC. Addiction motivation reformulated: an affective processing model of negative reinforcement. *Psychological Review*. 2004; 111(1):33–51. [pii]. 10.1037/0033-295X.111.1.332004-10332-002 [PubMed: 14756584]
- Bennett GG, Wolin KY, Robinson EL, Fowler S, Edwards CL. Perceived racial/ethnic harassment and tobacco use among African American young adults. *American Journal of Public Health*. 2005; 95(2):238–240. 95/2/238 [pii]. 10.2105/AJPH.2004.037812 [PubMed: 15671457]
- Brewer JA, Mallik S, Babuscio TA, Nich C, Johnson HE, Deleone CM, Rounsaville BJ. Mindfulness training for smoking cessation: results from a randomized controlled trial. *Drug and Alcohol Dependence*. 2011; 119(1–2):72–80. S0376-8716(11)00253-5 [pii]. 10.1016/j.drugalcdep.2011.05.027 [PubMed: 21723049]
- Britton WB, Shahar B, Szepsenwol O, Jacobs WJ. Mindfulness-based cognitive therapy improves emotional reactivity to social stress: results from a randomized controlled trial. *Behavior Therapy*. 2012; 43(2):365–380. S0005-7894(11)00131-6 [pii]. 10.1016/j.beth.2011.08.006 [PubMed: 22440072]

- Brown KW, Goodman RJ, Inzlicht M. Dispositional mindfulness and the attenuation of neural responses to emotional stimuli. *Social Cognitive and Affective Neuroscience*. 2012 nss004 [pii]. 10.1093/scan/nss004
- Brown KW, Ryan RM. The benefits of being present: mindfulness and its role in psychological well-being. *Journal of Personality and Social Psychology*. 2003; 84(4):822–848. [PubMed: 12703651]
- Bryant, FB.; Veroff, J. *Savoring: A new model of positive experience*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers; 2007.
- Businelle MS, Kendzor DE, Reitzel LR, Costello TJ, Cofta-Woerpel L, Li Y, Wetter DW. Mechanisms linking socioeconomic status to smoking cessation: a structural equation modeling approach. *Health Psychol*. 2010; 29(3):262–273. 2010-09923-005 [pii]. 10.1037/a0019285 [PubMed: 20496980]
- Cano MA, Mazas CA, Cofta-Woerpel L, Costello TJ, Li Y, Vidrine JJ, Wetter DW. Randomized Clinical Trial of a Palmtop Computer-Delivered Treatment for Smoking Cessation Among African American Smokers. in preparation.
- Catalino LI, Fredrickson BL. A Tuesday in the life of a flourisher: the role of positive emotional reactivity in optimal mental health. *Emotion*. 2011; 11(4):938–950.10.1037/a0024889 [PubMed: 21859208]
- CDC. Quitting smoking among adults--United States, 2001–2010. *Morbidity and Mortality Weekly Reports*. 2011; 60(44):1513–1519. mm6044a2 [pii].
- Chambers R, Gullone E, Allen NB. Mindful emotion regulation: An integrative review. *Clinical Psychology Review*. 2009; 29(6):560–572. S0272-7358(09)00086-5 [pii]. 10.1016/j.cpr.2009.06.005 [PubMed: 19632752]
- Chow SM, Ram N, Boker SM, Fujita F, Clore G. Emotion as a thermostat: representing emotion regulation using a damped oscillator model. *Emotion*. 2005; 5(2):208–225.10.1037/1528-3542.5.2.208 [PubMed: 15982086]
- Cofta-Woerpel L, McClure JB, Li Y, Urbauer D, Cinciripini PM, Wetter DW. Early cessation success or failure among women attempting to quit smoking: trajectories and volatility of urge and negative mood during the first postcessation week. *Journal of Abnormal Psychology*. 2011; 120(3):596–606. 2011-10194-001 [pii]. 10.1037/a0023755 [PubMed: 21574667]
- Davis JM, Fleming MF, Bonus KA, Baker TB. A pilot study on mindfulness based stress reduction for smokers. *BMC Complementary and Alternative Medicine*. 2007; 7:2. 1472-6882-7-2 [pii]. 10.1186/1472-6882-7-2 [PubMed: 17254362]
- Eaton LG, Funder DC. Emotional experience in daily life: valence, variability, and rate of change. *Emotion*. 2001; 1(4):413–421. [PubMed: 12901400]
- Fagan P, Moolchan ET, Lawrence D, Fernander A, Ponder PK. Identifying health disparities across the tobacco continuum. *Addiction*. 2007; 102(Suppl 2):5–29. ADD1952 [pii]. 10.1111/j.1360-0443.2007.01952.x [PubMed: 17850611]
- Hajek P, West R, Lee A, Foulds J, Owen L, Eiser JR, Main N. Randomized controlled trial of a midwife-delivered brief smoking cessation intervention in pregnancy. *Addiction*. 2001; 96(3):485–494.10.1080/0965214002005446 [PubMed: 11255587]
- Hayes, AF. *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. New York, NY: Guilford; 2013.
- Heatherton TF, Kozlowski LT, Frecker RC, Rickert W, Robinson J. Measuring the heaviness of smoking: using self-reported time to the first cigarette of the day and number of cigarettes smoked per day. *British Journal of Addiction*. 1989; 84(7):791–799. [PubMed: 2758152]
- Hendricks PS, Ditre JW, Drobes DJ, Brandon TH. The early time course of smoking withdrawal effects. *Psychopharmacology*. 2006; 187(3):385–396.10.1007/s00213-006-0429-9 [PubMed: 16752139]
- Heppner WL, Adams CE, Correa-Fernandez V, Castro Y, Li Y, Reitzel LR, Wetter DW. Dispositional mindfulness predicts enhanced smoking cessation and smoking lapse recovery. under review.
- Hill CL, Updegraff JA. Mindfulness and its relationship to emotional regulation. *Emotion*. 2012; 12(1):81–90. 2011-28774-001 [pii]. 10.1037/a0026355 [PubMed: 22148996]

- Jahng S, Wood PK, Trull TJ. Analysis of affective instability in ecological momentary assessment: Indices using successive difference and group comparison via multilevel modeling. *Psychological Methods*. 2008; 13(4):354–375. 2008-17368-004 [pii]. 10.1037/a0014173 [PubMed: 19071999]
- Kabat-Zinn, J. *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. New York: Delacourt; 1990.
- Kabat-Zinn, J. *Wherever You Go, There You Are: Mindfulness and Meditation in Everyday Life*. New York: Hyperion; 1994.
- Kendzor DE, Cofta-Woerpel LM, Mazas CA, Li Y, Vidrine JI, Reitzel LR, Wetter DW. Socioeconomic status, negative affect, and modifiable cancer risk factors in African-American smokers. *Cancer Epidemiology Biomarkers and Prevention*. 2008; 17(10):2546–2554. 17/10/2546 [pii]. 10.1158/1055-9965.EPI-08-0291
- Landrine H, Klonoff EA. Racial discrimination and cigarette smoking among Blacks: findings from two studies. *Ethnicity and Disease*. 2000; 10(2):195–202. [PubMed: 10892825]
- Larsen RJ. The Stability of Mood Variability - a Spectral Analytic Approach to Daily Mood Assessments. *Journal of Personality and Social Psychology*. 1987; 52(6):1195–1204.10.1037//0022-3514.52.6.1195
- Ludman EJ, Curry SJ, Grothaus LC, Graham E, Stout J, Lozano P. Depressive symptoms, stress, and weight concerns among African American and European American low-income female smokers. *Psychology of Addictive Behaviors*. 2002; 16(1):68–71. [PubMed: 11934089]
- McBride CM, Curry SJ, Lando HA, Pirie PL, Grothaus LC, Nelson JC. Prevention of relapse in women who quit smoking during pregnancy. *Am J Public Health*. 1999; 89(5):706–711. [PubMed: 10224982]
- Peeters F, Berkhof J, Delespaul P, Rottenberg J, Nicolson NA. Diurnal mood variation in major depressive disorder. *Emotion*. 2006; 6(3):383–391.10.1037/1528-3542.6.3.383 [PubMed: 16938080]
- Piasecki TM, Jorenby DE, Smith SS, Fiore MC, Baker TB. Smoking withdrawal dynamics: I. Abstinence distress in lapsers and abstainers. *Journal of Abnormal Psychology*. 2003a; 112(1):3–13. [PubMed: 12653409]
- Piasecki TM, Jorenby DE, Smith SS, Fiore MC, Baker TB. Smoking withdrawal dynamics: II. Improved tests of withdrawal-relapse relations. *Journal of Abnormal Psychology*. 2003b; 112(1):14–27. [PubMed: 12653410]
- Piasecki TM, Niaura R, Shadel WG, Abrams D, Goldstein M, Fiore MC, Baker TB. Smoking withdrawal dynamics in unaided quitters. *Journal of Abnormal Psychology*. 2000; 109(1):74–86. [PubMed: 10740938]
- Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1:385–401.
- Shavers VL, Fagan P, McDonald P. Health disparities across the cancer continuum. *Journal of Health Care for the Poor and Underserved*. 2007; 18(4 Suppl):1–5. S1548686907400018 [pii]. 10.1353/hpu.2007.0122 [PubMed: 18065848]
- Shiffman S, Paty JA, Gnys M, Kassel JA, Hickcox M. First lapses to smoking: within-subjects analysis of real-time reports. *Journal of Consulting and Clinical Psychology*. 1996; 64(2):366–379. [PubMed: 8871421]
- Smith BW, Ortiz JA, Steffen LE, Tooley EM, Wiggins KT, Yeater EA, Bernard ML. Mindfulness is associated with fewer PTSD symptoms, depressive symptoms, physical symptoms, and alcohol problems in urban firefighters. *Journal of Consulting and Clinical Psychology*. 2011; 79(5):613–617. 2011-19048-001 [pii]. 10.1037/a0025189 [PubMed: 21875175]
- Szanton SL, Wenzel J, Connolly AB, Piferi RL. Examining mindfulness-based stress reduction: perceptions from minority older adults residing in a low-income housing facility. *BMC Complementary and Alternative Medicine*. 2011; 11:44. 1472-6882-11-44 [pii]. 10.1186/1472-6882-11-44 [PubMed: 21627807]
- Teasdale JD, Moore RG, Hayhurst H, Pope M, Williams S, Segal ZV. Metacognitive awareness and prevention of relapse in depression: empirical evidence. *Journal of Consulting and Clinical Psychology*. 2002; 70(2):275–287. [PubMed: 11952186]

- Teasdale JD, Segal Z, Williams JM. How does cognitive therapy prevent depressive relapse and why should attentional control (mindfulness) training help? *Behaviour Research and Therapy*. 1995; 33(1):25–39. 0005-7967(94)E0011-7 [pii]. [PubMed: 7872934]
- Trull TJ, Solhan MB, Tragesser SL, Jahng S, Wood PK, Piasecki TM, Watson D. Affective instability: measuring a core feature of borderline personality disorder with ecological momentary assessment. *J Abnorm Psychol*. 2008; 117(3):647–661.10.1037/a0012532 [PubMed: 18729616]
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology*. 1988; 54(6): 1063–1070. [PubMed: 3397865]

Table 1

Predicting Affective Volatility Indices from Trait Mindfulness

Mindfulness Predicting Volatility of Negative Affect (PANAS NA)				
Volatility Index	MAAS coefficient (b)	SE	95% CI	p
Surrounding quit attempt	-10.30	4.91	-19.96, -.64	.037
Surrounding quit attempt (controlling for abstinence)	-9.85	4.93	-19.56, -.14	.047
Up to 1-month post-quit	-12.58	4.48	-21.41, -3.75	.005
Up to 1-month post-quit (controlling for abstinence)	-13.01	4.48	-21.84, -4.18	.004

Mindfulness Predicting Volatility of Positive Affect (PANAS PA)				
Volatility Index	MAAS coefficient (b)	SE	95% CI	p
Surrounding quit attempt	-2.42	4.16	-10.61, 5.76	.560
Surrounding quit attempt (controlling for abstinence)	-2.62	4.14	-10.78, 5.53	.527
Up to 1-month post-quit	-3.43	3.94	-11.19, 4.32	.384
Up to 1-month post-quit (controlling for abstinence)	-4.51	3.88	-12.15, 3.12	.246

Mindfulness Predicting Volatility of Depressive Symptoms (CES-D)				
Volatility Index	MAAS coefficient (b)	SE	95% CI	p
Surrounding quit attempt	-18.12	6.64	-31.18, -5.05	.007
Surrounding quit attempt (controlling for abstinence)	-17.34	6.68	-30.48, -4.19	.010
Up to 1-month post-quit	-20.36	6.12	-32.42, -8.30	.001
Up to 1-month post-quit (controlling for abstinence)	-19.59	6.19	-31.77, -7.40	.002

Notes. Linear regression models were fit to predict volatility indices (volatility of negative affect, positive affect, and depressive symptoms) from trait mindfulness.

MAAS = Mindful Attention Awareness Scale

PANAS NA = Positive and Negative Affect Schedule – Negative Affect

PANAS PA = Positive and Negative Affect Schedule – Positive Affect

CES-D = Center for Epidemiological Studies-Depression

All analyses control for baseline demographic characteristics (age, gender, education, income, partner status), dependence (baseline cigarettes per day and time to first cigarette), and treatment type (ST vs. CDT). Analyses controlling for abstinence surrounding the quit attempt also control for smoking abstinence at Day +3 and Day +10. Analyses controlling for abstinence up to 1-month post-quit also control for smoking abstinence at Day +3, Day +10, and Day +31.

Surrounding quit attempt = Day -12, Day -5, Day +3, Day +10

Up to 1-month post-quit = Day -12, Day -5, Day +3, Day +10, Day +31