

# MINIMAL POSITION AND CRITICAL MARTINGALE CONVERGENCE IN BRANCHING RANDOM WALKS, AND DIRECTED POLYMERS ON DISORDERED TREES

BY YUEYUN HU AND ZHAN SHI

*Université Paris XIII and Université Paris VI*

We establish a second-order almost sure limit theorem for the minimal position in a one-dimensional super-critical branching random walk, and also prove a martingale convergence theorem which answers a question of Biggins and Kyprianou [*Electron. J. Probab.* **10** (2005) 609–631]. Our method applies, furthermore, to the study of directed polymers on a disordered tree. In particular, we give a rigorous proof of a phase transition phenomenon for the partition function (from the point of view of convergence in probability), already described by Derrida and Spohn [*J. Statist. Phys.* **51** (1988) 817–840]. Surprisingly, this phase transition phenomenon disappears in the sense of upper almost sure limits.

## 1. Introduction.

1.1. *Branching random walk and martingale convergence.* We consider a branching random walk on the real line  $\mathbb{R}$ . Initially, a particle sits at the origin. Its children form the first generation; their displacements from the origin correspond to a point process on the line. These children have children of their own (who form the second generation), and behave—relative to their respective positions—like independent copies of the initial particle. And so on.

We write  $|u| = n$  if an individual  $u$  is in the  $n$ th generation, and denote its position by  $V(u)$ . [In particular, for the initial ancestor  $e$ , we have  $V(e) = 0$ .] We assume throughout the paper that, for some  $\delta > 0$ ,  $\delta_+ > 0$  and  $\delta_- > 0$ ,

$$(1.1) \quad \mathbf{E} \left\{ \left( \sum_{|u|=1} 1 \right)^{1+\delta} \right\} < \infty,$$

$$(1.2) \quad \mathbf{E} \left\{ \sum_{|u|=1} e^{-(1+\delta_+)V(u)} \right\} + \mathbf{E} \left\{ \sum_{|u|=1} e^{\delta_- V(u)} \right\} < \infty,$$

here  $\mathbf{E}$  denotes expectation with respect to  $\mathbf{P}$ , the law of the branching random walk.

---

Received April 2007; revised February 2008.

*AMS 2000 subject classification.* 60J80.

*Key words and phrases.* Branching random walk, minimal position, martingale convergence, spine, marked tree, directed polymer on a tree.

Let us define the (logarithmic) moment generating function

$$\psi(t) := \log \mathbf{E} \left\{ \sum_{|u|=1} e^{-tV(u)} \right\} \in (-\infty, \infty], \quad t \geq 0.$$

By (1.2),  $\psi(t) < \infty$  for  $t \in [-\delta_-, 1 + \delta_+]$ . Following Biggins and Kyprianou [9], we assume

$$(1.3) \quad \psi(0) > 0, \quad \psi(1) = \psi'(1) = 0.$$

Since the number of particles in each generation forms a Galton–Watson tree, the assumption  $\psi(0) > 0$  in (1.3) says that this Galton–Watson tree is super-critical.

In the study of the branching random walk, there is a fundamental martingale, defined as follows:

$$(1.4) \quad W_n := \sum_{|u|=n} e^{-V(u)}, \quad n = 0, 1, 2, \dots \left( \sum_{\emptyset} := 0 \right).$$

Since  $W_n \geq 0$ , it converges almost surely.

When  $\psi'(1) < 0$ , it is proved by Biggins and Kyprianou [7] that there exists a sequence of constants  $(a_n)$  such that  $\frac{W_n}{a_n}$  converges in probability to a nondegenerate limit which is (strictly) positive upon the survival of the system. This is called the Seneta–Heyde norming in [7] for branching random walk, referring to Seneta [35] and Heyde [22] on the rate of convergence in the classic Kesten–Stigum theorem for Galton–Watson processes.

The case  $\psi'(1) = 0$  is more delicate. In this case, it is known (Lyons [29]) that  $W_n \rightarrow 0$  almost surely. The following question is raised in Biggins and Kyprianou [9]: are there deterministic normalizers  $(a_n)$  such that  $\frac{W_n}{a_n}$  converges?

We aim at answering this question.

**THEOREM 1.1.** *Assume (1.1), (1.2) and (1.3). There exists a deterministic positive sequence  $(\lambda_n)$  with  $0 < \liminf_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} \leq \limsup_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} < \infty$ , such that, conditionally on the system’s survival,  $\lambda_n W_n$  converges in distribution to  $\mathscr{W}$ , with  $\mathscr{W} > 0$  a.s. The distribution of  $\mathscr{W}$  is given in (10.3).*

The limit  $\mathscr{W}$  in Theorem 1.1 turns out to satisfy a functional equation. Such functional equations are known to be closely related to (a discrete version of) the Kolmogorov–Petrovski–Piscounov (KPP) traveling wave equation; see Kyprianou [25] for more details.

The almost sure behavior of  $W_n$  is described in Theorem 1.3 below. The two theorems together give a clear image of the asymptotics of  $W_n$ .

1.2. *The minimal position in the branching random walk.* A natural question in the study of branching random walks is about  $\inf_{|u|=n} V(u)$ , the position of the leftmost individual in the  $n$ th generation. In the literature the concentration (in terms of tightness or even weak convergence) of  $\inf_{|u|=n} V(u)$  around its median/quantiles has been studied by many authors. See, for example, Bachmann [4] and Bramson and Zeitouni [14], as well as Section 5 of the survey paper by Aldous and Bandyopadhyay [2]. We also mention the recent paper of Lifshits [26], where an example of a branching random walk is constructed such that  $\inf_{|u|=n} V(u) - \text{median}(\{\inf_{|u|=n} V(u)\})$  is tight but does not converge weakly.

We are interested in the asymptotic speed of  $\inf_{|u|=n} V(u)$ . Under assumption (1.3), it is known that, conditionally on the system’s survival,

$$(1.5) \quad \frac{1}{n} \inf_{|u|=n} V(u) \rightarrow 0 \quad \text{a.s.},$$

$$(1.6) \quad \inf_{|u|=n} V(u) \rightarrow +\infty \quad \text{a.s.}$$

The “law of large numbers” in (1.5) is a classic result, and can be found in Hammersley [19], Kingman [23] and Biggins [5]. The system’s transience to the right, stated in (1.6), follows from the fact that  $W_n \rightarrow 0$ , a.s.

A refinement of (1.5) is obtained by McDiarmid [31]. Under the additional assumption  $\mathbf{E}\{(\sum_{|u|=1} 1)^2\} < \infty$ , it is proved in [31] that, for some constant  $c_1 < \infty$  and conditionally on the system’s survival,

$$\limsup_{n \rightarrow \infty} \frac{1}{\log n} \inf_{|u|=n} V(u) \leq c_1 \quad \text{a.s.}$$

We intend to determine the exact rate at which  $\inf_{|u|=n} V(u)$  goes to infinity.

**THEOREM 1.2.** *Assume (1.1), (1.2) and (1.3). Conditionally on the system’s survival, we have*

$$(1.7) \quad \limsup_{n \rightarrow \infty} \frac{1}{\log n} \inf_{|u|=n} V(u) = \frac{3}{2} \quad \text{a.s.},$$

$$(1.8) \quad \liminf_{n \rightarrow \infty} \frac{1}{\log n} \inf_{|u|=n} V(u) = \frac{1}{2} \quad \text{a.s.},$$

$$(1.9) \quad \lim_{n \rightarrow \infty} \frac{1}{\log n} \inf_{|u|=n} V(u) = \frac{3}{2} \quad \text{in probability.}$$

**REMARK.** (i) The most interesting part of Theorem 1.2 is (1.7)–(1.8). It reveals the presence of fluctuations of  $\inf_{|u|=n} V(u)$  on the logarithmic level, which is in contrast with known results of Bramson [13] and Dekking and Host [16] stating that, for a class of branching random walks,  $\frac{1}{\log \log n} \inf_{|u|=n} V(u)$  converges almost surely to a finite and positive constant.

(ii) Some brief comments on (1.3) are in order. In general [i.e., without assuming  $\psi(1) = \psi'(1) = 0$ ], the law of large numbers (1.5) reads  $\frac{1}{n} \inf_{|u|=n} V(u) \rightarrow c$ , a.s. (conditionally on the system's survival), where  $c := \inf\{a \in \mathbb{R} : g(a) \geq 0\}$ , with  $g(a) := \inf_{t \geq 0} \{ta + \psi(t)\}$ . If

$$(1.10) \quad t^* \psi'(t^*) = \psi(t^*)$$

for some  $t^* \in (0, \infty)$ , then the branching random walk associated with the point process  $\widehat{V}(u) := t^* V(u) + \psi(t^*)|u|$  satisfies (1.3). That is, as long as (1.10) has a solution [which is the case, e.g., if  $\psi(1) = 0$  and  $\psi'(1) > 0$ ], the study will boil down to the case (1.3).

It is, however, possible that (1.10) has no solution. In such a situation, Theorem 1.2 does not apply. For example, we have already mentioned a class of branching random walks exhibited in Bramson [13] and Dekking and Host [16], for which  $\inf_{|u|=n} V(u)$  has an exotic  $\log \log n$  behavior.

(iii) Under suitable assumptions, Addario–Berry [1] obtains a very precise asymptotic estimate of  $\mathbf{E}[\inf_{|u|=n} V(u)]$ , which implies (1.9).

(iv) In the case of branching Brownian motion, the analogue of (1.9) was proved by Bramson [12], by means of some powerful explicit analysis.

1.3. *Directed polymers on a disordered tree.* The following model is borrowed from the well-known paper of Derrida and Spohn [17]: Let  $\mathbb{T}$  be a rooted Cayley tree; we study all self-avoiding walks (= directed polymers) of  $n$  steps on  $\mathbb{T}$  starting from the root. To each edge of the tree is attached a random variable (= potential). We assume that these random variables are independent and identically distributed. For each walk  $\omega$ , its energy  $E(\omega)$  is the sum of the potentials of the edges visited by the walk. So the partition function is

$$Z_n := \sum_{\omega} e^{-\beta E(\omega)},$$

where the sum is over all self-avoiding walks of  $n$  steps on  $\mathbb{T}$ , and  $\beta > 0$  is the inverse temperature.

More generally, we take  $\mathbb{T}$  to be a Galton–Watson tree, and observe that the energy  $E(\omega)$  corresponds to (the partial sum of) the branching random walk described in the previous sections. The associated partition function becomes

$$(1.11) \quad W_{n,\beta} := \sum_{|u|=n} e^{-\beta V(u)}, \quad \beta > 0.$$

Clearly, when  $\beta = 1$ ,  $W_{n,1}$  is just the  $W_n$  defined in (1.4).

If  $0 < \beta < 1$ , the study of  $W_{n,\beta}$  boils down to the case  $\psi'(1) < 0$ , which was investigated by Biggins and Kyprianou [7]. In particular, conditionally on the system's survival,  $\frac{W_{n,\beta}}{\mathbf{E}\{W_{n,\beta}\}}$  converges almost surely to a (strictly) positive random variable.

We study the case  $\beta \geq 1$  in the present paper.

THEOREM 1.3. Assume (1.1), (1.2) and (1.3). Conditionally on the system's survival, we have

$$(1.12) \quad W_n = n^{-1/2+o(1)} \quad a.s.$$

THEOREM 1.4. Assume (1.1), (1.2) and (1.3), and let  $\beta > 1$ . Conditionally on the system's survival, we have

$$(1.13) \quad \limsup_{n \rightarrow \infty} \frac{\log W_{n,\beta}}{\log n} = -\frac{\beta}{2} \quad a.s.,$$

$$(1.14) \quad \liminf_{n \rightarrow \infty} \frac{\log W_{n,\beta}}{\log n} = -\frac{3\beta}{2} \quad a.s.,$$

$$(1.15) \quad W_{n,\beta} = n^{-3\beta/2+o(1)} \quad \text{in probability.}$$

Again, the most interesting part in Theorem 1.4 is (1.13) and (1.14), which describes a new fluctuation phenomenon. Also, there is no phase transition any more for  $W_{n,\beta}$  at  $\beta = 1$  from the point of view of upper almost sure limits.

The remark on (1.3), stated after Theorem 1.2, applies to Theorems 1.3 and 1.4 as well.

An important step in the proof of Theorems 1.3 and 1.4 is to estimate all small moments of  $W_n$  and  $W_{n,\beta}$ , respectively. This is done in the next theorems.

THEOREM 1.5. Assume (1.1), (1.2) and (1.3). For any  $\gamma \in [0, 1)$ , we have

$$(1.16) \quad 0 < \liminf_{n \rightarrow \infty} \mathbf{E}\{(n^{1/2}W_n)^\gamma\} \leq \limsup_{n \rightarrow \infty} \mathbf{E}\{(n^{1/2}W_n)^\gamma\} < \infty.$$

THEOREM 1.6. Assume (1.1), (1.2) and (1.3), and let  $\beta > 1$ . For any  $0 < r < \frac{1}{\beta}$ , we have

$$(1.17) \quad \mathbf{E}\{W_{n,\beta}^r\} = n^{-3r\beta/2+o(1)}, \quad n \rightarrow \infty.$$

The rest of the paper is as follows. In Section 2 we introduce a change-of-measures formula (Proposition 2.1) in terms of spines on marked trees. This formula will be of frequent use throughout the paper. Section 3 contains a few preliminary results of the lower tail probability of the martingale  $W_n$ . The proofs of the theorems are organized as follows:

- Section 4: upper bound in part (1.8) of Theorem 1.2.
- Section 5: Theorem 1.6.
- Section 6: Theorem 1.5.
- Section 7: Theorem 1.3, as well as parts (1.14) and (1.15) of Theorem 1.4.
- Section 8: (the rest of) Theorem 1.2.
- Section 9: part (1.13) of Theorem 1.4.

- Section 10: Theorem 1.1.

Section 4 relies on ideas borrowed from Bramson [12], and does not require the preliminaries in Sections 2 and 3.

Sections 5 and 6 are the technical part of the paper, where a common idea is applied in two different situations.

Throughout the paper we write

$$q := \mathbf{P}\{\text{the system's extinction}\} \in [0, 1).$$

The letter  $c$  with a subscript denotes finite and (strictly) positive constants. We also use the notation  $\sum_{\emptyset} := 0$ ,  $\prod_{\emptyset} := 1$ , and  $0^0 := 1$ . Moreover, we use  $a_n \sim b_n$ ,  $n \rightarrow \infty$ , to denote  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$ .

**2. Marked trees and spines.** This section is devoted to a change-of-measures result (Proposition 2.1) on marked trees in terms of spines. The material of this section has been presented in the literature in various forms; see, for example, Chauvin, Rouault and Wakolbinger [15], Lyons, Pemantle and Peres [30], Biggins and Kyprianou [8] and Hardy and Harris [20].

There is a one-to-one correspondence between branching random walks and marked trees. Let us first introduce some notation. We label individuals in the branching random walk by their line of descent, so if  $u = i_1 \cdots i_n \in \mathcal{U} := \{\emptyset\} \cup \bigcup_{k=1}^{\infty} (\mathbb{N}^*)^k$  (where  $\mathbb{N}^* := \{1, 2, \dots\}$ ), then  $u$  is the  $i_n$ th child of the  $i_{n-1}$ th child of... of the  $i_1$ th child of the initial ancestor  $e$ . It is sometimes convenient to consider an element  $u \in \mathcal{U}$  as a “word” of length  $|u|$ , with  $\emptyset$  corresponding to  $e$ . We identify an individual  $u$  with its corresponding word.

If  $u, v \in \mathcal{U}$ , we denote by  $uv$  the concatenated word, with  $u\emptyset = \emptyset u = u$ .

Let  $\overline{\mathcal{U}} := \{(u, V(u)) : u \in \mathcal{U}, V : \mathcal{U} \rightarrow \mathbb{R}\}$ . Let  $\Omega$  be Neveu’s space of marked trees, which consists of all the subsets  $\omega$  of  $\overline{\mathcal{U}}$  such that the first component of  $\omega$  is a tree. [Recall that a tree  $t$  is a subset of  $\mathcal{U}$  satisfying: (i)  $\emptyset \in t$ ; (ii) if  $uj \in t$  for some  $j \in \mathbb{N}^*$ , then  $u \in t$ ; (iii) if  $u \in t$ , then  $uj \in t$  if and only if  $1 \leq j \leq \nu_u(t)$  for some nonnegative integer  $\nu_u(t)$ .]

Let  $\mathbb{T} : \Omega \rightarrow \Omega$  be the identity application. According to Neveu [32], there exists a probability  $\mathbf{P}$  on  $\Omega$  such that the law of  $\mathbb{T}$  under  $\mathbf{P}$  is the law of the branching random walk described in the Introduction.

Let us make a more intuitive presentation. For any  $\omega \in \Omega$ , let

$$(2.1) \quad \mathbb{T}^{\text{GW}}(\omega) := \text{the set of individuals ever born in } \omega,$$

$$(2.2) \quad \mathbb{T}(\omega) := \{(u, V(u)), u \in \mathbb{T}^{\text{GW}}(\omega), V \text{ such that } (u, V(u)) \in \omega\}.$$

[Of course,  $\mathbb{T}(\omega) = \omega$ .] In words,  $\mathbb{T}^{\text{GW}}$  is a Galton–Watson tree, with the population members as the vertices, whereas the *marked tree*  $\mathbb{T}$  corresponds to the branching random walk. It is more convenient to write (2.2) in an informal way:

$$\mathbb{T} = \{(u, V(u)), u \in \mathbb{T}^{\text{GW}}\}.$$

For any  $u \in \mathbb{T}^{\text{GW}}$ , the *shifted* Galton–Watson subtree generated by  $u$  is

$$(2.3) \quad \mathbb{T}_u^{\text{GW}} := \{x \in \mathcal{U} : ux \in \mathbb{T}^{\text{GW}}\}.$$

[By shifted, we mean that  $\mathbb{T}_u^{\text{GW}}$  is also rooted at  $e$ .] For any  $x \in \mathbb{T}_u^{\text{GW}}$ , let

$$(2.4) \quad |x|_u := |ux| - |u|,$$

$$(2.5) \quad V_u(x) := V(ux) - V(u).$$

As such,  $|x|_u$  stands for the (relative) generation of  $x$  as a vertex of the Galton–Watson tree  $\mathbb{T}_u^{\text{GW}}$ , and  $(V_u(x), x \in \mathbb{T}_u^{\text{GW}})$  the branching random walk which corresponds to the *shifted marked subtree*

$$\mathbb{T}_u := \{(x, V_u(x)), x \in \mathbb{T}_u^{\text{GW}}\}.$$

Let  $\mathcal{F}_n := \sigma\{(u, V(u)), u \in \mathbb{T}^{\text{GW}}, |u| \leq n\}$ , which is the sigma-field induced by the first  $n$  generations of the branching random walk. Let  $\mathcal{F}_\infty$  be the sigma-field induced by the whole branching random walk.

Assume now  $\psi(0) > 0$  and  $\psi(1) = 0$ . Let  $\mathbf{Q}$  be a probability on  $\Omega$  such that, for any  $n \geq 1$ ,

$$(2.6) \quad \mathbf{Q}|_{\mathcal{F}_n} := W_n \bullet \mathbf{P}|_{\mathcal{F}_n}.$$

Fix  $n \geq 1$ . Let  $w_n^{(n)}$  be a random variable taking values in  $\{u \in \mathbb{T}^{\text{GW}}, |u| = n\}$  such that, for any  $|u| = n$ ,

$$(2.7) \quad \mathbf{Q}\{w_n^{(n)} = u | \mathcal{F}_\infty\} = \frac{e^{-V(u)}}{W_n}.$$

We write  $\llbracket e, w_n^{(n)} \rrbracket = \{e =: w_0^{(n)}, w_1^{(n)}, w_2^{(n)}, \dots, w_n^{(n)}\}$  for the shortest path in  $\mathbb{T}^{\text{GW}}$  relating the root  $e$  to  $w_n^{(n)}$ , with  $|w_k^{(n)}| = k$  for any  $1 \leq k \leq n$ .

For any individual  $u \in \mathbb{T}^{\text{GW}} \setminus \{e\}$ , let  $\overleftarrow{u}$  be the parent of  $u$  in  $\mathbb{T}^{\text{GW}}$ , and

$$\Delta V(u) := V(u) - V(\overleftarrow{u}).$$

For  $1 \leq k \leq n$ , we write

$$(2.8) \quad \mathcal{I}_k^{(n)} := \{u \in \mathbb{T}^{\text{GW}} : |u| = k, \overleftarrow{u} = w_{k-1}^{(n)}, u \neq w_k^{(n)}\}.$$

In words,  $\mathcal{I}_k^{(n)}$  is the set of children of  $w_{k-1}^{(n)}$  except  $w_k^{(n)}$  or, equivalently, the set of the brothers of  $w_k^{(n)}$ , and is possibly empty. Finally, let us introduce the following sigma-field:

$$(2.9) \quad \mathcal{G}_n := \sigma \left\{ \sum_{x \in \mathcal{I}_k^{(n)}} \delta_{\Delta V(x)}, V(w_k^{(n)}), w_k^{(n)}, \mathcal{I}_k^{(n)}, 1 \leq k \leq n \right\},$$

where  $\delta$  denotes the Dirac measure.

The promised change-of-measures result is as follows. For any marked tree  $\mathbb{T}$ , we define its truncation  $\mathbb{T}^n$  at level  $n$  by  $\mathbb{T}^n := \{(x, V(x)), x \in \mathbb{T}^{\text{GW}}, |x| \leq n\}$ ; see Figure 1.




FIG. 1. Spine; The truncated shifted subtrees  $\mathbb{T}_x^{n-|x|}, \mathbb{T}_y^{n-|y|}, \mathbb{T}_z^{n-|z|}, \dots$  are actually rooted at  $e$ .

PROPOSITION 2.1. Assume  $\psi(0) > 0$  and  $\psi(1) = 0$ , and fix  $n \geq 1$ . Under probability  $\mathbf{Q}$ ,

(i) the random variables  $(\sum_{x \in \mathcal{F}_k^{(n)}} \delta_{\Delta V(x)}, \Delta V(w_k^{(n)}))$ ,  $1 \leq k \leq n$ , are i.i.d., distributed as  $(\sum_{x \in \mathcal{F}_1^{(1)}} \delta_{\Delta V(x)}, \Delta V(w_1^{(1)}))$ ;

(ii) conditionally on  $\mathcal{G}_n$ , the truncated shifted marked subtrees  $\mathbb{T}_x^{n-|x|}$ , for  $x \in \bigcup_{k=1}^n \mathcal{F}_k^{(n)}$ , are independent; the conditional distribution of  $\mathbb{T}_x^{n-|x|}$  (for any  $x \in \bigcup_{k=1}^n \mathcal{F}_k^{(n)}$ ) under  $\mathbf{Q}$ , given  $\mathcal{G}_n$ , is identical to the distribution of  $\mathbb{T}^{n-|x|}$  under  $\mathbf{P}$ .

Throughout the paper, let  $((S_i, \sigma_i), i \geq 1)$  be such that  $(S_i - S_{i-1}, \sigma_i)$ , for  $i \geq 1$  (with  $S_0 = 0$ ), are i.i.d. random vectors under  $\mathbf{Q}$  and distributed as  $(V(w_1^{(1)}), \#\mathcal{F}_1^{(1)})$ .

COROLLARY 2.2. Assume  $\psi(0) > 0$  and  $\psi(1) = 0$ , and fix  $n \geq 1$ .

(i) Under  $\mathbf{Q}$ ,  $((V(w_k^{(n)}), \#\mathcal{F}_k^{(n)}), 1 \leq k \leq n)$  is distributed as  $((S_k, \sigma_k), 1 \leq k \leq n)$ . In particular, under  $\mathbf{Q}$ ,  $(V(w_k^{(n)}), 1 \leq k \leq n)$  is distributed as  $(S_k, 1 \leq k \leq n)$ .



(ii) For any measurable function  $F : \mathbb{R} \rightarrow \mathbb{R}_+$ ,

$$(2.10) \quad \mathbf{E}_{\mathbf{Q}}\{F(S_1)\} = \mathbf{E}\left\{\sum_{|u|=1} e^{-V(u)} F(V(u))\right\}.$$

In particular, we have  $\mathbf{E}_{\mathbf{Q}}\{S_1\} = 0$  under (1.2) and (1.3).

Corollary 2.2 follows immediately from Proposition 2.1, and can be found in several papers (e.g., Biggins and Kyprianou [9]).

We present two collections of probability estimates for  $(S_n)$  and for  $(V(u), |u| = 1)$ , respectively. They are simple consequences of Proposition 2.1, and will be of frequent use in the rest of the paper.

COROLLARY 2.3. Assume (1.2) and (1.3). Then

$$(2.11) \quad \mathbf{E}_{\mathbf{Q}}\{e^{aS_1}\} < \infty \quad \forall |a| \leq c_2,$$

$$(2.12) \quad \mathbf{Q}\{|S_n| \geq x\} \leq 2 \exp\left(-c_3 \min\left\{x, \frac{x^2}{n}\right\}\right) \quad \forall n \geq 1, \forall x \geq 0,$$

$$(2.13) \quad \mathbf{Q}\left\{\min_{1 \leq k \leq n} S_k > 0\right\} \sim \frac{c_4}{n^{1/2}}, \quad n \rightarrow \infty,$$

$$(2.14) \quad \sup_{n \geq 1} n^{1/2} \mathbf{E}_{\mathbf{Q}}\{e^{b \min_{0 \leq i \leq n} S_i}\} < \infty \quad \forall b \geq 0,$$

where  $c_2 := \min\{\delta_+, 1 + \delta_-\}$ . Furthermore, for any  $C \geq c > 0$ , we have

$$(2.15) \quad \mathbf{Q}\left\{\max_{0 \leq j, k \leq n, |j-k| \leq c \log n} |S_j - S_k| \geq C \log n\right\} \leq 2cn^{-(c_3 C - 1)} \log n \quad \forall n \geq 2.$$

COROLLARY 2.4. Assume (1.1), (1.2) and (1.3). Let  $0 < a \leq 1$ . Then

$$(2.16) \quad \mathbf{E}_{\mathbf{Q}}\left\{\left(\sum_{|u|=1} e^{-aV(u)}\right)^{\rho(a)}\right\} < \infty,$$

$$(2.17) \quad \mathbf{Q}\left\{\sup_{|u|=1} |V(u)| \geq x\right\} \leq c_5 e^{-c_6 x} \quad \forall x \geq 0,$$

with  $\rho(a) := \frac{\delta \delta_+}{1 + a \delta + \delta_+}$ , where  $\delta$  and  $\delta_+$  are the constants in (1.1) and (1.2), respectively.

PROOF OF COROLLARY 2.3. By Corollary 2.2 (ii),  $\mathbf{E}_{\mathbf{Q}}\{e^{aS_1}\} = \mathbf{E}\{\sum_{|u|=1} e^{(a-1)V(u)}\}$ , which, according to (1.2), is finite as long as  $|a| \leq c_2$ . This proves (2.11).

Once we have the exponential integrability in (2.11) for  $(S_n)$ , standard probability estimates for sums of i.i.d. random variables yield (2.12), (2.13) and (2.14); see Petrov [34]’s Theorem 2.7, Bingham [10] and Kozlov [24]’s Theorem A, respectively.

To check (2.15), we observe that the probability term on the left-hand side of (2.15) is bounded by  $\sum_{0 \leq j < k \leq n, k-j \leq c \log n} \mathbf{Q}\{|S_{k-j}| \geq C \log n\}$ . By (2.12),  $\mathbf{Q}\{|S_{k-j}| \geq C \log n\} \leq 2n^{-c_3 C}$  for  $k - j \leq c \log n$ . This yields (2.15).  $\square$

**PROOF OF COROLLARY 2.4.** Write  $\rho := \rho(a)$ . We have  $\mathbf{E}_{\mathbf{Q}}\{(\sum_{|u|=1} e^{-aV(u)})^\rho\} = \mathbf{E}_{\mathbf{Q}}\{W_{1,a}^\rho\} = \mathbf{E}\{W_{1,a}^\rho W_{1,1}\}$ . Let  $N := \sum_{|u|=1} 1$ . By Hölder’s inequality,  $W_{1,a} \leq W_{1,1+\delta_+}^{a/(1+\delta_+)} N^{(1-a+\delta_+)/(1+\delta_+)}$ , whereas  $W_{1,1} \leq W_{1,1+\delta_+}^{1/(1+\delta_+)} N^{\delta_+/(1+\delta_+)}$ . Therefore, by means of another application of Hölder’s inequality,  $\mathbf{E}\{W_{1,a}^\rho W_{1,1}\} \leq [\mathbf{E}(W_{1,1+\delta_+})]^{(1+a\rho)/(1+\delta_+)} [\mathbf{E}(N^{1+\delta})]^{(\delta_+-a\rho)/(1+\delta_+)}$ , which is finite [by (1.2) and (1.1)]. This implies (2.16).

To prove (2.17), we write  $A := \{\sup_{|u|=1} |V(u)| \geq x\}$ . By Chebyshev’s inequality,  $\mathbf{P}(A) \leq c_7 e^{-c_8 x}$ , where  $c_7 := \mathbf{E}(\sum_{|u|=1} e^{c_8 |V(u)|}) < \infty$  as long as  $0 < c_8 \leq \min\{\delta_-, 1 + \delta_+\}$  [by (1.2)]. Thus,  $\mathbf{Q}(A) = \mathbf{E}\{\sum_{|u|=1} e^{-V(u)} \mathbf{1}_A\} \leq c_9 [\mathbf{P}(A)]^{\rho(1)/(1+\rho(1))}$ , where  $c_9 := [\mathbf{E}\{(\sum_{|u|=1} e^{-V(u)})^{1+\rho(1)}\}]^{1/(1+\rho(1))} < \infty$ . Now (2.17) follows from (2.16), with  $c_6 := \frac{c_8 \rho(1)}{1+\rho(1)}$ .  $\square$

**3. Preliminary: small values of  $W_n$ .** This preliminary section is devoted to the study of the small values of  $W_n$ . Throughout the section, we assume (1.1), (1.2) and (1.3). We define two important events:

$$(3.1) \quad \mathcal{S} := \{\text{the system’s ultimate survival}\},$$

$$(3.2) \quad \mathcal{S}_n := \{\text{the system’s survival after } n \text{ generations}\} = \{W_n > 0\}.$$

Clearly,  $\mathcal{S} \subset \mathcal{S}_n$ . Recall (see, e.g., Harris [21], page 16) that, for some constant  $c_{10}$  and all  $n \geq 1$ ,

$$(3.3) \quad \mathbf{P}\{\mathcal{S}_n \setminus \mathcal{S}\} \leq e^{-c_{10} n}.$$

Here is the main result of the section.

**PROPOSITION 3.1.** *Assume (1.1), (1.2) and (1.3). For any  $\varepsilon > 0$ , there exists  $\vartheta > 0$  such that, for all sufficiently large  $n$ ,*

$$(3.4) \quad \mathbf{P}\{n^{1/2} W_n < n^{-\varepsilon} | \mathcal{S}\} \leq n^{-\vartheta}.$$

The proof of Proposition 3.1 relies on Neveu’s multiplicative martingale. Recall that under assumption (1.3), there exists a nonnegative random variable  $\xi^*$ , with  $\mathbf{P}\{\xi^* > 0\} > 0$ , such that

$$(3.5) \quad \xi^* \stackrel{\text{law}}{=} \sum_{|u|=1} \xi_u^* e^{-V(u)},$$

where, given  $\{(u, V(u)), |u| = 1\}$ ,  $\xi_u^*$  are independent copies of  $\xi^*$ , and “ $\stackrel{law}{=}$ ” stands for identity in distribution. Moreover, there is uniqueness of the distribution of  $\xi^*$  up to a scale change (see Liu [27]); in the rest of the paper we take the version of  $\xi^*$  as the unique one satisfying  $\mathbf{E}\{e^{-\xi^*}\} = \frac{1}{2}$ .

Let us introduce the Laplace transform of  $\xi^*$ :

$$(3.6) \quad \varphi^*(t) := \mathbf{E}\{e^{-t\xi^*}\}, \quad t \geq 0.$$

Let

$$(3.7) \quad W_n^* := \prod_{|u|=n} \varphi^*(e^{-V(u)}), \quad n \geq 1.$$

The process  $(W_n^*, n \geq 1)$  is also a martingale (Liu [27]). Following Neveu [33], we call  $W_n^*$  an associated “multiplicative martingale.”

The martingale  $W_n^*$  being bounded, it converges almost surely (when  $n \rightarrow \infty$ ) to, say,  $W_\infty^*$ . Let us recall from Liu [27] (see also Kyprianou [25]) that, for some  $c^* > 0$ ,

$$(3.8) \quad \log \frac{1}{W_\infty^*} \stackrel{law}{=} \xi^*,$$

$$(3.9) \quad \log\left(\frac{1}{\varphi^*(t)}\right) \sim c^* t \log\left(\frac{1}{t}\right), \quad t \rightarrow 0.$$

We first prove the following lemma:

LEMMA 3.2. Assume (1.1), (1.2) and (1.3). There exist  $\kappa > 0$  and  $a_0 \geq 1$  such that

$$(3.10) \quad \mathbf{E}\{(W_\infty^*)^a | W_\infty^* < 1\} \leq a^{-\kappa}, \quad \forall a \geq a_0,$$

$$(3.11) \quad \mathbf{E}\{(W_n^*)^a \mathbf{1}_{\mathcal{S}_n}\} \leq a^{-\kappa} + e^{-c_{10}n}, \quad \forall n \geq 1, \forall a \geq a_0.$$

PROOF. We are grateful to John Biggins for fixing a mistake in the original proof.

We first prove (3.10). In view of (3.8), it suffices to show that

$$(3.12) \quad \mathbf{E}\{e^{-a\xi^*} | \xi^* > 0\} \leq a^{-\kappa}, \quad a \geq a_0.$$

Let  $q \in [0, 1)$  be the system’s extinction probability. Let  $N := \sum_{|u|=1} 1$ . It is well known for Galton–Watson trees that  $q$  is the unique solution of  $\mathbf{E}(q^N) = q$  (for  $q \in [0, 1)$ ); see, for example, Harris [21], page 7. By (3.5),  $\varphi^*(t) = \mathbf{E}\{\prod_{|u|=1} \varphi^*(te^{-V(u)})\}$ . Therefore, by (3.6),  $\mathbf{P}\{\xi^* = 0\} = \varphi^*(\infty) = \lim_{t \rightarrow \infty} \mathbf{E}\{\prod_{|u|=1} \varphi^*(te^{-V(u)})\}$ , which, by dominated convergence, is  $= \mathbf{E}\{(\varphi^*(\infty))^N\} = \mathbf{E}\{(\mathbf{P}\{\xi^* = 0\})^N\}$ . Since  $\mathbf{P}\{\xi^* = 0\} < 1$ , this yields  $\mathbf{P}\{\xi^* = 0\} = q$ .

Following Biggins and Grey [6], we note that, for any  $t \geq 0$ ,

$$\mathbf{E}\{e^{-t\xi^*}\} = q + (1 - q)\mathbf{E}\{e^{-t\xi^*} \mid \xi^* > 0\}.$$

Let  $\widehat{\xi}$  be a random variable such that  $\mathbf{E}\{e^{-t\widehat{\xi}}\} = \mathbf{E}\{e^{-t\xi^*} \mid \xi^* > 0\}$  for any  $t \geq 0$ . Let  $Y$  be a random variable independent of everything else, such that  $\mathbf{P}\{Y = 0\} = q = 1 - \mathbf{P}\{Y = 1\}$ . Then  $\xi^*$  and  $Y\widehat{\xi}$  have the same law and, by (3.5), so do  $\xi^*$  and  $\sum_{|u|=1} e^{-V(u)} Y_u \widehat{\xi}_u$ , where, given  $\{u, |u| = 1\}$ ,  $(Y_u, \widehat{\xi}_u)$  are independent copies of  $(Y, \widehat{\xi})$ , independent of  $\{V(u), |u| = 1\}$ . Since  $\{\sum_{|u|=1} e^{-V(u)} Y_u \widehat{\xi}_u > 0\} = \{\sum_{|u|=1} Y_u > 0\}$ , this leads to

$$\mathbf{E}\{e^{-t\widehat{\xi}}\} = \mathbf{E}\left\{e^{-t \sum_{|u|=1} e^{-V(u)} Y_u \widehat{\xi}_u} \mid \sum_{|u|=1} Y_u > 0\right\}, \quad t \geq 0.$$

Let  $\widehat{\varphi}(t) := \mathbf{E}\{e^{-t\widehat{\xi}}\}$ ,  $t \geq 0$ . Then for any  $t \geq 0$  and  $c > 0$ ,

$$\widehat{\varphi}(t) = \mathbf{E}\left\{\prod_{|u|=1} \widehat{\varphi}(te^{-V(u)} Y_u) \mid \sum_{|u|=1} Y_u > 0\right\} \leq \mathbf{E}\left\{[\widehat{\varphi}(te^{-c})]^{N_c} \mid \sum_{|u|=1} Y_u > 0\right\},$$

where  $N_c := \sum_{|u|=1} \mathbf{1}_{\{Y_u=1, |V(u)| \leq c\}}$ . By monotone convergence,  $\lim_{c \rightarrow \infty} \mathbf{E}\{N_c \mid \sum_{|u|=1} Y_u > 0\} = \mathbf{E}\{\sum_{|u|=1} Y_u \mid \sum_{|u|=1} Y_u > 0\} > 1$  [because  $\mathbf{P}\{\sum_{|u|=1} Y_u \geq 2\} > 0$  by assumption (1.3)]. We can therefore choose and fix a constant  $c > 0$  such that  $\mathbf{E}\{N_c \mid \sum_{|u|=1} Y_u > 0\} > 1$ . By writing  $\widehat{f}(s) := \mathbf{E}\{s^{N_c} \mid \sum_{|u|=1} Y_u > 0\}$ , we have

$$\widehat{\varphi}(t) \leq \widehat{f}(\widehat{\varphi}(te^{-c})), \quad \forall t \geq 0.$$

Iterating the inequality yields that, for any  $t \geq 0$  and any  $n \geq 1$ ,

$$(3.13) \quad \mathbf{E}\{e^{-t\widehat{\xi}}\} \leq \widehat{f}^{(n)}(\mathbf{E}\{e^{-te^{-nc}\widehat{\xi}}\}), \quad \text{that is, } \mathbf{E}\{e^{-te^{-nc}\widehat{\xi}}\} \leq \widehat{f}^{(n)}(\mathbf{E}\{e^{-t\widehat{\xi}}\}),$$

where  $\widehat{f}^{(n)}$  denotes the  $n$ th iterate of  $\widehat{f}$ . It is well known for Galton–Watson trees (Athreya and Ney [3], Section I.11) that, for any  $s \in [0, 1)$ ,  $\lim_{n \rightarrow \infty} \gamma^{-n} \widehat{f}^{(n)}(s)$  converges to a finite limit, with  $\gamma := (\widehat{f})'(0) \leq \mathbf{P}\{\sum_{|u|=1} Y_u = 1 \mid \sum_{|u|=1} Y_u > 0\} < 1$ . Therefore, (3.13) yields (3.12), and thus (3.10).

It remains to check (3.11). Let  $a \geq 1$ . Since  $((W_n^*)^a, n \geq 0)$  is a bounded submartingale,  $\mathbf{E}\{(W_n^*)^a \mathbf{1}_{\mathcal{S}_n}\} \leq \mathbf{E}\{(W_\infty^*)^a \mathbf{1}_{\mathcal{S}_n}\}$ . Recall that  $W_\infty^* \leq 1$ ; thus,

$$\mathbf{E}\{(W_n^*)^a \mathbf{1}_{\mathcal{S}_n}\} \leq \mathbf{E}\{(W_\infty^*)^a \mathbf{1}_{\mathcal{S}}\} + \mathbf{P}\{\mathcal{S}_n \setminus \mathcal{S}\}.$$

By (3.3),  $\mathbf{P}\{\mathcal{S}_n \setminus \mathcal{S}\} \leq e^{-c10^n}$ . To estimate  $\mathbf{E}\{(W_\infty^*)^a \mathbf{1}_{\mathcal{S}}\}$ , we identify  $\mathcal{S}$  with  $\{W_\infty^* < 1\}$ : on the one hand,  $\mathcal{S}^c \subset \{W_n^* = 1\}$ , for all sufficiently large  $n\} \subset \{W_\infty^* = 1\}$ ; on the other hand, by (3.8),  $\mathbf{P}\{W_\infty^* < 1\} = \mathbf{P}\{\xi^* > 0\} = 1 - q = \mathbf{P}(\mathcal{S})$ . Therefore,  $\mathcal{S} = \{W_\infty^* < 1\}$ ,  $\mathbf{P}$ -a.s. Consequently,  $\mathbf{E}\{(W_\infty^*)^a \mathbf{1}_{\mathcal{S}}\} = \mathbf{E}\{(W_\infty^*)^a \mathbf{1}_{\{W_\infty^* < 1\}}\}$ , which, according to (3.10), is bounded by  $a^{-\kappa}$ , for  $a \geq a_0$ . Lemma 3.2 is proved.  $\square$

We are now ready for the proof of Proposition 3.1.

PROOF OF PROPOSITION 3.1. Let  $c_{11} > 0$  be such that  $\mathbf{P}\{\xi^* \leq c_{11}\} \geq \frac{1}{2}$ . Then  $\varphi^*(t) = \mathbf{E}\{e^{-t\xi^*}\} \geq e^{-c_{11}t}\mathbf{P}\{\xi^* \leq c_{11}\} \geq \frac{1}{2}e^{-c_{11}t}$  and, thus,  $\log(\frac{1}{\varphi^*(t)}) \leq c_{11}t + \log 2$ . Together with (3.9), this yields, on the event  $\mathcal{S}_n$ ,

$$\begin{aligned} \log\left(\frac{1}{W_n^*}\right) &= \sum_{|u|=n} \log\left(\frac{1}{\varphi^*(e^{-V(u)})}\right) \\ &\leq \sum_{|u|=n} \mathbf{1}_{\{V(u) \geq 1\}} c_{12} V(u) e^{-V(u)} + \sum_{|u|=n} \mathbf{1}_{\{V(u) < 1\}} (c_{11} e^{-V(u)} + \log 2). \end{aligned}$$

Since  $W_n = \sum_{|u|=n} e^{-V(u)}$ , we obtain, on  $\mathcal{S}_n$ , for any  $\lambda \geq 1$ ,

$$(3.14) \quad \log\left(\frac{1}{W_n^*}\right) \leq c_{13} \lambda W_n + c_{12} \sum_{|u|=n} \mathbf{1}_{\{V(u) \geq \lambda\}} V(u) e^{-V(u)},$$

where  $c_{13} := c_{11} + c_{12} + e \log 2$ . Note that  $c_{12}$  and  $c_{13}$  do not depend on  $\lambda$ .

Let  $0 < y \leq 1$ . Since  $\mathcal{S} \subset \mathcal{S}_n$ , it follows that, for  $c_{14} := c_{12} + c_{13}$ ,

$$\begin{aligned} (3.15) \quad \mathbf{P}\{\lambda W_n < y | \mathcal{S}_n\} &\leq \mathbf{P}\left\{\log\left(\frac{1}{W_n^*}\right) < c_{14} y | \mathcal{S}_n\right\} \\ &\quad + \mathbf{P}\left\{\sum_{|u|=n} \mathbf{1}_{\{V(u) \geq \lambda\}} V(u) e^{-V(u)} \geq y | \mathcal{S}_n\right\} \\ &=: \text{RHS}_{(3.15)}^1 + \text{RHS}_{(3.15)}^2, \end{aligned}$$

with obvious notation.

Recall that  $\mathbf{P}(\mathcal{S}_n) \geq \mathbf{P}(\mathcal{S}) = 1 - q$ . By Chebyshev's inequality,

$$\text{RHS}_{(3.15)}^1 \leq e^{c_{14}} \mathbf{E}\{(W_n^*)^{1/y} | \mathcal{S}_n\} \leq \frac{e^{c_{14}}}{1 - q} \mathbf{E}\{(W_n^*)^{1/y} \mathbf{1}_{\mathcal{S}_n}\}.$$

By (3.11), for  $n \geq 1$  and  $0 < y \leq \frac{1}{a_0}$ , with  $c_{15} := e^{c_{14}}/(1 - q)$ ,

$$(3.16) \quad \text{RHS}_{(3.15)}^1 \leq c_{15}(y^\kappa + e^{-c_{10}n}).$$

To estimate  $\text{RHS}_{(3.15)}^2$ , we observe that

$$\begin{aligned} \text{RHS}_{(3.15)}^2 &\leq \frac{1}{1 - q} \mathbf{P}\left\{\sum_{|u|=n} \mathbf{1}_{\{V(u) \geq \lambda\}} V(u) e^{-V(u)} \geq y\right\} \\ &\leq \frac{1}{(1 - q)y} \mathbf{E}\left\{\sum_{|u|=n} \mathbf{1}_{\{V(u) \geq \lambda\}} V(u) e^{-V(u)}\right\} \\ &= \frac{1}{(1 - q)y} \mathbf{E}\mathbf{Q}\left\{\sum_{|u|=n} \mathbf{1}_{\{V(u) \geq \lambda\}} \frac{V(u) e^{-V(u)}}{W_n}\right\} \\ &= \frac{1}{(1 - q)y} \mathbf{E}\mathbf{Q}\{V(w_n^{(n)}) \mathbf{1}_{\{V(w_n^{(n)}) \geq \lambda\}}\}. \end{aligned}$$

By Corollary 2.2(i),  $\mathbf{E}_{\mathbf{Q}}\{V(w_n^{(n)})\mathbf{1}_{\{V(w_n^{(n)}) \geq \lambda\}}\} = \mathbf{E}_{\mathbf{Q}}\{S_n \mathbf{1}_{\{S_n \geq \lambda\}}\} \leq (\mathbf{E}_{\mathbf{Q}}\{S_n^2\})^{1/2} \times (\mathbf{Q}\{S_n \geq \lambda\})^{1/2}$ , which, by (2.12), is bounded by  $c_{16}n \exp(-c_3 \min\{\lambda, \frac{\lambda^2}{n}\})$ . Accordingly,  $\text{RHS}_{(3.15)}^2 \leq \frac{c_{17}n}{y} \exp(-c_3 \min\{\lambda, \frac{\lambda^2}{n}\})$ . Together with (3.15) and (3.16), it yields that, for  $0 < y \leq \frac{1}{a_0}$ ,

$$\mathbf{P}\{\lambda W_n < y | \mathcal{S}_n\} \leq c_{15}(y^\kappa + e^{-c_{10}n}) + \frac{c_{17}n}{y} \exp\left(-c_3 \min\left\{\lambda, \frac{\lambda^2}{n}\right\}\right).$$

Let  $\lambda := n^{1/2}y^{-\kappa/2}$ . The inequality becomes, for  $0 < y \leq \frac{1}{a_0}$  and  $n \geq 1$ ,

$$\begin{aligned} \mathbf{P}\{n^{1/2}W_n < y^{(\kappa+2)/2} | \mathcal{S}_n\} \\ \leq c_{15}(y^\kappa + e^{-c_{10}n}) + \frac{c_{17}n}{y} \exp\left(-c_3 \frac{\min\{n^{1/2}y^{\kappa/2}, 1\}}{y^\kappa}\right). \end{aligned}$$

This readily yields Proposition 3.1.  $\square$

REMARK. Under the additional assumption that  $\{u, |u| = 1\}$  contains at least two elements almost surely, it is possible (Liu [28]) to improve (3.10):  $\mathbf{E}\{(W_\infty^*)^a | W_\infty^* < 1\} \leq \exp\{-a^\kappa\}$  for some  $\kappa_1 > 0$  and all sufficiently large  $a$ , from which one can deduce the stronger version of Proposition 3.1: for any  $\varepsilon > 0$ , there exists  $\vartheta_1 > 0$  such that  $\mathbf{P}\{n^{1/2}W_n < n^{-\varepsilon} | \mathcal{S}\} \leq \exp(-n^{\vartheta_1})$  for all sufficiently large  $n$ .

We complete this section with the following estimate which will be useful in the proof of Theorem 1.5.

LEMMA 3.3. Assume (1.1), (1.2) and (1.3). For any  $0 < s < 1$ ,

$$(3.17) \quad \sup_{n \geq 1} \mathbf{E}\left\{\left(\log \frac{1}{W_n^*}\right)^s\right\} < \infty.$$

PROOF. Let  $x > 1$ . By Chebyshev’s inequality,  $\mathbf{P}\{\log(\frac{1}{W_n^*}) \geq x\} = \mathbf{P}\{e^x W_n^* \leq 1\} \leq e \mathbf{E}\{e^{-e^x W_n^*}\}$ . Since  $W_n^*$  is a martingale, it follows from Jensen’s inequality that  $\mathbf{E}\{e^{-e^x W_n^*}\} \leq \mathbf{E}\{e^{-e^x W_\infty^*}\} \leq \mathbf{P}\{W_\infty^* \leq e^{-x/2}\} + \exp(-e^{x/2})$ . Therefore,

$$(3.18) \quad \mathbf{P}\left\{\log\left(\frac{1}{W_n^*}\right) \geq x\right\} \leq e \mathbf{P}\{W_\infty^* \leq e^{-x/2}\} + \exp(1 - e^{x/2}).$$

On the other hand, by integration by parts,  $\int_0^\infty e^{-ty} \mathbf{P}(\xi^* \geq y) dy = \frac{1 - \mathbf{E}(e^{-t\xi^*})}{t} = \frac{1 - \varphi^*(t)}{t}$ , which, according to (3.9), is  $\leq c_{18} \log(\frac{1}{t})$  for  $0 < t \leq \frac{1}{2}$ . Therefore, for  $a \geq 2$ ,  $c_{18} \log a \geq \int_0^\infty e^{-y/a} \mathbf{P}(\xi^* \geq y) dy \geq \int_0^a e^{-y/a} \mathbf{P}(\xi^* \geq a) dy = (1 -$

$e^{-1})a\mathbf{P}(\xi^* \geq a)$ . That is,  $\mathbf{P}(\xi^* \geq a) \leq \frac{c_{18}}{1-e^{-1}} \frac{\log a}{a}$  or, equivalently,  $\mathbf{P}(W_\infty^* \leq e^{-a}) \leq \frac{c_{18}}{1-e^{-1}} \frac{\log a}{a}$ , for  $a \geq 2$ . Substituting this in (3.18) gives that, for any  $x \geq 4$ ,

$$\mathbf{P}\left\{\log\left(\frac{1}{W_n^*}\right) \geq x\right\} \leq \frac{2ec_{18}}{1-e^{-1}} \frac{\log(x/2)}{x} + \exp(1 - e^{x/2}).$$

Lemma 3.3 follows immediately.  $\square$

**4. Proof of Theorem 1.2: upper bound in (1.8).** Assume (1.1), (1.2) and (1.3). This section is devoted to proving the upper bound in (1.8): conditionally on the system’s survival,

$$(4.1) \quad \liminf_{n \rightarrow \infty} \frac{1}{\log n} \inf_{|u|=n} V(u) \leq \frac{1}{2}, \quad \text{a.s.}$$

The proof borrows some ideas from Bramson [12]. We fix  $-\infty < a < b < \infty$  and  $\varepsilon > 0$ . Let  $\ell_1 \leq \ell_2 \leq 2\ell_1$  be integers; we are interested in the asymptotic case  $\ell_1 \rightarrow \infty$ . Consider  $n \in [\ell_1, \ell_2] \cap \mathbb{Z}$ . Let  $0 < c_{19} < 1$  be a constant, and let

$$g_n(k) := \min\{c_{19}k^{1/3}, c_{19}(n-k)^{1/3} + a \log \ell_1, n^\varepsilon\}, \quad 0 \leq k \leq n.$$

Let  $\mathbb{L}_n$  be the set of individuals  $x \in \mathbb{T}^{\text{GW}}$  with  $|x| = n$  such that

$$g_n(k) \leq V(x_k) \leq c_{20}k, \quad \forall 0 \leq k \leq n \quad \text{and} \quad a \log \ell_1 \leq V(x) \leq b \log \ell_1,$$

where  $x_0 := e, x_1, \dots, x_n := x$  are the vertices on the shortest path in  $\mathbb{T}^{\text{GW}}$  relating the root  $e$  and the vertex  $x$ , with  $|x_k| = k$  for any  $0 \leq k \leq n$ . We consider the measurable event

$$F_{\ell_1, \ell_2} := \bigcup_{n=\ell_1}^{\ell_2} \bigcup_{|x|=n} \{x \in \mathbb{L}_n\}.$$

We start by estimating the first moment of  $\#F_{\ell_1, \ell_2}$ :  $\mathbf{E}(\#F_{\ell_1, \ell_2}) = \sum_{n=\ell_1}^{\ell_2} \mathbf{E}\{\sum_{|x|=n} \mathbf{1}_{\{x \in \mathbb{L}_n\}}\}$ . Since  $\mathbf{E}\{\sum_{|x|=n} \mathbf{1}_{\{x \in \mathbb{L}_n\}}\} = \mathbf{E}_{\mathbf{Q}}\{\sum_{|x|=n} \frac{e^{-V(x)}}{W_n} e^{V(x)} \times \mathbf{1}_{\{x \in \mathbb{L}_n\}}\} = \mathbf{E}_{\mathbf{Q}}\{e^{V(w_n^{(n)})} \mathbf{1}_{\{w_n^{(n)} \in \mathbb{L}_n\}}\}$ , we can apply Corollary 2.2 to see that

$$\begin{aligned} \mathbf{E}(\#F_{\ell_1, \ell_2}) &= \sum_{n=\ell_1}^{\ell_2} \mathbf{E}_{\mathbf{Q}}\{e^{S_n} \mathbf{1}_{\{g_n(k) \leq S_k \leq c_{20}k, \forall 0 \leq k \leq n, a \log \ell_1 \leq S_n \leq b \log \ell_1\}}\} \\ &\geq \sum_{n=\ell_1}^{\ell_2} \ell_1^a \mathbf{Q}\{g_n(k) \leq S_k \leq c_{20}k, \forall 0 \leq k \leq n, a \log \ell_1 \leq S_n \leq b \log \ell_1\}. \end{aligned}$$

We choose (and fix) the constants  $c_{19}$  and  $c_{20}$  such that  $\mathbf{Q}\{c_{19} < S_1 < c_{20}\} > 0$ . Then,<sup>1</sup> the probability  $\mathbf{Q}\{\cdot\}$  on the right-hand side is  $\ell_1^{-(3/2)+o(1)}$ , for  $\ell_1 \rightarrow \infty$ .

<sup>1</sup>An easy way to see why  $-\frac{3}{2}$  should be the correct exponent for the probability is to split the event into three pieces: the first piece involving  $S_k$  for  $0 \leq k \leq \frac{n}{3}$ , the second piece for  $\frac{n}{3} \leq k \leq \frac{2n}{3}$ ,

Accordingly,

$$(4.2) \quad \mathbf{E}(\#F_{\ell_1, \ell_2}) \geq (\ell_2 - \ell_1 + 1)\ell_1^{a-(3/2)+o(1)}.$$

We now proceed to estimate the second moment of  $\#F_{\ell_1, \ell_2}$ . By definition,

$$\begin{aligned} \mathbf{E}[(\#F_{\ell_1, \ell_2})^2] &= \sum_{n=\ell_1}^{\ell_2} \sum_{m=\ell_1}^{\ell_2} \mathbf{E} \left\{ \sum_{|x|=n} \sum_{|y|=m} \mathbf{1}_{\{x \in \mathbb{L}_n, y \in \mathbb{L}_m\}} \right\} \\ &\leq 2 \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \mathbf{E} \left\{ \sum_{|x|=n} \sum_{|y|=m} \mathbf{1}_{\{x \in \mathbb{L}_n, y \in \mathbb{L}_m\}} \right\}. \end{aligned}$$

We look at the double sum  $\sum_{|x|=n} \sum_{|y|=m}$  on the right-hand side. By considering  $z$ , the youngest common ancestor of  $x$  and  $y$ , and writing  $k := |z|$ , we arrive at

$$\sum_{|x|=n} \sum_{|y|=m} \mathbf{1}_{\{x \in \mathbb{L}_n, y \in \mathbb{L}_m\}} = \sum_{k=0}^n \sum_{|z|=k} \sum_{(u,v)} \mathbf{1}_{\{zu \in \mathbb{L}_n, zv \in \mathbb{L}_m\}},$$

where the double sum  $\sum_{(u,v)}$  is over  $u, v \in \mathbb{T}_z^{\text{GW}}$  such that  $|u|_z = n - k$  and  $|v|_z = m - k$  and that the unique common ancestor of  $u$  and  $v$  in  $\mathbb{T}_z^{\text{GW}}$  is the root. Therefore,

$$\begin{aligned} \mathbf{E}[(\#F_{\ell_1, \ell_2})^2] &\leq 2 \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=0}^n \mathbf{E} \left\{ \sum_{|z|=k} \sum_{(u,v)} \mathbf{1}_{\{zu \in \mathbb{L}_n, zv \in \mathbb{L}_m\}} \right\} \\ &=: 2 \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=0}^n \Lambda_{k,n,m}. \end{aligned}$$

We estimate  $\Lambda_{k,n,m}$  according to three different situations.

*First situation:*  $0 \leq k \leq \lfloor n^\epsilon \rfloor$ . Let  $V_z(u) := V(zu) - V(z)$  as in Section 2. We have  $0 \leq g_n(k) \leq V(z) \leq c_{20}n^\epsilon$ , and  $V(zu_i) \geq 0$  for  $0 \leq i \leq n - k$  and  $V(zu_{n-k}) \leq b \log \ell_1$ , where  $u_0 := e, u_1, \dots, u_{n-k}$  are the vertices on the shortest path in  $\mathbb{T}_z^{\text{GW}}$  relating the root  $e$  and the vertex  $u$ , with  $|u_i|_z = i$  for any  $0 \leq i \leq n - k$ . Therefore,

---

and the third piece for  $\frac{2n}{3} \leq k \leq n$ . The probability of the first piece is  $n^{-(1/2)+o(1)}$  (it is essentially the probability of  $S_k$  being positive for  $1 \leq k \leq \frac{n}{3}$ , because conditionally on this,  $S_k$  converges weakly, after a suitable normalization, to a Brownian meander; see Bolthausen [11]). Similarly, the probability of the third piece is  $n^{-(1/2)+o(1)}$ . The second piece essentially says that after  $\frac{n}{3}$  steps, the random walk should lie in an interval of length of order  $\log n$ ; this probability is also  $n^{-(1/2)+o(1)}$ . Putting these pieces together yields the claimed exponent  $-\frac{3}{2}$ .

For a rigorous proof, the upper bound—not required here—is easier since we can only look at the event that the walk stays positive during  $n$  steps (with the same condition upon the random variable  $S_n$ ), whereas the lower bound needs some tedious but elementary writing, based on the Markov property. Similar arguments are used for the random walk  $(S_k)$  in several other places in the paper, without further mention.



$V_z(u_i) \geq -c_{20}n^\varepsilon$  for  $0 \leq i \leq n - k$ , and  $V_z(u) \leq b \log \ell_1$ . Accordingly,

$$\Lambda_{k,n,m} \leq \mathbf{E} \left\{ \sum_{|z|=k} \sum_{v \in \mathbb{T}_z^{\text{GW}}, |v|_z=m-k} \mathbf{1}_{\{zv \in \mathbb{L}_m\}} B_{n-k} \right\},$$

where

$$\begin{aligned} B_{n-k} &:= \mathbf{E} \left\{ \sum_{|x|=n-k} \mathbf{1}_{\{V(x_i) \geq -c_{20}n^\varepsilon, \forall 0 \leq i \leq n-k, V(x) \leq b \log \ell_1\}} \right\} \\ &= \mathbf{E} \mathbf{Q} \left\{ e^{V(w^{(n-k)})} \mathbf{1}_{\{V(w_i^{(n-k)}) \geq -c_{20}n^\varepsilon, \forall 0 \leq i \leq n-k, V(w_{n-k}^{(n-k)}) \leq b \log \ell_1\}} \right\} \\ &= \mathbf{E} \mathbf{Q} \left\{ e^{S_{n-k}} \mathbf{1}_{\{S_i \geq -c_{20}n^\varepsilon, \forall 0 \leq i \leq n-k, S_{n-k} \leq b \log \ell_1\}} \right\} \\ &\leq \ell_1^b \mathbf{Q} \left\{ S_i \geq -c_{20}n^\varepsilon, \forall 0 \leq i \leq n-k, S_{n-k} \leq b \log \ell_1 \right\} \\ &\leq \ell_1^{b-(3/2)+\varepsilon+o(1)} \leq \ell_1^{b-(3/2)+2\varepsilon}. \end{aligned}$$

Therefore,

$$\begin{aligned} \Lambda_{k,n,m} &\leq \ell_1^{b-(3/2)+2\varepsilon} \mathbf{E} \left\{ \sum_{|z|=k} \sum_{v \in \mathbb{T}_z^{\text{GW}}, |v|_z=m-k} \mathbf{1}_{\{zv \in \mathbb{L}_m\}} \right\} \\ &= \ell_1^{b-(3/2)+2\varepsilon} \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\} \end{aligned}$$

and, thus,

$$(4.3) \quad \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=0}^{\lfloor n^\varepsilon \rfloor} \Lambda_{k,n,m} \leq \ell_1^{b-(3/2)+2\varepsilon} (\ell_2 - \ell_1 + 1) (\ell_2^\varepsilon + 1) \mathbf{E}(\#F_{\ell_1, \ell_2}).$$

*Second situation:*  $\lfloor n^\varepsilon \rfloor + 1 \leq k \leq \min\{m - \lfloor n^\varepsilon \rfloor, n\}$ . In this situation, since  $V(z) \geq \max\{g_m(k), g_n(k)\} \geq c_{19}n^{\varepsilon/3}$ , we simply have  $V_z(u) \leq b \log \ell_1 - c_{19}n^{\varepsilon/3}$ . Exactly as in the first situation, we get

$$\Lambda_{k,n,m} \leq \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\} \mathbf{E} \left\{ \sum_{|x|=n-k} \mathbf{1}_{\{V(x) \leq b \log \ell_1 - c_{19}n^{\varepsilon/3}\}} \right\}.$$

The second  $\mathbf{E}\{\cdot\}$  on the right-hand side is

$$= \mathbf{E} \mathbf{Q} \left\{ e^{S_{n-k}} \mathbf{1}_{\{S_{n-k} \leq b \log \ell_1 - c_{19}n^{\varepsilon/3}\}} \right\} \leq \ell_1^b e^{-c_{19}n^{\varepsilon/3}}$$

and, thus,

$$(4.4) \quad \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=\lfloor n^\varepsilon \rfloor + 1}^{\min\{m - \lfloor n^\varepsilon \rfloor, n\}} \Lambda_{k,n,m} \leq \ell_1^b e^{-c_{19}\ell_1^{\varepsilon/3}} (\ell_2 - \ell_1 + 1) \ell_2 \mathbf{E}(\#F_{\ell_1, \ell_2}).$$

*Third and last situation:*  $m - \lfloor n^\varepsilon \rfloor + 1 \leq k \leq n$  (this situation may happen only if  $m \leq n + \lfloor n^\varepsilon \rfloor - 1$ ). This time  $V(z) \geq g_m(k) \geq a \log \ell_1$  and, thus,  $V_z(u) \leq (b -$

a)  $\log \ell_1$ ; consequently,

$$\begin{aligned} \Lambda_{k,n,m} &\leq \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\} \mathbf{E} \left\{ \sum_{|x|=n-k} \mathbf{1}_{\{V(x) \leq (b-a) \log \ell_1\}} \right\} \\ &\leq \ell_1^{b-a} \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\}. \end{aligned}$$

Therefore, in case  $m \leq n + \lfloor n^\varepsilon \rfloor - 1$ ,

$$\begin{aligned} \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=m-\lfloor n^\varepsilon \rfloor+1}^n \Lambda_{k,n,m} &\leq \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{n+\lfloor n^\varepsilon \rfloor-1} \ell_2^\varepsilon \ell_1^{b-a} \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\} \\ &\leq \sum_{m=\ell_1}^{\ell_2} \sum_{n=m-2\lfloor m^\varepsilon \rfloor}^m \ell_2^\varepsilon \ell_1^{b-a} \mathbf{E} \left\{ \sum_{|x|=m} \mathbf{1}_{\{x \in \mathbb{L}_m\}} \right\} \\ &\leq 2\ell_2^{2\varepsilon} \ell_1^{b-a} \mathbf{E}(\#F_{\ell_1, \ell_2}). \end{aligned}$$

Combining this with (4.3) and (4.4), and since

$$\mathbf{E}[(\#F_{\ell_1, \ell_2})^2] \leq 2 \sum_{n=\ell_1}^{\ell_2} \sum_{m=n}^{\ell_2} \sum_{k=0}^n \Lambda_{k,n,m},$$

we obtain

$$\begin{aligned} \frac{\mathbf{E}[(\#F_{\ell_1, \ell_2})^2]}{[\mathbf{E}(\#F_{\ell_1, \ell_2})]^2} &\leq (2\ell_1^{b-(3/2)+2\varepsilon} (\ell_2 - \ell_1 + 1)(\ell_2^\varepsilon + 1) \\ &\quad + 2\ell_1^b e^{-c_{19}\ell_1^{\varepsilon/3}} (\ell_2 - \ell_1 + 1)\ell_2 + 4\ell_2^{2\varepsilon} \ell_1^{b-a}) (\mathbf{E}(\#F_{\ell_1, \ell_2}))^{-1}. \end{aligned}$$

Since  $\ell_2 \leq 2\ell_1$ , we have  $2\ell_1^{b-(3/2)+2\varepsilon} (\ell_2^\varepsilon + 1) + 2\ell_1^b e^{-c_{19}\ell_1^{\varepsilon/3}} \ell_2 \leq \ell_1^{b-(3/2)+4\varepsilon}$  for all sufficiently large  $\ell_1$ . On the other hand,  $\mathbf{E}(\#F_{\ell_1, \ell_2}) \geq (\ell_2 - \ell_1 + 1)\ell_1^{a-(3/2)-\varepsilon}$  by (4.2) (for large  $\ell_1$ ). Therefore, when  $\ell_1$  is large, we have

$$\frac{\mathbf{E}[(\#F_{\ell_1, \ell_2})^2]}{[\mathbf{E}(\#F_{\ell_1, \ell_2})]^2} \leq \frac{\ell_1^{b-(3/2)+4\varepsilon} (\ell_2 - \ell_1 + 1) + \ell_1^{b-a+3\varepsilon}}{(\ell_2 - \ell_1 + 1)\ell_1^{a-(3/2)-\varepsilon}}.$$

By the Paley–Zygmund inequality,  $\mathbf{P}\{F_{\ell_1, \ell_2} \neq \emptyset\} \geq \frac{1}{4} \frac{[\mathbf{E}(\#F_{\ell_1, \ell_2})]^2}{\mathbf{E}[(\#F_{\ell_1, \ell_2})^2]}$ ; thus,

$$(4.5) \quad \mathbf{P} \left\{ \min_{\ell_1 \leq |x| \leq \ell_2} V(x) \leq b \log \ell_1 \right\} \geq \frac{1}{4} \frac{(\ell_2 - \ell_1 + 1)\ell_1^{a-(3/2)-\varepsilon}}{\ell_1^{b-(3/2)+4\varepsilon} (\ell_2 - \ell_1 + 1) + \ell_1^{b-a+3\varepsilon}}.$$

Of course, we can make  $a$  close to  $b$ , and  $\varepsilon$  close to 0, to see that, for any  $b \in \mathbb{R}$  and  $\varepsilon > 0$ , all sufficiently large  $\ell_1$  and all  $\ell_2 \in [\ell_1, 2\ell_1] \cap \mathbb{Z}$ ,

$$(4.6) \quad \mathbf{P} \left\{ \min_{\ell_1 \leq |x| \leq \ell_2} V(x) \leq b \log \ell_1 \right\} \geq \frac{\ell_2 - \ell_1 + 1}{\ell_1^\varepsilon (\ell_2 - \ell_1 + 1) + \ell_1^{(3/2)-b+\varepsilon}}.$$

[This is our basic estimate for the minimum of  $V(x)$ . In Section 5 we are going to apply (4.6) to  $\ell_2 := \ell_1$ .]

We now let  $b > \frac{1}{2}$  and take the subsequence  $n_j := 2^j$ ,  $j \geq j_0$  (with a sufficiently large integer  $j_0$ ). By (4.6) (and possibly by changing the value of  $\varepsilon$ ),

$$\mathbf{P}\left\{\min_{n_j \leq |x| \leq n_{j+1}} V(x) \leq b \log n_j\right\} \geq n_j^{-\varepsilon}.$$

Let  $\tau_j := \inf\{k : \#\{u : |u| = k\} \geq n_j^{2\varepsilon}\}$ . Then we have, for  $j \geq j_0$ ,

$$\begin{aligned} &\mathbf{P}\left\{\tau_j < \infty, \min_{\tau_j+n_j \leq |x| \leq \tau_j+n_{j+1}} V(x) > \max_{|y|=\tau_j} V(y) + b \log n_j\right\} \\ &\leq \left(\mathbf{P}\left\{\min_{n_j \leq |x| \leq n_{j+1}} V(x) > b \log n_j\right\}\right)^{\lfloor n_j^{2\varepsilon} \rfloor} \\ &\leq (1 - n_j^{-\varepsilon})^{\lfloor n_j^{2\varepsilon} \rfloor}, \end{aligned}$$

which is summable in  $j$ . By the Borel–Cantelli lemma, almost surely for all large  $j$ , we have either  $\tau_j = \infty$ , or  $\min_{\tau_j+n_j \leq |x| \leq \tau_j+n_{j+1}} V(x) \leq \max_{|y|=\tau_j} V(y) + b \log n_j$ .

By the well-known law of large numbers for the branching random walk (Hammersley [19], Kingman [23] and Biggins [5]), of which (1.5) was a special case, there exists a constant  $c_{21} > 0$  such that  $\frac{1}{n} \max_{|y|=n} V(y) \rightarrow c_{21}$  almost surely upon the system’s survival. In particular, upon survival,  $\max_{|y|=n} V(y) \leq 2c_{21}n$  almost surely for all large  $n$ . Consequently, upon the system’s survival, almost surely for all large  $j$ , we have either  $\tau_j = \infty$ , or  $\min_{\tau_j+n_j \leq |x| \leq \tau_j+n_{j+1}} V(x) < 2c_{21}\tau_j + b \log n_j$ .

Recall that the number of particles in each generation forms a Galton–Watson tree, which is super-critical under assumption (1.3) (because  $m := \mathbf{E}\{\sum_{|u|=1} 1\} > 1$ ). In particular, conditionally on the system’s survival,  $\frac{\#\{u:|u|=k\}}{m^k}$  converges a.s. to a (strictly) positive random variable when  $k \rightarrow \infty$ , which implies  $\tau_j \sim 2\varepsilon \frac{\log n_j}{\log m}$  a.s. ( $j \rightarrow \infty$ ). As a consequence, upon the system’s survival, we have, almost surely for all large  $j$ ,

$$\min_{n_j \leq |x| \leq 2n_{j+1}} V(x) \leq \frac{5\varepsilon c_{21}}{\log m} \log n_j + b \log n_j.$$

Since  $b$  can be as close to  $\frac{1}{2}$  as possible, this readily yields (4.1).

**5. Proof of Theorem 1.6.** Before proving Theorem 1.6, we need three estimates.

The first estimate, stated as Proposition 5.1, was proved by McDiarmid [31] under the additional assumption  $\mathbf{E}\{(\sum_{|u|=1} 1)^2\} < \infty$ .

PROPOSITION 5.1. Assume (1.1), (1.2) and (1.3). There exists  $c_{22} > 0$  such that, for any  $\varepsilon > 0$ , we can find  $c_{23} = c_{23}(\varepsilon) > 0$  satisfying

$$(5.1) \quad \mathbf{E} \left\{ \exp \left( c_{22} \inf_{|x|=n} V(x) \right) \mathbf{1}_{\mathcal{S}_n} \right\} \leq c_{23} n^{(3+\varepsilon)/2c_{22}}, \quad n \geq 1.$$

REMARK. Since  $W_n \geq \exp[-\inf_{|x|=n} V(x)]$ , it follows from (5.1) and Hölder’s inequality that, for any  $0 \leq s < c_{22}$  and  $\varepsilon > 0$ ,

$$(5.2) \quad \mathbf{E} \left\{ \frac{1}{W_n^s} \mathbf{1}_{\mathcal{S}_n} \right\} \leq c_{23}^{s/c_{22}} n^{(3+\varepsilon)/2s}, \quad n \geq 1.$$

This estimate will be useful in the proof of Theorem 1.5 in Section 6.

PROOF OF PROPOSITION 5.1. In the proof we write, for any  $k \geq 0$ ,

$$\underline{V}_k := \inf_{|u|=k} V(u).$$

Taking  $\ell_2 = \ell_1$  in (4.6) gives that, for any  $\varepsilon > 0$  and all sufficiently large  $\ell$  (say,  $\ell \geq \ell_0$ ), we have  $\mathbf{P}\{\underline{V}_\ell \geq \frac{3}{2} \log \ell\} \geq \ell^{-\varepsilon}$ ; thus,  $\mathbf{P}\{\underline{V}_\ell > \frac{3}{2} \log \ell\} \leq 1 - \ell^{-\varepsilon} \leq \exp(-\ell^{-\varepsilon})$ ,  $\forall \ell \geq \ell_0$ . For any  $r \in \mathbb{R}$  and integers  $k \geq 1$  and  $n > \ell \geq \ell_0$ , we have

$$\begin{aligned} & \mathbf{P}\{\underline{V}_n > \frac{3}{2} \log \ell + r\} \\ & \leq \mathbf{P}\{\#\{u : |u| = n - \ell, V(u) \leq r\} < k\} + (\mathbf{P}\{\underline{V}_\ell > \frac{3}{2} \log \ell\})^k \\ & \leq \mathbf{P}\{\#\{u : |u| = n - \ell, V(u) \leq r\} < k\} + \exp(-\ell^{-\varepsilon} k). \end{aligned}$$

By Lemma 1 of McDiarmid [31], there exist  $c_{24} > 0$ ,  $c_{25} > 0$  and  $c_{26} > 0$  such that, for any  $j \geq 1$ ,  $\mathbf{P}\{\#\{u : |u| = j, V(u) \leq c_{24}j\} \leq e^{c_{25}j}\} \leq q + e^{-c_{26}j}$ ,  $q$  being as before the probability of extinction. We choose  $j := \lfloor \frac{r}{c_{24}} \rfloor$  and  $\ell := n - \lfloor \frac{r}{c_{24}} \rfloor$  to see that, for all  $n \geq \ell_0$  and all  $0 \leq r \leq c_{24}(n - \ell_0)$ ,

$$\mathbf{P}\{\underline{V}_n > \frac{3}{2} \log n + r\} \leq q + e^{-c_{26}\lfloor r/c_{24} \rfloor} + \exp(-n^{-\varepsilon} \lfloor e^{c_{25}\lfloor r/c_{24} \rfloor} \rfloor).$$

Noting that  $\{\underline{V}_n > \frac{3}{2} \log n + r\} \cap \mathcal{S}_n^c = \mathcal{S}_n^c$  and that  $\mathbf{P}\{\mathcal{S}_n^c\} \geq q - e^{-c_{10}n}$  [see (3.3)], we obtain, for  $0 \leq r \leq c_{24}(n - \ell_0)$ ,

$$(5.3) \quad \begin{aligned} & \mathbf{P}\{\underline{V}_n > \frac{3}{2} \log n + r, \mathcal{S}_n\} \\ & \leq e^{-c_{10}n} + e^{-c_{26}\lfloor r/c_{24} \rfloor} + \exp(-n^{-\varepsilon} \lfloor e^{c_{25}\lfloor r/c_{24} \rfloor} \rfloor). \end{aligned}$$

This implies that, for any  $0 < c_{27} < \min\{\frac{c_{26}}{c_{24}}, \frac{2c_{10}}{c_{24}}\}$ , there exists a constant  $c_{28} > 0$  such that  $\mathbf{E}\{e^{c_{27}\underline{V}_n} \mathbf{1}_{\{\frac{3}{2} \log n < \underline{V}_n \leq c_{24}/2n\} \cap \mathcal{S}_n}\} \leq c_{28} n^{c_{29}}$ , with  $c_{29} := (\frac{3}{2} + \frac{c_{24}}{c_{25}} \varepsilon) c_{27}$ . Therefore,

$$(5.4) \quad \mathbf{E}\{e^{c_{27}\underline{V}_n} \mathbf{1}_{\{\underline{V}_n \leq c_{24}/2n\} \cap \mathcal{S}_n}\} \leq c_{30} n^{c_{29}}, \quad n \geq 1,$$

where  $c_{30} := c_{28} + 1$ .

On the other hand, letting  $\delta_- > 0$  be as in (1.2), we have  $e^{\delta_- V_n} \mathbf{1}_{\mathcal{S}_n} \leq \sum_{|u|=n} e^{\delta_- V(u)}$ . Since  $\psi(-\delta_-) := \log \mathbf{E}\{\sum_{|u|=n} e^{\delta_- V(u)}\} < \infty$  by (1.2), we can choose and fix  $c_{31} > 0$  sufficiently large (in particular,  $c_{31} > \frac{c_{24}}{2}$ ) such that, for any  $x \geq c_{31}$ ,

$$\mathbf{P}\{\underline{V}_n > xn, \mathcal{S}_n\} \leq e^{-\delta_- xn + \psi(-\delta_-)n} \leq e^{-\delta_- xn/2}, \quad \forall n \geq 1.$$

Therefore, for any  $c_{32} < \frac{\delta_-}{2}$ , we have

$$(5.5) \quad \sup_{n \geq 1} \mathbf{E}\{e^{c_{32} \underline{V}_n} \mathbf{1}_{\{\underline{V}_n > c_{31}n\} \cap \mathcal{S}_n}\} < \infty.$$

Finally, (5.3) also implies that, for  $n \geq \ell_0$ ,

$$\begin{aligned} \mathbf{P}\left\{\underline{V}_n > \frac{c_{24}}{2}n, \mathcal{S}_n\right\} &\leq e^{-c_{10}n} + e^{-c_{26}\lfloor n/2-3/(2c_{24})\log n \rfloor} \\ &\quad + \exp(-n^{-\varepsilon} \lfloor e^{c_{25}\lfloor n/2-3/(2c_{24})\log n \rfloor} \rfloor). \end{aligned}$$

Therefore, for any  $c_{33} < \min\{\frac{c_{10}}{c_{31}}, \frac{c_{26}}{2c_{31}}\}$ ,

$$\sup_{n \geq 1} \mathbf{E}\{e^{c_{33} \underline{V}_n} \mathbf{1}_{\{c_{24}/2n < \underline{V}_n \leq c_{31}n\} \cap \mathcal{S}_n}\} < \infty,$$

which, combined with (5.4) and (5.5), completes the proof of Proposition 5.1, with  $c_{22} := \min\{c_{27}, c_{32}, c_{33}\}$ .  $\square$

LEMMA 5.2. *Let  $X_1, X_2, \dots, X_N$  be independent nonnegative random variables, and let  $T_N := \sum_{i=1}^N X_i$ . For any nonincreasing function  $F : (0, \infty) \rightarrow \mathbb{R}_+$ , we have*

$$\mathbf{E}\{F(T_N) \mathbf{1}_{\{T_N > 0\}}\} \leq \max_{1 \leq i \leq N} \mathbf{E}\{F(X_i) | X_i > 0\}.$$

Moreover,

$$\mathbf{E}\{F(T_N) \mathbf{1}_{\{T_N > 0\}}\} \leq \sum_{i=1}^N b^{i-1} \mathbf{E}\{F(X_i) \mathbf{1}_{\{X_i > 0\}}\},$$

where  $b := \max_{1 \leq i \leq N} \mathbf{P}\{X_i = 0\}$ .

PROOF. Let  $\tau := \min\{i \geq 1 : X_i > 0\}$  (with  $\min \emptyset := \infty$ ). Then  $\mathbf{E}\{F(T_N) \times \mathbf{1}_{\{T_N > 0\}}\} = \sum_{i=1}^N \mathbf{E}\{F(T_N) \mathbf{1}_{\{\tau=i\}}\}$ . Since  $F$  is nonincreasing, we have  $F(T_N) \times \mathbf{1}_{\{\tau=i\}} \leq F(X_i) \mathbf{1}_{\{\tau=i\}} = F(X_i) \mathbf{1}_{\{X_i > 0\}} \mathbf{1}_{\{X_j=0, \forall j < i\}}$ . By independence, this leads to

$$\mathbf{E}\{F(T_N) \mathbf{1}_{\{T_N > 0\}}\} \leq \sum_{i=1}^N \mathbf{E}\{F(X_i) \mathbf{1}_{\{X_i > 0\}}\} \mathbf{P}\{X_j = 0, \forall j < i\}.$$

This yields immediately the second inequality of the lemma, since  $\mathbf{P}\{X_j = 0, \forall j < i\} \leq b^{i-1}$ .

To prove the first inequality of the lemma, we observe that  $\mathbf{E}\{F(X_i)\mathbf{1}_{\{X_i>0\}}\} \leq \mathbf{P}\{X_i > 0\} \max_{1 \leq k \leq N} \mathbf{E}\{F(X_k)|X_k > 0\}$ . Therefore,

$$\mathbf{E}\{F(T_N)\mathbf{1}_{\{T_N>0\}}\} \leq \max_{1 \leq k \leq N} \mathbf{E}\{F(X_k)|X_k > 0\} \sum_{i=1}^N \mathbf{P}\{X_i > 0\} \mathbf{P}\{X_j = 0, \forall j < i\}.$$

The  $\sum_{i=1}^N \dots$  expression on the right-hand side is  $= \sum_{i=1}^N \mathbf{P}\{X_i > 0, X_j = 0, \forall j < i\} = \sum_{i=1}^N \mathbf{P}\{\tau = i\} = \mathbf{P}\{T_N > 0\} \leq 1$ . This yields the first inequality of the lemma.  $\square$

To state our third estimate, let  $\underline{w}^{(n)} \in \llbracket e, w_n^{(n)} \rrbracket$  be a vertex such that

$$(5.6) \quad V(\underline{w}^{(n)}) = \min_{u \in \llbracket e, w_n^{(n)} \rrbracket} V(u).$$

[If there are several such vertices, we choose, say, the oldest.] The following estimate gives a (stochastic) lower bound for  $\frac{1}{W_{n,\beta}}$  under  $\mathbf{Q}$  outside a “small” set. We recall that  $W_{n,\beta} > 0$ ,  $\mathbf{Q}$ -almost surely (but not necessarily  $\mathbf{P}$ -almost surely).

LEMMA 5.3. *Assume (1.1), (1.2) and (1.3). For any  $K > 0$ , there exist  $\theta > 0$  and  $n_0 < \infty$  such that, for any  $n \geq n_0$ , any  $\beta > 0$ , and any nondecreasing function  $G : (0, \infty) \rightarrow (0, \infty)$ ,*

$$(5.7) \quad \mathbf{E}_{\mathbf{Q}} \left\{ G \left( \frac{e^{-\beta V(\underline{w}^{(n)})}}{W_{n,\beta}} \right) \mathbf{1}_{E_n} \right\} \leq \frac{1}{1 - q} \max_{0 \leq k < n} \mathbf{E} \left\{ G \left( \frac{n^{\theta\beta}}{W_{k,\beta}} \right) \mathbf{1}_{\mathcal{S}_k} \right\},$$

where  $E_n$  is a measurable event such that

$$\mathbf{Q}\{E_n\} \geq 1 - \frac{1}{n^K}, \quad n \geq n_0.$$

PROOF. Recall from (2.8) that  $\mathcal{S}_k^{(n)}$  is the set of the brothers of  $w_k^{(n)}$ . For any pair  $0 \leq k < n$ , we say that *the level  $k$  is  $n$ -good* if

$$\mathcal{S}_k^{(n)} \neq \emptyset \quad \text{and} \quad \mathbb{T}_u^{\text{GW}}$$
 survives at least  $n - k$  generations,  $\forall u \in \mathcal{S}_k^{(n)}$ ,

where  $\mathbb{T}_u^{\text{GW}}$  is the shifted Galton–Watson subtree generated by  $u$  [see (2.3)]. By  $\mathbb{T}_u^{\text{GW}}$  surviving at least  $n - k$  generations, we mean that there exists  $v \in \mathbb{T}_u^{\text{GW}}$  such that  $|v|_u = n - k$  [see (2.4) for notation].

In words,  $k$  is  $n$ -good means any subtree generated by any of the brothers of  $w_k^{(n)}$  has offspring for at least  $n - k$  generations.

Let  $\mathcal{G}_n$  be the sigma-field defined in (2.9). By Proposition 2.1,

$$\mathbf{Q}\{k \text{ is } n\text{-good} | \mathcal{G}_n\} = \mathbf{1}_{\{\mathcal{S}_k^{(n)} \neq \emptyset\}} (\mathbf{P}\{\mathcal{S}_{n-k}\})^{\#\mathcal{S}_k^{(n)}},$$

where  $\mathcal{S}_n$  denotes the system's survival after  $n$  generations [see (3.2)]. Since  $\mathbf{P}\{\mathcal{S}_{n-k} \geq \mathbf{P}\{\mathcal{S}\}\} = 1 - q$ , whereas  $\#\mathcal{S}_k^{(n)}$  and  $\#\mathcal{S}_1^{(1)}$  have the same distribution under  $\mathbf{Q}$  (Proposition 2.1), we have

$$\mathbf{Q}\{k \text{ is } n\text{-good}\} \geq \mathbf{E}_{\mathbf{Q}}\{\mathbf{1}_{\{\#\mathcal{S}_1^{(1)} \geq 1\}}(1 - q)^{\#\mathcal{S}_1^{(1)}}\} = c_{34} \in (0, 1).$$

As a consequence, for all  $1 \leq \ell < n$ , by Proposition 2.1 again,

$$\begin{aligned} \mathbf{Q}\left\{\bigcup_{k=1}^n \bigcap_{j:1 \leq j \leq n, |j-k| \leq \ell} \{j \text{ is not } n\text{-good}\}\right\} &\leq \sum_{k=1}^n \prod_{j:1 \leq j \leq n, |j-k| \leq \ell} \mathbf{Q}\{j \text{ is not } n\text{-good}\} \\ &\leq n(1 - c_{34})^{\ell+1}, \end{aligned}$$

which is bounded by  $ne^{-c_{34}(\ell+1)}$  (using the inequality  $1 - x \leq e^{-x}$ , for  $x \geq 0$ ). Let  $K > 0$ . We take  $\ell = \ell(n) := \lfloor c_{35} \log n \rfloor$  with  $c_{35} := \frac{K+2}{c_{34}}$ . Let  $c_{36} := \frac{K+2}{c_6}$  [where  $c_6$  is as in (2.17)] and  $c_{37} := \max\{\frac{K+2}{c_3}, c_{35}\}$  [ $c_3$  being the constant in (2.12)]. Let

$$(5.8) \quad E_n^{(1)} := \bigcap_{k=1}^n \bigcup_{j:1 \leq j \leq n, |j-k| \leq \lfloor c_{35} \log n \rfloor} \{j \text{ is } n\text{-good}\},$$

$$(5.9) \quad E_n^{(2)} := \left\{ \max_{1 \leq j \leq n} \sup_{u \in \mathcal{S}_j^{(n)}} |V(u) - V(w_{j-1}^{(n)})| \leq c_{36} \log n \right\},$$

$$(5.10) \quad E_n^{(3)} := \left\{ \max_{0 \leq j, k \leq n, |j-k| \leq c_{35} \log n} |V(w_j^{(n)}) - V(w_k^{(n)})| \leq c_{37} \log n \right\}.$$

We have

$$\mathbf{Q}\{E_n^{(1)}\} \geq 1 - ne^{-c_{34}c_{35} \log n} = 1 - \frac{1}{n^{K+1}}.$$

On the other hand, by Corollary 2.2,

$$\mathbf{Q}\{(E_n^{(2)})^c\} \leq n\mathbf{Q}\left\{\sup_{u \in \mathcal{S}_1^{(1)}} |V(u)| > c_{36} \log n\right\} \leq n\mathbf{Q}\left\{\sup_{|u|=1} |V(u)| > c_{36} \log n\right\}.$$

Applying (2.17) yields that

$$\mathbf{Q}\{E_n^{(2)}\} \geq 1 - c_5 n^{-(c_{36}c_6 - 1)} = 1 - \frac{c_5}{n^{K+1}}.$$

To estimate  $\mathbf{Q}\{E_n^{(3)}\}$ , we note that, by Corollary 2.2,

$$\mathbf{Q}\{(E_n^{(3)})^c\} = \mathbf{Q}\left\{\max_{0 \leq j, k \leq n, |j-k| \leq c_{35} \log n} |S_j - S_k| > c_{37} \log n\right\},$$

which, in view of (2.15), is bounded by  $2c_{35}n^{-(c_3c_{37}-1)} \log n$ . Consequently, if

$$(5.11) \quad E_n := E_n^{(1)} \cap E_n^{(2)} \cap E_n^{(3)},$$

then  $\mathbf{Q}\{E_n\} \geq 1 - \frac{1}{n^K}$  for all large  $n$ .

It remains to check (5.7). By definition,

$$\begin{aligned}
 W_{n,\beta} &= \sum_{j=1}^n \sum_{u \in \mathcal{I}_j^{(n)}} e^{-\beta V(u)} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-\beta V_u(x)} + e^{-\beta V(w^{(n)})} \\
 (5.12) \quad &\geq \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} e^{-\beta V(u)} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-\beta V_u(x)}
 \end{aligned}$$

for any  $\mathcal{L} \subset \{1, 2, \dots, n\}$ . We choose  $\mathcal{L} := \{1 \leq j \leq n : |j - |\underline{w}^{(n)}|| < c_{35} \log n\}$ .

On the event  $E_n$ , for  $u \in \mathcal{I}_j^{(n)}$  with some  $j \in \mathcal{L}$ , we have  $V(u) \leq V(\underline{w}^{(n)}) + (c_{36} + c_{37}) \log n$ . Writing  $\theta := c_{36} + c_{37}$ , this leads to  $W_{n,\beta} \geq n^{-\theta\beta} e^{-\beta V(\underline{w}^{(n)})} \times \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} \xi_u$ , where

$$\xi_u := \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-\beta V_u(x)}.$$

Since  $\sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} \xi_u > 0$  on  $E_n$ , we arrive at

$$\frac{e^{-\beta V(\underline{w}^{(n)})}}{W_{n,\beta}} \mathbf{1}_{E_n} \leq \frac{n^{\theta\beta}}{\sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} \xi_u} \mathbf{1}_{\{\sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} \xi_u > 0\}}.$$

Let  $\mathcal{G}_n$  be the sigma-field in (2.9). We observe that  $\mathcal{L}$  and  $\mathcal{I}_j^{(n)}$  are  $\mathcal{G}_n$ -measurable. Moreover, an application of Proposition 2.1 tells us that under  $\mathbf{Q}$ , conditionally on  $\mathcal{G}_n$ , the random variables  $\xi_u, u \in \mathcal{I}_j^{(n)}, j \in \mathcal{L}$ , are independent, and are distributed as  $W_{n-j,\beta}$  under  $\mathbf{P}$ . We are thus entitled to apply Lemma 5.2 to see that, if  $G$  is nondecreasing,

$$\begin{aligned}
 \mathbf{E}_{\mathbf{Q}} \left\{ G \left( \frac{e^{-\beta V(\underline{w}^{(n)})}}{W_{n,\beta}} \right) \mathbf{1}_{E_n} \mid \mathcal{G}_n \right\} &\leq \max_{j \in \mathcal{L}} \mathbf{E} \left\{ G \left( \frac{n^{\theta\beta}}{W_{n-j,\beta}} \right) \mid W_{n-j,\beta} > 0 \right\} \\
 &\leq \max_{0 \leq k < n} \mathbf{E} \left\{ G \left( \frac{n^{\theta\beta}}{W_{k,\beta}} \right) \mid W_{k,\beta} > 0 \right\}.
 \end{aligned}$$

Since  $\mathbf{P}\{W_{k,\beta} > 0\} = \mathbf{P}\{\mathcal{S}_k\} \geq \mathbf{P}\{\mathcal{S}\} = 1 - q$ , this yields Lemma 5.3.  $\square$

We are now ready for the proof of Theorem 1.6. For the sake of clarity, the upper and lower bounds are proved in distinct parts. Let us start with the upper bound.

**PROOF OF THEOREM 1.6. The upper bound.** We assume (1.1), (1.2) and (1.3), and fix  $\beta > 1$ .

For any  $Z \geq 0$  which is  $\mathcal{F}_n$ -measurable, we have  $\mathbf{E}\{W_{n,\beta}Z\} = \mathbf{E}_{\mathbf{Q}}\{\sum_{|u|=n} \frac{e^{-\beta V(u)}}{W_n} Z\} = \mathbf{E}_{\mathbf{Q}}\{\sum_{|u|=n} \mathbf{1}_{\{w^{(n)}=u\}} e^{-(\beta-1)V(u)} Z\}$  and, thus,

$$(5.13) \quad \mathbf{E}\{W_{n,\beta}Z\} = \mathbf{E}_{\mathbf{Q}}\{e^{-(\beta-1)V(w^{(n)})} Z\}.$$



Let  $s \in (\frac{\beta-1}{\beta}, 1)$ , and  $\lambda > 0$ . (We will choose  $\lambda = \frac{3}{2}$ .) Then

$$\begin{aligned} \mathbf{E}\{W_{n,\beta}^{1-s}\} &\leq n^{-(1-s)\beta\lambda} + \mathbf{E}\{W_{n,\beta}^{1-s} \mathbf{1}_{\{W_{n,\beta} > n^{-\beta\lambda}\}}\} \\ &= n^{-(1-s)\beta\lambda} + \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{\{W_{n,\beta} > n^{-\beta\lambda}\}}\right\}. \end{aligned}$$

Since  $e^{-\beta V(w_n^{(n)})} \leq W_{n,\beta}$ , we have  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq \frac{1}{W_{n,\beta}^{s-(\beta-1)/\beta}}$ ; thus, on the event

$\{W_{n,\beta} > n^{-\beta\lambda}\}$ , we have  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq n^{[\beta s - (\beta-1)]\lambda}$ .

Let  $K := [\beta s - (\beta - 1)]\lambda + (1 - s)\beta\lambda$ , and let  $E_n$  be the event in Lemma 5.3. Since  $\mathbf{Q}(E_n^c) \leq n^{-K}$  for all sufficiently large  $n$  (see Lemma 5.3), we obtain, for large  $n$ ,

$$\begin{aligned} \mathbf{E}\{W_{n,\beta}^{1-s}\} &\leq n^{-(1-s)\beta\lambda} + n^{[\beta s - (\beta-1)]\lambda - K} \\ (5.14) \quad &+ \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{\{W_{n,\beta} > n^{-\beta\lambda}\} \cap E_n}\right\} \\ &= 2n^{-(1-s)\beta\lambda} + \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{\{W_{n,\beta} > n^{-\beta\lambda}\} \cap E_n}\right\}. \end{aligned}$$

We now estimate the expectation expression  $\mathbf{E}\mathbf{Q}\{\cdot\}$  on the right-hand side. Let  $a > 0$  and  $\varrho > b > 0$  be constants such that  $(\beta - 1)a > s\beta\lambda + \frac{3}{2}$  and  $[\beta s - (\beta - 1)]b > \frac{3}{2}$ . (The choice of  $\varrho$  will be made precise later on.) We recall that  $\underline{w}_n^{(n)} \in \llbracket e, w_n^{(n)} \rrbracket$  satisfies  $V(\underline{w}_n^{(n)}) = \min_{u \in \llbracket e, w_n^{(n)} \rrbracket} V(u)$ , and consider the following events:

$$\begin{aligned} E_{1,n} &:= \{V(w_n^{(n)}) > a \log n\} \cup \{V(w_n^{(n)}) \leq -b \log n\}, \\ E_{2,n} &:= \{V(\underline{w}_n^{(n)}) < -\varrho \log n, V(w_n^{(n)}) > -b \log n\}, \\ E_{3,n} &:= \{V(\underline{w}_n^{(n)}) \geq -\varrho \log n, -b \log n < V(w_n^{(n)}) \leq a \log n\}. \end{aligned}$$

Clearly,  $\mathbf{Q}(\bigcup_{i=1}^3 E_{i,n}) = 1$ .

On the event  $E_{1,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}$ , we have either  $V(w_n^{(n)}) > a \log n$ , in which case  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq n^{s\beta\lambda - (\beta-1)a}$ , or  $V(w_n^{(n)}) \leq -b \log n$ , in which case we use the trivial inequality  $W_{n,\beta} \geq e^{-\beta V(w_n^{(n)})}$  to see that  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq e^{[\beta s - (\beta-1)]V(w_n^{(n)})} \leq n^{-[\beta s - (\beta-1)]b}$  (recalling that  $\beta s > \beta - 1$ ). Since  $s\beta\lambda - (\beta - 1)a < -\frac{3}{2}$  and  $[\beta s - (\beta - 1)]b > \frac{3}{2}$ , we obtain

$$(5.15) \quad \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{E_{1,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}}\right\} \leq n^{-3/2}.$$

We now study the integral on  $E_{2,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\} \cap E_n$ . Since  $s > 0$ , we can choose  $s_1 > 0$  and  $0 < s_2 \leq \frac{c_{22}}{\beta}$  [where  $c_{22}$  is the constant in (5.1)] such that  $s = s_1 + s_2$ . We have, on  $E_{2,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}$ ,

$$\begin{aligned} \frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} &= \frac{e^{\beta s_2 V(w_n^{(n)}) - (\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^{s_1}} \frac{e^{-\beta s_2 V(w_n^{(n)})}}{W_{n,\beta}^{s_2}} \\ &\leq n^{-\beta s_2 \varrho + (\beta-1)b + \beta \lambda s_1} \frac{e^{-\beta s_2 V(w_n^{(n)})}}{W_{n,\beta}^{s_2}}. \end{aligned}$$

Therefore, by an application of Lemma 5.3 to  $G(x) := x^{s_2}$ ,  $x > 0$ , we obtain, for all sufficiently large  $n$ ,

$$\begin{aligned} \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{E_{2,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\} \cap E_n}\right\} \\ \leq \frac{n^{-\beta s_2 \varrho + (\beta-1)b + \beta \lambda s_1}}{1 - q} \max_{0 \leq k < n} \mathbf{E}\left\{\frac{n^{s_2 \theta \beta}}{W_{k,\beta}^{s_2}} \mathbf{1}_{\mathcal{I}_k}\right\}. \end{aligned}$$

By definition,  $\frac{1}{W_{k,\beta}^{s_2}} \leq \exp(\beta s_2 \inf_{|x|=k} V(x))$ ; thus, by (5.1),  $\mathbf{E}\left\{\frac{n^{s_2 \theta \beta}}{W_{k,\beta}^{s_2}} \mathbf{1}_{\mathcal{I}_k}\right\} \leq c_{23}^{\beta s_2 / c_{22}} n^{s_2 \theta \beta + (3+\varepsilon)/2 \beta s_2}$  for all  $0 \leq k < n$ . We choose (and fix) the constant  $\varrho$  so large that  $-\beta s_2 \varrho + (\beta - 1)b + \beta \lambda s_1 + s_2 \theta \beta + \frac{3+\varepsilon}{2} \beta s_2 < -\frac{3}{2}$ . Therefore, for all large  $n$ ,

$$(5.16) \quad \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{E_{2,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\} \cap E_n}\right\} \leq n^{-3/2}.$$

We make a partition of  $E_{3,n}$ : let  $M \geq 2$  be an integer, and let  $a_i := -b + \frac{i(a+b)}{M}$ ,  $0 \leq i \leq M$ . By definition,

$$\begin{aligned} E_{3,n} &= \bigcup_{i=0}^{M-1} \{V(\underline{w}^{(n)}) \geq -\varrho \log n, a_i \log n < V(w_n^{(n)}) \leq a_{i+1} \log n\} \\ &=: \bigcup_{i=0}^{M-1} E_{3,n,i}. \end{aligned}$$

Let  $0 \leq i \leq M - 1$ . There are two possible situations. First situation:  $a_i \leq \lambda$ . In this case, we argue that, on the event  $E_{3,n,i}$ , we have  $W_{n,\beta} \geq e^{-\beta V(w_n^{(n)})} \geq n^{-\beta a_{i+1}}$  and  $e^{-(\beta-1)V(w_n^{(n)})} \leq n^{-(\beta-1)a_i}$ , thus,  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq n^{\beta s a_{i+1} - (\beta-1)a_i} = n^{\beta s a_i - (\beta-1)a_i + \beta s(a+b)/M} \leq n^{[\beta s - (\beta-1)]\lambda + \beta s(a+b)/M}$ . Accordingly, in this situation,

$$\mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \mathbf{1}_{E_{3,n,i}}\right\} \leq n^{[\beta s - (\beta-1)]\lambda + \beta s(a+b)/M} \mathbf{Q}(E_{3,n,i}).$$

Second (and last) situation:  $a_i > \lambda$ . We have, on  $E_{3,n,i} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}$ ,  $\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s} \leq n^{\beta\lambda s - (\beta-1)a_i} \leq n^{[\beta s - (\beta-1)]\lambda}$ ; thus, in this situation,

$$\mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s}\mathbf{1}_{E_{3,n,i} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}}\right\} \leq n^{[\beta s - (\beta-1)]\lambda}\mathbf{Q}(E_{3,n,i}).$$

We have therefore proved that

$$\begin{aligned} &\mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s}\mathbf{1}_{E_{3,n} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}}\right\} \\ &= \sum_{i=0}^{M-1} \mathbf{E}\mathbf{Q}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s}\mathbf{1}_{E_{3,n,i} \cap \{W_{n,\beta} > n^{-\beta\lambda}\}}\right\} \\ &\leq n^{[\beta s - (\beta-1)]\lambda + \beta s(a+b)/M}\mathbf{Q}(E_{3,n}). \end{aligned}$$

By Corollary 2.2,  $\mathbf{Q}(E_{3,n}) = \mathbf{P}\{\min_{0 \leq k \leq n} S_k \geq -\varrho \log n, -b \log n \leq S_n \leq a \times \log n\} = n^{-(3/2)+o(1)}$ . Combining this with (5.14), (5.15) and (5.16) yields

$$\mathbf{E}\{W_{n,\beta}^{1-s}\} \leq 2n^{-(1-s)\beta\lambda} + 2n^{-3/2} + n^{[\beta s - (\beta-1)]\lambda + \beta s(a+b)/M - (3/2)+o(1)}.$$

We choose  $\lambda := \frac{3}{2}$ . Since  $M$  can be as large as possible, this yields the upper bound in Theorem 1.6 by posing  $r := 1 - s$ .  $\square$

**PROOF OF THEOREM 1.6. The lower bound.** Assume (1.1), (1.2) and (1.3). Let  $\beta > 1$  and  $s \in (1 - \frac{1}{\beta}, 1)$ . By means of (5.12) and the elementary inequality  $(a + b)^{1-s} \leq a^{1-s} + b^{1-s}$  (for  $a \geq 0$  and  $b \geq 0$ ), we have

$$\begin{aligned} W_{n,\beta}^{1-s} &\leq \sum_{j=1}^n \sum_{u \in \mathcal{I}_j^{(n)}} e^{-(1-s)\beta V(u)} \left( \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-\beta V_u(x)} \right)^{1-s} + e^{-(1-s)\beta V(w_n^{(n)})} \\ &= \sum_{j=1}^n e^{-(1-s)\beta V(w_{j-1}^{(n)})} \sum_{u \in \mathcal{I}_j^{(n)}} e^{-(1-s)\beta[V(u) - V(w_{j-1}^{(n)})]} \\ &\quad \times \left( \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-\beta V_u(x)} \right)^{1-s} \\ &\quad + e^{-(1-s)\beta V(w_n^{(n)})}. \end{aligned}$$

Let  $\mathcal{G}_n$  be the sigma-field defined in (2.9), and let

$$\Xi_j = \Xi_j(n, s, \beta) := \sum_{u \in \mathcal{I}_j^{(n)}} e^{-(1-s)\beta[V(u) - V(w_{j-1}^{(n)})]}, \quad 1 \leq j \leq n.$$

Since  $V(w_j^{(n)})$  and  $\mathcal{F}_j^{(n)}$ , for  $1 \leq j \leq n$ , are  $\mathcal{G}_n$ -measurable, it follows from Proposition 2.1 that

$$\mathbf{E}_{\mathbf{Q}}\{W_{n,\beta}^{1-s} | \mathcal{G}_n\} \leq \sum_{j=1}^n e^{-(1-s)\beta V(w_{j-1}^{(n)})} \Xi_j \mathbf{E}\{W_{n-j,\beta}^{1-s}\} + e^{-(1-s)\beta V(w_n^{(n)})}.$$

Let  $\varepsilon > 0$  be small, and let  $r := \frac{3}{2}(1-s)\beta - \varepsilon$ . By means of the already proved upper bound for  $\mathbf{E}(W_{n,\beta}^{1-s})$ , this leads to, with  $c_{38} \geq 1$ ,

$$(5.17) \quad \begin{aligned} &\mathbf{E}_{\mathbf{Q}}\{W_{n,\beta}^{1-s} | \mathcal{G}_n\} \\ &\leq c_{38} \sum_{j=1}^n e^{-(1-s)\beta V(w_{j-1}^{(n)})} (n-j+1)^{-r} \Xi_j + e^{-(1-s)\beta V(w_n^{(n)})}. \end{aligned}$$

Since  $\mathbf{E}(W_{n,\beta}^{1-s}) = \mathbf{E}_{\mathbf{Q}}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{W_{n,\beta}^s}\right\}$  [see (5.13)], we have, by Jensen’s inequality [noticing that  $V(w_n^{(n)})$  is  $\mathcal{G}_n$ -measurable],

$$\mathbf{E}(W_{n,\beta}^{1-s}) \geq \mathbf{E}_{\mathbf{Q}}\left\{\frac{e^{-(\beta-1)V(w_n^{(n)})}}{\{\mathbf{E}_{\mathbf{Q}}(W_{n,\beta}^{1-s} | \mathcal{G}_n)\}^{s/(1-s)}}\right\},$$

which, in view of (5.17), yields

$$\begin{aligned} \mathbf{E}(W_{n,\beta}^{1-s}) &\geq \frac{1}{c_{38}^{s/(1-s)}} \\ &\times \mathbf{E}_{\mathbf{Q}}\left\{e^{-(\beta-1)V(w_n^{(n)})}\right. \\ &\quad \times \left.\left(\left(\sum_{j=1}^n e^{-(1-s)\beta V(w_{j-1}^{(n)})} (n-j+1)^{-r} \Xi_j\right.\right.\right. \\ &\quad \left.\left.\left.+ e^{-(1-s)\beta V(w_n^{(n)})}\right)^{s/(1-s)}\right)^{-1}\right\}. \end{aligned}$$

By Proposition 2.1, if  $(S_j - S_{j-1}, \xi_j)$ , for  $j \geq 1$  (with  $S_0 := 0$ ), are i.i.d. random variables under  $\mathbf{Q}$  and distributed as  $(V(w_1^{(1)}), \sum_{u \in \mathcal{F}_1^{(1)}} e^{-(1-s)\beta V(u)})$ , then the  $\mathbf{E}_{\mathbf{Q}}\{\cdot\}$  expression on the right-hand side is

$$\begin{aligned} &= \mathbf{E}_{\mathbf{Q}}\left\{\frac{e^{-(\beta-1)S_n}}{\{\sum_{j=1}^n (n-j+1)^{-r} e^{-(1-s)\beta S_{j-1}} \xi_j + e^{-(1-s)\beta S_n}\}^{s/(1-s)}}\right\} \\ &= \mathbf{E}_{\mathbf{Q}}\left\{\frac{e^{[\beta s - (\beta-1)]\tilde{S}_n}}{\{\sum_{k=1}^n k^{-r} e^{(1-s)\beta \tilde{S}_k} \tilde{\xi}_k + 1\}^{s/(1-s)}}\right\}, \end{aligned}$$

where

$$\tilde{S}_\ell := S_n - S_{n-\ell}, \quad \tilde{\xi}_\ell := \xi_{n+1-\ell}, \quad 1 \leq \ell \leq n.$$

Consequently,

$$\mathbf{E}(W_{n,\beta}^{1-s}) \geq \frac{1}{c_{38}^{s/(1-s)}} \mathbf{E} \mathbf{Q} \left\{ \frac{e^{[\beta s - (\beta - 1)]\tilde{S}_n}}{\left\{ \sum_{k=1}^n k^{-r} e^{(1-s)\beta \tilde{S}_k \tilde{\xi}_k} + 1 \right\}^{s/(1-s)}} \right\}.$$

Let  $c_{39} > 0$  be a constant, and define

$$E_{n,1}^{\tilde{S}} := \bigcap_{k=1}^{\lfloor n^\varepsilon \rfloor - 1} \{ \tilde{S}_k \leq -c_{39} k^{1/3} \} \cap \{ -2n^{\varepsilon/2} \leq \tilde{S}_{\lfloor n^\varepsilon \rfloor} \leq -n^{\varepsilon/2} \},$$

$$E_{n,2}^{\tilde{S}} := \bigcap_{k=\lfloor n^\varepsilon \rfloor + 1}^{n - \lfloor n^\varepsilon \rfloor - 1} \{ \tilde{S}_k \leq -[k^{1/3} \wedge (n - k)^{1/3}] \} \cap \{ -2n^{\varepsilon/2} \leq \tilde{S}_{n - \lfloor n^\varepsilon \rfloor} \leq -n^{\varepsilon/2} \},$$

$$E_{n,3}^{\tilde{S}} := \bigcap_{k=n - \lfloor n^\varepsilon \rfloor + 1}^{n-1} \left\{ \tilde{S}_k \leq \frac{3}{2} \log n \right\} \cap \left\{ \frac{3 - \varepsilon}{2} \log n \leq \tilde{S}_n \leq \frac{3}{2} \log n \right\}.$$

Let  $\rho := \rho((1 - s)\beta)$  be the constant in Corollary 2.4, and let

$$E_{n,1}^{\tilde{\xi}} := \bigcap_{k=1}^{\lfloor n^\varepsilon \rfloor} \{ \tilde{\xi}_k \leq n^{2\varepsilon/\rho} \},$$

$$E_{n,2}^{\tilde{\xi}} := \bigcap_{k=\lfloor n^\varepsilon \rfloor + 1}^{n - \lfloor n^\varepsilon \rfloor} \{ \tilde{\xi}_k \leq e^{n^{\varepsilon/4}} \},$$

$$E_{n,3}^{\tilde{\xi}} := \bigcap_{k=n - \lfloor n^\varepsilon \rfloor + 1}^n \{ \tilde{\xi}_k \leq n^{2\varepsilon/\rho} \}.$$

On  $\bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}})$ , we have  $\sum_{k=1}^n k^{-r} e^{(1-s)\beta \tilde{S}_k \tilde{\xi}_k} + 1 \leq c_{40} n^{2\varepsilon + (2\varepsilon/\rho)}$ , while  $e^{[\beta s - (\beta - 1)]\tilde{S}_n} \geq n^{(3-\varepsilon)[\beta s - (\beta - 1)]/2}$  (recalling that  $\beta s > \beta - 1$ ). Therefore, with  $c_{41} := (2 + \frac{2}{\rho})^{\frac{s}{1-s}}$ ,

$$\begin{aligned} \mathbf{E}(W_{n,\beta}^{1-s}) &\geq (c_{38} c_{40})^{-s/(1-s)} n^{-c_{41}\varepsilon} n^{(3-\varepsilon)[\beta s - (\beta - 1)]/2} \\ (5.18) \quad &\times \mathbf{Q} \left\{ \bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}}) \right\}. \end{aligned}$$

We need to bound  $\mathbf{Q}(\bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}}))$  from below. Let  $\tilde{S}_0 := 0$ . Note that, under  $\mathbf{Q}$ ,  $(\tilde{S}_\ell - \tilde{S}_{\ell-1}, \tilde{\xi}_\ell)$ ,  $1 \leq \ell \leq n$ , are i.i.d., distributed as  $(S_1, \xi_1)$ . For  $j \leq n$ , let

$\tilde{\mathcal{G}}_j$  be the sigma-field generated by  $(\tilde{S}_k, \tilde{\xi}_k)$ ,  $1 \leq k \leq j$ . Then  $E_{n,1}^{\tilde{S}}, E_{n,2}^{\tilde{S}}, E_{n,1}^{\tilde{\xi}}$  and  $E_{n,2}^{\tilde{\xi}}$  are  $\tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor}$ -measurable, whereas  $E_{n,3}^{\tilde{\xi}}$  is independent of  $\tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor}$ . Therefore,

$$\begin{aligned} & \mathbf{Q} \left( \bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}}) \mid \tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor} \right) \\ & \geq [\mathbf{Q}(E_{n,3}^{\tilde{S}} \mid \tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor}) + \mathbf{Q}(E_{n,3}^{\tilde{\xi}}) - 1] \mathbf{1}_{E_{n,1}^{\tilde{S}} \cap E_{n,2}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}} \cap E_{n,2}^{\tilde{\xi}}}. \end{aligned}$$

We have  $c_{42} := \mathbf{E}\mathbf{Q}(\xi_1^\rho) < \infty$  [by (2.16)]; thus,  $\mathbf{Q}\{\xi_1 > n^{2\varepsilon/\rho}\} \leq c_{42}n^{-2\varepsilon}$ , which entails  $\mathbf{Q}(E_{n,3}^{\tilde{\xi}}) = (\mathbf{Q}\{\xi_1 \leq n^{2\varepsilon/\rho}\})^{\lfloor n^\varepsilon \rfloor} \geq (1 - c_{42}n^{-2\varepsilon})^{\lfloor n^\varepsilon \rfloor} \geq 1 - c_{43}n^{-\varepsilon}$ . To estimate  $\mathbf{Q}(E_{n,3}^{\tilde{S}} \mid \tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor})$ , we use the Markov property to see that, if  $\tilde{S}_{n-\lfloor n^\varepsilon \rfloor} \in I_n := [-2n^{\varepsilon/2}, -n^{\varepsilon/2}]$ , the conditional probability is (writing  $N := \lfloor n^\varepsilon \rfloor$ )

$$\begin{aligned} & \geq \inf_{z \in I_n} \mathbf{Q} \left\{ S_i \leq \frac{3}{2} \log n - z, \forall 1 \leq i \leq N - 1, \right. \\ & \qquad \left. \frac{3 - \varepsilon}{2} \log n - z \leq S_N \leq \frac{3}{2} \log n - z \right\}, \end{aligned}$$

which is greater than  $N^{-(1/2)+o(1)}$ . Therefore,

$$\mathbf{Q}(E_{n,3}^{\tilde{S}} \mid \tilde{\mathcal{G}}_{n-\lfloor n^\varepsilon \rfloor}) + \mathbf{Q}(E_{n,3}^{\tilde{\xi}}) - 1 \geq n^{-(\varepsilon/2)+o(1)} - c_{43}n^{-\varepsilon} = n^{-(\varepsilon/2)+o(1)}.$$

As a consequence,

$$(5.19) \quad \mathbf{Q} \left\{ \bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}}) \right\} \geq n^{-(\varepsilon/2)+o(1)} \mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,2}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}} \cap E_{n,2}^{\tilde{\xi}}).$$

To estimate  $\mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,2}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}} \cap E_{n,2}^{\tilde{\xi}})$ , we condition on  $\tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}$ , and note that  $E_{n,1}^{\tilde{S}}$  and  $E_{n,1}^{\tilde{\xi}}$  are  $\tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}$ -measurable, whereas  $E_{n,2}^{\tilde{\xi}}$  is independent of  $\tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}$ . Since  $\mathbf{Q}(E_{n,2}^{\tilde{S}} \mid \tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}) \geq n^{-(3-\varepsilon)/2+o(1)}$ , whereas  $\mathbf{Q}(E_{n,2}^{\tilde{\xi}}) = [\mathbf{Q}\{\xi_1 \leq e^{n^\varepsilon/4}\}]^{n-2\lfloor n^\varepsilon \rfloor} \geq [1 - c_{42}e^{-\rho n^{\varepsilon/4}}]^{n-2\lfloor n^\varepsilon \rfloor} \geq 1 - e^{-n^{\varepsilon/5}}$  (for large  $n$ ), we have

$$\begin{aligned} \mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,2}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}} \cap E_{n,2}^{\tilde{\xi}} \mid \tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}) & \geq [\mathbf{Q}(E_{n,2}^{\tilde{S}} \mid \tilde{\mathcal{G}}_{\lfloor n^\varepsilon \rfloor}) + \mathbf{Q}(E_{n,2}^{\tilde{\xi}}) - 1] \mathbf{1}_{E_{n,1}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}}} \\ & \geq n^{-(3-\varepsilon)/2+o(1)} \mathbf{1}_{E_{n,1}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}}}. \end{aligned}$$

Thus,  $\mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,2}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}} \cap E_{n,2}^{\tilde{\xi}}) \geq n^{-(3-\varepsilon)/2+o(1)} \mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}})$ . Going back to (5.19), we have

$$\begin{aligned} & \mathbf{Q} \left\{ \bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}}) \right\} \geq n^{-(3/2)+o(1)} \mathbf{Q}(E_{n,1}^{\tilde{S}} \cap E_{n,1}^{\tilde{\xi}}) \\ & \geq n^{-(3/2)+o(1)} [\mathbf{Q}(E_{n,1}^{\tilde{S}}) + \mathbf{Q}(E_{n,1}^{\tilde{\xi}}) - 1]. \end{aligned}$$

We choose the constant  $c_{39} > 0$  sufficiently small so that  $\mathbf{Q}(E_{n,1}^{\tilde{S}}) \geq n^{-(\varepsilon/2)+o(1)}$ , whereas  $\mathbf{Q}(E_{n,1}^{\tilde{\xi}}) = \mathbf{Q}(E_{n,3}^{\tilde{\xi}}) \geq 1 - c_{43}n^{-\varepsilon}$ . Accordingly,

$$\mathbf{Q}\left\{\bigcap_{i=1}^3 (E_{n,i}^{\tilde{S}} \cap E_{n,i}^{\tilde{\xi}})\right\} \geq n^{-(3+\varepsilon)/2+o(1)}, \quad n \rightarrow \infty.$$

Substituting this into (5.18) yields

$$\mathbf{E}(W_{n,\beta}^{1-s}) \geq n^{-c_{41}\varepsilon} n^{(3-\varepsilon)[\beta s - (\beta-1)]/2} n^{-(3+\varepsilon)/2+o(1)}.$$

Since  $\varepsilon$  can be as small as possible, this implies the lower bound in Theorem 1.6. □

**6. Proof of Theorem 1.5.** The basic idea in the proof of Theorem 1.5 is the same as in the proof of Theorem 1.6. Again, we prove the upper and lower bounds in distinct parts, for the sake of clarity. Throughout the section, we assume (1.1), (1.2) and (1.3).

*PROOF OF THEOREM 1.5: The upper bound.* Clearly,  $n^{1/2}W_n \leq \bar{Y}_n$ , where

$$\bar{Y}_n := \sum_{|u|=n} (n^{1/2} \vee V(u)^+) e^{-V(u)}.$$

Recall  $W_n^*$  from (3.7). Applying (3.14) to  $\lambda = 1$ , we see that  $\bar{Y}_n \geq \frac{1}{c_{44}} \log(\frac{1}{W_n^*})$ , with  $c_{44} := c_{12} + c_{13}$ . Thus,  $\mathbf{P}\{\bar{Y}_n < x, \mathcal{S}_n\} \leq \mathbf{P}\{\log(\frac{1}{W_n^*}) < c_{44}x, \mathcal{S}_n\} \leq e^{c_{44}} \times \mathbf{E}\{(W_n^*)^{1/x} \mathbf{1}_{\mathcal{S}_n}\}$ , which, according to (3.11), is bounded by  $e^{c_{44}}(x^\kappa + e^{-c_{10}n})$  for  $0 < x \leq \frac{1}{a_0}$ . Thus, for any fixed  $c > 0$  and  $0 < s < \min\{\frac{c_{10}}{c}, \kappa\}$ , we have  $\sup_{n \geq 1} \mathbf{E}\{\frac{1}{\bar{Y}_n} \mathbf{1}_{\{\bar{Y}_n \geq e^{-cn}\} \cap \mathcal{S}_n}\} < \infty$ . On the other hand, let  $c_{31}$  and  $c_{32}$  be as in (5.5); since  $\bar{Y}_n \geq \exp\{-\inf_{|u|=n} V(u)\}$ , it follows from (5.5) that  $\sup_{n \geq 1} \mathbf{E}\{\frac{1}{\bar{Y}_n^{c_{32}}} \mathbf{1}_{\{\bar{Y}_n < e^{-c_{31}n}\} \cap \mathcal{S}_n}\} < \infty$ . As a consequence,

$$(6.1) \quad \sup_{n \geq 1} \mathbf{E}\left\{\frac{1}{\bar{Y}_n^s} \mathbf{1}_{\mathcal{S}_n}\right\} < \infty, \quad 0 < s < \min\left\{c_{32}, \frac{c_{10}}{c_{31}}, \kappa\right\}.$$

We now fix  $0 < s < \min\{\frac{1}{2}, c_{32}, \frac{c_{10}}{c_{31}}, \kappa\}$ . Let  $K \geq 1$  and let  $E_n$  be the event in (5.11), satisfying  $\mathbf{Q}\{E_n\} \geq 1 - n^{-K}$  for  $n \geq n_0$ . We write

$$\mathbf{E}\{(n^{1/2}W_n)^{1-s}\} = \mathbf{E}\{(n^{1/2}W_n)^{1-s} \mathbf{1}_{E_n}\} + \mathbf{E}\{(n^{1/2}W_n)^{1-s} \mathbf{1}_{E_n^c}\}.$$

For  $n \geq n_0$ ,  $\mathbf{E}\{W_n^{1-s} \mathbf{1}_{E_n^c}\} \leq [\mathbf{E}\{W_n^{1-2s}\}]^{1/2} [\mathbf{E}\{W_n \mathbf{1}_{E_n^c}\}]^{1/2} = [\mathbf{E}\{W_n^{1-2s}\}]^{1/2} \times [\mathbf{Q}\{E_n^c\}]^{1/2} \leq [\mathbf{E}\{W_n\}]^{(1/2)-s} n^{-K/2}$ , which equals  $n^{-K/2}$  (since  $\mathbf{E}\{W_n\} = 1$ ).

Therefore, for  $n \rightarrow \infty$ ,

$$\mathbf{E}\{(n^{1/2}W_n)^{1-s}\} \leq \mathbf{E}\{\bar{Y}_n^{1-s} \mathbf{1}_{E_n}\} + o(1).$$

Exactly as in (5.13), we have  $\mathbf{E}\{\bar{Y}_n^{1-s} \mathbf{1}_{E_n}\} = \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \vee V(w_n^{(n)+})\bar{Y}_n^{-s} \mathbf{1}_{E_n}\}$ . Thus, for  $n \rightarrow \infty$ ,

$$(6.2) \quad \mathbf{E}\{(n^{1/2}W_n)^{1-s}\} \leq \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} + V(w_n^{(n)+})\bar{Y}_n^{-s} \mathbf{1}_{E_n}\} + o(1).$$

For any subset  $\mathcal{L} \subset \{1, 2, \dots, n\}$ , we have

$$\begin{aligned} \bar{Y}_n &\geq \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} \max\{n^{1/2}, V(x)^+\} e^{-V(x)} \\ &= \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} e^{-V(u)} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} \max\{n^{1/2}, [V(u) + V_u(x)]^+\} e^{-V_u(x)}. \end{aligned}$$

Recall that  $\underline{w}^{(n)}$  is the oldest vertex in  $\llbracket e, w_n^{(n)} \rrbracket$  such that  $V(\underline{w}^{(n)}) = \min_{u \in \llbracket e, w_n^{(n)} \rrbracket} V(u)$ . Let  $c_{35}$  be the constant in (5.8). We choose

$$\mathcal{L} := \begin{cases} \{j \leq n : \mathcal{I}_j^{(n)} \neq \emptyset, |\underline{w}^{(n)}| < j < |\underline{w}^{(n)}| + c_{35} \log n, \\ \quad \text{if } n - |\underline{w}^{(n)}| \geq 2c_{35} \log n, \\ \{j \leq n : \mathcal{I}_j^{(n)} \neq \emptyset, |\underline{w}^{(n)}| - c_{35} \log n < j < |\underline{w}^{(n)}|\}, \\ \quad \text{otherwise.} \end{cases}$$

On the event  $E_n$ , it is clear that  $\mathcal{L} \neq \emptyset$  and that, for any  $u \in \mathcal{I}_j^{(n)}$  (with  $j \in \mathcal{L}$ ),

$$(6.3) \quad |V(u) - V(\underline{w}^{(n)})| \leq c_{45} \log n,$$

where  $c_{45} := c_{36} + c_{37}$ , with  $c_{36}$  and  $c_{37}$  as in (5.9) and (5.10), respectively.

We distinguish two possible situations, depending on whether  $V(\underline{w}^{(n)}) \geq -c_{46} \log n$ , where  $c_{46} := \frac{1}{s} + c_{45}$ . In both situations, we consider a sufficiently large  $n$  and an arbitrary  $u \in \mathcal{I}_j^{(n)}$  (with  $j \in \mathcal{L}$ ).

On  $\{V(\underline{w}^{(n)}) \geq -c_{46} \log n\} \cap E_n$ , we have  $\max\{n^{1/2}, [V(u) + V_u(x)]^+\} \geq \frac{1}{2}(n^{1/2} \vee V_u(x)^+)$  [this holds trivially in case  $V_u(x) \leq n^{1/2}$ ; otherwise  $[V(u) + V_u(x)]^+ \geq V_u(x) - (c_{46} + c_{45}) \log n \geq \frac{1}{2}V_u(x)^+$ ] and, thus,

$$\begin{aligned} \bar{Y}_n &\geq \frac{1}{2} \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} e^{-V(u)} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} \max\{(n-j)^{1/2}, V_u(x)^+\} e^{-V_u(x)} \\ &=: \frac{1}{2} \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{I}_j^{(n)}} e^{-V(u)} \xi_u. \end{aligned}$$



If, however,  $V(\underline{w}^{(n)}) < -c_{46} \log n$ , then on  $E_n$ ,  $V(u) \leq V(\underline{w}^{(n)}) + c_{45} \log n < -\frac{1}{s} \log n$  and, since  $\max\{n^{1/2}, [V(u) + V_u(x)]^+\} \geq n^{1/2}$ , we have, in this case,

$$\begin{aligned} \bar{Y}_n &\geq n^{(1/s)+(1/2)} \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} e^{-V_u(x)} \\ &=: n^{(1/s)+(1/2)} \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} \eta_u. \end{aligned}$$

Therefore, in both situations we have

$$\begin{aligned} \bar{Y}_n^{-s} \mathbf{1}_{E_n} &\leq 2^s \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} e^{-V(u)} \xi_u \right)^{-s} \mathbf{1}_{E_n} \\ (6.4) \quad &+ n^{-(s/2)-1} \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} \eta_u \right)^{-s} \mathbf{1}_{E_n}. \end{aligned}$$

[Since  $\sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} 1 > 0$  on  $E_n$ , the  $(\cdot)^{-s}$  expressions on the right-hand side are well defined.]

We claim that there exists  $0 < s_0 < 1$  such that, for any  $\varepsilon > 0$  and  $s \in (0, s_0)$ ,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}} \left\{ (n^{1/2} + V(w_n^{(n)})^+) \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} e^{-V(u)} \xi_u \right)^{-s} \mathbf{1}_{E_n} \right\} \\ (6.5) \quad \leq c_{48}, \end{aligned}$$

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}} \left\{ (n^{1/2} + V(w_n^{(n)})^+) \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{F}_j^{(n)}} \eta_u \right)^{-s} \mathbf{1}_{E_n} \right\} \\ (6.6) \quad \leq c_{47} n^{1/2+(3+\varepsilon)/2s}. \end{aligned}$$

We admit (6.5) and (6.6) for the time being. In view of (6.4), we obtain, for  $0 < s < s_* := \min\{\frac{1}{2}, s_0, c_{32}, \frac{c_{10}}{c_{31}}, \kappa\}$ ,

$$\mathbf{E}_{\mathbf{Q}} \{ (n^{1/2} + V(w_n^{(n)})^+) \bar{Y}_n^{-s} \mathbf{1}_{E_n} \} \leq 2^s c_{48} + o(1).$$

Substituting this in (6.2), we see that  $\sup_{n \geq 1} \mathbf{E}\{(n^{1/2} W_n)^{1-s}\} < \infty$  for any  $s \in (0, s_*)$ . This yields the last inequality in (1.16) when  $\gamma$  is close to 1. By Jensen’s inequality, it holds for all  $\gamma \in [0, 1)$ . This will complete the proof of the upper bound in Theorem 1.5.

It remains to check (6.5) and (6.6). We only present the proof of (6.5), because the proof of (6.6) is similar and slightly easier, using (5.2) in place of (6.1).

Recall  $\mathcal{G}_n$  from (2.9). By Proposition 2.1, under  $\mathbf{Q}$  and conditionally on  $\mathcal{G}_n$ , the random variables  $\xi_u$ , for  $u \in \mathcal{J}_j^{(n)}$  and  $j \in \mathcal{L}$ , are independent. We write  $\mathcal{L} := \{j(1), \dots, j(N)\}$ , with  $j(1) < \dots < j(N)$ . It follows from the second part of Lemma 5.2 that

$$\begin{aligned} & \mathbf{E}_{\mathbf{Q}} \left\{ \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{J}_j^{(n)}} e^{-V(u)} \xi_u \right)^{-s} \mathbf{1}_{E_n} \middle| \mathcal{G}_n \right\} \\ & \leq \sum_{i=1}^N b^{i-1} \mathbf{E}_{\mathbf{Q}} \left\{ \left( \sum_{u \in \mathcal{J}_{j(i)}^{(n)}} e^{-V(u)} \xi_u \right)^{-s} \mathbf{1}_{\{\sum_{u \in \mathcal{J}_{j(i)}^{(n)}} e^{-V(u)} \xi_u > 0\}} \middle| \mathcal{G}_n \right\}, \end{aligned}$$

where  $b := \max_{j \in \mathcal{L}} \mathbf{Q}\{\sum_{u \in \mathcal{J}_j^{(n)}} e^{-V(u)} \xi_u = 0 \mid \mathcal{G}_n\}$ . We note that  $b \leq \max_{1 \leq j \leq n} \mathbf{P}\{\mathcal{S}_{n-j}^c\} \leq q$ , and that, for any  $i \leq N$ , the  $\mathbf{E}_{\mathbf{Q}}\{\cdot\}$  expression on the right-hand side is, according to the first part of Lemma 5.2, bounded by

$$\frac{1}{1-q} \max_{u \in \mathcal{J}_{j(i)}^{(n)}} \mathbf{E}_{\mathbf{Q}} \left\{ \frac{e^{sV(u)}}{\xi_u^s} \mathbf{1}_{\{\xi_u > 0\}} \middle| \mathcal{G}_n \right\}.$$

By Proposition 2.1,  $\mathbf{E}_{\mathbf{Q}}\{\frac{1}{\xi_u^s} \mathbf{1}_{\{\xi_u > 0\}} \mid \mathcal{G}_n\} = \mathbf{E}\{\frac{1}{Y_{n-j}^s} \mathbf{1}_{\mathcal{S}_{n-j}}\}$ , which is bounded in  $n$  and  $j$  [by (6.1)]. Summarizing, we have proved that

$$\mathbf{E}_{\mathbf{Q}} \left\{ \left( \sum_{j \in \mathcal{L}} \sum_{u \in \mathcal{J}_j^{(n)}} e^{-V(u)} \xi_u \right)^{-s} \mathbf{1}_{E_n} \middle| \mathcal{G}_n \right\} \leq c_{49} \sum_{i=1}^N q^{i-1} \max_{u \in \mathcal{J}_{j(i)}^{(n)}} e^{sV(u)}.$$

As a consequence, the expression on the left-hand side of (6.5) is bounded by  $c_{49} \mathbf{E}_{\mathbf{Q}}\{\Lambda_n\}$ , where

$$\begin{aligned} \Lambda_n & := (n^{1/2} + V(w_n^{(n)})^+) \sum_{i=1}^N q^{i-1} \max_{u \in \mathcal{J}_{j(i)}^{(n)}} e^{sV(u)} \mathbf{1}_{\{|V(u) - V(\underline{w}^{(n)})| \leq c_{45} \log n\}} \\ & \leq \tilde{\Lambda}_n := (n^{1/2} + V(w_n^{(n)})^+) \sum_{i=1}^N q^{i-1} \max_{u \in \mathcal{J}_{j(i)}^{(n)}} e^{sV(u)}. \end{aligned}$$

The proof of (6.5) now boils down to verifying the following estimates: there exists  $0 < s_0 < 1$  such that, for any  $s \in (0, s_0)$ ,

$$(6.7) \quad \sup_n \mathbf{E}_{\mathbf{Q}} \{ \tilde{\Lambda}_n \mathbf{1}_{\{n - |\underline{w}^{(n)}| \geq 2c_{35} \log n\}} \} < \infty,$$

$$(6.8) \quad \lim_{n \rightarrow \infty} \mathbf{E}_{\mathbf{Q}} \{ \Lambda_n \mathbf{1}_{\{n - |\underline{w}^{(n)}| < 2c_{35} \log n\}} \} = 0.$$

Let us first check (6.7). Let  $S_0 := 0$  and let  $(S_j - S_{j-1}, \sigma_j, \Delta_j)$ ,  $j \geq 1$ , be i.i.d. random variables under  $\mathbf{Q}$  and distributed as  $(V(w^{(1)}), \#\mathcal{S}_1^{(1)}, \max_{u \in \mathcal{S}_1^{(1)}} e^{sV(u)})$ .

Let

$$\underline{S}_n := \min_{0 \leq i \leq n} S_i, \quad \vartheta_n := \inf\{k \geq 0 : S_k = \underline{S}_n\}.$$

[The random variable  $\vartheta_n$  has nothing to do with the constant  $\vartheta$  in Proposition 3.1.] Writing LHS<sub>(6.7)</sub> for  $\mathbf{E}_{\mathbf{Q}}\{\tilde{\Lambda}_n \mathbf{1}_{\{n - |\underline{w}^{(n)}| \geq 2c_{35} \log n\}}\}$ , it follows from Proposition 2.1 that

$$\begin{aligned} \text{LHS}_{(6.7)} &= \mathbf{E}_{\mathbf{Q}}\left\{ [n^{1/2} + S_n^+] \sum_{i=1}^M q^{i-1} e^{s S_{\ell(i)-1}} \Delta_{\ell(i)} \mathbf{1}_{\{n - \vartheta_n \geq 2c_{35} \log n\}} \right\} \\ &= \mathbf{E}_{\mathbf{Q}}\left\{ [n^{1/2} + S_n^+] e^{s \underline{S}_n} \sum_{i=1}^M q^{i-1} e^{s [S_{\ell(i)-1} - S_{\ell(0)}]} \Delta_{\ell(i)} \mathbf{1}_{\{n - \vartheta_n \geq 2c_{35} \log n\}} \right\}, \end{aligned}$$

where  $\ell(i) := \inf\{k > \ell(i - 1) : \sigma_k \geq 1\}$  with  $\ell(0) := \vartheta_n$ , and  $M := \sup\{i : \ell(i) < \vartheta_n + c_{35} \log n\}$ .

At this stage, we use a standard trick for random walks: let  $\nu_0 := 0$  and let

$$\nu_i := \inf\left\{k > \nu_{i-1} : S_k < \min_{0 \leq j \leq \nu_{i-1}} S_j\right\}, \quad i \geq 1.$$

In words,  $0 = \nu_0 < \nu_1 < \dots$  are strict descending ladder times. On the event  $\{\nu_k \leq n < \nu_{k+1}\}$  (for  $k \geq 0$ ), we have  $\vartheta_n = \nu_k$  and  $\underline{S}_n = S_{\nu_k}$ . Thus, LHS<sub>(6.7)</sub> equals

$$\begin{aligned} \sum_{k=0}^{\infty} \mathbf{E}_{\mathbf{Q}}\left\{ \mathbf{1}_{\{n - \nu_k \geq 2c_{35} \log n\}} \mathbf{1}_{\{\nu_k \leq n < \nu_{k+1}\}} [n^{1/2} + S_n^+] e^{s S_{\nu_k}} \right. \\ \left. \times \sum_{i=1}^M q^{i-1} e^{s [S_{\ell(i)-1} - S_{\ell(0)}]} \Delta_{\ell(i)} \right\}. \end{aligned}$$

For any  $k$ , we look at the expectation  $\mathbf{E}_{\mathbf{Q}}\{\cdot\}$  on the right-hand side. By conditioning upon  $(S_j, \sigma_j, \Delta_j, 1 \leq j \leq \nu_k)$ , and since  $S_n^+ = [S_{\nu_k} + (S_n - S_{\nu_k})]^+ \leq (S_n - S_{\nu_k})^+ = S_n - S_{\nu_k}$  on  $\{\nu_k \leq n < \nu_{k+1}\}$ , we obtain

$$(6.9) \quad \text{LHS}_{(6.7)} \leq \sum_{k=0}^{\infty} \mathbf{E}_{\mathbf{Q}}\{\mathbf{1}_{\{n - \nu_k \geq 2c_{35} \log n\}} e^{s S_{\nu_k}} f_n(n - \nu_k)\},$$

where, for any  $1 \leq j \leq n$ ,

$$f_n(j) := \mathbf{E}_{\mathbf{Q}}\left\{ \mathbf{1}_{\{\nu_1 > j\}} [n^{1/2} + S_j] \sum_{i=1}^{M'} q^{i-1} e^{s S_{m(i)-1}} \Delta_{m(i)} \right\},$$

and  $m(i) := \inf\{k > m(i - 1) : \sigma_k \geq 1\}$  with  $m(0) := 0$ , and  $M' := \sup\{i : m(i) < c_{35} \log n\}$ . For brevity, we write  $L_n := \sum_{i=1}^{M'} q^{i-1} e^{s S_{m(i)-1}} \Delta_{m(i)} = \sum_{i=1}^{\infty} q^{i-1} \times e^{s S_{m(i)-1}} \Delta_{m(i)} \mathbf{1}_{\{m(i) < c_{35} \log n\}}$  for the moment. By the Cauchy–Schwarz inequality,

$$f_n(j) \leq [\mathbf{Q}\{\nu_1 > j\}]^{1/2} [\mathbf{E}_{\mathbf{Q}}\{(n^{1/2} + S_j)^2 | \nu_1 > j\}]^{1/2} [\mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{\nu_1 > j\}}\}]^{1/2}.$$

By (2.13),  $\mathbf{Q}\{v_1 > j\} \leq c_{50}j^{-1/2}$  for some  $c_{50} > 0$  and all  $j \geq 1$ . On the other hand,  $(n^{1/2} + S_j)^2 \leq 2(n + S_j^2)$ , and it is known (Bolthausen [11]) that  $\mathbf{E}_{\mathbf{Q}}\{\frac{S_j^2}{j} | v_1 > j\} \rightarrow c_{51} \in (0, \infty)$  for  $j \rightarrow \infty$ . Therefore,  $\mathbf{E}_{\mathbf{Q}}\{(n^{1/2} + S_j)^2 | v_1 > j\} \leq c_{52}n$  for some  $c_{52} > 0$  and all  $n \geq j \geq 1$ . Accordingly, with  $c_{53} := c_{50}^{1/2} c_{52}^{1/2}$ , we have

$$f_n(j) \leq c_{53}j^{-1/4}n^{1/2}[\mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{v_1 > j\}}\}]^{1/2}, \quad 1 \leq j \leq n.$$

By the Cauchy–Schwarz inequality,  $L_n^2 \leq (\sum_{i=1}^{\infty} q^{i-1}) \sum_{i=1}^{\infty} q^{i-1} e^{2sS_{m(i)-1}} \Delta_{m(i)}^2 \times \mathbf{1}_{\{m(i) < c_{35} \log n\}}$ . Therefore, for  $j \geq 2c_{35} \log n$ ,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{v_1 > j\}}\} &\leq \frac{1}{1-q} \sum_{i=1}^{\infty} q^{i-1} \mathbf{E}_{\mathbf{Q}}\{e^{2sS_{m(i)-1}} \Delta_{m(i)}^2 \mathbf{1}_{\{m(i) < c_{35} \log n\}} \mathbf{1}_{\{v_1 > j\}}\} \\ &\leq \frac{1}{1-q} \sum_{i=1}^{\infty} q^{i-1} \mathbf{E}_{\mathbf{Q}}\{e^{2sS_{m(i)-1}} \Delta_{m(i)}^2 \mathbf{1}_{\{m(i) \leq j/2\}} \mathbf{1}_{\{v_1 > j\}}\}. \end{aligned}$$

For any  $i \geq 1$ , to estimate the expectation  $\mathbf{E}_{\mathbf{Q}}\{\cdot\}$  on the right-hand side, we apply the strong Markov property at time  $m(i)$  to see that

$$\mathbf{E}_{\mathbf{Q}}\{\cdot\} = \mathbf{E}_{\mathbf{Q}}\{e^{2sS_{m(i)-1}} \Delta_{m(i)}^2 \mathbf{1}_{\{m(i) \leq j/2\}} \mathbf{1}_{\{v_1 > m(i)\}} g(S_{m(i)}, j - m(i))\},$$

where  $g(z, k) := \mathbf{Q}\{z + S_i \geq 0, \forall 1 \leq i \leq k\}$  for any  $z \geq 0$  and  $k \geq 1$ . By (13) of Kozlov [24],  $g(z, k) \leq c_{54}(z + 1)/k^{1/2}$  for some  $c_{54} > 0$  and all  $z \geq 0$  and  $k \geq 1$ . Since  $z + 1 \leq c_{55}e^{sz}$  for all  $z \geq 0$ , this yields, with  $c_{56} := \frac{c_{55}}{1-q}$ ,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{v_1 > j\}}\} &\leq c_{56} \sum_{i=1}^{\infty} q^{i-1} \mathbf{E}_{\mathbf{Q}}\left\{e^{2sS_{m(i)-1}} \Delta_{m(i)}^2 \mathbf{1}_{\{m(i) \leq j/2\}} \frac{e^{sS_{m(i)}}}{(j - m(i))^{1/2}}\right\} \\ &\leq \frac{c_{56}}{(j/2)^{1/2}} \sum_{i=1}^{\infty} q^{i-1} \mathbf{E}_{\mathbf{Q}}\{e^{2sS_{m(i)-1} + sS_{m(i)}} \Delta_{m(i)}^2\} \\ &= \frac{2^{1/2}c_{56}}{j^{1/2}} \mathbf{E}_{\mathbf{Q}}\left\{\sum_{i=1}^{\infty} q^{i-1} e^{2sS_{m(i)-1} + sS_{m(i)}} \Delta_{m(i)}^2\right\}. \end{aligned}$$

We observe that  $\sum_{i=1}^{\infty} q^{i-1} e^{2sS_{m(i)-1} + sS_{m(i)}} \Delta_{m(i)}^2 \leq \sum_{k=1}^{\infty} q^{R(k)-1} e^{2sS_{k-1} + sS_k} \Delta_k^2$ , where  $R(k) := \#\{1 \leq j \leq k : s_j \geq 1\}$ . Therefore, with  $c_{57} := 2^{1/2}c_{56}$ ,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{v_1 > j\}}\} &\leq \frac{c_{57}}{j^{1/2}} \sum_{k=1}^{\infty} \mathbf{E}_{\mathbf{Q}}\{q^{R(k)-1} e^{2sS_{k-1} + sS_k} \Delta_k^2\} \\ &\leq \frac{c_{57}}{j^{1/2}} \sum_{k=1}^{\infty} [\mathbf{E}_{\mathbf{Q}}\{q^{2[R(k)-1]}\}]^{1/2} [\mathbf{E}_{\mathbf{Q}}\{e^{4sS_{k-1} + 2sS_k} \Delta_k^4\}]^{1/2}. \end{aligned}$$

By definition,  $\mathbf{E}_{\mathbf{Q}}\{q^{2[R(k)-1]}\} = q^{-2r^k}$ , with  $r := \mathbf{Q}(\sigma_1 = 0) + q^2\mathbf{Q}(\sigma_1 \geq 1) < 1$  [because  $q < 1$  and  $\mathbf{Q}(\sigma_1 = 0) < 1$ ]. On the other hand,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{e^{4sS_{k-1}+2sS_k} \Delta_k^4\} &= \mathbf{E}_{\mathbf{Q}}\{e^{6sS_{k-1}}\} \mathbf{E}_{\mathbf{Q}}\{e^{2s(S_k-S_{k-1})} \Delta_k^4\} \\ &= [\mathbf{E}_{\mathbf{Q}}\{e^{6sS_1}\}]^{k-1} \mathbf{E}_{\mathbf{Q}}\{e^{2sS_1} \Delta_1^4\}. \end{aligned}$$

By (2.11), there exists  $s_{\#} > 0$  sufficiently small such that  $\mathbf{E}_{\mathbf{Q}}\{e^{6sS_1}\} < \frac{1}{r}$  for all  $0 < s < s_{\#}$ . On the other hand,  $\mathbf{E}_{\mathbf{Q}}\{\Delta_1^8\} < \infty$  for  $0 < s < \frac{c_6}{8}$  [by (2.17)], and  $\mathbf{E}_{\mathbf{Q}}\{e^{4sS_1}\} < \infty$  for  $0 < s \leq \frac{c_2}{4}$  [by (2.11)]; thus,  $\mathbf{E}_{\mathbf{Q}}\{e^{2sS_1} \Delta_1^4\} < \infty$  for  $0 < s < \min\{\frac{c_6}{8}, \frac{c_2}{4}\}$ . As a consequence, for any  $0 < s < \min\{s_{\#}, \frac{c_6}{8}, \frac{c_2}{4}\}$ , we have  $\mathbf{E}_{\mathbf{Q}}\{L_n^2 \mathbf{1}_{\{v_1 > j\}}\} \leq \frac{c_{58}}{j^{1/2}}$ , for some  $c_{58} > 0$  and all  $n \geq j \geq 1$  with  $j \geq 2c_{35} \log n$ , which yields

$$f_n(j) \leq c_{53} c_{58}^{1/2} j^{-1/2} n^{1/2}.$$

Going back to (6.9), we obtain, for any  $0 < s < \min\{s_{\#}, \frac{c_6}{8}, \frac{c_2}{4}\}$  and  $c_{59} := c_{53} c_{58}^{1/2}$ ,

$$\text{LHS}_{(6.7)} \leq c_{59} n^{1/2} \sum_{k=0}^{\infty} \mathbf{E}_{\mathbf{Q}}\left\{ \mathbf{1}_{\{n-v_k \geq 2c_{35} \log n\}} \frac{e^{sS_{v_k}}}{(n-v_k)^{1/2}} \right\}.$$

By (2.13) again,  $\frac{1}{j^{1/2}} \leq c_{60} \mathbf{Q}\{v_1 > j\}$  for all  $j \geq 1$ . Thus, with  $c_{61} := c_{59} c_{60}$ ,

$$\begin{aligned} \text{LHS}_{(6.7)} &\leq c_{61} n^{1/2} \sum_{k=0}^{\infty} \mathbf{E}_{\mathbf{Q}}\{ \mathbf{1}_{\{n-v_k \geq 2c_{35} \log n\}} e^{sS_{v_k}} \mathbf{1}_{\{v_{k+1} > n\}} \} \\ &\leq c_{61} n^{1/2} \sum_{k=0}^{\infty} \mathbf{E}_{\mathbf{Q}}\{ \mathbf{1}_{\{v_k \leq n < v_{k+1}\}} e^{sS_{v_k}} \}, \end{aligned}$$

which equals  $c_{61} n^{1/2} \mathbf{E}_{\mathbf{Q}}\{e^{s \min_{0 \leq i \leq n} S_i}\}$ , and, according to (2.14), is bounded in  $n$ . This completes the proof of (6.7).

It remains to check (6.8). By definition,

$$\Lambda_n \leq [n^{1/2} + V(w_n^{(n)})^+] n^{sc_{45}} e^{sV(w^{(n)})} \sum_{i=1}^N q^{i-1}.$$

Since  $\sum_{i=1}^N q^{i-1} \leq \frac{1}{1-q}$ , this leads to, by an application of Proposition 2.1,

$$\mathbf{E}_{\mathbf{Q}}\{ \Lambda_n \mathbf{1}_{\{n-|w^{(n)}| < 2c_{35} \log n\}} \} \leq \frac{n^{sc_{45}}}{1-q} \mathbf{E}_{\mathbf{Q}}\{ [n^{1/2} + S_n^+] e^{s\underline{S}_n} \mathbf{1}_{\{n-\vartheta_n < 2c_{35} \log n\}} \},$$

where  $(S_i)$  is as in Proposition 2.1 and, as before,  $\underline{S}_n := \min_{0 \leq i \leq n} S_i$ ,  $\vartheta_n := \inf\{k \geq 0 : S_k = \underline{S}_n\}$ .

Let  $0 < \varepsilon < \frac{1}{2}$ ; let  $A_n := \{S_n > n^{1/2+\varepsilon}\}$  and  $B_n := \{S_n \leq n^{1/2+\varepsilon}\} = A_n^c$ .

Since  $\mathbf{E}_{\mathbf{Q}}\{e^{aS_1}\} < \infty$  for  $|a| < c_2$  [see (2.11)] and  $\mathbf{Q}(A_n) \leq 2 \exp(-c_3 n^{2\varepsilon})$  [see (2.12)], the Cauchy–Schwarz inequality yields  $n^{sc_{45}} \mathbf{E}_{\mathbf{Q}}\{[n^{1/2} + S_n^+]e^{sS_n} \mathbf{1}_{A_n}\} \rightarrow 0$ ,  $n \rightarrow \infty$ .

On  $B_n$ , we have  $n^{1/2} + S_n^+ \leq 2n^{1/2+\varepsilon}$ ; thus,  $\mathbf{E}_{\mathbf{Q}}\{[n^{1/2} + S_n^+]e^{sS_n} \times \mathbf{1}_{B_n \cap \{n - \vartheta_n < 2c_{35} \log n\}}\} \leq 2n^{1/2+\varepsilon} \mathbf{E}_{\mathbf{Q}}\{e^{sS_n} \mathbf{1}_{\{n - \vartheta_n < 2c_{35} \log n\}}\}$ . It is clear that  $\underline{S}_n \leq \underline{S}_{\lfloor n/2 \rfloor} := \min_{0 \leq i \leq \lfloor n/2 \rfloor} S_i$ , and that  $\{n - \vartheta_n < 2c_{35} \log n\} \subset \{\frac{n}{2} - \tilde{\vartheta}_{n/2} < 2c_{35} \log n\}$ , where  $\tilde{\vartheta}_{n/2} := \min\{k \geq 0: \tilde{S}_k = \min_{0 \leq i \leq n - \lfloor n/2 \rfloor} \tilde{S}_i\}$ , with  $\tilde{S}_i := S_{i + \lfloor n/2 \rfloor} - S_{\lfloor n/2 \rfloor}$ ,  $i \geq 0$ . Since  $\underline{S}_{\lfloor n/2 \rfloor}$  and  $\tilde{\vartheta}_{n/2}$  are independent, we have  $\mathbf{E}_{\mathbf{Q}}\{e^{s\underline{S}_n} \mathbf{1}_{\{n - \vartheta_n < 2c_{35} \log n\}}\} \leq \mathbf{E}_{\mathbf{Q}}\{e^{s\underline{S}_{\lfloor n/2 \rfloor}}\} \mathbf{Q}\{\frac{n}{2} - \tilde{\vartheta}_{n/2} < 2c_{35} \log n\}$ . By (2.14),  $\mathbf{E}_{\mathbf{Q}}\{e^{s\underline{S}_{\lfloor n/2 \rfloor}}\} \leq c_{62} n^{-1/2}$ ; on the other hand,  $\mathbf{Q}\{\frac{n}{2} - \tilde{\vartheta}_{n/2} < 2c_{35} \log n\} \leq c_{63} \frac{(\log n)^{1/2}}{n^{1/2}}$  (see Feller [18], page 398). Therefore,  $\mathbf{E}_{\mathbf{Q}}\{[n^{1/2} + S_n^+]e^{s\underline{S}_n} \mathbf{1}_{B_n \cap \{n - \vartheta_n < 2c_{35} \log n\}}\} \leq c_{64} n^{-1/2+\varepsilon} (\log n)^{1/2}$ .

Summarizing, we have proved that, for any  $s > 0$  and  $0 < \varepsilon < \frac{1}{2}$ , when  $n \rightarrow \infty$ ,

$$\mathbf{E}_{\mathbf{Q}}\{\Lambda_n \mathbf{1}_{\{n - \underline{w}^{(n)} < 2c_{35} \log n\}}\} \leq o(1) + \frac{c_{64}}{1 - q} n^{sc_{45} - 1/2 + \varepsilon} (\log n)^{1/2},$$

which yields (6.8), as long as  $0 < s < \frac{1}{2c_{45}}$ .  $\square$

PROOF OF THEOREM 1.5. *The lower bound.* We start with

$$n^{1/2} W_n \geq \underline{Y}_n := \sum_{|u|=n} (n^{1/2} \wedge V(u)^+) e^{-V(u)}.$$

Let  $s \in (0, 1)$ . Exactly as in (5.13), we have

$$(6.10) \quad \mathbf{E}\{\underline{Y}_n^{1-s}\} = \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)}))^+ \underline{Y}_n^{-s}\}.$$

By definition,

$$\begin{aligned} \underline{Y}_n &= \sum_{j=1}^n \sum_{u \in \mathcal{J}_j^{(n)}} e^{-V(u)} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} \min\{n^{1/2}, [V(u) + V_u(x)]^+\} e^{-V_u(x)} \\ &\quad + \min\{n^{1/2}, V(w_n^{(n)})^+\} e^{-V(w_n^{(n)})} \\ &\leq \sum_{j=1}^n e^{-V(w_{j-1}^{(n)})} \sum_{u \in \mathcal{J}_j^{(n)}} e^{-\Delta_u} \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j} [V(w_{j-1}^{(n)})^+ + \Delta_u^+ + V_u(x)^+] \\ &\quad \times e^{-V_u(x)} + \Theta_n, \end{aligned}$$

where  $\Delta_u := V(u) - V(w_{j-1}^{(n)})$  [for  $u \in \mathcal{J}_j^{(n)}$ ], and  $\Theta_n := V(w_n^{(n)})^+ e^{-V(w_n^{(n)})}$ .

By means of the elementary inequality  $(\sum_i a_i)^{-s} \geq (\sum_i a_i^s)^{-1}$  and  $(\sum_i b_i)^s \leq \sum_i b_i^s$  for nonnegative  $a_i$  and  $b_i$ , we obtain  $\underline{Y}_n^{-s} \geq \frac{1}{Z_n}$  on  $\mathcal{S}_n$ , with  $Z_n$  being defined

as

$$\sum_j e^{-sV(w_{j-1}^{(n)})} \sum_u e^{-s\Delta_u} \left\{ [(V(w_{j-1}^{(n)}))^+]^s + (\Delta_u^+)^s \right\} \left( \sum_x e^{-V_u(x)} \right)^s + \left[ \sum_x V_u(x)^+ e^{-V_u(x)} \right]^s \Big\} + \Theta_n^s,$$

where  $\sum_j := \sum_{j=1}^n$ ,  $\sum_u := \sum_{u \in \mathcal{S}_j^{(n)}}$ , and  $\sum_x := \sum_{x \in \mathbb{T}_u^{\text{GW}}, |x|_u = n-j}$ . We now condition upon  $\mathcal{G}_n$ , and note that  $V(w_j^{(n)})$  and  $\mathcal{S}_j^{(n)}$  are  $\mathcal{G}_n$ -measurable. By Proposition 2.1,

$$\mathbf{E}_{\mathbf{Q}}\{Z_n | \mathcal{G}_n\} = \sum_j e^{-sV(w_{j-1}^{(n)})} \sum_u e^{-s\Delta_u} \{ ((V(w_{j-1}^{(n)}))^+)^s + (\Delta_u^+)^s \} \times \mathbf{E}(W_{n-j}^s + U_{n-j}^s) \Big\} + \Theta_n^s,$$

where, for any  $k \geq 0$ ,  $U_k := \sum_{|y|=k} V(y)^+ e^{-V(y)}$ . By Jensen’s inequality,  $\mathbf{E}(W_{n-j}^s) \leq [\mathbf{E}(W_{n-j})]^s = 1$ . On the other hand, by (3.9),  $U_k \leq c_{65} \log \frac{1}{W_k^*}$  and, thus, by Lemma 3.3,  $\mathbf{E}(U_k^s) \leq c_{65}^s \mathbf{E}\{[\log \frac{1}{W_k^*}]^s\} \leq c_{66}$ . Therefore, the  $\sum_u$  sum on the right-hand side (without  $\Theta_n^s$ , of course) is

$$\begin{aligned} &\leq \sum_u e^{-s\Delta_u} \{ (V(w_{j-1}^{(n)}))^+)^s + (\Delta_u^+)^s + c_{67} \} \\ &= [V(w_{j-1}^{(n)})^+]^s \sum_u e^{-s\Delta_u} + \sum_u e^{-s\Delta_u} \{ (\Delta_u^+)^s + c_{67} \}. \end{aligned}$$

There exists  $c_{68} = c_{68}(s) < \infty$  such that  $e^{-sa} \{ (a^+)^s + c_{67} \} \leq c_{68} (e^{-sa} + e^{-sa/2})$  for all  $a \in \mathbb{R}$ . As a consequence,

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{Z_n | \mathcal{G}_n\} &\leq c_{69} \sum_{j=1}^n e^{-sV(w_{j-1}^{(n)})} \{ [V(w_{j-1}^{(n)})^+]^s + 1 \} \\ &\quad \times \sum_{u \in \mathcal{S}_j^{(n)}} [e^{-s\Delta_u} + e^{-s/2\Delta_u}] + \Theta_n^s. \end{aligned}$$

By Jensen’s inequality again,  $\mathbf{E}_{\mathbf{Q}}\{\frac{1}{Z_n} | \mathcal{G}_n\} \geq \frac{1}{\mathbf{E}_{\mathbf{Q}}\{Z_n | \mathcal{G}_n\}}$ . Since  $Y_n^{-s} \geq \frac{1}{Z_n}$  on  $\mathcal{S}_n$ , this leads to

$$\begin{aligned} &\mathbf{E}_{\mathbf{Q}}\{Y_n^{-s} | \mathcal{G}_n\} \\ &\geq \frac{c_{70}}{\sum_{j=1}^n e^{-sV(w_{j-1}^{(n)})} \{ [V(w_{j-1}^{(n)})^+]^s + 1 \} \sum_{u \in \mathcal{S}_j^{(n)}} [e^{-s\Delta_u} + e^{-\frac{s}{2}\Delta_u}] + \Theta_n^s}. \end{aligned}$$

We apply Proposition 2.1: if  $(S_j - S_{j-1}, \eta_j)$ , for  $j \geq 1$  (with  $S_0 := 0$ ), are i.i.d. random variables (under  $\mathbf{Q}$ ) and distributed as  $(V(w^{(1)}), \sum_{u \in \mathcal{J}_1^{(1)}} [e^{-sV(u)} + e^{-s/2V(u)}])$ , then

$$\begin{aligned} & \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} \\ & \geq c_{70} \mathbf{E}_{\mathbf{Q}} \left\{ \frac{n^{1/2} \wedge S_n^+}{\sum_{j=1}^n e^{-sS_{j-1}} [(S_{j-1}^+)^s + 1] \eta_j + e^{-sS_n} (S_n^+)^s} \right\} \\ & \geq c_{70} \mathbf{E}_{\mathbf{Q}} \left\{ \frac{(n^{1/2} \wedge S_n) \mathbf{1}_{\{\min_{1 \leq j \leq n} S_j > 0\}}}{\sum_{j=1}^n e^{-sS_{j-1}} (S_{j-1}^s + 1) \eta_j + e^{-sS_n} S_n^s} \right\}. \end{aligned}$$

Note that if  $S_j > 0$ , then  $e^{-sS_j} [S_j^s + 1] \leq c_{71} e^{-tS_j}$  with  $t := \frac{s}{2}$ . Therefore, by writing

$$\mathbf{Q}^{(n)}\{\cdot\} := \mathbf{Q}\left\{\cdot \mid \min_{1 \leq j \leq n} S_j > 0\right\},$$

and  $\mathbf{E}_{\mathbf{Q}}^{(n)}$  the expectation with respect to  $\mathbf{Q}^{(n)}$ , and  $\hat{\eta}_j := \eta_j + 1$  for brevity, we get that

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} & \geq c_{72} \mathbf{Q}\left\{\min_{1 \leq j \leq n} S_j > 0\right\} \mathbf{E}_{\mathbf{Q}}^{(n)} \left\{ \frac{n^{1/2} \wedge S_n}{\sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j} \right\} \\ & \geq c_{72} \mathbf{Q}\left\{\min_{1 \leq j \leq n} S_j > 0\right\} \mathbf{E}_{\mathbf{Q}}^{(n)} \left\{ \frac{\varepsilon n^{1/2} \mathbf{1}_{\{S_n > \varepsilon n^{1/2}\}}}{\sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j} \right\}. \end{aligned}$$

Since  $\mathbf{Q}\{\min_{1 \leq j \leq n} S_j > 0\} \geq c_{73} n^{-1/2}$  [see (2.13)], this leads to

$$\begin{aligned} & \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} \\ & \geq c_{74} \varepsilon \mathbf{E}_{\mathbf{Q}}^{(n)} \left\{ \frac{\mathbf{1}_{\{S_n > \varepsilon n^{1/2}\}}}{\sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j} \right\} \\ & \geq c_{74} \varepsilon \left[ \mathbf{E}_{\mathbf{Q}}^{(n)} \left\{ \frac{1}{\sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j} \right\} - \mathbf{Q}^{(n)}\{S_n \leq \varepsilon n^{1/2}\} \right]. \end{aligned}$$

Let  $\rho(s) > 0$  be as in Corollary 2.4. We have  $\mathbf{E}_{\mathbf{Q}}\{(\sum_{u \in \mathcal{J}_1^{(1)}} e^{-sV(u)})^{\rho(s)}\} < \infty$  by (2.16). Since  $\rho(s) \leq \rho(\frac{s}{2})$ , we also have  $\mathbf{E}_{\mathbf{Q}}\{(\sum_{u \in \mathcal{J}_1^{(1)}} e^{-s/2V(u)})^{\rho(s)}\} < \infty$ . Therefore,  $\mathbf{E}_{\mathbf{Q}}\{\hat{\eta}_1^{\rho(s)}\} < \infty$ . We are thus entitled to apply Lemma 6.1 (stated and proved below) to see that  $\mathbf{E}_{\mathbf{Q}}^{(n)}\left\{\frac{1}{1 + \sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j}\right\} \geq c_{75}$  for some  $c_{75} \in (0, \infty)$  and all  $n \geq n_0$ . Since  $\frac{1}{\sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j} \geq \frac{1}{1 + \sum_{j=1}^{n+1} e^{-tS_{j-1}} \hat{\eta}_j}$ , this yields

$$\mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} \geq c_{74} \varepsilon [c_{75} - \mathbf{Q}^{(n)}\{S_n \leq \varepsilon n^{1/2}\}], \quad n \geq n_0.$$



On the other hand,  $S_n/n^{1/2}$  under  $\mathbf{Q}^{(n)}$  converges weakly to the terminal value of a Brownian meander (see Bolthausen [11]); in particular,  $\lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \mathbf{Q}^{(n)}\{S_n \leq \varepsilon n^{1/2}\} = 0$ . We can thus choose (and fix) a small  $\varepsilon > 0$  such that  $\mathbf{Q}^{(n)}\{S_n \leq \varepsilon n^{1/2}\} \leq \frac{c_{75}}{2}$  for all  $n \geq n_1$ . Therefore, for  $n \geq n_0 + n_1$ ,

$$\mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} \geq c_{74}\varepsilon \left[ c_{75} - \frac{c_{75}}{2} \right].$$

As a consequence, we have proved that, for  $0 < s < 1$ ,

$$\liminf_{n \rightarrow \infty} \mathbf{E}_{\mathbf{Q}}\{(n^{1/2} \wedge V(w_n^{(n)})^+) \underline{Y}_n^{-s}\} > 0,$$

which, in view of (6.10), yields the first inequality in (1.16), and thus completes the proof of the lower bound in Theorem 1.5.  $\square$

We complete the proof of Theorem 1.5 by proving the following lemma, which is a very simple variant of a result of Kozlov [24].

LEMMA 6.1. *Let  $\{(X_k, \eta_k), k \geq 1\}$  be a sequence of i.i.d. random vectors defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  with  $\mathbb{P}\{\eta_1 \geq 0\} = 1$ , such that  $\mathbb{E}\{\eta_1^\theta\} < \infty$  for some  $\theta > 0$ . We assume  $\mathbb{E}(X_1) = 0$  and  $0 < \mathbb{E}(X_1^2) < \infty$ . Let  $S_0 := 0$  and  $S_n := X_1 + \dots + X_n$ , for  $n \geq 1$ . Then*

$$(6.11) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left\{ \frac{1}{1 + \sum_{k=1}^{n+1} \eta_k e^{-S_{k-1}}} \middle| \min_{1 \leq k \leq n} S_k > 0 \right\} = c_{76} \in (0, \infty).$$

PROOF. The lemma is an analogue of the identity (26) of Kozlov [24], except that the distribution of our  $\eta_1$  is slightly different from that of Kozlov’s, which explains the moment condition  $\mathbb{E}\{\eta_1^\theta\} < \infty$ : this condition will be seen to guarantee

$$(6.12) \quad \lim_{j \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbb{E} \left\{ \frac{1}{1 + \sum_{k=1}^j \eta_k e^{-S_{k-1}}} - \frac{1}{1 + \sum_{k=1}^{n+1} \eta_k e^{-S_{k-1}}} \middle| \min_{1 \leq k \leq n} S_k > 0 \right\} = 0.$$

The identity (6.12), which plays the role of Kozlov’s Lemma 1 in [24], is the key ingredient in the proof of (6.11). Since the rest of the proof goes along the lines of [24] with obvious modifications, we only prove (6.12) here.

Without loss of generality, we assume  $\theta \leq 2$  (otherwise, we can replace  $\theta$  by 2). We observe that, for  $n > j$ , the integrand in (6.12) is nonnegative, and is

$$\leq \frac{\sum_{k=j+1}^{n+1} \eta_k e^{-S_{k-1}}}{1 + \sum_{k=1}^{n+1} \eta_k e^{-S_{k-1}}} \leq \left( \frac{\sum_{k=j+1}^{n+1} \eta_k e^{-S_{k-1}}}{1 + \sum_{k=1}^{n+1} \eta_k e^{-S_{k-1}}} \right)^{\theta/2} \leq \left( \sum_{k=j+1}^{n+1} \eta_k e^{-S_{k-1}} \right)^{\theta/2},$$

which is bounded by  $\sum_{k=j+1}^{n+1} \eta_k^{\theta/2} e^{-\theta/2 S_{k-1}}$ . Since  $\mathbb{P}\{\min_{1 \leq k \leq n} S_k > 0\} \sim c_4/n^{1/2}$  [see (2.13)], we only need to check that

$$(6.13) \quad \lim_{j \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{\eta_k^{\theta/2} e^{-\theta/2 S_{k-1}} \mathbf{1}_{\{\min_{1 \leq i \leq n} S_i > 0\}}\} = 0.$$

Let  $\text{LHS}_{(6.13)}$  denote the  $n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{\cdot\}$  expression on the left-hand side. Let  $\widehat{S}_i = \widehat{S}_i(k) := S_{i+k} - S_k$ ,  $i \geq 0$ . It is clear that  $(\widehat{S}_i, i \geq 0)$  is independent of  $(\eta_k, X_1, \dots, X_k)$ , and is distributed as  $(S_i, i \geq 0)$ . Write  $\underline{S}_{k-1} := \min_{1 \leq j \leq k-1} S_j$  and  $\widehat{\underline{S}}_{n-k} := \min_{1 \leq i \leq n-k} \widehat{S}_i$ . Then

$$\text{LHS}_{(6.13)} \leq n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{\eta_k^{\theta/2} e^{-\theta/2 S_{k-1}} \mathbf{1}_{\{\underline{S}_{k-1} > 0, \widehat{\underline{S}}_{n-k} > -S_{k-1} - X_k\}}\}.$$

To estimate  $\mathbb{E}\{\cdot\}$  on the right-hand side, we first condition upon  $(\eta_k, S_{k-1}, \underline{S}_{k-1}, X_k)$ , which leaves us to estimate the tail probability of  $\widehat{\underline{S}}_{n-k}$ . At this stage, it is convenient to recall (see (13) of Kozlov [24]) that  $\mathbb{P}\{\widehat{\underline{S}}_{n-k} > -y\} \leq c_{54} \frac{1+y^+}{(n-k+1)^{1/2}}$  for some  $c_{54} > 0$  and all  $y \in \mathbb{R}$ . Accordingly,

$$\begin{aligned} \text{LHS}_{(6.13)} &\leq c_{54} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\left\{\eta_k^{\theta/2} e^{-\theta/2 S_{k-1}} \mathbf{1}_{\{\underline{S}_{k-1} > 0\}} \frac{1 + (S_{k-1} + X_k)^+}{(n-k+1)^{1/2}}\right\} \\ &\leq c_{54} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\left\{\eta_k^{\theta/2} e^{-\theta/2 S_{k-1}} \mathbf{1}_{\{\underline{S}_{k-1} > 0\}} \frac{1 + S_{k-1} + X_k^+}{(n-k+1)^{1/2}}\right\}. \end{aligned}$$

On the right-hand side,  $(\eta_k, X_k)$  is independent of  $(\underline{S}_{k-1}, S_{k-1})$ . We condition upon  $(\underline{S}_{k-1}, S_{k-1})$ : for any  $z \geq 1$ , an application of the Cauchy–Schwarz inequality gives

$$\mathbb{E}\{\eta_k^{\theta/2} (z + X_k^+)\} \leq [\mathbb{E}(\eta_k^\theta)]^{1/2} [\mathbb{E}\{(z + X_k^+)^2\}]^{1/2}.$$

Of course,  $\mathbb{E}(\eta_k^\theta) = \mathbb{E}(\eta_1^\theta) < \infty$  by assumption, and  $\mathbb{E}\{(z + X_k^+)^2\} \leq 2\mathbb{E}(z^2 + X_k^2) = 2[z^2 + \mathbb{E}(X_1^2)]$ . Thus,  $\mathbb{E}\{\eta_k^{\theta/2} (z + X_k^+)\} \leq c_{77} z$  for  $z \geq 1$ . Consequently, with  $c_{78} := c_{54} c_{77}$ ,

$$\begin{aligned} \text{LHS}_{(6.13)} &\leq c_{78} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\left\{e^{-\theta/2 S_{k-1}} \mathbf{1}_{\{\underline{S}_{k-1} > 0\}} \frac{1 + S_{k-1}}{(n-k+1)^{1/2}}\right\} \\ &\leq c_{79} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\left\{e^{-\theta/3 S_{k-1}} \mathbf{1}_{\{\underline{S}_{k-1} > 0\}} \frac{1}{(n-k+1)^{1/2}}\right\}, \end{aligned}$$

the last inequality following from the fact that  $\sup_{x>0} (1+x)e^{-\theta/6x} < \infty$ .

We use once again the estimate (2.13), which implies  $\frac{1}{(n-k+1)^{1/2}} \leq c_{80} \mathbb{P}\{S_i > S_{k-1}, \forall k \leq i \leq n\}$ . Since  $(S_i - S_{k-1}, k \leq i \leq n)$  is independent of  $(S_{k-1}, \underline{S}_{k-1})$ , this implies, with  $c_{81} := c_{79}c_{80}$ ,

$$\begin{aligned} \text{LHS}_{(6.13)} &\leq c_{81}n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{S_{k-1}>0, S_i>S_{k-1}, \forall k \leq i \leq n\}}\} \\ &\leq c_{81}n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{\underline{S}_n>0\}}\}, \end{aligned}$$

where  $\underline{S}_n := \min_{1 \leq i \leq n} S_i$ . It remains to check that

$$(6.14) \quad \lim_{j \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{1/2} \sum_{k=j+1}^{n+1} \mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{\underline{S}_n>0\}}\} = 0.$$

This would immediately follow from Lemma 1 of Kozlov [24], but we have been kindly informed by Gerold Alsmeyer (to whom we are grateful) of a flaw in its proof, on page 800, line 3 of [24], so we need to proceed differently. Since  $\mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{\underline{S}_n>0\}}\} \leq n^{-(3/2)+o(1)}(n-k+2)^{-1/2}$  (for  $n \rightarrow \infty$ ) uniformly in  $k \in [\frac{n}{2}, n+1]$ , we have  $n^{1/2} \sum_{k=\lfloor n/2 \rfloor}^{n+1} \mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{\underline{S}_n>0\}}\} \rightarrow 0, n \rightarrow \infty$ . On the other hand, (36) of Kozlov [24] (applied to  $\delta = \frac{1}{2}$  and  $\eta_i = 1$  there) implies that  $\lim_j \limsup_n n^{1/2} \sum_{k=j+1}^{\lfloor n/2 \rfloor} \mathbb{E}\{e^{-\theta/3S_{k-1}} \mathbf{1}_{\{\underline{S}_n>0\}}\} = 0$ . Therefore, (6.14) holds: Lemma 6.1 is proved.  $\square$

**7. Proof of Theorem 1.3 and (1.14)–(1.15) of Theorem 1.4.** In this section we prove Theorem 1.3, as well as parts (1.14)–(1.15) of Theorem 1.4. We assume (1.1), (1.2) and (1.3) throughout the section.

**PROOF OF THEOREM 1.3 AND (1.14) AND (1.15) OF THEOREM 1.4.** *Upper bounds.* Let  $\varepsilon > 0$ . By Theorem 1.6 and Chebyshev’s inequality,  $\mathbf{P}\{W_{n,\beta} > n^{-(3\beta/2)+\varepsilon}\} \rightarrow 0$ . Therefore,  $W_{n,\beta} \leq n^{-(3\beta/2)+o(1)}$  in probability, yielding the upper bound in (1.15).

The upper bound in (1.14) follows trivially from the upper bound in (1.15).

It remains to prove the upper bound in Theorem 1.3. Fix  $\gamma \in (0, 1)$ . Since  $W_n^\gamma$  is a nonnegative supermartingale, the maximal inequality tells that, for any  $n \leq m$  and any  $\lambda > 0$ ,

$$\mathbf{P}\left\{\max_{n \leq j \leq m} W_j^\gamma \geq \lambda\right\} \leq \frac{\mathbf{E}(W_n^\gamma)}{\lambda} \leq \frac{c_{82}}{\lambda n^{\gamma/2}},$$

the last inequality being a consequence of Theorem 1.5. Let  $\varepsilon > 0$  and let  $n_k := \lfloor k^{2/\varepsilon} \rfloor$ . Then  $\sum_k \mathbf{P}\{\max_{n_k \leq j \leq n_{k+1}} W_j^\gamma \geq n_k^{-(\gamma/2)+\varepsilon}\} < \infty$ . By the Borel–Cantelli lemma, almost surely for all large  $k$ ,  $\max_{n_k \leq j \leq n_{k+1}} W_j < n_k^{-(1/2)+(\varepsilon/\gamma)}$ . Since

$\frac{\varepsilon}{\gamma}$  can be arbitrarily small, this yields the desired upper bound:  $W_n \leq n^{-(1/2)+o(1)}$  a.s.  $\square$

PROOF OF THEOREM 1.3 AND (1.14) AND (1.15) OF THEOREM 1.4. *Lower bounds.* To prove the lower bound in (1.14) and (1.15), we use the Paley–Zygmund inequality and Theorem 1.6 to see that

$$(7.1) \quad \mathbf{P}\{W_{n,\beta} > n^{-(3\beta/2)+o(1)}\} \geq n^{o(1)}, \quad n \rightarrow \infty.$$

This is the analogue of (4.5) for  $W_n$ . From here, the argument follows the lines in the proof of the upper bound in (1.8) of Theorem 1.2 (Section 4), and goes as follows: let  $\varepsilon > 0$  and let  $\tau_n := \inf\{k \geq 1 : \#\{u : |u| = k\} \geq n^{2\varepsilon}\}$ . Then

$$\begin{aligned} & \mathbf{P}\left\{\tau_n < \infty, \min_{k \in [n/2, n]} W_{k+\tau_n, \beta} \leq n^{-(3\beta/2)-\varepsilon} \exp\left[-\beta \max_{|x|=\tau_n} V(x)\right]\right\} \\ & \leq \sum_{k \in [n/2, n]} \mathbf{P}\left\{\tau_n < \infty, W_{k+\tau_n, \beta} \leq n^{-(3\beta/2)-\varepsilon} \exp\left[-\beta \max_{|x|=\tau_n} V(x)\right]\right\} \\ & \leq \sum_{k \in [n/2, n]} (\mathbf{P}\{W_{k, \beta} \leq n^{-(3\beta/2)-\varepsilon}\})^{\lfloor n^{2\varepsilon} \rfloor}, \end{aligned}$$

which, according to (7.1), is bounded by  $n \exp(-n^{-\varepsilon} \lfloor n^{2\varepsilon} \rfloor)$  (for all sufficiently large  $n$ ), thus summable in  $n$ . By the Borel–Cantelli lemma, almost surely for all sufficiently large  $n$ , we have either  $\tau_n = \infty$ , or  $\min_{k \in [n/2, n]} W_{k+\tau_n, \beta} > n^{-(3\beta/2)-\varepsilon} \exp[-\beta \max_{|x|=\tau_n} V(x)]$ . Conditionally on the system’s ultimate survival, we have  $\frac{1}{n} \max_{|x|=n} V(x) \rightarrow c_{21}$  a.s.,  $\tau_n \sim \frac{2\varepsilon \log n}{\log m}$  a.s.,  $n \rightarrow \infty$ , and  $W_{n, \beta} \geq \min_{k \in [n/2, n]} W_{k+\tau_n, \beta}$  for all sufficiently large  $n$ . This readily yields lower bounds in (1.14) and (1.15): conditionally on the system’s survival,  $W_{n, \beta} \geq n^{-(3\beta/2)+o(1)}$  almost surely (and a fortiori, in probability).

The lower bound in Theorem 1.3 is along exactly the same lines, but using Theorem 1.5 instead of Theorem 1.6.  $\square$

**8. Proof of Theorem 1.2.** Assume (1.1), (1.2) and (1.3). Let  $\beta > 1$ . We trivially have  $W_{n, \beta} \leq W_n \exp\{-(\beta - 1) \inf_{|u|=n} V(u)\}$  and  $W_{n, \beta} \geq \exp\{-\beta \times \inf_{|u|=n} V(u)\}$ . Therefore,  $\frac{1}{\beta} \log \frac{1}{W_{n, \beta}} \leq \inf_{|u|=n} V(u) \leq \frac{1}{\beta-1} \log \frac{W_n}{W_{n, \beta}}$  on  $\mathcal{S}_n$ . Since  $\beta$  can be as large as possible, by means of Theorem 1.3 and of parts (1.14) and (1.15) of Theorem 1.4, we immediately get (1.7) and (1.9).

Since  $W_n \geq \exp\{-\inf_{|u|=n} V(u)\}$ , the lower bound in (1.8) follows immediately from Theorem 1.3, whereas the upper bound in (1.8) was already proved in Section 4.

**9. Proof of part (1.13) of Theorem 1.4.** The upper bound follows from Theorem 1.3 and the elementary inequality  $W_{n, \beta} \leq W_n^\beta$ , the lower bound from (1.8) and the relation  $W_{n, \beta} \geq \exp\{-\beta \inf_{|u|=n} V(u)\}$ .

**10. Proof of Theorem 1.1.** The proof of Theorem 1.1 relies on Theorem 1.5 and a preliminary result, stated below as Proposition 10.1. Theorem 1.5 ensures the tightness of  $(n^{1/2}W_n, n \geq 1)$ , whereas Proposition 10.1 implies that  $\frac{W_{n+1}}{W_n}$  converges to 1 in probability (conditionally on the system’s survival).

PROPOSITION 10.1. *Assume (1.1), (1.2) and (1.3). For any  $\gamma > 0$ , there exists  $\gamma_1 > 0$  such that, for all sufficiently large  $n$ ,*

$$(10.1) \quad \mathbf{P}\left\{\left|\frac{W_{n+1}}{W_n} - 1\right| \geq n^{-\gamma} \mid \mathcal{S}\right\} \leq n^{-\gamma_1}.$$

PROOF. Let  $1 < \beta \leq \min\{2, 1 + \rho(1)\}$ , where  $\rho(1)$  is the constant in Corollary 2.4.

We use a probability estimate of Petrov [34], page 82: for centered random variables  $\xi_1, \dots, \xi_\ell$  with  $\mathbf{E}\{|\xi_i|^\beta\} < \infty$  (for  $1 \leq i \leq \ell$ ), we have  $\mathbf{E}\{|\sum_{i=1}^\ell \xi_i|^\beta\} \leq 2 \sum_{i=1}^\ell \mathbf{E}\{|\xi_i|^\beta\}$ .

By definition, on the set  $\mathcal{S}_n$ , we have

$$\frac{W_{n+1}}{W_n} - 1 = \sum_{|u|=n} \frac{e^{-V(u)}}{W_n} \left( \sum_{x \in \mathbb{T}_u^{\text{GW}}: |x|_u=1} e^{-V_u(x)} - 1 \right),$$

where  $\mathbb{T}^{\text{GW}}$  and  $|x|_u$  are as in (2.1) and (2.4), respectively. Conditioning on  $\mathcal{F}_n$ , and applying Proposition 2.1 and Petrov’s probability inequality recalled above, we see that, on  $\mathcal{S}_n$ ,

$$(10.2) \quad \begin{aligned} \mathbf{E}\left\{\left|\frac{W_{n+1}}{W_n} - 1\right|^\beta \mid \mathcal{F}_n\right\} &\leq 2 \sum_{|u|=n} \frac{e^{-\beta V(u)}}{W_n^\beta} \mathbf{E}\left\{\left|\sum_{|y|=1} e^{-V(y)} - 1\right|^\beta\right\} \\ &= c_{83} \frac{W_{n,\beta}}{W_n^\beta}, \end{aligned}$$

where  $c_{83} := 2\mathbf{E}\{|\sum_{|v|=1} e^{-V(v)} - 1|^\beta\} < \infty$  [see (2.16)], and  $W_{n,\beta}$  is as in (1.11).

Let  $\varepsilon > 0$  and  $b > 0$ . Let  $s \in (\frac{\beta-1}{\beta}, 1)$ . Define  $D_n := \{W_n \geq n^{-(1/2)-\varepsilon}\} \cap \{W_{n,\beta} \leq n^{-(3\beta/2)+b}\}$ . By Proposition 3.1,  $\mathbf{P}\{W_n < n^{-(1/2)-\varepsilon}, \mathcal{S}\} \leq n^{-\vartheta}$  for some  $\vartheta > 0$  and all large  $n$ , whereas, by Theorem 1.6,  $\mathbf{P}\{W_{n,\beta} > n^{-(3\beta/2)+b}\} \leq n^{3\beta(1-s)/2 - (1-s)b} \mathbf{E}\{W_{n,\beta}^{1-s}\} = n^{-(1-s)b+o(1)}$ . Therefore,

$$\mathbf{P}\{\mathcal{S} \setminus D_n\} \leq n^{-\vartheta} + n^{-(1-s)b+o(1)}, \quad n \rightarrow \infty.$$

On the other hand, since  $\mathcal{S} \subset \mathcal{S}_n$ , it follows from (10.2) and Chebyshev’s inequality that, for  $n \rightarrow \infty$ ,

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{W_{n+1}}{W_n} - 1\right| \geq n^{-\gamma}, D_n, \mathcal{S}\right\} &\leq n^{\gamma\beta} \mathbf{E}\left\{c_{83} \frac{W_{n,\beta}}{W_n^\beta} \mathbf{1}_{D_n \cap \mathcal{S}_n}\right\} \\ &\leq c_{83} n^{\gamma\beta - (3\beta/2) + b + [(1/2) + \varepsilon]\beta}. \end{aligned}$$

As a consequence, when  $n \rightarrow \infty$ ,

$$\mathbf{P}\left\{\left|\frac{W_{n+1}}{W_n} - 1\right| \geq n^{-\gamma}, \mathcal{S}\right\} \leq n^{-\vartheta} + n^{-(1-s)b+o(1)} + c_{83}n^{\gamma\beta-\beta+b+\varepsilon\beta}.$$

We choose  $\varepsilon$  and  $b$  sufficiently small such that  $\gamma\beta - \beta + b + \varepsilon\beta < 0$ . Proposition 10.1 is proved.  $\square$

We now have all of the ingredients needed for the proof of Theorem 1.1.

PROOF OF THEOREM 1.1. Once Proposition 10.1 is established, the proof of Theorem 1.1 follows the lines of Biggins and Kyprianou [7].

Assume (1.1), (1.2) and (1.3). Let  $\lambda_n > 0$  satisfy  $\mathbf{E}\{(\lambda_n W_n)^{1/2}\} = 1$ . That is,

$$\lambda_n := \{\mathbf{E}(W_n^{1/2})\}^{-2}.$$

By Theorem 1.5, we have  $0 < \liminf_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} \leq \limsup_{n \rightarrow \infty} \frac{\lambda_n}{n^{1/2}} < \infty$ , and  $(\lambda_n W_n)$  is tight. Let  $\widehat{\mathcal{W}}$  be any possible (weak) limit of  $(\lambda_n W_n)$  along a subsequence. By Theorem 1.5 and dominated convergence,  $\mathbf{E}(\widehat{\mathcal{W}}^{1/2}) = 1$ . We now prove the uniqueness of  $\widehat{\mathcal{W}}$ .

By definition,

$$W_{n+1} = \sum_{|v|=1} e^{-V(v)} \sum_{x \in \mathbb{T}_v^{\text{GW}}, |x|_v=n} e^{-V_v(x)}.$$

By assumption,  $\lambda_n W_n \rightarrow \widehat{\mathcal{W}}$  in distribution when  $n$  goes to infinity along a certain subsequence. Thus,  $\lambda_n W_{n+1}$  converges weakly (when  $n$  goes along the same subsequence) to  $\sum_{|v|=1} e^{-V(v)} \widehat{\mathcal{W}}_v$ , where, conditionally on  $(v, V(v), |v|=1)$ ,  $\widehat{\mathcal{W}}_v$  are independent copies of  $\widehat{\mathcal{W}}$ .

On the other hand, by Proposition 10.1,  $\lambda_n W_{n+1}$  also converges weakly (along the same subsequence) to  $\widehat{\mathcal{W}}$ . Therefore,

$$\widehat{\mathcal{W}} \stackrel{\text{law}}{=} \sum_{|v|=1} e^{-V(v)} \widehat{\mathcal{W}}_v.$$

This is the same equation for  $\xi^*$  in (3.5). Recall that (3.5) has a unique solution up to a scale change (Liu [27]), and since  $\mathbf{E}(\widehat{\mathcal{W}}^{1/2}) = 1$ , we have  $\widehat{\mathcal{W}} \stackrel{\text{law}}{=} c_{84}\xi^*$ , with  $c_{84} := [\mathbf{E}\{(\xi^*)^{1/2}\}]^{-2}$ . The uniqueness (in law) of  $\widehat{\mathcal{W}}$  shows that  $\lambda_n W_n$  converges weakly to  $\widehat{\mathcal{W}}$  when  $n \rightarrow \infty$ .

By (3.3),  $\mathbf{P}\{W_n > 0\} = \mathbf{P}\{\mathcal{S}_n\} \rightarrow \mathbf{P}\{\mathcal{S}\} = \mathbf{P}\{\xi^* > 0\}$ . Let  $\mathcal{W} > 0$  be a random variable such that

$$(10.3) \quad \mathbf{E}(e^{-a\mathcal{W}}) = \mathbf{E}(e^{-a\widehat{\mathcal{W}}} | \widehat{\mathcal{W}} > 0), \quad \forall a \geq 0.$$

It follows that, conditionally on the system’s survival,  $\lambda_n W_n$  converges in distribution to  $\mathcal{W}$ .  $\square$

**Acknowledgments.** We are grateful to Bruno Jaffuel, Gerold Alsmeyer, John Biggins and Alain Rouault for pointing out errors in the first versions of the manuscript, and to John Biggins for fixing Lemma 3.2. We also wish to thank two referees for their careful reading and for very helpful suggestions, without which we would not have had enough courage to remove the uniform ellipticity condition in the original manuscript.

## REFERENCES

- [1] ADDARIO-BERRY, L. (2006). Ballot theorems and the heights of trees. Ph.D. thesis, McGill Univ.
- [2] ALDOUS, D. J. and BANDYOPADHYAY, A. (2005). A survey of max-type recursive distributional equations. *Ann. Appl. Probab.* **15** 1047–1110. [MR2134098](#)
- [3] ATHREYA, K. B. and NEY, P. E. (1972). *Branching Processes. Die Grundlehren der mathematischen Wissenschaften* **196**. Springer, New York. [MR0373040](#)
- [4] BACHMANN, M. (2000). Limit theorems for the minimal position in a branching random walk with independent logconcave displacements. *Adv. in Appl. Probab.* **32** 159–176. [MR1765165](#)
- [5] BIGGINS, J. D. (1976). The first- and last-birth problems for a multitype age-dependent branching process. *Adv. in Appl. Probab.* **8** 446–459. [MR0420890](#)
- [6] BIGGINS, J. D. and GREY, D. R. (1979). Continuity of limit random variables in the branching random walk. *J. Appl. Probab.* **16** 740–749. [MR549554](#)
- [7] BIGGINS, J. D. and KYPRIANOU, A. E. (1997). Seneta-Heyde norming in the branching random walk. *Ann. Probab.* **25** 337–360. [MR1428512](#)
- [8] BIGGINS, J. D. and KYPRIANOU, A. E. (2004). Measure change in multitype branching. *Adv. in Appl. Probab.* **36** 544–581. [MR2058149](#)
- [9] BIGGINS, J. D. and KYPRIANOU, A. E. (2005). Fixed points of the smoothing transform: The boundary case. *Electron. J. Probab.* **10** 609–631 (electronic). [MR2147319](#)
- [10] BINGHAM, N. H. (1973). Limit theorems in fluctuation theory. *Adv. in Appl. Probab.* **5** 554–569. [MR0348843](#)
- [11] BOLTHAUSEN, E. (1989). A central limit theorem for two-dimensional random walks in random sceneries. *Ann. Probab.* **17** 108–115. [MR972774](#)
- [12] BRAMSON, M. D. (1978). Maximal displacement of branching Brownian motion. *Comm. Pure Appl. Math.* **31** 531–581. [MR0494541](#)
- [13] BRAMSON, M. D. (1978). Minimal displacement of branching random walk. *Z. Wahrsch. Verw. Gebiete* **45** 89–108. [MR510529](#)
- [14] BRAMSON, M. D. and ZEITOUNI, O. (2009). Tightness for a family of recursion equations. *Ann. Probab.* **37** 615–653.
- [15] CHAUVIN, B., ROUAULT, A. and WAKOLBINGER, A. (1991). Growing conditioned trees. *Stochastic Process. Appl.* **39** 117–130. [MR1135089](#)
- [16] DEKKING, F. M. and HOST, B. (1991). Limit distributions for minimal displacement of branching random walks. *Probab. Theory Related Fields* **90** 403–426. [MR1133373](#)
- [17] DERRIDA, B. and SPOHN, H. (1988). Polymers on disordered trees, spin glasses, and traveling waves. *J. Statist. Phys.* **51** 817–840. [MR971033](#)
- [18] FELLER, W. (1971). *An Introduction to Probability Theory and Its Applications. Vol. II*, 2nd ed. Wiley, New York. [MR0270403](#)
- [19] HAMMERSLEY, J. M. (1974). Postulates for subadditive processes. *Ann. Probab.* **2** 652–680. [MR0370721](#)
- [20] HARDY, R. and HARRIS, S. C. (2004). A new formulation of the spine approach to branching diffusions. Mathematics Preprint, Univ. Bath.

- [21] HARRIS, T. E. (1963). *The Theory of Branching Processes. Die Grundlehren der Mathematischen Wissenschaften* **119**. Springer, Berlin. [MR0163361](#)
- [22] HEYDE, C. C. (1970). Extension of a result of Seneta for the super-critical Galton–Watson process. *Ann. Math. Statist.* **41** 739–742. [MR0254929](#)
- [23] KINGMAN, J. F. C. (1975). The first birth problem for an age-dependent branching process. *Ann. Probab.* **3** 790–801. [MR0400438](#)
- [24] KOZLOV, M. V. (1976). The asymptotic behavior of the probability of non-extinction of critical branching processes in a random environment. *Teor. Veroyatnost. i Primenen.* **21** 813–825. [MR0428492](#)
- [25] KYPRIANOU, A. E. (1998). Slow variation and uniqueness of solutions to the functional equation in the branching random walk. *J. Appl. Probab.* **35** 795–801. [MR1671230](#)
- [26] LIFSHITS, M. A. (2007). Some limit theorems on binary trees. (In preparation.)
- [27] LIU, Q. S. (2000). On generalized multiplicative cascades. *Stochastic Process. Appl.* **86** 263–286. [MR1741808](#)
- [28] LIU, Q. S. (2001). Asymptotic properties and absolute continuity of laws stable by random weighted mean. *Stochastic Process. Appl.* **95** 83–107. [MR1847093](#)
- [29] LYONS, R. (1997). A simple path to Biggins’ martingale convergence for branching random walk. In *Classical and Modern Branching Processes (Minneapolis, MN, 1994)*. *IMA Vol. Math. Appl.* **84** 217–221. Springer, New York. [MR1601749](#)
- [30] LYONS, R., PEMANTLE, R. and PERES, Y. (1995). Conceptual proofs of  $L \log L$  criteria for mean behavior of branching processes. *Ann. Probab.* **23** 1125–1138. [MR1349164](#)
- [31] MCDIARMID, C. (1995). Minimal positions in a branching random walk. *Ann. Appl. Probab.* **5** 128–139. [MR1325045](#)
- [32] NEVEU, J. (1986). Arbres et processus de Galton–Watson. *Ann. Inst. H. Poincaré Probab. Statist.* **22** 199–207. [MR850756](#)
- [33] NEVEU, J. (1988). Multiplicative martingales for spatial branching processes. In *Seminar on Stochastic Processes, 1987 (Princeton, NJ, 1987)*. *Progr. Probab. Statist.* **15** 223–242. Birkhäuser Boston, Boston. [MR1046418](#)
- [34] PETROV, V. V. (1995). *Limit Theorems of Probability Theory: Sequences of independent random variables*. *Oxford Studies in Probability* **4**. Clarendon, Oxford. [MR1353441](#)
- [35] SENETA, E. (1968). On recent theorems concerning the supercritical Galton–Watson process. *Ann. Math. Statist.* **39** 2098–2102. [MR0234530](#)

DÉPARTEMENT DE MATHÉMATIQUES  
UNIVERSITÉ PARIS XIII  
99 AVENUE J-B CLÉMENT  
F-93430 VILLETANEUSE  
FRANCE  
E-MAIL: [yueyun@math.univ-paris13.fr](mailto:yueyun@math.univ-paris13.fr)

LABORATOIRE DE PROBABILITÉS  
ET MODÈLES ALÉATOIRES  
UNIVERSITÉ PARIS VI  
4 PLACE JUSSIEU  
F-75252 PARIS CEDEX 05  
FRANCE  
E-MAIL: [zhan@proba.jussieu.fr](mailto:zhan@proba.jussieu.fr)