

Minimal surfaces and weak gravity

Mehmet Demirtas,^a Cody Long,^b Liam McAllister^a and Mike Stillman^c

^a*Department of Physics, Cornell University,
East Avenue, Ithaca, NY 14853, U.S.A.*

^b*Department of Physics, Northeastern University,
110 Forsyth St, Boston, MA 02115, U.S.A.*

^c*Department of Mathematics, Cornell University,
Tower Road, Ithaca, NY 14853, U.S.A.*

E-mail: md775@cornell.edu, co.long@northeastern.edu,
mcallister@cornell.edu, mike@math.cornell.edu

ABSTRACT: We show that the Weak Gravity Conjecture (WGC) implies a nontrivial upper bound on the volumes of the minimal-volume cycles in certain homology classes that admit no calibrated representatives. In compactification of type IIB string theory on an orientifold X of a Calabi-Yau threefold, we consider a homology class $[\Sigma] \in H_4(X, \mathbb{Z})$ represented by a union Σ_{\cup} of holomorphic and antiholomorphic cycles. The instanton form of the WGC applied to the axion charge $[\Sigma]$ implies an upper bound on the action of a non-BPS Euclidean D3-brane wrapping the minimal-volume representative Σ_{\min} of $[\Sigma]$. We give an explicit example of an orientifold X of a hypersurface in a toric variety, and a hyperplane $\mathcal{H} \subset H_4(X, \mathbb{Z})$, such that for any $[\Sigma] \in \mathcal{H}$ that satisfies the WGC, the minimal volume obeys $\text{Vol}(\Sigma_{\min}) \ll \text{Vol}(\Sigma_{\cup})$: the holomorphic and antiholomorphic components recombine to form a much smaller cycle. In particular, the sub-Lattice WGC applied to X implies large recombination, no matter how sparse the sublattice. Non-BPS instantons wrapping Σ_{\min} are then more important than would be predicted from a study of BPS instantons wrapping the separate components of Σ_{\cup} . Our analysis hinges on a novel computation of effective divisors in X that are not inherited from effective divisors of the toric variety.

KEYWORDS: Flux compactifications, D-branes, Solitons Monopoles and Instantons

ARXIV EPRINT: [1906.08262](https://arxiv.org/abs/1906.08262)

Contents

1	Introduction	1
2	Axions, orientifolds, and weak gravity	5
3	An orientifold where weak gravity implies large recombination	7
4	Discussion	11
A	Geometry of the example	14
B	Effective divisors in the example	16
	B.1 Cohomology of line bundles on toric varieties	17
	B.2 The explicit example	18

1 Introduction

In quantum theories of extended objects, such as string theories, there can be contributions to the path integral from extended objects wrapping cycles in spacetime. Famous examples include worldsheet instantons, in which the Euclidean worldsheet of a string wraps a two-cycle, and Dp -brane instantons, in which the Euclidean worldvolume of a Dp -brane wraps a $p + 1$ -cycle.

Computing instanton contributions to the four-dimensional action is an important problem, doubly so in theories in which the leading interactions of a given type are produced by instantons. For example, in a compactification of type IIB string theory on an orientifold X of a Calabi-Yau threefold, the axions that result from reduction of the Ramond-Ramond four-form C_4 on four-cycles in X have no non-derivative interactions at any order in perturbation theory, but acquire a potential from Euclidean D3-branes wrapping four-cycles [1].

The semiclassical action of a Euclidean D3-brane includes a term proportional to the volume of the wrapped cycle, and so the Euclidean D3-brane will wrap a cycle that is at least locally volume-minimizing in its homology class.¹ Thus, to compute instanton effects in the four-dimensional action, a first step is to compute the volumes of minimal cycles. That is, one should find, for each class $[\Sigma] \in H_4(X, \mathbb{Z})$, the minimum-volume representative $\Sigma_{\min} \in [\Sigma]$, as well as its volume $\text{Vol}(\Sigma_{\min})$. This is still a far cry from an exact computation of the potential from instantons, which would require a more refined study of the other degrees of freedom on the instanton worldvolume, including a counting of fermion zero modes. Even so, computing $\text{Vol}(\Sigma_{\min})$ gives insight about the relative importance of *possible* instanton contributions.

¹We neglect for the moment the effects of the worldvolume gauge field, cf. [2].

When X is a Kähler threefold, the computation of $\text{Vol}(\Sigma_{\min})$ is almost trivial for a special set of classes $[\Sigma] \in H_4(X, \mathbb{Z})$: if $[\Sigma]$ can be represented by an effective divisor $\Sigma_{\mathcal{E}}$, then $\Sigma_{\mathcal{E}}$ is calibrated by the Kähler form J , and is absolutely volume-minimizing in its class, with volume

$$\text{Vol}(\Sigma_{\mathcal{E}}) = \text{Vol}(\Sigma_{\min}) = \frac{1}{2} \int_{\Sigma_{\mathcal{E}}} J \wedge J, \tag{1.1}$$

which is readily evaluated in terms of the Kähler parameters and intersection numbers of X . One can similarly compute $\text{Vol}(\Sigma_{\min})$ when $[\Sigma]$ can be represented by an anti-effective divisor² $\Sigma_{\bar{\mathcal{E}}}$:

$$\text{Vol}(\Sigma_{\bar{\mathcal{E}}}) = \text{Vol}(\Sigma_{\min}) = -\frac{1}{2} \int_{\Sigma_{\bar{\mathcal{E}}}} J \wedge J. \tag{1.2}$$

Thus, for any effective or anti-effective divisor, the semiclassical action is easily computed, in the above approximation.

However, many classes in $H_4(X, \mathbb{Z})$ are neither effective nor anti-effective! Computing the minimum-volume representative is then highly nontrivial: in fact, it is an instance of one of the fundamental problems in geometric measure theory, the *Plateau problem*, which we now review in slightly more general terms.

Suppose that one is given a Riemannian manifold M of real dimension n , with fixed metric g . Given a homology class $[\Sigma] \in H_p(M, \mathbb{Z})$, $0 < p < n$, what representative of $[\Sigma]$ has minimal volume? A crucial subtlety is that the volume functional may not attain a minimum on any *smooth* representative of $[\Sigma]$. To make the variational problem well-posed, one should — heuristically — include appropriate limit points corresponding to mildly singular representatives.

The modern theory of this problem was founded by Federer and Fleming [3]. They defined integral p -currents, which roughly correspond to formal sums of p -dimensional submanifolds, except for sets of p -dimensional Hausdorff measure zero. Federer and Fleming showed that for any M and $[\Sigma]$, there exists an integral p -current Σ_{\min} representing $[\Sigma]$ that has minimal volume. With some laxity of language, we refer to this minimal current as a minimal cycle. Applied to string theory compactified on M , the minimal cycle Σ_{\min} plausibly describes the configuration of a Euclidean brane in the class $[\Sigma]$, up to corrections from worldvolume fields beyond the embedding coordinates.

Geometric measure theory thus provides a sound framework for analyzing volume-minimization. Even so, actually computing the volume $\text{Vol}(\Sigma_{\min})$ of the minimal cycle in a given class remains difficult for general M and $[\Sigma]$. One knows that Σ_{\min} exists [3], and much is known about its degree of singularity [4], but not much more can be said at this level of generality. However, if M is a Kähler manifold, the minimal cycles in effective classes are readily obtained from calibration data, as reviewed above. One might therefore hope to characterize minimal cycles in general classes in terms of the properties of calibrated minimal cycles in effective classes.

²By an anti-effective divisor, we mean a finite formal sum of irreducible antiholomorphic hypersurfaces, with nonnegative integer coefficients.

Suppose, then, that X is a Kähler threefold with $h^{2,0}(X) = 0$, such as a Calabi-Yau threefold or an orientifold thereof.³ Then any $[\Sigma] \in H_4(X, \mathbb{Z})$ can be represented by a union of irreducible holomorphic and antiholomorphic four-cycles. We refer to such a representative Σ_{\cup} as a *piecewise-calibrated* representative, because each irreducible component is either calibrated by J , or would be calibrated by J following a reversal of orientation, cf. (1.2). The volume $\text{Vol}(\Sigma_{\cup})$ is then the sum of the volumes of the constituent holomorphic and antiholomorphic four-cycles, with each of these obtained by applying (1.1) or (1.2), respectively.⁴ Thus, for given $[\Sigma]$, $\text{Vol}(\Sigma_{\cup})$ is readily computed in terms of the Kähler parameters of X .

To connect to the problem of Euclidean D3-branes, we note that in compactifications preserving $\mathcal{N} = 1$ supersymmetry, only Euclidean D3-branes wrapping effective divisors can contribute to the superpotential [1], and so the cycle volumes associated to superpotential terms are given in terms of calibration data. On the other hand, a non-BPS Euclidean D3-brane wrapping the minimal-volume representative Σ_{\min} of a non-effective class $[\Sigma]$ could contribute to the Kähler potential or to a higher F-term [5], depending on the number of fermion zero modes [6–9]. The piecewise-calibrated representative Σ_{\cup} can be understood as an unstable collection of BPS and anti-BPS Euclidean D3-branes that can recombine and fuse, reducing their volume, until they arrive at the non-BPS volume-minimizing configuration Σ_{\min} .

If we define the *recombination fraction*

$$\tau_{\Sigma} := \frac{\text{Vol}(\Sigma_{\cup}) - \text{Vol}(\Sigma_{\min})}{\text{Vol}(\Sigma_{\min})}, \tag{1.3}$$

then $\text{Vol}(\Sigma_{\cup})$ gives a useful approximation to $\text{Vol}(\Sigma_{\min})$ if and only if $\tau_{\Sigma} \ll 1$. When $\tau_{\Sigma} \gtrsim 1$, the minimal cycle and piecewise-calibrated cycle are quite different, and the action of a non-BPS instanton cannot be accurately estimated by adding up the action of its piecewise-calibrated BPS and anti-BPS ‘constituents’. When $\tau_{\Sigma} \gg 1$, we say that *large recombination* has occurred.

For the purpose of controlling the α' expansion in string theory, it is quite important to know whether $\text{Vol}(\Sigma_{\min}) \approx \text{Vol}(\Sigma_{\cup})$ is a fair approximation, both for $[\Sigma] \in H_4(X, \mathbb{Z})$ as we have discussed, and for the analogous situation when $[\Sigma] \in H_2(X, \mathbb{Z})$ is a non-holomorphic curve class. The reason is as follows. For X an orientifold of a Calabi-Yau threefold, let $\{\sigma_i\}$, $i = 1, \dots, h^{1,1}(X)$, be a basis of $H_2(X, \mathbb{Z})$, and define the Kähler parameters $t_i := \int_{\sigma_i} J$. Many authors take $t_i \gg 1$ to be a *sufficient* condition for ensuring that X is large enough so that perturbative and nonperturbative corrections in the α' expansion are well-controlled. There is some motivation for this condition: the actions of worldsheet instantons and Euclidean D3-branes wrapping effective curve and divisor classes, respectively, are indeed determined by the t_i and by the topological data of X . Likewise, for non-effective classes $[\Sigma]$, the volume $\text{Vol}(\Sigma_{\cup})$ of the piecewise-calibrated representative

³In more general Kähler manifolds X with $h^{2,0}(X) \neq 0$, our analysis is only relevant to cycles dual to elements in the Néron-Severi lattice of X .

⁴In general, a class $[\Sigma]$ may have multiple piecewise-calibrated representatives $\{\Sigma_{\cup}^I\}$, with I an index set, as we will explain further below. By Σ_{\cup} we mean the one of these that has the smallest volume.

is again determined by the t_i . However, when $\tau_\Sigma \gg 1$ for some $[\Sigma]$, the corresponding non-BPS instanton is far more important than the computation of $\text{Vol}(\Sigma_\cup)$ would suggest.

To put this in stark terms, one can envision a ‘vulnerable’ Calabi-Yau threefold X for which, even with all $t_i \gg 1$, there are non-BPS instantons with actions of order unity, even though all BPS instantons have large action. The resulting corrections would invalidate the α' expansion.

Whether such vulnerable threefolds actually exist is a purely geometric question, amounting to the existence of a pair $\{X, [\Sigma]\}$ with τ_Σ sufficiently large. In principle this could be answered within geometric measure theory, without any input from physics. However, to the best of our knowledge, this question is not settled for Calabi-Yau n -folds for any n . Perhaps the closest result is this: Micallef and Wolfson have exhibited a pair $\{X, [\Sigma]\}$ with X a K3 surface near an orbifold limit, and $[\Sigma] \in H_2(X, \mathbb{Z})$, for which $\tau_\Sigma \geq \varepsilon$, with ε parameterizing the distance from the orbifold limit [10]. However, this establishes only that $\tau_\Sigma > 0$ can occur in a K3 surface, not that $\tau_\Sigma \gg 1$ can occur there. In other words, Micallef and Wolfson proved that nonzero recombination can occur, not that large recombination can occur.⁵

The goal of this work is to derive results about minimal surfaces that follow from the *Weak Gravity Conjecture* (WGC) [18]. The instanton form of the WGC applied to an axion charge vector \vec{Q} asserts an upper bound on the action $S_{\min}[\vec{Q}]$ of the smallest-action instanton of charge \vec{Q} , in terms of a certain quadratic norm $\|\vec{Q}\|$. Schematically,

$$\text{WGC} \Rightarrow S_{\min}[\vec{Q}] \leq \text{const.} \times \|\vec{Q}\|. \tag{1.4}$$

We present a precise version of this relation, applied to the case of interest, in (2.10).⁶

We will consider compactification of type IIB string theory on an orientifold X of a complex threefold, and study the axions resulting from reduction of the Ramond-Ramond four-form C_4 on four-cycles. For a given class $[\Sigma]$, the axion charge vector is then $[\Sigma]$ itself, understood as a vector in $H_4(X, \mathbb{Z})$. The corresponding instantons are Euclidean D3-branes wrapping the minimal surface Σ_{\min} , with Euclidean action whose real part is proportional to $\text{Vol}(\Sigma_{\min})$. The quadratic norm $\|\Sigma\|$ is determined by the metric on Kähler moduli space, see (2.9).

In this situation, the WGC applied to the class $[\Sigma]$ asserts a lower bound

$$\text{WGC} \Rightarrow \tau_\Sigma \geq \tau_\Sigma^{\min}, \tag{1.5}$$

with τ_Σ^{\min} computable in terms of $\|\Sigma\|$, cf. (2.11). We will give an explicit example of an orientifold X of a Calabi-Yau threefold hypersurface in a toric variety, and a hyperplane $\mathcal{H} \subset H_4(X, \mathbb{Z})$, for which $\tau_\Sigma^{\min} \gg 1$ for all $[\Sigma] \in \mathcal{H}$. Thus, if the WGC applies to *any* $[\Sigma] \in \mathcal{H}$, it implies $\tau_\Sigma \gg 1$. In particular, if the sub-Lattice WGC [20] holds in X , it follows that $\tau_\Sigma \gg 1$, and so certain four-cycles necessarily undergo large recombination.

The evidence that the sub-Lattice WGC is a fact about quantum gravity is not conclusive, and for the purpose of this work we are agnostic about its truth value. Our result

⁵See also, for example, [11–17] for related ideas and further references.

⁶The possibility of using the WGC to bound the volumes of non-holomorphic curves in compactifications on K3 surfaces was noted in [19].

can on the one hand be read as preparing a purely geometric test, the condition $\tau_\Sigma \geq \tau_\Sigma^{\min}$ of (1.5), failure of which in a single example of a string compactification would disprove the sub-Lattice WGC. It would be striking if the tools of geometric measure theory could be used in this way to discern properties of quantum gravity. On the other hand, if the sub-Lattice WGC is taken to be true, either provisionally or based on further evidence about quantum gravity, then our result can be read as an upper bound on the volumes of certain minimal surfaces.

The organization of this paper is as follows. In section 2 we collect the necessary results about the WGC. In section 3 we present an explicit example where $\tau_\Sigma^{\min} \gg 1$. We discuss some implications, and conclude, in section 4. Appendix A gives more details about the geometry of the example. In appendix B, we prove a useful lemma about effective divisors in the example.

2 Axions, orientifolds, and weak gravity

We begin by explaining how the WGC constrains the volumes of non-holomorphic four cycles. Let us first describe our normalization conventions. We define the string length ℓ_s as

$$\ell_s = 2\pi\sqrt{\alpha'}. \tag{2.1}$$

Following the conventions of [21] we take

$$\frac{1}{2\kappa_{10}^2} = \frac{2\pi}{\ell_s^8}, \quad T_p = \frac{2\pi}{g_s \ell_s^{p+1}}, \tag{2.2}$$

where T_p is the tension of a Dp-brane. We measure all cycle volumes using the ten-dimensional Einstein-frame metric, and express these volumes in units of ℓ_s . The four-dimensional Planck mass is given by

$$M_{\text{pl}}^2 = \frac{4\pi\mathcal{V}}{\ell_s^2 g_s^{1/2}}, \tag{2.3}$$

where \mathcal{V} is the compactification volume. Henceforth, we set $\ell_s = 1$, so that the only dimensionful parameter appearing in our equations is M_{pl} .

We consider type IIB string theory compactified on an O3/O7 orientifold X of a Calabi-Yau threefold \tilde{X} .⁷ Given a basis $\{D_i\}$, $i = 1 \dots h^{1,1}$ for $H_4(X, \mathbb{Z})$, the Kähler moduli are written as

$$T^i = \frac{1}{2} \int_{D_i} J \wedge J + i \int_{D_i} C_4 \equiv \tau^i + i\theta^i, \quad i = 1 \dots h^{1,1}, \tag{2.4}$$

where C_4 is the Ramond-Ramond four-form. The effective Lagrangian for the axions θ^i has the form

$$\mathcal{L} = \frac{M_{\text{pl}}^2}{2} \mathcal{R}_4 - \frac{M_{\text{pl}}^2}{2} K_{ij} \partial^\mu \theta^i \partial_\mu \theta^j - V(\theta). \tag{2.5}$$

⁷In this case we have $h^{2,0}(X) = 0$, see e.g. [22], and therefore all elements of $H_4(X, \mathbb{Z})$ admit piecewise-calibrated representatives.

At large volume, the Kähler metric K_{ij} on the complexified Kähler moduli space of X is independent of the θ^i and is obtained from the tree level Kähler potential $\mathcal{K} = -2 \log \mathcal{V}$. In perturbation theory, the axions θ^i enjoy continuous shift symmetries due to the ten-dimensional gauge invariance of C_4 . These are broken to discrete shift symmetries by non-perturbative contributions to the scalar potential. The discrete shift symmetries generate the period lattice, denoted Γ^* , so the general axion potential can be written as

$$V(\theta) = \sum_{[\Sigma] \in \Gamma^*} Z_\Sigma e^{2\pi i [\Sigma] \cdot \vec{\theta}}, \tag{2.6}$$

where $Z_{-\Sigma} = Z_\Sigma^*$ as $V(\theta)$ is real. We will consider the potential $V(\theta)$ generated by Euclidean D3-branes, which wrap minimum-volume representatives Σ_{\min} of homology classes $[\Sigma] \in H_4(X, \mathbb{Z})$, and thus $\Gamma^* = H_4(X, \mathbb{Z})$. In the semiclassical regime, the coefficients Z_Σ are given by

$$Z_\Sigma = A_\Sigma e^{-S_\Sigma}, \tag{2.7}$$

where A_Σ is the one-loop determinant and the semiclassical action S_Σ of the instanton is determined by the cycle volume,

$$\text{Re } S_\Sigma = 2\pi \text{Vol}(\Sigma_{\min}), \tag{2.8}$$

measured with respect to the metric g on X .

The instanton form of the WGC applied to a charge vector $[\Sigma]$ states that there exists an instanton with charge $[\Sigma]$ that satisfies

$$\text{Re } S_\Sigma \leq c \|\Sigma\|, \quad \text{where } \|\Sigma\| := 2\pi \sqrt{\Sigma_i (K^{-1})^{ij} \Sigma_j}, \tag{2.9}$$

for some $c \sim 1$. Various forms of the WGC constrain various subsets of the charge lattice $H_4(X, \mathbb{Z})$. In particular, the sub-Lattice WGC states that there exists a sublattice $\Gamma_{\text{sub}}^* \subseteq \Gamma^*$ of finite index⁸ such that at every site in Γ_{sub}^* there is an instanton that satisfies (2.9).⁹

Using (2.8), the WGC constraint (2.9) becomes

$$\text{Vol}(\Sigma_{\min}) \leq c \sqrt{\Sigma_i (K^{-1})^{ij} \Sigma_j}. \tag{2.10}$$

Equivalently, using (1.3), the WGC implies a lower bound on the recombination fraction

$$\tau_\Sigma \geq \tau_\Sigma^{\min} := \frac{2\pi \text{Vol}(\Sigma_{\min})}{c \|\Sigma\|} - 1. \tag{2.11}$$

In the next section, we will present an example in which — as a purely geometric statement, making no assumption concerning the WGC — for any charge vector $[\Sigma]$ lying in a specific hyperplane $\mathcal{H} \subset H_4(X, \mathbb{Z})$, the Kähler form J can be chosen so as to make the lower bound τ_Σ^{\min} arbitrarily large. If the WGC applies to any such $[\Sigma]$, it implies $\tau_\Sigma \gg 1$.

⁸The index of a sublattice Γ_{sub}^* is the smallest integer n such that $n[\Sigma] \in \Gamma_{\text{sub}}^*$ for all $[\Sigma] \in \Gamma^*$.

⁹The foundational paper on the Weak Gravity Conjecture is [18]. The sub-Lattice WGC was formulated by Heidenreich, Reece, and Rudelius in [20], building on [23], and closely related work by Montero, Shiu, and Soler appears in [24]. Other related recent developments include [20, 25–42].

We remark that given an example of large recombination — i.e., a pair $\{[\Sigma], J\}$ for which $\mathfrak{r}_\Sigma^{\min} \gg 1$ — one can always rescale $J \rightarrow \lambda J$, for $\lambda \in \mathbb{R}^+$, without affecting the relation (2.11). Thus, by an appropriate rigid dilation of X , we can ensure that J is as far as we like from the walls of the Kähler cone, so that α' corrections do not impact our analysis.

Before proceeding to the example, let us first establish that for Euclidean D3-branes wrapping effective or anti-effective cycles, (2.9) is always satisfied with $c = \sqrt{3}/2$. The proof is as follows. We expand the Kähler form J in terms of the Poincaré duals ω_i of a set of basis divisors D_i ,

$$J = t_i \omega_i. \tag{2.12}$$

In terms of the Kähler parameters t_i , the divisor volumes and the volume of X are

$$\begin{aligned} \tau_i &= \frac{1}{2} \kappa_{ijk} t_j t_k, \\ \mathcal{V} &= \frac{1}{6} \kappa_{ijk} t_i t_j t_k, \end{aligned} \tag{2.13}$$

where $\kappa_{ijk} = \#D_i \cap D_j \cap D_k$ are the triple intersection numbers. The inverse Kähler metric $(K^{-1})^{ij}$ has the form

$$(K^{-1})^{ij} = 4\tau_i \tau_j - 4\mathcal{V} \kappa_{ijk} t_k. \tag{2.14}$$

Using (2.13) and (2.14) we can write

$$\frac{\text{Re } S_\Sigma}{\|\Sigma\|} = \frac{2\pi \text{Vol}(\Sigma)}{\|\Sigma\|} = \frac{2\pi t \cdot K^{-1} \cdot \Sigma}{8\pi \mathcal{V} \sqrt{\Sigma \cdot K^{-1} \cdot \Sigma}} = \frac{t \cdot K^{-1} \cdot \hat{\Sigma}}{4\mathcal{V}}, \tag{2.15}$$

where we defined $\hat{\Sigma} = \Sigma / \|\Sigma\|$. The ratio in (2.15) is maximized when $\Sigma \propto t$, and since $t \cdot K^{-1} \cdot t = 12\mathcal{V}^2$ we have

$$\frac{\text{Re } S_\Sigma}{\|\Sigma\|} \leq \frac{12\mathcal{V}^2}{4\mathcal{V} \sqrt{12\mathcal{V}^2}} = \frac{\sqrt{3}}{2}. \tag{2.16}$$

3 An orientifold where weak gravity implies large recombination

We now give an explicit example of an orientifold X of a Calabi-Yau threefold hypersurface, and a hyperplane $\mathcal{H} \subset H_4(X, \mathbb{Z})$, such that for any $[\Sigma] \in H$, the ratio $\mathfrak{r}_\Sigma^{\min}$ defined in (2.11) can be made arbitrarily large by a choice of the Kähler form J .¹⁰

We consider the product $V = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{F}_4$, where \mathbb{F}_4 is the fourth Hirzebruch surface. Realized as a toric fourfold, V has degrees in the Cox ring

x_0	x_1	x_2	x_3	x_4	h	x_6	η
1	1	0	0	0	0	0	0
0	0	1	1	0	0	0	0
0	0	0	0	0	1	0	1
0	0	0	0	1	-4	1	0

(3.1)

¹⁰Further details are presented in appendix A.

and Stanley-Reisner ideal

$$\langle x_0x_1, x_4x_6, x_2x_3, h\eta \rangle . \tag{3.2}$$

The orientifold $X \subset V$ is defined by the vanishing of a polynomial F of the form

$$F = hP_{(2,2,0,4)} - \eta P_{(2,2,0,0)} , \tag{3.3}$$

where $P_{(2,2,0,4)}$ and $P_{(2,2,0,0)}$ are polynomials independent of h and η . We define a basis $\{D_a, D_b, D_c, D_d\}$ for $H_4(X, \mathbb{Z})$ with degrees

$$\begin{aligned} [D_a] &= (1, 0, 0, 0) , \\ [D_b] &= (0, 1, 0, 0) , \\ [D_c] &= (0, 0, 1, -4) , \\ [D_d] &= (0, 0, 0, 1) , \end{aligned} \tag{3.4}$$

and expand the Kähler form J as

$$J = t_a\omega_a + t_b\omega_b + t_c\omega_c + t_d\omega_d , \tag{3.5}$$

where ω_A , $A \in \{a, b, c, d\}$, are Poincaré dual to D_A . The Mori cone of V is simplicial, and the volumes of its generators are

$$\{t_a, t_b, t_c, t_d - 4t_c\} . \tag{3.6}$$

We now make the notational replacement

$$t_d \rightarrow t_e + 4t_c , \tag{3.7}$$

so that the Kähler cone conditions read $t_a > 0, t_b > 0, t_c > 0$, and $t_e > 0$. With this parameterization the volume \mathcal{V} of X takes the form

$$\mathcal{V} = 4t_b t_c (t_a + t_c) + t_b (t_a + 2t_c) t_e + 2t_a t_c (2t_c + t_e) . \tag{3.8}$$

To show that the WGC implies large recombination, we need to find a non-effective divisor class $[\Sigma] \in H_4(X, \mathbb{Z})$, and a point in the Kähler cone of X for which τ_Σ^{\min} is large. To find such a non-effective divisor class and characterize its piecewise calibrated representatives, we must first identify the effective and anti-effective divisor classes on X .

Recall that for Y an algebraic n -fold, a homology class $[\Sigma] \in H_{2m}(Y, \mathbb{Z})$, for $1 \leq m \leq n$, is called (anti-)effective if it can be represented by a (anti-)holomorphic $2m$ -cycle. The cone of effective $2m$ -cycles, which we denote by $\mathcal{E}_m(Y)$, is by definition the cone in $H_{2m}(Y, \mathbb{R})$ generated by the effective classes in $H_{2m}(Y, \mathbb{Z})$. Likewise, the cone $\bar{\mathcal{E}}_m(Y)$ of anti-effective $2m$ -cycles is generated by the anti-effective classes in $H_{2m}(Y, \mathbb{Z})$.

When Y is a hypersurface of dimension n in a toric variety V of dimension $n+1$, some effective divisors D on Y are inherited from effective divisors \hat{D} on V , with $D = Y \cap \hat{D}$. Enumerating these inherited divisors is straightforward, because $\mathcal{E}_n(V)$ is readily computed from toric data. However, a general such Y may also contain *autochthonous* effective divisors, i.e. effective divisors D that are not of the form $Y \cap \hat{D}$ for any effective divisor

\widehat{D} on V , and so are not simply inherited from the divisors of V . Certain autochthonous divisors can be identified by attempting to factor the defining equation of Y [43].

Finding all autochthonous divisors on X and computing the generators of $\mathcal{E}_2(X)$ exactly is difficult and beyond the scope of this work. Instead, we will construct a cone $\mathcal{E}_2^{\text{out}}(X)$ that contains $\mathcal{E}_2(X)$. We first note that any effective divisor in X is an effective surface in V , i.e. $\mathcal{E}_2(X) \subseteq \mathcal{E}_2(V)$.¹¹ Moreover, by a theorem of Fulton, MacPherson, Sottile, and Sturmfels [44], the cone of effective n -cycles $\mathcal{E}_n(V)$ in V is generated by the toric n -cycles of V . We consider a general line bundle \mathcal{L} on V , and demand $\mathcal{L}|_X \in \mathcal{E}_2(V)$. We find¹² that any such line bundle is generated over $\mathbb{Z}_{\geq 0}$ by:

$$\begin{aligned}
 [D_a] &= (1, 0, 0, 0), \\
 [D_b] &= (0, 1, 0, 0), \\
 [D_c] &= (0, 0, 1, -4), \\
 [D_d] &= (0, 0, 0, 1), \\
 [D_e] &= (-2, 2, 1, 0), \\
 [D_f] &= (2, 2, -1, 4), \\
 [D_g] &= (2, -2, 1, 0).
 \end{aligned} \tag{3.9}$$

The divisors $[D_a], \dots, [D_g]$ are the extremal generators of $\mathcal{E}_2^{\text{out}}(X)$. Their volumes¹³ $\tau_a := \int_{D_a} J \wedge J, \dots, \tau_g := \int_{D_g} J \wedge J$ are

$$\begin{aligned}
 \tau_a &= 4t_b t_c + 4t_c^2 + t_b t_e + 2t_c t_e, \\
 \tau_b &= 4t_a t_c + 4t_c^2 + t_a t_e + 2t_c t_e, \\
 \tau_c &= 2t_a t_e + 2t_b t_e, \\
 \tau_d &= t_a t_b + 2t_b t_c + 2t_a t_c, \\
 \tau_e &= 4t_a t_b + 16t_a t_c + 4t_a t_e, \\
 \tau_f &= 8t_a t_c + 8t_b t_c + 16t_c^2 + 8t_c t_e, \\
 \tau_g &= 4t_a t_b + 16t_b t_c + 4t_b t_e.
 \end{aligned} \tag{3.10}$$

Next, we consider the hyperplane $\mathcal{H} \subset H_4(X, \mathbb{Z})$ spanned by charge vectors of the form

$$[\Sigma^{(k)}] := (k, -k, c, d), \tag{3.11}$$

where $k \in \mathbb{Z}_+$ and $c, d \in \mathbb{Z}$. In appendix B, we prove that every such $[\Sigma^{(k)}]$ is neither effective nor anti-effective. We will now demonstrate that any WGC that constrains a class $[\Sigma^{(k)}]$ implies large recombination.

¹¹The converse is not true in general. We will show below that $\mathcal{E}_2(X) \neq \mathcal{E}_2(V)$ in our example.

¹²Further details will appear in [43].

¹³While the extremal generators of $\mathcal{E}_2^{\text{out}}(X)$ are not necessarily effective divisors on X , they are effective divisors on V , and it is in the latter sense that τ_a, \dots, τ_g are volumes.

As $[\Sigma^{(k)}]$ is non-effective, we express it as a sum of effective and anti-effective divisor classes, and construct a piecewise-calibrated representative $\Sigma_{\cup}^{(k)}$:

$$[\Sigma^{(k)}] = \sum_{[\Sigma_i] \in \mathcal{E}_2(X)} \alpha_i [\Sigma_i] + \sum_{[\bar{\Sigma}_i] \in \bar{\mathcal{E}}_2(X)} \beta_i [\bar{\Sigma}_i], \quad (3.12)$$

$$\Sigma_{\cup}^{(k)} = \bigcup_{[\Sigma_i] \in \mathcal{E}_2(X)} \alpha_i \Sigma_i \cup \bigcup_{[\bar{\Sigma}_i] \in \bar{\mathcal{E}}_2(X)} \beta_i \bar{\Sigma}_i, \quad (3.13)$$

where $\alpha_i, \beta_i \in \mathbb{Z}_{\geq 0}$. The volume of $\Sigma_{\cup}^{(k)}$ is the sum of the volumes of its holomorphic and anti-holomorphic constituents,

$$\text{Vol}(\Sigma_{\cup}^{(k)}) = \sum_{[\Sigma_i] \in \mathcal{E}_2(X)} \alpha_i \text{Vol}(\Sigma_i) + \sum_{[\bar{\Sigma}_i] \in \bar{\mathcal{E}}_2(X)} \beta_i \text{Vol}(\bar{\Sigma}_i). \quad (3.14)$$

Note that the expansion (3.12) is not unique: any given $[\Sigma^{(k)}]$ can be expanded in infinitely many inequivalent ways. We define $\Sigma_{\cup}^{(k)}$ to be the piecewise-calibrated representative with the smallest volume. In general, finding the smallest-volume piecewise-calibrated representative is difficult. However, we will work in a region of the Kähler cone where the problem is simplified considerably. Consider the locus where $t_a = t_b = \sigma$, $t_c = t_e = \delta$ for some $\delta, \sigma > 0$. The volumes of the generators of $\mathcal{E}_2^{\text{out}}(X)$, cf. (3.10), become

$$\begin{aligned} \tau_a &= 6\delta^2 + 5\sigma\delta, \\ \tau_b &= 6\delta^2 + 5\sigma\delta, \\ \tau_c &= 4\sigma\delta, \\ \tau_d &= 4\sigma\delta + \sigma^2, \\ \tau_e &= 20\sigma\delta + 4\sigma^2, \\ \tau_f &= 24\delta^2 + 16\sigma\delta, \\ \tau_g &= 20\sigma\delta + 4\sigma^2. \end{aligned} \quad (3.15)$$

The quadratic norm $\|\Sigma^{(k)}\|$, cf. (2.9), is computed using (2.13), (2.14) and (3.8):¹⁴

$$\begin{aligned} \|\Sigma^{(k)}\|^2 &= 8\pi^2 \left(60k^2\delta^3\sigma + (128c^2 + 16cd + 32d^2 + 25k^2)\delta^2\sigma^2 \right. \\ &\quad \left. + (40c^2 - 4cd + 16d^2)\delta\sigma^3 + 4d^2\sigma^4 \right). \end{aligned} \quad (3.16)$$

We will now show that the volume of any piecewise-calibrated representative of $[\Sigma^{(k)}]$ obeys $\text{Vol}(\Sigma_{\cup}^{(k)}) > 12\delta^2$. As $\mathcal{E}_2(X) \subset \mathcal{E}_2^{\text{out}}(X)$, the volume of any effective or anti-effective divisor can be written as a nonnegative integer sum of the volumes given in (3.10). We can therefore write

$$\text{Vol}(\Sigma_{\cup}^{(k)}) = \sum_A \gamma^A \tau_A, \quad (3.17)$$

where $A \in \{a, b, c, d, e, f, g\}$ and $\gamma^A \in \mathbb{Z}_{\geq 0}$. Comparing to (3.10), we can write

$$\text{Vol}(\Sigma_{\cup}^{(k)}) = m_1\delta^2 + m_2\delta\sigma + m_3\sigma^2, \quad (3.18)$$

¹⁴Intersection numbers κ_{ijk} can be calculated by differentiating (3.8).

with $m_1, m_2, m_3 \in \mathbb{Z}_{\geq 0}$. Let us now show that $m_1 \geq 12$. We can represent each homology class by the corresponding degrees, as in (3.4), and rewrite the sum (3.12) as

$$[\Sigma^{(k)}] = \sum_i C_i^{(k)}(a_i, b_i, c_i, d_i) \tag{3.19}$$

where i runs over effective and anti-effective divisors, and $C_i^{(k)} \in \mathbb{Z}_{\geq 0}$. We then break up the general sum (3.19) into three subclasses:

$$\begin{aligned} [\Sigma^{(k)}] &= \sum_{a_i=b_i=0} A_i^{(k)}(a_i, b_i, c_i, d_i) \\ &+ \sum_{a_i=-b_i \neq 0} B_i^{(k)}(a_i, b_i, c_i, d_i) \\ &+ \sum_{a_i \neq -b_i} C_i^{(k)}(a_i, b_i, c_i, d_i), \end{aligned} \tag{3.20}$$

where $A_i^{(k)}, B_i^{(k)}, C_i^{(k)} \in \mathbb{Z}_{\geq 0}$. By Theorem 1 of appendix B, any divisor class with $a_i = -b_i \neq 0$ is non-effective and cannot contribute to the sum, so we have $B_i^{(k)} = 0$. By assumption $k \neq 0$, so there must be at least one term where $a_i \neq 0$, and at least one other term where $b_i \neq 0$: that is, $C_i^{(k)} \neq 0$ for at least two values of i . From (3.4) and (3.15), we see that the volume of any divisor with $a_i \neq -b_i$ is greater than $6\delta^2$. It follows that $\text{Vol}(\Sigma_{\cup}^{(k)}) > 12\delta^2$.

The quadratic norm $\|\Sigma^{(k)}\|$ grows only as $\mathcal{O}(\delta^{3/2})$ in the limit $\delta \gg \sigma$. Thus, for $\delta \gg \sigma$, there is a lower bound on the recombination fraction:

$$\mathfrak{r}_{\Sigma^{(k)}}^{\min} = \frac{2\pi \text{Vol}(\Sigma_{\cup}^{(k)})}{c \|\Sigma^{(k)}\|} - 1 \sim \mathcal{O}(\delta^{1/2}). \tag{3.21}$$

By taking δ large while holding σ fixed, we can make $\mathfrak{r}_{\Sigma^{(k)}}^{\min}$ as large as desired.

As a concrete example, take $\delta = 100$, $\sigma = 1$ and $\Sigma = (-k, k, 0, 0)$, for some $k \in \mathbb{Z}_{\geq 0}$. We find $\|\Sigma\| = 1000\pi k\sqrt{482}$ and $\text{Vol}(\Sigma_{\cup}^{(k)}) > 120000k$. Setting $c = 1$ in (2.9), we get

$$\mathfrak{r}_{\Sigma^{(k)}}^{\min} > 9.9. \tag{3.22}$$

The argument presented above applies for any $k \in \mathbb{Z}_+$ and any $c, d \in \mathbb{Z}$. Thus, any version of the WGC that constrains at least one charge vector lying in the hyperplane \mathcal{H} spanned by the $[\Sigma^{(k)}]$ necessarily implies that large recombination occurs. In particular, because the vector $(1, -1, c, d)$ lies in Γ^* , for any sublattice $\Gamma_{\text{sub}}^* \subset \Gamma^*$ there exists a finite $n \in \mathbb{Z}_+$ such that $(n, -n, nc, nd) \in \Gamma_{\text{sub}}^*$. Thus, the sub-Lattice WGC for any sublattice Γ_{sub}^* , no matter how sparse, implies that some of the charges in the hyperplane spanned by $[\Sigma^{(k)}]$ must obey (2.9). It follows that the sub-Lattice WGC requires large recombination in our example.

4 Discussion

We have argued that if the Weak Gravity Conjecture is true, the volumes of certain minimal surfaces must obey a nontrivial upper bound. The bound requires that each minimal surface

is sufficiently small compared to any union of holomorphic and antiholomorphic surfaces representing the same homology class.

Specifically, in a compactification of type IIB string theory on an O3/O7 orientifold of any Calabi-Yau threefold X , and for any $[\Sigma] \in H_4(X, \mathbb{Z})$ that is neither effective nor anti-effective on X , but is constrained by the WGC, the volume $\text{Vol}(\Sigma_{\min})$ of the minimal-volume representative of $[\Sigma]$ must obey (2.10). This amounts to the statement that the minimal cycle representing $[\Sigma]$ must be sufficiently small in comparison to any union of holomorphic and antiholomorphic cycles representing $[\Sigma]$.

The outline of our argument was as follows. The real part of the action of a Euclidean D3-brane in a class $[\Sigma]$, ignoring the effects of magnetization and of the fluctuation determinant, is $2\pi\text{Vol}(\Sigma_{\min})$. The instanton form of the WGC then gives an upper bound on $\text{Vol}(\Sigma_{\min})$ in terms of the quadratic norm $\|\Sigma\|$, cf. (2.9), of the axion charge $[\Sigma]$. Although $\text{Vol}(\Sigma_{\min})$ is difficult to compute, one can instead compare $\|\Sigma\|$ to the volume $\text{Vol}_{\cup}(\Sigma)$ of a piecewise-calibrated representative of $[\Sigma]$, which corresponds to the sum of the actions of a collection of BPS and anti-BPS Euclidean D3-branes whose total charge is $[\Sigma]$. Given any such piecewise-calibrated representative, one can think of fusing and recombining the holomorphic and antiholomorphic constituents to form a non-holomorphic cycle with smaller total volume. Any version of the WGC that nontrivially constrains $[\Sigma]$ asserts that $\text{Vol}(\Sigma_{\min}) < \text{Vol}_{\cup}(\Sigma)$, with (2.11) giving the precise relation.

The physical origin of our statement is simply that the WGC prescribes a minimum amount of binding energy that must be released when certain collections of BPS and anti-BPS states bind to form a non-BPS state. We have mapped this requirement into a statement about minimal surfaces in orientifolds of Calabi-Yau threefolds, i.e. about an infinite class of instances of the Plateau problem.

We should comment that if one is given the topological data of an orientifold X of a Calabi-Yau threefold, together with the Kähler cone $\mathcal{K}(X)$ of X and the cone $\mathcal{E}_2(X)$ of effective divisors on X , it is straightforward to evaluate $\text{Vol}_{\cup}(\Sigma)$ and $\|\Sigma\|$ at any point in $\mathcal{K}(X)$. Even so, it turned out to be nontrivial to exhibit such an X , as an orientifold of a hypersurface in a toric variety V , for which it was possible to show $\text{Vol}_{\cup}(\Sigma) \gg \|\Sigma\|$. The catch is that $\mathcal{E}_2(X)$ is *not* simply inherited from V by intersecting X with divisors of V — see [45] for a related discussion — and with incomplete knowledge of $\mathcal{E}_2(X)$ it is difficult to compute $\text{Vol}_{\cup}(\Sigma)$. We overcame this limitation by showing, in appendix B, that a particular family of divisor classes are neither effective nor anti-effective, and so arrived at the example of section 3.

Some cautionary remarks are necessary. We have ignored degrees of freedom on the Euclidean D3-brane worldvolume other than the embedding coordinates, we have omitted the Pfaffian prefactor, and we have ignored the effects of magnetization. These simplifying assumptions led to (2.8), which we used to relate the defining statement (2.9) of the instanton WGC to the version (2.10) used in our analysis. It is therefore logically possible that there exist examples of pairs $\{X, [\Sigma]\}$ in which $\text{Vol}(\Sigma_{\min})$ violates (2.10), and yet the WGC (2.9) holds nonetheless, because (2.8) is strongly violated. We find such large violations of (2.8) to be implausible in the regime where all Kähler parameters are large, but this deserves more detailed study.

We have also neglected the possibility of binding energy of non-recombined instantons. It is conceivable that the binding energy between a BPS instanton and an anti-BPS instanton could reduce their joint action so much that the WGC is satisfied without the need for recombination. Determining whether this occurs in our example, and more generally understanding the role of bound states in the WGC, are interesting problems that lie beyond the scope of this work.¹⁵

As in most of the literature, we have assumed that the WGC applies at generic points in the moduli space of X . It might be that the WGC only applies on-shell, i.e., at the points in the moduli space where the scalar potential is minimized. One would then need to stabilize the moduli, and check the relation (2.11) at the vacua of the resulting effective field theory. Furthermore, in theories with light scalar fields the WGC can imply constraints that differ from (2.9) [46].

An important application concerns the validity of the α' expansion. As we explained in the introduction, a common assumption is that α' corrections to the effective action are small when the Kähler form J is ‘big’ enough so that every holomorphic curve and holomorphic divisor is large in string units. However, if τ_Σ is sufficiently large for some $[\Sigma]$, then a non-BPS instanton with charge $[\Sigma]$ can give corrections to the effective action that are large enough to invalidate the α' expansion. One should therefore ask whether there exist pairs $\{X, [\Sigma]\}$ with dangerously large τ_Σ .

It would also be interesting to understand the degree to which the WGC implies recombination in other types of four-dimensional $\mathcal{N} = 1$ compactifications, such as type IIA string theory on a Calabi-Yau orientifold, or M-theory on a G_2 manifold.

This work opens the possibility of proving or disproving various versions of the WGC using the methods of geometric measure theory. More broadly, we anticipate that geometric measure theory can be used to illuminate the study of non-BPS instantons.

Acknowledgments

We thank Chris Beasley, Naomi Gendler, Jim Halverson, Ben Heidenreich, Manki Kim, Tom Rudelius, John Stout, Ben Sung, Irene Valenzuela, and Max Zimet for helpful conversations. We are grateful to Camillo De Lellis for discussions about geometric measure theory, and we thank the anonymous referee for comments that led to clarification of our arguments. The work of M.D. and L.M. was supported in part by NSF grant PHY-1719877. The work of C.L. was supported in part by NSF grant PHY-1620526. The work of M.S. was supported in part by NSF grant DMS-1502294. Portions of this work were completed at the Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1607611.

¹⁵We thank the referee for helpful comments on this possibility.

A Geometry of the example

We consider the product $\tilde{V} = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{F}_2$. It can be realized as a toric variety obtained from the 4d reflexive polytope Δ° with boundary points

$$\begin{pmatrix} 1 & -1 & 1 & -1 & -1 & -1 & -1 & 1 \\ 1 & -1 & -1 & 1 & -1 & -1 & -1 & 1 \\ -1 & 1 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 2 & 1 & 0 & -1 \end{pmatrix} \tag{A.1}$$

There is a unique triangulation \mathcal{T} of the boundary of Δ° , which corresponds to a smooth toric fourfold \tilde{V} . The maximal simplices of \mathcal{T} are

$$\begin{array}{cccc} [[0, 2, 4, 5], & [0, 2, 4, 7], & [0, 2, 5, 6], & [0, 2, 6, 7], \\ [0, 3, 4, 5], & [0, 3, 4, 7], & [0, 3, 5, 6], & [0, 3, 6, 7], \\ [1, 2, 4, 5], & [1, 2, 4, 7], & [1, 2, 5, 6], & [1, 2, 6, 7], \\ [1, 3, 4, 5], & [1, 3, 4, 7], & [1, 3, 5, 6], & [1, 3, 6, 7]. \end{array} \tag{A.2}$$

The degrees in the Cox ring of \tilde{V} take the form

x_0	x_1	x_2	x_3	x_4	x_5	x_6	x_7	$-K$
1	1	0	0	0	0	0	0	2
0	0	1	1	0	0	0	0	2
0	0	0	0	0	1	0	1	2
0	0	0	0	1	-2	1	0	0

(A.3)

Here x_i are the toric coordinates of \tilde{V} , and $-K$ is the anticanonical class. The Stanley-Reisner ideal takes the form

$$\langle x_0x_1, x_2x_3, x_4x_6, x_5x_7 \rangle. \tag{A.4}$$

We consider an anticanonical hypersurface \tilde{X} in \tilde{V} , defined by the vanishing of a polynomial F . A generic F takes the form

$$F = x_5^2 P_{(2,2,0,4)} + x_5 x_7 P_{(2,2,0,2)} - x_7^2 P_{(2,2,0,0)}, \tag{A.5}$$

where the P 's are generic polynomials in the variables $(x_0, x_1, x_2, x_3, x_4, x_6)$. We will now construct an orientifold X of this Calabi-Yau manifold by choosing a \mathbb{Z}_2 involution σ , such that $X \simeq \tilde{X}/\sigma$. The general procedure is outlined in [47, 48]. We choose an orientifold involution of the form $\sigma : x_5 \rightarrow -x_5$. To ensure the Calabi-Yau is invariant under this involution we require only even powers of x_5 to appear in F , so we take

$$F = x_5^2 P_{(2,2,0,4)} - x_7^2 P_{(2,2,0,0)}. \tag{A.6}$$

In order to identify the fixed loci, we need to take into account the projective scalings of the coordinates in (A.3). Under the action $x_5 \rightarrow -x_5$, from (A.3) the fixed point loci are solutions of

$$\begin{aligned}
 x_0 &= \lambda_1 x_0 \\
 x_1 &= \lambda_1 x_1 \\
 x_2 &= \lambda_3 x_2 \\
 x_3 &= \lambda_3 x_3 \\
 x_4 &= \lambda_4 x_4 \\
 x_5 &= -\lambda_2 \lambda_4^{-2} x_5 \\
 x_6 &= \lambda_4 x_6 \\
 x_7 &= \lambda_2 x_7,
 \end{aligned}
 \tag{A.7}$$

where $\lambda_i \in \mathbb{C}^*$. Via (A.4) we see that x_0 and x_1 cannot vanish simultaneously, and so we can set $\lambda_1 = 1$. In addition, x_2 and x_3 cannot vanish simultaneously, and so $\lambda_3 = 1$. Finally, x_4 and x_6 cannot vanish simultaneously, and so $\lambda_4 = 1$. We are then left with

$$\begin{aligned}
 x_5 &= -\lambda_2 x_5 \\
 x_7 &= \lambda_2 x_7.
 \end{aligned}
 \tag{A.8}$$

Since x_5 and x_7 cannot vanish simultaneously the solutions to this are $\lambda_2 = 1, x_5 = 0$ and $\lambda_2 = -1, x_7 = 0$. We will therefore have O7-planes on the subloci $x_5 = 0$ and $x_7 = 0$, but these O7-planes will not intersect. Following [48], we construct a new toric variety $V \simeq \tilde{V}/\mathbb{Z}_2$ by writing down a basis of sections of the line bundles in (A.3) that are invariant under σ , which will be the toric coordinates of V . This is achieved by taking

$$(x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_7) \rightarrow (x_0, x_1, x_2, x_3, x_4, x_5^2, x_6, x_7^2).
 \tag{A.9}$$

We then define $h \equiv x_5^2$ and $\eta \equiv x_7^2$. Such a map is $2 \rightarrow 1$ everywhere away from the fixed loci, and $1 \rightarrow 1$ along the fixed loci. V then has degrees in the Cox ring of the form

x_0	x_1	x_2	x_3	x_4	h	x_6	η	F
1	1	0	0	0	0	0	0	2
0	0	1	1	0	0	0	0	2
0	0	0	0	0	1	0	1	1
0	0	0	0	1	-4	1	0	0

(A.10)

The Stanley-Reisner ideal takes the same form as before, with $x_2 \rightarrow h$, and $x_7 \rightarrow \eta$:

$$\langle x_0 x_1, x_4 x_6, x_2 x_3, h \eta \rangle.
 \tag{A.11}$$

The new toric variety V is also smooth, and has the same cone structure as given in (A.2). From the degrees one can construct the rays of the corresponding fan,

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 4 & -1 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \tag{A.12}$$

The orientifold $X \subset V$ is then defined by the vanishing of the polynomial F , with x_5^2 replaced by h and x_7^2 replaced by η :

$$F = hP_{(2,2,0,4)} - \eta P_{(2,2,0,0)} = 0. \tag{A.13}$$

Note that $\{F = 0\}$ is not a Calabi-Yau hypersurface in V .

Let us briefly discuss the F-theory lift of this example. The canonical bundle on X can be computed from the adjunction formula:

$$K_X = (K_V + X)|_X. \tag{A.14}$$

We have $\mathcal{O}_{\tilde{V}}(K_V) = \mathcal{O}_V(-2, -2, -2, 2)$, and $\mathcal{O}_V(X) = \mathcal{O}_V(2, 2, 1, 0)$, and therefore $K_X \simeq \mathcal{O}_V(0, 0, -1, 2)|_X$. The Weierstrass model is written as $y^2 = x^3 + fx + g$, with $f \in \Gamma(\mathcal{O}(-4K_X))$ and $g \in \Gamma(\mathcal{O}(-6K_X))$. In our example f and g both take particularly simple forms

$$\begin{aligned} f &= h^2 (a_1 h^2 P_8(x_4, x_6) + a_2 h \eta P_4(x_4, x_6) + a_3 \eta^2), \\ g &= h^3 (b_1 h^3 P_{12}(x_4, x_6) + b_2 h^2 \eta P'_8(x_4, x_6) + b_3 h \eta^2 P'_4(x_4, x_6) + b_4 \eta^3), \end{aligned} \tag{A.15}$$

where the $a_i, b_i \in \mathbb{C}$. There is a non-Higgsable I_0^* fiber on $h = 0$, consistent with the fact that D_h is rigid. In the particular case

$$\begin{aligned} f &= h^2 \eta^2 \\ g &= h^3 \eta^3, \end{aligned} \tag{A.16}$$

we have an $SO(8)$ gauge group on both $h = 0$ and $\eta = 0$, and no additional 7-branes in X . Such a case allows us to tune to arbitrarily weak coupling globally on X , and so we expect the effective theory to be well-controlled. The Euler characteristic can be computed by the method of [49], and we find $\chi(Y) = 576$. The D3-brane tadpole of this Calabi-Yau fourfold with gauge group $SO(8)^2$ is therefore $\chi(Y)/24 = 24$.

B Effective divisors in the example

In this appendix, we show that on $X = \{F_{2,2,1,0} = 0\} \subset V = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{F}_4$, if $D = (-a, a, c, d)$ is an induced divisor class on X , for $a \in \mathbb{Z}_+, c, d \in \mathbb{Z}$, then D is not effective on X . Recall that a divisor class D on X is called *effective* if $H^0(X, \mathcal{O}_X(D)) \neq 0$.

It is more convenient, and hopefully makes the computations more clear, to generalize this to a more natural setting. In fact, we show the following: suppose that Y is any smooth toric variety, that $\alpha \in \text{Pic}(Y)$ is any effective divisor class, and that $F = F_{2,2,\alpha}$ is a generic polynomial in the Cox ring S of $V = \mathbb{P}^1 \times \mathbb{P}^1 \times Y$ of the given triple degree. Let $X \subset V$ be the zero locus of F . We assume that X is smooth. Given (a, b, γ) , where $a, b \in \mathbb{Z}$, and $\gamma \in \text{Pic}(Y)$, we obtain the pullback $\mathcal{O}_X(a, b, \gamma)$ on X .

In this appendix, we show

Theorem 1 *Fix an integer $a > 0$. For all F in a dense Zariski-open subset of $S_{2,2,\alpha}$, and for all $\gamma \in \text{Pic}(Y)$, the divisor class $(-a, a, \gamma)$ is not effective on X .*

We will first recall some basic facts about the cohomology of line bundles on toric varieties, then we analyze the relevant examples.

B.1 Cohomology of line bundles on toric varieties

We recall the description of the cohomology of line bundles on a toric variety. For simplicity, we now restrict to the case of simplicial toric varieties, but there is no problem generalizing to arbitrary complete normal toric varieties.

Let $V \subset \mathbb{P}_\Sigma$ be the simplicial toric variety corresponding to a complete simplicial fan Σ , with r rays. Let D_1, \dots, D_r denote the corresponding divisors on V . Suppose that $S = \mathbb{C}[x_1, \dots, x_r]$ is the homogeneous coordinate ring of V (Cox ring), and that $I \subset S$ is the Stanley-Reisner ideal of S corresponding to the fan Σ . The polynomial ring S is graded by $\text{Cl}(V)$, with $\deg x_i = [D_i]$.

We often identify Σ with a set of subsets of $[\mathbf{r}] := \{1, \dots, r\}$. Given a subset $\lambda \subset [\mathbf{r}]$, define the simplicial complex Σ_λ to be

$$\Sigma_\lambda := \{\tau \subset \Sigma \mid \tau \subset \lambda\}. \tag{B.1}$$

Consider the Laurent ring $S_{\mathbf{x}} := \mathbb{C}[\mathbb{Z}^r] = \mathbb{C}[x_1^{\pm 1}, \dots, x_r^{\pm 1}]$. It turns out that the cohomology groups of line bundles can often be identified with the \mathbb{C} -span of a finite number of monomials in $S_{\mathbf{x}}$, or at least are closely related to such sets.

For $u \in \mathbb{Z}^r$, let $\text{neg}(u) := \{i \in [\mathbf{r}] \mid u_i < 0\}$. For $\lambda \subset [\mathbf{r}]$ and $\alpha \in \text{Cl}(V)$, define

$$\mathcal{B}(\lambda, \alpha) := \{x^u \mid u \in \mathbb{Z}^r, \text{neg}(u) = \lambda, \deg(u) = \alpha\}, \tag{B.2}$$

and denote the \mathbb{C} -vector space they generate by

$$\text{span}(\lambda, \alpha) := \text{span}_{\mathbb{C}} \mathcal{B}(\lambda, \alpha). \tag{B.3}$$

The result we use is the following (see [50–52]):

$$H^i(\mathcal{O}_V(\alpha)) = \bigoplus_{\lambda \subset [\mathbf{r}]} \text{span}(\lambda, \alpha) \otimes \tilde{H}^{i-1}(\Sigma_\lambda, \mathbb{C}). \tag{B.4}$$

Let

$$\Sigma^i = \Sigma_V^i := \{\lambda \subset [\mathbf{r}] \mid \dim \tilde{H}^{i-1}(\Sigma_\lambda) \neq 0\}. \tag{B.5}$$

If $x^v \in S_\beta$ has degree $\beta \in \text{Cl}(V)$, then x^v defines a map of vector spaces

$$H^i(\mathcal{O}_V(\alpha - \beta)) \xrightarrow{x^v} H^i(\mathcal{O}_V(\alpha)). \tag{B.6}$$

Suppose $\lambda \in \Sigma^i$ and $x^u \in \mathcal{B}(\lambda, \alpha - \beta)$. Let $\mu = \text{neg}(v + u)$. Notice that $\mu \subset \lambda$, since all entries of v are non-negative, so $\Sigma_\mu \subset \Sigma_\lambda$, and therefore there is a natural map

$$\pi : \tilde{H}^{i-1}(\Sigma_\lambda) \longrightarrow \tilde{H}^{i-1}(\Sigma_\mu). \tag{B.7}$$

With the identification in (B.4), the map (B.6) sends

$$x^u \otimes e \in \text{span}(\lambda, \alpha - \beta) \otimes \tilde{H}^{i-1}(\Sigma_\lambda) \tag{B.8}$$

to

$$x^{u+v} \otimes \pi(e). \tag{B.9}$$

In the examples that we consider in this appendix, it turns out that $\lambda \in \Sigma^1$ if and only if $\dim \tilde{H}^0(\Sigma_\lambda) = 1$. In this case, a basis for $H^1(\mathcal{O}_V(\alpha))$ is given by the set of monomials

$$\mathcal{B}(\alpha) := \bigcup_{\lambda \in \Sigma^1} \mathcal{B}(\lambda, \alpha). \tag{B.10}$$

B.2 The explicit example

Recall the example of interest: let Y be any smooth toric variety, let $\alpha \in \text{Pic}(Y)$ be any effective divisor class, and let $F = F_{2,2,\alpha}$ be a generic polynomial in the Cox ring S of $V = \mathbb{P}^1 \times \mathbb{P}^1 \times Y$ of the given triple degree. Let $X \subset V$ be the zero locus of F , and assume that X is smooth.

Since $\text{Cl}(V) = \text{Pic}(V) = \mathbb{Z}^2 \oplus \text{Pic}(Y)$, we write degrees as triples (a, b, γ) , where $a, b \in \mathbb{Z}$ and $\gamma \in \text{Pic}(Y)$. The Cox ring of V can be written as $S = R[x_1, x_2, x_3, x_4]$, where R is the Cox ring of Y , and degrees (over S) are: $\deg(x_1) = \deg(x_2) = (1, 0, 0)$, $\deg(x_3) = \deg(x_4) = (0, 1, 0)$, and if $f \in R_\gamma$, then $\deg_S(f) = (0, 0, \gamma)$.

Given $(a, b, \gamma) \in \text{Pic}(V)$, where $a, b \in \mathbb{Z}$, and $\gamma \in \text{Pic}(Y)$, we obtain the pullback $\mathcal{O}_X(a, b, \gamma)$ on X . Using the long exact sequence in cohomology arising from the short exact sequence

$$0 \longrightarrow \mathcal{O}_V(a - 2, b - 2, \gamma - \alpha) \xrightarrow{F} \mathcal{O}_V(a, b, \gamma) \longrightarrow \mathcal{O}_X(a, b, \gamma) \longrightarrow 0, \tag{B.11}$$

it follows that (a, b, γ) is an effective divisor class on X if and only if either it is an effective divisor class on V , or the kernel of the map

$$\phi_{a,b,\gamma} : H^1(\mathcal{O}_V(a - 2, b - 2, \gamma - \alpha)) \longrightarrow H^1(\mathcal{O}_V(a, b, \gamma)) \tag{B.12}$$

is nonzero.

Our strategy for proving Theorem 1 is the following. We fix $a > 0$, and identify bases for the source and target H^1 modules for the map $\phi_{-a,a,\gamma}$. We sum over all γ , obtaining a map $\phi_{-a,a,*}$ of graded R -modules, which turns out to be of size $(a^2 - 1) \times (a^2 - 1)$, i.e. a square matrix. We show that $\det(\phi_{-a,a,*}) \neq 0$, which implies that for all γ the kernel of

$\phi_{-a,a,\gamma}$ is zero, and so $(-a, a, \gamma)$ is not effective on X .¹⁶ We show that the determinant is nonzero by choosing a specialization of F , and bases of the source and target, such that $\phi_{-a,a,*}$ is in block diagonal form, and each block is (at worst) a tridiagonal matrix, whose determinant is nonzero.

Proof of Theorem 1. Define

$$H^1(\mathcal{O}_V(a, b, *)) := \bigoplus_{\gamma \in \text{Pic}(Y)} H^1(\mathcal{O}_V(a, b, \gamma)). \quad (\text{B.13})$$

This is a finitely generated graded R -module. Let $\phi_{a,b,*} := \bigoplus_{\gamma \in \text{Pic}(Y)} \phi_{a,b,\gamma}$. Then

$$\phi_{a,b,*} : H^1(\mathcal{O}_V(a-2, b-2, *)) \longrightarrow H^1(\mathcal{O}_V(a, b, *)) \quad (\text{B.14})$$

is a graded R -module map, whose entries (once a basis is chosen) all lie in R_α .

The set Σ_V^1 introduced in (B.5) is

$$\Sigma_V^1 = \{\{1, 2\}, \{3, 4\}\} \cup \Sigma_Y^1, \quad (\text{B.15})$$

where Σ_Y^1 is a set of subsets only involving the indices of the variables in R .

Fixing $a > 0$, we will now consider degrees of the form $(-a, b, \gamma)$. Given an integer $a > 0$ and an integer b , define the set of (Laurent) monomials

$$\mathcal{B}_{a,b} := \left\{ \frac{x_3^k x_4^\ell}{x_1^i x_2^j} \mid k + \ell = b, i + j = a, k, \ell \geq 0, i, j \geq 1 \right\}. \quad (\text{B.16})$$

Lemma 2 For $a > 0$, $\mathcal{B}_{a,b}$ is a (free) basis over R of $H^1(\mathcal{O}_V(-a, b, *))$. If $b < 0$, then $H^1(\mathcal{O}_V(-a, b, *)) = 0$.

Proof. This follows from the discussion in the last subsection by the observation that if $d = (-a, b, \gamma)$, for $a > 0$, then there is only one $\lambda \in \Sigma^1$ for which $\mathcal{B}(\lambda, d) \neq \emptyset$, namely $\lambda = \{1, 2\}$. In this case, Σ_λ consists of two disjoint points, and so $\tilde{H}^0(\Sigma_\lambda, \mathbb{C}) = \mathbb{C}$. A basis of $H^1(\mathcal{O}_V(-a, b, \gamma))$ is given by the Laurent monomials in $\mathcal{B}_{a,b}$, multiplied by any monomial in R_γ . \square

Note that for $a > 0$ and $b \geq 0$, we have $\dim_{\mathbb{C}} H^1(\mathcal{O}_V(-a, b, \gamma)) = (a-1)(b+1) \dim R_\gamma$ and $\text{rank}_R H^1(\mathcal{O}_V(-a, b, *)) = (a-1)(b+1)$. Also note that, for $a > 0$,

$$\dim_{\mathbb{C}} H^1(\mathcal{O}_V(-a-2, b-2, \gamma)) = \begin{cases} 0 & \text{if } b \leq 1 \\ (a+1)(b-1) \dim R_\gamma & \text{if } b \geq 1 \end{cases} \quad (\text{B.17})$$

Computing the matrix of the map

$$\phi_{-a,b,*} : H^1(\mathcal{O}_V(-a-2, b-2, *)) \longrightarrow H^1(\mathcal{O}_V(-a, b, *)) \quad (\text{B.18})$$

¹⁶The determinant of a square matrix M over an integral domain R — such as a polynomial ring — vanishes if and only if M has nonvanishing kernel. To see this, we work over the field of fractions of R , where from linear algebra one has $\det(M) = 0 \Leftrightarrow \ker(M) \neq 0$. Clearing denominators then gives the desired result for R .

induced by multiplication by the element $F \in S_{2,2,\alpha}$ with respect to these bases is straightforward: if x^u is a monomial in $S_{2,2,0}$, and m is a monomial in $B_{-a-2,b-2}$, then m maps to either the monomial $x^u m$, if that is in $\mathcal{B}_{-a,b}$, or 0, if not. By linearity, we obtain the matrix for the map (B.18).

The following proposition is all that is needed to finish the proof of Theorem 1.

Proposition 3 *Fix $a > 0$. Then $\phi_{-a,a,*}$ is a graded matrix of size $(a^2 - 1) \times (a^2 - 1)$ over the ring R , and for F in a dense Zariski-open subset of $S_{2,2,\alpha}$, this determinant is nonzero.*

Proof. Notice that a basis (over R) for the source of the map $\phi_{-a,a,*}$ is $\mathcal{B}_{a+2,a-2}$, a basis for the target is $\mathcal{B}_{a,a}$, and by inspection both of these have rank $a^2 - 1$.

Let $F \in S_{2,2,\alpha}$. We will show that the matrix $\phi_{-a,a,*}$ (with entries in R) has nonzero determinant, for at least one F of this degree, which implies that the determinant is nonzero for a Zariski-open subset of $F \in S_{2,2,\alpha}$, proving the result.

To do this, we choose a specific F of the form

$$F = x_1^2 x_3^2 A + x_1 x_2 x_3 x_4 B + x_2^2 x_4^2 C, \tag{B.19}$$

where $A, B, C \in R_\alpha$ are chosen generically (actually, if x^v is any monomial in R_α , then we could choose A, B, C to be \mathbb{C} -multiples of this monomial).

We now partition the bases of the source and target of $\phi_{-a,a,*}$ in a manner which will place the resulting matrix in block form, with blocks that are square matrices whose determinants are evidently nonzero. This is enough to prove the result.

To this end, let $t = \frac{x_2 x_4}{x_1 x_3}$, which allows us to write F as:

$$F = x_1 x_2 x_3 x_4 (t^{-1} A + B + t C). \tag{B.20}$$

We define an equivalence relation on the set of monomials $m \in \mathcal{B}_{a,b}$ by setting $m \sim n$ if there is an $\ell \in \mathbb{Z}$ such that $n = t^\ell m$. Define $\mathcal{B}_{a,b,m} := \{n \in \mathcal{B}_{a,b} \mid n \sim m\}$. Let $\widehat{\mathcal{B}}_{a,b} := \{m \in \mathcal{B}_{a,b} \mid mt^{-1} \notin \mathcal{B}_{a,b}\}$. It is easy to check that $\widehat{\mathcal{B}}_{a,b}$ is a full set of representatives under this equivalence relation, and that if $m \in \widehat{\mathcal{B}}_{a,b}$, then $\mathcal{B}_{a,b,m} = \{m, tm, t^2 m, \dots, t^i m\}$, for some integer i depending on m (and also on a and b). It is also clear that $\phi_{-a,b,*}$ maps the span of $\mathcal{B}_{a+2,b+2,m}$ into the span of $\mathcal{B}_{a,b,x_1 x_2 x_3 x_4 m}$, and so becomes block diagonal with these partitions of the bases. The matrix of F , on each m -set, has a tridiagonal structure. When $a = b$, in fact, one checks that each block is a square matrix, and is (at most) tridiagonal, with determinant a nonzero polynomial in A, B , and C , thus proving the proposition and hence the theorem. \square

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Witten, *Nonperturbative superpotentials in string theory*, *Nucl. Phys. B* **474** (1996) 343 [[hep-th/9604030](#)] [[INSPIRE](#)].
- [2] M. Mariño, R. Minasian, G.W. Moore and A. Strominger, *Nonlinear instantons from supersymmetric p -branes*, *JHEP* **01** (2000) 005 [[hep-th/9911206](#)] [[INSPIRE](#)].
- [3] H. Federer and W.H. Fleming, *Normal and integral currents*, *Annals Math.* **72** (1960) 458.
- [4] F.J. Almgren Jr., *Q valued functions minimizing Dirichlet's integral and the regularity of area minimizing rectifiable currents up to codimension two*, *Bull. Amer. Math. Soc.* **8** (1983) 327.
- [5] C. Beasley and E. Witten, *New instanton effects in string theory*, *JHEP* **02** (2006) 060 [[hep-th/0512039](#)] [[INSPIRE](#)].
- [6] M. Cvetič, R. Richter and T. Weigand, *(Non-)BPS bound states and D-brane instantons*, *JHEP* **07** (2008) 012 [[arXiv:0803.2513](#)] [[INSPIRE](#)].
- [7] I. Garcia-Etxebarria, F. Marchesano and A.M. Uranga, *Non-perturbative F -terms across lines of BPS stability*, *JHEP* **07** (2008) 028 [[arXiv:0805.0713](#)] [[INSPIRE](#)].
- [8] M. Buican and S. Franco, *SUSY breaking mediation by D-brane instantons*, *JHEP* **12** (2008) 030 [[arXiv:0806.1964](#)] [[INSPIRE](#)].
- [9] R. Blumenhagen, M. Cvetič, S. Kachru and T. Weigand, *D-brane instantons in type II orientifolds*, *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 269 [[arXiv:0902.3251](#)].
- [10] M. Micallef and J. Wolfson, *Area minimizers in a $K3$ surface and holomorphicity*, *Geom. Funct. Anal.* **16** (2006) 437.
- [11] H. Federer, *Geometric measure theory*, Grundlehren der mathematischen Wissenschaften, Springer, Berlin, Heidelberg, Germany (1969).
- [12] H.B. Lawson and J. Simons, *On stable currents and their application to global problems in real and complex geometry*, *Annals Math.* **98** (1973) 427.
- [13] S.X.-D. Chang, *Two dimensional area minimizing integral currents are classical minimal surfaces*, *J. Amer. Math. Soc.* **1** (1988) 699.
- [14] F. Almgren, *Questions and answers about area-minimizing surfaces and geometric measure theory*, in *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, U.S.A., 1990)*, *Proc. Sympos. Pure Math.* **54**, Amer. Math. Soc., Providence, RI, U.S.A. (1993), pg. 29.
- [15] C. Arezzo and G.L. Nave, *Minimal two spheres in Kähler-Einstein Fano manifolds*, *Adv. Math.* **191** (2005) 209.
- [16] C. Arezzo and J. Sun, *A variational characterization of J -holomorphic curves in symplectic manifolds*, [arXiv:1211.7016](#).
- [17] L. Foscolo, *ALF gravitational instantons and collapsing Ricci-flat metrics on the $K3$ surface*, [arXiv:1603.06315](#) [[INSPIRE](#)].

- [18] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The string landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060 [[hep-th/0601001](#)] [[INSPIRE](#)].
- [19] H. Ooguri and C. Vafa, *Non-supersymmetric AdS and the swampland*, *Adv. Theor. Math. Phys.* **21** (2017) 1787 [[arXiv:1610.01533](#)] [[INSPIRE](#)].
- [20] B. Heidenreich, M. Reece and T. Rudelius, *Evidence for a sublattice weak gravity conjecture*, *JHEP* **08** (2017) 025 [[arXiv:1606.08437](#)] [[INSPIRE](#)].
- [21] F. Denef, M.R. Douglas, B. Florea, A. Grassi and S. Kachru, *Fixing all moduli in a simple F-theory compactification*, *Adv. Theor. Math. Phys.* **9** (2005) 861 [[hep-th/0503124](#)] [[INSPIRE](#)].
- [22] T. Weigand, *F-theory*, *PoS(TASI2017)016* (2018) [[arXiv:1806.01854](#)] [[INSPIRE](#)].
- [23] B. Heidenreich, M. Reece and T. Rudelius, *Sharpening the weak gravity conjecture with dimensional reduction*, *JHEP* **02** (2016) 140 [[arXiv:1509.06374](#)] [[INSPIRE](#)].
- [24] M. Montero, G. Shiu and P. Soler, *The weak gravity conjecture in three dimensions*, *JHEP* **10** (2016) 159 [[arXiv:1606.08438](#)] [[INSPIRE](#)].
- [25] C. Cheung and G.N. Remmen, *Naturalness and the weak gravity conjecture*, *Phys. Rev. Lett.* **113** (2014) 051601 [[arXiv:1402.2287](#)] [[INSPIRE](#)].
- [26] T. Rudelius, *Constraints on axion inflation from the weak gravity conjecture*, *JCAP* **09** (2015) 020 [[arXiv:1503.00795](#)] [[INSPIRE](#)].
- [27] J. Brown, W. Cottrell, G. Shiu and P. Soler, *Fencing in the swampland: quantum gravity constraints on large field inflation*, *JHEP* **10** (2015) 023 [[arXiv:1503.04783](#)] [[INSPIRE](#)].
- [28] A. Hebecker, P. Mangat, F. Rompineve and L.T. Witkowski, *Winding out of the swamp: evading the weak gravity conjecture with F-term winding inflation?*, *Phys. Lett. B* **748** (2015) 455 [[arXiv:1503.07912](#)] [[INSPIRE](#)].
- [29] J. Brown, W. Cottrell, G. Shiu and P. Soler, *On axionic field ranges, loopholes and the weak gravity conjecture*, *JHEP* **04** (2016) 017 [[arXiv:1504.00659](#)] [[INSPIRE](#)].
- [30] A. Hebecker, F. Rompineve and A. Westphal, *Axion monodromy and the weak gravity conjecture*, *JHEP* **04** (2016) 157 [[arXiv:1512.03768](#)] [[INSPIRE](#)].
- [31] G. Shiu, P. Soler and W. Cottrell, *Weak gravity conjecture and extremal black holes*, *Sci. China Phys. Mech. Astron.* **62** (2019) 110412 [[arXiv:1611.06270](#)] [[INSPIRE](#)].
- [32] B. Heidenreich, M. Reece and T. Rudelius, *The weak gravity conjecture and emergence from an ultraviolet cutoff*, *Eur. Phys. J. C* **78** (2018) 337 [[arXiv:1712.01868](#)] [[INSPIRE](#)].
- [33] Y. Hamada and G. Shiu, *Weak gravity conjecture, multiple point principle and the Standard Model landscape*, *JHEP* **11** (2017) 043 [[arXiv:1707.06326](#)] [[INSPIRE](#)].
- [34] C. Cheung, J. Liu and G.N. Remmen, *Proof of the weak gravity conjecture from black hole entropy*, *JHEP* **10** (2018) 004 [[arXiv:1801.08546](#)] [[INSPIRE](#)].
- [35] S. Andriolo, D. Junghans, T. Noumi and G. Shiu, *A tower weak gravity conjecture from infrared consistency*, *Fortsch. Phys.* **66** (2018) 1800020 [[arXiv:1802.04287](#)] [[INSPIRE](#)].
- [36] B. Heidenreich, M. Reece and T. Rudelius, *Emergence of weak coupling at large distance in quantum gravity*, *Phys. Rev. Lett.* **121** (2018) 051601 [[arXiv:1802.08698](#)] [[INSPIRE](#)].
- [37] Y. Hamada, T. Noumi and G. Shiu, *Weak gravity conjecture from unitarity and causality*, *Phys. Rev. Lett.* **123** (2019) 051601 [[arXiv:1810.03637](#)] [[INSPIRE](#)].

- [38] T.W. Grimm, E. Palti and I. Valenzuela, *Infinite distances in field space and massless towers of states*, *JHEP* **08** (2018) 143 [[arXiv:1802.08264](#)] [[INSPIRE](#)].
- [39] S.-J. Lee, W. Lerche and T. Weigand, *Tensionless strings and the weak gravity conjecture*, *JHEP* **10** (2018) 164 [[arXiv:1808.05958](#)] [[INSPIRE](#)].
- [40] S.-J. Lee, W. Lerche and T. Weigand, *A stringy test of the scalar weak gravity conjecture*, *Nucl. Phys. B* **938** (2019) 321 [[arXiv:1810.05169](#)] [[INSPIRE](#)].
- [41] T.W. Grimm and D. Van De Heisteeg, *Infinite distances and the axion weak gravity conjecture*, [arXiv:1905.00901](#) [[INSPIRE](#)].
- [42] T.W. Grimm, F. Ruehle and D. van de Heisteeg, *Classifying Calabi-Yau threefolds using infinite distance limits*, [arXiv:1910.02963](#) [[INSPIRE](#)].
- [43] M. Demirtas, C. Long, L. McAllister and M. Stillman, *Effective divisors of Calabi-Yau hypersurfaces*, work in progress.
- [44] W. Fulton, R. Macpherson, F. Sottile and B. Sturmfels, *Intersection theory on spherical varieties*, *J. Alg. Geom.* **4** (2000) 181.
- [45] M. Demirtas, C. Long, L. McAllister and M. Stillman, *The Kreuzer-Skarke axiverse*, [arXiv:1808.01282](#) [[INSPIRE](#)].
- [46] E. Palti, *The weak gravity conjecture and scalar fields*, *JHEP* **08** (2017) 034 [[arXiv:1705.04328](#)] [[INSPIRE](#)].
- [47] A. Collinucci, *New F-theory lifts*, *JHEP* **08** (2009) 076 [[arXiv:0812.0175](#)] [[INSPIRE](#)].
- [48] A. Collinucci, *New F-theory lifts. II. Permutation orientifolds and enhanced singularities*, *JHEP* **04** (2010) 076 [[arXiv:0906.0003](#)] [[INSPIRE](#)].
- [49] M. Esole, P. Jefferson and M.J. Kang, *Euler characteristics of crepant resolutions of Weierstrass models*, *Commun. Math. Phys.* **371** (2019) 99 [[arXiv:1703.00905](#)] [[INSPIRE](#)].
- [50] D. Eisenbud, M. Mustața and M. Stillman, *Cohomology on toric varieties and local cohomology with monomial supports*, *J. Symbol. Comput.* **29** (2000) 583.
- [51] D. Maclagan and G.G. Smith, *Multigraded Castelnuovo-Mumford regularity*, *J. Reine Angew. Math.* **571** (2004) 179.
- [52] R. Blumenhagen, B. Jurke, T. Rahn and H. Roschy, *Cohomology of line bundles: a computational algorithm*, *J. Math. Phys.* **51** (2010) 103525 [[arXiv:1003.5217](#)] [[INSPIRE](#)].