

RESEARCH ARTICLE

miR-71 and miR-263 Jointly Regulate Target Genes *Chitin synthase* and *Chitinase* to Control Locust Molting

Meiling Yang¹✉, Yanli Wang¹✉, Feng Jiang², Tianqi Song¹, Huimin Wang², Qing Liu¹, Jie Zhang², Jianzhen Zhang^{1*}, Le Kang^{1,2*}

1 Institute of Applied Biology, Shanxi University, Taiyuan, Shanxi, China and State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing, China, **2** Beijing Institutes of Life Science, Chinese Academy of Sciences, Beijing, China

✉ These authors contributed equally to this work.

* zjz@sxu.edu.cn (JiaZ); lkang@ioz.ac.cn (LK)



CrossMark
click for updates

 OPEN ACCESS

Citation: Yang M, Wang Y, Jiang F, Song T, Wang H, Liu Q, et al. (2016) miR-71 and miR-263 Jointly Regulate Target Genes *Chitin synthase* and *Chitinase* to Control Locust Molting. *PLoS Genet* 12(8): e1006257. doi:10.1371/journal.pgen.1006257

Editor: Lynn M Riddiford, Howard Hughes Medical Institute, UNITED STATES

Received: March 30, 2016

Accepted: July 23, 2016

Published: August 17, 2016

Copyright: © 2016 Yang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB11010000), National Basic Research Program of China (2012CB114102) and National Natural Science Foundation of China Grants (31472051, 31301915, 31430023, 31272380). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Chitin synthase and chitinase play crucial roles in chitin biosynthesis and degradation during insect molting. Silencing of Dicer-1 results in reduced levels of mature miRNAs and severely blocks molting in the migratory locust. However, the regulatory mechanism of miRNAs in the molting process of locusts has remained elusive. In this study, we found that in chitin metabolism, two crucial enzymes, chitin synthase (CHS) and chitinase (CHT) were regulated by miR-71 and miR-263 during nymph molting. The coding sequence of *CHS1* and the 3'-untranslated region of *CHT10* contain functional binding sites for miR-71 and miR-263, respectively. miR-71/miR-263 displayed cellular co-localization with their target genes in epidermal cells and directly interacted with *CHS1* and *CHT10* in the locust integument, respectively. Injections of miR-71 and miR-263 agomirs suppressed the expression of *CHS1* and *CHT10*, which consequently altered chitin production of new and old cuticles and resulted in a molting-defective phenotype in locusts. Unexpectedly, reduced expression of miR-71 and miR-263 increased *CHS1* and *CHT10* mRNA expression and led to molting defects similar to those induced by miRNA delivery. This study reveals a novel function and balancing modulation pattern of two miRNAs in chitin biosynthesis and degradation, and it provides insight into the underlying molecular mechanisms of the molting process in locusts.

Author Summary

Molting is a crucial process in the growth and development in insects. Disturbing the molting process represents an attractive strategy for developing safe and effective insecticides. The migratory locust is a hemimetabolous pest that undergoes five molting stages in its life cycle. Similar molting defects can be observed in expression silencing of the key genes both in miRNA processing and in chitin metabolism. However, any link between a specific miRNA to chitin metabolism has not yet been established. In this study, we

Competing Interests: The authors have declared that no competing interests exist.

elucidated a mechanism by which two miRNAs regulate chitin metabolism related to the molting process. We found that miR-71 and miR-263 directly repress two genes, *chitin synthase1* (*CHS1*) and *chitinase10* (*CHT10*), which are required for chitin biosynthesis and degradation in chitin metabolism. Manipulation of miR-71 and miR-263 expression blocked molting and resulted in abnormal molting by negatively regulating the expression of *LmCHS1* and *LmCHT10*. Furthermore, both up-regulation and down-regulation of *LmCHS1* and *LmCHT10* by miRNA manipulation altered the chitin content of the new cuticle and old cuticles, leading to a similar aberrant molting phenotype. Our results demonstrate a balancing modulation pattern of two miRNAs in chitin biosynthesis and degradation that controlled the precise molting process in locusts.

Introduction

Molting is a crucial process in insect growth and development [1,2]. Chitin, as a vital component of the cuticle of the epidermis, plays key roles in maintaining morphology and the molting process [3]. Because chitin is absent in plants and vertebrates, and insect growth and development are strictly dependent on chitin biosynthesis and degradation, chitin metabolism represents an attractive target for developing safe and effective insecticides [4].

The migratory locust *Locusta migratoria*, a worldwide insect pest species, undergoes five molting stages in its life cycle [5,6]. The chitin-mediated molting process is considered to depend on two crucial genes, *chitin synthase* (*CHS*) and *chitinase* (*CHT*), which are regulated by molting hormone 20-hydroxyecdysone (20E) and juvenile hormone [7,8,9,10,11]. Chitin synthases are the key regulatory enzymes for chitin synthesis in insects and represent a specific target of insecticides [12]. The *LmCHS1* gene cloned from the migratory locust is expressed specifically in the epidermis during the molting stage. Knockdown of the *LmCHS1* gene increases the number of non-molting and abnormal molting nymphs [6]. However, another paralog *LmCHS2* contributes to the biosynthesis of chitin associated with the peritrophic matrix [13]. Moreover, chitinases are hydrolytic enzymes that are required for the degradation of glycosidic bonds of chitin [14]. *TcCHT10* prevents larval molting and plays a vital role during the molting process at all developmental stages; the other paralogs, *CHT5* and *CHT7*, prevent molting and wings from folding properly only in adults [15,16]. An interesting feature of *CHS1* and *CHT10* in locusts is that the abrupt increase and decrease in transcript levels at the end of every nymph stage (before molting) suggest that the two key enzymes are likely precisely modulated in the molting process. However, the underlying regulatory molecular mechanisms of enzyme-dependent chitin metabolism and the molting process have remained elusive.

MicroRNAs (miRNAs), small non-coding regulatory RNAs, have emerged as key posttranscriptional regulators of gene expression in multiple biological processes [17] because they can directly trigger translational repression or mRNA degradation by low complementary base-pairing with the 3'UTRs of the target genes [18,19]. However, recent studies have shown that miRNAs can extensively target the protein-coding region of mRNAs in animals or insects [20,21,22]. Many studies have shown that miRNAs critically affect the molting of insects, thus resulting in molting defect phenotypes. For example, miR-8-5p and miR-2a-3p negatively regulate membrane-bound trehalase and phosphoacetylglucosamine mutase of the chitin biosynthesis pathway, leading to a significant reduction in survival rate along with a molting defect phenotype in the hemipteran insect *Nilaparvata lugens* [23]. Several distinct miRNAs have been approved in the regulation of insect metamorphosis. The loss of miR-2 up-regulates Kr-

h1 mRNA, thereby leading to impaired metamorphosis [24,25]. Additionally, let-7 and miR-125 mutants induce temporal mis-regulation of specific metamorphic processes in *Drosophila* [26]. In the migratory locust, we reported that depletion of Dicer-1, the enzyme that catalyzes the final step of miRNA biosynthesis, induced a molting defect [27]. Results indicated that miRNAs play a crucial role in regulating the molting process of locusts. However, the mechanism regarding how miRNAs affect posttranscriptional modifications in the molting process has not yet been fully elucidated.

Considering that *CHS1* and *CHT10* are crucial molt-dependent enzymes that balance chitin metabolism in many insect species [15,16,28,29], we chose *CHS1* and *CHT10* as candidate genes. We hypothesized that miRNAs might play essential roles in the regulation of *CHS1*- and *CHT10*-mediated molting processes. In this study, we performed small RNA transcriptome sequencing to identify expressed miRNAs in the integument of locusts. We found that the integument-expressed miR-71 and miR-263 directly target the two key genes *CHS1* and *CHT10* and regulate chitin production during the molting process, resulting in the successful molting of the migratory locust. Our results reveal a molecular mechanism by which miRNAs play a role in balancing the modulation of *CHS1*- and *CHT10*-dependent chitin metabolism during molting.

Results

miRNA identification

To identify the miRNAs associated with molting, we sequenced a transcriptome of small RNAs of the locust integument, which is an important tissue during the molting process in insects. A total of 15,459,187 sequencing reads were obtained, of which 4,590,268 (29.7%) corresponding to mature and star strands were mapped to the known miRNA precursors of locusts [30]. Forty-five conserved miRNAs showed transcriptional activities (reads per million threshold 1) in the integument of locusts. Their expression levels varied over several orders of magnitude. The top ten most highly expressed miRNAs were miR-9b, miR-184, miR-14, miR-100, bantam, miR-71, miR-275, miR-305, miR-263 and miR-279b (Fig 1A). All of the expressed miRNAs were used for further miRNA candidate screening.

Potential miRNAs targeting *CHS1* and *CHT10*

CHS1 and *CHT10* involved in chitin metabolism have been confirmed to regulate the insect molting process [15,16,28,29]. Using the miRanda software, we predicted the expressed miRNAs that could potentially bind to *LmCHS1* and *LmCHT10*. Thirteen miRNAs exhibited potential target sites in the 3'UTR and CDS regions of *LmCHS1*, and 6 miRNAs possessed target sites located in the 3'UTR of *LmCHT10* in locusts (Fig 1B, S1 Table, and S2 Table). An additional prediction software, RNAhybrid, was used to further improve the target prediction efficiency. The RNAhybrid program also identified *LmCHS1* and *LmCHT10* as potential targets for miR-71 and miR-263, respectively (S1 Fig). Furthermore, we confirmed the absence of miR-71 and miR-263 binding sites in the *Tweedle*, *Cryptocephal*, *Obstructor*, *Knickkopf*, and *ecd1* genes to exclude the other possible miR-71/miR-263 targets [31,32,33,34,35], which can lead to molting defects similar to those caused by *LmCHS1* and *LmCHT10*.

To confirm the correlation of the expression pattern between *LmCHS1*, *LmCHT10* and their target miRNAs, we performed stem-loop quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR) to quantify the expression levels of these predicted miRNAs and target genes in the integument of second-instar nymphs (S2 Fig). The overall expression of miRNAs (miR-71 or miR-263) and that of the target genes (*LmCHS1* or *LmCHT10*) exhibited opposite patterns during the second nymph stage (Fig 1C and 1D). The miR-71 expression




Fig 1. miR-71 and miR-263 may target chitin synthase (*CHS1*) and chitinase (*CHT10*), two key enzymes of chitin metabolism. (A) miRNAs identified in the locust integument by small RNA transcriptome. Only the miRNAs with high/moderate expression level (reads per million, RPM>15) are shown. (B) miR-71 and miR-263 target sites were predicted in the *CHS1* and *CHT10* genes of *Locusta migratoria*. (C) The expression levels of *CHS1* and *CHT10* were determined in the integuments every day (D1–5: Days 1–5) during the second-instar nymph stage using qPCR. (D) The expression levels of miR-71 and miR-263 were determined in the integuments every day (D1–5: Days 1–5) during the second-instar nymph stage using qPCR. The qPCR data are presented as means \pm SEM ($n = 6$); the same letter indicates data that are not significantly different.

doi:10.1371/journal.pgen.1006257.g001

levels showed the opposite wave-like pattern of miR-263 expression levels, with the highest level occurring at the mid-stage for miR-71, whereas miR-263 expression decreased to the lowest level (Fig 1D). In contrast, the mRNA expression of *LmCHS1* was down-regulated at the mid-stage and up-regulated at the early and late stages. However, the mRNA expression of *LmCHT10* was suppressed at the early and late stages and was promoted at the mid-stage (Fig 1C). These data indicate that miR-71/miR263 expression is negatively correlated with *LmCHS1* and *LmCHT10* expression during new integument formation in the nymph stages. The results imply that there is a possible regulatory relationship between the miRNAs and the genes.

miR-71 and miR-263 regulate *LmCHS1* and *LmCHT10*, respectively

To confirm the interactions of miR-71, miR-263 and their targeting genes *in vitro*, we performed reporter assays using luciferase constructs fused to the coding region of *LmCHS1* and the 3'UTR of *LmCHT10*. Compared with the agomir control (agomir-NC), the constructs with either the *LmCHS1* or *LmCHT10* binding sites produced lower luciferase activity when co-transfected with miR-71 or miR-263 agomir, respectively, in S2 cells (Fig 2A and 2B). When the regions homologous to the “seed” sequence of miR-71 and miR-263 were mutated in the *LmCHS1* and *LmCHT10* reporter constructs, the luciferase activity returned to levels similar to those produced by mock transfection with the empty reporter plasmid (Fig 2A and 2B). However, the luciferase activity of sites transfected with miR-252, whose expression is negatively correlated with *LmCHS1*, showed no change compared with the control (S3 Fig). To further validate the effect of endogenous miR-71 and miR-263 in S2 cells on the luciferase activity, we investigated miRNA-71 and miR-263 levels in S2 cells. The miR-71 homolog was not detected




Fig 2. Interactions between miRNAs (miR-71 and miR-263) and their targets *CHS1* and *CHT10* in locusts. (A, B) Luciferase reporter assays were analyzed in S2 cells co-transfected with miR-71 or miR-263 overexpression vectors and psi-CHECK2 vectors containing wild-type (WT) or mutant (MT, the 8 nt of the region that corresponds to the miRNA seed mutated) target gene sequences of *CHS1* (A) and *CHT10* (B) ($n = 6$). (C, D) RIP was performed with an anti-Ago-1 antibody. qPCR analysis was performed to amplify the *CHS1* and *CHT10* mRNAs from the Ago-1 immunoprecipitates from extracts of integument tissue treated with agomir-71 or agomir-263 48 h later compared with the agomir controls (agomir-NC). The data for the luciferase activities and qPCR analyses are presented as means \pm SEM ($n = 6$). * $p < 0.05$; ** $p < 0.01$. (E) miRNAs and their target genes were co-labeled to determine the co-localization between these molecules in the locust integument on the third day and the fifth day of the 2nd instar stage by FISH. Where green (*CHS1* and *CHT10*) and red signals (miR-71 and miR-263) overlap, a yellow signal is observed, indicating the co-localization of miRNAs and their targets. Control 1, scrambled miRNA and *CHS1* antisense probe; Control 2, scrambled miRNA and *CHT10* antisense probe; Control 3, scrambled miRNA, *CHS1* and *CHT10* sense probes. The images were visualized using an LSM 710 confocal fluorescence microscope (Zeiss) at a magnification of 40X.

doi:10.1371/journal.pgen.1006257.g002

in the fly genome [36]. The small RNA transcriptome data indicated that only a few reads for miR-263 (7 counts in GEO accession GSM272651 and 1 count in GEO accession GSM272652) were detected in S2 cells, implying a limited expression of miR-263 in S2 cells. We examined luciferase activity in S2 cells with antagomir-263. The luciferase signals of the *CHT10* construct incubated with antagomir-263 did not vary significantly compared with those of the control (S4 Fig). The data suggested that the endogenously expressed miR-263 did not affect the luciferase assay results for the locust miR-263. Thus, the predicted miRNA binding sites in *LmCHS1* and *LmCHT10* are functional and might be targeted by miR-71 and miR-263, respectively, in S2 cells.

Ago1, as a RNA binding protein, is a core component of RISC involved in miRNA-mediated gene silencing. Anti-Ago1 RIP is a biochemical approach to identify the composition and organization of endogenous mRNAs, miRNAs associated with Ago1 proteins. This approach is widely used in interaction validation between miRNA and its target *in vivo*. We then performed an RNA immunoprecipitation assay in the integument to examine the interactions of miR-71 and miR-263 with their targeting genes *in vivo* (Fig 2C and 2D). *LmCHS1* or *LmCHT10* were significantly enriched in the Ago1-immunoprecipitated RNAs from the integuments treated with agomir-71 or agomir-263 compared with those treated with agomir-NC. These results indicated that miR-71 and miR-263 directly regulate *LmCHS1* and *LmCHT10* in the locust integument, respectively.

Co-localization of the miRNAs and targeting genes

To determine whether miR-71/miR-263 were co-localized in the locust integument, we performed *in situ* analyses of miRNA-71/miR-263 and their targets by miRNA/mRNA fluorescence *in situ* hybridization (FISH). Indeed, we found that miR-71 and *LmCHS1* as well as miR-263 and *LmCHT10* were both widely detected in the epidermal cells of the locust integument (Fig 2E). Specifically, miR-71 is co-localized with *LmCHS1* and miR-263 is co-localized with *LmCHT10* in cells of the integument. The results suggest that in the locust integument, *LmCHS1* and *LmCHT10* interact directly with miR-71 and miR-263, respectively, in a spatial manner.

miR-71 controls chitin synthesis by targeting *LmCHS1*, and miR-263 inhibits chitin degradation by targeting *LmCHT10*

To determine the effects of miR-71 and miR-263 on their target genes *in vivo*, we detected the expression levels of *LmCHS1* and *LmCHT10* after miRNA agomir (overexpression) or antagomir (knockdown) administration in the locust integument. We first assessed the miRNA expression changes by injecting locusts with miRNA agomir or antagomir *in vivo* to confirm the delivery efficiency of miRNA administration. The qPCR results showed that miR-71 and miR-263 levels were significantly induced and depleted by their agomir and antagomir treatments, respectively. As expected, the treatment had no effect on the expression of the negative control, let-7 (S5 Fig). Moreover, the mRNA levels of *LmCHS1* decreased by approximately 60% compared to those in the control locusts after agomir-71 injection (Fig 3A). Additionally, antagomir-71 injection resulted in a significant up-regulation of the mRNA expression level of *LmCHS1* (Fig 3B). In contrast, inhibition of approximately 55% of *LmCHT10* expression was observed after agomir-263 injection. Additionally, the mRNA levels of *LmCHT10* were up-regulated by miR-263 knockdown (Fig 3A and 3B). No significant effects on the expression of paralog genes *LmCHS2* and *LmCHT5* upon miR-71/miR-263 administration were observed, suggesting that the agomir/antagomir injection specifically acted on the target genes (S6 Fig). Additionally, we sought to determine whether the administration of miR-71 had any effects on




Fig 3. Effect of miR-71 and miR-263 administration on *CHS1* and *CHT10* expression and chitin content in the locust integument. (A, B) The effects of 210 pmol of agomir- and antagomir-71 or agomir- and antagomir-263 treatment for 48 h after injection on the mRNA expression levels of *CHS1* and *CHT10* in locust integuments were studied using qPCR. (C, D) Chitin production was evaluated in the integuments of locusts after treatment with 210 pmol agomir- or antagomir-71/agomir- or antagomir-263 treatment. The qPCR and chitin content data are presented as means \pm SEM ($n = 6$). * $p < 0.05$; ** $p < 0.01$.

doi:10.1371/journal.pgen.1006257.g003

the expression of miR-263, or vice versa. The results indicated that there was no interaction between miR-71 and miR-263, which were involved in the regulation of two distinct processes (chitin synthesis and degradation) (S7 Fig). Since 20E is believed to primarily regulate insect growth and development processes, including molting [2,37], we further used 20E treatments to investigate whether 20E might play a relevant role in the regulation of the expression of miR-71 and miR-263. The results indicated that 20E treatment depressed the expression levels of miR-263, but did not have significant effects on miR-71 expression level (S8 Fig).

Because *LmCHS1* and *LmCHT10* are essential genes in chitin synthesis and degradation, we determined the effects of miR-71 and miR-263 on chitin production in the integuments after miRNA administration *in vivo*. The administration of the miR-71 agomir reduced chitin production by approximately 48% (Fig 3C). In contrast, miR-71 knockdown significantly

increased the chitin content (by approximately 18%, $p < 0.05$) of the locusts. Accordingly, a significant increase in chitin content was observed after agomir-263 manipulation, and antagomir-263 injection decreased chitin content (Fig 3D). Thus, administration of miR-71 and miR-263 may have profound regulatory effects on chitin production and content *in vivo* in locusts.

miR-71 and miR-263 regulate the molting process of locusts by inhibiting chitin metabolism

To determine the function of miR-71 and miR-263 during the molting process, we monitored the molting of locusts after agomir-71 or agomir-263 injection, respectively. The locusts injected with agomir-71 or agomir-263 displayed a distinct molting defect phenotype. In total, of 25 nymphs injected with agomir-71, 12 (48%) died during the molting process from second instar to third instar, whereas only 8.7% (2 out of 23) of the control nymphs died during this process (Fig 4A). Similarly, after injection of agomir-263, the mortality reached 40.7%, which was significantly higher than that of the control (only 7.4%) during the molting process (Fig 4C). In parallel, miR-71 and miR263 knockdown caused by injection of antagomir-71 or antagomir-263 resulted in incomplete ecdysis, with 24.0% and 20.7% mortality, respectively (Fig 4B and 4D).

After nymphs were injected with the miR-71 and miR-263 agomir or antagomir, the nymphs exhibited abnormal and unsuccessful molting (Fig 4E–4H), in which a certain amount of the old cuticle was separated from the body but not detached from the body to any extent. Moreover, some nymphs showed a molting defect, in which legs failed to slough from the old cuticle, and other nymphs died without obvious molting defects due to molting arrest (Fig 4E–4H).

The chitin contents of new/old cuticle were regulated by miR-71 and miR-263 in molting

Treatment of locusts with agomir-71 and agomir-263 resulted in the down-regulation of *LmCHS1* and *LmCHT10* expression and a corresponding change in chitin content, thereby generating a significant molting defect (Fig 4I). To further determine whether the abnormal layer of the cuticle was responsible for the molting defect induced by miR-71 and miR-263, we performed hematoxylin and eosin staining and chitin staining in the integument by injecting agomir-71 or agomir-263 into the locusts (Fig 5A and 5C). A significant decrease in chitin content occurred in response to miR-71 overexpression in the newly formed cuticle, which exhibited a severe deficiency due to diminished chitin synthesis (Fig 5A). Similarly, RNAi for *LmCHS1* prevented the synthesis of cuticle chitin, as expected in the newly formed cuticle (Fig 5B and 5D). Conversely, miR-263 overexpression inhibited the degradation of the old cuticle, the chitin of which was not diminished compared with that of the agomir controls (Fig 5A and 5C). Consistent with the miR-263-mediated phenotype, knockdown of *LmCHT10* transcripts hindered chitin degradation of the old cuticle, leading to a dramatically thickened layer of the old cuticle and impeding the shedding of the old cuticle during molting (Fig 5B and 5D). Therefore, the change in chitin content of new/old cuticle regulated by miR-71 and miR-263 is a key mediator of defective molting.

Discussion

Our previous studies confirmed that depletion of Dicer-1 prevented the molting process [38] and chitin-metabolic genes, including *LmCHS* and *LmCHT*, are related to molting in the migratory locust [7]. The results presented herein explored the link between the Dicer-




Fig 4. miR-71 and miR-263 hinder molting of the migratory locust. (A-D) The effects of 210 pmol agomir-71 or agomir-263 and antagomir-71 or antagomir-263 treatment on the mortality of locusts were studied after injection. N, the number of animals treated. (E-H) The effects of 210 pmol agomir-71 or agomir-263 and antagomir-71 or antagomir-263 treatment on the molt phenotype of locusts were studied after injection. The red arrow indicates the shedding old cuticle in the defective molting process. OC, old cuticle, NC, new cuticle. (I) Model of the miR-71- and miR-263-mediated chitin metabolism pathway associated with the molting of the migratory locust. miR-71 controls chitin production by regulating *CHS1*, and miR-263 controls chitin degradation by modulating *CHT10* in the locust integument.

doi:10.1371/journal.pgen.1006257.g004




Fig 5. Effect of miR-71 and miR-263 administration on the chitin metabolism of the new and old cuticle. (A, B) Hematoxylin and eosin staining of the integument was performed by injecting agomir-71/agomir-263 (A) or by injecting dsRNA against *CHS1*/*CHT10* (B) into the locusts. (C, D) The chitin staining experiment in the integument was performed by injecting agomir-71/-263 (C) or injecting dsRNA against *CHS1*/*CHT10* (D) into the locusts. The chitin and cuticle thicknesses of the new cuticle/old cuticle were quantified using ZEN 2.1 software and expressed as means \pm SEM ($n = 4$). OC, old cuticle, NC, new cuticle. * $p < 0.05$; ** $p < 0.01$.

doi:10.1371/journal.pgen.1006257.g005

mediated phenotype and *LmCHS/LmCHT*-associated molting. In this study, we found that miR-71 and miR-263 control chitin synthesis and degradation by targeting *LmCHS1* and *LmCHT10*, resulting in post-transcriptional regulation of the molting process in locusts (Fig 4I). This miRNA-mediated mechanism of chitin metabolism provides insight into the molecular basis of the molting process in locusts.

We found that miR-71 targets *LmCHS1* and miR-263 targets *LmCHT10* in the chitin metabolic pathway of locusts. Chitin synthases and chitinases are responsible for the synthesis or degradation of chitins, which represent two inverse processes [39]. Thus, we suspected that the two processes of chitin synthesis and degradation affect one another. Therefore, we tested whether these two miRNAs interacted with each other using agomir-71/263 treatment. However, this was not the case (S7 Fig). The expression levels of the two miRNAs were very similar to those of the respective controls. In addition, 20-hydroxyecdysone (20E), as a key steroid hormone, coordinates multiple developmental events involving insect molting and metamorphosis [40]. Although 20E treatment is considered to be correlated with *CHS* and *CHT* expression, its roles in the regulation of *CHS* and *CHT* remain a matter of controversy. *DmeCHS-1* and *DmeCHS-2* transcription is activated by 20E during *Drosophila* metamorphosis [41]. *MsCHS-1* gene is negatively controlled by 20E, reflecting a dual effect of 20E [42]. *LmCHT5* and *TmCHT5* gene expression can be induced by 20E during the molting process [43,44]. However, the pathway linking 20E to *CHS* or *CHT* is still largely unknown. miR-8-5p and miR-2a-3p act as molecular regulators that tune the chitin biosynthesis pathway in response to 20E [45]. Thereby, we wondered whether 20E might play a role in the regulation of the expression of miR-71 and miR-263, leading to the precise expression of *LmCHS1* and *LmCHT10*. The 20E treatment inhibited miR-263 expression and induced *LmCHT10* expression but did not affect the expression levels of miR-71 or *LmCHS1* (S1 Fig). Thus, the 20E-miR-263-*CHT10* axis may switch the degradation of chitin on and off, whereas miR-71-*CHS1* axis-mediated chitin synthesis is regulated by mechanisms other than 20E.

The regulatory function of miR-263 is conserved across a broad range of insect species. We performed miRNA target prediction in other insect species for which *CHT10/CHS1* UTR sequences were available in the NCBI GenBank. The prediction results revealed that miRNA-263 binding sites of *CHT10* are present in several holometabolous insects including flies, beetles and mosquitoes, whereas no binding sites for miR-71 were identified in other insect species (S3 Table). This finding suggests that these regulatory interactions have been evolutionarily conserved, indicating that there is selective pressure to maintain the regulatory interactions of miR-263 and *CHT10* across species. Previous studies have shown that the expression patterns of *CHT10* during developmental instar stages are quite similar among insects [15,46], suggesting a common regulatory mechanism of the miRNA-263-dependent molting process. Our study provided experimental evidence that this regulatory mechanism is also present in hemimetabolous locusts. Members of Orthoptera occupy a more basal position in the insect lineage relative to holometabolous insects [47,48]. Thus, we presume that the regulatory roles of miR-263 in the molting process represent an ancestral function in insects that perhaps originated with the emergence of the common ancestor of hemimetabolous and holometabolous insects.

miRNAs are precise regulators that are able to sharpen developmental transcription by increasing and reducing target expression to meet developmental demands [49]. Due to the reduced expression of *CHS1* and *CHT10*, the agomir treatments of miRNA-71 and miR-263 can cause severe deficiency of new cuticle synthesis and failed degradation of the old cuticle, respectively. Unexpectedly, the antagomir treatments of miR-71/miR-263 and the resulting up-regulation of their target genes resulted in a similar aberrant phenotype, indicating that the elevated ectopic expression of *CHS1/CHT10* is also detrimental to the molting process. These results indicate that extremely high or low expression of *CHS1* and *CHT10* during a critical

period of the molting process can result in the development of an aberrant molting phenotype. miRNAs could tune the transcriptional activities of target genes to physiologically relevant levels [50]. miR-71/miR-263 could directly interact with *CHS1*/*CHT10* and play a role in ensuring an accurate level of their expression. The precise interactions of miR-71/miR-263 and *CHS1*/*CHT10* regulate the molting process in a spatio-temporal manner. Taken together, our results show that the transcriptional activities of *CHS1* and *CHT10* are tuned to a precise level at which they can execute proper function, emphasizing the important roles of miRNA-mediated precise regulation in the molting process.

The conclusion that emerges is that two miRNAs control the molting process by precisely regulating chitin metabolism. miR-71 and miR-263 suppress *CHS1* and *CHT10* transcript levels, thus preventing the progression of the molt to the next stage. This miRNA-mediated post-transcriptional regulation of chitin metabolism is particularly significant for understanding the molting process of locusts and potentially provides new targets for controlling locust plagues worldwide.

Materials and Methods

Insects

Locusts were obtained from the same locust colonies, which were maintained at the Institute of Zoology, Chinese Academy of Sciences, China. Nymphs were reared under a 14:10 light/dark photo regime at $30 \pm 2^\circ\text{C}$ and were fed fresh wheat seedlings and bran.

Small RNA transcriptome sequencing

Total RNA was extracted using TRIzol (Invitrogen) and treated with DNase I following the manufacturer's instructions. The RNA concentration and purity were assessed in an Agilent 2100 Bioanalyzer (Agilent) to verify RNA integrity. Small RNA libraries were constructed using a TruSeq small RNA sample preparation kit (Illumina). Briefly, the 3' and 5' RNA adapters were ligated to the corresponding ends of small RNAs. Following adapter ligation, the ligated RNA fragments were reverse transcribed using M-MLV reverse transcriptase (Invitrogen). The resulting cDNA products were PCR amplified with two primers that were complementary to the ends of the adapter sequences. The PCR amplicons were separated by size in 6% Novex polyacrylamide gel for miRNA enrichment and sequenced on an Illumina Genome Analyzer IIx sequencing system. Using the Cutadapt software, we trimmed the low-quality reads and the reads showing sequence similarity to adaptor sequences at the start or end terminals. The quantifier module in the miRDeep (version 2.0.0.5) software package was used to measure expression levels based on read counts.

In vitro luciferase validation

The ~300-bp sequences of the CDS and the 3' UTR surrounding the predicted miR-71 and miR-263 target sites in *CHS1* and *CHT10*, respectively, were separately cloned into the psi-CHECK-2 vector (Promega) using the XhoI and NotI sites. To generate the mutation version, the 8 nt of binding sites were mutated (GTTTTTCA for *CHS1*; GTGCTATT for *CHT10*), which include the region complementary to the miR-71 and miR-263 seed. S2 cells were co-transfected with 800 ng of the luciferase reporter vector or the empty vector and agomir-71 (-263) at a 1:4 ratio using the Lipofectamine™ 2000 reagent (Invitrogen) according to the manufacturer's instructions. The activities of the firefly and *Renilla* luciferases were measured 48 h after transfection with the Dual-Glo Luciferase Assay System (Promega) using a luminometer

(Promega). Results are expressed as the ratio Renilla/firefly luciferase activity (mean \pm SEM) based on six independent replicates.

miRNA agomir and antagomir treatment *in vivo*

The miRNA agomir or antagomir, each of which is a stable miRNA mimic or inhibitor, was used to validate the function of the miRNA *in vivo*. Briefly, 210 pmol of agomir-71 (-263) or antagomir-71 (-263) (500 μ M; RiboBio) was injected into the thoracic hemocoels of second-stadium nymphs two times at 48 h intervals. The agomir or antagomir negative controls (500 μ M) were also injected into the locust thoracic hemocoels (RiboBio). All injections were administered using a nanoliter injector (World Precision Instruments) with a glass micropipette tip. Treated nymphs were subjected to phenotypic observation of molting process. Their integuments were harvested, snap-frozen, and stored at -80°C .

Assays of quantitative PCR for mRNA and miRNA

Total RNA enriched for small RNAs was isolated from integuments using the mirVana miRNA isolation kit (Ambion). Moloney murine leukemia virus (M-MLV) reverse transcriptase (Promega) and a miRNA first-strand cDNA synthesis kit (Ambion) were used to prepare the Oligo (dT)-primed cDNA and stem-loop cDNA, respectively. The miRNAs and mRNAs were subjected to qPCR using the SYBR Green miRNA expression and gene expression assays, respectively, according to the manufacturer's instructions (Tiangen); qPCR was performed on a LightCycler[®] 480 instrument (Roche). The PCR data were analyzed using the $2^{-\Delta\Delta C_t}$ method of relative quantification. As endogenous controls, U6 snRNA and ribosomal protein *RP49* were used to quantify the miRNA and mRNA expression levels, respectively. Dissociation curves were determined for each miRNA and mRNA to confirm unique amplification. The qPCR primers are listed in [S4 Table](#). All the qRT-PCR reactions were performed in six biological replicates. Four integuments were involved in one biological replicate.

Co-localization of miRNA and its targets by fluorescence *in situ* hybridization

A combined two-color fluorescence *in situ* analysis of miRNA-71 (-263) and its targets was performed on the integuments of second-instar nymphs by co-labeling of the miRNA and its target according to the method described by Nuovo et al. [51]. An antisense locked nucleic acid (LNA) detection probe for miR-71, miR-263 or a scrambled control (Exiqon) was labeled with double digoxigenin. Biotin-labeled antisense and sense probes of *CHS1* and *CHT10* were generated from linearized recombinant pGEM-T Easy plasmids using the T7/SP6 RNA transcription system (Roche, Basel, Switzerland) following the recommended protocols. Based on the time-point at which higher expression activities were observed for miR-71 and miR-263, we selected the nymphs on the third day of the 2nd instar stage for miR-71 and *CHS1* co-localization detection and the nymphs on the fifth day of the 2nd instar stage for miR-263 and *CHT10* co-localization detection, respectively. The integuments were fixed in 4% paraformaldehyde overnight. The paraffin-embedded integument tissue slides (5 μ m thick) were deparaffinized in xylene, rehydrated with an ethanol gradient, digested with 20 μ g/mL proteinase K (Roche) at 37°C for 15 min, and incubated with the LNA miRNA probes and its target RNA probe at 60°C for 5 min. The slides were then hybridized for 7–15 h at 37°C and washed in $0.2\times$ SSC and 2% BSA at 4°C for 5 min. The slides were incubated in anti-digoxigenin-alkaline phosphatase conjugate (1:150 dilution) for 30 min at 37°C , followed by incubation with the HNPP substrate. For biotin-labeled probes, a TSA kit (Perkin Elmer, MA, USA) including a streptavidin horse radish peroxidase-conjugate and fluorescein tyramide substrate was used. The signals of the miRNA

and its target were detected using an LSM 710 confocal fluorescence microscope (Zeiss). The primers for probe synthesis of *CHS1* and *CHT10* are listed in [S4 Table](#).

RNA immunoprecipitation (RIP) assays

The RIP assay was performed using a Magna RIP Quad kit (Millipore) according to the manufacturer's instructions, with slight modifications. The 2-day-old second instar nymphs were microinjected with agomir-71 or agomir-263. A scrambled miRNA agomir was used as a negative control. Treated nymphs were subjected to RIP analysis 48 h later. Eight integuments of abdomen were collected and homogenized in ice-cold RIP lysis buffer. The homogenates were stored at -80°C overnight. A total of 5 μg of Ago-1 antibody (Abmart) or normal mouse IgG (Millipore), which was used as a negative control, was pre-incubated with magnetic beads. The frozen homogenates in the RIP lysates were thawed and centrifuged, and the supernatants were incubated with the magnetic bead-antibody complex at 4°C overnight. The immunoprecipitated RNAs were reverse-transcribed into cDNA using random hexamers. qPCR was performed to quantify *LmCHS1* and *LmCHT10*. The supernatants of the RIP lysates (input) and the IgG controls were assayed to normalize the relative expression levels of the target genes.

Quantification of chitin in the integument tissue extracts

The chitin content of the locust integuments was quantified after miR-71 or miR-263 administration. The integument tissues of the locust nymphs were immediately dissected and stored in liquid nitrogen. Three integuments of locust abdomens were homogenized in liquid nitrogen and transferred to 3% SDS. The homogenates were incubated at 100°C for 15 min; then, 120% KOH was added. The pellets were re-suspended and incubated at 130°C for 1 h. After cooling, 0.8 ml ice-cold 75% ethanol was added, and the samples were shaken until the KOH and ethanol formed a single phase. The homogenates were then centrifuged at 4°C , and the supernatants were discarded. The pellets were washed with 40% cold ethanol containing insoluble chitosan. Approximately 50 μl of 10% NaNO_2 and 50 μl of 10% KHSO_4 were added to each sample, and the samples were centrifuged at 4°C . The supernatants were combined with 20 μl of 12.5% $\text{NH}_4\text{SO}_3\text{NH}_2$ and 20 μl freshly prepared 0.5% (wt/vol) 3-methyl-2-benzothiazolone hydrazone hydrochloride hydrate solution, and the reaction was heated to 99.9°C for 3 min. After cooling, 20 μl of 0.83% $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ solution was added to the reaction. Measurements of the reaction mixture were performed using a microplate reader at 650 nm using glucosamine as a standard.

RNA interference

To knock down *CHS1* and *CHT10*, double-stranded RNA (dsRNA) was synthesized using T7 RiboMAXTM Express RNAi System (Promega, USA) following the manufacturer's instructions. Each insect was injected with 3 μg of dsRNAs at day 3 of the second instar nymphs. Control nymphs were injected with equivalent volumes of dsGFP alone. Total 25 nymphs were injected with dsRNA for each gene. Nymphs were observed carefully after injection. The nymphs that typically showed abnormal ecdysis were used for subsequent extraction. Six abnormal nymphs and six control nymphs were used for hematoxylin and eosin staining and chitin staining.

Statistical analysis

The SPSS 17.0 software (SPSS Inc.) was used for statistical analysis. The differences between treatments were compared using either Student's *t*-test or one-way analysis of variance

(ANOVA) followed by Tukey's test for multiple comparisons. The Mann-Whitney U test was used to analyze the behavioral data due to its non-normal distribution characteristics. $p < 0.05$ was considered statistically significant. All results are expressed as means \pm SEM.

Supporting Information

S1 Fig. Target prediction of miR-71 and miR-263 in the *CHS1* and *CHT10* genes of *Locusta migratoria* by RNAhybrid program.

(TIF)

S2 Fig. Expression of miRNAs containing potential binding sites with *CHS1* and *CHT10* by prediction using miRanda bioinformatics software were determined in the integuments of locusts by qRT-PCR. The data are presented as means \pm SEM ($n = 6$).

(TIF)

S3 Fig. Validation of miR-252 site not in *CHS1* mRNA of the locust. Luciferase reporter assays were analyzed in S2 cells co-transfected with miR-252 agomir and psi-CHECK2 vectors containing the target gene sequence of *CHS1* ($n = 6$). The data for the luciferase activities are presented as means \pm SEM ($n = 6$).

(TIF)

S4 Fig. Validation of miR-263 site in *CHT10* mRNA of the locust. Luciferase reporter assays were analyzed in S2 cells co-transfected with miR-263 antagomir and psi-CHECK2 vectors containing target gene sequences of *CHT10* ($n = 6$). The data for the luciferase activities are presented as means \pm SEM ($n = 6$).

(TIF)

S5 Fig. Effect of miR-71/miR-263 overexpression or silencing on the expression of miR-71/miR-263. (A, B) The expression levels of miR-71 and miR-263 were determined 48 h after treatment with 210 pmol agomir-71/ agomir-263 (A) or antagomir-71/antagomir-263 (B) using qPCR respectively. (C) The expression levels of let-7 were determined 48 h after treatment with 210 pmol agomir-71/agomir-263 or antagomir-71/antagomir-263 using qPCR, respectively.

(TIF)

S6 Fig. Effect of miR-71/miR-263 overexpression on the expression of other members of their target gene family. (A) *CHS2* expression was quantified using qRT-PCR 48 h after treatment of locusts with 210 pmol agomir-71. (B) *CHT5* expression was quantified using qRT-PCR 48 h after treatment of locusts with 210 pmol agomir-263. The data are presented as means \pm SEM ($n = 6$). * $p < 0.05$; ** $p < 0.01$.

(TIF)

S7 Fig. Effect of miR-71/miR-263 overexpression or silencing on the expression of miR-263/miR-71. (A) The expression levels of miR-263 were quantified using qPCR 48 h after the locusts were treated with 210 pmol of agomir- and antagomir-71. (B) The expression levels of miR-71 were quantified using qPCR 48 h after the locusts were treated with 210 pmol of agomir- and antagomir-263. All data are presented as means \pm SEM ($n = 6$).

(TIF)

S8 Fig. Effect of 20E on the expression levels of miR-71, miR-263 and their targets genes. (A) miR-71 and miR-263 expression was quantified using qRT-PCR 2 h and 6 h after the locusts were treated with 20E. (B) The expression levels of *CHS1* and *CHT10* were quantified using qRT-PCR 2 h and 6 h after the locusts were treated with 20E. The data are presented as

means \pm SEM ($n = 6$). $**p < 0.01$.
(TIF)

S1 Table. miRNAs identified in the integument of *Locusta migratoria* that contained the potential binding site with *CHS1* as predicted by the miRanda software.
(XLSX)

S2 Table. miRNAs identified in the integument of *Locusta migratoria* that contained the potential binding site with *CHT10* as predicted by the miRanda software.
(XLSX)

S3 Table. miRNAs contained the potential binding site with *CHT10* as predicted by the miRanda software in other species.
(XLSX)

S4 Table. The primary primers used in the study.
(XLS)

Author Contributions

Conceptualization: MY YW FJ LK.

Formal analysis: MY YW FJ TS HW QL JieZ.

Methodology: MY YW FJ TS HW QL.

Resources: MY YW FJ TS JieZ LK.

Supervision: LK JiaZ.

Writing - original draft: MY YW FJ LK.

Writing - review & editing: MY YW FJ JiaZ LK.

References

1. Niwa YS, Niwa R (2016) Transcriptional regulation of insect steroid hormone biosynthesis and its role in controlling timing of molting and metamorphosis. *Development Growth & Differentiation* 58: 94–105.
2. Riddiford LM, Hiruma K, Zhou XF, Nelson CA (2003) Insights into the molecular basis of the hormonal control of molting and metamorphosis from *Manduca sexta* and *Drosophila melanogaster*. *Insect Biochemistry and Molecular Biology* 33: 1327–1338. PMID: [14599504](#)
3. Merzendorfer H (2006) Insect chitin synthases: a review. *J Comp Physiol B* 176: 1–15. PMID: [16075270](#)
4. Cohen E (2001) Chitin synthesis and inhibition: a revisit. *Pest Management Science* 57: 946–950. PMID: [11695188](#)
5. Kang L, Chen XY, Zhou Y, Liu BW, Zheng W, et al. (2004) The analysis of large-scale gene expression correlated to the phase changes of the migratory locust. *Proceedings of the National Academy of Sciences of the United States of America* 101: 17611–17615. PMID: [15591108](#)
6. Zhang JZ, Liu XJ, Zhang JQ, Li DQ, Sun Y, et al. (2010) Silencing of two alternative splicing-derived mRNA variants of chitin synthase 1 gene by RNAi is lethal to the oriental migratory locust, *Locusta migratoria manilensis* (Meyen). *Insect Biochemistry and Molecular Biology* 40: 824–833. doi: [10.1016/j.ibmb.2010.08.001](#) PMID: [20713155](#)
7. Liu X, Zhang H, Li S, Zhu KY, Ma E, et al. (2012) Characterization of a midgut-specific chitin synthase gene (LmCHS2) responsible for biosynthesis of chitin of peritrophic matrix in *Locusta migratoria*. *Insect Biochem Mol Biol* 42: 902–910. doi: [10.1016/j.ibmb.2012.09.002](#) PMID: [23006725](#)
8. Zhang J, Liu X, Li D, Sun Y, Guo Y, et al. (2010) Silencing of two alternative splicing-derived mRNA variants of chitin synthase 1 gene by RNAi is lethal to the oriental migratory locust, *Locusta migratoria manilensis* (Meyen). *Insect Biochem Mol Biol* 40: 824–833. doi: [10.1016/j.ibmb.2010.08.001](#) PMID: [20713155](#)

9. Rong S, Li DQ, Zhang XY, Li S, Zhu KY, et al. (2013) RNA interference to reveal roles of beta-N-acetylglucosaminidase gene during molting process in *Locusta migratoria*. *Insect Sci* 20: 109–119. doi: [10.1111/j.1744-7917.2012.01573.x](https://doi.org/10.1111/j.1744-7917.2012.01573.x) PMID: [23955831](https://pubmed.ncbi.nlm.nih.gov/23955831/)
10. Ampasala DR, Zheng SC, Zhang DY, Ladd T, Doucet D, et al. (2011) An epidermis-specific chitin synthase cDNA in *Choristoneura fumiferana*: cloning, characterization, developmental and hormonal-regulated expression. *Archives of Insect Biochemistry and Physiology* 76: 83–96. doi: [10.1002/arch.20404](https://doi.org/10.1002/arch.20404) PMID: [21181720](https://pubmed.ncbi.nlm.nih.gov/21181720/)
11. Takahashi M, Kiuchi M, Kamimura M (2002) A new chitinase-related gene, BmChiR1, is induced in the *Bombyx mori* anterior silk gland at molt and metamorphosis by ecdysteroid. *Insect Biochemistry and Molecular Biology* 32: 147–151. PMID: [11755056](https://pubmed.ncbi.nlm.nih.gov/11755056/)
12. Zhuo WW, Fang Y, Kong LF, Li X, Sima YH, et al. (2014) Chitin synthase A: a novel epidermal development regulation gene in the larvae of *Bombyx mori*. *Molecular Biology Reports* 41: 4177–4186. doi: [10.1007/s11033-014-3288-1](https://doi.org/10.1007/s11033-014-3288-1) PMID: [24577751](https://pubmed.ncbi.nlm.nih.gov/24577751/)
13. Liu XJ, Zhang HH, Li S, Zhu KY, Ma EB, et al. (2012) Characterization of a midgut-specific chitin synthase gene (LmCHS2) responsible for biosynthesis of chitin of peritrophic matrix in *Locusta migratoria*. *Insect Biochemistry and Molecular Biology* 42: 902–910. doi: [10.1016/j.ibmb.2012.09.002](https://doi.org/10.1016/j.ibmb.2012.09.002) PMID: [23006725](https://pubmed.ncbi.nlm.nih.gov/23006725/)
14. Yang WJ, Xu KK, Zhang RY, Dou W, Wang JJ (2013) Transcriptional Regulation of a Chitinase Gene by 20-Hydroxyecdysone and Starvation in the Oriental Fruit Fly, *Bactrocera dorsalis*. *International Journal of Molecular Sciences* 14: 20048–20063. doi: [10.3390/ijms141020048](https://doi.org/10.3390/ijms141020048) PMID: [24113584](https://pubmed.ncbi.nlm.nih.gov/24113584/)
15. Zhu QS, Arakane Y, Beeman RW, Kramer KJ, Muthukrishnan S (2008) Functional specialization among insect chitinase family genes revealed by RNA interference. *Proceedings of the National Academy of Sciences of the United States of America* 105: 6650–6655. doi: [10.1073/pnas.0800739105](https://doi.org/10.1073/pnas.0800739105) PMID: [18436642](https://pubmed.ncbi.nlm.nih.gov/18436642/)
16. Xi Y, Pan PL, Ye YX, Yu B, Xu HJ, et al. (2015) Chitinase-like gene family in the brown planthopper, *Nilaparvata lugens*. *Insect Molecular Biology* 24: 29–40. doi: [10.1111/imb.12133](https://doi.org/10.1111/imb.12133) PMID: [25224926](https://pubmed.ncbi.nlm.nih.gov/25224926/)
17. Krol J, Loedige I, Filipowicz W (2010) The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* 11: 597–610. doi: [10.1038/nrg2843](https://doi.org/10.1038/nrg2843) PMID: [20661255](https://pubmed.ncbi.nlm.nih.gov/20661255/)
18. Lai EC (2003) microRNAs: Runts of the genome assert themselves. *Current Biology* 13: R925–R936. PMID: [14654021](https://pubmed.ncbi.nlm.nih.gov/14654021/)
19. Flynt AS, Lai EC (2008) Biological principles of microRNA-mediated regulation: shared themes amid diversity. *Nature Reviews Genetics* 9: 831–842. doi: [10.1038/nrg2455](https://doi.org/10.1038/nrg2455) PMID: [18852696](https://pubmed.ncbi.nlm.nih.gov/18852696/)
20. Rigoutsos I (2009) New tricks for animal microRNAs: targeting of amino acid coding regions at conserved and nonconserved sites. *Cancer Res* 69: 3245–3248. doi: [10.1158/0008-5472.CAN-09-0352](https://doi.org/10.1158/0008-5472.CAN-09-0352) PMID: [19351814](https://pubmed.ncbi.nlm.nih.gov/19351814/)
21. Duursma AM, Kedde M, Schrier M, le Sage C, Agami R (2008) miR-148 targets human DNMT3b protein coding region. *RNA* 14: 872–877. doi: [10.1261/ma.972008](https://doi.org/10.1261/ma.972008) PMID: [18367714](https://pubmed.ncbi.nlm.nih.gov/18367714/)
22. Yang M, Wei Y, Jiang F, Wang Y, Guo X, et al. (2014) MicroRNA-133 inhibits behavioral aggregation by controlling dopamine synthesis in locusts. *PLoS Genet* 10: e1004206. doi: [10.1371/journal.pgen.1004206](https://doi.org/10.1371/journal.pgen.1004206) PMID: [24586212](https://pubmed.ncbi.nlm.nih.gov/24586212/)
23. Chen J, Liang Z, Liang Y, Pang R, Zhang W (2013) Conserved microRNAs miR-8-5p and miR-2a-3p modulate chitin biosynthesis in response to 20-hydroxyecdysone signaling in the brown planthopper, *Nilaparvata lugens*. *Insect Biochem Mol Biol* 43: 839–848. doi: [10.1016/j.ibmb.2013.06.002](https://doi.org/10.1016/j.ibmb.2013.06.002) PMID: [23796434](https://pubmed.ncbi.nlm.nih.gov/23796434/)
24. Gomez-Orte E, Belles X (2009) MicroRNA-dependent metamorphosis in hemimetabolan insects. *Proceedings of the National Academy of Sciences of the United States of America* 106: 21678–21682. doi: [10.1073/pnas.0907391106](https://doi.org/10.1073/pnas.0907391106) PMID: [19966227](https://pubmed.ncbi.nlm.nih.gov/19966227/)
25. Lozano J, Montanez R, Belles X (2015) MiR-2 family regulates insect metamorphosis by controlling the juvenile hormone signaling pathway. *Proceedings of the National Academy of Sciences of the United States of America* 112: 3740–3745. doi: [10.1073/pnas.1418522112](https://doi.org/10.1073/pnas.1418522112) PMID: [25775510](https://pubmed.ncbi.nlm.nih.gov/25775510/)
26. Caygill EE, Johnston LA (2008) Temporal regulation of metamorphic processes in *Drosophila* by the let-7 and miR-125 heterochronic microRNAs. *Current Biology* 18: 943–950. doi: [10.1016/j.cub.2008.06.020](https://doi.org/10.1016/j.cub.2008.06.020) PMID: [18571409](https://pubmed.ncbi.nlm.nih.gov/18571409/)
27. Wang YL, Yang ML, Jiang F, Zhang JZ, Kang L (2013) MicroRNA-dependent development revealed by RNA interference-mediated gene silencing of LmDicer1 in the migratory locust. *Insect Sci* 20: 53–60. doi: [10.1111/j.1744-7917.2012.01542.x](https://doi.org/10.1111/j.1744-7917.2012.01542.x) PMID: [23955825](https://pubmed.ncbi.nlm.nih.gov/23955825/)
28. Mansur JE, Alvarenga ESL, Figueira-Mansur J, Franco TA, Ramos IB, et al. (2014) Effects of chitin synthase double-stranded RNA on molting and oogenesis in the Chagas disease vector *Rhodnius*

- prolixus. *Insect Biochemistry and Molecular Biology* 51: 110–121. doi: [10.1016/j.ibmb.2013.12.006](https://doi.org/10.1016/j.ibmb.2013.12.006) PMID: [24398146](https://pubmed.ncbi.nlm.nih.gov/24398146/)
29. Wang Y, Fan HW, Huang HJ, Xue J, Wu WJ, et al. (2012) Chitin synthase 1 gene and its two alternative splicing variants from two sap-sucking insects, *Nilaparvata lugens* and *Laodelphax striatellus* (Hemiptera: Delphacidae). *Insect Biochemistry and Molecular Biology* 42: 637–646. doi: [10.1016/j.ibmb.2012.04.009](https://doi.org/10.1016/j.ibmb.2012.04.009) PMID: [22634163](https://pubmed.ncbi.nlm.nih.gov/22634163/)
 30. Wang Y, Jiang F, Wang H, Song T, Wei Y, et al. (2015) Evidence for the expression of abundant microRNAs in the locust genome. *Sci Rep* 5: 13608. doi: [10.1038/srep13608](https://doi.org/10.1038/srep13608) PMID: [26329925](https://pubmed.ncbi.nlm.nih.gov/26329925/)
 31. Soares MPM, Silva-Torres FA, Elias-Neto M, Nunes FMF, Simoes ZLP, et al. (2011) Ecdysteroid-Dependent Expression of the Tweedle and Peroxidase Genes during Adult Cuticle Formation in the Honey Bee, *Apis mellifera*. *PLoS One* 6.
 32. Hewes RS, Schaefer AM, Taghert PH (2000) The cryptocephal gene (ATF4) encodes multiple basic-leucine zipper proteins controlling molting and metamorphosis in *Drosophila*. *Genetics* 155: 1711–1723. PMID: [10924469](https://pubmed.ncbi.nlm.nih.gov/10924469/)
 33. Petkau G, Wingen C, Jussen LCA, Radtke T, Behr M (2012) Obstructor-A Is Required for Epithelial Extracellular Matrix Dynamics, Exoskeleton Function, and Tubulogenesis. *Journal of Biological Chemistry* 287: 21396–21405. doi: [10.1074/jbc.M112.359984](https://doi.org/10.1074/jbc.M112.359984) PMID: [22544743](https://pubmed.ncbi.nlm.nih.gov/22544743/)
 34. Chaudhari SS, Arakane Y, Specht CA, Moussian B, Kramer KJ, et al. (2013) Retroactive Maintains Cuticle Integrity by Promoting the Trafficking of Knickkopf into the Procuticle of *Tribolium castaneum*. *Plos Genetics* 9.
 35. Garen A, Kauvar L, Lepesant JA (1977) Roles of Ecdysone in *Drosophila* Development. *Proceedings of the National Academy of Sciences of the United States of America* 74: 5099–5103. PMID: [16592466](https://pubmed.ncbi.nlm.nih.gov/16592466/)
 36. Marco A, Hui JHL, Ronshaugen M, Griffiths-Jones S (2010) Functional Shifts in Insect microRNA Evolution. *Genome Biology and Evolution* 2: 686–696. doi: [10.1093/gbe/evq053](https://doi.org/10.1093/gbe/evq053) PMID: [20817720](https://pubmed.ncbi.nlm.nih.gov/20817720/)
 37. Cai MJ, Zhao WL, Jing YP, Song Q, Zhang XQ, et al. (2016) 20-hydroxyecdysone activates Forkhead box O to promote proteolysis during *Helicoverpa armigera* molting. *Development* 143: 1005–1015. doi: [10.1242/dev.128694](https://doi.org/10.1242/dev.128694) PMID: [26893349](https://pubmed.ncbi.nlm.nih.gov/26893349/)
 38. Wang YL, Yang ML, Jiang F, Zhang JZ, Kang L (2013) MicroRNA-dependent development revealed by RNA interference-mediated gene silencing of *LmDicer1* in the migratory locust. *Insect Science* 20: 53–60. doi: [10.1111/j.1744-7917.2012.01542.x](https://doi.org/10.1111/j.1744-7917.2012.01542.x) PMID: [23955825](https://pubmed.ncbi.nlm.nih.gov/23955825/)
 39. Merzendorfer H, Zimoch L (2003) Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *Journal of Experimental Biology* 206: 4393–4412. PMID: [14610026](https://pubmed.ncbi.nlm.nih.gov/14610026/)
 40. Dhadialla TS, Carlson GR, Le DP (1998) New insecticides with ecdysteroidal and juvenile hormone activity. *Annual Review of Entomology* 43: 545–569. PMID: [9444757](https://pubmed.ncbi.nlm.nih.gov/9444757/)
 41. Gagou ME, Kapsetaki M, Turberg A, Kafetzopoulos D (2002) Stage-specific expression of the chitin synthase *DmeChSA* and *DmeChSB* genes during the onset of *Drosophila* metamorphosis. *Insect Biochemistry and Molecular Biology* 32: 141–146. PMID: [11755055](https://pubmed.ncbi.nlm.nih.gov/11755055/)
 42. Zhu YC, Specht CA, Dittmer NT, Muthukrishnan S, Kanost MR, et al. (2002) Sequence of a cDNA and expression of the gene encoding a putative epidermal chitin synthase of *Manduca sexta*. *Insect Biochemistry and Molecular Biology* 32: 1497–1506. PMID: [12530217](https://pubmed.ncbi.nlm.nih.gov/12530217/)
 43. Royer W, Fraichard S, Bouhin H (2002) A novel putative insect chitinase with multiple catalytic domains: hormonal regulation during metamorphosis. *Biochemical Journal* 366: 921–928. PMID: [12059786](https://pubmed.ncbi.nlm.nih.gov/12059786/)
 44. Li D, Zhang JQ, Wang Y, Liu XJ, Ma EB, et al. (2015) Two chitinase 5 genes from *Locusta migratoria*: Molecular characteristics and functional differentiation. *Insect Biochemistry and Molecular Biology* 58: 46–54. doi: [10.1016/j.ibmb.2015.01.004](https://doi.org/10.1016/j.ibmb.2015.01.004) PMID: [25623241](https://pubmed.ncbi.nlm.nih.gov/25623241/)
 45. Chen J, Liang ZK, Liang YK, Pang R, Zhang WQ (2013) Conserved microRNAs miR-8-5p and miR-2a-3p modulate chitin biosynthesis in response to 20-hydroxyecdysone signaling in the brown planthopper, *Nilaparvata lugens*. *Insect Biochemistry and Molecular Biology* 43: 839–848. doi: [10.1016/j.ibmb.2013.06.002](https://doi.org/10.1016/j.ibmb.2013.06.002) PMID: [23796434](https://pubmed.ncbi.nlm.nih.gov/23796434/)
 46. Zhang JZ, Zhang X, Arakane Y, Muthukrishnan S, Kramer KJ, et al. (2011) Comparative Genomic Analysis of Chitinase and Chitinase-Like Genes in the African Malaria Mosquito (*Anopheles gambiae*). *PLoS One* 6.
 47. Misof B, Liu SL, Meusemann K, Peters RS, Donath A, et al. (2014) Phylogenomics resolves the timing and pattern of insect evolution. *Science* 346: 763–767. doi: [10.1126/science.1257570](https://doi.org/10.1126/science.1257570) PMID: [25378627](https://pubmed.ncbi.nlm.nih.gov/25378627/)
 48. Giribet G, Edgecombe GD, Wheeler WC (2001) Arthropod phylogeny based on eight molecular loci and morphology. *Nature* 413: 157–161. PMID: [11557979](https://pubmed.ncbi.nlm.nih.gov/11557979/)

49. Ebert MS, Sharp PA (2012) Roles for microRNAs in conferring robustness to biological processes. *Cell* 149: 515–524. doi: [10.1016/j.cell.2012.04.005](https://doi.org/10.1016/j.cell.2012.04.005) PMID: [22541426](https://pubmed.ncbi.nlm.nih.gov/22541426/)
50. Hobert O (2007) miRNAs play a tune. *Cell* 131: 22–24. PMID: [17923083](https://pubmed.ncbi.nlm.nih.gov/17923083/)
51. Nuovo GJ, Elton TS, Nana-Sinkam P, Volinia S, Croce CM, et al. (2009) A methodology for the combined in situ analyses of the precursor and mature forms of microRNAs and correlation with their putative targets. *Nat Protoc* 4: 107–115. doi: [10.1038/nprot.2008.215](https://doi.org/10.1038/nprot.2008.215) PMID: [19131963](https://pubmed.ncbi.nlm.nih.gov/19131963/)