

miRNA malfunction causes spinal motor neuron disease

Sharon Haramati^a, Elik Chapnik^{b,1}, Yehezkel Sztainberg^{a,c,1}, Raya Eilam^d, Raaya Zwang^a, Noga Gershoni^b, Edwina McGlinn^e, Patrick W. Heiser^f, Anne-Marie Wills^g, Itzhak Wirguin^h, Lee L. Rubin^f, Hidemi Misawaⁱ, Clifford J. Tabin^{e,2}, Robert Brown, Jr.^j, Alon Chen^{a,2}, and Eran Hornstein^{b,2}

Departments of ^aNeurobiology and ^bMolecular Genetics and ^cVeterinary Resources, Weizmann Institute of Science, Rehovot 76100, Israel; ^dThe Leslie and Susan Gonda (Goldschmied) Multidisciplinary Brain Research Center, Bar-Ilan University, Ramat-Gan 52900, Israel; ^eDepartment of Genetics, Harvard Medical School, Boston, MA 02115; ^fDepartment of Stem Cell and Regenerative Biology, Harvard Stem Cell Institute, Harvard University, Cambridge, MA 02138; ^gDepartment of Neurology, Massachusetts General Hospital, Boston, MA 02114; ^hDepartment of Neurology, Soroka Medical Center, Ben-Gurion University of the Negev, Beer-Sheva 84101, Israel; ⁱDepartment of Pharmacology, Keio University Faculty of Pharmacy, Tokyo 160-8582, Japan; and ^jDepartment of Neurology, University of Massachusetts School of Medicine, Worcester, MA 01655

Contributed by Clifford J. Tabin, May 5, 2010 (sent for review November 30, 2009)

Defective RNA metabolism is an emerging mechanism involved in ALS pathogenesis and possibly in other neurodegenerative disorders. Here, we show that microRNA (miRNA) activity is essential for long-term survival of postmitotic spinal motor neurons (SMNs) in vivo. Thus, mice that do not process miRNA in SMNs exhibit hallmarks of spinal muscular atrophy (SMA), including sclerosis of the spinal cord ventral horns, aberrant end plate architecture, and myofiber atrophy with signs of denervation. Furthermore, a neurofilament heavy subunit previously implicated in motor neuron degeneration is specifically up-regulated in miRNA-deficient SMNs. We demonstrate that the heavy neurofilament subunit is a target of miR-9, a miRNA that is specifically down-regulated in a genetic model of SMA. These data provide evidence for miRNA function in SMN diseases and emphasize the potential role of miR-9-based regulatory mechanisms in adult neurons and neurodegenerative states.

ALS | Dicer | microRNA | motor neuron | neurodegeneration

Regulation by micro-RNA (miRNA) appears to be the most abundant mode of posttranscriptional regulation (1). This is because hundreds of miRNA genes, each regulating a diverse set of downstream targets, take part in practically all cellular processes, whether in health or disease.

Genome-encoded miRNAs are transcribed as long RNA transcripts that fold back on themselves to form distinctive hairpin structures. The long miRNA precursor is first digested by the Drosha microprocessor complex (2–4) and then by Dicer1 (5). The mature miRNA is loaded onto the Argonaute silencing complex (6) that directs posttranscriptional repression through miRNA:mRNA pairing. The two main mechanisms for repression of gene expression by miRNA are miRNA-directed translational repression and mRNA destabilization [reviewed in (7, 8)].

Work over the past years has documented a crucial role for miRNA-dependent posttranscriptional gene regulation in the development and function of neurons [e.g., (9–14)]; recently reviewed in (15–17)]. For example, miR-9 is an ancient neuronal gene involved in flies in selection of neuronal precursors from the neuroepithelium (10). The miR-9 gene is conserved to vertebrates, wherein it specifies the midbrain-hindbrain boundary (18) and, with miR-124, plays a role in neuronal differentiation (19, 20).

Furthermore, alterations in the function of miRNA contribute to susceptibility to neuronal disease. Although this may be associated with loss of neurons (21–27), behavioral and neuroanatomical phenotypes in the absence of neurodegeneration were also reported (28). The expression of specific miRNA was also linked to neurodegeneration; for example, a significant decrease in miR-9 and miR-9* expression was noted in patients with Huntington's disease (20), miR-9 and miR-132 are downregulated in Alzheimer's disease brains (29) and loss of miR-133 expression were suggested to play a role in Parkinson's disease (23).

ALS is a neurodegenerative disease that specifically affects upper and lower motor neurons (MNs), leading to progressive paralysis and death. Recently discovered mutations in the genes encoding the RNA-binding proteins FUS/TLS [ALS6 locus (30, 31)] and TARDBP/TDP43 [ALS10 locus (32, 33)] suggest important roles for regulatory RNA in the pathogenesis of ALS (34). Intriguingly, these disease-related RNA-binding proteins were identified in neuronal RNA granules (35, 36) and with miRNA-associated complexes (3, 37). Similarly, juvenile forms of motor neuron diseases (MND) are related to posttranscriptional regulators of gene expression, namely, SETX [ALS4 locus; (38)], IGHMBP2 (39) and SMN1 (40, 41), the latter functionally engaged in miRNA-protein complexes (42, 43). Plausibly, a considerable portion of the MND spectrum may be directly related to RNA metabolism and posttranscriptional regulation of gene expression.

This emerging appreciation of RNA regulatory function in neurons encouraged us to hypothesize that miRNA may be involved in the pathogenesis of MNDs. In this work, we show that miRNA dysfunction causes spinal muscular atrophy (SMA). Furthermore, we show that the neurofilament heavy subunit (NEFH) previously implicated in MND is specifically up-regulated in Dicer1-deficient MNs.

We additionally relate the down-regulation of the miR-9 gene to changes in neurofilament stoichiometry in both the Dicer1 model and in a murine SMN1 model of SMA. These data provide direct evidence for miRNA malfunction in MNDs and promote further evaluation of miR-9 in neurodegeneration.

Results

Loss of miRNA Activity in the MNDicer^{mut} Causes Progressive Locomotor Dysfunction. Because miRNA makes up the largest group of regulatory RNA (1) and has previously been associated with neurodegenerative states (22, 23, 25, 28), we sought to evaluate its involvement in MN pathologies. To this end, we specifically ablated Dicer1 in postmitotic postnatal MNs, crossing a Dicer1 conditional allele (44) with a Cre-recombinase transgene driven by a cholinergic-specific promoter [vesicular acetyl-choline trans-

Author contributions: S.H., E.C., C.J.T., A.C., and E.H. designed research; S.H., E.C., H.S., R.E., R.Z., N.G., E.M., P.W.H., A.-M.W., I.W., L.L.R., H.M., R.B., and E.H. performed research; P.W.H., A.-M.W., L.L.R., and H.M. contributed new reagents/analytic tools; S.H., E.C., H.S., R.E.-A., A.C., and E.H. analyzed data; and S.H., A.C., and E.H. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

¹E.C. and H.S. contributed equally to this work.

²To whom correspondence may be addressed. E-mail: tabin@genetics.med.harvard.edu, alon.chen@weizmann.ac.il, or eran.hornstein@weizmann.ac.il.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1006151107/-DCSupplemental.

porter (VChT-Cre.fast (45–47)). This transgene is expressed in postmitotic somatic MNs as early as postnatal day 7 but is not expressed during development. Because Dicer1 activity is required for miRNA processing in vivo (1, 25, 28), VChT-Cre.fast;Dicer1^{flx/flx} animals (referred to below as “MNDicer^{mut}”) lose the ability to make functional miRNA in a subset of postmitotic SMNs, and therefore provide a compelling model for miRNA loss of function in SMNs (details of the mouse genetic system are shown in Fig. S1).

Whereas VChT-Cre.fast;Dicer1^{flx/+} heterozygous animals (“controls”) are apparently normal and their survival is comparable with that of WT mice, the median life of MNDicer^{mut} mice is 29 wk, with the last moribund animal euthanized at 50 wk of age. In addition, MNDicer^{mut} mice gained weight slower than control littermates (Fig. 1*A* and *B*). To understand the pathology of these mice better, we conducted a broad series of functional tests to evaluate their locomotor activity. From the age of 7 wk, the MNDicer^{mut} mice did not perform as well as controls on a “vertical pole test” (Fig. 1*C*). Furthermore, a video-monitored “open-field” assay revealed that MNDicer^{mut} mice progressively travel shorter distances and rear less than controls (Fig. 1*D* and *E*). In addition, a home-cage study across the circadian cycle indicated that the locomotor activity of MNDicer^{mut} mice gradually deteriorates compared with that of controls (Fig. 1*F* and *H*). We hypothesized that this apparent deterioration in activity and locomotor parameters is likely the consequence of muscular atrophy.

MNDicer^{mut} Mouse Exhibits Denervation Muscular Atrophy. To characterize the muscle phenotype directly, we performed a needle electromyography (EMG) study on the hind-limb interosseus and gastrocnemius muscles. EMG performed on anaesthetized animals

revealed frequent fibrillation potentials in the MNDicer^{mut} mouse relative to controls (Fig. 2*A* and *B*). These data are consistent with an ongoing denervation process, which probably underlies the progressive locomotive deterioration of the MNDicer^{mut} animals and also their observable tremor.

Histological examination of the MNDicer^{mut} tibialis anterior muscle by myosin ATPase reaction further supports neuropathy, because MNDicer^{mut} mice frequently exhibited fiber type grouping. This finding is characteristic of a denervation/reinnervation pathology (48) and was not seen in the controls (Fig. 2*C*). Further, muscle fibers with a large cross-sectional area are specifically lost in the MNDicer^{mut} mouse, and the total fiber diameter of the MNDicer^{mut} mouse is reduced relative to controls, although this failed to reach statistical significance (Fig. S2). Finally, we depicted angular myofibers on muscle histology, a pathognomonic sign of denervation-related muscular atrophy (Fig. 2*D*). Taken together, we conclude that MNDicer^{mut} animals suffer from denervation muscular atrophy, which suggests loss of spinal motor neurons (SMNs).

MNs Are Lost in the MNDicer^{mut} Mouse. On spinal cord histology, we studied by Nissl staining the numbers of large perikaryon numbers (>20 μ m in diameter) in the ventral horn of the lumbar (L4–L5) level of the spinal cord. We noted a significant decrease in large MN numbers in multiple sections throughout the lumbar regions of the MNDicer^{mut} mouse relative to controls (average of 7 and 10.1 MNs per section, respectively; depicted in 15 lumbar sections per animal; $n = 5$ and $n = 5$, respectively; Fig. 3*A*).

Negative immunoreactivity for both TUNEL and activated caspase-3 was consistent with a typical slow death profile of SMNs encountered in many MNDs.




Fig. 1. Inferior survival and motor activity of the MNDicer^{mut} mice. (A) Kaplan–Meier survival curve for controls and conditional Dicer1 KO mice (control, $n = 12$; MNDicer^{mut}, $n = 12$). The median survival of MNDicer^{mut} mice is 29 wk, and it is >60 wk for controls. (B) Weight gain of controls and MNDicer^{mut} mice. (C) Time to complete a turn in the pole task for controls and MNDicer^{mut} mice. (D and E) Open-field measures at 8, 16, and 31 wk of age. (D) Ratio of distance traveled in the open-field arena compared with the mean of controls (control, $n = 12$; MNDicer^{mut}, $n = 12$). (E) Ratio of rearing events performed in the open-field arena compared with the mean of controls (control, $n = 12$; MNDicer^{mut}, $n = 12$). Home-cage locomotion of MNDicer^{mut} and controls at 11 (F), 21 (G), and 50 (H) wk of age (control, $n = 12$; MNDicer^{mut}, $n = 12$). (Inset) Average of activity throughout the measured period. * $P < 0.05$; ** $P < 0.01$.




Fig. 2. MNDicer^{mut} mice exhibit muscular atrophy with signs of denervation. Hind-limb interosseous and gastrocnemius muscle bipolar EMG recording. (A) Representative EMG traces of control (Upper) and MNDicer^{mut} (Lower) mice under anesthesia. Frequent fibrillation potentials are annotated by red arrows. (B) EMG Pathology Index was evaluated for individual controls (O; $n = 5$) and MNDicer^{mut} mice (●; $n = 5$). This scale (range: 0–7) reflects the intensity and frequency of fibrillation potentials in coded mice, noting that the electromyographer was blinded as to the genotype of the mouse tested. (C) Basic ATPase staining of transverse section through control and MNDicer^{mut} tibialis anterior muscles. Fiber grouping events were observed only in mutant muscles. (D) H&E staining of transverse section through the tibialis anterior MNDicer^{mut} muscle. Angular fibers are marked by arrows. (Scale bar: 100 μm .) * $P < 0.05$; ** $P < 0.01$.

Reactive astrocytosis is often taken as an indication of neuronal toxicity or neuronal death (49); therefore, we immunohistochemically quantified GFAP expression levels by immunofluorescence and Western blotting. We detected enhanced GFAP immunoreactivity in sections of the lateroventral aspect of the lumbar spinal cord of the MNDicer^{mut} mice and substantiated this by Western blot analysis that revealed higher levels of GFAP expression in spinal cord extracts of the MNDicer^{mut} animals relative to controls (Fig. 3B). These data support reactive astrocytosis and MN loss.

We further evaluated the discrete population of proximal motor axons at the ventral root before they are joined by sensory axons. MNDicer^{mut} mice exhibit a significant decrease in MN axon numbers when compared with controls, whereas dorsal root sensory axons remain intact as expected (Fig. 3C).

Signs of Axonopathy in the MNDicer^{mut} Mouse. Dysfunction and/or degeneration of the neuromuscular junction (NMJ) accompanies or even precedes the loss of MN bodies in a few models of ALS (50–52). We went on to evaluate potential distal axonal defects in the MNDicer^{mut} mouse. Detailed evaluation of the pre- and postsynaptic compartments of 350 individual NMJs in the hind-limb tibialis anterior revealed that aberrant architecture was twice as frequent in the NMJs of MNDicer^{mut} mice relative to controls (Fig. 3D). This is intriguing, because miRNA is known to have distal perisynaptic functions (11, 53–55), suggesting that miRNA-related neuropathy may exist while the axons are still occupying the end plate.

MNDicer^{mut} Mouse Fails to Coordinate Neurofilament Subunit Stoichiometry. Dysregulation of the coordinated expression of the light neurofilament (NEFL), medium neurofilament (NEFM), and NEFH subunits causes axonal cytoskeletal defects (56, 57). For example, NEFL mutations cause type 2E Charcot–Marie–Tooth motor neuropathy (58). Furthermore, experimental perturbation of the fine neurofilament balance in mouse models results in phenotypes closely resembling human MN pathologies (59, 60) and has previously been suggested as a component of human ALS (61–63).

More specifically, posttranscriptional regulation of neurofilament gene expression plays a key role in neuronal well-being




Fig. 3. MNDicer^{mut} mice exhibit spinal cord ventral horn sclerosis and axonopathy. (A) (Left) Representative Nissl staining of lumbar (L4–L5) sections from a MNDicer^{mut} mouse and a control littermate. (Insets) Enlargements of a ventral horn area in each section. The dashed line represents the border under which large-diameter cells (>20 μm) were counted. (Right) Average number of MNs counted per ventral horn section in lumbar spinal cord of 4-mo-old MNDicer^{mut} mice and controls (average of 7 and 10.1 MNs per section, respectively; 15 lumbar sections per animal; $n = 5$ and $n = 5$, respectively). (B) (Left) Representative lumbar section from 4-mo-old MNDicer^{mut} mice and controls immunostained for GFAP. (Right) Quantification of the GFAP immunofluorescence (IF) signal (arbitrary units, 3 lumbar sections per animal; $n = 5$ and $n = 5$, respectively) and quantification of GFAP by Western blot analysis of lower spinal cord extracts from MNDicer^{mut} mice (“mut”) and controls, normalized to the expression of β -tubulin (arbitrary units; $n = 3$ and $n = 4$, respectively). A representative capture from the Western blot analysis is provided for two mutants and two controls. (C) (Left) Representative dorsal (sensory, Left) and ventral (motor, Right) roots used for axon number measurements, stained with anti-NEFM antibody. (Right) Average axon number in dorsal and ventral roots of MNDicer^{mut} mice and controls ($n = 2$ and $n = 2$, respectively). (D) (Left) Representative hind-limb tibialis anterior NMJ, demonstrating complete overlap (Upper) or partial overlap (Lower) between the postsynapse (red, rhodamine-labeled bungarotoxin) and presynapse (green, mixture of anti-neurofilament and synaptophysin antibodies; yellow, merged channels) components. (Right) Percentage of pathological end plates in MNDicer^{mut} mice and controls. These represent 17 aberrant NMJs of 651 NMJs that were individually screened in control mice and 43 aberrant NMJs of 760 NMJs in MNDicer^{mut} mice ($n = 2$ and $n = 2$, respectively). * $P < 0.05$; ** $P < 0.01$.

(64), and deletion of the NEFH tail (was suggested as a component of ALS (61).

Previous work revealed that neurofilament expression is regulated by the 3'UTR of the mRNA. Further, the 3'UTR appears to interact with an uncharacterized *trans*-acting factor (65–67) that is attenuated in ALS (68). We reasoned that this ill-characterized *trans*-acting factor may, in fact, be a miRNA. Thus, we analyzed the relative expression levels of the neurofilament subunit proteins in MNDicer^{mut} mice and sibling controls. Quantification of

the neurofilament immunofluorescent signal in approximately 2,000 lumbar axons revealed that the expression levels of NEFL and NEFM were comparable with the WT. However, the expression of the heavy subunit (NEFH) is specifically up-regulated in the MNDicer^{mut} mouse (Fig. 4A).

Coordinated Expression of the Neurofilament Subunits Is Achieved by miR-9. The up-regulation of NEFH in the MNDicer^{mut} mouse, and consequent loss of the coordinated expression of the three neurofilament subunits could, in principle, have been attributable to a direct or indirect requirement for miRNA to mediate the levels of these genes. To assess the possibility of direct miRNA regulation, we searched the neurofilament sequences for potential miRNA-binding sites. We found a single miR-9-binding site on the NEFL mRNA. In contrast, the NEFH mRNA harbors nine miR-9-binding sites, dispersed over the 3'UTR of NEFH mRNA and the 3'-portion of the coding region (Fig. 4B). Next, we obtained a heterologous reporter assay to demonstrate that miR-9 is able to modulate the expression of NEFH and that this is dependent on the presence of miR-9-binding sites at the NEFH

mRNA (Fig. 4C). These data strongly suggest a model in which the loss of miR-9 expression or activity may result in derepression of NEFH and, subsequently, dysregulation of neurofilament stoichiometry.

miR-9 Is Specifically Down-Regulated in a Model of SMA. To relate these results to the pathogenesis observed in other models of MND, we profiled miRNA expression levels in MNs carrying an SMN1^{mut} allele, which is characteristic of the pediatric form of SMA (69). Notably, dysregulation of neurofilament expression in the Dicer1 model is reminiscent of the SMN1 mutant phenotype (70), and SMN1 is functionally engaged in miRNA-protein complexes in human cells (42, 43). Thus, we have carried out *in vitro* differentiation of ES cells harboring an SMN1 mutation into MNs (71, 72). Next, we screened a miRNA microarray (LNA oligo platform; Exiqon) with labeled RNA extracted from FACS-purified SMN1^{mut} MNs. Direct comparison of RNA from WT and SMN1^{mut} MNs revealed that the expression of only a few miRNAs is significantly decreased in SMN1^{mut} MNs. Intriguingly, the most significantly down-regulated miRNAs were miR-9 and miR-9* (Fig. 4D). These two miRNA species are processed from the same hairpin, and quantitative PCR assay revealed up to a 15-fold decrease in the expression of both miR-9 and miR-9* in SMN1^{mut} MNs relative to control (Fig. 4E).

Taken together, we present here a model for SMN disease based on Dicer1 loss of function. In this model, SMN-specific loss of miRNA activity results in denervation muscular atrophy. Additionally, changes in the expression levels of the neurofilament subunits likely contribute to the disease. This phenotype is attributed to dysregulation of miR-9, an upstream regulator of the neurofilament mRNAs. The relevance of miR-9 to MNDs originates from its neuron-specific expression and its dramatic down-regulation in SMN1-deficient MNs. It will be important to explore how miR-9 acts as an effector gene downstream of SMN1 and what are the specific subsets of SMA phenotypes governed by miR-9.

Discussion

The role of miRNA in neurons and the ways by which miRNA is involved in neurological diseases are gradually being uncovered. Thus, the loss of miRNA activity, through recombination of a Dicer1 conditional allele, was shown to cause progressive neurodegeneration in several neuronal systems (21–26, 28). Additionally, changes in the expression of specific miRNAs were reported in neurodegenerative states such as Huntington's chorea and Parkinson's disease.

Consistently, the data presented in this work suggest that miRNA dysfunction results in neurodegeneration of SMNs. As a result of miRNA malfunction in SMNs, mice develop denervation-dependent muscle atrophy. Although MNs die in all forms of ALS and SMA, Dicer1 inactivation in neurons does not always lead to cell death. For example, targeted deletion of Dicer1 in striatal neurons did not result in overt neuronal loss (28).

Early changes in animal activity were noted as early as 2 mo, and signs of the disease can be documented by EMG or histology at 4 mo of age. This suggests relatively slow progression of the disease, which is consistent with the slow onset of cell death in other Dicer1 KO models (25).

The loss of MNs and their dysfunction are often related to aggregation pathologies and to defects in their intermediate filament system (60, 64, 73). In order for the neuron to function properly, neurofilament gene expression must be tightly coordinated. We demonstrated that the coordinated expression of the neurofilament subunits is perturbed by Dicer1 loss of function, because there is specific up-regulation of just the NEFH. We were further able to link this observation to dysregulation of miR-9. The unequal affinity of miR-9 to the three neurofilament subunit mRNAs is apparently attributable to the differing number of




Fig. 4. miR-9 is specifically down-regulated in a model of SMA and is located upstream of coordinated expression of the neurofilament subunits. (A) Binned distribution of neurofilament subunit expression intensity. The percentage of axons at any intensity bin is mentioned on the y axis. NEFL (Left), NEFM (Center), and NEFH (Right). Black and gray lines represent the global mean intensity of control and MNDicer^{mut} axons, respectively. (B) Illustration of sequences cloned into luciferase reporter constructs used for functional evaluation of miR-9 interactions with neurofilament subunit mRNAs, wherein NEFH^{mut} stands for seed-mutated NEFH. Gray boxes represent miR-9-binding sites (C) Heterologous luciferase reporter assay reveals that miR-9 may function upstream of the NF subunits. Levels of luciferase activity in HEK293 cells transfected with either an empty vector or a vector over-expressing miR-9. Data are normalized to the activity of a cotransfected β -galactosidase reporter and presented as the percentage of luciferase activity in the absence of miR-9. OC-2 (a fragment of the Onecut2 3'UTR) is used as a positive control. (D and E) WT control mouse ES cells (mESCs) and SMN1^{mut} mESCs harboring a homozygous mSMN1 mutation and two copies of an hSMN2 transgene were differentiated *in vitro* into MNs. The cells were FACS-purified according to the expression of GFP transgene, driven by the Hlx9 promoter. (D) Volcano plot exemplifying the log₂ ratio of SMN1^{mut}/WT miRNA expression on the x axis and the log₁₀ P value obtained by a two-tailed Student's *t* test on the y axis. (E) Quantitative PCR analysis of miR-9 and miR-9* expression in MNs derived from SMN1^{mut} mESCs (gray bars) and WT mESCs (empty bars). **P* < 0.05; ***P* < 0.01.

miRNA-binding sites (seed matches) on these mRNAs. Thus, the NEFH mRNA harbors nine miR-9-binding sites, whereas only a single seed match is positioned within the NEFL mRNA. Consequentially, loss of miRNA activity affects these target genes differentially, as is revealed by direct axon immunostaining in the Dicer1 model and by a reporter assay, wherein miR-9 overexpression affects NEFH expression only in the presence of miR-9 seed-match sequences on NEFH 3'UTR. Together, these data imply a unique role for miRNA in parallel fine-tuning of related genes, whose coordinated expression should be tightly controlled. Our data on miR-9 may explain previous observations describing the probable presence of a *trans*-acting factor, acting at a post-transcriptional level in the regulation of proper neurofilament stoichiometry (61, 65–68).

miR-9 is a highly conserved neuronal-specific miRNA that has been shown to be involved in many facets of neurobiology. In flies, miR-9 was shown to be important for proper selection of neuronal precursors from the neuroepithelium (10), and in zebrafish, it was shown to be involved in the setting of the midbrain-hindbrain boundary (18). In mammals, the miR-9 gene is involved in a feedback loop with the repressor element-1 silencing transcription factor and its co-factor complex (REST/Co-REST) complex and acts alongside miR-124 in switching of BAF chromatin-remodeling complexes in neural development (19, 20).

The potential role of miR-9 in coordinated regulation of the neurofilament subunits corresponds to previous observations. Specifically, the neurofilament 3'UTR is essential for regulation of their expression (64). Further, we have shown that miR-9 is located downstream of SMN1, and may therefore mediate intermediate filament defects reported in SMA (70).

In the future, gain- and loss-of-function studies may help to clarify whether the neurofilament defects that are observed in many MN diseases may be modified by manipulations of miR-9 expression and what additional facets of SMN1-dependent SMA are attributed to miR-9 function.

A functional role for miRNA in specific neurological processes emerges when our observations of miR-9 action upstream of neurofilament expression are considered together with reports suggesting roles for other miRNAs, such as miR-132, miR-134, miR-124 and miR-138 (9, 11, 13, 14), in mature neurons. Importantly, Williams et al. (74) recently showed that deleting the gene encoding miR-206 in G93A-*Sod1* mice accelerated the progression of ALS symptoms and shortened survival, suggesting that miR-206 has a neuroprotective role in the postsynaptic compartment after nerve damage. This is likely, because miR-206 normally represses histone deacetylase 4, an established inhibitor of muscle reinnervation. Thus, miRNA dysfunction has direct relevance for our understanding of neurodegenerative disorders perturbing the regulation of specific target genes at the neuron or in the innervated myofiber.

The RNA-binding capability of proteins involved in MN pathologies (i.e., TDP-43 and FUS/TLS as well as SETX, SMN1, and IGHMBP2) implies that a considerable number of diseases within the MN spectrum may be directly related to RNA metabolism and posttranscriptional regulation of gene expression. Indeed, in our study, we were able to show that loss of SMN1 activity in cultured ES cell-derived MNs affects the specific ex-

pression of a subset of miRNAs, including expression of the neuronal miR-9 gene. Taken together with SMN1 physical engagement in miRNA-protein complexes (7, 71), it is plausible that SMN1 functions in miRNA bioprocessing in neurons.

The proteins TDP43 and FUS/TLS have recently revolutionized the way in which ALS is viewed, implying a pivotal role for defects in RNA regulation (30–34). Strikingly, these two proteins appear to interact physically with Drosha (3). Therefore, one possibility is that either FUS/TLS, TDP-43, or both are involved in microprocessing. This is further supported by a recent report suggesting direct role for TDP-43 in the processing of a few miRNA (37). However, these proteins were reported to be associated with RNA transport in neurons (35, 36), suggesting that they may have RNA-related regulatory roles in the cytoplasm. Finally, TDP-43 binds and regulates expression of the NEFL subunit through its 3'UTR (75), providing an intriguing hypothesis that TDP-43 may work as a cofactor of the Argonaute silencing complex.

In summary, the data presented in this work provide direct evidence for the role of miRNA in MND and substantiate our understanding of miRNA-related neurodegenerative states in general. Initial support for a functional relationship of miRNA with proteins, such as TDP-43, FUS/TLS, and SMN1, should encourage revision of MN pathologies and further exploration of miRNA-based mechanisms in ALS pathogenesis and related diseases.

Materials and Methods

Animals. We crossed a Dicer1 conditional allele (43) with a Cre-recombinase transgene driven by a cholinergic-specific promoter (VAcT-Cre) (45) (Fig. S1). Protocols for the behavioral examinations are described in *SI Text*. Needle EMG was performed with a bipolar electrode inserted into the hind-limb interosseous and gastrocnemius muscles. A scale (range: 1–7) designated the “EMG Pathology Index,” which reflects the intensity and frequency of fibrillation potentials, is described in *SI Text*. Spinal cord, ventral root, and muscle tissue preparation; staining protocols; and the antibodies used are described in *SI Text*.

Differentiation of MNs in Culture. Mouse ES cells from a Tg(Hlx9-GFP)1Tmj Tg (SMN2)89Ahmb Snn1tm1Msd/J mouse (stock no. 006570; Jackson Laboratory) (69) were differentiated into MNs as previously described (71, 72). Labeled RNA was hybridized onto a miRCURY LNA microarray (Exiqon). Quantitative PCR assays for miR-9 and miR-9* were performed with Taqman (Applied Biosystems). Constructs for the neurofilament luciferase assays are described in *SI Text*.

ACKNOWLEDGMENTS. We thank Menachem Segal, Eithan Galun, Avraham Yaron, and Yoram Groner for feedback on the work; Mike Fainzilber and Benny Shilo for remarks on the manuscript; Tali Zimmermann and Judith Chermesh for mouse husbandry; Dena Leshkowitz and Ester Feldmesser for statistics; and Cherill Banks for editorial assistance. E.H. is the incumbent of the Helen and Milton A. Kimmelman Career Development Chair. A.C. is the incumbent of the Philip Harris and Gerald Ronson Career Development Chair. This work was supported by research grants (to E.H.) from the Israel Science Foundation Legacy program, Britain-Israel Research and Academic Partnership, Nella and Leon Benozio Center for Neurological Disease, Estate of Florence Blau, and Wolfson Family Charitable Trust for miRNA and by research grants (to A.C.) from the Israel Science Foundation, Israel Ministry of Health, Nella and Leon Benozio Center for Neurosciences, Roberto and Renata Ruhman, Mr. and Mrs. Mike Kahn, Mr. Jorge David Ashkenazi, and Mr. and Mrs. Barry Wolfe. Work at the lab of L.L.R. is supported by the Spinal Muscular Atrophy Foundation and by the Harvard Stem Cell Institute.

- Bartel DP (2009) MicroRNAs: Target recognition and regulatory functions. *Cell* 136:215–233.
- Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ (2004) Processing of primary microRNAs by the Microprocessor complex. *Nature* 432:231–235.
- Gregory RI, et al. (2004) The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432:235–240.
- Lee Y, et al. (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425:415–419.
- Grishok A, et al. (2001) Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell* 106:23–34.
- Hutvagner G, Zamore PD (2002) A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 297:2056–2060.
- Kim VN, Jinju Han J, Siom MC (2009) Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol* 10:126–139.
- Filipowicz W, Bhattacharyya SN, Sonenberg N (2008) Mechanisms of post-transcriptional regulation by microRNAs: Are the answers in sight? *Nat Rev Genet* 9:102–114.
- Vo N, et al. (2005) A cAMP-response element binding protein-induced microRNA regulates neuronal morphogenesis. *Proc Natl Acad Sci USA* 102:16426–16431.
- Li Y, Wang F, Lee JA, Gao FB (2006) MicroRNA-9a ensures the precise specification of sensory organ precursors in *Drosophila*. *Genes Dev* 20:2793–2805.

11. Schrott GM, et al. (2006) A brain-specific microRNA regulates dendritic spine development. *Nature* 439:283–289.
12. Makeyev EV, Zhang J, Carrasco MA, Maniatis T (2007) The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell* 27:435–448.
13. Yu JY, Chung KH, Deo M, Thompson RC, Turner DL (2008) MicroRNA miR-124 regulates neurite outgrowth during neuronal differentiation. *Exp Cell Res* 314:2618–2633.
14. Siegel G, et al. (2009) A functional screen implicates microRNA-138-dependent regulation of the depalmitoylation enzyme APT1 in dendritic spine morphogenesis. *Nat Cell Biol* 11:705–716.
15. Makeyev EV, Maniatis T (2008) Multilevel regulation of gene expression by microRNAs. *Science* 319:1789–1790.
16. Schrott G (2009) microRNAs at the synapse. *Nat Rev Neurosci* 10:842–849.
17. Coolen M, Bally-Cuif L (2009) MicroRNAs in brain development and physiology. *Curr Opin Neurobiol* 19:461–470.
18. Leucht C, et al. (2008) MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary. *Nat Neurosci* 11:641–648.
19. Yoo AS, Staahl BT, Chen L, Crabtree GR (2009) MicroRNA-mediated switching of chromatin-remodelling complexes in neural development. *Nature* 460:642–646.
20. Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL (2008) The bifunctional microRNA miR-9/miR-9* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* 28:14341–14346.
21. Bilen J, Liu N, Burnett BG, Pittman RN, Bonini NM (2006) MicroRNA pathways modulate polyglutamine-induced neurodegeneration. *Mol Cell* 24:157–163.
22. Hebert SS, De Strooper B (2007) Molecular biology: miRNAs in neurodegeneration. *Science* 317:1120–1124.
23. Kim J, et al. (2007) A MicroRNA feedback circuit in midbrain dopamine neurons. *Science* 317:1179–1180.
24. Nelson PT, Keller JN (2007) RNA in brain disease: No longer just "the messenger in the middle." *J Neuropathol Exp Neurol* 66:461–468.
25. Schaefer A, et al. (2007) Cerebellar neurodegeneration in the absence of microRNAs. *J Exp Med* 204:1553–1558.
26. Davis TH, et al. (2008) Conditional loss of Dicer disrupts cellular and tissue morphogenesis in the cortex and hippocampus. *J Neurosci* 28:4322–4330.
27. Eacker SM, Dawson TM, Dawson VL (2009) Understanding microRNAs in neurodegeneration. *Nat Rev Neurosci* 10:837–841.
28. Cuellar TL, et al. (2008) Dicer loss in striatal neurons produces behavioral and neuroanatomical phenotypes in the absence of neurodegeneration. *Proc Natl Acad Sci USA* 105:5614–5619.
29. Cogswell, et al. (2008) Identification of miRNA changes in Alzheimer's disease brain and CSF yields putative biomarkers and insights into disease pathways. *J Alzheimer's Dis* 14:27–41.
30. Kwiatkowski TJ, Jr, et al. (2009) Mutations in the FUS/ALS gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323:1205–1208.
31. Vance C, et al. (2009) Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323:1208–1211.
32. Sreedharan J, et al. (2008) TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science* 319:1668–1672.
33. Kabashi E, et al. (2008) TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet* 40:572–574.
34. Lagier-Tourenne C, Cleveland DW (2009) Rethinking ALS: The FUS about TDP-43. *Cell* 136:1001–1004.
35. Kiebler MA, Bassell GJ (2006) Neuronal RNA granules: Movers and makers. *Neuron* 51:685–690.
36. Wang IF, Wu LS, Chang HY, Shen CK (2008) TDP-43, the signature protein of FTLD-U, is a neuronal activity-responsive factor. *J Neurochem* 105:797–806.
37. Buratti E, et al. (2010) Nuclear factor TDP-43 can affect selected microRNA levels. *FEBS J* 277:2268–2281.
38. Chen YZ, et al. (2004) DNA/RNA helicase gene mutations in a form of juvenile amyotrophic lateral sclerosis (ALS4). *Am J Hum Genet* 74:1128–1135.
39. Grohmann K, et al. (2001) Mutations in the gene encoding immunoglobulin mubinding protein 2 cause spinal muscular atrophy with respiratory distress type 1. *Nat Genet* 29:75–77.
40. Lorson CL, Hahnen E, Androphy EJ, Wirth B (1999) A single nucleotide in the SMN gene regulates splicing and is responsible for spinal muscular atrophy. *Proc Natl Acad Sci USA* 96:6307–6311.
41. Lefebvre S, et al. (1995) Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* 80:155–165.
42. Dostie J, et al. (2003) Numerous microRNPs in neuronal cells containing novel microRNAs. *RNA* 9:180–186.
43. Mourelatos Z, et al. (2002) miRNPs: A novel class of ribonucleoproteins containing numerous microRNAs. *Genes Dev* 16:720–728.
44. Harfe BD, McManus MT, Mansfield JH, Hornstein E, Tabin CJ (2005) The RNaseIII enzyme Dicer is required for morphogenesis but not patterning of the vertebrate limb. *Proc Natl Acad Sci USA* 102:10898–10903.
45. Misawa H, et al. (2003) VChT-Cre.Fast and VChT-Cre.Slow: Postnatal expression of Cre recombinase in somatomotor neurons with different onset. *Genesis* 37:44–50.
46. Yamanaka K, et al. (2008) Astrocytes as determinants of disease progression in inherited amyotrophic lateral sclerosis. *Nat Neurosci* 11:251–253.
47. Misawa H, et al. (2006) Conditional knockout of Mn superoxide dismutase in postnatal motor neurons reveals resistance to mitochondrial generated superoxide radicals. *Neurobiol Dis* 23:169–177.
48. Morris CJ, Woolf AL (1970) Mechanism of type 1 muscle fibre grouping. *Nature* 226:1061–1062.
49. Hall ED, Oostveen JA, Gurney ME (1998) Relationship of microglial and astrocytic activation to disease onset and progression in a transgenic model of familial ALS. *Glia* 23:249–256.
50. Kennel PF, Finiels F, Revah F, Mallet J (1996) Neuromuscular function impairment is not caused by motor neurone loss in FALS mice: An electromyographic study. *NeuroReport* 7:1427–1431.
51. Frey D, et al. (2000) Early and selective loss of neuromuscular synapse subtypes with low sprouting competence in motoneuron diseases. *J Neurosci* 20:2534–2542.
52. Fischer LR, et al. (2004) Amyotrophic lateral sclerosis is a distal axonopathy: Evidence in mice and man. *Exp Neurol* 185:232–240.
53. Corbin R, Olsson-Carter K, Slack F (2009) The role of microRNAs in synaptic development and function. *BMB Rep* 42:131–135.
54. Sokol NS, Xu P, Jan YN, Ambros V (2008) Drosophila let-7 microRNA is required for remodeling of the neuromusculature during metamorphosis. *Genes Dev* 22:1591–1596.
55. Fiore R, et al. (2009) Mef2-mediated transcription of the miR379-410 cluster regulates activity-dependent dendritogenesis by fine-tuning Pumilio2 protein levels. *EMBO J* 28:697–710.
56. Julien JP (1999) Neurofilament functions in health and disease. *Curr Opin Neurobiol* 9:554–560.
57. Liem RK, Messing A (2009) Dysfunctions of neuronal and glial intermediate filaments in disease. *J Clin Invest* 119:1814–1824.
58. Mersiyanova IV, et al. (2000) A new variant of Charcot-Marie-Tooth disease type 2 is probably the result of a mutation in the neurofilament-light gene. *Am J Hum Genet* 67:37–46.
59. Côté F, Collard JF, Julien JP (1993) Progressive neuronopathy in transgenic mice expressing the human neurofilament heavy gene: A mouse model of amyotrophic lateral sclerosis. *Cell* 73:35–46.
60. Larivière RC, Julien JP (2004) Functions of intermediate filaments in neuronal development and disease. *J Neurobiol* 58:131–148.
61. Al-Chalabi A, et al. (1999) Deletions of the heavy neurofilament subunit tail in amyotrophic lateral sclerosis. *Hum Mol Genet* 8:157–164.
62. Figlewicz DA, et al. (1994) Variants of the heavy neurofilament subunit are associated with the development of amyotrophic lateral sclerosis. *Hum Mol Genet* 3:1757–1761.
63. Tomkins J, et al. (1998) Novel insertion in the KSP region of the neurofilament heavy gene in amyotrophic lateral sclerosis (ALS). *NeuroReport* 9:3967–3970.
64. Thyagarajan A, Strong MJ, Szaro BG (2007) Post-transcriptional control of neurofilaments in development and disease. *Exp Cell Res* 313:2088–2097.
65. Lin H, Zhai J, Schlaepfer WW (2005) RNA-binding protein is involved in aggregation of light neurofilament protein and is implicated in the pathogenesis of motor neuron degeneration. *Hum Mol Genet* 14:3643–3659.
66. Nie Z, et al. (2002) Untranslated element in neurofilament mRNA has neuropathic effect on motor neurons of transgenic mice. *J Neurosci* 22:7662–7670.
67. Cañete-Soler R, Silberg DG, Gershon MD, Schlaepfer WW (1999) Mutation in neurofilament transgene implicates RNA processing in the pathogenesis of neurodegenerative disease. *J Neurosci* 19:1273–1283.
68. Ge WW, Leystra-Lantz C, Wen W, Strong MJ (2003) Selective loss of trans-acting instability determinants of neurofilament mRNA in amyotrophic lateral sclerosis spinal cord. *J Biol Chem* 278:26558–26563.
69. Monani UR, et al. (2000) The human centromeric survival motor neuron gene (SMN2) rescues embryonic lethality in *Smn(-/-)* mice and results in a mouse with spinal muscular atrophy. *Hum Mol Genet* 9:333–339.
70. Cifuentes-Diaz C, et al. (2002) Neurofilament accumulation at the motor endplate and lack of axonal sprouting in a spinal muscular atrophy mouse model. *Hum Mol Genet* 11:1439–1447.
71. Wichterle H, Lieberam I, Porter JA, Jessell TM (2002) Directed differentiation of embryonic stem cells into motor neurons. *Cell* 110:385–397.
72. Di Giorgio FP, Carrasco MA, Siao MC, Maniatis T, Eggan K (2007) Non-cell autonomous effect of glia on motor neurons in an embryonic stem cell-based ALS model. *Nat Neurosci* 10:608–614.
73. Xiao S, McLean J, Robertson J (2006) Neuronal intermediate filaments and ALS: A new look at an old question. *Biochim Biophys Acta* 1762:1001–1012.
74. Williams AH, et al. (2009) MicroRNA-206 delays ALS progression and promotes regeneration of neuromuscular synapses in mice. *Science* 326:1549–1554.
75. Strong MJ, et al. (2007) TDP43 is a human low molecular weight neurofilament (hNFL) mRNA-binding protein. *Mol Cell Neurosci* 35:320–327.