

# Mitochondrial Dysfunction and Oxidative Stress in Seizure-Induced Neuronal Cell Death

Yao-Chung Chuang

**Abstract-** Epilepsy is considered one of the most common neurological disorders worldwide. The burst firing associated with prolonged epileptic discharges could lead to a large number of changes and cascades of events at the cellular level. From its role as the cellular powerhouse, the mitochondrion is emerging as a key participant in cell death because of its association with an ever-growing list of apoptosis-related proteins. Prolonged seizures may result in the mitochondrial dysfunction and increased production of reactive oxygen species and nitric oxide (NO) precede neuronal cell death and cause subsequent epileptogenesis. Emerging evidences also showed that intrinsic mitochondrial apoptotic pathway may contribute to the neuropathology of human epilepsy, particularly in the hippocampus. Subsequent laboratory studies in the animal model of status epilepticus provide credence to the notion that activation of nuclear factor- $\kappa$ B upregulates NO synthase (NOS) II gene expression with temporal correlation of NOS II derived NO-, superoxide anion- and peroxynitrite-dependent reduction in mitochondrial Complex I activity, leading to apoptotic neuronal cell death in the hippocampus. These results will broaden our understanding on the intimate link between mitochondrial function, oxidative stress and mitochondria-dependent apoptotic signaling triggered by epileptic seizures. It will open a new vista in the development of more effective neuroprotective strategies against seizure-induced brain damage by modification of bioenergetic failure in the mitochondria and in the design of novel treatment perspectives for therapy-resistant forms of epilepsy.

**Key Words:** Mitochondrial dysfunction, Oxidative stress, Epileptic seizures, Cell death, Hippocampus

*Acta Neurol Taiwan 2010;19:3-15*

## INTRODUCTION

Epilepsy is considered one of the most common neurological disorders worldwide, with a prevalence of 0.5-1% in the general population<sup>(1)</sup>. It is a chronic dynamic medical problem characterized by recurrent

unprovoked seizures that often requires long-term antiepileptic drug therapy<sup>(2)</sup>. The burst firing associated with prolonged epileptic discharges could lead to a large number of changes and cascades of events at the cellular level, such as activation of glutamate receptors, changes in composition of glutamate and  $\gamma$ -aminobutyric acid

From the Department of Neurology and Center for Translational Research in Biomedical Sciences, Chang Gung Memorial Hospital-Kaohsiung Medical Center, Chang Gung University College of Medicine, Kaohsiung, Taiwan.

Received October 15, 2009. Revised November 2, 2009.

Accepted November 2, 2009.

Reprint requests and correspondence to: Yao-Chung Chuang, MD, PhD. Department of Neurology, Chang Gung Memorial Hospital-Kaohsiung, Kaohsiung County 83301, Taiwan.  
E-mail: ycchuang@adm.cgmh.org.tw

receptor, cytokine activation, oxidative stress, neurogenesis, changes in plasticity or activation of some late cell death pathways<sup>(3-5)</sup>. Clinical and epidemiological studies suggest that patients with chronic epilepsy undergo progressive brain degeneration that is accompanied by long-term behavioral changes and cognitive declines despite optimal antiepileptic drug therapy<sup>(6,7)</sup>. Status epilepticus, or the condition of prolonged epileptic seizures, is a major neurological and medical emergency that is associated with significant morbidity and mortality<sup>(8)</sup>. Status epilepticus in humans<sup>(9,10)</sup> and animal models<sup>(3,11,12)</sup> results in significant cerebral damage and increases the risk of subsequent seizures, alongside a characteristic pattern of neuronal cell loss preferentially in the hippocampus. The hippocampus is especially vulnerable with selective neuronal loss in the hilus, CA1 and CA3 regions<sup>(11,12)</sup>. As a result, there has been considerable interest in defining the molecular pathways involved in seizure-induced neuronal cell death in the vulnerable hippocampus. Oxidative stress and mitochondrial dysfunction could be acute consequences of status epilepticus that are related to the mechanism of seizure-induced neuronal cell death and subsequent epileptogenesis<sup>(13)</sup>. This article will focus on the potential role of mitochondrial dysfunction and oxidative stress in seizure-associated apoptotic cell death in the hippocampus.

## INSIGHTS FROM ANIMAL MODELS

Although animal models have many limitations, studies in animal models have made important contributions to our understanding of seizure-related neuronal injury. Animal studies provide the opportunity to examine anatomical, chemical, cellular, molecular and functional changes after seizures, as well as to identify age- or sex-specific consequences from seizures<sup>(14-16)</sup>. A wide variety of animal models of epilepsy and status epilepticus had been used, including electrical stimulation models, chemoconvulsant-induced models (e.g. kainic acid; KA, pilocarpine, picrotoxin or bicucullin), physical models (e.g. hyperthermia, or photic or auditory stimulation), genetic models (e.g. mutant, transgenic or knockout) and spontaneous seizure models (e.g. post-kindling or post-

chemoconvulsant). In the future, we can anticipate that animal models will move into a new millennium. A new trend in animal models is emerging that promises to offer powerful insights into the cause and effect of seizures. However, extrapolation of conclusion from animal data to human beings must be done with caution, despite apparent neuropathological similarities between experimental and human epilepsies<sup>(3,15,16)</sup>.

Systemic or intracerebral injection of KA, a powerful excitotoxin which stimulates a subtype of the ionotropic receptor of neurotransmitter glutamate, can result in sustained epileptic activity in the hippocampus followed by a selective pattern of brain damage that neuropathological changes are similar to human temporal lobe epilepsy<sup>(12,17)</sup>. Microinjection of KA into the CA3 subfield of hippocampus in anesthetized rats elicited seizure-like hippocampal electroencephalography (hEEG) activity and simultaneous power spectral changes in hEEG<sup>(18,19)</sup>. Based on this experimental model, we carried out electrophysiological, biochemical, immunohistochemical, anatomical and electron microscopic investigations on the ipsilateral (injection side for KA) and the contralateral (recording side for hEEG) hippocampal CA3 subfield in our recent studies<sup>(19,22)</sup>. This allowed us to ascertain that results from those analyses were consequential directly to experimental temporal lobe status epilepticus and not indirectly to KA excitotoxicity.

## NEURONAL CELL DEATH IN THE HIPPOCAMPUS FOLLOWING STATUS EPILEPTICUS

Seizure-induced neuronal cell death is no exception to the emerging complexities of the molecular control of neurodegeneration, and there is controversy as to whether cell death occurs in a programmed/controlled (apoptotic) or uncontrolled/passive (necrotic) manner. The nature of hippocampal neuronal cell death following prolonged seizures has been reported as either apoptotic<sup>(23-25)</sup>, necrotic<sup>(5,26)</sup> or both<sup>(27,28)</sup>. Necrosis is generally taken as the principal morphological phenotype of dying cells after seizures<sup>(11,29)</sup>, based at least on classical definitions and morphological criteria. However, programmed


cell death mechanisms associated with cellular apoptosis have been shown to be activated by experimental status epilepticus that supports apoptotic cell death plays an important role in seizure-induced brain damages<sup>(22-25,30)</sup>. Factors such as the variation in duration and severity of seizures, metabolic disturbances, bioenergetic failure during or after seizures and age- or genetic- specific factors may all contribute to determining the eventual pathway of cell death<sup>(3,22,23)</sup>. It is probably that severe or prolonged seizure discharges trigger both forms of neuronal cell death, whereas brief seizures may result in apoptosis in the same or different neuronal population<sup>(23)</sup>. Apoptotic neuronal cell death is much prominent after seizure-induced excitotoxic injury in immature brain compared with mature brain<sup>(3)</sup>.

Apoptosis emerged as a focus of research on seizure-induced neuronal cell death in the mid-1990s. In the earlier work, researchers detected in situ 'apoptotic' DNA fragmentation (detected by terminal deoxynucleotidyl dUTP nick end labelling; TUNEL) and DNA laddering in tissue samples from the rat brain after prolonged seizures<sup>(31)</sup>. Based on an animal model of experimental status epilepticus, our recent studies<sup>(20-22)</sup> revealed that seizure-induced apoptotic cell death was detected in the vulnerable CA1 and CA3 neurons 1-7 days after a low

dose of intrahippocampal administration of KA (Fig. 1). We recognize that during a prolonged seizure, neuronal cells may exhibit a temporary drop in adenosine triphosphate (ATP) production<sup>(22)</sup>. A critical determinant of the eventual cell death fate resides in intracellular ATP concentration, the production of which depends on the structural and functional integrity of the mitochondria. Whereas ATP depletion is associated with necrosis, ATP is required for the development of apoptosis<sup>(32)</sup>. Our recent research noted that preserved mitochondrial ultrastructural integrity and maintained energy metabolism following experimental status epilepticus is associated specifically with apoptotic, not necrotic, cell death in hippocampal CA3 or CA1 neurons<sup>(22)</sup>. Since intermediate forms of cell death with both necrotic and apoptotic features have been found after seizures<sup>(26-28)</sup>, further investigations for the detailed mechanisms of different pathways of cell death is needed.

### MITOCHONDRIA AND APOPTOTIC SIGNALLING PATHWAYS IN SEIZURE-INDUCED NEURONAL CELL DEATH

Mitochondria are ubiquitous intracellular organelles enclosed by a double membrane-bound structure. The



**Figure 1.** (A) Sections stained with cresyl violet showing neuronal cell loss in the bilateral hippocampal CA3 subfields 7 days after microinjection of kainic acid (KA, 0.5 nmol) into the left hippocampal CA3 subfield in rats. DG: dentate gyrus. Scale bar, 5 mm. (B) Laser scanning confocal microscopic images of bilateral CA3 subfield of hippocampus showing pyramidal cells that were immunoreactive to a neuronal marker, NeuN (red fluorescence), or were additionally stained positively for TUNEL (green fluorescence), 7 days after microinjection of PBS or KA (0.5 nmol) into the left hippocampal CA3 subfield in rats. Note that double-labeled neurons displayed yellow fluorescence and were denoted by arrows. Scale bar, 20  $\mu$ m.

primary function of mitochondria is the production of cellular energy in the form of ATP by the mitochondrial respiratory chain through oxidative phosphorylation<sup>(33)</sup>. Mitochondrial oxidative phosphorylation consists of five multienzyme complexes (Complexes I-V) located in the mitochondrial inner membrane. Biochemical evidence suggested that the majority of cerebral ATP consumption is used for operation of the electrogenic activity of neurons<sup>(34)</sup>. Adequate energy supply by mitochondria is essential for neuronal excitability and neuronal survival.

From its role as the cellular powerhouse, the mitochondrion is emerging as a key participant in cell death because of its association with an ever-growing list of apoptosis-related proteins<sup>(35,36)</sup>. A variety of key events in apoptosis focuses on mitochondria, including the release of several apoptogenic factors (such as cytochrome c, apoptosis-inducing factor; AIF, endonuclease G, Smac/DIABLO and HtrA2/OMI), changes in electron transport, loss of mitochondrial transmembrane potential, altered cellular oxidation-reduction, and participation of pro- and antiapoptotic Bcl-2 family proteins<sup>(35-38)</sup>. One of the decisive steps of the apoptotic cascade is related to the mitochondrial permeability transition pore (MPTP)<sup>(39)</sup>. Transient opening of these non-specific pores in the mitochondrial inner membrane under conditions of cellular stress causes the mitochondrial transmembrane potential to collapse, and triggers the release of cytochrome c and other proapoptotic molecules that initiate the apoptotic cascade. Growing evidence also suggests that cytochrome c participates in mitochondrial pathways of apoptosis by translocation to the cytoplasm where it activates caspase-3 via triggering the caspase-9 pathway<sup>(37,39)</sup>.

Emerging evidences suggest that the intrinsic mitochondrial apoptotic pathway plays an important role in neuronal cell death after seizures<sup>(21,22,25,40)</sup>. Mitochondrial calcium loading has long been known to be related with acute neuronal pathological changes following status epilepticus<sup>(41)</sup>. Using a rat model of focally evoked status epilepticus we demonstrated that cytochrome c releases from the damaged mitochondria in the hippocampal neurons within 1 day following seizures<sup>(20)</sup>. Moreover, characteristic biochemical (DNA fragmentation), histochemical (TUNEL or activated caspase-3 staining) or ultra-

structural (electron microscopy) features of apoptotic cell death were presented bilaterally in the hippocampus 1-7 days after the elicitation of sustained hippocampal seizure activity by microinjection of KA into the unilateral CA3 subfield<sup>(20-22)</sup>. Other studies have corroborated that seizures up-regulate caspase-3 within affected neuronal and glial populations in limbic regions such as the entorhinal cortex, amygdala and hippocampus<sup>(30,42-44)</sup>. Cytochrome c release in the damaged hippocampus following seizures, whereupon it associated with Apaf-1, commensurate with the appearance of activated caspases-9 and -3 and subsequently DNA fragmentation<sup>(20,44,45)</sup>.

In addition to caspase-dependent cell death pathways, mitochondria release proteins also propagate caspase-independent neuronal apoptotic cascade. While AIF is an important mitochondria release protein via caspase-independent cell death pathway, there is evidence that the calpain-mediated release of AIF is important in seizure-induced neuronal cell death<sup>(46-48)</sup>.

Bcl-2 family proteins, like caspases, are involved in regulating seizure-induced neuronal cell death. The evidence of Bcl-2 family involvement in seizure-induced neuronal cell death has also been demonstrated<sup>(49-51)</sup>. Upstream pro-apoptotic BH3 domain (Bcl-2 homology domain 3)-only members BAD, Bid and Bim, can be activated via calcium-dependent mechanisms and each was found to be activated by seizures<sup>(49,50)</sup>. In our recent studies (unpublished data), we also noted that translocation of the Bax particle from cytosol to mitochondria in hippocampal CA3 neurons 3 h after experimental status epilepticus, and this coincides with the timing of cytochrome c release. Moreover, the level of serum Bcl-2 significantly increased in patients with uncontrolled epilepsy<sup>(52)</sup>. In patients with intractable temporal lobe epilepsy, tissues from temporal neocortex expressed raised levels of Bcl-2, Bcl-XL and activated caspase-3<sup>(53)</sup>. Correlative analysis showed the expression of p53, fas and caspase-3 were positively correlated with seizure frequency in the resected samples of sclerotic hippocampi from patients with mesial temporal sclerosis<sup>(54)</sup>. These clinical evidences also suggest that intrinsic mitochondrial apoptotic pathway may contribute to the neuropathology of human epilepsy, particularly in the hippocampus.

## MITOCHONDRIAL DYSFUNCTION AND OXIDATIVE STRESS FOLLOWING EPILEPTIC SEIZURES

### Mitochondrial Dysfunction Following Epileptic Seizures

Mitochondrial oxidative phosphorylation provides the major source of ATP in the cortical neurons<sup>(34)</sup>. Sustained epileptic seizures will change the redox potential and decline the ATP content that may lead to collapse of energy production and supply in the brain<sup>(11)</sup>. However, whether mitochondrial dysfunction occurs following epileptic seizure is still under debate. There are only limited evidences for mitochondrial dysfunction associated with epilepsy and status epilepticus by both using animal models and human samples (see Table 1 for review)<sup>(19,55-61)</sup>. In our studies<sup>(19)</sup>, enzyme assay for the key enzymes of mitochondrial respiratory chain revealed significant depression of the activity of nicotinamide ade-

nine dinucleotide cytochrome c reductase (Complex I+III) in the dentate gyrus, CA1 and CA3 subfields of hippocampus following 180 minutes after KA-induced temporal lobe status epilepticus. On the other hand, the activities of succinate cytochrome c reductase (Complex II+III) and cytochrome c oxidase (Complex IV) remained unaltered. After 180 minutes of microinjection of KA into the CA3 subfield, significant mitochondrial ultrastructural injury was observed in hippocampus (Fig. 2). It follows that the prolonged epileptic seizures probably lead to a dysfunction of Complex I in the mitochondrial electron transport chain and mitochondrial ultrastructural injury in the hippocampus.


Complex I is markedly more susceptible to oxidative stress and glutathionylation than other respiratory chain complexes<sup>(62)</sup>. It is a major source of superoxide anion ( $O_2^{\cdot-}$ ), making it a candidate for increased mitochondrial reactive oxygen species (ROS) production and redox signaling<sup>(62,63)</sup>. Dysfunction of Complex I may lead to

**Table 1.** Evidences of mitochondrial dysfunction following epileptic seizures from animal and human studies

Reference	Model	Samples	Experimental findings
Kunz et al. <sup>(55)</sup>	KA-treated rats	Hippocampal slices	Increased basal energy turnover with glucose as substrate Higher uncoupled rate of respiration
Kunz et al. <sup>(56)</sup>	Temporal lobe epilepsy (human)	Hippocampal specimens	Mitochondrial Complex I deficiency and ultrastructural abnormalities of mitochondria in the epileptic focus
Cock et al. <sup>(57)</sup>	Perforant path stimulation model of rats	Whole brain tissues	Reduction of brain aconitase and $\alpha$ -ketoglutarate dehydrogenase activities Decrease in reduced glutathione levels
Kudin et al. <sup>(61)</sup>	Pilocarpine-treated rats	Hippocampal tissues and slices	Decline of the activities of Complexes I and IV and lower mitochondrial membrane potential in CA1 and CA3 subfields Decrease in mitochondrial DNA copy number in CA3
Chuang et al. <sup>(19)</sup>	Microinjection of KA into the hippocampus of rats	Hippocampal tissues	Dysfunction of Complex I in the mitochondrial electron transport chain and mitochondrial ultrastructural injury
Gibbs et al. <sup>(60)</sup>	Perforant path stimulation model of rats	Hippocampal tissues	Reductions in glutathione, $\alpha$ -ketoglutarate dehydrogenase, aconitase, citrate synthase, and Complex I activities
Gao et al. <sup>(59)</sup>	Pilocarpine-treated rats	Hippocampal tissues	Depression of mitochondrial- and nuclear-encoded COX activity and COX III expression Mitochondrial ultrastructural damage
Folbergrová et al. <sup>(58)</sup>	Intracerebroventricular infusion of homocysteic acid in rats	Cerebral cortex	Mitochondrial Complex I inhibition

KA: kainic acid, COX: cytochrome oxidase





**Figure 2.** Representative electron photomicrographs of mitochondrial ultrastructure in hippocampal CA3 subfield. (A) Intact mitochondrial ultrastructure 30 min after microinjection of kainic acid (KA) into the hippocampus. (B) Mild mitochondrial damage 180 min after microinjection of KA into the hippocampus. Note swelling of all mitochondrial spaces, particular in cristae (asterisk). (C) Severe mitochondrial damage 180 min after hippocampal application of KA. Note severe mitochondrial swelling accompanied by a disruption in membrane integrity (arrows). Scale bar: 200 nm.

incomplete mitochondrial electron transport and decreased ATP production<sup>(63)</sup>. The selective loss of Complex I activity contributes to neurodegenerative diseases such as Parkinson's disease and Huntington's disease<sup>(64)</sup>. Based on this animal model, we have demonstrated that the time-dependent selective dysfunction in activity of Complex I. Whereas the pilocarpine-treated rats with spontaneous seizures exhibited a selective decline of the activities of Complexes I and IV in the hippocampal CA1 and CA3 subfields<sup>(61)</sup>, corroborating reports of mitochondrial Complex I inhibition after seizures have since emerged from other laboratories<sup>(58,60)</sup>. This pattern of mitochondrial respiratory chain dysfunction is strengthened by the finding of patients with refractory temporal lobe epilepsy showing Complex I deficiency in the CA3 subfield<sup>(56)</sup>. Thus, we proposed that the selective dysfunction of Complex I may be linked to seizure-induced neuronal cell death in the hippocampus and play an important role in the intrinsic mitochondrial apoptotic pathway.

#### **Oxidative Stress Following Epileptic Seizures**

ROS and reactive nitrogen species has been implicated in neuronal cell death in both acute and chronic neurological diseases such as stroke, trauma, spinal cord injury, Parkinson disease, Alzheimer disease, Huntington disease, Friedreich ataxia, and amyotrophic lateral sclerosis<sup>(64,65)</sup>. Oxidative stress is thought to be an important consequence of glutamate receptor activation and excitotoxicity<sup>(13,65)</sup>, which play a critical role in epileptic brain damage<sup>(13,65)</sup>. Data from animal studies suggested that prolonged seizure activity might result in the increased production of ROS and generation of nitric oxide (NO) and peroxynitrite preceded neuronal cell death in vulnerable brain regions<sup>(20,21,66-69)</sup>.

Under normal physiological conditions, 1-2% of molecular oxygen consumed by mammalian cells is metabolized to ROS via electron leakage from mitochondrial electron transport chain. Therefore, the mitochondrial respiratory chain, in particular Complex I, is a primary site for leakage of electrons from the transport chain, leading to an increase in ROS generation<sup>(70,71)</sup>. Since mitochondria are a major source of ROS production, impaired mitochondrial respiratory chain function

and calcium-dependent depolarization of mitochondrial membrane potential may further lead to incomplete O<sub>2</sub> consumption, reduced production of ATP and exacerbated overproduction of ROS<sup>(13,63)</sup>. Free radicals can damage all cell structures, including lipids, proteins, DNA and mitochondrial membrane structure<sup>(63)</sup>. As inhibition of mitochondrial respiratory chain by prolonged seizures results in excess free radical production, and free radicals themselves are direct inhibitors of the mitochondrial respiratory chain, this can result in a vicious cycle that leads to oxidative cell damage<sup>(13,63,72)</sup>.

Nitric oxide (NO) is a free radical that is widely regarded as a molecular messenger that participates in diverse physiological processes in the central nervous system (CNS), including brain development, pain perception, neuronal plasticity, memory and behavior<sup>(73)</sup>. A pathological role for NO in neurological diseases, including epilepsy, has also been described<sup>(20,21,73-75)</sup>. The functional significance of NO in epileptic seizures, however, remains controversial. Hippocampal neuronal damage caused by excessive formation of NO has been implicated in experimental epilepsy models using acetylcholinesterase inhibitors<sup>(74)</sup> or glutamatergic receptor agonists<sup>(69,75-77)</sup> as the seizure-inducing agents. However, the roles of NO synthase (NOS) isoforms in the seizure-induced neuronal cell death are also unsettled. The mechanism that triggers limbic seizures and delayed excitotoxic damage in the hippocampus is generally attributed to NO generated by neuronal NOS (NOS I or nNOS)<sup>(74-76,78)</sup>. The role of inducible NOS (NOS II or iNOS) or endothelial NOS (NOS III or eNOS) in seizure-induced neuronal cell death is unclear. Based on the experimental model of status epilepticus, our recent studies provided credence to the notion that NO-, O<sub>2</sub><sup>•-</sup> and peroxynitrite-activated mitochondrial apoptotic signaling underlies neuronal damage induced by status epilepticus<sup>(20,21)</sup>. The repertoire of cellular events elicited by sustained seizure activity in the hippocampal CA3 subfield entails overproduction in NOS II-derived NO, increase in O<sub>2</sub><sup>•-</sup>-production, formation of peroxynitrite and depression of Complex I in the mitochondrial respiratory chain, followed by translocation of cytochrome c from mitochondria to the cytosol and activation of cas-

pase-3, leading to apoptotic cell death<sup>(19,21)</sup>. More recently, we noted that transcriptional upregulation of NOS II gene by nuclear factor- $\kappa$ B (NF- $\kappa$ B) promotes apoptotic neuronal cell death in hippocampal CA3 neurons (unpublished data).

The mitochondrial respiratory chain is sensitive to both NO- and peroxynitrite-mediated mitochondrial damage<sup>(79)</sup>. NO is known to depress mitochondrial respiratory functions, and its transnitrosylation product, S-nitrosothiol, depresses Complex I activity<sup>(79,80)</sup>. In addition, peroxynitrite can inhibit mitochondrial Complex I by tyrosine nitration<sup>(81)</sup>. Blockade of NOS II activity or reduction of peroxynitrite in the hippocampal CA3 subfield, which antagonized the reduced mitochondrial Complex I activity, also blunted apoptotic cell death induced by experimental temporal lobe status epilepticus<sup>(20,21)</sup>. Complex I of the respiratory chain has also been suggested to be a constituent of the MPTP<sup>(82)</sup>, the opening of which purportedly causes the mitochondrial transmembrane potential to collapse, and triggers the mitochondrial pathways of apoptosis by releasing cytochrome c to the cytoplasm where it activates caspase dependent death pathway<sup>(83)</sup>. It is thus reasonable to speculate that inhibition of mitochondrial respiratory Complex I by NO and peroxynitrite may trigger apoptosis following prolonged seizures by eliciting a reduction in mitochondrial transmembrane potential as a consequence of MPTP opening.

Mitochondrial production of O<sub>2</sub><sup>•-</sup> is a known contributor to apoptosis<sup>(70)</sup>, and O<sub>2</sub><sup>•-</sup>-dependent mitochondrial signaling in apoptotic cell death after brain damage has been reported<sup>(84)</sup>. We also demonstrated recently that O<sub>2</sub><sup>•-</sup> production in the hippocampal CA3 subfield manifested a significant elevation 3-24 h following induction of experimental temporal lobe status epilepticus<sup>(21)</sup>. Compounds with O<sub>2</sub><sup>•-</sup> scavenger, Tempol, or an electron carrier protein, coenzyme Q10 ameliorated seizure-induced O<sub>2</sub><sup>•-</sup> production and the depression of Complex I, and prevent apoptotic cell death in the hippocampus further suggest an association between mitochondrial respiratory activities and the generation of O<sub>2</sub><sup>•-</sup> in the hippocampus during experimental status epilepticus<sup>(21)</sup>. In addition, NO augments the generation of reactive oxy-

gen species through interaction with components of the mitochondrial electron transport chain, thereby triggering mechanisms of cell death<sup>(85)</sup>.

### THE ROLE OF MITOCHONDRIAL DYSFUNCTION AND OXIDATIVE STRESS IN EPILEPTOGENESIS

Because neuronal cell death may be an important factor contributing to epileptogenesis, mechanisms that influence neuronal viability may also play a role in the process of epileptogenesis. As impairment of mitochondrial function and increased ROS have recently been observed in the seizure focus of human and experimental epilepsy<sup>(13,86)</sup>, the crucial question is whether seizure-induced free radical production and mitochondrial dysfunction results in chronic redox alterations in neurons that increase seizure susceptibility and lead to the development of subsequent epilepsy. The most prominent example of mitochondrial dysfunction causing epilepsy is the occurrence of epileptic seizures in mitochondrial diseases arising from mutations in mitochondrial DNA (mtDNA) or nuclear DNA<sup>(86-90)</sup>. Defects in the process of oxidative phosphorylation in the CNS are a characteristic sign of mitochondrial encephalomyopathies. The most common mitochondrial disorders presenting with an epileptic phenotype has been well reviewed by Kudin et al<sup>(86)</sup>. A well known mitochondrial disorder with generalized seizures which is linked to point mutations in the mitochondrial tRNA<sup>Leu</sup> gene<sup>(90)</sup> is the myoclonus epilepsy with ragged red fibers (MERRF) syndrome. Partial seizures are frequently noticed in mitochondrial encephalopathy with lactic acidosis and stroke-like episodes (MELAS) syndrome, which is associated with mutations in the mitochondrial tRNA<sup>Leu</sup> gene<sup>(88,89)</sup>. In addition, systemic administration of mitochondrial toxins, such as 3-nitropropionic acid<sup>(91)</sup> and cyanide<sup>(92)</sup>, inhibits the functions of mitochondrial respiratory chain that can compromise cellular energy metabolism and induce seizures in animal models. These accumulating evidences implicated that both mtDNA mutations and exogenous mitochondrial toxins cause mitochondrial respiratory chain dysfunction which is associated with at

least some of the mechanisms of epileptogenesis.

Several common neurological conditions such as hypoxia, stroke, traumatic brain injury, aging and neurodegenerative diseases render the brain susceptible to epileptic seizures<sup>(13,93)</sup>. In fact, increased oxidative stress and mitochondrial dysfunction is the common cellular events under these neuropathologic conditions. Mice with partial deficiency of the mitochondrial superoxide dismutase show increased incidence of spontaneous and handling-induced seizures that correlates with chronic mitochondrial oxidative stress<sup>(13)</sup>. Increased oxidative mtDNA damage, mitochondrial H<sub>2</sub>O<sub>2</sub> production and alterations in the mitochondrial base excision repair pathway have been noted in the rat hippocampus after a period of 3 months following status epilepticus. The data provide evidence for mitochondrial oxidative stress in epilepsy and suggest that mitochondrial injury may contribute to epileptogenesis<sup>(94)</sup>. These evidences raise an intriguing possibility that mitochondrial dysfunction initiated by free radical production could increase susceptibility of seizure<sup>(13)</sup>.

The mechanisms of mitochondrial dysfunction and oxidative stress during epileptogenesis remain unclear. Since mitochondrial oxidative phosphorylation provides the major source of ATP in neurons and mitochondria participate in intracellular calcium homeostasis, their dysfunction strongly affects neuronal excitability and synaptic transmission<sup>(13,86)</sup>. Thus, decreased intracellular ATP levels and alterations of neuronal calcium homeostasis may be potential factors contributing to increased susceptibility of epileptic seizures associated with mitochondrial dysfunction. These changes strongly affect neuronal excitability and synaptic transmission, which is proposed to be highly relevant for seizure generation<sup>(13,86)</sup>. Further studies are mandatory in the future to confirm this implication.

### CONCLUSION

Oxidative stress and mitochondrial dysfunction occur as a consequence of prolonged epileptic seizures and contribute seizure-induced neuronal cell death. Subsequent laboratory studies<sup>(19-22)</sup> in the animal model of



status epilepticus provide credence to the notion that activation of NF- $\kappa$ B in hippocampal CA3 neurons upregulates NOS II gene expression with temporal correlation of NOS II derived NO-, O<sub>2</sub><sup>•-</sup> and peroxynitrite-dependent reduction in mitochondrial respiratory enzyme Complex I activity, leading to apoptotic neuronal cell death in the hippocampus (summary in Fig. 3). Recent studies suggested that mitochondrial dysfunction and chronic oxidative stress can render the brain more susceptible to epileptic seizures<sup>(13,94,95)</sup>. Therefore, both of mitochondrial dysfunction and oxidative stress are important causes and consequences of prolonged seizures. In keeping with the role of mitochondrial dysfunction and oxidative stress in the pathogenesis of apoptosis, protection of the mitochondrion from NO-,


O<sub>2</sub><sup>•-</sup> or peroxynitrite-promoted neuronal stress in the hippocampus may therefore become a novel target for therapeutic strategy against seizure-induced brain damage and epileptogenesis<sup>(13,37,83,85)</sup>.

## ACKNOWLEDGEMENTS

The work from our laboratories is supported by research grants from the National Science Council, Taiwan.

## REFERENCES

1. Hauser WA, Annegers JF, Kurland LT. Prevalence of epilepsy in Rochester, Minnesota: 1940-1980. *Epilepsia* 1991;32: 429-45.
2. Kwan P, Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med* 2000;342:314-9.
3. Haut SR, Velískova J, Moshé SL. Susceptibility of immature and adult brains to seizure effects. *Lancet Neurol* 2004; 3:608-17.
4. Macdonald RL, Kapur J. Acute cellular alterations in the hippocampus after status epilepticus. *Epilepsia* 1999; 40(Suppl 1):S9-20.
5. Fujikawa DG. Prolonged seizures and cellular injury: understanding the connection. *Epilepsy Behav* 2005; 7(Suppl 3):S3-11.
6. Cendes F. Progressive hippocampal and extrahippocampal atrophy in drug resistant epilepsy. *Curr Opin Neurol* 2005; 18:173-7.
7. Sutula TP, Hagen J, Pitkänen A. Do epileptic seizures damage the brain? *Curr Opin Neurol* 2003;16:189-95.
8. Lowenstein DH, Alldredge BK. Status epilepticus. *N Engl J Med* 1998;338:970-6.
9. DeGiorgio CM, Tomiyasu U, Gott PS, et al. Hippocampal pyramidal cell loss in human status epilepticus. *Epilepsia* 1992;33:23-7.
10. Duncan JS. Seizure-induced neuronal injury: human data. *Neurology* 2002;59:S15-20.
11. Wasterlain CG, Fujikawa DG, Penix L, et al. Pathophysiological mechanisms of brain damage from status epilepticus. *Epilepsia* 1993;34(Suppl 1):S37-53.
12. Ben-Ari Y. Limbic seizure and brain damage produced by



**Figure 3.** A schematic outline of the proposed role of mitochondrial dysfunction and oxidative stress in seizure-induced neuronal cell death from subsequent laboratory studies in the animal model of status epilepticus.

- kainic acid: mechanisms and relevance to human temporal lobe epilepsy. *Neuroscience* 1985;14:375-403.
13. Patel M. Mitochondrial dysfunction and oxidative stress: cause and consequence of epileptic seizures. *Free Radic Biol Med* 2004;37:1951-62.
  14. Fisher RS. Animal models of the epilepsies. *Brain Res Rev* 1989;14:245-78.
  15. Cole AJ, Koh S, Zheng Y. Are seizures harmful: what can we learn from animal models? *Prog Brain Res* 2002;135:13-23.
  16. Löscher W. Animal models of intractable epilepsy. *Prog Neurobiol* 1997;53:239-58.
  17. Schwob JE, Fuller T, Price JL, et al. Widespread patterns of neuronal damage following systemic or intracerebral injections of kainic acid: a histological study. *Neuroscience* 1980;5:991-1014.
  18. Lin YY, Yen SH, Pan JT, et al. Transient elevation in plasma prolactin level in rats with temporal lobe status epilepticus. *Neurology* 1999;53:885-7.
  19. Chuang YC, Chang AYW, Lin JW, et al. Mitochondrial dysfunction and ultrastructural damage in the hippocampus during kainic acid-induced status epilepticus in the rat. *Epilepsia* 2004;45:1202-9.
  20. Chuang YC, Chen SD, Lin TK, et al. Upregulation of nitric oxide synthase II contributes to apoptotic cell death in the hippocampal CA3 subfield via a cytochrome *c*/caspase-3 signaling cascade following induction of experimental temporal lobe status epilepticus in the rat. *Neuropharmacology* 2007;52:1263-73.
  21. Chuang YC, Chen SD, Liou CW, et al. Contribution of nitric oxide, superoxide anion, and peroxynitrite to activation of mitochondrial apoptotic signaling in hippocampal CA3 subfield following experimental temporal lobe status epilepticus. *Epilepsia* 2009;50:731-46.
  22. Chuang YC, Lin JW, Chen SD, et al. Preservation of mitochondrial integrity and energy metabolism during experimental status epilepticus leads to neuronal apoptotic cell death in the hippocampus of the rat. *Seizure* 2009;18:420-8.
  23. Bengzon J, Mohapel P, Ekdahl CT, et al. Neuronal apoptosis after brief and prolonged seizures. *Prog Brain Res* 2002;135:111-9.
  24. Mikati MA, Abi-Habib RJ, El Sabban ME, et al. Hippocampal programmed cell death after status epilepticus: evidence for NMDA-receptor and ceramide-mediated mechanisms. *Epilepsia* 2003;44:282-91.
  25. Henshall DC, Simon RP. Epilepsy and apoptosis pathways. *J Cereb Blood F MET* 2005;25:1557-72.
  26. Fujikawa DG, Shinmei SS, Cai B. Kainic acid-induced seizures produce necrotic, not apoptotic, neurons with internucleosomal DNA cleavage: implications for programmed cell death mechanisms. *Neuroscience* 2000;98:41-53.
  27. Covolán L, Smith RL, Mello LEAM. Ultrastructural identification of dentate granule cell death from pilocarpine-induced seizures. *Epilepsy Res* 2000;41:9-21.
  28. Sloviter RS, Dean E, Sollas AL, et al. Apoptosis and necrosis induced in different hippocampal neuron populations by repetitive perforant path stimulation in the rat. *J Comp Neurol* 1996;366:516-33.
  29. Fujikawa DG, Shinmei SS, Cai B. Seizure-induced neuronal necrosis: implications for programmed cell death mechanisms. *Epilepsia* 2000;41(Suppl 6):S9-13.
  30. Weise J, Engelhorn T, Dorfler A, et al. Expression time course and spatial distribution of activated caspase-3 after experimental status epilepticus: contribution of delayed neuronal cell death to seizure-induced neuronal injury. *Neurobiol Dis* 2005;18:582-90.
  31. Pollard H, Charriaut-Marlangue C, Cantagrel S, et al. Kainate-induced apoptotic cell death in hippocampal neurons. *Neuroscience* 1994;63:7-18.
  32. Leist M, Single B, Castoldi AF, et al. Intracellular adenosine triphosphate (ATP) concentration: a switch in the decision between apoptosis and necrosis. *J Exp Med* 1997;185:1481-6.
  33. Hatefi Y. The mitochondrial electron transport and oxidative phosphorylation system. *Annu Rev Biochem* 1985;54:1015-69.
  34. Ames III A. CNS energy metabolism as related to function. *Brain Res Rev* 2000;34:42-68.
  35. Kroemer G. Mitochondrial control of apoptosis: an overview. *Biochem Soc Symp* 1999;66:1-15.
  36. Daugas E, Nochy D, Ravagnan L, et al. Apoptosis-inducing factor (AIF): a ubiquitous mitochondrial oxidoreductase involved in apoptosis. *FEBS Lett* 2000;476:118-23.
  37. Green DR, Reed JC. Mitochondria and apoptosis. *Science* 1998;281:1309-12.
  38. Saelens X, Festjens N, Vande Walle L, et al. Toxic proteins released from mitochondria in cell death. *Oncogene* 2004;

- 23:2861-74.
39. Crompton M. Mitochondrial intermembrane junctional complexes and their role in cell death. *J Physiol* 2000; 529(Pt 1):11-21.
  40. Liou AKF, Clark RS, Henshall DC, et al. To die or not to die for neurons in ischemia, traumatic brain injury and epilepsy: a review on the stress-activated signaling pathways and apoptotic pathways. *Prog Neurobiol* 2003;69: 103-42.
  41. Griffiths T, Evans MC, Meldrum BS. Status epilepticus: the reversibility of calcium loading and acute neuronal pathological changes in the rat hippocampus. *Neuroscience* 1984;12:557-67.
  42. Kondratyev A, Gale K. Intracerebral injection of caspase-3 inhibitor prevents neuronal apoptosis after kainic acid-evoked status epilepticus. *Brain Res Mol Brain Res* 2000; 75:216-24.
  43. Narkilahti S, Pirttilä TJ, Lukasiuk K, et al. Expression and activation of caspase 3 following status epilepticus in the rat. *Eur J Neurosci* 2003;18:1486-96.
  44. Henshall DC, Chen J, Simon RP. Involvement of caspase-3-like protease in the mechanism of cell death following focally evoked limbic seizures. *J Neurochem* 2000;74: 1215-23.
  45. Henshall DC, Bonislawski DP, Skradski SL, et al. Formation of the Apaf-1/cytochrome c complex precedes activation of caspase-9 during seizure-induced neuronal death. *Cell Death Differ* 2001;8:1169-81.
  46. Schindler CK, Pearson EG, Bonner HP, et al. Caspase-3 cleavage and nuclear localization of caspase-activated DNase in human temporal lobe epilepsy. *J Cereb Blood Flow Metab* 2006;26:583-9.
  47. Cheung EC, Melanson-Drapeau L, Cregan SP, et al. Apoptosis-inducing factor is a key factor in neuronal cell death propagated by BAX-dependent and BAX-independent mechanisms. *J Neurosci* 2005;25:1324-34.
  48. Lankiewicz S, Marc Luetjens C, Truc Bui N, et al. Activation of calpain I converts excitotoxic neuron death into a caspase-independent cell death. *J Biol Chem* 2000; 275:17064-71.
  49. Henshall DC, Araki T, Schindler CK, et al. Activation of Bcl-2-associated death protein and counter-response of Akt within cell populations during seizure-induced neuronal death. *J Neurosci* 2002;22:8458-65.
  50. Shinoda S, Schindler CK, Meller R, et al. Bim regulation may determine hippocampal vulnerability after injurious seizures and in temporal lobe epilepsy. *J Clin Invest* 2004; 113:1059-68.
  51. Korhonen L, Belluardo N, Mudo G, et al. Increase in Bcl-2 phosphorylation and reduced levels of BH3-only Bcl-2 family proteins in kainic acid-mediated neuronal death in the rat brain. *Eur J Neurosci* 2003;18:1121-34.
  52. El-Hodhod MA, Tomoum HY, Abd Al-Aziz MM, et al. Serum Fas and Bcl-2 in patients with epilepsy. *Acta Neurol Scand* 2006;113:315-21.
  53. Henshall DC, Clark RS, Adelson PD, et al. Alterations in bcl-2 and caspase gene family protein expression in human temporal lobe epilepsy. *Neurology* 2000;55:250-7.
  54. Xu S, Pang Q, Liu Y, et al. Neuronal apoptosis in the resected sclerotic hippocampus in patients with mesial temporal lobe epilepsy. *J Clin Neurosci* 2007;14:835-40.
  55. Kunz WS, Goussakov IV, Beck H, et al. Altered mitochondrial oxidative phosphorylation in hippocampal slices of kainate-treated rats. *Brain Res* 1999;826:236-42.
  56. Kunz WS, Kudin AP, Vielhaber S, et al. Mitochondrial complex I deficiency in the epileptic focus of patients with temporal lobe epilepsy. *Ann Neurol* 2000;48:766-73.
  57. Cock HR, Tong X, Hargreaves IP, et al. Mitochondrial dysfunction associated with neuronal death following status epilepticus in rat. *Epilepsy Res* 2002;48:157-68.
  58. Folbergrová J, Ješina P, Drahotka Z, et al. Mitochondrial complex I inhibition in cerebral cortex of immature rats following homocysteic acid-induced seizures. *Exp Neurol* 2007;204:597-609.
  59. Gao J, Chi ZF, Liu XW, et al. Mitochondrial dysfunction and ultrastructural damage in the hippocampus of pilocarpine-induced epileptic rat. *Neurosci Lett* 2007;411:152-7.
  60. Gibbs JE, Walker MC, Cock HR. Levetiracetam: antiepileptic properties and protective effects on mitochondrial dysfunction in experimental status epilepticus. *Epilepsia* 2006;47:469-78.
  61. Kudin AP, Kudina TA, Seyfried J, et al. Seizure-dependent modulation of mitochondrial oxidative phosphorylation in rat hippocampus. *Eur J Neurosci* 2002;15:1105-14.
  62. Taylor ER, Hurrell F, Shannon RJ, et al. Reversible glutathionylation of complex I increases mitochondrial superoxide formation. *J Biol Chem* 2003;278:19603-10.

63. Cadenas E, Davies KJ. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med* 2000;29:222-30.
64. Beal MF. Energetics in the pathogenesis of neurodegenerative diseases. *Trends Neurosci* 2000;23:298-304.
65. Coyle JT, Puttfarcken P. Oxidative stress, glutamate, and neurodegenerative disorders. *Science* 1993;262:689-95.
66. Frantseva MV, Perez Velazquez JL, Tsoraklidis G, et al. Oxidative stress is involved in seizure-induced neurodegeneration in the kindling model of epilepsy. *Neuroscience* 2000;97:431-5.
67. Liang LP, Ho YS, Patel M. Mitochondrial superoxide production in kainate-induced hippocampal damage. *Neuroscience* 2000;101:563-70.
68. Bruce AJ, Baudry M. Oxygen free radicals in rat limbic structures after kainate-induced seizures. *Free Radic Biol Med* 1995;18:993-1002.
69. Milatovic D, Gupta RC, Dettbarn WD. Involvement of nitric oxide in kainic acid-induced excitotoxicity in rat brain. *Brain Res* 2002;957:330-7.
70. Raha S, Robinson BH. Mitochondria, oxygen free radicals, disease and ageing. *Trends Biochem Sci* 2000;25:502-8.
71. Li N, Ragheb K, Lawler G, et al. Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J Biol Chem* 2003;278:8516-25.
72. Cock HR. The role of mitochondria and oxidative stress in neuronal damage after brief and prolonged seizures. *Prog Brain Res* 2002;135:187-96.
73. Guix FX, Uribesalgo I, Coma M, et al. The physiology and pathophysiology of nitric oxide in the brain. *Prog Neurobiol* 2005;76:126-52.
74. Bagetta G, Paoletti AM, Leta A, et al. Abnormal expression of neuronal nitric oxide synthase triggers limbic seizures and hippocampal damage in rat. *Biochem Biophys Res Commun* 2002;291:255-60.
75. Yasuda H, Fujii M, Fujisawa H, et al. Changes in nitric oxide synthesis and epileptic activity in the contralateral hippocampus of rats following intrahippocampal kainate injection. *Epilepsia* 2001;42:13-20.
76. Gupta RC, Dettbarn WD. Prevention of kainic acid seizures-induced changes in levels of nitric oxide and high-energy phosphates by 7-nitroindazole in rat brain regions. *Brain Res* 2003;981:184-92.
77. Montcot C, Rondi-Reig L, Springhetti V, et al. Inhibition of neuronal (type 1) nitric oxide synthase prevents hyperaemia and hippocampal lesions resulting from kainate-induced seizures. *Neuroscience* 1998;84:791-800.
78. Rajasekaran K. Seizure-induced oxidative stress in rat brain regions: blockade by nNOS inhibition. *Pharmacol Biochem Behav* 2005;80:263-72.
79. Stewart VC, Heales SJ. Nitric oxide-induced mitochondrial dysfunction: implications for neurodegeneration. *Free Radic Biol Med* 2003;34:287-303.
80. Clementi E, Brown GC, Feelisch M, et al. Persistent inhibition of cell respiration by nitric oxide: crucial role of S-nitrosylation of mitochondrial complex I and protective action of glutathione. *Proc Natl Acad Sci USA* 1998;95:7631-6.
81. Yamamoto T, Maruyama W, Kato Y, et al. Selective nitration of mitochondrial complex I by peroxynitrite: involvement in mitochondria dysfunction and cell death of dopaminergic SH-SY5Y cells. *J Neural Transm* 2002;109:1-13.
82. Chauvin C, De Oliveira F, Ronot X, et al. Rotenone inhibits the mitochondrial permeability transition-induced cell death in U937 and KB cells. *J Biol Chem* 2001;276:41394-8.
83. Kroemer G, Reed JC. Mitochondrial control of cell death. *Nat Med* 2000;6:513-9.
84. Lossi L, Merighi A. In vivo cellular and molecular mechanisms of neuronal apoptosis in the mammalian CNS. *Prog Neurobiol* 2003;69:287-312.
85. Moncada S, Erusalimsky JD. Does nitric oxide modulate mitochondrial energy generation and apoptosis? *Nat Rev Mol Cell Biol* 2002;3:214-20.
86. Kudin AP, Zsurka G, Elger CE, et al. Mitochondrial involvement in temporal lobe epilepsy. *Exp Neurol* 2009;218:326-32.
87. Simon DK, Johns DR. Mitochondrial disorders: clinical and genetic features. *Annu Rev Med* 1999;50:111-27.
88. DiMauro S, Kulikova R, Tanji K, et al. Mitochondrial genes for generalized epilepsies. *Adv Neurol* 1999;79:411-9.
89. Canafoglia L, Franceschetti S, Antozzi C, et al. Epileptic phenotypes associated with mitochondrial disorders. *Neurology* 2001;56:1340-6.
90. Shoffner JM, Lott MT, Lezza AM, et al. Myoclonic epilep-

- sy and ragged-red fiber disease (MERRF) is associated with a mitochondrial DNA tRNA(Lys) mutation. *Cell* 1990; 61:931-7.
91. Urbanska EM, Blaszcak P, Saran T, et al. Mitochondrial toxin 3-nitropropionic acid evokes seizures in mice. *Eur J Pharmacol* 1998;359:55-8.
92. Yamamoto H. A hypothesis for cyanide-induced tonic seizures with supporting evidence. *Toxicology* 1995;95:19-26.
93. Hauser WA, Annegers JF. Risk factors for epilepsy. *Epilepsy Res Suppl* 1991;4:45-52.
94. Jarrett SG, Liang LP, Hellier JL, et al. Mitochondrial DNA damage and impaired base excision repair during epileptogenesis. *Neurobiol Dis* 2008;30:130-8.
95. Liang LP, Patel M. Mitochondrial oxidative stress and increased seizure susceptibility in Sod2(-/+) mice. *Free Radic Biol Med* 2004;36:542-4.