

## MIXED PROBLEM WITH BOUNDARY INTEGRAL CONDITIONS FOR A CERTAIN PARABOLIC EQUATION

ABDELFATAH BOUZIANI  
*Institut National d'Enseignement  
Supérieur de Mécanique  
Oum El Bouaghi, Algérie*

(Received October, 1994; Revised December, 1995)

### ABSTRACT

The present article is devoted to a proof of the existence and uniqueness of a solution of a mixed problem with boundary integral conditions for a certain parabolic equation. The proof is based on an energy inequality and on the fact that the range of the operator generated by the problem is dense.

**Key words:** Parabolic Equation, Boundary Integral Conditions, Energy Inequality.

**AMS (MOS) subject classifications:** 34B99, 34B15.

### 1. Introduction

In the rectangle  $Q = (0, b) \times (0, T)$ , we consider the equation

$$\mathcal{L}u = \frac{\partial u}{\partial t} + (-1)^m a(t) \frac{\partial^{2m} u}{\partial x^{2m}} = f(x, t), \quad (1.1)$$

where  $a(t)$  is bounded,  $0 < a_0 \leq a(t) \leq a_1$ , and  $a(t)$  has the bounded derivative such that  $0 < c_0 \leq a'(t) \leq c_1$  for  $t \in [0, T]$ .

We adhere to equation (1.1) the initial condition

$$\ell u = u(x, 0) = \varphi(x) \quad (1.2)$$

and the boundary conditions

$$\int_0^b x^k \cdot u(x, t) dx = 0, \quad k = \overline{0, 2m-1}. \quad (1.3)$$

The importance of problems with integral conditions has been pointed out by Samarskii [9]. Problems which combine local and integral condition for second order parabolic equations are investigated by the potential method [2, 7], by Fourier's method [4-6], and by the energy inequalities method [1, 8, 10].

In this paper, the existence and uniqueness of a solution of problem (1.1)-(1.3) is proved. The proof is based on the method of energy inequalities, presented in [1]. Such problems have not been studied previously.

## 2. Preliminaries

First, we introduce the appropriate function spaces which will be used in the paper. We denote  $B_2^m(0, b)$  by:

$$B_2^m(0, b) := \begin{cases} L^2(0, b) & \text{for } m = 0, \\ \{u/\mathfrak{J}^m u \in L^2(0, b)\} & \text{for } m \geq 1, \end{cases} \quad (2.1)$$

where  $\mathfrak{J}^m u := \int_0^x \frac{(x-\xi)^{m-1}}{(m-1)!} u(\xi, t) d\xi$ ,  $m \geq 1$ . For  $m \geq 1$ , the scalar product in  $B_2^m(0, b)$  is defined by:

$$(u, v)_{B_2^m(0, b)} = \int_0^b \mathfrak{J}^m u \mathfrak{J}^m v dx.$$

The associated norm is:

$$\|u\|_{B_2^m(0, b)} = \|\mathfrak{J}^m u\|_{L^2(0, b)} \text{ for } m \geq 1.$$

**Lemma 1:** For  $m \in \mathbb{N}$ , we have

$$\|u\|_{B_2^m(0, b)}^2 \leq \frac{b^2}{2} \|u\|_{B_2^{m-1}(0, b)}^2. \quad (2.2)$$

**Proof:** The Cauchy-Schwarz inequality gives

$$\begin{aligned} |\mathfrak{J}^m u|^2 &\leq \left| \int_0^x \mathfrak{J}^{m-1} u(\xi, t) d\xi \right|^2 \leq \left( \int_0^x d\xi \right) \cdot \left( \int_0^x |\mathfrak{J}^{m-1} u(\xi, t)|^2 d\xi \right) \\ &\leq x \cdot \int_0^x |\mathfrak{J}^{m-1} u(\xi, t)|^2 d\xi \leq x \cdot \int_0^b |\mathfrak{J}^{m-1} u(\xi, t)|^2 d\xi. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \|u\|_{B_2^m(0, b)}^2 &\leq \int_0^b |\mathfrak{J}^{m-1} u(\xi, t)|^2 d\xi \cdot \int_0^b x dx \\ &= \frac{b^2}{2} \|u\|_{B_2^{m-1}(0, b)}^2. \quad \square \end{aligned}$$

**Corollary:** For  $m \in \mathbb{N}$ , we have

$$\|u\|_{B_2^m(0, b)}^2 \leq \left(\frac{b^2}{2}\right)^m \cdot \|u\|_{L^2(0, b)}^2. \quad (2.3)$$

**Remark:** Inequalities (2.2) and (2.3) remain valid, if we replace the interval  $(0, b)$  by a bounded region  $\Omega$  of  $\mathbb{R}^n$ . It suffices to replace  $b$  by  $\text{meas}(\Omega)$  (measure of  $\Omega$ ) in (2.2) and (2.3).  $\square$

The space  $B_2^{m, k}(Q)$  is the space with the finite norm

$$\|u\|_{B_2^{m, k}(Q)}^2 = \int_0^T \|u(\cdot, t)\|_{B_2^m(0, b)}^2 dt + \int_0^b \|u(x, \cdot)\|_{B_2^k(0, T)}^2 dx.$$

The space  $B_2^{0,0}(Q)$  coincides with  $L^2(Q)$ .

We associate with problem (1.1)-(1.3), the operator  $L = (\mathfrak{L}, \ell)$  with domain denoted by  $D(L) = : E$ . The operator  $L$  is from  $E$  to  $F$ ;  $E$  is Banach space of the functions  $u \in L^2(0, b)$  satisfying (1.3), with the finite norm

$$\|u\|_E^2 = \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q)}^2 + \left\| \frac{\partial^{2m} u}{\partial x^{2m}} \right\|_{B_2^{m,0}(Q)}^2 + \sup_{0 \leq \tau \leq T} \|u(x, \tau)\|_{L^2(0, b)}^2, \tag{2.4}$$

where  $F$  is the Hilbert space obtained by completing the space  $B_2^{m,0}(Q) \times L^2(0, b)$  equipped with the norm

$$\|\mathfrak{F}\|_F^2 = \|f\|_{B_2^{m,0}(Q)}^2 + \|\varphi\|_{L^2(0, b)}^2, \mathfrak{F} = (f, \varphi). \tag{2.5}$$

Here, we assumed that the function  $\varphi$  satisfies the conditions in the form (1.3), i.e.,

$$\int_0^b x^k \cdot \varphi dx = 0, \quad k = \overline{0, 2m-1}. \tag{2.7}$$

### 3. Two-Sided A Priori Estimates

**Theorem 1:** *The following a priori estimate*

$$\|Lu\|_F \leq c \|u\|_E \tag{3.1}$$

holds for any function  $u \in E$ , where constant  $c$  is independent of  $u$ .

**Proof:** Equation (1.1) implies that

$$\|\ell u\|_{B_2^{m,0}(Q)}^2 \leq 2 \left( \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q)}^2 + a_1^2 \left\| \frac{\partial^{2m} u}{\partial x^{2m}} \right\|_{B_2^{m,0}(Q)}^2 \right) \tag{3.2}$$

and initial condition (1.2) yields

$$\|\mathfrak{L}u\|_{L^2(0, b)}^2 \leq \sup_{0 \leq \tau \leq T} \|u(x, \tau)\|_{L^2(0, b)}^2. \tag{3.3}$$

Combining inequality (3.2) with (3.3), we obtain (3.1) for  $u \in E$ , with  $c := \max(2^{1/2}, 2^{1/2}a_1)$ .  $\square$

**Theorem 2:** *For any function  $u \in E$ , we have the inequality*

$$\|u\|_E \leq c \|Lu\|_F, \tag{3.4}$$

where constant  $c > 0$  does not depend on  $u$ .

**Proof:** We consider the scalar product in  $L^2(Q^\tau)$ , where  $Q^\tau := (0, b) \times (0, \tau)$  and  $0 \leq \tau \leq T$ . Observe that

$$\begin{aligned} & 2 \int_{Q^\tau} \left| \mathfrak{T}^m \frac{\partial u}{\partial t} \right|^2 dx dt + \int_0^b a(\tau) |u(x, \tau)|^2 dx \\ &= 2 \operatorname{Re} \left( \mathfrak{L}u, (-1)^m \mathfrak{T}^{2m} \frac{\partial \bar{u}}{\partial t} \right)_{0, Q^\tau} + \int_0^b a(0) |\varphi|^2 dx + \int_{Q^\tau} a'(t) |u|^2 dx dt \end{aligned} \tag{3.5}$$

We estimate the first term on the right-hand side of (3.5). By applying an elementary inequality we have

$$2 \operatorname{Re} \left( \mathcal{L}u, (-1)^m \mathcal{F}^{2m} \frac{\partial \bar{u}}{\partial t} \right)_{0, Q^\tau} \leq \| \mathcal{L}u \|_{B_2^{m,0}(Q^\tau)}^2 + \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q^\tau)}^2. \tag{3.6}$$

From equation (1.1), we obtain

$$\frac{1}{4} a_0^2 \left\| \frac{\partial^{2m} u}{\partial x^{2m}} \right\|_{B_2^{m,0}(Q^\tau)}^2 \leq \frac{1}{2} \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q^\tau)}^2 + \frac{1}{2} \| \mathcal{L}u \|_{B_2^{m,0}(Q^\tau)}^2. \tag{3.7}$$

Therefore, by formulas (3.5)-(3.7),

$$\begin{aligned} & \frac{1}{2} \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q^\tau)}^2 + \frac{1}{4} a_0^2 \left\| \frac{\partial^{2m} u}{\partial x^{2m}} \right\|_{B_2^{m,0}(Q^\tau)}^2 + a_0 \| u(x, \tau) \|_{L^2(0,b)}^2 \\ & \leq \frac{3}{2} \| \mathcal{L}u \|_{B_2^{m,0}(Q^\tau)}^2 + a_1 \| \ell u \|_{L^2(0,b)}^2 + c_1 \| u \|_{L^2(Q^\tau)}^2. \end{aligned}$$

Applying Lemma 7.1 from [3] to the above inequality we get

$$\begin{aligned} & \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q^\tau)}^2 + \left\| \frac{\partial^{2m} u}{\partial x^{2m}} \right\|_{B_2^{m,0}(Q^\tau)}^2 + \| u(x, \tau) \|_{L^2(0,b)}^2 \\ & \leq c_2 \left( \| \mathcal{L}u \|_{B_2^{m,0}(Q)}^2 + \| \ell u \|_{L^2(0,b)}^2 \right), \end{aligned}$$

where

$$c_2 := \frac{\max(3/2, a_1)}{\min(1/2, 1/4 a_0^2, a_0)} \exp(c_1 T).$$

Since the right-hand side of the above inequality does not depend on  $\tau$ , we can take the least upper bound of the left side with respect to  $\tau$  from 0 to  $T$ . Thus, inequality (3.4) holds, where  $c := c_2^{1/2}$ . □

### 4. Solvability of the Problem

From inequality (3.1), it follows that operator  $L: E \rightarrow F$  is continuous, while from inequality (3.4) it follows that the range of operator  $L$  is closed in  $F$  and, therefore, there is the continuous inverse operator  $L^{-1}$  yielding the solution. In other words, this means that operator  $L$  is a linear homeomorphism from the space  $E$  on the closed set  $R(L) \subset F$ . To prove that problem (1.1)-(1.3) has a unique solution, it remains to show that  $R(L) = F$ .

**Theorem 3:** *Let the conditions of Theorem 2 hold, and let the coefficient  $a(t)$  have bounded derivatives up to the second order. Then, for any functions  $f \in B_2^{m,0}(Q)$  and  $\varphi \in L^2(0,b)$ , there is a unique solution  $u = L^{-1} \mathcal{F}$  of problem (1.1)-(1.3), where  $\mathcal{F} = (f, \varphi)$ , and*

$$\| u \|_E \leq c \left( \| f \|_{B_2^{m,0}(Q)} + \| \varphi \|_{L^2(0,b)} \right),$$

where constant  $c$  is independent of  $u$ .

**Proof:** To prove Theorem 3, we need the following proposition.

**Proposition:** *Let  $D_0(L) = \{u/u \in D(L), \ell u = 0\}$  and let the conditions of Theorem 3 hold. If for  $v \in B_2^{m,0}(Q)$  and for all  $u \in D_0(L)$ ,*

$$(\mathcal{L}u, v)_{B_2^{m,0}(Q)} = 0, \tag{4.1}$$

then  $v$  vanishes almost everywhere on  $Q$ .

**Proof of the Proposition:** Assume that relation (4.1) holds for any function  $u \in D_0(L)$ . Using this fact we can express (4.1) in a special form. First define  $h$  by the formula

$$h := - \int_t^T \frac{\partial}{\partial \tau} \left( a(\tau) \frac{\partial u}{\partial \tau} \right) d\tau.$$

Let  $\frac{\partial u}{\partial t}$  be a solution of

$$a(t) \frac{\partial u}{\partial t} = h \tag{4.2}$$

and let

$$D_s(L) := \{u/u \in D(L) : u = 0 \text{ for } t \leq s\}. \tag{4.3}$$

We, now, have

$$v = - \frac{\partial}{\partial t} \left( a(t) \frac{\partial u}{\partial t} \right). \tag{4.4}$$

Relations (4.2) and (4.3) imply that  $u$  is in  $D_0(L)$ . It possesses, in fact, a higher order of smoothness, and we have the following result:

**Lemma 2:** *If the conditions of the proposition are met, then the function  $u$  defined by (4.2) and (4.3) has derivatives with respect to  $t$  up to the second order belonging to the space  $B_2^{m,0}(Q_s)$ , where  $Q_s = (0, b) \times (s, T)$ .*

**Proof of Lemma 2:** To prove Lemma 2, we will use the following  $t$ -averaging operators: Let  $\omega \in C^\infty(\mathbb{R})$ ,  $\omega \geq 0$ ;  $\omega = 0$  in a neighborhood of  $t = 0$  and  $t = T$ , and outside the interval  $(0, T)$ , and let  $\int_{\mathbb{R}} \omega(t) dt = 1$ . We consider the operators  $\rho_\epsilon$  defined by the formula

$$(\rho_\epsilon w)(x, t) = \frac{1}{\epsilon} \int_0^T \omega\left(\frac{s-t}{\epsilon}\right) w(x, s) ds \text{ for } w \in B_2^{m,0}(Q).$$

The above operators have the following properties:

**P1:** The function  $\rho_\epsilon w \in C^\infty(Q)$  and it vanishes in a neighborhood of  $t = T$  if  $w \in B_2^{m,0}(Q)$ , and  $\rho_\epsilon u \in D_s(L)$  if  $u \in D_s(L)$ .

**P2:** If  $w \in B_2^{m,0}(Q)$ , then  $\|\rho_\epsilon w - w\|_{B_2^{m,0}(Q)} \rightarrow 0$  when  $\epsilon \rightarrow 0$ , and  $\|\rho_\epsilon w\|_{B_2^{m,0}(Q)} \leq \|w\|_{B_2^{m,0}(Q)}$ .

**P3:**  $\frac{d^k}{dt^k} \rho_\epsilon u = \rho_\epsilon \frac{d^k u}{dt^k}$  for  $k = 1, 2$  if  $u \in D_s(L)$ .

**P4:** If  $w \in B_2^{m,0}(Q)$  then,

$$\left\| \frac{\partial}{\partial t} (a(t) \rho_\epsilon w - \rho_\epsilon a(t) w) \right\|_{B_2^{m,0}(Q)} \rightarrow 0, \text{ when } \epsilon \rightarrow 0.$$

Proofs of properties P1-P4 are similar to the proofs of the corresponding properties obtained in [3] (see Lemma 9.1). □

Applying the operators  $\rho_\epsilon$  and  $\frac{\partial}{\partial t}$  to equation (4.2), we obtain

$$a(t) \frac{\partial}{\partial t} \rho_\epsilon \frac{\partial u}{\partial t} = \frac{\partial}{\partial t} \left( a(t) \rho_\epsilon \frac{\partial u}{\partial t} - \rho_\epsilon a(t) \frac{\partial u}{\partial t} \right) - a'(t) \rho_\epsilon \frac{\partial u}{\partial t} + \frac{\partial}{\partial t} \rho_\epsilon h.$$

It follows that

$$\begin{aligned} & \left\| a(t) \frac{\partial}{\partial t} \rho_\epsilon \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q)}^2 \leq c_3 \left( \left\| \rho_\epsilon \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q)}^2 \right. \\ & \left. + \left\| \frac{\partial}{\partial t} \rho_\epsilon h \right\|_{B_2^{m,0}(Q)}^2 + \left\| \frac{\partial}{\partial t} \left( a(t) \rho_\epsilon \frac{\partial u}{\partial t} - \rho_\epsilon a(t) \frac{\partial u}{\partial t} \right) \right\|_{B_2^{m,0}(Q)}^2 \right), \end{aligned}$$

where  $c_3 = \max(3c_1, 3)$ .

By virtue of properties *P1-P4* of the  $t$ -averaging operators and by inequality (2.3), we have

$$\left( \left\| \frac{\partial^2 u}{\partial t^2} \right\|_{B_2^{m,0}(Q)}^2 \leq c_4 \left\| \frac{\partial u}{\partial t} \right\|_{L^2(Q)}^2 + \left\| \frac{\partial}{\partial t} \rho_\epsilon h \right\|_{B_2^{m,0}(Q)}^2 \right),$$

where  $c_4 = \max(c_3 b^{2m}/(a_0^2 2^m), 1/a_0^2)$ . This yields the proof of Lemma 2.  $\square$

Now, we will prove the proposition. Replace  $v$  in (4.1) by its representation (4.4). We have

$$\begin{aligned} & -2 \operatorname{Re} \left( \frac{\partial u}{\partial t}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)} \\ & -2 \operatorname{Re} \left( (-1)^m a(t) \frac{\partial^{2m} u}{\partial x^{2m}}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)} = 0. \end{aligned} \quad (4.5)$$

We write the remaining two terms of (4.5) in the form:

$$\begin{aligned} & -2 \operatorname{Re} \left( \frac{\partial u}{\partial t}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)} \\ & = \left\| a^{1/2}(s) \mathfrak{F}^m \frac{\partial u(x,s)}{\partial t} \right\|_{L^2(0,b)}^2 - \left\| a^{1/2}(t) \mathfrak{F}^m \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2, \end{aligned} \quad (4.6)$$

$$\begin{aligned} & -2 \operatorname{Re} \left( (-1)^m a(t) \frac{\partial^{2m} u}{\partial x^{2m}}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)} \\ & = 2 \left\| a(t) \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2 + \operatorname{Re} \left( a'(T) u(x, T), a(T) \bar{u}(x, T) \right)_{L^2(0,b)} \\ & - \left\| a'(t) u \right\|_{L^2(Q_s)}^2 - \operatorname{Re} \left( a''(t) u, a(t) \bar{u} \right)_{L^2(Q_s)}. \end{aligned} \quad (4.7)$$

Elementary calculations, starting from (4.6) and (4.7), yield the inequalities

$$\begin{aligned} & a_0 \left\| \frac{\partial u(x,s)}{\partial t} \right\|_{B_2^{m,0}(b)}^2 \leq c_1 \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m,0}(Q_s)}^2 - 2 \operatorname{Re} \left( \frac{\partial u}{\partial t}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)}, \\ & 2a_0^2 \left\| \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2 + a_0 c_0 \left\| u(x, T) \right\|_{L^2(0,b)}^2 \\ & \leq -2 \operatorname{Re} \left( (-1)^m a(t) \frac{\partial^{2m} u}{\partial x^{2m}}, \frac{\partial}{\partial t} \left( a(t) \frac{\partial \bar{u}}{\partial t} \right) \right)_{B_2^{m,0}(Q_s)} + (1/2a_1^2 + c_1^2 + 1/2c_5^2) \left\| u \right\|_{L^2(Q_s)}^2, \end{aligned}$$

where  $c_5 = \sup_{0 \leq t \leq T} |a''(t)|$ .

Consequently,

$$\begin{aligned} & \left\| \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2 + \|u(x, T)\|_{L^2(0, b)}^2 + \left\| \frac{\partial u(x, s)}{\partial t} \right\|_{B_2^m(0, b)}^2 \\ & \leq c_6 \left( \|u\|_{L^2(Q_s)}^2 + \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m, 0}(Q_s)}^2 \right), \end{aligned} \tag{4.8}$$

where  $c_6 := \max(c_1, 1/2a_1^2 + c_1^2 + 1/2c_5^2)/\min(a_0, 2a_0^2, a_0c_0)$ .

Inequality (4.8) is the basic of our proof. To use (4.8), we note that constant  $c_6$  is independent of  $s$ . However, function  $u$  in (4.8) depends on  $s$ . To avoid this difficulty we introduce a new function  $\theta$  by the formula

$$\theta(x, t) := \int_t^T \frac{\partial u}{\partial \tau} d\tau.$$

Then,  $u(x, t) = \theta(x, s) - \theta(x, t)$ ,  $u(x, T) = \theta(x, s)$ , and we have

$$\|u\|_{L^2(Q_s)}^2 \leq 2 \left( \|\theta(x, t)\|_{L^2(Q_s)}^2 + (T - s) \|\theta(x, s)\|_{L^2(0, b)}^2 \right).$$

Hence, if  $s_0 > 0$  satisfies  $0 < 2c_6(T - s_0) \leq 1/2$ , then (4.8) implies that

$$\begin{aligned} & \left\| \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2 + \left\| \frac{\partial u(x, s)}{\partial t} \right\|_{B_2^m(0, b)}^2 + \|\theta(x, s)\|_{L^2(0, b)}^2 \\ & \leq 4c_6 \left( \left\| \frac{\partial u}{\partial t} \right\|_{B_2^{m, 0}(Q_s)}^2 + \|\theta(x, t)\|_{L^2(Q_s)}^2 \right), \end{aligned} \tag{4.9}$$

for all  $s \in [T - s_0, T]$ .

We denote the sum of the two terms on the right of (4.9) by  $\beta(s)$ . Hence, we obtain

$$\left\| \frac{\partial u}{\partial t} \right\|_{L^2(Q_s)}^2 - \frac{d\beta(s)}{ds} \leq 4c_6\beta(s),$$

and, consequently,

$$- \frac{d}{ds}(\beta(s) \exp(4c_6s)) \leq 0. \tag{4.10}$$

Integrating (4.10) over  $(s, T)$  and taking into account that  $\beta(T) = 0$ , we obtain

$$\beta(s) \exp(4c_6s) \leq 0. \tag{4.11}$$

It follows from (4.11), that  $v = 0$  almost everywhere on  $Q_{T-s_0}$ . Proceeding this way step by step along the rectangle with side  $s_0$ , we prove that  $v = 0$  almost everywhere on  $Q$ . This completes the proof of the proposition.  $\square$

Now, we will prove Theorem 3. For this purpose it is sufficient to prove that the range  $R(L)$  of  $L$  is dense in  $F$ .

Suppose that, for some  $V = (v, v_0) \in {}^\perp R(L)$ ,

$$(\mathcal{L}u, v)_{B_2^{m, 0}(Q)} + (\ell u, v_0)_{L^2(0, b)} = 0. \tag{4.12}$$

We must prove that  $V = 0$ . Putting  $u \in D_0(L)$  into (4.12) we obtain

$$(\mathcal{L}u, v)_{B_2^{m, 0}(Q)} = 0, \quad u \in D(L).$$

Hence, the proposition implies that  $v = 0$ . Thus, (4.12) takes the form


$$(\ell u, v_0)_{L^2(0,b)} = 0, \quad u \in D(L).$$

Since the range of operator  $\ell$  is everywhere dense in  $L^2(0,b)$ , the above relation implies that  $v_0 = 0$ . Hence,  $V = 0$ . This proves Theorem 3.  $\square$

## References

- [1] Benouar, N.E. and Yurchuk, Mixed problem with an integral condition for parabolic equations with an integral condition for parabolic equations with the Bessel operator, *Differents. Uravn.* **27**:12 (1991), 2094-2098.
- [2] Cannon, J.R., The solution of the heat equation subject to the specification of energy, *Quart. Appl. Math.* **21**:2 (1963), 155-160.
- [3] Gårding, L., *Cauchy's Problem for Hyperbolic Equations*, University of Chicago 1957.
- [4] Ionkin, N.I., Solution of boundary value problems in heat conduction theory with nonlocal boundary conditions, *Differents. Uravn.* **13**:2 (1977), 294-304.
- [5] Ionkin, N.I., Stability of a problem in heat conduction theory with nonlocal boundary conditions, *Differents. Uravn.* **15**:7 (1979), 1279-1283.
- [6] Ionkin, N.I. and Moiseev, E.I., A problem for the heat conduction equation with two-point boundary condition, *Differents. Uravn.* **15**:7 (1979), 1284-1295.
- [7] Kamynin, N.I., A boundary value problem in the theory of the heat conduction with non-classical boundary condition, *Th., Vychisl., Mat., Fiz.* **4**:6 (1964), 1006-1024.
- [8] Kartynnik, A.V., Three point boundary value problem with an integral space variables conditions for second order parabolic equations, *Differents. Uravn.* **26** (1990), 1568-1575.
- [9] Samarskii, A.A., Some problems in differential equations theory, *Differents. Uravn.* **16**:11 (1980), 1925-1935.
- [10] Yurchuk, N.I., Mixed problem with an integral condition for certain parabolic equations, *Differents. Uravn.* **22**:12 (1986), 2117-2126.





# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

