

MÖBIUS METRIC IN SECTOR DOMAINS

OONA RAINIO AND MATTI VUORINEN

ABSTRACT. The Möbius metric δ_G is studied in the cases where its domain G is an open sector of the complex plane. We introduce upper and lower bounds for this metric in terms of the hyperbolic metric and the angle of the sector, and then use these results to find bounds for the distortion of the Möbius metric under quasiregular mappings defined in sector domains. Furthermore, we numerically study the Möbius metric and its connection to the hyperbolic metric in polygon domains.

Author information.

Oona Rainio⁽¹⁾ email: ormrai@utu.fi ORCID: 0000-0002-7775-7656

Matti Vuorinen⁽¹⁾ email: vuorinen@utu.fi ORCID: 0000-0002-1734-8228

(1): Department of Mathematics and Statistics, University of Turku, FI-20014 Turku, Finland

Funding. Oona Rainio's research was funded by the University of Turku Graduate School UTUGS.

Data availability statement. Not applicable, no new data was generated.

Conflict of interest statement. No conflict of interest.

1. INTRODUCTION

One of the most important concepts in the geometric function theory is the *intrinsic distance*. It means that, given two points in a domain, we do not only consider how close these points are to each other but also how they are located with respect to the boundary of the domain. In order to measure these kinds of distances, we need to use suitable intrinsic or hyperbolic type metrics, which have been recently studied, for instance, in [1, 5, 6, 7, 9, 12, 13, 14].

In this article, we focus on one of these intrinsic metrics, which is defined as follows: For any domain $G \subset \overline{\mathbb{R}}^n = \mathbb{R}^n \cup \{\infty\}$ whose complement $(\overline{\mathbb{R}}^n \setminus G)$ contains at least two points, let the *Möbius metric* be the function $\delta_G : G \times G \rightarrow [0, \infty)$,

$$(1.1) \quad \delta_G(x, y) = \sup_{a, b \in \partial G} \log(1 + |a, x, b, y|),$$

where $|a, x, b, y|$ is the cross-ratio defined in (2.1). This metric was first introduced in [17, pp. 115-116] and then later studied more extensively by P. Seittenranta in his PhD

File: `seit20210514.tex`, printed: 2023-1-10, 4.11

2010 *Mathematics Subject Classification*. Primary 51M10; Secondary 30C62.

Key words and phrases. Hyperbolic geometry, hyperbolic metric, intrinsic geometry, Möbius metric, quasiregular mappings, triangular ratio metric.

thesis [15, Def. 1.1, p. 511], which is why it is sometimes also referred to as Seittenranta's metric.

Due to the Möbius invariance of the cross-ratio, the distances defined with the Möbius metric are preserved under Möbius transformations, which is one of the most useful properties of this metric. However, there are still numerous open questions concerning this metric. For instance, while it is known that the value of this metric is equal to that of the hyperbolic metric ρ or the distance ratio metric j in some special cases, see Theorems 2.3 and 3.1, the Möbius metric is studied very little in other kinds of domains. To fill this gap, our aim here is to find more information about the Möbius metric in the cases where the domain G is either an open sector of the complex plane or a polygon.

The main result of this article is as follows:

Theorem 1.2. *For all points x, y in an open sector S_θ with an angle $0 < \theta < 2\pi$, the following inequalities hold:*

$$(1) \quad \rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y) \leq \min \left\{ 2, \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2 \right\} \rho_{S_\theta}(x, y), \quad \text{if } \theta < \pi,$$

$$(2) \quad \delta_{S_\theta}(x, y) = \rho_{S_\theta}(x, y), \quad \text{if } \theta = \pi,$$

$$(3) \quad \max \left\{ 2 \operatorname{arth} \frac{\operatorname{th}(\rho_{S_\theta}(x, y)/2)}{2}, \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2 \rho_{S_\theta}(x, y) \right\} \leq \delta_{S_\theta}(x, y) \leq 4\psi, \quad \text{if } \theta > \pi,$$

where

$$\psi = \begin{cases} \min \{ \rho_{S_\theta}(x, y), \operatorname{arth}((\theta/\pi)\operatorname{th}(\rho_{S_\theta}(x, y)/2)) \}, & \text{if } (\theta/\pi)\operatorname{th}(\rho_{S_\theta}(x, y)/2) < 1, \\ \rho_{S_\theta}(x, y) & \text{otherwise.} \end{cases}$$

The structure of this article is as follows. First, in Section 3, we combine some already known inequalities to create some initial bounds for the Möbius metric in a general domain. Then, in Section 4, we study the Möbius metric defined in an open sector by showing how the supremum of the cross-ratio in its definition (1.1) can be found. These results will be used in Section 5, where we introduce bounds for the Möbius metric in terms of the hyperbolic metric in a sector and prove Theorem 1.2. In Section 6, we apply these results and prove bounds for the distortion of the Möbius metric under quasiregular mappings of the unit disk into sector domains. Finally, in Section 7, we utilise the recent computational methods from [11] to experimentally study the inequalities between the Möbius and hyperbolic metric in polygon domains and formulate a few conjectures.

2. PRELIMINARIES

First, define the following notations for the Euclidean metric. Let the distance from a point $x \in \mathbb{R}^n$ to a non-empty set $F \subset \mathbb{R}^n$ be $d(x, F) = \inf\{|x - z| \mid z \in F\}$. For a domain $G \subsetneq \mathbb{R}^n$, denote $d_G(x) = d(x, \partial G)$ for all $x \in G$. Let the Euclidean diameter of a non-empty set F be $d(F)$ and the Euclidean distance between two non-empty separate sets

F_0, F_1 be $d(F_0, F_1)$. Furthermore, denote the Euclidean open ball with a center $x \in \mathbb{R}^n$ and a radius $r > 0$ by $B^n(x, r)$, the corresponding closed ball by $\overline{B}^n(x, r)$ and its boundary sphere by $S^{n-1}(x, r)$.

Let $\overline{\mathbb{R}}^n = \mathbb{R}^n \cup \{\infty\}$ be as in the introduction, and also denote $\overline{\mathbb{C}}^n = \mathbb{C}^n \cup \{\infty\}$. For all distinct points $x, y \in \overline{\mathbb{R}}^n$, define the *spherical (chordal) metric* as in [5, (3.6), p. 29]:

$$q(x, y) = \frac{|x - y|}{\sqrt{1 + |x|^2}\sqrt{1 + |y|^2}}, \quad \text{if } x \neq \infty \neq y; \quad q(x, \infty) = \frac{1}{\sqrt{1 + |x|^2}}.$$

For any four distinct points $a, b, c, d \in \overline{\mathbb{R}}^n$, define the *cross-ratio* as [5, (3.10), p. 33]

$$(2.1) \quad |a, b, c, d| = \frac{q(a, c)q(b, d)}{q(a, b)q(c, d)},$$

and note that, if $\infty \notin \{a, b, c, d\}$, then this definition yields

$$|a, b, c, d| = \frac{|a - c||b - d|}{|a - b||c - d|}.$$

Other than the Möbius metric, we will be needing a few other hyperbolic type metrics. Define the upper half-space $\mathbb{H}^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n > 0\}$, the unit ball $\mathbb{B}^n = B^n(0, 1)$ and the open sector $S_\theta = \{x \in \mathbb{C} \setminus \{0\} \mid 0 < \arg(x) < \theta\}$ with an angle $\theta \in (0, 2\pi)$. Here, $\arg(x) \in [0, 2\pi)$ denotes the principal branch of the argument of a complex number $x \in \mathbb{C} \setminus \{0\}$. Now, we can define the *hyperbolic metric* in these three domains by using the following formulas, respectively [5, (4.8), p. 52 & (4.14), p. 55]:

$$\begin{aligned} \text{ch}\rho_{\mathbb{H}^n}(x, y) &= 1 + \frac{|x - y|^2}{2d_{\mathbb{H}^n}(x)d_{\mathbb{H}^n}(y)}, \quad x, y \in \mathbb{H}^n, \\ \text{sh}^2 \frac{\rho_{\mathbb{B}^n}(x, y)}{2} &= \frac{|x - y|^2}{(1 - |x|^2)(1 - |y|^2)}, \quad x, y \in \mathbb{B}^n, \\ \rho_{S_\theta}(x, y) &= \rho_{\mathbb{H}^2}(x^{\pi/\theta}, y^{\pi/\theta}), \quad x, y \in S_\theta, \end{aligned}$$

From these formulas, it follows that:

$$(2.2) \quad \text{th} \frac{\rho_{\mathbb{H}^2}(x, y)}{2} = \left| \frac{x - y}{x - \bar{y}} \right|, \quad \text{th} \frac{\rho_{\mathbb{B}^2}(x, y)}{2} = \left| \frac{x - y}{1 - x\bar{y}} \right|,$$

where \bar{y} is the complex conjugate of y .

For a domain $G \subsetneq \mathbb{R}^n$, the *distance ratio metric* [17, p. 25] introduced by Gehring and Osgood [3] is the function $j_G : G \times G \rightarrow [0, \infty)$,

$$j_G(x, y) = \log\left(1 + \frac{|x - y|}{\min\{d_G(x), d_G(y)\}}\right).$$

As noted in [7, 2.2, p. 1123 & Lemma 2.1, p. 1124], this metric can be used to define another metric, the so called j_G^* -metric $j_G^* : G \times G \rightarrow [0, 1]$,

$$j_G^*(x, y) = \operatorname{th} \frac{j_G(x, y)}{2} = \frac{|x - y|}{|x - y| + 2 \min\{d_G(x), d_G(y)\}}.$$

Furthermore, the *quasihyperbolic metric* introduced by Gehring and Palka in [4] is defined as the function $k_G : G \times G \rightarrow [0, \infty)$

$$k_G(x, y) = \inf_{\gamma \in \Gamma_{xy}} \int_{\gamma} \frac{|dx|}{d_G(x)},$$

where Γ_{xy} consists of all the rectifiable curves in G joining x and y . Consider yet the *triangular ratio metric* [1, (1.1), p. 683] $s_G : G \times G \rightarrow [0, 1]$,

$$s_G(x, y) = \frac{|x - y|}{\inf_{z \in \partial G} (|x - z| + |z - y|)},$$

which was originally introduced by P. Hästö in 2002 [8].

The following result expresses the main property of the Möbius metric:

Theorem 2.3. [15], [5, Thm 5.16, p. 75] *The Möbius metric δ_G is Möbius invariant: If $G \subset \mathbb{R}^n$ is a domain such that $\mathbb{R}^n \setminus G$ contains at least two points and $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Möbius transformation, then for all $x, y \in G$,*

$$\delta_{h(G)}(h(x), h(y)) = \delta_G(x, y).$$

Furthermore, $\delta_G = \rho_G$ for $G \in \{\mathbb{B}^n, \mathbb{H}^n\}$.

3. GENERAL INEQUALITIES

In this section, we will briefly review a few already existing inequalities and show how they can be used to create bounds for the Möbius metric. Note that the inequalities found here concern mostly the situation, in which the shape of the domain G is not known. For instance, Corollary 3.5 gives us an inequality for a simply connected uniform domain G , but its constants are probably not very sharp when compared to those that could be obtained when knowing the exact shape of the domain G .

Theorem 3.1. [5, Thm 5.16, p. 75] *For all points x, y in a domain $G \subsetneq \mathbb{R}^n$,*

$$j_G(x, y) \leq \delta_G(x, y) \leq 2j_G(x, y)$$

and, in the special case $G = \mathbb{R}^n \setminus \{0\}$, $\delta_G = j_G$.

Theorem 3.2. [2, (3.2.1), p. 35] *For all points x, y in a simply connected domain $G \subsetneq \mathbb{R}^2$,*

$$\frac{1}{2}k_G(x, y) \leq \rho_G(x, y) \leq 2k_G(x, y).$$

Theorem 3.3. [5, Cor. 5.6, p. 69] *For all points x, y in a domain $G \subsetneq \mathbb{R}^n$, $j_G(x, y) \leq k_G(x, y)$.*

Definition 3.4. [5, Def. 6.1, p. 84],[10, Def. 2.4, p. 8] A domain $G \subset \mathbb{R}^n$ is *uniform* if there exists a number $A \geq 1$ such that the inequality $k_G(x, y) \leq Aj_G(x, y)$ holds for all $x, y \in G$ and the smallest such number A fulfilling this condition is called the *uniformity constant* of G .

Corollary 3.5. *If a domain $G \subsetneq \mathbb{R}^2$ is simply connected and uniform with the uniformity constant A_G , then*

$$\rho_G(x, y)/(2A_G) \leq \delta_G(x, y) \leq 4\rho_G(x, y)$$

for all $x, y \in G$.

Proof. Follows from Theorems 3.3 and 3.2, and Definition 3.4. \square

Now, let us find some bounds for the Möbius metric with the triangular ratio metric and the j^* -metric in the cases of both a convex domain G and a non-convex one.

Lemma 3.6. [7, Lemma 2.1, p. 1124; Lemma 2.2, p. 1125 & Thm 2.9(i), p. 1129] *For all points x, y in a domain $G \subsetneq \mathbb{R}^n$,*

$$j_G^*(x, y) \leq s_G(x, y) \leq 2j_G^*(x, y)$$

and, if G is convex, the constant 2 above can be replaced by $\sqrt{2}$.

Lemma 3.7. [7, Lemma 2.7(ii), p. 1128] *For all points x, y in a convex domain $G \subsetneq \mathbb{R}^n$,*

$$\operatorname{th} \frac{j_G(x, y)}{2} \leq s_G(x, y) \leq \operatorname{th} j_G(x, y).$$

Corollary 3.8. *For a domain $G \subsetneq \mathbb{R}^n$ such that $\overline{\mathbb{R}^n} \setminus G$ contains at least two points and for all $x, y \in G$, the following inequalities hold:*

- (1) $j_G^*(x, y) \leq \operatorname{th}(\delta_G(x, y)/2) \leq \operatorname{th}(2\operatorname{arth}(j_G^*(x, y))) \leq 2j_G^*(x, y)$,
- (2) $s_G(x, y)/2 \leq \operatorname{th}(\delta_G(x, y)/2) \leq \operatorname{th}(2\operatorname{arth}(s_G(x, y))) \leq 2s_G(x, y)$.

Furthermore, if G is convex, then for all $x, y \in G$

- (3) $s_G(x, y)/\sqrt{2} \leq \operatorname{th}(\delta_G(x, y)/2)$,
- (4) $s_G(x, y) \leq \operatorname{th}(\delta_G(x, y))$.

Proof. (1) Follows from Theorem 3.1 and the definition of j^* -metric.

(2) Follows from the first inequality and Lemma 3.6.

(3) Follows from the first inequality and Lemma 3.6.

(4) Follows from Theorem 3.1 and Lemma 3.7. \square

Finally, let us consider the case where the domain G is an open sector. Note that neither the inequalities of Corollary 3.10 nor Corollary 3.12 have the best possible constants. In fact, they are only used to prove our main result, Theorem 1.2, in Section 5.

Theorem 3.9. [14, Cor. 4.9, p. 9] *For a fixed angle $\theta \in (0, 2\pi)$ and for all $x, y \in S_\theta$, the following results hold:*

- (1) $s_{S_\theta}(x, y) \leq \operatorname{th}(\rho_{S_\theta}(x, y)/2) \leq (\pi/\theta) \sin(\theta/2) s_{S_\theta}(x, y)$ if $\theta \in (0, \pi)$,
- (2) $s_{S_\theta}(x, y) = \operatorname{th}(\rho_{S_\theta}(x, y)/2)$ if $\theta = \pi$,

(3) $(\pi/\theta)s_{S_\theta}(x, y) \leq \text{th}(\rho_{S_\theta}(x, y)/2) \leq s_{S_\theta}(x, y)$ if $\theta \in (\pi, 2\pi)$.
Furthermore, these bounds are also sharp.

Corollary 3.10. *For all points $x, y \in S_\theta$, the following inequalities hold:*

- (1) $\max \left\{ 2\text{arth} \left(\frac{\theta}{\sqrt{2}\pi \sin(\theta/2)} \text{th} \frac{\rho_{S_\theta}(x, y)}{2} \right), \text{arth} \left(\frac{\theta}{\pi \sin(\theta/2)} \text{th} \frac{\rho_{S_\theta}(x, y)}{2} \right) \right\}$
 $\leq \delta_{S_\theta}(x, y) \leq 2\rho_{S_\theta}(x, y), \quad \text{if } 0 < \theta < \pi,$
- (2) $2\text{arth} \frac{\text{th}(\rho_{S_\theta}(x, y)/2)}{2} \leq \delta_{S_\theta}(x, y), \quad \text{if } \pi < \theta < 2\pi,$
- (3) $\delta_{S_\theta}(x, y) \leq 4\text{arth} \left(\frac{\theta}{\pi} \text{th} \frac{\rho_{S_\theta}(x, y)}{2} \right), \quad \text{if } \pi < \theta < 2\pi \quad \text{and} \quad \frac{\theta}{\pi} \text{th} \frac{\rho_{S_\theta}(x, y)}{2} < 1.$

Proof. Follows from Corollary 3.8 and Theorem 3.9. □

Theorem 3.11. [10, Thm 1.7 & Thm 1.8, p. 6] *An open sector S_θ is uniform with the constant A_θ that fulfills*

$$A_\theta = \frac{1}{\sin(\theta/2)} + 1, \quad \text{if } 0 < \theta \leq \pi, \quad \text{and}$$

$$\max \left\{ 2, \frac{2 \log(\tan(\theta/4)) + \theta - \pi}{\log(1 - 2 \cos(\theta/2))} \right\} \leq A_\theta \leq 4 \left(\frac{\theta}{2\pi - \theta} \right)^2 \left(\frac{1}{\sin(\theta/2)} + 1 \right),$$

if $\pi < \theta < 2\pi$.

Corollary 3.12. *For all points $x, y \in S_\theta$,*

- (1) $\frac{\sin(\theta/2)}{2(1 + \sin(\theta/2))} \rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y) \leq 4\rho_{S_\theta}(x, y), \quad \text{if } 0 < \theta \leq \pi,$
- (2) $\frac{1}{8} \left(\frac{2\pi}{\theta} - 1 \right)^2 \frac{\sin(\theta/2)}{1 + \sin(\theta/2)} \rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y) \leq 4\rho_{S_\theta}(x, y) \quad \text{if } \pi < \theta < 2\pi.$

Proof. Follows from Corollary 3.5 and Theorem 3.11. □

4. MÖBIUS METRIC IN OPEN SECTOR

In this section, our aim is to find ways to estimate the value of the Möbius metric defined in an open sector S_θ . To do this, we will study the supremum of the cross-ratio needed in the definition of the metric δ_{S_θ} , in both cases where the angle θ is less than π and greater than π . The main result of this section is Corollary 4.13 but, in order to prove it, we need to consider several other results first.

Proposition 4.1. (1) *If $x, y \in \mathbb{H}^2$ such that $|x| = |y| = r > 0$ and $\arg(x) \leq \arg(y)$, then $\sup_{a, b \in \mathbb{R}} |a, x, b, y| = |r, x, -r, y|$.*

(2) *If $x, y \in i\mathbb{R} \cap \mathbb{B}^2$ such that $\text{Im}(x) \leq \text{Im}(y)$, then $\sup_{a, b \in S^1} |a, x, b, y| = |-i, x, i, y|$.*

Proof. Since $\delta_{\mathbb{H}^2}(x, y) = \sup_{a, b \in \mathbb{R}} \log(1 + |a, x, b, y|)$, both results can be verified by the fact that $\delta_G = \rho_G$ for $G \in \{\mathbb{B}^n, \mathbb{H}^n\}$ according to Theorem 2.3. \square

Lemma 4.2. *For all points x, y in an open sector S_θ with an angle $0 < \theta < \pi$ such that $\arg(x) \leq \arg(y)$ and $|x| = |y| = r > 0$, there is a Möbius transformation f that maps S_θ onto the lens-shaped domain*

$$B^2((1-u)i, u) \cap B^2((u-1)i, u), \quad u = \frac{1}{1 - \cos(\theta/2)} > 1,$$

and x, y into $f(x), f(y) \in i\mathbb{R} \cap \mathbb{B}^2$ so that $\text{Im}(f(x)) \leq \text{Im}(f(y))$.

Proof. Define the function $f : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$,

$$(4.3) \quad f(z) = \frac{-i(1 + e^{\theta i/2})(z - re^{\theta i/2})}{(1 - e^{\theta i/2})(z + re^{\theta i/2})}.$$

Clearly, f is the Möbius transformation that fulfills

$$(4.4) \quad f(r) = -i, \quad f(re^{\theta i/2}) = 0, \quad f(re^{\theta i}) = i, \quad f(0) = \frac{-\sin(\theta/2)}{1 - \cos(\theta/2)} = -f(\infty).$$

By the general properties of Möbius transformations, f preserves the angles and must turn a line into a circle if any three points chosen from it are no longer collinear after the transformation. Thus, f maps two sides of the sector S_θ onto two circular arcs that are symmetric with respect to the both coordinate axes, meet each other in the points $f(0)$ and $f(\infty)$ at an angle of θ , and out of which one contains the point i and the other one the point $-i$. See Figure 1.

Consider a circle $S^1((1-u)i, u)$, $u > 1$. Clearly, it is symmetric with respect to the coordinate axes and $i \in S^1((1-u)i, u)$. Using simple trigonometry, it can be calculated that the two interior angles of the figure consisting of the real axis and the circular arc $S^1((1-u)i, u) \cap \mathbb{H}^2$ are

$$\frac{\pi}{2} - \arcsin\left(\frac{u-1}{u}\right) \in \left(0, \frac{\pi}{2}\right).$$

If the value of this angle is $\theta/2$, we can solve that

$$(4.5) \quad u = \frac{1}{1 - \cos(\theta/2)}.$$

By combining all our observations made above, we will have that, for all $0 < \theta < \pi$,

$$\begin{aligned} f(\partial S_\theta) &= (S^1((1-u)i, u) \cap \mathbb{H}^2) \cup (S^1((u-1)i, u) \setminus \mathbb{H}^2), \\ f(S_\theta) &= B^2((1-u)i, u) \cap B^2((u-1)i, u), \end{aligned}$$

where u is as in (4.5). The final part of the lemma is very trivial: From the behaviour of the points in (4.4), we see that the transformation f maps the circle $S^1(0, r)$ onto the imaginary axis and, if $x, y \in S^1(0, r)$ such that $0 < \arg(x) \leq \arg(y) < \theta$, then clearly $f(x), f(y) \in [-i, i]$ so that $-1 < \text{Im}(f(x)) \leq \text{Im}(f(y)) < 1$. \square




FIGURE 1. Sector S_θ before and after the Möbius transformation f defined in (4.3), when $\theta = 3\pi/4$.

The result of Lemma 4.2 is very useful because it follows from the Möbius invariance of the cross-ratio that the value of the Möbius metric between $x, y \in S_\theta$ can be calculated in the lens-shaped symmetric domain $f(S_\theta)$ for $f(x), f(y)$, see Figure 1.

Theorem 4.6. *For all $0 < \theta < \pi$ and $x, y \in S_\theta$ such that $\arg(x) \leq \arg(y)$ and $|x| = |y| = r > 0$, the supremum $\sup_{a, b \in \partial S_\theta} |a, x, b, y|$ is given by the points $a = r$ and $b = re^{\theta i}$.*

Proof. Let f be the Möbius transformation defined in (4.3), under which the open sector S_θ with an angle $0 < \theta < \pi$ is mapped onto a lens-shaped domain $f(S_\theta)$. For all points $x, y \in i\mathbb{R} \cap \mathbb{B}^2$, choose $a, b \in f(\partial S_\theta)$ so that the cross-ratio $|a, x, b, y|$ is at greatest. Note that, for all points $u \in i\mathbb{R} \cap \mathbb{B}^2$ and $v \in \mathbb{C} \setminus \mathbb{B}^2$, the inequality

$$(4.7) \quad \max \left\{ \frac{|i - v|}{|u - i|}, \frac{|-i - v|}{|u - (-i)|} \right\} \geq 1,$$

holds. It follows from this that $|x - a| \leq |a - b|$ holds, because otherwise replacing a either by i or $-i$ would give a greater value for the cross-ratio $|a, x, b, y|$.

Fix now $a' \in [x, a] \cap S^1$. By the inequality $|x - a| \leq |a - b|$ and the triangle inequality,

$$(4.8) \quad \frac{|a - b|}{|x - a|} \leq \frac{|a - b| - |a - a'|}{|x - a| - |a - a'|} = \frac{|a - b| - |a - a'|}{|x - a'|} \leq \frac{|a' - b|}{|x - a'|}.$$

Let us yet show that there is a point $b' \in S^1$ such that

$$(4.9) \quad \frac{|a' - b|}{|y - b|} \leq \frac{|a' - b'|}{|y - b'|}.$$

If $|y - b| \leq |a' - b|$ holds for the point $b' \in [y, b] \cap S^1$, then the inequality (4.9) follows from the triangle inequality just like (4.8). If $|y - b| > |a' - b|$ instead, then there is $b \in \{i, -i\}$ such that $|y - b'| \leq |a' - b'|$ by the inequality (4.7) and the inequality (4.9) clearly holds for this choice of b' .

Thus, if $a, b \in f(\partial S_\theta)$ give the supremum of $|a, x, b, y|$ for given points $x, y \in i\mathbb{R} \cap \mathbb{B}^2$ and a', b' are chosen like above, it follows from the inequalities (4.8) and (4.9) that

$$|a, x, b, y| \leq |a', x, b, y| \leq |a', x, b', y| \leq \sup_{a', b' \in S^1} |a', x, b', y|.$$

By Proposition 4.1(2), if $x, y \in i\mathbb{R} \cap \mathbb{B}^2$ such that $\text{Im}(x) \leq \text{Im}(y)$, then $\sup_{a', b' \in S^1} |a', x, b', y|$ is given by $a = -i$ and $b = i$. Since $i, -i \in f(\partial S_\theta) \cap S^1$, it must hold that

$$\sup_{a, b \in f(\partial S_\theta)} |a, x, b, y| = |-i, x, i, y|.$$

Because f preserves the cross-ratio as a Möbius transformation, we can now show that, for all $x, y \in S_\theta$ such that $|x| = |y| = r$ and $\arg(x) \leq \arg(y)$,

$$\sup_{a, b \in \partial S_\theta} |a, x, b, y| = \sup_{a, b \in f(\partial S_\theta)} |a, f(x), b, f(y)| = |-i, f(x), i, f(y)| = |r, x, r e^{\theta i}, y|.$$

□

Note that Theorem 4.6 does not hold in the case $\theta > \pi$, as the following example shows.

Example 4.10. For $x = e^{(1-k)\theta i/2}$ and $y = e^{(1+k)\theta i/2}$ with $0 < k < 1$ and $\pi < \theta < 2\pi$,

$$\lim_{k \rightarrow 0^+} \frac{|1, x, e^{\theta i}, y|}{|0, x, \infty, y|} = \lim_{k \rightarrow 0^+} \frac{\sin(\theta/2)}{2 \sin^2((1-k)\theta/4)} = \frac{\sin(\theta/2)}{2 \sin^2(\theta/4)} = \frac{\cos(\theta/4)}{\sin(\theta/4)} < 1$$

and it follows that $\sup_{a, b \in \partial S_\theta} |a, x, b, y|$ is not attained with $a = 1$ and $b = e^{\theta i}$.

However, we can still use Lemma 4.2 to calculate the supremum of the cross-ratio in the Möbius metric for points x, y in a sector S_θ with $\theta > \pi$, as can be seen from the following result.

Corollary 4.11. *For any open sector S_θ with an angle $\pi < \theta < 2\pi$, there is a Möbius transformation f that maps S_θ onto the domain*

$$B^2((1-u)i, u) \cup B^2((u-1)i, u), \quad u = \frac{1}{1 - \cos(\theta/2)} \in \left(\frac{1}{2}, 1\right),$$




FIGURE 2. Sector S_θ before and after the Möbius transformation f defined in (4.3), when $\theta = 5\pi/4$.

and, for all $x, y \in S_\theta$,

$$\sup_{a, b \in \partial S_\theta} |a, x, b, y| = \sup_{a, b \in f(\partial S_\theta)} |a, f(x), b, f(y)|.$$

Proof. Let the Möbius transformation f be as in (4.3) with, for instance, $r = 1$. The proof now goes just like that of Theorem 4.6, but it must be noted that f maps the sides of S_θ onto circular arcs that meet each other at an angle $\theta > \pi$. Thus, $f(S_\theta)$ must be an union of two disks $B^2((1-u)i, u)$ and $B^2((u-1)i, u)$ instead of their intersection and $1/2 < u < 1$ now, see Figure 2. The final part of the proof follows from the Möbius invariance of the cross-ratio. \square

Several computational experiments support the next conjecture.

Conjecture 4.12. *If $\pi < \theta < 2\pi$ and $x = re^{(1-k)\theta i/2}$ and $y = re^{(1+k)\theta i/2}$ with $r > 0$ and $0 < k < 1$, then*

$$\sup_{a, b \in \partial S_\theta} |a, x, b, y| = \max\{|r, x, re^{\theta i}, y|, |0, x, \infty, y|\}.$$

The results of this section about the supremum of the cross-ratio give us information about the values of the Möbius metric δ_{S_θ} defined in a sector domain.

Corollary 4.13. *For all $x, y \in S_\theta$ with $0 < \theta < 2\pi$ such that $|x| = |y|$ and $\arg(x) \leq \arg(y)$,*

$$\delta_{S_\theta}(x, y) \geq \log \left(1 + \frac{\sin(\theta/2) \sin((\arg(y) - \arg(x))/2)}{\sin(\arg(x)/2) \sin((\theta - \arg(y))/2)} \right),$$

where the equality holds whenever $\theta \leq \pi$.

Proof. Let $x = re^{ui}$ and $y = re^{vi}$ with $0 < u \leq v < \theta$. By Theorem 4.6 and Proposition 4.1(1), the supremum $\sup_{a, b \in \partial S_\theta} |a, x, y, b|$ is now found by choosing $a = r$ and $b = re^{\theta i}$ if

$\theta \leq \pi$, and these choices of a, b give a lower limit for the supremum if $\theta > \pi$. The result follows now directly from the definition of $\delta_{S_\theta}(x, y)$. \square

5. MÖBIUS METRIC AND HYPERBOLIC METRIC IN OPEN SECTOR

In this section, we will study the connection between the Möbius metric and the hyperbolic metric in an open sector S_θ with an angle $0 < \theta < 2\pi$. The main result of this section is Corollary 5.11, which will be used to prove Theorem 1.2. However, in order to derive these results, we need to define the following quotient.

For all $0 < k < 1$ and $0 < \theta < 2\pi$, define

$$(5.1) \quad Q(k, \theta) \equiv \log \left(1 + \frac{\sin(\theta/2) \sin(k\theta/2)}{\sin^2((1-k)\theta/4)} \right) / \log \left(1 + \frac{\sin(k\pi/2)}{\sin^2((1-k)\pi/4)} \right).$$

The quotient above is very much needed here because it equals to the value of the quotient between the Möbius metric and the hyperbolic metric in certain cases, as will be shown in the next lemma.

Lemma 5.2. *For all $x, y \in S_\theta$ such that $x = re^{(1-k)\theta i/2}$ and $y = re^{(1+k)\theta i/2}$ with $r > 0$ and $0 < k < 1$,*

$$\frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)} = Q(k, \theta), \quad \text{if } 0 < \theta < \pi; \quad \text{and} \quad \frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)} \geq Q(k, \theta), \quad \text{if } \pi < \theta < 2\pi.$$

Proof. Recall the trigonometric identities $\sin(u) = \cos(\pi/2 - u)$ and $\cos(2v) = 1 - 2\sin^2(v)$. It follows from these that

$$1 - \sin(k\pi/2) = 1 - \cos((1-k)\pi/2) = 2\sin^2((1-k)\pi/4).$$

By using this formula, we will have

$$\begin{aligned} \rho_{S_\theta}(x, y) &= \rho_{S_\theta}(re^{(1-k)\theta i/2}, re^{(1+k)\theta i/2}) = \rho_{\mathbb{H}^2}(r^{\pi/\theta} e^{(1-k)\pi i/2}, r^{\pi/\theta} e^{(1+k)\pi i/2}) \\ &= \log \left(\frac{|e^{(1-k)\pi i/2} - e^{-(1+k)\pi i/2}| + |e^{(1-k)\pi i/2} - e^{(1+k)\pi i/2}|}{|e^{(1-k)\pi i/2} - e^{-(1+k)\pi i/2}| - |e^{(1-k)\pi i/2} - e^{(1+k)\pi i/2}|} \right) = \log \left(\frac{1 + \sin(k\pi/2)}{1 - \sin(k\pi/2)} \right) \\ &= \log \left(1 + \frac{2\sin(k\pi/2)}{1 - \sin(k\pi/2)} \right) = \log \left(1 + \frac{\sin(k\pi/2)}{\sin^2((1-k)\pi/4)} \right). \end{aligned}$$

Combining the expression above and Corollary 4.13, our result follows. \square

While the result of Lemma 5.2 only holds for distinct points $x, y \in S_\theta$ that are symmetric with respect to the angle bisector of the sector and fulfill $|x| = |y|$, Corollary 5.5 shows us why studying the quotient $Q(k, \theta)$ is useful also outside these restrictions.

Lemma 5.3. [14, Lemma 4.2, pp. 7-8] *For given two distinct points $x, y \in \mathbb{H}^2$, there exists a Möbius transformation $g : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ such that $|g(x)| = |g(y)| = 1$ and $\text{Im}(g(x)) = \text{Im}(g(y))$.*

(1) *If $\text{Im}(x) = \text{Im}(y)$, then $g(z) = (z - a)/r$ where $a = \text{Re}((x + y)/2)$ and $r = |x - a|$.*

- (2) If $\operatorname{Re}(x) = \operatorname{Re}(y) = a$ and $r = \sqrt{\operatorname{Im}(x)\operatorname{Im}(y)}$, then g is the Möbius transformation fulfilling $g(a - r) = 0$, $g(a) = 1$ and $g(a + r) = \infty$.
- (3) In the remaining case, the angle $\alpha = \angle(L(x, y), \mathbb{R})$ belongs to $(0, \pi/2)$. Let $S^1(c_1, r_1)$ and $S^1(c_2, r_2)$ be two circles centered at the real axis and orthogonal to each other, such that $x, y \in S^1(c_1, r_1)$ and $c_2 = L(x, y) \cap \mathbb{R}$. Then g is determined by $g(B^2(c_1, r_1) \cap \mathbb{H}^2) = \mathbb{B}^2 \cap \mathbb{H}^2$, $g(c_1 - r_1) = -1$, $g(c_1 + r_1) = 1$ and $g(S^1(c_2, r_2) \cap \mathbb{H}^2) = \{yi \mid y > 0\}$.

Lemma 5.4. [14, Lemma 4.5, p. 8] For all distinct points $x, y \in S_\theta$ with $0 < \theta < 2\pi$, there is a conformal mapping $f : S_\theta \rightarrow S_\theta$ such that $f(x) = e^{(1-k)\theta i/2}$ and $f(y) = e^{(1+k)\theta i/2}$ for some $k \in (0, 1)$.

Proof. Consider a conformal map $h : S_\theta \rightarrow \mathbb{H}^2$, $h(z) = z^{\pi/\theta}$. Fix $g : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ as the Möbius invariant map of Lemma 5.3 for the points $h(x), h(y)$. Define a conformal mapping $f = h^{-1} \circ g \circ h$. Since $|g(h(x))| = |g(h(y))| = 1$ and $\operatorname{Im}(g(h(x))) = \operatorname{Im}(g(h(y)))$, we can write $g(h(x)) = e^{(1-k)\pi i/2}$ and $g(h(y)) = e^{(1+k)\pi i/2}$ for some $0 < k < 1$. By using this k , we will have the points $f(x) = e^{(1-k)\theta i/2}$ and $f(y) = e^{(1+k)\theta i/2}$. \square

Corollary 5.5. For all $0 < \theta < 2\pi$ and distinct $x, y \in S_\theta$,

$$\inf_{0 < k < 1} Q(k, \theta) \leq \frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)} \leq \sup_{0 < k < 1} Q(k, \theta), \quad \text{if } 0 < \theta \leq \pi$$

$$\inf_{0 < k < 1} Q(k, \theta) \leq \frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)}, \quad \text{if } \pi < \theta < 2\pi$$

where $f(x) = e^{(1-k)\theta i/2}$ and $f(y) = e^{(1+k)\theta i/2}$.

Proof. Let the mappings f, g, h be as in Lemma 5.3 and the proof of Lemma 5.4. Note that, even though the mapping f does not necessarily preserve the distance $\delta_{S_\theta}(x, y)$, by Theorem 2.3 and the conformal invariance of the hyperbolic metric,

$$\delta_{\mathbb{H}^2}(h(x), h(y)) = \delta_{\mathbb{H}^2}(g(h(x)), g(h(y))) = \rho_{\mathbb{H}^2}(g(h(x)), g(h(y))) = \rho_{\mathbb{H}^2}(h(x), h(y)),$$

$$\rho_{S_\theta}(x, y) = \rho_{S_\theta}(f(x), f(y)),$$

for all points $x, y \in S_\theta$. It follows from this that

$$\inf_{x, y \in S_\theta} \frac{\delta_{S_\theta}(f(x), f(y))}{\rho_{S_\theta}(f(x), f(y))} \leq \frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)} \leq \sup_{x, y \in S_\theta} \frac{\delta_{S_\theta}(f(x), f(y))}{\rho_{S_\theta}(f(x), f(y))},$$

which leads to the result of our corollary by Lemmas 5.2 and 5.4. \square

Before studying the values of the quotient $Q(k, \theta)$, consider yet the following proposition.

Proposition 5.6. (1) For all constants $u, v \in (0, \pi]$, the quotients $\sin(uk)/\sin(vk)$ and $\sin((1-k)v)/\sin((1-k)u)$ are increasing with respect to $0 < k < 1$ if and only if $u \leq v$, and decreasing if $u > v$ instead.

(2) The quotient $\sin(t/2)/t$ is decreasing with respect to $0 < t < 2\pi$.

- (3) The quotient $t \sin(t)/\sin^2(t/2)$ is decreasing with respect to $0 < t < 2\pi$.
(4) The quotient $\log(1 + \mu q)/\log(1 + q)$ is increasing with respect to $q > 0$ if $0 < \mu \leq 1$, and decreasing if $\mu > 1$.

Proof. (1) First, define a function $f_1 : [0, \pi] \rightarrow \mathbb{R}$, $f_1(t) = \sin(t/2) - 2t$, and note that by calculus $f_1(t) \leq 0$ for all $0 \leq t \leq \pi$. Now, consider the function $f_2 : (0, \pi] \rightarrow \mathbb{R}$, $f_2(u) = u \cos(uk)/\sin(uk)$ with $0 < k < 1$. By differentiation and simple trigonometric identities,

$$f_2'(u) = \frac{\sin(uk/2) - 2uk}{2 \sin^2(uk)} = \frac{f_1(uk)}{2 \sin^2(uk)} \leq 0$$

so f_2 is decreasing with respect to $0 < u \leq \pi$. Finally, denote $f_3 : (0, 1) \rightarrow \mathbb{R}$, $f_3(k) = \sin(uk)/\sin(vk)$, where $u, v \in (0, \pi]$ are constants. By differentiation,

$$\begin{aligned} f_3'(k) &= \frac{u \cos(uk) \sin(vk) - v \sin(uk) \cos(vk)}{\sin^2(vk)} \geq 0 \quad \Leftrightarrow \quad \frac{u \cos(uk)}{\sin(uk)} \geq \frac{v \cos(vk)}{\sin(vk)} \\ \Leftrightarrow f_2(u) &\geq f_2(v) \quad \Leftrightarrow \quad u \leq v. \end{aligned}$$

This is enough to prove the result because $1/f_3(1 - k)$ is increasing (or decreasing) with respect to $0 < k < 1$ whenever f_3 is.

(2) Define $f_4, f_5 : (0, 2\pi) \rightarrow \mathbb{R}$, $f_4(t) = \sin(t/2)$, $f_5(t) = t$. Since $f_4(0) = f_5(0) = 0$ and $f_4'(t)/f_5'(t) = \cos(t/2)/2$ is decreasing with respect to t , by [5, Thm B.2, p. 465], $f_4(t)/f_5(t)$ is decreasing, too.

(3) Denote $f_6 : (0, 2\pi) \rightarrow \mathbb{R}$, $f_6(t) = t \sin(t)/\sin^2(t/2)$. By differentiation and some trigonometric identities,

$$f_6'(t) = \frac{(\sin(t) + t \cos(t))(1 - \cos(t)) - t \sin^2(t)}{2 \sin^4(t/2)} = \frac{(\sin(t) - t)(1 - \cos(t))}{2 \sin^4(t/2)} \leq 0,$$

so f_6 is decreasing for $0 < t < 2\pi$.

(4) Define $f_7, f_8 : (0, \infty) \rightarrow \mathbb{R}$, $f_7(q) = \log(1 + \mu q)$, $f_8(q) = \log(1 + q)$. Note that $f_7(0) = f_8(0) = 0$ and $f_7'(q)/f_8'(q) = \mu(1 + q)/(1 + \mu q)$ is increasing with respect to $q > 0$ if $0 < \mu \leq 1$ and decreasing if $\mu > 1$. By [5, Thm B.2, p. 465], the quotient $f_7(q)/f_8(q)$ is increasing (or decreasing) whenever $f_7'(q)/f_8'(q)$ is, so the result follows. \square

Theorem 5.7. *For all $0 < k < 1$ and $0 < \theta < 2\pi$, the quotient $Q(k, \theta)$ defined in (5.1) fulfills*

$$\begin{aligned} 1 \leq Q(k, \theta) &\leq (\pi \sin(\theta/2)/\theta)^2, \quad \text{if } \theta < \pi, \\ (\pi \sin(\theta/2)/\theta)^2 &\leq Q(k, \theta) \leq 1, \quad \text{if } \theta > \pi. \end{aligned}$$

Proof. Note that the quotient $Q(k, \theta)$ can be written as $\log(1 + q_0(k, \theta))/\log(1 + q_1(k))$, where

$$(5.8) \quad q_0(k, \theta) = \frac{\sin(\theta/2) \sin(k\theta/2)}{\sin^2((1 - k)\theta/4)}, \quad q_1(k) = \frac{\sin(k\pi/2)}{\sin^2((1 - k)\pi/4)}.$$

Clearly,

$$\frac{q_0(k, \theta)}{q_1(k)} = \sin(\theta/2) \frac{\sin(k\theta/2)}{\sin(k\pi/2)} \left(\frac{\sin((1-k)\pi/4)}{\sin((1-k)\theta/4)} \right)^2.$$

Trivially, $q_0(k, \theta)/q_1(k) = 1$, whenever $\theta = \pi$. It follows from Proposition 5.6(1) that $q_0(k, \theta)/q_1(k)$ is increasing with respect to k if $0 < \theta < \pi$, and decreasing if $\pi < \theta < 2\pi$. Thus, this quotient is bounded by its limit values

$$\mu_0(\theta) \equiv \lim_{k \rightarrow 0^+} \frac{q_0(k, \theta)}{q_1(k)} = \frac{\theta \sin(\theta/2)}{2\pi \sin^2(\theta/4)}, \quad \mu_1(\theta) \equiv \lim_{k \rightarrow 1^-} \frac{q_0(k, \theta)}{q_1(k)} = \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2.$$

By Proposition 5.6(2)-(3), both of the functions $\mu_0(\theta)$ and $\mu_1(\theta)$ are decreasing with respect to $0 < \theta < 2\pi$. Since $\mu_0(\pi) = \mu_1(\pi) = 1$, this means that the functions μ_0, μ_1 are greater than or equal to 1 for $0 < \theta < \pi$, and less than or equal to 1 for $\pi < \theta < 2\pi$. It follows that

$$(5.9) \quad \begin{aligned} q_1(k) &\leq \mu_0(\theta)q_1(k) \leq q_0(k, \theta) \leq \mu_1(\theta)q_1(k), \text{ if } 0 < \theta < \pi, \text{ and} \\ \mu_1(\theta)q_1(k) &\leq q_0(k, \theta) \leq \mu_0(\theta)q_1(k) \leq q_1(k), \text{ if } \pi < \theta < 2\pi. \end{aligned}$$

Recall now the expression for the quotient $q_1(k)$ from (5.8). This quotient $q_1(k)$ must be strictly increasing with respect to $0 < k < 1$, because it has a strictly increasing positive numerator $\sin(k\pi/2)$ and a strictly decreasing positive denominator $\sin^2((1-k)\pi/4)$. Since $q_1(k)$ has limit values $\lim_{k \rightarrow 0^+} q_1(k) = 0$ and $\lim_{k \rightarrow 1^-} q_1(k) = \infty$, it maps the interval $(0, 1)$ onto $(0, \infty)$. Furthermore, we already earlier noted that $\mu_1(\theta) \geq 1$ if $0 < \theta < \pi$, and $\mu_1(\theta) \leq 1$ if $\pi < \theta < 2\pi$. It follows from these observations, inequalities in (5.9) and Proposition 5.6(4) that, if $0 < \theta < \pi$,

$$\begin{aligned} \inf_{0 < k < 1} Q(k, \theta) &= \inf_{0 < k < 1} \frac{\log(1 + q_0(k, \theta))}{\log(1 + q_1(k))} \geq \inf_{0 < k < 1} \frac{\log(1 + q_1(k))}{\log(1 + q_1(k))} = 1, \\ \sup_{0 < k < 1} Q(k, \theta) &\leq \sup_{0 < k < 1} \frac{\log(1 + \mu_1(\theta)q_1(k))}{\log(1 + q_1(k))} = \lim_{q_1 \rightarrow 0^+} \frac{\log(1 + \mu_1(\theta)q_1)}{\log(1 + q_1)} = \mu_1(\theta), \end{aligned}$$

and, if $\pi < \theta < 2\pi$,

$$\begin{aligned} \inf_{0 < k < 1} Q(k, \theta) &\geq \inf_{0 < k < 1} \frac{\log(1 + \mu_1(\theta)q_1(k))}{\log(1 + q_1(k))} = \lim_{q_1 \rightarrow 0^+} \frac{\log(1 + \mu_1(\theta)q_1)}{\log(1 + q_1)} = \mu_1(\theta), \\ \sup_{0 < k < 1} Q(k, \theta) &\leq \sup_{0 < k < 1} \frac{\log(1 + q_1(k))}{\log(1 + q_1(k))} = 1, \end{aligned}$$

which proves the theorem. \square

It can be verified that the number 1 in the inequalities of Theorem 5.7 is the best possible constant by showing that it is the limit value of the quotient $Q(k, \theta)$ whenever $k \rightarrow 1^-$. However, the bound $(\pi \sin(\theta/2)/\theta)^2$ in Theorem 5.7 does not seem to be sharp. According to several numerical test, the quotient $Q(k, \theta)$ is monotonic with respect to k , either decreasing when $0 < \theta \leq \pi$ or increasing when $\pi \leq \theta < 2\pi$, which would lead into the following result.

Conjecture 5.10. For all $0 < k < 1$ and $0 < \theta < 2\pi$, the quotient $Q(k, \theta)$ fulfills

$$1 = \lim_{k \rightarrow 1^-} Q(k, \theta) \leq Q(k, \theta) \leq \lim_{k \rightarrow 0^+} Q(k, \theta) = \frac{\theta \sin(\theta/2)}{2\pi \sin^2(\theta/4)}, \quad \text{if } \theta < \pi,$$

$$\frac{\theta \sin(\theta/2)}{2\pi \sin^2(\theta/4)} = \lim_{k \rightarrow 0^+} Q(k, \theta) \leq Q(k, \theta) \leq \lim_{k \rightarrow 1^-} Q(k, \theta) = 1, \quad \text{if } \theta > \pi.$$

Finally, we will have the following result.

Corollary 5.11. For all $0 < \theta < 2\pi$ and $x, y \in S_\theta$, the following inequalities hold:

- (1) $\rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y) \leq (\pi \sin(\theta/2)/\theta)^2 \rho_{S_\theta}(x, y),$ if $\theta < \pi,$
- (2) $\delta_{S_\theta}(x, y) = \rho_{S_\theta}(x, y),$ if $\theta = \pi,$
- (3) $(\pi \sin(\theta/2)/\theta)^2 \rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y),$ if $\theta > \pi.$

Proof. Follows from Corollary 5.5, Theorem 5.7 and the fact that $Q(k, \pi) = 1$. \square

Remark 5.12. If Conjecture 5.10 holds, then the coefficient $(\pi \sin(\theta/2)/\theta)^2$ in Corollary 5.11 can be replaced by $\theta \sin(\theta/2)/(2\pi \sin^2(\theta/4))$, which gives us even sharper bounds for the Möbius metric.

Note that Corollary 5.11 does not offer an upper bound for the metric δ_{S_θ} in terms of ρ_{S_θ} in the case $\theta > \pi$. As stated in Corollary 5.5, $Q(k, \theta)$ is only a lower limit for the quotient $\delta_{S_\theta}(x, y)/\rho_{S_\theta}(x, y)$, so the result of Theorem 5.7 gives us this upper bound. Specifically, even though $Q(k, \theta) \leq 1$ for $\pi < \theta < 2\pi$, the inequality $\delta_{S_\theta}(x, y) \leq \rho_{S_\theta}(x, y)$ does not hold, as will be shown next.

Lemma 5.13. For all $\pi < \theta < 2\pi$, there are some points $x, y \in S_\theta$ such that $\delta_{S_\theta}(x, y) > \rho_{S_\theta}(x, y)$.

Proof. For $x = e^{(1-k)\theta i/2}$ and $y = e^{(1+k)\theta i/2}$ with $0 < k < 1$, the distance ρ_{S_θ} is as in the proof of Lemma 5.2 and, consequently,

$$\lim_{k \rightarrow 0^+} \frac{\delta_{S_\theta}(x, y)}{\rho_{S_\theta}(x, y)} \geq \lim_{k \rightarrow 0^+} \frac{|0, x, \infty, y|}{\rho_{S_\theta}(x, y)} = \lim_{k \rightarrow 0^+} \frac{\log(1 + 2 \sin(k\theta/2))}{\log(1 + \sin(k\pi/2)/\sin^2((1-k)\pi/4))} = \frac{\theta}{\pi} > 1.$$

\square

Finally, we can combine the inequalities of Corollary 5.11 with our earlier results from Section 3 in order to show that Theorem 1.2 holds.

5.14. Proof of Theorem 1.2. Follows directly from Corollaries 3.10, 3.12 and 5.11. Note that Theorem 1.2 only contains the best ones out of the bounds found and, for instance, the lower bound for $\delta_{S_\theta}(x, y)$ in Corollary 3.10(1) is never better than the one in Corollary 5.11(1). Similarly, it can be shown that Corollary 5.11 has always better lower bounds than Corollary 3.12.

6. MÖBIUS METRIC UNDER QUASIREGULAR MAPPINGS

In this section, we will yet briefly consider the behaviour of the Möbius metric under K -quasiregular mappings. This topic has already been researched in [15], see for instance [15, Thm 5.12, pp. 528-529], but we can improve the existing results with our new bounds for the Möbius metric in sector domains. However, let us first define all the concepts needed.

Definition 6.1. [5, pp. 289-288] Choose a domain $G \subset \mathbb{R}^n$ and let the function $f : G \rightarrow \mathbb{R}^n$ be ACLⁿ, as defined in [5, p. 150]. Suppose that there exists a constant $K \geq 1$ such that the inequality

$$(6.2) \quad |f'(x)|^n \leq K J_f(x), \quad |f'(x)| = \max_{|h|=1} |f'(x)h|,$$

where $J_f(x)$ is the Jacobian determinant of f at point $x \in G$, holds a.e. in G . Then the function f is called *quasiregular* and the smallest constant $K \geq 1$ fulfilling the inequality (6.2) is the *outer dilatation* of f . Similarly, the *inner dilatation* of f is the smallest constant $K \geq 1$ such that the inequality

$$(6.3) \quad J_f(x) \leq K \ell(f'(x))^n, \quad \ell(f'(x)) = \min_{|h|=1} |f'(x)h|$$

holds a.e. in G . The function f is K -*quasiregular*, if $\max\{K_I(f), K_O(f)\} \leq K$, where $K_I(f)$ and $K_O(f)$ are the inner and the outer dilatation of f , respectively.

See [5, (7.1), p. 104] and [16] for the definition of the conformal modulus of a curve family Γ and denote it by $M(\Gamma)$. For any non-empty subsets $F_0, F_1 \subsetneq \mathbb{R}^n$, let $\Delta(F_0, F_1; \mathbb{R}^n)$ be the family of all the closed non-constant curves joining these two subsets F_0 and F_1 . Furthermore, denote the Euclidean line segment between two points $x, y \in (\mathbb{R}^n \cup \{\infty\})$ by $[x, y]$ and let e_k be the k th unit vector of the n -dimensional space, $k = 1, \dots, n$. Now, we can define the *Grötzsch capacity* [5, (7.17), p. 121] as the decreasing homeomorphism $\gamma_n : (1, \infty) \rightarrow (0, \infty)$,

$$\gamma_n(s) = M(\Delta(\overline{\mathbb{B}^n}, [se_1, \infty]; \mathbb{R}^n)), \quad s > 1.$$

Note that, if $n = 2$, we have the following explicit formulas [5, (7.18), p. 122]

$$\gamma_2(1/r) = \frac{2\pi}{\mu(r)}, \quad \mu(r) = \frac{\pi}{2} \frac{\mathfrak{K}(\sqrt{1-r^2})}{\mathfrak{K}(r)}, \quad \mathfrak{K}(r) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-r^2x^2)}}.$$

By using the definition of the Grötzsch capacity, we can define also an increasing homeomorphism $\varphi_{K,n} : [0, 1] \rightarrow [0, 1]$, [5, (9.13), p. 167]

$$\varphi_{K,n}(r) = \frac{1}{\gamma_n^{-1}(K\gamma_n(1/r))} \quad \text{for } 0 < r < 1, K > 0; \quad \varphi_{K,n}(0) = 0, \quad \varphi_{K,n}(1) = 1,$$

and a number λ_n [5, (9.5) p. 157 & (9.6), p. 158]

$$\log \lambda_n = \lim_{t \rightarrow \infty} ((\gamma_n(t)/\omega_{n-1})^{1/(1-n)} - \log t),$$

where ω_{n-1} is the $(n-1)$ -dimensional surface area of the unit sphere $S^{n-1}(0, 1)$. For every $n \geq 2$, $4 \leq \lambda_n < 2e^{n-1}$, and $\lambda_2 = 4$.

The Schwarz lemma is one of the most well-known results in the distortion theory and, while its original version is about the distortion of the Euclidean metric under holomorphic functions, there exists the following modified version of the Schwarz lemma that tells about the distortion of the hyperbolic metric under K -quasiregular mappings.

Theorem 6.4. [5, Thm 16.2, p. 300 & Thm 16.39, p. 313] *Let $G, G' \in \{\mathbb{H}^n, \mathbb{B}^n\}$, $f : G \rightarrow f(G) \subset G'$ be a non-constant K -quasiregular mapping and $\alpha = K_I(f)^{1/(1-n)}$. Now,*

$$(1) \quad \operatorname{th} \frac{\rho_{G'}(f(x), f(y))}{2} \leq \varphi_{K,n} \left(\operatorname{th} \frac{\rho_G(x, y)}{2} \right) \leq \lambda_n^{1-\alpha} \left(\operatorname{th} \frac{\rho_G(x, y)}{2} \right)^\alpha,$$

$$(2) \quad \rho_{G'}(f(x), f(y)) \leq K_I(f)(\rho_G(x, y) + \log 4)$$

holds for all $x, y \in G$. Furthermore, in the two-dimensional case $n = 2$,

$$(3) \quad \rho_{G'}(f(x), f(y)) \leq c(K) \max\{\rho_G(x, y), \rho_G(x, y)^{1/K}\}$$

for all $x, y \in G$, where

$$c(K) = 2\operatorname{arth}(\varphi_{K,2}(\operatorname{th}(1/2))) \leq v(K-1) + K, \quad v = \log(2(1 + \sqrt{1 - 1/e^2})) < 1.3507,$$

as in [5, Thm 16.39, p. 313]. Here, $c(K) \rightarrow 1$ when $K \rightarrow 1$ and, by the conformal invariance of the hyperbolic metric, the result (3) also holds for any two simply connected planar domains G, G' because they can be mapped conformally onto the unit disk \mathbb{B}^2 .

Corollary 6.5. *If S_θ is a sector with angle $0 < \theta \leq \pi$ and $f : S_\theta \rightarrow S_\theta$ is a non-constant K -quasiregular mapping, then*

$$\delta_{S_\theta}(f(x), f(y)) \leq c(K) \min \left\{ 2, \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2 \right\} \max\{\delta_{S_\theta}(x, y), \delta_{S_\theta}(x, y)^{1/K}\}$$

for all $x, y \in S_\theta$.

Proof. Follows from Theorems 1.2(1)-(2) and 6.4(3). □

A similar result holds for a non-convex sector.

Corollary 6.6. *If S_θ is a sector with angle $\pi < \theta < 2\pi$ and $f : S_\theta \rightarrow S_\theta$ is a non-constant K -quasiregular mapping, then*

$$\delta_{S_\theta}(f(x), f(y)) \leq 4c(K) \max \left\{ \left(\frac{\theta}{\pi \sin(\theta/2)} \right)^2 \delta_{S_\theta}(x, y), \left(\frac{\theta}{\pi \sin(\theta/2)} \right)^{2/K} \delta_{S_\theta}(x, y)^{1/K} \right\}$$

for all $x, y \in S_\theta$.

Proof. Follows from Theorems 1.2(3) and 6.4(3). □

Corollary 6.7. *If S_θ is a sector with angle $\pi < \theta < 2\pi$ and $f : S_\theta \rightarrow S_\theta$ is a non-constant K -quasiregular mapping, then*

$$\operatorname{th} \frac{\delta_{S_\theta}(f(x), f(y))}{4} \leq c(K) \frac{\theta}{\pi} \left(2 \operatorname{th} \frac{\delta_{S_\theta}(x, y)}{2} \right)^{1/K}$$

for all $x, y \in S_\theta$.

Proof. By Theorems 1.2(3), 6.4(3) and 3.9(3),

$$\begin{aligned} \operatorname{th} \frac{\delta_{S_\theta}(f(x), f(y))}{4} &\leq \frac{\theta}{\pi} \operatorname{th} \frac{\rho_{S_\theta}(f(x), f(y))}{2} \leq \frac{\theta}{\pi} \operatorname{th} \left(\frac{c(K)}{2} \max\{\rho_{S_\theta}(x, y), \rho_{S_\theta}(x, y)^{1/K}\} \right) \\ &\leq \frac{\theta}{\pi} \operatorname{th} \left(\frac{c(K)}{2} \max\{2 \operatorname{arth}(s_{S_\theta}(x, y)), (2 \operatorname{arth}(s_{S_\theta}(x, y)))^{1/K}\} \right). \end{aligned}$$

It follows from [14, Thm 5.3, p. 11] that

$$\operatorname{th} \left(\frac{C}{2} \max\{2 \operatorname{arth}(t), (2 \operatorname{arth}(t))^{1/K}\} \right) \leq Ct^{1/K}$$

for all $0 < t < 1$, $K \geq 1$ and $C \geq 1$. Consequently,

$$\operatorname{th} \frac{\delta_{S_\theta}(f(x), f(y))}{4} \leq c(K) \frac{\theta}{\pi} s_{S_\theta}(x, y)^{1/K}.$$

By Corollary 3.8(2), $s_{S_\theta}(x, y) \leq 2 \operatorname{th}(\delta_{S_\theta}(x, y)/2)$, so the result follows. \square

Remark 6.8. Neither Corollary 6.6 nor Corollary 6.7 offers a bound for the distortion that is always better than the result of the other corollary, which can be seen by studying the case where $\theta \rightarrow \pi^-$ and $c(K) = K = 1$ for varying points $x, y \in S_\theta$.

Corollary 6.9. *If S_θ is a sector with angle $0 < \theta < 2\pi$ and $f : \mathbb{B}^2 \rightarrow S_\theta$ is a non-constant K -quasiregular mapping, then, for all $x, y \in \mathbb{B}^2$,*

$$\delta_{S_\theta}(f(x), f(y)) \leq c(K) \min \left\{ 2, \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2 \right\} \max\{\delta_{\mathbb{B}^2}(x, y), \delta_{\mathbb{B}^2}(x, y)^{1/K}\},$$

if $0 < \theta \leq \pi$, and

$$\delta_{S_\theta}(f(x), f(y)) \leq 4c(K) \max\{\delta_{\mathbb{B}^2}(x, y), \delta_{\mathbb{B}^2}(x, y)^{1/K}\} \quad \text{if } \pi < \theta < 2\pi.$$

Proof. Follows from Theorems 2.3, 1.2 and 6.4(3). \square

Corollary 6.10. *If S_θ is a sector with angle $0 < \theta \leq \pi$ and $f : \mathbb{B}^2 \rightarrow S_\theta$ is a non-constant K -quasiregular mapping, then for all $x \in \mathbb{B}^2$ such that $|x| \geq (e-1)/(e+1)$,*

$$|f(x)| \leq |f(0)| \left(\frac{1+|x|}{1-|x|} \right)^{c(K)u(\theta)} \quad \text{with } u(\theta) = \min \left\{ 2, \left(\frac{\pi \sin(\theta/2)}{\theta} \right)^2 \right\}.$$

Proof. By the triangle inequality, Theorems 3.1, 1.2 and 6.4(3), and [5, (4.14), p. 55],

$$\begin{aligned} \log \frac{|f(x)|}{|f(0)|} &\leq \log \frac{|f(x) - f(0)| + |f(0)|}{|f(0)|} = \log \left(1 + \frac{|f(x) - f(0)|}{|f(0)|} \right) \\ &\leq \log \left(1 + \frac{|f(x) - f(0)|}{\min\{d_{S_\theta}(f(x)), d_{S_\theta}(f(0))\}} \right) = j_{S_\theta}(f(x), f(0)) \leq \delta_{S_\theta}(f(x), f(0)) \\ &\leq u(\theta) \rho_{S_\theta}(f(x), f(0)) \leq c(K)u(\theta) \max\{\rho_{\mathbb{B}^2}(x, 0), \rho_{\mathbb{B}^2}(x, 0)^{1/K}\} \\ &= c(K)u(\theta) \max \left\{ \log \frac{1 + |x|}{1 - |x|}, \left(\log \frac{1 + |x|}{1 - |x|} \right)^{1/K} \right\}, \end{aligned}$$

and, if the inequality

$$\log \frac{1 + |x|}{1 - |x|} \geq \left(\log \frac{1 + |x|}{1 - |x|} \right)^{1/K} \quad \Leftrightarrow \quad \frac{1 + |x|}{1 - |x|} \geq e \quad \Leftrightarrow \quad |x| \geq \frac{e - 1}{1 + e}$$

holds, then we will have

$$\log \frac{|f(x)|}{|f(0)|} \leq c(K)u(\theta) \log \frac{1 + |x|}{1 - |x|} \quad \Leftrightarrow \quad |f(x)| \leq |f(0)| \left(\frac{1 + |x|}{1 - |x|} \right)^{c(K)u(\theta)}.$$

□

Remark 6.11. Note that Corollary 6.10 refines [5, Thm 16.19(1), p. 306] when $n = 2$.

7. MÖBIUS METRIC IN POLYGON

In this section, we will introduce a few open questions related to the Möbius metric inside a polygon domain. Especially, we are interested in the inequality between the Möbius metric and the hyperbolic metric defined in a polygon. All the computational findings and Figure 3 have been made with MATLAB programs from [11].

Even though the inequality $\rho_{S_\theta}(x, y) \leq \delta_{S_\theta}(x, y)$ holds in all convex sectors by Theorem 1.2 and these metrics are equivalent in such convex domains as the unit disk and the upper half-plane, our computer experiments verify that neither of metrics is always greater than or equal to the other in all polygonal domains, not even in all convex polygons.

Conjecture 7.1. *If $G \subsetneq \mathbb{R}^2$ is any bounded polygonal domain, there are always some points $x, y, u, v \in G$ such that $\rho_G(x, y) < \delta_G(x, y)$ and $\rho_G(u, v) > \delta_G(u, v)$.*

However, the values of these two metrics do not differ very much from each other in the domain G_k shaped like a regular convex k -gon with k vertices $e^{p2\pi i/k}$, $p = 0, 1, \dots, k - 1$, especially as the value of k grows and this domain resembles more and more the unit disk, where these metrics are equivalent.

Conjecture 7.2. *If G_k is the above regular k -gon and $x, y \in \cap G_k$ are distinct points, then*

$$\lim_{k \rightarrow \infty} (\delta_{G_k}(x, y) / \rho_{G_k}(x, y)) \rightarrow 1.$$




FIGURE 3. Contour plot of the quotient $\delta_G(x, y)/\rho_G(x, y)$, when G is the polygon with vertices $1, e^{0.95\pi i/3}, 0.1e^{\pi i/3}, e^{1.05\pi i/3}, e^{2\pi i/3}, e^{\pi i}, e^{4\pi i/3}, e^{5\pi i/3}$, the point x is fixed as 0 , and y varies inside G .

Furthermore, recall from Corollary 3.5 that the inequality $\delta_G(x, y) \leq 4\rho_G(x, y)$ holds for all points x, y in a simply-connected domain G , and our computer tests suggests that this result can be improved in the case of polygonal domains.

Conjecture 7.3. *For all polygonal domains $G \subsetneq \mathbb{R}^2$, the inequality $\rho_G(x, y)/2 \leq \delta_G(x, y) \leq 2\rho_G(x, y)$ holds for all $x, y \in G$.*

Figure 3 contains an example of a non-convex polygon where the values of the quotient $\delta_G(x, y)/\rho_G(x, y)$ vary at least on the interval $[0.73, 1.64]$. Note that, based on our computer tests, the latter constant in the inequality of Conjecture 7.3 can be replaced with a smaller one when considering only convex domains. Another interesting notion is that by Corollary 3.5 the uniformity constant A_G of a domain G is fulfills

$$A_G \geq \rho_G(x, y)/(2\delta_G(x, y))$$

for all points x, y in the domain x, y , so computing the maximum value of the quotient $\rho_G(x, y)/(2\delta_G(x, y))$ gives an lower bound for the uniformity constant A_G .

REFERENCES

1. J. CHEN, P. HARIRI, R. KLÉN AND M. VUORINEN, Lipschitz conditions, triangular ratio metric, and quasiconformal maps. *Ann. Acad. Sci. Fenn. Math.*, 40 (2015), 683-709. Zbl 1374.30069, MR3409699.
2. F.W. GEHRING AND K. HAG, *The ubiquitous quasidisk*. With contributions by Ole Jacob Broch. Mathematical Surveys and Monographs, 184. American Mathematical Society, Providence, RI, 2012. xii+171 pp. Zbl 1267.30003, MR2933660.

3. F.W. GEHRING AND B.G. OSGOOD, Uniform domains and the quasi-hyperbolic metric, *J. Analyse Math.*, *36* (1979), 50–74. Zbl 0449.30012, MR0581801.
4. F.W. GEHRING AND B. P. PALKA, Quasiconformally homogeneous domains, *J. Analyse Math.* *30* (1976), 172–199. Zbl 0349.30019, MR0437753.
5. P. HARIRI, R. KLÉN, AND M. VUORINEN, *Conformally Invariant Metrics and Quasiconformal Mappings*, Springer Monographs in Mathematics, Springer, Berlin, 2020. Zbl 1450.30003 MR4179585.
6. P. HARIRI, R. KLÉN, AND M. VUORINEN, Local convexity of metric balls. *Monatsh. Math.*, *186*, (2018), 281–298. Zbl 1395.51019, MR3808654.
7. P. HARIRI, M. VUORINEN AND X. ZHANG, Inequalities and Bilipschitz Conditions for Triangular Ratio Metric. *Rocky Mountain J. Math.*, *47*, 4 (2017), 1121–1148. Zbl 1376.30019, MR3689948.
8. P. HÄSTÖ, A new weighted metric, the relative metric I. *J. Math. Anal. Appl.*, *274* (2002), 38–58. Zbl 1019.54011, MR1936685.
9. P. HÄSTÖ, Z. IBRAGIMOV, D. MINDA, S. PONNUSAMY, AND S. SWADESH, Isometries of some hyperbolic-type path metrics, and the hyperbolic medial axis. (English summary) In the tradition of Ahlfors-Bers. IV, 63–74, *Contemp. Math.*, *432*, Amer. Math. Soc., Providence, RI, 2007. Zbl 1147.30029, MR2342807.
10. H. LINDÉN, Quasihyperbolic geodesics and uniformity in elementary domains. Dissertation, University of Helsinki, Helsinki, 2005. Ann. Acad. Sci. Fenn. Math. Diss. No. 146 (2005), 50 pp. Zbl 1092.51003, MR2183008.
11. M.M.S. NASSER, O. RAINIO, AND M. VUORINEN, Condenser capacity and hyperbolic perimeter. *Comput. Math. Appl.* *105* (2022) 54–74. Zbl 07447596, MR4345260.
12. O. RAINIO, Intrinsic quasi-metrics. *Bull. Malays. Math. Sci. Soc.* *44*, 5 (2021), 2873–2891. Zbl 1476.51019, MR4296316.
13. O. RAINIO AND M. VUORINEN, Introducing a new intrinsic metric. *Result. Math.* *77*, 2 (2022), 71. Zbl 07490182,
14. O. RAINIO AND M. VUORINEN, Triangular ratio metric under quasiconformal mappings in sector domains. *Comput. Methods Func. Theory* (to appear), arXiv: 2005.11990.
15. P. SEITTENRANTA, Möbius-invariant metrics. *Math. Proc. Cambridge Philos. Soc.* *125* (1999), 511–533. Zbl 0917.30015, MR1656825.
16. J. VÄISÄLÄ, *Lectures on n -Dimensional Quasiconformal Mappings*. Lecture Notes in Mathematics, 229. Springer-Verlag, Berlin, 1971. Zbl 0221.30031, MR0454009.
17. M. VUORINEN, *Conformal geometry and quasiregular mappings*. Lecture Notes in Mathematics, 1319. Springer-Verlag, Berlin, 1988. Zbl 0646.30025, MR0950174.