

# Modeling and structural analysis of cellulases using *Clostridium thermocellum* as template

Nathan Vinod Kumar<sup>1</sup>, Mary Esther Rani<sup>1\*</sup>, Rathinasamy Gunaseeli<sup>2</sup>, Narayanan Dhiraviam Kannan<sup>3</sup> & Jayavel Sridhar<sup>4</sup>

<sup>1</sup>Research Centre, Department of Botany and Microbiology, Lady Doak College, Madurai – 625002; <sup>2</sup>Center for Environmental Studies, Lady Doak College, Madurai -625002; <sup>3</sup>Department of Plant Biotechnology, School of Biotechnology, Madurai Kamaraj University, Madurai-625021; <sup>4</sup>UGC-NRCBS, School of Biological Sciences, Madurai Kamaraj University, Madurai-625021; Mary Esther Rani – Email: ranimam@yahoo.com; Phone: +91 9443064133; \*Corresponding author

Received October 16, 2012; Accepted October 26, 2012; Published November 13, 2012

## Abstract:

Cellulase is one of the most widely distributed enzymes with wide application. They are involved in conversion of biomass into simpler sugars. Cellulase of *Trichoderma longibrachiatum*, a known cellulolytic fungus was compared with *Clostridium thermocellum* [AAA23226.1] cellulase. Blastp was performed with AAA23226.1 as query sequence to obtain nine similar sequences from NCBI protein data bank. The physicochemical properties of cellulase were analyzed using ExPASy's ProtParam tool namely ProtParam, SOPMA and GOR IV. Homology modeling was done using SWISS MODEL and checked quality by RMSD values using VMD1.9.1. Active sites of each model were predicted using automated active site prediction server of SCFBio. Study revealed instability of cellulase of two eukaryotic strains namely *Trichoderma longibrachiatum* [CAA43059.1] and *Melanocarpus albomyces* [CAD56665.1]. The negative GRAVY score value of cellulases ensured better interaction and activity in aqueous phase. It was found that molecular weight (M. Wt) ranges between 25-127.56 kDa. Iso-electric point (pI) of cellulases was found to be acidic in nature. GOR IV and SOPMA were used to predict secondary structure of cellulase, which showed that random coil, was dominated. Neighbor joining tree with *C. thermocellum* [AAA23226.1] cellulase as root showed that cellulases of *Thermoaerobacter subterraneus* [ZP\_07835928] and *C. thermocellum* [CAA4305.1] were more similar to eukaryotic cellulases supported by least boot strap values. *Pseudoalteromonas haloplanktis* cellulase was found to be the ideal model supported by least RMSD score among the predicted structures. *Trichoderma longibrachiatum* cellulase was found to be the best compared to other cellulases, which possess high number of active sites with ASN and THR rich active sites. CYS residues were also present ensuring stable interaction and better bonding. Hydrophilic residues were found high in active sites of all analyzed models and template.

## Background:

Cellulases are important enzymes in many proposed processes for producing fuels and chemicals from plant biomass [1]. They are multienzyme complexes, comprising three major components; endo-  $\beta$ -glucanase (EC 3.2.1.4), exo-  $\beta$  - glucanase (EC 3.2.1.9.1) and  $\beta$ - glucosidase (EC 3.2.1.2.1), which have been shown to act synergistically in the hydrolysis of cellulose [2]. These enzymes are being isolated from many microbial sources and characterized. Cellulase plays a major role in carbon cycling in the biosphere by contributing towards the major carbon source [3]. The role of cellulase in host- pathogen interaction is quite important when considering the cellulase producing ability of pathogenic strains [4]. These enzymes help

in hydrolysis of substrates available and for the utilization of microbial growth and for their metabolism. There are no much reports available on computational characterization of the cellulase enzyme. *Clostridium thermocellum* is a thermophilic bacterial strain with high cellulolytic activity of about 5.32IU/L [5]. However, when eukaryotic cellulases are concern, *Trichoderma sp.* are widely used as a cellulase source and reported to possess very high activity. It was found to be the best strain for conversion of substrate into glucose of about 0.92mg/0.5ml which shows its higher cellulolytic activity [6].

Cellulase being an enzyme of wide application needs to be characterized in all aspects to understand the structural and

functional relations. Present study is to identify more efficient cellulolytic enzyme producing microorganism for bioprospecting using the computational analysis. Protein sequences of cellulase retrieved from NCBI and were subjected to ProtParam to analyze physicochemical parameter, secondary structure prediction using GOR IV and SOPMA, homology modeling (Swiss model), phylogenetic analysis and active site prediction by SCFBio.

## Methodology:

### Sequence retrieval and alignment

Cellulase protein sequence of *Clostridium thermocellum* [AAA23226.1] was retrieved from the National Center for Biotechnology (NCBI) and made as the query sequence for the structure, properties prediction and modeling. Blastp was performed and obtained nine similar sequences of different strains. Clustal W multiple sequence alignment was done for those sequences using BioEdit5.0.

### Secondary structure and physicochemical characterization cellulose

The sequences obtained were analyzed using various softwares available in the ExPASy server [7]. The GOR IV analysis was performed to understand the presence of helices, beta turns and coils in the protein structure [8]. Self-optimized prediction method with alignment (SOPMA) analysis was done for analyzing the structural components [9]. Comparison was made between the GOR IV and SOPMA analysis results. ProtParam software analysis was done to understand about the amino acid composition, molecular weight, instability index, aliphatic index and grand average of hydropathicity (GRAVY) [7]. Hydropathy plot analysis for all cellulase sequences was performed and the nature of amino acid residues were studied using ProtScale [7] based on Kyte and Doolittle scale.

### Homology modeling of cellulase

Homology models were predicted using SWISS- MODEL [10-12] and the quality was analysed using VMD 1. 9.1 [13]. RMSD values were calculated using the RMSD calculator and the best homology model was selected. Ramachandran plot for the best predicted model was depicted by RAMPAGE software [14].

### Phylogenetic analysis

Phylogenetic relation among the aligned cellulase sequences obtained from Blastp were analyzed based on neighbor joining method [15] using MEGA 4.0 [16]. The cellulase sequence of *C. thermocellum* [AAA23226.1] was considered as the root taxon for the analysis. Confidence level was analyzed using bootstrap of 1000 replications.

### Activity validation by active site comparison

Active sites of the predicted models and the template were analyzed using Automated Active Site prediction AADS server of SCFBio [17]. Amino acid compositions of all the cavities were analyzed and the frequency of amino acid occurrence in the cavities of each models were analyzed.

## Discussion:

### Blast analysis and sequence retrieval

The cellulase protein sequence of *Clostridium thermocellum* [AAA23226.1] was used as query sequence and nine sequences were obtained by performing Blastp. Multiple sequence


alignment was done in BioEdit software and further used for phylogenetic analysis in MEGA.

### Secondary structure and physicochemical analysis

SOPMA and GOR IV were used to predict the secondary structure, percentage of alpha, extended and random coils of cellulase producing microorganism are presented **Table 1 (see supplementary material)**. SOPMA analysis for the structure prediction was also done and obtained the percentage of alpha, extended, beta and random coils (**Table 1**). The secondary structure indicates whether a given amino acid lies in a helix, strand or coil [18, 19]. SOPMA was used for structure prediction of cellulase protein [20]. Random coil dominates the other forms in the cellulase analyzed by SOPMA and GOR IV. It was identified that random coils of *M. albomyces* (58.72%) and *T. longibrachiatum* (57.88%) were dominant compared to other forms. However, followed by random coils, extended forms ranging from (10%-27%) was dominant over  $\alpha$  and  $\beta$  helix. All the cellulases analyzed,  $\alpha$ -helix was ranging from (13%-37%) dominates  $\beta$ -helix, which had less percentage of conformation (4%-10%).

ProtParam analysis was performed and the number of amino acid residues, molecular weight, pI value, aliphatic index and GRAVY index was obtained for each sequence **Table 2 (see supplementary material)**. Comparison of the amino acid residue occurrence in cellulase sequences were done and the most dominant residues were highlighted **Table 3 (see supplementary material)**. It was found that molecular weight ranging from 25-127 kDa and it was higher in *C. thermocellum* (83 kDa) and lower in *M. albomyces* (25kDa). Comparing to the eukaryotic cellulase available, the higher aliphatic index of up to 97.51 was noted in *T. subterraneus* strains which indicate their stability over a wide range of temperatures. GRAVY value was negative in all species studied. It was notable that the bacterial strains had lower GRAVY values indicating the better possibilities of aqueous interaction. pI value showed that cellulase is acidic in nature. *T. subterraneus* had a slightly neutral pI value and the highest GRAVY value. Generally it was observed that towards acidic pI values the GRAVY tends to be low. In eukaryotic cellulases, the occurrence of  $\alpha$  helices was found to be too low. In case of *A. bisporus*,  $\alpha$  helices was similar to that of lower taxonomic groups. Moreover these cellulases possess higher percentages of random coils. A general pattern of inverse relationship between the percentage of occurrence of  $\alpha$  helices and random coils were observed in both higher and lower taxonomic levels.

Cellulase of *M. albomyces*, *T. longibrachiatum* and *R. flavefaciens FD-1* was classified as unstable ( $II > 40$ ) with an instability index (II) of 53.54, 55.23 and 54.34 respectively. It is notable that the *M. albomyces* and *T. longibrachiatum* are eukaryotic isolates and possess the least percentage of alpha helices in their structure. *P. haloplanktis* and *R. flavefaciens FD-1* with dominant amino acid residues Asn (10.1%) and Ser (11.6%) respectively which are hydrophilic residues, all the other sequences had ALA and GLY as dominant residues which are hydrophobic in nature. ALA was dominant in cellulases of *A. bisporus*, *C. thermocellum*, *P. carotovorum*, *Saccharophagus sp.* and *T. subterraneus* whereas, Gly was dominant for *C. thermocellum*, *M. albomyces* and *T. longibrachiatum*.



**Figure 1:** Homology Model of *Pseudoalteromonas haloplanktis* cellulase based on template Itvn predicted using SWISS MODEL. The model showed least RMSD value compared to other models


### Homology model validation

SWISS MODEL was used to predict the homology model of the cellulase sequences and the protein structure quality was analyzed. RMSD values for the models were calculated and the model with least value i.e. the best predicted model is shown in **(Figure 1)**. Ramachandran plot for the model was constructed using RAMPAGE software. Residue B 169 -LEU belonged to outlier region and the number of residues in the allowed and favoured region was very close to the expected values. It was observed that 94.8% of residues were in favoured region and 5.5% in allowed region. It was found that 0.2% was found in outlier region.

### Phylogenetic analysis

Phylogenetic tree was constructed using the ten sequences based on neighbour joining method with reference sequence *C. thermocellum* [AAA23226.1] as a root **(Figure 2)**. It was observed that the cellulase of *T. subterraneus* [ZP\_07835928.1] was found to be more related to the eukaryotic cellulases. *T. longibrachiatum* [CAA43059.1], *T. subterraneus* [ZP\_07835928.1], *M. albomyces* [CAD56665.1], *A. bisporus* [CAA83971.1], *C. thermocellum* [CAA43035.1] were belonging to same group. It can be implied that cellulase sequence of *T. subterraneus* and *C. thermocellum* were much similar to eukaryotic cellulase and it is not much evolved from the *C. thermocellum* [AAA23226.1] cellulase sequence. But the higher boot strap values for the other sequences supports its divergence from the root sequence. However, all the taxa of the group belonged to prokaryotic origin. There was no much influence for evolutionary divergence of the sequence with respect to variations in secondary structure.

Compared to bacterial cellulases, fungal cellulases are widely used. Moreover, the cellulolytic activities are high for fungal cellulases. Highest cellulase activity for *C. thermocellum* was 12.05IU/ml [5]. *P. haloplanktis* being a psychrophilic bacterium the cellulase obtained is cold adaptable. Cellulase from the former has conserved five amino acid residues in their active sites [21]. *C. thermocellum* is a thermophilic bacteria and its cellulase has a better heat stability. It is known to be ethanogenic strain and cellulase from this source has high commercial applications [22]. Cysteine residues contribute to protein thermal stability [22]. Amongst fungi, species of *Trichoderma* and *Aspergillus* are well known for cellulolytic potential [23]. Apart from the above, other fungi used for cellulase production are *Humicola* and *Aspergillus sp.* [24]. Hydrophathy plot for the cellulase sequence was constructed using ProtScale based on Kyte and Doolittle and the hydrophilicity and hydrophobicity nature was observed from the plot. It was observed that the majority of the residues were belonging to the hydrophilic regions confirming the interaction of the enzymes in aqueous medium. Aliphatic residues namely ALA, LEU, ILE and VAL were among the hydrophobic residues in the profile. Similarly, Phe which is an aromatic residue and sulfur containing residues MET and CYS were the other residues belonging to hydrophobic regions of ProtScale profile.



**Figure 2:** Neighbor joining tree showing evolutionary relationship among cellulase sequences of different origin were depicted using MEGA 4.0. Boot strap values are depicted at the nodes and branch lengths are also shown. *Clostridium thermocellum* CAA43035.1 and *T. subterraneus* ZP07835928.1 was grouped among eukaryotic cellulase sequences.

### Active site prediction based on active site

Active sites for each model and template were predicted using Active Site prediction server and tabulated **Table 4 (see supplementary material)**. It was found that *T. longibrachiatum* had most number of cavities (192). *C. thermocellum* [AAA23226.1] had 84 cavities which were very close to template with 85 cavities. Comparison of amino acid residues present in the cavities of each models were made. It was inferred that THR rich active sites may be favouring the enzyme activity in

extreme environments and ASN rich cavities may be contributing towards better enzyme activity. Among the analysed models, 4 models and the template was found to possess ASN as the dominant residue in its active sites. Both *C. thermocellum* and *R. flavefaciens* FD-1 cellulases had LYS rich active sites. ARG was dominant in active sites of *M. albomyces* [CAD56665.1] and *T. subterraneus* DSM 13965[ZP\_07835928.1] cellulases. However *P. haloplanktis*, an extremophile had THR dominant active sites. In *T. longibrachiatum* ASN and THR was found to be dominant in active sites with a frequency of 10.58. It is clearly notable that the hydrophilic amino acid residues are high in the active sites of these enzyme structures ensuring their interaction with substrate in aqueous phase. However the least found residue was CYS which assures stable interaction and bonding. Though the frequency of CYS was too low, it was found in both *C. thermocellum* and 3 eukaryotic cellulases. So this result validates the higher cellulolytic activity and *T. longibrachiatum* could be the source of most active cellulase from the present study.

### Conclusion:

These studies provide an insight for better prospecting of cellulolytic isolates from the environment for various industrial applications. Among the microbial cellulase used in the present work, *T. longibrachiatum* cellulase was found to be best with high number of active sites.

### Acknowledgement:

Authors are thankful to Department of Science and Technology, Government of India and TNSCST for providing the grant to facilitate the research. Moreover, we thank the Management, Lady Doak College, Madurai for providing us facilities and support for our work. JS thank the UGC Govt. of India for sponsoring the Networking Resource Centre in Biological Sciences at Madurai Kamaraj University, Madurai.

### References:

- [1] Wilson DB, *Appl Microbiol Biotechnol.* 2012 **93**: 497
- [2] Ryu DDY & Mandels M, *Enzym Microb Tech.* 1980 **2**: 91
- [3] Nowak J *et al. Mater Sci Eng.* 2005 **C25**: 119
- [4] Gibson DM *et al. Curr Opin Microbiol.* 2011 **14**: 264
- [5] Otajevwo FD & Aluyi HAS, *Modern Appl Sci.* 2011 **5**: 141
- [6] Omojasola P *et al. Nat Sci.* 2008 **6**: 64
- [7] <http://www.expasy.org/tools/>
- [8] [http://npsa-pbil.ibcp.fr/cgi-bin/npsa\\_automat.pi?page=npsa\\_gor4.html](http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pi?page=npsa_gor4.html).
- [9] Geourjon C & Deleage G, *Prot Eng.* 1994 **7**: 157
- [10] Arnold K *et al. Bioinformatics.* 2006 **22**: 195
- [11] Kiefer F *et al. Nucleic Acids Res.* 2009 **37**: 387 [PMID: 18931379]
- [12] Peitsch MC, *Bio/Technol.* 1995 **13**: 658
- [13] Humphrey W *et al. J Mol Graphics.* 1996 **14**: 33 [PMID: 8744570]
- [14] Lovell SC *et al. Proteins: Struct Funct Genet.* 2002 **50**: 437
- [15] Saitou N & Nei M, *Mol Biol & Evol.* 1987 **4**: 406 [PMID: 3447015]
- [16] Tamura K *et al. Mol Biol & Evol.* 2007 **24**: 1596 [PMID: 17486738]
- [17] Tanya S *et al. J Chem Inf Model.* 2011 **51**: 2515 [PMID: 21877713]
- [18] Jyotsna C *et al. Int J Eng Sci Tech.* 2010 **2**: 3468
- [19] Ojeiru FE *et al. Yonago Acta Medica.* 2010 **53**: 25
- [20] Pradeep NV *et al. Adv Life Sci Tech.* 2012 **4**: 8
- [21] Garsoux G *et al. Biochem J.* 2004 **384**: 247 [PMCID: PMC 3421452]
- [22] Xu B *et al. Eur J Biochem.* 2001 **268**: 3718
- [23] Lynd LR *et al. Microbiol Molec Biol Rev.* 2002 **66**: 506 [PMCID: PMC 120791]
- [24] Ghori MI *et al. Afr J Biotechnol.* 2011 **10**: 5878

Edited by P Kanguane

Citation: Kumar *et al.* Bioinformation 8(22): 1105-1110 (2012)

**License statement:** This is an open-access article, which permits unrestricted use, distribution, and reproduction in any medium, for non-commercial purposes, provided the original author and source are credited

## Supplementary material:

**Table 1:** Percentage of amino acids sequence forming secondary structure in GOR IV and SOPMA prediction

Sl. No	Organisms	Accession Number	GOR IV Analysis			SOPMA Predictions			
			α-helix (Hh) (%)	Extended strand (Ee) (%)	Random coils (%)	α-helix (Hh) (%)	β-turns (Tt) (%)	Extended strand (Ee) (%)	Random coils (%)
1	<i>Clostridium thermocellum</i>	AAA23226.1	25.24	22.54	52.23	37.52	4.72	11.61	46.15
2	<i>Agaricus bisporus</i>	CAA83971.1	25.57	17.12	57.31	30.82	7.08	10.5	51.6
3	<i>Melanocarpus albomyces</i>	CAD56665.1	9.36	24.26	66.38	13.62	5.96	21.7	58.72
4	<i>Clostridium thermocellum</i>	CAA43035.1	23.14	25.71	51.15	35.45	2.84	13.8	47.9
5	<i>Pectobacterium carotovorum</i>	AAC37033.1	20.27	26.8	52.93	29.95	6.53	22.52	40.99
6	<i>Trichoderma longibrachiatum</i>	CAA43059.1	6.91	25.27	67.82	15.77	7.34	19.01	57.88
7	<i>Pseudoalteromonas haloplanktis</i>	CAA76775.1	24.09	23.28	52.63	23.89	8.1	17.81	50.2
8	<i>Ruminococcus flavefaciens FD-1</i>	AAA26468.1	26.56	19.38	54.06	31.56	6.25	23.44	38.75
9	<i>Saccharophagus sp. Myt-1</i>	BAL42331.1	18.96	27.24	53.8	23.06	10.85	27.84	38.26
10	<i>Thermoaerobacter subterraneus</i>	ZP_07835928.1	48.62	11.6	39.78	37.02	6.63	22.38	33.98

**Table 2:** ProtParam analysis of cellulase sequences

Sl.No	Organisms	Accession Number	No. of Amino acids	M. Wt	pI	Aliphatic Index	GRAVY	Instability Index (II)	Stability
1	<i>Clostridium thermocellum</i>	AAA23226.1	741	83558.3	5.26	63.36	-0.497	29.82	Stable
2	<i>Agaricus bisporus</i>	CAA83971.1	438	46209.5	5.01	74.73	-0.166	34.57	Stable
3	<i>Melanocarpus albomyces</i>	CAD56665.1	235	25000.8	5.31	55.32	-0.251	53.54	Unstable
4	<i>Clostridium thermocellum</i>	CAA43035.1	739	82088.9	5.19	74.22	-0.324	29.93	Stable
5	<i>Pectobacterium carotovorum</i>	AAC37033.1	444	48300.6	5.32	77.77	-0.369	27.89	Stable
6	<i>Trichoderma longibrachiatum</i>	CAA43059.1	463	48337.0	4.80	55.23	-0.374	41.07	Unstable
7	<i>Pseudoalteromonas haloplanktis</i>	CAA76775.1	494	52873.4	4.21	69.13	-0.403	18.68	Stable
8	<i>Ruminococcus flavefaciens FD-1</i>	AAA26468.1	320	35938.0	4.43	85.94	-0.374	54.34	Unstable
9	<i>Saccharophagus sp. Myt-1</i>	BAL42331.1	1171	127561.5	4.77	71.85	-0.380	23.35	Stable
10	<i>Thermoaerobacter subterraneus</i>	ZP_07835928.1	362	39191.0	6.28	97.51	-0.067	38.89	Stable

**Table 3:** Occurrence percentage of amino acid residues (Highlighted are the highest amino acid frequencies)

Amino acid Residues	AAA23226.1	CAA83971.1	CAD56665.1	CAA43035.1	AAC37033.1	CAA43059.1	CAA76775.1	AAA26468.1	BAL42331.1	ZP_07835928.1
	Percentage of Occurrence in %									
Ala	8.60	13.20	10.20	9.60	8.60	6.50	9.30	5.90	10.20	11.90
Arg	3.50	2.30	5.10	3.50	4.50	2.40	1.40	2.50	3.10	8.80
Asn	4.50	6.80	3.80	4.90	7.40	7.60	10.10	7.80	7.90	0.60
Asp	6.70	5.70	6.00	6.80	7.00	4.50	8.90	7.20	6.00	5.80
Cys	0.30	2.10	6.00	0.80	0.00	4.80	1.80	0.60	1.00	0.30
Gln	2.80	5.70	4.30	2.70	3.80	3.70	3.00	0.90	4.70	3.00
Glu	5.10	2.10	3.40	4.90	3.20	2.20	3.80	8.80	5.10	6.90
Gly	8.80	8.70	12.30	8.40	9.50	12.10	8.90	2.50	9.20	10.20
His	1.60	1.10	1.70	1.90	1.60	1.10	1.80	1.60	2.70	3.90
Ile	4.20	3.90	1.70	4.60	5.40	3.00	5.10	10.60	5.20	4.10

Leu	5.50	6.60	6.40	7.00	7.00	6.30	4.90	5.90	5.40	9.40
Lys	5.90	3.40	2.10	5.30	3.60	1.90	3.40	5.30	2.80	2.50
Met	2.80	0.70	0.90	1.40	1.60	1.90	1.20	2.80	1.20	2.50
Phe	3.80	3.90	6.40	4.50	2.90	1.70	3.00	2.20	3.60	2.50
Pro	5.10	7.10	7.20	5.40	5.00	6.50	3.20	4.10	4.50	7.20
Ser	6.70	7.50	8.10	6.80	7.70	11.90	8.50	11.60	5.70	2.80
Thr	7.00	6.60	5.10	6.00	7.90	11.00	8.30	8.10	8.20	3.60
Trp	3.80	2.30	2.60	2.60	2.90	1.50	2.60	0.30	3.40	1.10
Tyr	7.30	3.20	2.10	6.50	3.40	5.20	3.20	5.90	3.00	1.70
Val	5.80	7.10	4.70	6.60	7.20	4.30	7.30	5.30	7.00	11.30

**Table 4:** Amino acid frequencies in active sites of predicted cellulase models and template

Percentage of amino acids in active sites											
Amino acid residues	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7	Model 8	Model 9	Model 10	Template
ALA	3.91	5.34	5.89	3.85	6.22	5.38	7.2	4.12	6.82	8.63	6.7
ARG	5.12	1.90	9.55	7.74	6.94	3.73	3.64	9.28	5.08	13.4	3.81
ASN	4.48	10.5	4.59	4.05	8.41	10.58	9.34	5.38	11.26	0.22	9.63
ASP	6.80	5.57	7.98	7.62	6.52	5.63	7.15	4.22	6.18	4.71	6.96
CYS	0.13	0.01	2.50	1.1	0	3.11	0	0	0	0.14	0
GLN	3.46	1.71	5.38	4.03	7.14	5.22	4.68	7.47	6.64	3.77	5.34
GLU	7.30	7.41	6.12	6.97	6.74	3.07	7.86	6.69	6.64	8.36	7.75
GLY	5.41	8.78	6.49	4.67	6.06	6.71	6.87	3.83	3.6	5.22	6.28
HIE	1.75	4.05	2.50	2.37	1.92	0.64	2.86	2.89	4.43	5.89	3.22
ILE	3.20	5.63	2.55	3.44	3.84	2.69	5.28	3.24	6.7	3.9	4.9
LEU	2.97	6.71	4.03	5.44	7.46	3.03	2.77	4.22	5.84	8.75	3.32
LYS	10.57	3.62	3.10	10.54	6.39	4.48	6.93	9.34	6.8	4.06	7.68
MET	1.46	3.62	0.00	0.97	0.48	1.58	0.69	6.42	1.74	1.88	0.8
PHE	4.11	3.19	5.80	3.96	3.13	1.86	2.46	6.65	2.38	3.9	2.79
PRO	5.55	6.67	7.79	4.91	5.15	6.43	2.43	3.15	2.23	6.9	3.06
SER	5.26	3.59	6.40	4.13	6.68	10.81	4.49	3.46	4.46	1.97	4.59
THR	9.11	5.34	5.05	4.84	5.11	10.58	10.79	4.58	4.49	2.43	9.51
TRP	6.09	8.27	5.15	5.7	5.69	3.39	6.15	8.19	5.78	1.7	5.7
TYR	9.82	3.87	4.22	9.7	3.26	7.11	4.14	2.52	4.16	3.67	3.84
VAL	3.40	4.22	4.91	3.88	2.86	3.97	4.27	4.35	4.77	10.5	4.12
<b>Total Cavities</b>	<b>84</b>	<b>34</b>	<b>26</b>	<b>58</b>	<b>39</b>	<b>192</b>	<b>71</b>	<b>29</b>	<b>37</b>	<b>49</b>	<b>85</b>