

1994

## Modeling Scattered Function Data on Curved Surfaces

Chandrajit L. Bajaj

Guoliang Xu

Report Number:  
94-050

---

Bajaj, Chandrajit L. and Xu, Guoliang, "Modeling Scattered Function Data on Curved Surfaces" (1994).  
*Department of Computer Science Technical Reports*. Paper 1150.  
<https://docs.lib.purdue.edu/cstech/1150>

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries.  
Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

**MODELING SCATTERED FUNCTION  
DATA ON CURVED SURFACES**

**Chandrajit Bajaj  
Guoliang Xu**

**CSD TR-94-050  
July 1994**

# MODELING SCATTERED FUNCTION DATA ON CURVED SURFACES\*

Chandrajit L. Bajaj

Department of Computer Science,  
Purdue University,  
West Lafayette, IN 47907  
email: bajaj@cs.purdue.edu

and

Guoliang Xu<sup>†</sup>

Computing Center,  
Chinese Academy of Sciences,  
Beijing, 100080  
email: xuguo@cs.purdue.edu

## Abstract

We present efficient algorithms to model a collection of scattered function data defined on a given smooth domain surface  $D$  in three dimensional real space ( $\mathbf{R}^3$ ), by a  $C^1$  cubic or a  $C^2$  quintic piecewise trivariate polynomial approximation  $F$  (a mapping from  $D$  into  $\mathbf{R}^4$ ). The smooth polynomial pieces or finite elements of  $F$  are defined on a three dimensional triangulation called the simplicial hull and defined over the domain surface  $D$ . Our smooth polynomial approximations allows one to additionally control the local geometry of the modeled function  $F$ . We also present two different techniques for visualizing the graph of the function  $F$ .

## 1 Introduction

In this paper, we consider the following problem: Given an arbitrary collection of points  $P = \{(x_i, y_i, z_i, F_i) \in \mathbf{R}^4\}_{i=1}^M$  with  $(x_i, y_i, z_i) \in \mathbf{R}^3$  on a given smooth surface  $D$ , called the *do-*

---

\*Supported in part by NSF grants CCR 92-22467, DMS 91-01424, AFOSR grants F49620-93-10138, F49620-94-1-0080, ONR grant N00014-94-1-0370 and NASA grant NAG-93-1-1473.

<sup>†</sup>While visiting the Computer Science Department, Purdue University, West Lafayette, Indiana, USA

main surface, construct a  $C^1/C^2$  (“/” stands for “or”) piecewise smooth function  $F$ , known as the *function-on-surface*, such that  $F(x_i, y_i, z_i) = F_i$ ,  $i = 1, \dots, M$ . Also visualize the graph of the function-on-surface  $F$ .

The problem of modeling and visualizing functions sampled on physical objects arises in several application areas: characterizing the rain fall on the earth, the pressure on the wing of an airplane and the temperature on a human body. A number of methods have been developed for dealing with this problem (for surveys see [3], [7]). Currently known approaches for approximating function-on-surface data however possess restrictions either on the domain surfaces or the function-on-surfaces. The domain surfaces are usually assumed to be spherical, convex or genus zero. The function-on-surface are not always polynomial [4], [8] or rather higher order polynomial [9] or a large number of pieces [1] compared to the approach of this paper. The method of [1] is a  $C^1$  Clough-Tocher scheme that splits a tetrahedron into 4 subtetrahedra, uses degree 5 polynomials and requires  $C^2$  data on the vertices of each subtetrahedron. Another Clough-Tocher scheme [10] requires only  $C^1$  data at the vertices, for again constructing a  $C^1$  function which is a cubic polynomial over each subtetrahedron, however splits the original tetrahedron into 12 pieces. A  $C^1$  scheme [9] that does not split each tetrahedron uses degree 9 polynomials and requires  $C^4$  data at the vertices. In extending the method of [9] to a  $C^2$  scheme, requires degree 17 polynomials and  $C^8$  data at the vertices of each tetrahedron. Compared to these approaches, our  $C^1/C^2$  construction has no splitting and uses much lower degree polynomials (cubic/quintic) requiring only  $C^1/C^2$  data respectively, at the vertices of each tetrahedron.

Our solution to the modeling problem involves the following steps: (a). Construct a planar triangular approximation  $T$  of the domain surface  $D$  in the region of the points  $(x_i, y_i, z_i)$  on  $D$ . (b). Generate  $C^1/C^2$  data at the vertices of the triangulation  $T$  for a desired  $C^1/C^2$  smooth approximation, respectively. (c). Construct a simplicial hull (defined below)  $\Sigma$  surrounding the triangulation  $T$ . (d). Build the  $C^1/C^2$  function-on-surface  $F$  over  $\Sigma$  by locally interpolating the  $C^1/C^2$  data, respectively. (e). Visualize the graph of the function-on-surface  $F$ . We shall not address the first two steps (a) and (b) in this paper. A algorithm for the construction of the triangulation  $T$  of the given surface is given in [5]. See also Figure 1.1. However, we require our triangulation to satisfy certain conditions which will be discussed in §3. The problem of estimating the  $C^1/C^2$  data at the vertices of  $T$  is studied in a separate paper [2]. In this paper, we detail the steps (c), (d) and (e) in §3, §4, and §5 respectively, after the notation and preliminary section §2.

## 2 Notation and Preliminary Details

**Bernstein-Bezier (BB) Form:** Let  $p_1, p_2, p_3, p_4 \in \mathbb{R}^3$  be affine independent. Then the tetrahedron with vertices  $p_1, p_2, p_3$ , and  $p_4$  is the convex hull defined by  $[p_1 p_2 p_3 p_4] = \{p \in \mathbb{R}^3 : p = \sum_{i=1}^4 \alpha_i p_i, \alpha_i \geq 0, \sum_{i=1}^4 \alpha_i = 1\}$ . For any  $p = \sum_{i=1}^4 \alpha_i p_i \in [p_1 p_2 p_3 p_4]$ ,  $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)^T$  denotes the barycentric coordinates of  $p$ . Any polynomial  $f(p)$  of degree  $n$  can be expressed as Bernstein-Bezier (BB) form over  $[p_1 p_2 p_3 p_4]$  as  $f(p) = \sum_{|\lambda|=n} b_\lambda B_\lambda^n(\alpha)$ ,  $\lambda \in \mathcal{Z}_+^4$ , where  $B_\lambda^n(\alpha) = \frac{n!}{\lambda_1! \lambda_2! \lambda_3! \lambda_4!} \alpha_1^{\lambda_1} \alpha_2^{\lambda_2} \alpha_3^{\lambda_3} \alpha_4^{\lambda_4}$  is Bernstein polynomial,  $|\lambda| = \sum_{i=1}^4 \lambda_i$  with  $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)^T =$




Figure 1.1: A piecewise smooth domain surface  $D_1$  and a triangulation on it.

$\sum_{i=1}^4 \lambda_i e_i$ ,  $b_\lambda = b_{\lambda_1 \lambda_2 \lambda_3 \lambda_4}$  (as a subscript, we simply write  $\lambda$  as  $\lambda_1 \lambda_2 \lambda_3 \lambda_4$ ) are called control points or weights, and  $\mathcal{Z}_+^4$  stands for the set of all four dimensional vectors with nonnegative integer components. The following basic facts about the BB form will be used in this paper.

**Lemma 2.1.** *Let  $f(p) = F(\alpha) = \sum_{|\lambda|=n} b_\lambda B_\lambda^n(\alpha)$  where  $\alpha$  denotes the barycentric coordinates of  $p$ . Then for any pair of points  $p^{(1)}$  and  $p^{(2)}$ , with  $\alpha^{(1)}$  and  $\alpha^{(2)}$  as their barycentric coordinates, we have*

$$\begin{aligned} \nabla f(p)^T (p^{(1)} - p^{(2)}) &= n \sum_{|\lambda|=n-1} b_\lambda^1 (\alpha^{(1)} - \alpha^{(2)}) B_\lambda^{n-1}(\alpha) \\ (p^{(1)} - p^{(2)})^T \nabla^2 f(p) (p^{(1)} - p^{(2)}) &= n(n-1) \sum_{|\lambda|=n-2} b_\lambda^2 (\alpha^{(1)} - \alpha^{(2)}) B_\lambda^{n-2}(\alpha) \end{aligned}$$

where  $\nabla f(p) = [\frac{\partial f(p)}{\partial x} \quad \frac{\partial f(p)}{\partial y} \quad \frac{\partial f(p)}{\partial z}]^T$ ,  $\nabla^2 f(p) = [\nabla \frac{\partial f(p)}{\partial x} \quad \nabla \frac{\partial f(p)}{\partial y} \quad \nabla \frac{\partial f(p)}{\partial z}]$  and  $b_\lambda^r (\alpha^{(1)} - \alpha^{(2)}) = \sum_{|j|=r} b_{\lambda+j} B_j^r(\alpha^{(1)} - \alpha^{(2)})$

See [6] for the two dimensional case of the above lemma. From this lemma we have

**Corollary 2.2.** *Let  $f(p) = \sum_{|\lambda|=n} b_\lambda B_\lambda^n(\alpha)$  be defined on the tetrahedron  $[p_1 p_2 p_3 p_4]$ , then*

$$b_{(n-1)e_i + e_j} = b_{ne_i} + \frac{1}{n} (p_j - p_i)^T \nabla f(p_i), \quad j \neq i \quad (2.1)$$

$$\begin{aligned} b_{(n-2)e_i + e_j + e_k} &= -b_{ne_i} + b_{(n-1)e_i + e_j} + b_{(n-1)e_i + e_k} \\ &+ \frac{1}{n(n-1)} (p_j - p_i)^T \nabla^2 f(p_i) (p_k - p_i), \quad j \neq i, k \neq i \end{aligned} \quad (2.2)$$




Figure 2.1: The related control points of  $C^1$  (a) and  $C^2$  (b) conditions

The corollary tell us that the weights around a vertex can be computed from the given  $C^2$  data.

**Lemma 2.3** ([6]). *Let  $f(p) = \sum_{|\lambda|=n} a_\lambda B_\lambda^n(\alpha)$  and  $g(p) = \sum_{|\lambda|=n} b_\lambda B_\lambda^n(\alpha)$  be two polynomials defined on two tetrahedra  $[p_1 p_2 p_3 p_4]$  and  $[p'_1 p_2 p_3 p_4]$ , respectively. Then*

(i)  *$f$  and  $g$  are  $C^0$  continuous at the common face  $[p_2 p_3 p_4]$  if and only if*

$$a_\lambda = b_\lambda, \text{ for any } \lambda = 0\lambda_2\lambda_3\lambda_4, \quad |\lambda| = n \quad (2.3)$$

(ii)  *$f$  and  $g$  are  $C^1$  continuous at the common face  $[p_2 p_3 p_4]$  if and only if (2.3) holds and*

$$b_{1\lambda_2\lambda_3\lambda_4} = \beta_1 a_{1\lambda_2\lambda_3\lambda_4} + \beta_2 a_{0\lambda_2\lambda_3\lambda_4+0100} + \beta_3 a_{0\lambda_2\lambda_3\lambda_4+0010} + \beta_4 a_{0\lambda_2\lambda_3\lambda_4+0001} \quad (2.4)$$

(iii)  *$f$  and  $g$  are  $C^2$  continuous at the common face  $[p_2 p_3 p_4]$  if and only if (2.3)-(2.4) holds and*

$$\begin{aligned} b_{2\lambda_2\lambda_3\lambda_4} &= \beta_1^2 a_{2\lambda_2\lambda_3\lambda_4} + 2\beta_1\beta_2 a_{0\lambda_2\lambda_3\lambda_4+1100} + 2\beta_1\beta_3 a_{0\lambda_2\lambda_3\lambda_4+1010} + 2\beta_1\beta_4 a_{0\lambda_2\lambda_3\lambda_4+1001} \\ &+ \beta_2^2 a_{0\lambda_2\lambda_3\lambda_4+0200} + 2\beta_2\beta_3 a_{0\lambda_2\lambda_3\lambda_4+0110} + 2\beta_2\beta_4 a_{0\lambda_2\lambda_3\lambda_4+0101} \\ &+ \beta_3^2 a_{0\lambda_2\lambda_3\lambda_4+0020} + 2\beta_3\beta_4 a_{0\lambda_2\lambda_3\lambda_4+0011} + \beta_4^2 a_{0\lambda_2\lambda_3\lambda_4+0002} \end{aligned} \quad (2.5)$$

where  $\beta = (\beta_1, \beta_2, \beta_3, \beta_4)^T$  are defined by the relation  $p'_1 = \beta_1 p_1 + \beta_2 p_2 + \beta_3 p_3 + \beta_4 p_4$ ,  $|\beta| = 1$ .

In Lemma 2.3, if we divide (2.4) and (2.5) by  $\beta_4^2$ , then the  $C^1$  and  $C^2$  conditions become

$$a_{0\lambda_2\lambda_3\lambda_4+0001} = \mu_1 a_{1\lambda_2\lambda_3\lambda_4} + \mu_2 b_{1\lambda_2\lambda_3\lambda_4} + \mu_3 a_{0\lambda_2\lambda_3\lambda_4+0100} + \mu_4 a_{0\lambda_2\lambda_3\lambda_4+0010} \quad (2.6)$$

$$\begin{aligned} &\mu_1 (\mu_1 a_{2\lambda_2\lambda_3\lambda_4} + \mu_3 a_{0\lambda_2\lambda_3\lambda_4+1100} + \mu_4 a_{0\lambda_2\lambda_3\lambda_4+1010} - a_{0\lambda_2\lambda_3\lambda_4+1001}) \\ &= \mu_2 (\mu_2 b_{2\lambda_2\lambda_3\lambda_4} + \mu_3 b_{0\lambda_2\lambda_3\lambda_4+1100} + \mu_4 b_{0\lambda_2\lambda_3\lambda_4+1010} - b_{0\lambda_2\lambda_3\lambda_4+1001}) \end{aligned} \quad (2.7)$$

respectively, where  $\mu_1 = -\frac{\beta_1}{\beta_4}$ ,  $\mu_2 = \frac{1}{\beta_4}$ ,  $\mu_3 = -\frac{\beta_2}{\beta_4}$ ,  $\mu_4 = -\frac{\beta_3}{\beta_4}$ , that is  $p_4 = \mu_1 p_1 + \mu_2 p'_1 + \mu_3 p_2 + \mu_4 p_3$ .

It is not difficult to show the following from Corollary 2.2 :

**Lemma 2.4.** *Let  $f(p)$  and  $g(p)$  be defined as Lemma 2.3. If the coefficients of  $f$  and  $g$  around the vertices are determined by (2.1)-(2.2), then the  $C^1$  and  $C^2$  conditions (2.4)-(2.5) related only to these coefficients are satisfied.*

**Degree Elevation.** The polynomial  $f(p) = \sum_{|\lambda|=n} b_\lambda B_\lambda^n(\alpha)$  can be written as one of degree  $n + 1$  (see e.g. [6]).  $f(p) = \sum_{|\lambda|=n+1} (Eb)_\lambda B_\lambda^{n+1}(\alpha)$ ,  $\lambda \in \mathcal{Z}_+^4$  where  $(Eb)_\lambda = \frac{1}{n+1} \sum_{i=1}^4 \lambda_i b_{\lambda - e_i}$ . We shall use these formulas in approximating lower degree polynomials, in §4.

### 3 Simplicial Hull

Given a planar triangular approximation  $T$  of  $D$  containing (and not necessarily as vertices) the points  $(x_i, y_i, z_i)$  on  $D$ , a *simplicial hull* of  $D$  and  $T$ , denoted by  $\Sigma$ , is a collection of non-degenerate tetrahedra which satisfies the following:

(1) Each tetrahedron in  $\Sigma$  has either a single edge of  $T$  (then it will be called an *edge tetrahedron*) or a single face of  $T$  (then it will be called a *face tetrahedron*).

(2) For each face  $f$  of  $T$  there are at most two face tetrahedra (above and below  $f$ ) in  $\Sigma$  that share the face  $f$ .

(3) Two face tetrahedra that share a common edge do not intersect in any other region. This condition is referred to in this paper as *non-self-intersection*.

(4) For each edge there are two pairs of common face sharing edge tetrahedra in  $\Sigma$ , such that each pair blends the two adjacent face tetrahedra on the same side.

(5) The surface  $D$  is contained in  $\Sigma$ . This condition is referred to in this paper as the *surface containment condition*.

Therefore, a simplicial hull of  $D$  and  $T$  is in a neighborhood surrounding  $D$ . It should be noted that, for the given triangulation  $T$  of  $D$ , there may exist infinitely many simplicial hulls or perhaps no simplicial hull may exist. However under the following conditions on  $T$ , we can always construct a simplicial hull.

**Condition 1.** *The triangulation  $T$  is locally even. That is for every face of  $T$ , say  $[p_1 p_2 p_3]$ , the angle between the surface normal  $n_i$  at the vertex  $p_i$  and the normal of the face  $[p_1 p_2 p_3]$  is less than*

$$\tan^{-1} \left( \frac{2s \tan(\frac{1}{2} \min\{\alpha_1, \alpha_2, \alpha_3\})}{\| \|p_j - p_i\| (p_k - p_i) + \|p_k - p_i\| (p_j - p_i) \|} \right)$$

for  $i = 1, 2, 3$  and distinct  $1 \leq i, j, k \leq 3$ . Here  $s$  is the area of the face  $[p_1 p_2 p_3]$ , and  $\alpha_1, \alpha_2, \alpha_3$  are the dihedral angles of the three edges of the face  $[p_1 p_2 p_3]$ .

**Condition 2.** *The surface  $D$  is single sheeted on  $T$ . That is, for every face of  $T$ , say  $[p_1 p_2 p_3]$  let  $L$  be a straight line that is perpendicular to the face  $f$  and passes through the center  $c$  of the inscribed circle of  $f$ . Let  $p_4$  and  $q_4$  be the center's nearest points on  $L$  off each side of  $f$  such that  $\|p_4 - c\| = \|q_4 - c\|$  and the three tangent planes at the three vertices are contained in  $[p_4 p_1 p_2 p_3 q_4]$ . Then for any  $p \in f$  the broken line  $[p_4 p q_4]$  intersects the surface  $D$  only once.*

**Condition 3.** *Any two adjacent faces are not coplanar.*

Since the given surface is curved and smooth, by adding additional points on  $D$ , we can modify the algorithm of [5] to achieve a  $T$  satisfying the above conditions.

For such a  $T$  we now show how to construct a simplicial hull  $\Sigma$  in two easy steps.

**1. Build Face Tetrahedra.** For each face  $f = [p_1 p_2 p_3]$  of  $T$ , let  $L$  be a straight line that is perpendicular to the face  $f$  and passes through the center  $c$  of the inscribed circle of  $f$ . Let  $p_4$

and  $q_4$  be the center's nearest points on  $L$  off each side of  $f$  such that  $\|p_4 - c\| = \|q_4 - c\|$  and the three tangent planes at the three vertices are contained in  $[p_4 p_1 p_2 p_3 q_4]$ , then construct two face tetrahedra  $[p_1 p_2 p_3 p_4]$  and  $[p_1 p_2 p_3 q_4]$ .

**2. Build Edge Tetrahedra.** Let  $[p_2 p_3]$  be an edge of  $T$  and  $[p_1 p_2 p_3]$  and  $[p'_1 p_2 p_3]$  be the two adjacent faces. Let  $[p_1 p_2 p_3 p_4]$  and  $[p_1 p_2 p_3 q_4]$ , and  $[p'_1 p_2 p_3 p'_4]$  and  $[p'_1 p_2 p_3 q'_4]$  be the face tetrahedra built for the faces  $[p_1 p_2 p_3]$  and  $[p'_1 p_2 p_3]$ , respectively. Now two pairs of tetrahedra are constructed. The first pair  $[p''_1 p_2 p_3 p_4]$  and  $[p''_1 p_2 p_3 p'_4]$  is between  $[p'_1 p_2 p_3 p'_4]$  and  $[p_1 p_2 p_3 p_4]$ . The second pair  $[q''_1 p_2 p_3 q_4]$  and  $[q''_1 p_2 p_3 q'_4]$  is between  $[p'_1 p_2 p_3 q'_4]$  and  $[p_1 p_2 p_3 q_4]$ . Here  $p''_1 \in (p_4 p'_4)$  or above  $(p_4, p'_4)$ , say  $p''_1 = \frac{(1-t)}{2}(p_2 + p_3) + \frac{t}{2}(p'_4 + p_4)$ ,  $t \geq 1$ , so that  $p''_1$  is above  $[p_2, p_3]$  and the surface containment condition is satisfied. Similarly,  $q''_1 \in (q_4 q'_4)$  or below  $(q_4, q'_4)$ , say  $q''_1 = \frac{(1-t)}{2}(p_2 + p_3) + \frac{t}{2}(q'_4 + q_4)$ ,  $t \geq 1$ , so that  $q''_1$  is below  $[p_2, p_3]$  and the surface containment condition is satisfied.

The locally even condition guarantees that the face tetrahedron constructed has height (the distance between the top vertex  $p_4$  or  $q_4$  to the face) at most  $r \tan(\frac{1}{2} \min\{\alpha_0, \alpha_1, \alpha_2\})$ , where  $r$  is the radius of the inscribed circle. Hence the dihedral angles at the bottom edges of the tetrahedron are less than  $\frac{1}{2} \min\{\alpha_0, \alpha_1, \alpha_2\}$ . Therefore, there is no additional intersection between two adjacent face tetrahedra.

## 4 $C^1/C^2$ Interpolation by Cubic/Quintic

Suppose we have established a simplicial hull  $\Sigma$  for the given triangulation  $T$  of  $D$ . Now we construct a  $C^1/C^2$  function  $f$  over  $\Sigma$  such that  $f$  has the given  $C^1/C^2$  data, respectively at each vertex. Let  $V_1 = [p_1 p_2 p_3 p_4]$ ,  $V_2 = [p'_1 p_2 p_3 p'_4]$ ,  $W_1 = [p''_1 p_2 p_3 p_4]$ ,  $W_2 = [p''_1 p_2 p_3 p'_4]$ ,  $V'_1 = [p_1 p_2 p_3 q_4]$ ,  $V'_2 = [p'_1 p_2 p_3 q'_4]$ ,  $W'_1 = [q''_1 p_2 p_3 q_4]$ ,  $W'_2 = [q''_1 p_2 p_3 q'_4]$  and the cubic/quintic polynomials  $f_i$  over  $V_i$ ,  $g_i$  over  $W_i$ ,  $f'_i$  over  $V'_i$  and  $g'_i$  over  $W'_i$  be expressed in Bernstein-Bezier form with coefficients  $a_\lambda^{(i)}$ ,  $b_\lambda^{(i)}$ ,  $c_\lambda^{(i)}$  and  $d_\lambda^{(i)}$ , respectively. Now we shall determine these coefficients step by step. Denote

$$\begin{aligned} p''_1 &= \beta_1^{(1)} p_1 + \beta_2^{(1)} p_2 + \beta_3^{(1)} p_3 + \beta_4^{(1)} p_4, & \beta_1^{(1)} + \beta_2^{(1)} + \beta_3^{(1)} + \beta_4^{(1)} &= 1 \\ p'_1 &= \beta_1^{(2)} p'_1 + \beta_2^{(2)} p_2 + \beta_3^{(2)} p_3 + \beta_4^{(2)} p'_4, & \beta_1^{(2)} + \beta_2^{(2)} + \beta_3^{(2)} + \beta_4^{(2)} &= 1 \\ p'_1 &= \mu_1 p_4 + \mu_2 p'_4 + \mu_3 p_2 + \mu_4 p_3, & \mu_1 + \mu_2 + \mu_3 + \mu_4 &= 1 \end{aligned} \quad (4.1)$$

### $C^1$ Cubic Scheme

- (1) The number 0 weights (see Figure 4.1) are given by the function values at the vertices.
- (2) The number 1 weights are determined by formula (2.1) from  $C^1$  data.
- (3) The number 2 weights, that is  $a_{1110}^{(i)}$ , are free.
- (4) The number 3 weights are determined by  $C^1$  conditions (2.4) and (2.6). More precisely,

$$a_{0111}^{(i)} = \theta_1^{(i)} a_{1110}^{(1)} + \theta_2^{(i)} a_{0210}^{(i)} + \theta_3^{(i)} a_{0120}^{(i)} + \theta_4^{(i)} a_{1110}^{(2)}, \quad i = 1, 2$$

where

$$\begin{aligned} p_4 &= \theta_1^{(1)} p_1 + \theta_2^{(1)} p_2 + \theta_3^{(1)} p_3 + \theta_4^{(1)} p'_1, & \theta_1^{(1)} + \theta_2^{(1)} + \theta_3^{(1)} + \theta_4^{(1)} &= 1 \\ p'_4 &= \theta_1^{(2)} p_1 + \theta_2^{(2)} p_2 + \theta_3^{(2)} p_3 + \theta_4^{(2)} p'_1, & \theta_1^{(2)} + \theta_2^{(2)} + \theta_3^{(2)} + \theta_4^{(2)} &= 1 \end{aligned}$$





- (5) The number 4 weights are free.
- (6) The number 5 weights are determined by  $C^1$  conditions (2.4).
- (7) The number 6 weights are free.
- (8) The number 7 weights are determined by  $C^1$  conditions (2.6).

The remaining weights with index  $\lambda_1\lambda_2\lambda_3\lambda_4$  are determined by  $C^1$  condition (2.4) for  $\lambda_4 \leq 1$  and freely chosen for  $\lambda_4 > 1$ .

### $C^2$ Quintic Scheme

(1) The number 0 weights(see Figure 4.2) are given by the function values at the vertices. For examples,  $a_{5e_i}^{(1)} = f(p_i)$ ,  $i = 1, 2, 3$ .

- (2) The number 1 weights are determined by formula (2.1).
- (3) The number 2 weights are determined by formula (2.2).
- (4) The number 3 weights, that is  $a_{1220}^{(i)}$ ,  $a_{2210}^{(i)}$  and  $a_{2120}^{(i)}$ , are free.
- (5) The number 4 weights are determined by  $C^1$  conditions (2.4), that is

$$a_{0221}^{(i)} = \theta_1^{(i)} a_{1220}^{(1)} + \theta_2^{(i)} a_{0320}^{(i)} + \theta_3^{(i)} a_{0230}^{(i)} + \theta_4^{(i)} a_{1220}^{(2)}$$

$$b_{1220}^{(1)} = \mu_1 a_{0221}^{(1)} + \mu_2 a_{0221}^{(2)} + \mu_3 a_{0320}^{(1)} + \mu_4 a_{0230}^{(1)}$$

(6) The number 5 and 6 weights have to be determined simultaneously. In determining these weights, we need to consider all the  $C^1$  and  $C^2$  conditions related to the tetrahedra surrounding the vertex  $p_2$ . Suppose there are  $k$  triangles(hence  $k$  edges) around  $p_2$ , then by  $C^1$  and  $C^2$  conditions, we have  $6k$  equations. That is, crossing each face, we have two equations. The number of related unknowns is also  $6k$ . That is,  $k$  number 5 weights and  $5k$  number 6 weights. Now we investigate these equations. It follows from (2.4) and (2.5) that

$$b_{1211}^{(i)} = \beta_1^{(i)} a_{1211}^{(i)} + \beta_2^{(i)} a_{0311}^{(i)} + \beta_3^{(i)} a_{0221}^{(i)} + \beta_4^{(i)} a_{0212}^{(i)} \quad (4.2)$$

$$b_{2210}^{(i)} = \beta_1^{(i)} \beta_1^{(i)} a_{2210}^{(i)} + 2\beta_1^{(i)} \beta_2^{(i)} a_{1310}^{(i)} + 2\beta_1^{(i)} \beta_3^{(i)} a_{1220}^{(i)} + 2\beta_1^{(i)} \beta_4^{(i)} a_{1211}^{(i)} + \beta_2^{(i)} \beta_2^{(i)} a_{0410}^{(i)} \\ + 2\beta_2^{(i)} \beta_3^{(i)} a_{0320}^{(i)} + 2\beta_2^{(i)} \beta_4^{(i)} a_{0311}^{(i)} + \beta_3^{(i)} \beta_3^{(i)} a_{0230}^{(i)} + 2\beta_3^{(i)} \beta_4^{(i)} a_{0221}^{(i)} + \beta_4^{(i)} \beta_4^{(i)} a_{0212}^{(i)} \quad (4.3)$$

for  $i = 1, 2$ . (4.2) and (4.3) can be written briefly as

$$b_{1211}^{(i)} = \beta_1^{(i)} a_{1211}^{(i)} + \beta_4^{(i)} a_{0212}^{(i)} + \gamma_0^{(i)} \quad (4.4)$$

$$b_{2210}^{(i)} = 2\beta_1^{(i)} \beta_4^{(i)} a_{1211}^{(i)} + \beta_4^{(i)} \beta_4^{(i)} a_{0212}^{(i)} + \gamma_1^{(i)} \quad (4.5)$$

where  $\gamma_0^{(i)}$  and  $\gamma_1^{(i)}$  are the known terms in (4.2) and (4.3). Since (see (2.6) and (2.7) )

$$b_{2210}^{(1)} = \mu_1 b_{1211}^{(1)} + \mu_2 b_{1211}^{(2)} + \gamma_2 \quad (4.6)$$

$$\mu_1^2 b_{0212}^{(1)} - \mu_1 b_{1211}^{(1)} = \mu_2^2 b_{0212}^{(2)} - \mu_2 b_{1211}^{(2)} + \gamma_3 \quad (4.7)$$

where  $\gamma_2 = \mu_3 b_{1310}^{(i)} + \mu_4 b_{1220}^{(i)}$  and  $\gamma_3 = \mu_2(\mu_3 b_{0311}^{(2)} + \mu_4 b_{0221}^{(2)}) - \mu_1(\mu_3 b_{0311}^{(1)} + \mu_4 b_{0221}^{(1)})$ , then by substituting (4.4) into (4.6) and (4.7) and then eliminating  $b_{2210}^{(i)}$  from (4.5) and (4.6) we get three equations related to four unknowns which could be written as:

$$= - \begin{bmatrix} \beta_4^{(1)} - \mu_1 & -\mu_2 \\ -\mu_1 & \beta_4^{(2)} - \mu_2 \end{bmatrix} \begin{bmatrix} \beta_4^{(1)} & 0 \\ 0 & \beta_4^{(2)} \end{bmatrix} \begin{bmatrix} a_{0212}^{(1)} \\ a_{0212}^{(2)} \end{bmatrix} \\ = - \begin{bmatrix} 2\beta_4^{(1)} - \mu_1 & -\mu_2 \\ -\mu_1 & 2\beta_4^{(2)} - \mu_2 \end{bmatrix} \begin{bmatrix} \beta_1^{(1)} & 0 \\ 0 & \beta_1^{(2)} \end{bmatrix} \begin{bmatrix} a_{1211}^{(1)} \\ a_{1211}^{(2)} \end{bmatrix} + \begin{bmatrix} \gamma_4^{(1)} \\ \gamma_4^{(2)} \end{bmatrix} \quad (4.8)$$

$$[-\mu_1(\beta_4^{(1)} - \mu_1) \quad \mu_2(\beta_4^{(2)} - \mu_2)] \begin{bmatrix} a_{0212}^{(1)} \\ a_{0212}^{(2)} \end{bmatrix} - [\mu_1\beta_1^{(1)}, -\mu_2\beta_1^{(2)}] \begin{bmatrix} a_{1211}^{(1)} \\ a_{1211}^{(2)} \end{bmatrix} = \gamma_5 \quad (4.9)$$

where  $\gamma_4^{(1)} = \mu_1\gamma_0^{(1)} + \mu_2\gamma_0^{(2)} + \gamma_2 - \gamma_1^{(1)}$ ,  $\gamma_4^{(2)} = \mu_1\gamma_0^{(1)} + \mu_2\gamma_0^{(2)} + \gamma_2 - \gamma_1^{(2)}$ , and  $\gamma_5 = \gamma_3 + \mu_1\gamma_0^{(1)} - \mu_2\gamma_0^{(2)}$ . Since the coefficient matrix of (4.8) is nonsingular, by solving  $[a_{0212}^{(1)} \ a_{0212}^{(2)}]^T$  from (4.8) and then substituting it into (4.9), we get one equation relating to the unknowns  $a_{1211}^{(1)}$ ,  $a_{1211}^{(2)}$ . Let the equation be in the form

$$\phi_i a_{1211}^{(1)} + \psi_i a_{1211}^{(2)} = \omega_i \quad (4.10)$$

Then, these unknowns form a closed chain around the vertex  $p_2$ . The coefficient matrix of all these equations related to the vertex  $p_2$  is in the form of

$$A = \begin{bmatrix} \phi_1 & \psi_1 & & \\ & \phi_2 & \psi_2 & \\ & & \ddots & \\ \psi_k & & & \phi_k \end{bmatrix}$$

The system (4.10) is a solvable in general with one degree of freedom. That is the rank of matrix  $A$  is  $k - 1$ . Hence the system can be solved. However, if the surrounding tetrahedra

at the same side at  $p_2$  are not closed, the matrix  $A$  is in the form of  $A = \begin{bmatrix} \phi_1 & \psi_1 & & \\ & \ddots & \ddots & \\ & & \phi_k & \psi_k \end{bmatrix}$

which can be changed to  $A = \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$  if one of the unknowns, say the  $l$ -th is chosen to be a free parameter. Hence the system of equations can be decomposed into two sub-systems. Each of the sub-systems can be easily solved.

(7) The number 7 weights are similarly determined as that of number 6.

(8) The number 8 weight  $a_{1112}^{(i)}$  are free.

(9) The number 9 weights are determined by the  $C^1$  and  $C^2$  conditions. Both the number of equations and the number of unknowns are  $6k$ . That is for  $i = 1, 2$

$$b_{1202}^{(i)} = \beta_1^{(i)} a_{1202}^{(i)} + \beta_2^{(i)} a_{0302}^{(i)} + \beta_3^{(i)} a_{0212}^{(i)} + \beta_4^{(i)} a_{0203}^{(i)} \quad (4.11)$$

$$b_{2201}^{(i)} = \beta_1^{(i)} \beta_1^{(i)} a_{2201}^{(i)} + 2\beta_1^{(i)} \beta_2^{(i)} a_{1301}^{(i)} + 2\beta_1^{(i)} \beta_3^{(i)} a_{1211}^{(i)} + 2\beta_1^{(i)} \beta_4^{(i)} a_{1202}^{(i)} + \beta_2^{(i)} \beta_2^{(i)} a_{0401}^{(i)} \\ + 2\beta_2^{(i)} \beta_3^{(i)} a_{0311}^{(i)} + 2\beta_2^{(i)} \beta_4^{(i)} a_{0302}^{(i)} + \beta_3^{(i)} \beta_3^{(i)} a_{0221}^{(i)} + 2\beta_3^{(i)} \beta_4^{(i)} a_{0212}^{(i)} + \beta_4^{(i)} \beta_4^{(i)} a_{0203}^{(i)} \quad (4.12)$$

$$b_{3200}^{(1)} = \mu_1 b_{2201}^{(1)} + \mu_2 b_{2201}^{(2)} + \gamma_6 \quad (4.13)$$

$$\mu_1^2 b_{1202}^{(1)} - \mu_1 b_{2201}^{(1)} = \mu_2^2 b_{1202}^{(2)} - \mu_2 b_{2201}^{(2)} + \gamma_7 \quad (4.14)$$

where  $\gamma_6 = \mu_3 b_{2300}^{(i)} + \mu_4 b_{2210}^{(i)}$  and  $\gamma_7 = \mu_2(\mu_3 b_{1301}^{(2)} + \mu_4 b_{1211}^{(2)}) - \mu_1(\mu_3 b_{1301}^{(1)} + \mu_4 b_{1211}^{(1)})$ . Substitute (4.11) and (4.12) into (4.14), so that we have

$$\mu_1 \beta_4^{(1)} (\mu_1 - \beta_4^{(1)}) b_{0203}^{(1)} - \mu_2 \beta_4^{(2)} (\mu_2 - \beta_4^{(2)}) b_{0203}^{(2)} = \dots$$

This is a system that is in the same form as (4.10). The coefficient matrix of this system is nonsingular, in general.

**(10)** For the number 10 weights, we have six equations parallel to the equations (4.11)–(4.14) with all the indices changed by the rule:

$$\text{The index of the number 10 weight} = \text{The index of the number 9 weight} - e_2 + e_3$$

and seven independent weights. By choosing one of them, say  $b_{3110}^{(i)}$ , to be a free parameter, the system can be solved.

**(11)** The number 11 weights are determined in the same way as the number 9.

**(12)** The number 12 and 13 weights are free, while the number 14 are determined by  $C^1$  and  $C^2$  conditions. That is  $b_{1103}^{(i)}$  are defined by (2.4).  $b_{2102}^{(i)}$  are defined by (2.5). For  $b_{3101}^{(i)}$ , we have by (2.6) and (2.7) that

$$\mu_1 b_{3101}^{(1)} + \mu_2 b_{3101}^{(2)} = b_{4100}^{(1)} + \gamma_8, \quad -\mu_1 b_{3101}^{(1)} + \mu_2 b_{3101}^{(2)} = \mu_2^2 b_{2102}^{(2)} - \mu_1^2 b_{2102}^{(2)} + \gamma_9$$

where  $\gamma_8 = -\mu_3 b_{3200}^{(i)} - \mu_4 b_{3110}^{(i)}$  and  $\gamma_9 = \mu_2(\mu_3 b_{2201}^{(2)} + \mu_4 b_{2111}^{(2)}) - \mu_1(\mu_3 b_{2201}^{(1)} + \mu_4 b_{2111}^{(1)})$ .

$$b_{3101}^{(1)} = \frac{b_{4100}^{(1)} - \mu_2^2 b_{2102}^{(2)} + \mu_1^2 b_{2102}^{(2)} + \gamma_8 - \gamma_9}{2\mu_1}, \quad b_{3101}^{(2)} = \frac{b_{4100}^{(1)} + \mu_2^2 b_{2102}^{(2)} - \mu_1^2 b_{2102}^{(2)} + \gamma_8 + \gamma_9}{2\mu_1}$$

**(13)** The number 15 weights are similar to that of number 14, the index being changed by the same rule as above.

**(14)** The number 16 weights are free, the number 17's are determined by  $C^1$  and  $C^2$  conditions.

**(15)** The number 0 to number 8 weights of the lower tetrahedra, below faces of  $T$  (see Figure 4.2) are determined by  $C^0$ ,  $C^1$  and  $C^2$  conditions (2.3), (2.4) and (2.5) from weights in the upper tetrahedron.

**16** The number 9 to 17 weights of the lower tetrahedra are determined in a fashion similar to the  $C^0$ ,  $C^1$  and  $C^2$  conditions between the face and edge tetrahedra.

In summary, the construction steps **1–14** and **16** is according to the  $C^0$ ,  $C^1$  and  $C^2$  conditions across the common faces between face and edge tetrahedra that are both above or both below the original triangulation  $T$ . Step **15** is according to the  $C^0$ ,  $C^1$  and  $C^2$  conditions across the

faces of  $T$  and between the upper and lower tetrahedra. Therefore, the composite function is global  $C^2$  continuous in  $\Sigma$ .

### The Use of Free Weights

In both of the  $C^1$  and  $C^2$  schemes described above, there are some free weights which can be freely determined to control the local geometry of  $F$  without affecting the continuity. We suggest three approaches or their combinations to achieve this local control. The first is to modify the shape of  $F$  by interactively adjusting the free weights. The second is to locally interpolate some of the function-on-surface data earlier approximated by the polynomial in each tetrahedron. The third approach is to least-square approximate some additional lower degree polynomial (acting as a controlling function) by use of the degree elevation formula of §2. For example, in the  $C^1$  scheme, the number 2 weights can be determined by

$$a_{1110}^{(i)} = \frac{1}{4}(a_{1200}^{(i)} + a_{2100}^{(i)} + a_{2010}^{(i)} + a_{1020}^{(i)} + a_{0210}^{(i)} + a_{0120}^{(i)}) - \frac{1}{6}(a_{3000}^{(i)} + a_{0300}^{(i)} + a_{0030}^{(i)})$$

and the number 4 weights are determined by

$$a_{0003}^{(i)} = \frac{1}{3}[2(q_{0101}^{(i)} + q_{1001}^{(i)} + q_{0011}^{(i)}) - (a_{0300}^{(i)} + a_{3000}^{(i)} + a_{0030}^{(i)})]$$

$$a_{0102}^{(i)} = \frac{1}{3}(2q_{0101}^{(i)} + a_{0003}^{(i)}), \quad a_{1002}^{(i)} = \frac{1}{3}(2q_{1001}^{(i)} + a_{0003}^{(i)}), \quad a_{0012}^{(i)} = \frac{1}{3}(2q_{0011}^{(i)} + a_{0003}^{(i)})$$

where

$$q_{0101}^{(i)} = \frac{3}{4}(a_{1101}^{(i)} - a_{1011}^{(i)} + a_{0111}^{(i)} + a_{0201}^{(i)}) - \frac{1}{4}(q_{1100}^{(i)} - q_{1010}^{(i)} + q_{0110}^{(i)} + a_{0300}^{(i)})$$

$$q_{1001}^{(i)} = \frac{3}{4}(a_{1101}^{(i)} + a_{1011}^{(i)} - a_{0111}^{(i)} + a_{2001}^{(i)}) - \frac{1}{4}(q_{1100}^{(i)} + q_{1010}^{(i)} - q_{0110}^{(i)} + a_{3000}^{(i)})$$

$$q_{0011}^{(i)} = \frac{3}{4}(-a_{1101}^{(i)} + a_{1011}^{(i)} + a_{0111}^{(i)} + a_{0021}^{(i)}) - \frac{1}{4}(-q_{1100}^{(i)} + q_{1010}^{(i)} + q_{0110}^{(i)} + a_{0030}^{(i)})$$

$$q_{1100}^{(i)} = \frac{1}{4}(3a_{1200}^{(i)} + 3a_{2100}^{(i)} - a_{0300}^{(i)} - a_{3000}^{(i)})$$

$$q_{1010}^{(i)} = \frac{1}{4}(3a_{2010}^{(i)} + 3a_{1020}^{(i)} - a_{0030}^{(i)} - a_{3000}^{(i)})$$

$$q_{0110}^{(i)} = \frac{1}{4}(3a_{0210}^{(i)} + 3a_{0120}^{(i)} - a_{0300}^{(i)} - a_{0030}^{(i)})$$

## 5 Visualization and Examples

We can visualize the graph of the constructed function  $F$  on the domain surface  $D$  either by projecting the iso-contours onto the surface  $D$ , or by directly displaying iso-contours or the surface graph of the function  $F$  in space.

### Displaying Iso-contours of $F$ on $D$




Figure 5.1: Iso-contours of a  $C^1$  approximated function  $F$  shown on a domain torus  $D$




Figure 5.2: Iso-contours of a  $C^2$  approximated function  $F$  shown on and surrounding a domain torus  $D$  using a normal projection

We display the iso-contours on the domain surface by showing different colors in the region between two iso-contours. In our approach, we achieve this by first generating a planar triangular approximation of the domain surface, and then generating the corresponding four dimensional triangles on  $F$ , and finally intersecting these triangles with the iso-values to get the line segments of the iso-contours. Let  $w$  be a given iso-value,  $[p_1 p_2 p_3]$  be a triangle on  $D$ . Without loss of generality, we may assume  $F(p_1) \leq F(p_2) \leq F(p_3)$ . Then if  $w < F(p_1)$  or  $w > F(p_3)$ , the triangle does not intersect the iso-value. If  $w \in [F(p_1), F(p_3)]$ , say  $w \in [F(p_1), F(p_2)]$ , let  $t_1 = \frac{w-F(p_1)}{F(p_2)-F(p_1)}$ ,  $t_2 = \frac{w-F(p_1)}{F(p_3)-F(p_1)}$ ,  $q_1 = t_1 p_1 + (1-t_1)p_2$ ,  $q_2 = t_2 p_1 + (1-t_2)p_3$ , then  $[q_1 q_2]$  is one segment of the contour  $F(p) = w$ . The collection of all of these line segments form a piecewise approximation to the iso-contours. By increasing the resolution of the triangulation of the domain surface, we can get better approximations of the iso-contours. Figure 5.1 (left and right) shows the iso-contours of a  $C^1$  approximated function  $F$ , on a domain torus  $D$ . Figure 5.2 (left and right) shows the iso-contours of a  $C^2$  approximated function  $F$ , on a domain torus  $D$ . The iso-contours of the  $C^2$  approximated function  $F$  are also shown surrounding the domain torus using the normal projection scheme given below.

### Displaying Iso-contours and the graph of $F$ in $\mathbb{R}^3$

Since the iso-contours may not clearly indicate the geometric shape of the function-on-surface, one often plot the function-on-surface in one way or another. One approach is to use a radial projection from some center of the domain. However, if the domain surface is not convex or has non-zero genus, this projection scheme has difficulties caused by self-intersection. Another more natural way is to use the normal projection, that is, project the point  $p$  on the domain surface  $D$  to a distance proportional to  $F(p)$  in the normal direction of  $D$  at  $p$ :  $G(p) = p + L \frac{\nabla f(p)(F(p)-F_{min})}{\|\nabla f(p)\|(F_{max}-F_{min})}$  where  $L$  is a positive scalar,  $F_{min}$  and  $F_{max}$  are minimum and maximum values of  $F$  on  $D$ . Here  $L$  has to be chosen properly so that the projected surface  $G$  does not self-intersect.

Figures 5.3, 5.4, (left and right) shows the iso-contours of a  $C^2$  approximated function  $F$ , on a domain  $D$ . The iso-contours of the  $C^2$  approximated function  $F$  are also shown surrounding the domain using the normal projection scheme.

## References

- [1] P. Alfeld. A trivariate clough-tocher scheme for tetrahedral data. *Computer Aided Geometric Design*, 1:169–181, 1984.
- [2] C. Bajaj and G. Xu. *The Estimation of the Gradients and Hessian for Functions defined on Surfaces*. Computer Science Technical Report, Purdue University, 1994.
- [3] R.E. Barnhill and T.A. Foley. Methods for constructing surfaces on surfaces. In G.Farin, editor, *Geometric Modeling: Methods and their Applications*, pages 1–15. Springer, Berlin, 1991.
- [4] R.E. Barnhill, K. Opitz, and H. Pottmann. Fat surfaces: a trivariate approach to triangle-based interpolation on surfaces. *Computer Aided Geometric Design*, 9:365–378, 1992.




Figure 5.3: Iso-contours of a  $C^2$  approximated function  $F_1$  shown on and surrounding a domain surface  $D_1$ .




Figure 5.4: The surface and iso-contours in  $\mathbf{R}^3$  of the  $C^2$  approximated function  $F_1$  surrounding the domain  $D_1$ . These are a normal projection from the domain  $D_1$



- [5] L. P. Chew. Guaranteed-quality mesh generation for curved surface. In *9th Annual Computational Geometry*, pages 274–280, CA,USA, May 1993.
- [6] G. Farin. *Curves and Surfaces for Computer Aided Geometric Design: A Practical Guide*. Academic Press Inc., 1990.
- [7] R. Franke. Recent advances in the approximation of surfaces from scattered data. In C.K.Chui, L.L.Schumaker, and F.I.Utreras, editors, *Multivariate Approximation*, pages 275–335. Academic Press, New York, 1987.
- [8] G. Nielson, T. Foley, B. Hamann, and D. Lane. Visualizing and Modeling Scattered Multivariate Data. *IEEE Computer Graphics And Applications*, 11:47–55, 1991.
- [9] K.L. Rescorla.  $C^1$  Trivariate Polynomial Interpolation. *Computer Aided Geometric Design*, 4:237–244, 1987.
- [10] A. J. Worsey and G. Farin. An n-dimensional clough-tocher interpolant. *Constructive Approximation*, 3:99–110, 1987.