

LETTER • OPEN ACCESS

Modelling public health improvements as a result of air pollution control policies in the UK over four decades—1970 to 2010

To cite this article: E Carnell *et al* 2019 *Environ. Res. Lett.* **14** 074001

View the [article online](#) for updates and enhancements.

You may also like

- [Scoping review to understand the potential for public health impacts of transitioning to lower carbon emission technologies and policies](#)
Rachel Tham, Geoff Morgan, Shyamali Dharmage *et al.*
- [Hazardous air pollutant exposure as a contributing factor to COVID-19 mortality in the United States](#)
Michael Petroni, Dustin Hill, Lylla Younes *et al.*
- [Examining the impact of inter-provincial migration on environmental health in China](#)
Yu-Ou Yang, Juan-Juan Hou, Xin-Yu Chen *et al.*

Environmental Research Letters



LETTER

Modelling public health improvements as a result of air pollution control policies in the UK over four decades—1970 to 2010

OPEN ACCESS

RECEIVED

8 January 2019

REVISED

29 March 2019

ACCEPTED FOR PUBLICATION

2 April 2019

PUBLISHED

26 June 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



E Carnell¹ , M Vieno¹ , S Vardoulakis^{2,3} , R Beck¹, C Heaviside^{4,5} , S Tomlinson¹ , U Dragosits¹ , MR Heal⁶ and S Reis^{1,3,7}

¹ Centre for Ecology & Hydrology, Bush Estate, Penicuik, EH26 0QB, United Kingdom

² Institute of Occupational Medicine, Research Avenue North, Edinburgh, EH14 4AP, United Kingdom

³ University of Exeter Medical School, Knowledge Spa, Truro, TR1 3HD, United Kingdom

⁴ Environmental Change Institute, University of Oxford, South Parks Road, Oxford, OX1 3QY, United Kingdom

⁵ School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁶ School of Chemistry, University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh, EH9 3FJ, United Kingdom

⁷ Author to whom any correspondence should be addressed.

E-mail: srei@ceh.ac.uk

Keywords: air pollution, long-term trends, attributable mortality, respiratory health

Supplementary material for this article is available [online](#)

Abstract

In much of the industrialised world, policy interventions to address the challenges of wide-spread air pollution as resulting from development and economic progress in the 2nd half of the 20th century have overall led to reductions in air pollution levels and related health effects since the 1970s. While overall improvements towards reducing health effects from ambient air pollution are recorded, comprehensive and consistent assessments of the long-term impact of policy interventions are still scarce. In this paper, we conduct a model assessment over a 40 year period of air pollution in the UK. In order to correct for the short and longer term variability of meteorological factors contributing to trends in ambient concentrations of priority air pollutants (nitrogen dioxide, sulphur dioxide, fine particulate matter and ozone), we use a fixed meteorological year for all model simulations. Hence, the modelled changes in air pollutant concentrations and related health effects are solely a function of the changes in emissions since 1970. These changes in emissions are primarily driven by policy interventions, ranging from phasing out of specific fuels or substances, to regulating the use of chemicals and driving the development of cleaner, more efficient technologies. Over the 40 year period, UK attributable mortality due to exposure to PM_{2.5} and NO₂ have declined by 56% and 44% respectively, while ozone attributable respiratory mortality increased by 17% over the same period (however, with a slight decrease by 14% between 2000 and 2010).

1. Introduction

1.1. Rationale for this paper

Outdoor air pollution presents a major challenge for public health and ranks among the main causes of adverse health effects globally (Lim *et al* 2012, Cohen *et al* 2017, Landrigan *et al* 2017). It is estimated to account for about 4.2 million deaths and to rank 5th worldwide among all risk factors (HEI 2017). Long-term exposure to air pollution reduces life expectancy, mainly due to an increased risk of cardiovascular and respiratory disease and lung cancer, while short-term

exposure can as well aggravate respiratory and cardiovascular illness and trigger asthma attacks (RCP 2016). The World Health Organization (WHO) has classified outdoor air pollution as carcinogenic (Loomis *et al* 2013), and there is emerging evidence of possible links of air pollution with diabetes, obesity, cognitive decline, dementia, and adverse birth outcomes (RCP 2016).

In most industrialised countries, concentrations of air pollutants substantially increased in the first half of the 20th century because of the rise of fossil fuel combustion to satisfy the fast-growing needs for energy and mobility. This was in particular the case for sulphur

dioxide (SO₂), nitrogen dioxide (NO₂), carbon monoxide (CO), particulate matter (both in the form of 'soot', and as defined by PM_{2.5}, the mass concentration of particles with an aerodynamic diameter of <2.5 μm) and non-methane volatile organic compounds (NMVOCs). During recent decades, however, increasingly stringent air pollution control policies, and improved fuels and pollution abatement technology (e.g. introduction of three-way catalytic converters in cars), have generally resulted in declines in emissions in high-income countries. Consequently, there has been a decline in the concentrations of these air pollutants in most high-income countries. However, for some pollutants the situation is more complex. For example, emissions of NO_x and NO₂ concentrations did rise with the increase of road transport, until catalytic converters and increasingly stringent emissions standards led to a decoupling of emission trends from transport development. In parallel, concentrations of ground-level ozone (O₃), a secondary pollutant formed from chemical reactions between NO_x and NMVOCs, have continued rising in many areas. Also, emissions of ammonia (NH₃), mainly originating from agriculture, have not been declining substantially and more recently showed signs of a slight increase. NH₃ is a precursor to the formation of secondary inorganic aerosols (SIA), a major component of PM_{2.5} concentrations.

Long-term trends of air pollution and its effects on human and ecosystem health are usually assessed through analysis of time series of measurements collected through regulatory monitoring networks. However, robust and reliable monitoring data have only been available since the 1970s for some air pollutants, and as late as the 1990s for others. In addition, monitoring networks are typically sparse, which is a challenge when using monitoring sites to represent population exposure (Willocks *et al* 2012). The emergence of earth observation products provides wider spatial coverage (van Donkelaar *et al* 2015, Pope *et al* 2018), but observational air quality data, whether ground-based or from remote sensing can only reveal the combined effect of changes in emissions, land use, and meteorological conditions on the formation, transport and environmental fate of pollutants. Atmospheric chemistry transport models (ACTMs) explicitly consider all of these processes, and can be used to forecast as well as to reconstruct historic scenarios. Most model assessments aim to deliver as realistic as possible representations of air pollutant concentrations for the quantification of health effects, including comparison with observations for evaluation purposes. In many cases, complex steps towards data integration are required, as exemplified by the Global Burden of Disease study (Shaddick *et al* 2018). As a consequence, it is a challenge to disentangle the size of the effect of emissions reductions (as a result of policy implementation) from the inter-annual variability due to meteorological variability (Pope *et al* 2014, Huang *et al* 2017). It is information on the effects of emissions

reductions on long-term air pollution trends that gives insight on effectiveness of past policies and helps inform future policy development, as for instance presented by EPA (2011), Matus *et al* (2008, 2012) and Zhang *et al* (2018) for the USA and China, as well as WHO (2013a, 2013b) with a focus on Europe. Hennehan *et al* (2016) have made a case for the evaluation of policy effectiveness in their review and developed frameworks for accountability studies more generally.

To address this evidence gap, we modelled the effect of four decades of emission reductions across Europe on UK annual mean concentrations of regulated air pollutants and their health impacts at ten-year time intervals of 1970, 1980, 1990, 2000, and 2010. In order to focus on the impact of emission changes brought about by policy measures, all model runs were conducted using the same meteorological input data, varying only the emissions using available historic emission inventory data. To our knowledge, this presents a novel approach and yields a unique insight into the effectiveness of historic emission control policies, and provides a basis for designing ex-ante assessments of future policy interventions.

1.2. AQ trends and policies—a UK perspective

Although Public Health Acts in 1875 and 1926 contained measures to abate emissions of smoke, the first major modern legislation in the UK to focus on mitigation of the harmful effects of air pollution was the 1956 Clean Air Act, introduced in response to the notorious sulphurous 'smog' episodes in London and elsewhere (Brimblecombe 1987, Malley *et al* 2016) and their impacts on morbidity and mortality (Anderson 2009). Subsequent concern on the effects of air pollution was then predominantly stimulated by the observation during the 1970s of forest dieback ('Waldsterben') and landscape acidification across much of Europe. The implementation of Europe-wide emission control policies therefore initially targeted SO₂ emissions from large combustion plants and resulted in a de-coupling of continuing economic growth and SO₂ emissions, with the latter showing a very significant downward trend since the 1980s (Vestreng *et al* 2007). From that time, substantial evidence has also accumulated for wide-spread public health effects associated with emissions of nitrogen oxides (NO_x), including their contribution, in combination with emissions of NMVOCs, to the formation of O₃, and to the formation of SIA. Consequently, from the 1990s onwards a range of additional pan-European emission control policies (see below) have led to reductions of NO_x and NMVOC (and primary PM) emissions (Vestreng *et al* 2009).

The current UK approach to air quality management is outlined in its National Air Quality Strategy, first developed in 1997, as an outcome of the 1995 Environment Act. Air quality standards regulations set into UK law objectives for the concentrations of a

number of air pollutants that are comparable to (or stricter than) those set out in relevant European Union Ambient Air Quality Directives, the most recent of which is Directive 2008/50/EC which replaced a number of previous Directives. Air pollutants with UK air quality objectives include PM₁₀, PM_{2.5}, NO₂, O₃, SO₂, CO, benzene, 1,3-butadiene, polycyclic aromatic hydrocarbons (PAHs) and a number of metals. These standards are based on expert-group considerations of the epidemiological and toxicological evidence for the harmful effects of these pollutants on human health.

A component part of the UK Air Quality Strategy is Local Air Quality Management whereby local authorities are required to assess the air quality in their area. Areas where air quality objectives are not likely to be achieved must be designated as air quality management areas and an action plan to improve the air quality developed. At the national scale, the UK abides by EU Directives and international protocols that aim to reduce exposure to air pollutants through regulation of pollutant emissions from individual point or sector sources, and through a cap on total national emissions of primary pollutants. Examples of the former regulations include emissions from combustion plants, road and off-road vehicles and machinery, and industrial solvent usage, whilst an example of the latter is the National Emissions Ceilings Directive. Actions taken under these regulations have led to substantial reductions in the largest ambient concentrations of most of the pollutants with air quality objectives, particularly the primary pollutants such as organic compounds, metals, CO and SO₂. No local air quality management areas in the UK are currently declared due to concern over meeting air quality objectives for these pollutants.

Instead, the two pollutants currently causing the greatest concern in Europe, and for which UK air quality management areas have been declared, are PM_{2.5} and NO₂ (COMEAP 2009, 2010, WHO 2013a, 2013b, RCP 2016). The corresponding objectives are, for PM_{2.5}, an annual mean of 25 $\mu\text{g m}^{-3}$ and a target to reduce population-weighted average concentrations by 15% between 2010 and 2020, and, for NO₂, an annual mean of 40 $\mu\text{g m}^{-3}$ and an hourly mean of 200 $\mu\text{g m}^{-3}$ not to be exceeded for more than 18 h per year. (The objective for PM_{2.5} is more stringent in Scotland than in the rest of the UK.) The UK Committee on the Medical Effects of Air Pollutants (COMEAP) have recently emphasised the strengthening evidence for mortality effects from long-term (i.e. annual average) exposure to NO₂ (COMEAP 2018). NO₂ concentrations in many urban areas are declining only slowly because road traffic volume is increasing and the share of diesel vehicles in the UK fleet has been increasing over the past decades. The same expert group has also reported on the health effects from exposure to O₃ (COMEAP 2015), which is of particular note given the currently rising levels of urban O₃ (where the bulk of the population are exposed) due to the slowly

decreasing urban NO_x emissions. The exposure reduction target for PM_{2.5} marks the first appearance in legislation of an aim to improve overall population health burden rather than solely to attain a particular target concentration.


1.3. An overview of this study

In this paper, we focus on NO₂, PM_{2.5}, SO₂ and O₃ as pollutants which have been in the spotlight due to their large environmental and public health effects over time. For NO₂ and PM_{2.5}, exceedances of current air quality objectives, respectively of WHO guideline values, remain major policy challenges. The considerably higher than pre-industrial level of O₃ is also a concern for public health professionals (WHO 2013a, 2013b, COMEAP 2015). Sulphur dioxide was previously a pollutant of major concern. The spatially-resolved modelled pollutant concentrations produced for this paper reflect the four decades of changes in anthropogenic emissions only (i.e. natural sources such as volcanoes were left unchanged for the purpose of the study); all other model parameters (including meteorology and land use) remain constant throughout the model time series. Linking these concentration maps with population data at each 10 year time step to calculate population weighted concentrations, we assess the changes in public health impact due to the change in emissions and concentrations over this time. As the model calculations are based on a constant meteorology year (2015), the health impacts do not match those estimated in any of the specific years. However, this approach allows for a consistent evaluation of the effects of emission reductions on changes to human exposure and health burden over a 40 year period via the effect of those emissions reductions on ambient concentrations of specific air pollutants.

2. Methods

2.1. Emissions data

In order to assess the effect of policies to reduce air pollution, the model required a consistent set of emission data for the whole period. For the UK, anthropogenic emission data as compiled by the National Atmospheric Emissions Inventory (NAEI 2018) were obtained for each of the modelled years (1970, 1980, 1990, 2000 and 2010) based on the 2015 NAEI release (latest available inventory year at the time of this work). The anthropogenic emissions include domestic aviation (i.e. landing and take-off cycles, as well as airport operations) and shipping, but emissions from natural phenomena (wild fires, dust events, volcanoes etc), NO_x releases from soil, and biological/natural VOC were set at the 2015 NAEI values. For non-UK emissions, Centre for Emission Inventories and Projections (CEIP 2018) data was used for each corresponding year. For the UK, the 2012 spatial distribution was used and rescaled with the respective national total



emissions for each year because historic spatial distributions of national emissions are not available once recalculated national and sectoral total emissions are released, and at the time of conducting the model runs, was the most up-to-date version available. Since these model calculations were undertaken, updated UK and European emission datasets have been released, with some differences between the emissions used in this paper and recalculated emissions for previous years. This is inevitable, as new findings and corrections of emission factors emerge, which are then used to backcast historic emission time series to provide consistent reporting under different national and international requirements. A key change to emission data is a result of corrections to emission factors of some road vehicles following the 2015 ‘Dieselgate’ scandal, where real world emissions from diesel vehicles were under-reported by manufacturers due to manipulated testing procedures (e.g. Brand 2016). However, as these recalculations will typically correct errors and are applied consistently to specific sources and sectors, the relative changes in emissions on which our model results are based should not be subject to substantive changes.

Figure 1 illustrates the changes in the national emission estimates of NH₃, SO₂, NO_x, NMVOC and primary PM_{2.5} at 10 year intervals between 1970 and 2010. SO₂ emissions have declined substantially during this period as a consequence of policy regulations aiming to address acid rain in the UK and Europe. Primary PM_{2.5} emissions have also been effectively reduced in this period. For NO_x emissions, the increase in road transport volume offset reductions in other sectors until the early 1990s, when implementation of increasingly stringent EURO vehicle emission standards yielded declines in total NO_x emissions. The NMVOC emissions are observed to have reduced only in the latter part of the time period and, overall, the relatively modest reductions in NMVOC and NO_x mean there is still substantial contribution to O₃

formation. Reductions in NH₃ emissions have been very small because very few regulations to date have addressed agricultural emissions—a key contributor to total NH₃ emissions in the UK and in Europe. Latest indications suggest a slight increase in NH₃ emissions (EEA 2018).

2.2. Atmospheric chemistry transport modelling

The ACTM EMEP4UK rv4.3 was used for this work and is described in Vieno *et al* (2016a and 2016b). The EMEP4UK model is a regional application of the EMEP MSC-W model (Simpson *et al* 2012). The EMEP MSC-W model rv4.3 is freely accessible at https://github.com/metno/emep-ctm/releases/tag/rv4_3. The meteorological data to drive the EMEP4UK ACTM are generated by the Weather and Research Forecast model (WRF). Both the ACTM and WRF model use the same model domain covering Europe at a horizontal resolution of 50 km × 50 km with a nested higher resolution for the UK at a horizontal resolution of 5 km × 5 km.

As described previously, only changes to anthropogenic emissions are incorporated into each historical simulation; all other model parameters, including the meteorological year (2015) for each simulation year, are kept constant. This ensures that changes to modelled concentrations reflect only the trends in anthropogenic emissions. While this does not allow for a direct validation of model outputs against observations, it provides a consistent method to assess the actual changes in ambient modelled concentrations as a function of changes in emission input data.

In-depth information about model setup and validation is provided in suppl. mat. Section S1 is available online at stacks.iop.org/ERL/14/074001/mmedia.

2.3. Health impact assessment

The attributable fraction (AF) of all-cause mortality in the UK associated with PM_{2.5}, NO₂ and O₃ (AF of respiratory mortality only for O₃) for each of the years

1970, 1980, 1990, 2000 and 2010 was calculated as

$$AF = [(RR - 1)/RR],$$

where RR (relative risk) = $\beta^{(x/10)}$, β is the concentration-response coefficient for each pollutant (per $10 \mu\text{g m}^{-3}$) and x is the population-weighted annual average pollutant concentration. We do not provide an estimate of changes in absolute mortality figures from air pollution, since this would also depend on changes in baseline mortality rates and population sizes over time.

The values of β for estimating the annual attributable mortality from long-term exposure to air pollution used here were based on the UK Committee on the Medical Effects of Air Pollution recommendations. Specifically, the β coefficient for $\text{PM}_{2.5}$ used here was 1.060 (95% CI: 1.04–1.08) (COMEAP 2010), and for NO_2 the value of 1.023 (95% CI: 1.008, 1.037) was used (COMEAP 2018). We assumed that there is no threshold for the mortality effects of these two pollutants (i.e. we quantified burdens to zero concentrations).

The mortality estimates presented here are based on quantification using either $\text{PM}_{2.5}$ or NO_2 as the primary indicator of the air pollution mixture. The size of the overlap between burdens derived from associations of mortality with NO_2 and $\text{PM}_{2.5}$ is unknown, therefore single-pollutant burden estimates, $\text{PM}_{2.5}$ or NO_2 in this case, should not be added together as doing so would probably lead to over-estimation of the effects of the air pollution mixture. For O_3 , despite some evidence for long-term effects, COMEAP do not recommend a β value (COMEAP 2015). Therefore the value used here is the 1.014 (95% CI = 1.005, 1.024) per $10 \mu\text{g m}^{-3}$ of the warm season (April–September) average of daily maximum 8 h mean O_3 concentration recommended by the WHO Health Risks of Air Pollution in Europe (HRAPIE) project for the effects of long-term exposure to O_3 on respiratory mortality, with a cut-off value of $70 \mu\text{g m}^{-3}$ (WHO 2013a), although we accept that evidence on the mortality effects associate with long-term exposure to O_3 is not conclusive (Atkinson *et al* 2016).

3. Results

3.1. Trends in air quality 1970–2010

Figure 2 illustrates the modelled concentrations of SO_2 , NO_2 , O_3 and $\text{PM}_{2.5}$ for the five 10 year snapshots in time. For SO_2 and $\text{PM}_{2.5}$, the concentration maps indicate a consistent downward trend over the whole period as observed for emissions of SO_2 and primary $\text{PM}_{2.5}$ (figure 1). For NO_2 , the increase in NO_x emissions due to a substantial increase in road transport emissions over the period between 1970 and 1990 is highlighted by a marked increase in NO_2 concentrations in urban areas and along the major UK transport corridors. After 1990, NO_2 concentrations decline across the whole of the UK. In contrast, concentrations of O_3 have seen a steady increase over the 40 year


period, with modest increases from 1970 to 1990 but more substantial increases thereafter in the areas where NO_x emissions (and consequently NO_2 concentrations) decrease most. The modelled trends for O_3 are the result of the combination of the emission changes in both NO_x and NMVOC, as well as changes in amounts of transboundary transport of O_3 . The increase in the second half of the study period indicates that in order to achieve a reduction in O_3 , further substantial reductions in NO_x (and NMVOC) emissions would be required across the UK, with O_3 formation being increasingly NO_x -limited in the UK (i.e. NO_x emission control will produce a greater reduction in O_3 concentrations, Monks *et al* 2015).

The spatial distribution of concentration changes primarily reflects the main population centres and transport corridors, where emission reductions are predominantly occurring. In the case of $\text{PM}_{2.5}$ concentration changes, the resulting reductions are a combination of UK and European emission reductions, as the formation of SIA (ammonium nitrates and sulphates) over continental Europe has declined due to Europe-wide emission controls over the same period. As a result, advection of secondary $\text{PM}_{2.5}$ to the UK has declined as well, contributing to lower $\text{PM}_{2.5}$ concentrations over the UK. For ground-level ozone, the increase in global background concentrations, in parallel to the reduced destruction of ozone by titration due to decreasing NO_2 concentrations act together in showing an increasing trend (Read *et al* 2008, Yan *et al* 2018).

Figure 3 illustrates the changes in average, 1st and 99th percentile modelled annual-average concentrations across all model land-based UK grid cells over the study period. For SO_2 , both high and average annual-average concentrations have substantially declined. High annual-average concentrations of $\text{PM}_{2.5}$ have also consistently declined, with low concentrations showing a slight decrease, and average concentrations having fallen to less than $10 \mu\text{g m}^{-3}$ over the period. While average SO_2 concentrations show by far the largest reduction at 94%, average $\text{PM}_{2.5}$ concentrations declined by 53%. The situation for NO_2 and O_3 is more complex, with an increase in 99th percentile annual-average NO_2 concentrations until 1990, followed by a steep decline amounting to a total of 48% reduction between 1970 and 2010. 1st and 99th percentiles and annual-average O_3 concentrations have increased throughout the period, with the average and high concentrations showing, respectively, a slowing increase and a slight decrease since 1990. Compared to 1970, minimum, average and maximum modelled concentrations of O_3 are 72%, 18% and 12% higher in 2010, respectively.

3.2. Trends in attributable mortality 1970–2010


Figure 4 shows the trends in spatial patterns and magnitudes of population-weighted concentrations



for SO_2 , NO_2 , $\text{PM}_{2.5}$ and O_3 for each 10 year time interval. This illustrates the spatial variability of emission reductions and resulting absolute changes in population weighted concentrations for each pollutant, which provides the base for the assessment of attributable mortality changes over the study period (see suppl. mat. section S2 for a detailed description of the population mapping method applied).

Estimates of the UK population-weighted average AF of all-cause mortality for long-term exposure to NO_2 and $\text{PM}_{2.5}$ (respiratory mortality for O_3) at each time point are shown in table 1 and figure 5. The trends in AF of mortality reflect the spatial integration

of the trends in population weighted concentrations shown in figure 4. Over the entire period considered (1970–2010), the population-weighted average AF of mortality for $\text{PM}_{2.5}$ fell by 56%, from 11.83% to 5.21%, and for NO_2 it fell by 44%, from 5.32% to 2.96% (table 1). Over the same period, the population-weighted average AF of respiratory mortality due to O_3 increased by 17%. However, the AF of mortality for O_3 and NO_2 effects rose slightly from 1970 to 1990, when they were 2.33% and 5.76%, respectively, and then fell by 2010, with the AF of respiratory mortality for O_3 still higher in 2010 than it was in 1970 (figure 5). In contrast, the population-weighted average AF of



mortality associated with PM_{2.5} has declined steadily each 10 year time point, from 11.83% in 1970 to 5.21% in 2010 in the UK.


4. Discussion

4.1. Implications for policy development

The overarching aim of this study was to simulate how pollutant concentrations, together with an indicator of associated health impacts (AF of mortality), have changed in the UK over the 40 year period 1970–2010 in response to the effect of air quality policies on anthropogenic pollutant emissions. The values and trends presented in section 3 therefore do not represent actual values for a given year because all historical simulations used a common meteorological year. This was a deliberate choice to ensure that inter-annual variability in meteorology did not confound interpretation of the effects of the anthropogenic emissions alone. However, this is the advantage

of our modelling approach. Observational trends will show the combined effect of the range of factors that contribute to atmospheric pollutant concentration changes, which makes it challenging to isolate the effect of one factor, in this case emissions, alone. Yet, although the modelling does not specifically simulate the concentrations in a given year, all other aspects of the modelling (although set to be constant for each year of modelling) are based on real meteorological conditions.

While it is not possible to attribute specific changes in concentrations and associated mortality burdens to individual policies, as their implementations overlap in time and have locally/regionally different effects, figure 6 illustrates when the main UK, EU and UNECE policy interventions aiming at reducing air pollutant emissions entered into force over the investigated time period. For SO₂, the 1968 UK Clean Air Act had already set in motion substantial reductions in emissions from coal-fired power plants. The UNECE Air






Figure 5. Population-weighted UK average attributable fraction of mortality for O₃ (blue), NO₂ (red) and PM_{2.5} (green) for 1970, 1980, 1990, 2000 and 2010.

Table 1. Population-weighted UK average attributable fraction of respiratory mortality associated with long-term exposure to O₃, and all-cause mortality associated with long-term exposure to NO₂ and PM_{2.5}, for 1970, 1980, 1990, 2000 and 2010.

Year	AF respiratory mortality (%)		AF all-cause mortality (%)	
	O ₃	NO ₂	NO ₂	PM _{2.5}
1970	1.51	5.32	5.32	11.83
1980	1.93	5.44	5.44	10.50
1990	2.33	5.76	5.76	9.50
2000	2.06	4.19	4.19	6.90
2010	1.77	2.96	2.96	5.21

Convention, which originally focused on curbing emissions contributing to the transboundary effects of SO₂ emissions on forest dieback, contributed further to a decline in these emissions. Emissions of NO_x (and, to a lesser extent, NMVOC) have been affected mainly by EC Directives setting emission limits for petrol and diesel vehicles, including the wide-spread introduction of three-way catalytic converters since the early 1970s. Emissions of PM_{2.5} have continuously declined due to introduction of UK and EU emissions regulations on both stationary sources (as a prerequisite for effective control of SO₂ emissions) and mobile sources (with a focus on diesel exhaust emissions). Emissions of NH₃ have declined only moderately since 1980, when the inventory was first compiled, because no UK or pan-European emission control policies have so far achieved substantive reductions of agricultural emissions. The Netherlands and Denmark have unilaterally implemented stringent agricultural NH₃ emissions reduction measures, which will have had some impact on the amount of transboundary SIA into the UK.

As section 3.2 illustrates, public health benefits of emission control policies over the past 40 years have been substantial, with the fraction of all-cause mortality attributable to long-term exposure to both NO₂ and PM_{2.5} showing a marked decline. Emission control regulations by definition can only target emissions of primary pollutants. Hence, the majority of policy interventions which have led to substantive reductions

in emissions first targeted large point sources (SO₂, PM_{2.5} and NO_x) followed by more distributed point sources, most notably the tailpipe emissions of NO_x, NMVOC and PM_{2.5} from road transport. The trends shown in section 3 indicate the additional challenges associated with a secondary pollutant such as O₃. Concentrations of O₃, and associated impact on mortality, increased in the early part of the period due to the reduction in NO_x emissions and the resulting reduction in the chemical removal of O₃ in high-NO_x areas. Since the 1990s, however, AF of mortality due to exposure to O₃ has started to decline as both NO_x and NMVOC emission reductions policies take effect and summer-time O₃ levels have declined (AQEG 2009). The decline in O₃ in the UK has not been as large as it could have been for the reductions in NO_x and NMVOC, because of a slow rise in northern hemisphere background O₃ from rising methane, NO_x and NMVOC emissions further afield transported into the UK (AQEG 2009). However, mortality attributable to ozone is only calculated based on summer-time O₃ levels, therefore trends in mortality may not match annual trends in O₃ concentrations.

The emission reductions described could be considered as 'low-hanging fruit', as they have been achieved primarily through command-and-control approaches targeting large industries, either by implementing stringent control measures e.g. in power plants, or requiring type approval at production for large volumes of motor vehicles (AQEG 2004, 2012). The case of ammonia emissions, which show only small variations over the period for which emission data are available, illustrates the challenges of designing and implementing policies for a large number of small, varied sources with a wide range of technological and behavioural aspects (AQEG 2018). Similar challenges should be expected for further emission reductions in source sectors such as residential combustion (AQEG 2017), or through mitigating pollutant emissions through behavioural change (e.g. changing private transport behaviour) (de Nazelle *et al* 2011, Rojas-Rueda *et al* 2016, Vardoulakis *et al* 2018). The influence of European emissions on UK PM_{2.5} concentrations and the




Figure 6. Timeline of emission changes (based on the UK NAEI) from 1970 to 2010, and of the coming into force of selected air quality legislation in a UK (black), EU (red) and UNECE Air Convention (purple) context. Percentage values show changes in emissions in 2010 with respect to those in 1970 (in 1980 for NH_3).

effect of different potential emissions reductions approaches have been discussed in detail by Vieno *et al* (2014, 2016a and 2016b), respectively.

4.2. Limitations and uncertainties

Although a strength of this modelling study is the elimination of the influence of inter-annual meteorological variability on the time series of spatially-resolved pollutant concentrations, the actual meteorological year selected (2015) may have specific patterns of seasonal effects which differ from a multi-year average. However, as the assessment focuses on the changes in concentrations and health impacts, keeping all other influencing factors constant, the overall results presented in this study are likely not affected by the choice of meteorological year. A detailed discussion of the limitations and uncertainties can be found in section S3 of the suppl. mat.

5. Conclusions

5.1. Effectiveness of AQ policies to reduce public health impacts in the UK

Policies aimed at reducing air pollutant emissions and improving air quality in the UK have been effective in

leading to a substantial decline in concentrations of priority air pollutants over the past 40 + years. Consequently, mortality effects due to exposure to air pollution (as quantified by AF of all-cause mortality) have dropped by 44% and 56% for NO_2 and $\text{PM}_{2.5}$, respectively between 1970 and 2010. For O_3 , attributable respiratory mortality in 2010 was 17% higher than it was in 1970, despite a decline by 24% compared to 1990.

These public health benefits cannot easily be attributed to individual policies and represent a mix of European/regional and UK or local interventions, which have all contributed to a decline in emissions over the study period. At the same time, the data illustrates that changes in emissions of NH_3 from the agricultural sector in particular have been modest. This is because no stringent policies have so far been implemented for agricultural emissions in the UK, with the observed emissions primarily arising from structural changes in UK animal production.

In addition to the reduction in primary $\text{PM}_{2.5}$ emissions in the UK, European policies leading to a decline in SO_2 and NO_x (and, to a lesser extent, NH_3) emissions have contributed to a decline in ambient concentrations of $\text{PM}_{2.5}$ due to reducing precursors of SIA.

5.2. Pollutant-specific aspects and future challenges

Current UK policies in place and in the pipeline are projected to lead to further reductions in NO_x emissions from mobile sources (road transport, shipping, off-road machinery), while at the same time local interventions in the UK such as low-emission zones or clean air zones are expected to contribute to alleviate public health effects by reducing population exposure. Further policy measures are directed at small combustion sources, addressing NO_x as well as primary $\text{PM}_{2.5}$ emissions. In the draft national Air Quality Strategy, which is currently under consultation, the UK Department for Food, Environment and Rural Affairs has set out targets for the reduction of NH_3 emissions from agriculture, designed to attain the National Emission Ceilings target for 2030 of a 16% reduction compared with 2005 levels. Overall, policies addressing agricultural NH_3 emissions are being implemented across Europe, which will likely lead to a marked decrease of contributions to UK $\text{PM}_{2.5}$ from long-range transported SIAs, in particular ammonium nitrate, in the future. Finally, the recent decline in attributable respiratory mortality due to O_3 exposure indicates that further additional reductions of precursors in similar ratios would lead to additional benefits. Nonlinear chemistry in O_3 production, and the impacts of precursor emissions over hemispheric scales, precludes pronouncements on significant reductions and requires further specific study. This should as well investigate to what extent future policies to address ground-level ozone effects on human health, ecosystems and agricultural crops would require more refined spatial and temporal assessments to identify the most effective control strategies for ozone precursor emissions.

Acknowledgments

This work was supported by the Natural Environment Research Council award number NE/R016429/1 as part of the UK-SCaPE programme delivering National Capability.

ORCID iDs

E Carnell  <https://orcid.org/0000-0003-0870-1955>

M Vieno  <https://orcid.org/0000-0001-7741-9377>

S Vardoulakis  <https://orcid.org/0000-0003-3944-7128>

C Heaviside  <https://orcid.org/0000-0002-0263-4985>

S Tomlinson  <https://orcid.org/0000-0002-3237-7596>

U Dragosits  <https://orcid.org/0000-0002-9283-6467>

MR Heal  <https://orcid.org/0000-0001-5539-7293>

S Reis  <https://orcid.org/0000-0003-2428-8320>

References

- Anderson H R 2009 Air pollution and mortality: a history *Atmos. Environ.* **43** 142–52
- AQEG 2004 *Nitrogen dioxide in the United Kingdom* Air Quality Expert Group, UK Department for Environment, Food and Rural Affairs, London PB 9025 (<https://uk-air.defra.gov.uk/library/aeqeg/publications>) (Accessed: 20 December 2018)
- AQEG 2009 *Ozone in the United Kingdom* Air Quality Expert Group, UK Department for Environment, Food and Rural Affairs, London PB13216 (http://uk-air.defra.gov.uk/library/reports?report_id=544) (Accessed: 20 December 2018)
- AQEG 2012 *Fine Particulate Matter (PM_{2.5}) in the United Kingdom* Air Quality Expert Group, UK Department for Environment, Food and Rural Affairs, London PB13837 (http://uk-air.defra.gov.uk/library/reports?report_id=727) (Accessed: 20 December 2018)
- AQEG 2017 *The Potential Air Quality Impacts from Biomass Combustion* Air Quality Expert Group, UK Department for Environment, Food and Rural Affairs, London PB14465 (https://uk-air.defra.gov.uk/library/reports.php?report_id=935) (Accessed: 20 December 2018)
- AQEG 2018 *Air Pollution from Agriculture* Air Quality Expert Group, UK Department for Environment, Food and Rural Affairs, London PB14509 (https://uk-air.defra.gov.uk/library/reports.php?report_id=969) (Accessed: 20 December 2018)
- Atkinson R W *et al* 2016 Long-term exposure to ambient ozone and mortality: a quantitative systematic review and meta-analysis of evidence from cohort studies *BMJ Open* **6** e009493
- Brand C 2016 Beyond 'Dieselgate': implications of unaccounted and future air pollutant emissions and energy use for cars in the United Kingdom *Energy Policy* **97** 1–12
- Brimblecombe P 1987 *The Big Smoke. A History of Air Pollution in London Since Medieval Times* (Abingdon-on-Thames: Routledge)
- Centre for Emission Inventories and Projections (CEIP) 2018 (<http://ceip.at/>) (Accessed: 3 October 2018)
- Cohen A J *et al* 2017 Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the global burden of diseases study 2015 *Lancet* **389** 1907–18
- COMEAP 2009 *Long-Term Exposure to Air Pollution: Effect on Mortality* (London: UK Department of Health, Committee on the Medical Effects of Air Pollutants) (<https://gov.uk/government/publications/comeap-long-term-exposure-to-air-pollution-effect-on-mortality>)
- COMEAP 2010 *The Mortality Effects of Long-Term Exposure to Particulate Air Pollution in the United Kingdom* (London: UK Department of Health, Committee on the Medical Effects of Air Pollutants) (<http://comeap.org.uk/documents/reports.html>)
- COMEAP 2015 *Quantification of Mortality and Hospital Admissions Associated with Ground-Level Ozone* (London: UK Department of Health, Committee on the Medical Effects of Air Pollutants) (<https://gov.uk/government/publications/comeap-quantification-of-mortality-and-hospital-admissions-associated-with-ground-level-ozone>)
- COMEAP 2018 *Associations of Long-Term Average Nitrogen Dioxide with Mortality* PHE Report No. 2018238 (London: UK Department of Health, Committee on the Medical Effects of Air Pollutants) (https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/734799/COMEAP_NO2_Report.pdf)
- de Nazelle A *et al* 2011 Improving health through policies that promote active travel: a review of evidence to support integrated health impact assessment *Environ. Int.* **37** 766–77
- Department for Environment, Food and Rural Affairs 2007 *The Air Quality Strategy for England, Scotland, Wales and Northern Ireland* (London: Defra)
- EEA 2018 Air Quality in Europe - 2018 Report *European Environment Agency Report 12/2018* (Copenhagen, Denmark:

- European Environment Agency) (<https://www.eea.europa.eu/publications/air-quality-in-europe-2018>)
- EPA 2011 *The Benefits and Costs of the Clean Air Act from 1990 to 2020* US Environmental Protection Agency (<https://epa.gov/sites/production/files/2015-07/documents/summaryreport.pdf>) (Accessed: 22 March 2019)
- HEI (Health Effects Institute) 2017 *State of Global Air 2017. Special Report* Boston, MA: Health Effects Institute (https://stateofglobalair.org/sites/default/files/SOGA2017_report.pdf)
- Henneman L R F, Liu C, Mulholland J A and Russell A G 2016 Evaluating the effectiveness of air quality regulations: a review of accountability studies and frameworks *J. Air Waste Manage. Assoc.* **67** 144–72
- Huang Q, Wang T, Chen P, Huang X, Zhu J and Zhuang B 2017 Impacts of emissions reduction and meteorological conditions on air quality improvement during the 2014 Youth Olympic Games in Nanjing, China *Atmos. Chem. Phys.* **17** 13457–71
- Landrigan P J *et al* 2017 The lancet commission on pollution and health *Lancet* **391** 462–512
- Lim S S *et al* 2012 A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010 *Lancet* **380** 2224–60
- Loomis D *et al* 2013 The carcinogenicity of outdoor air pollution *Lancet Oncol.* **14** 1262–3
- Malley C S, Heal M R and Braban C F 2016 Insights from a chronology of the development of atmospheric composition monitoring networks since the 1800s *Atmosphere* **7** 160
- Matus K, Nam K M, Selin N E, Lamsal L N, Reilly J M and Paltsev S 2012 Health Damages from Air Pollution in China *Glob. Environ. Change* **22** 55–66
- Matus K, Yang T, Paltsev S, Reilly J M and Nam K M 2008 Toward integrated assessment of environmental change: air pollution health effects in the USA *Clim. Change* **88** 59–92
- Monks P S *et al* 2015 Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer *Atmos. Chem. Phys.* **15** 8889–973
- National Atmospheric Emission Inventory 2018 (<http://naei.beis.gov.uk/>) (Accessed: 03 October 2018)
- Pope R J, Arnold S R, Chipperfield M P, Latter B G, Siddans R and Kerridge B J 2018 Widespread changes in UK air quality observed from space *Atmos. Sci. Lett.* **19** e817
- Pope R J, Savage N H, Chipperfield M P, Arnold S R and Osborn T J 2014 The influence of synoptic weather regimes on UK air quality: analysis of satellite column NO₂ *Atmos. Sci. Lett.* **15** 211–7
- RCP 2016 *Every Breath We Take: The Lifelong Impact of Air Pollution* (London: Royal College of Physicians) (<https://rcplondon.ac.uk/projects/outputs/every-breath-we-take-lifelong-impact-air-pollution>)
- Reid N, Yap D and Bloxam R 2008 The potential role of background ozone on current and emerging air issues: an overview *Air Qual. Atmos. Health* **1** 19–29
- Rojas-Rueda D *et al* 2016 Health impacts of active transportation in Europe *PLoS One* **11** e0149990
- Shaddick G *et al* 2018 Data Integration for the assessment of population exposure to ambient air pollution for global burden of disease assessment *Environ. Sci. Technol.* **52** 9069–78
- Simpson D *et al* 2012 The EMEP MSC-W chemical transport model—technical description *Atmos. Chem. Phys.* **12** 7825–65
- van Donkelaar A, Martin R V, Brauer M and Boys B L 2015 Use of satellite observations for long-term exposure assessment of global concentrations of fine particulate matter *Environ. Health Perspect.* **123** 135–43
- Vardoulakis S, Kettle R, Cosford P, Lincoln P, Holgate S, Grigg J, Kelly F and Pencheon D 2018 Local action on outdoor air pollution to improve health *Int. J. Public Health* **63** 557–65
- Vestreng V, Myhre G, Fagerli H, Reis S and Tarrasón L 2007 Twenty-five years of continuous sulphur dioxide emission reduction in Europe *Atmos. Chem. Phys.* **7** 3663–81
- Vestreng V, Ntziachristos L, Semb A, Reis S, Isaksen I S A and Tarrasón L 2009 Evolution of NO_x emissions in Europe with focus on road transport control measures *Atmos. Chem. Phys.* **9** 1503–20
- Vieno M *et al* 2014 The role of long-range transport and domestic emissions in determining atmospheric secondary inorganic particle concentrations across the UK *Atmos. Chem. Phys.* **14** 8435–47
- Vieno M *et al* 2016a The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust *Environ. Res. Lett.* **11** 044004
- Vieno M, Heal M R, Williams M L, Carnell E J, Nemitz E, Stedman J R and Reis S 2016b The sensitivities of emissions reductions for the mitigation of UK PM_{2.5} *Atmos. Chem. Phys.* **16** 265–76
- WHO 2013a *Health Risks of Air Pollution in Europe—HRAPIE Project* World Health Organisation, Copenhagen, Denmark (http://euro.who.int/__data/assets/pdf_file/0006/238956/Health_risks_air_pollution_HRAPIE_project.pdf)
- WHO 2013b *Review of Evidence on Health Aspects of Air Pollution—REVIHAAP Project: Technical Report* World Health Organisation, Copenhagen, Denmark (http://www.euro.who.int/__data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf)
- Willocks L J, Bhaskar A, Ramsay C N, Lee D, Brewster D H, Fischbacher C M, Chalmers J, Morris G and Scott E M 2012 Cardiovascular disease and air pollution in Scotland: no association or insufficient data and study design? *BMC Public Health* **12** 227
- Yan Y, Pozzer A, Ojha N, Lin J and Lelieveld J 2018 Analysis of European ozone trends in the period 1995–2014 *Atmos. Chem. Phys.* **18** 5589–605
- Zhang Y, West J J, Mathur R, Xing J, Hogrefe C, Roselle S J, Bash J O, Pleim J E, Gan C-M and Wong D C 2018 Long-term trends in the ambient PM_{2.5}- and O₃-related mortality burdens in the United States under emission reductions from 1990 to 2010 *Atmos. Chem. Phys.* **18** 15003–16