



## **Modelling the global atmospheric transport and deposition of radionuclides from the Fukushima Dai-ichi nuclear accident.**

Theodoros Christoudias (1) and Jos Lelieveld (2)

(1) Cyprus Institute, Cyprus (christoudias@cyi.ac.cy), (2) Max Planck Institute of Chemistry, Mainz, Germany (jos.lelieveld@mpic.de)

We modeled the global atmospheric dispersion and deposition of radionuclides released from the Fukushima Dai-ichi nuclear power plant accident. The EMAC atmospheric chemistry – general circulation model was used, with circulation dynamics nudged towards ERA-Interim reanalysis data. We applied a resolution of approximately 0.5 degrees in latitude and longitude (T255). The model accounts for emissions and transport of the radioactive isotopes  $^{131}\text{I}$  and  $^{137}\text{Cs}$ , and removal processes through precipitation, particle sedimentation and dry deposition. In addition, we simulated the release of  $^{133}\text{Xe}$ , a noble gas that can be regarded as a passive transport tracer of contaminated air. The source terms are based on Chino et al. (2011) and Stohl et al. (2012); especially the emission estimates of  $^{131}\text{I}$  are associated with a high degree of uncertainty. The calculated concentrations have been compared to station observations by the Comprehensive Nuclear-Test-Ban Treaty Organisation (CTBTO).