

RECEIVED: December 30, 2020

REVISED: April 8, 2021

ACCEPTED: April 20, 2021

PUBLISHED: May 13, 2021

Modular invariant A_4 models for quarks and leptons with generalized CP symmetry

Chang-Yuan Yao,^a Jun-Nan Lu^{b,c} and Gui-Jun Ding^{b,c}

^a*School of Physics, Nankai University,
No. 94 Weijin Road, Tianjin 300071, China*

^b*Peng Huanwu Center for Fundamental Theory, University of Science and Technology of China,
No. 96 Jinzhai Road, Hefei, Anhui 230026, China*

^c*Interdisciplinary Center for Theoretical Study and Department of Modern Physics,
University of Science and Technology of China, No. 96 Jinzhai Road, Hefei, Anhui 230026, China*

*E-mail: yaocy@nankai.edu.cn, hitman@mail.ustc.edu.cn,
dinggj@ustc.edu.cn*

ABSTRACT: We perform a systematical analysis of the A_4 modular models with generalized CP for the masses and flavor mixing of quarks and leptons, and the most general form of the quark and lepton mass matrices is given. The CP invariance requires all couplings real in the chosen basis and thus the vacuum expectation value of the modulus τ uniquely breaks both the modular symmetry and CP symmetry. The phenomenologically viable models with minimal number of free parameters and the results of fit are presented. We find 20 models with 7 real free parameters that can accommodate the experimental data of lepton sector. We then apply A_4 modular symmetry to the quark sector to explain quark masses and CKM mixing matrix, the minimal viable quark model is found to contain 10 free real parameters. Finally, we give two predictive quark-lepton unification models which use only 16 real free parameters to explain the flavor patterns of both quarks and leptons.

KEYWORDS: Beyond Standard Model, CP violation, Discrete Symmetries, Neutrino Physics

ARXIV EPRINT: [2012.13390](https://arxiv.org/abs/2012.13390)

Contents

1	Introduction	1
2	Modular symmetry and modular forms of level $N = 3$	3
2.1	Modular forms of level $N = 3$	4
3	Generalized CP consistent with A_4 symmetry	6
4	Lepton models based on A_4 modular symmetry with gCP	7
4.1	Charged lepton sector	8
4.2	Neutrino sector	12
4.3	Phenomenologically viable lepton models	13
5	Quark models based on A_4 modular symmetry with gCP	17
6	Unified models of leptons and quarks	23
6.1	Unified model of L_7 - Q_{11}	23
6.2	Unified model of L_8 - Q_{10}	25
7	Conclusion	28

1 Introduction

The masses and mixing parameters of both quarks and leptons have been measured precisely, however the origin of the hierarchical masses and different mixing patterns of quarks and leptons remain elusive. Although there has been intense theoretical activity in the past decades, there is still no leading candidate for a flavor theory of quarks and leptons. From the theoretical point of view, a very appealing possibility is flavor symmetry as the guiding principle. In this setup, the lepton mixing arises from the mismatched flavor symmetry breaking patterns in the charged lepton and neutrino sectors, and similarly for the quark CKM mixing matrix. See [1] for the latest review.

In conventional flavor symmetry models, the scalar potential of the flavon fields has to be cleverly designed to get the correct vacuum alignment, and generally certain auxiliary symmetry is necessary to forbid dangerous operators. As a consequence, the models look rather complex. Recently the modular invariance as flavor symmetry is suggested to overcome these drawbacks [2]. The lepton mass matrices would appear as the combinations of Yukawa couplings and modular forms which are holomorphic functions of the modulus τ . The vacuum expectation value of τ is the unique source of flavor symmetry breaking. In modular invariant lepton models, the neutrino masses and mixing parameters can be

predicted in terms of a few input parameters and thus these models have strong predictive power.

The phenomenologies of modular invariance have widely studied in the literature, and many modular invariant models for lepton masses and mixing have been constructed by using the inhomogeneous finite modular group Γ_N for $\Gamma_2 \cong S_3$ [3–6], $\Gamma_3 \cong A_4$ [2–4, 7–31], $\Gamma_4 \cong S_4$ [20, 32–39], $\Gamma_5 \cong A_5$ [37, 40, 41] and $\Gamma_7 \cong \text{PSL}(2, Z_7)$ [42]. The formalism of the homogeneous finite modular group Γ'_N which is the double covering of Γ_N have been explored in [43]. The homogeneous finite modular groups $\Gamma'_3 \cong T'$ [43, 44], $\Gamma'_4 \cong S'_4$ [45, 46] and $\Gamma'_5 \cong A'_5$ [47, 48] have been utilized to build lepton and quark models. Moreover, the modular invariance approach has been extended to include the rational weight modular forms, then the modular group should be extended to its metaplectic covering group and the modular forms can be decomposed into irreducible multiplets of the finite metaplectic group $\tilde{\Gamma}_N$ [49]. The superstring theory requires six compact space dimensions with multiple moduli, the modular invariant supersymmetric theories with single modulus has been extended to more general automorphic supersymmetric theory where several moduli can occur naturally [50, 51]. The exchange of moduli between electrons and neutrinos can induce a non-standard neutrino interactions which can leads to a shift of the neutrino mass matrix [26]. Thus the presence of moduli can potentially be tested in neutrino oscillation experiments [26]. The predictive power of the modular invariance approach would be improved further by including the generalized CP symmetry (gCP) which acts on the complex modulus τ as $\tau \xrightarrow{CP} -\tau^*$ up to modular transformations [52–56]. In the symmetric basis where the generators S and T are represented by unitary and symmetric matrices, invariance under gCP would require all coupling constants real.

It is known that A_4 is the smallest finite modular group which admits a three-dimensional irreducible representation such that the three generations of fermion fields can be embedded into a triplet. Many A_4 modular models have been constructed [2–4, 7–31], and most predictive models without gCP use eight independent real parameters to describe the neutrino masses, mixing angles and CP violation phases. In this paper, we intend to perform a systematical analysis of lepton and quark models based on $\Gamma_3 \cong A_4$ modular symmetry with gCP, and we aim at minimizing the number of free parameters. For lepton models, we find that 20 viable models can successfully describe the experimental data of lepton masses and mixing parameters in terms of seven real parameters including the complex modulus τ . In the quark sector, in order to accommodate the measured values of quark masses and CKM mixing matrix, at least 10 free parameters are necessary and we obtain thousands of viable quark models with 10 free real parameters. By combining the lepton and quark sectors, we find that the quark-lepton unification can be achieved with a common value of the modulus τ .

The paper is organized as follows. In section 2, we give a brief review of the basic concepts of modular symmetry and modular groups, the even weight modular forms of level 3 are constructed up to weight 8 and they are organized into irreducible multiplets of A_4 . The generalized CP symmetry compatible with A_4 modular symmetry is discussed in section 3. In section 4, we perform a systematical classification of A_4 modular lepton models with gCP symmetry, the phenomenologically viable models with minimal number

of free parameters and the numerical results of the fit are presented. In section 5, we utilize the A_4 modular symmetry and the gCP symmetry to explain the quark masses and mixing. The complete models for quarks and leptons are presented in section 6. We conclude the paper in section 7.

2 Modular symmetry and modular forms of level $N = 3$

In this section, we will firstly introduce some basic concepts of modular symmetry. The modular group $\bar{\Gamma}$ can be defined from a 2-dimensional special linear group $\Gamma = \text{SL}(2, \mathbb{Z})$ with integer entries and determinant equals to 1 [57, 58]:

$$\text{SL}(2, \mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\}. \quad (2.1)$$

The $\text{SL}(2, \mathbb{Z})$ group acts on the complex modulus τ in the upper half complex plane $\Im\tau > 0$ via fractional linear transformations,

$$\tau \mapsto \gamma\tau = \gamma(\tau) = \frac{a\tau + b}{c\tau + d}, \quad \gamma \in \text{SL}(2, \mathbb{Z}). \quad (2.2)$$

It is easy to find that γ and $-\gamma$ induce the same linear fractional transformation and thus one typically defines the projective special linear group $\text{PSL}(2, \mathbb{Z})$ by the quotient group $\text{PSL}(2, \mathbb{Z}) \equiv \text{SL}(2, \mathbb{Z})/\{I, -I\}$ with I is the identity matrix. The modular group $\bar{\Gamma}$ is isomorphic to $\text{PSL}(2, \mathbb{Z})$ and it has two generators S and T satisfying [57]

$$S^2 = (ST)^3 = \mathbf{1}. \quad (2.3)$$

The matrix forms of S and T are

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}. \quad (2.4)$$

Under the actions of S and T , the modulus τ transform as

$$S : \tau \mapsto -\frac{1}{\tau}, \quad T : \tau \mapsto \tau + 1. \quad (2.5)$$

For a positive integer N , the principal congruence subgroup of level N is defined as

$$\Gamma(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}), \quad a - 1 = d - 1 = b = c = 0 \pmod{N} \right\}, \quad (2.6)$$

which is a normal subgroup of the special linear group $\text{SL}(2, \mathbb{Z})$. One can find that T^N is an element of $\Gamma(N)$. The projective principal congruence subgroup is defined as

$$\bar{\Gamma}(N) = \begin{cases} \Gamma(N)/\{\pm I\} & \text{for } N = 1, 2 \\ \Gamma(N) & \text{for } N \geq 3. \end{cases} \quad (2.7)$$

Notice that $-I$ does not belong to $\Gamma(N)$ for $N \geq 3$. The finite modular group Γ_N is the quotient group $\Gamma_N \equiv \bar{\Gamma}/\bar{\Gamma}(N)$ which can be obtained from $\bar{\Gamma}(1)$ by imposing the condition $T^N = 1$. Consequently the generators S and T of Γ_N satisfy the relations

$$S^2 = (ST)^3 = T^N = 1. \tag{2.8}$$

The finite modular groups Γ_N with $N = 2, 3, 4, 5$ are isomorphic to S_3, A_4, S_4 and A_5 respectively [59]. For $N > 5$, additional relations besides those in eq. (2.8) are necessary in order to render the group finite [59].

The modular form $f(\tau)$ of weight k and level N is a holomorphic function of the complex modulus τ and it transforms under the action of $\bar{\Gamma}(N)$ as follows,

$$f(\tau) \rightarrow f(\gamma\tau) = (c\tau + d)^k f(\tau) \quad \text{for } \forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \bar{\Gamma}(N). \tag{2.9}$$

The modular forms of weight k and level N span a finite dimensional linear space. The modular forms $f_i(\tau)$ can be arranged into some modular multiplets $Y_{\mathbf{r}} \equiv (f_1(\tau), f_2(\tau), \dots)^T$ which transform as certain irreducible representation \mathbf{r} of the Γ_N for even k [2, 43] i.e.

$$Y_i^{(k)}(\gamma\tau) = (c\tau + d)^k \rho_{ij}(\gamma) Y_j^{(k)}(\tau) \quad \text{for } \forall \gamma \in \bar{\Gamma}, \tag{2.10}$$

where γ is the representative element of the coset $\gamma\bar{\Gamma}(N)$ in Γ_N , and $\rho_{ij}(\gamma)$ is the representation matrix of γ .

2.1 Modular forms of level $N = 3$

The linear space of modular forms of level 3 and weight $2k$ has dimension $2k + 1$, and the modular forms of level 3 can be organized into different irreducible representation of the inhomogeneous finite modular group $\Gamma_3 \cong A_4$ up to the automorphy factor [2]. It is known that A_4 is the discrete symmetry group of the rotations that leave a tetrahedron invariant, or the group of the even permutations of four objects. The A_4 group has three one-dimensional irreducible representations denoted by $\mathbf{1}, \mathbf{1}', \mathbf{1}''$ and a three-dimensional representation denoted by $\mathbf{3}$. The representation matrices of the generators S and T are

$$\begin{aligned} \mathbf{1} : S &= 1, & T &= 1, \\ \mathbf{1}' : S &= 1, & T &= \omega, \\ \mathbf{1}'' : S &= 1, & T &= \omega^2, \\ \mathbf{3} : S &= \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, & T &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \end{aligned} \tag{2.11}$$

where $\omega = e^{2\pi i/3} = -1/2 + i\sqrt{3}/2$. The decomposition of the tensor product of two triplets α and β is

$$\begin{aligned}
 (\alpha \otimes \beta)_{\mathbf{1}} &= \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2, \\
 (\alpha \otimes \beta)_{\mathbf{1}'} &= \alpha_3\beta_3 + \alpha_1\beta_2 + \alpha_2\beta_1, \\
 (\alpha \otimes \beta)_{\mathbf{1}''} &= \alpha_2\beta_2 + \alpha_1\beta_3 + \alpha_3\beta_1, \\
 (\alpha \otimes \beta)_{\mathbf{3}_S} &= \begin{pmatrix} 2\alpha_1\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_2 \\ 2\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2\alpha_2\beta_2 - \alpha_1\beta_3 - \alpha_3\beta_1 \end{pmatrix}, \\
 (\alpha \otimes \beta)_{\mathbf{3}_A} &= \begin{pmatrix} \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_1\beta_2 - \alpha_2\beta_1 \\ \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix},
 \end{aligned} \tag{2.12}$$

where $(\alpha \otimes \beta)_{\mathbf{r}}$ denotes the contraction of α and β into the representation \mathbf{r} , $\mathbf{3}_S$ and $\mathbf{3}_A$ stand for the symmetric and the antisymmetric triplet combinations respectively.

The ring of the modular forms of level 3 can be generated by three linearly independent modular forms of weight 2 which are given by [2]:

$$\begin{aligned}
 Y_1(\tau) &= \frac{i}{2\pi} \left[\frac{\eta'(\tau/3)}{\eta(\tau/3)} + \frac{\eta'((\tau+1)/3)}{\eta((\tau+1)/3)} + \frac{\eta'((\tau+2)/3)}{\eta((\tau+2)/3)} - \frac{27\eta'(3\tau)}{\eta(3\tau)} \right], \\
 Y_2(\tau) &= \frac{-i}{\pi} \left[\frac{\eta'(\tau/3)}{\eta(\tau/3)} + \omega^2 \frac{\eta'((\tau+1)/3)}{\eta((\tau+1)/3)} + \omega \frac{\eta'((\tau+2)/3)}{\eta((\tau+2)/3)} \right], \\
 Y_3(\tau) &= \frac{-i}{\pi} \left[\frac{\eta'(\tau/3)}{\eta(\tau/3)} + \omega \frac{\eta'((\tau+1)/3)}{\eta((\tau+1)/3)} + \omega^2 \frac{\eta'((\tau+2)/3)}{\eta((\tau+2)/3)} \right],
 \end{aligned} \tag{2.13}$$

where $\eta(\tau)$ is the Dedekind eta-function

$$\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n), \quad q = e^{2\pi i\tau}. \tag{2.14}$$

We can arrange the three modular functions into a vector $Y_{\mathbf{3}}^{(2)}(\tau) = (Y_1(\tau), Y_2(\tau), Y_3(\tau))^T$ transforming as a triplet $\mathbf{3}$ of A_4 ,

$$Y_{\mathbf{3}}^{(2)}(-1/\tau) = \tau^2 \rho_{\mathbf{3}}(S) Y_{\mathbf{3}}^{(2)}(\tau), \quad Y_{\mathbf{3}}^{(2)}(\tau+1) = \rho_{\mathbf{3}}(T) Y_{\mathbf{3}}^{(2)}(\tau), \tag{2.15}$$

where $\rho_{\mathbf{3}}(S)$ and $\rho_{\mathbf{3}}(T)$ are the representation matrices of S and T in the triplet representation $\mathbf{3}$ given in eq. (2.11). Multiplets of higher weight modular forms can be constructed from the tensor products of $Y_{\mathbf{3}}^{(2)}$. At weight $k = 4$, we find five independent modular forms which decompose as $\mathbf{3} \oplus \mathbf{1} \oplus \mathbf{1}'$ under A_4 ,

$$\begin{aligned}
 Y_{\mathbf{3}}^{(4)} &= \frac{1}{2} (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}}^{(2)})_{\mathbf{3}} = \begin{pmatrix} Y_1^2 - Y_2 Y_3 \\ Y_3^2 - Y_1 Y_2 \\ Y_2^2 - Y_1 Y_3 \end{pmatrix}, \\
 Y_{\mathbf{1}}^{(4)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}}^{(2)})_{\mathbf{1}} = Y_1^2 + 2Y_2 Y_3, \\
 Y_{\mathbf{1}'}^{(4)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}}^{(2)})_{\mathbf{1}'} = Y_3^2 + 2Y_1 Y_2.
 \end{aligned} \tag{2.16}$$

Similarly there are seven modular forms of weight 6, and they can be arranged into a singlet **1** and two triplets **3** of A_4 ,

$$\begin{aligned}
 Y_{\mathbf{1}}^{(6)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}}^{(4)})_{\mathbf{1}} = Y_1^3 + Y_2^3 + Y_3^3 - 3Y_1 Y_2 Y_3, \\
 Y_{\mathbf{3}I}^{(6)} &= Y_{\mathbf{3}}^{(2)} Y_{\mathbf{1}}^{(4)} = (Y_1^2 + 2Y_2 Y_3) \begin{pmatrix} Y_1 \\ Y_2 \\ Y_3 \end{pmatrix}, \\
 Y_{\mathbf{3}II}^{(6)} &= Y_{\mathbf{3}}^{(2)} Y_{\mathbf{1}'}^{(4)} = (Y_3^2 + 2Y_1 Y_2) \begin{pmatrix} Y_3 \\ Y_1 \\ Y_2 \end{pmatrix}.
 \end{aligned} \tag{2.17}$$

Finally the linearly independent weight 8 modular forms of level 3 can be decomposed into three singlets **1**, **1'**, **1''** and two triplets **3** under A_4 ,

$$\begin{aligned}
 Y_{\mathbf{1}}^{(8)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}I}^{(6)})_{\mathbf{1}} = (Y_1^2 + 2Y_2 Y_3)^2, \\
 Y_{\mathbf{1}'}^{(8)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}I}^{(6)})_{\mathbf{1}'} = (Y_1^2 + 2Y_2 Y_3)(Y_3^2 + 2Y_1 Y_2), \\
 Y_{\mathbf{1}''}^{(8)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}II}^{(6)})_{\mathbf{1}''} = (Y_3^2 + 2Y_1 Y_2)^2, \\
 Y_{\mathbf{3}I}^{(8)} &= Y_{\mathbf{3}}^{(2)} Y_{\mathbf{1}}^{(6)} = (Y_1^3 + Y_2^3 + Y_3^3 - 3Y_1 Y_2 Y_3) \begin{pmatrix} Y_1 \\ Y_2 \\ Y_3 \end{pmatrix}, \\
 Y_{\mathbf{3}II}^{(8)} &= (Y_{\mathbf{3}}^{(2)} Y_{\mathbf{3}II}^{(6)})_{\mathbf{3}A} = (Y_3^2 + 2Y_1 Y_2) \begin{pmatrix} Y_2^2 - Y_1 Y_3 \\ Y_1^2 - Y_2 Y_3 \\ Y_3^2 - Y_1 Y_2 \end{pmatrix}.
 \end{aligned} \tag{2.18}$$

3 Generalized CP consistent with A_4 symmetry

The modular group $SL(2, Z)$ can consistently combine with the generalized CP symmetry by introducing another new generator which is represented by [52]

$$\mathcal{CP} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{3.1}$$

Thus the modular group $\Gamma \equiv SL(2, Z)$ is enhanced to $\Gamma^* \equiv GL(2, Z)$. Under the action of \mathcal{CP} , the complex modulus τ transforms as

$$\tau \xrightarrow{\mathcal{CP}} -\tau^*. \tag{3.2}$$

As a consequence, the action of Γ^* on the upper-half complex plane is

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma^* : \begin{cases} \tau \rightarrow \frac{a\tau + b}{c\tau + d} & \text{for } ad - bc = 1, \\ \tau \rightarrow \frac{a\tau^* + b}{c\tau^* + d} & \text{for } ad - bc = -1. \end{cases} \tag{3.3}$$

Considering the CP transformation, then a modular transformation γ and subsequently the inverse CP transformation on the modulus τ , we have

$$\tau \xrightarrow{\mathcal{CP}} -\tau^* \xrightarrow{\gamma} -\frac{a\tau^* + b}{c\tau^* + d} \xrightarrow{\mathcal{CP}^{-1}} \frac{a\tau - b}{-c\tau + d}. \tag{3.4}$$

Hence the consistency condition chain $\mathcal{CP} \rightarrow \gamma \rightarrow \mathcal{CP}^{-1}$ maps the modular group element γ into another element $u(\gamma)$,¹

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow u(\gamma) = \begin{pmatrix} a & -b \\ -c & d \end{pmatrix} = \mathcal{CP}\gamma\mathcal{CP}^{-1}. \quad (3.5)$$

It is straightforward to check that u is an automorphism and it preserves the structure of modular group, i.e., $u(\gamma_1)u(\gamma_2) = u(\gamma_1\gamma_2)$. Moreover there is no group element $\gamma' \in \Gamma$ such that $u(\gamma) = \gamma'\gamma'^{-1}$, therefore u is an outer automorphism of the modular group. It is remarkable that $u(S) = S^{-1}$ and $u(T) = T^{-1}$.

On the other hand, the CP transformation acts on the matter field and the modular form multiplets as follow,

$$\psi(x) \xrightarrow{\mathcal{CP}} X_{\mathbf{r}}\bar{\psi}(x_{\mathcal{P}}), \quad Y(\tau) \xrightarrow{\mathcal{CP}} Y(-\tau^*) = X_{\mathbf{r}}Y^*(\tau), \quad (3.6)$$

where $x = (t, \vec{x})$ and $x_{\mathcal{P}} = (t, -\vec{x})$. Applying the consistency condition chain $\mathcal{CP} \rightarrow \gamma \rightarrow \mathcal{CP}^{-1}$ to the matter field, we can obtain the constraint on the generalized CP transformation $X_{\mathbf{r}}$,

$$X_{\mathbf{r}}\rho_{\mathbf{r}}^*(\gamma)X_{\mathbf{r}}^{-1} = \rho_{\mathbf{r}}(u(\gamma)), \quad \gamma \in \Gamma, \quad (3.7)$$

which has to be fulfilled for consistent implementation of the generalized CP symmetry in the context of modular symmetry. It is sufficient to impose the above consistency condition on the generators S and T ,

$$X_{\mathbf{r}}\rho_{\mathbf{r}}^*(S)X_{\mathbf{r}}^{-1} = \rho_{\mathbf{r}}^{-1}(S), \quad X_{\mathbf{r}}\rho_{\mathbf{r}}^*(T)X_{\mathbf{r}}^{-1} = \rho_{\mathbf{r}}^{-1}(T), \quad (3.8)$$

In the present work, we study the finite modular group A_4 , and both S and T are represented by unitary and symmetric matrices in our working basis, as shown in eq. (2.11). Hence the CP transformation $X_{\mathbf{r}}$ is determined to take the canonical form,

$$X_{\mathbf{r}} = \mathbf{1}_{\mathbf{r}}, \quad (3.9)$$

up to an overall phase. Furthermore, eq. (2.12) implies that the Clebsch-Gordan coefficients are real in the chosen basis, consequently all coupling constants would be real if the theory is required invariant under the CP.

4 Lepton models based on A_4 modular symmetry with gCP

In this section, we shall perform a systematical classification of all minimal lepton models based on the $\Gamma_3 \cong A_4$ modular symmetry with gCP. We shall formulate our models in the framework of $\mathcal{N} = 1$ global supersymmetry. In the present work, the Kähler potential is taken to be the minimal form as in the original work [2]. Note that the Kähler

¹A second CP transformation can possibly be defined with $u(\gamma) = \chi(\gamma) \begin{pmatrix} a & -b \\ -c & d \end{pmatrix}$, where $\chi(\gamma) = \pm 1$ is a homomorphism or more concretely $\chi(S) = \chi(T) = -1$ [45]. In the present work, we consider the even weight modular forms and the inhomogeneous finite modular group $\Gamma_3 \cong A_4$, the modular transformation γ is identified with $-\gamma$. Thus two generalized CP symmetries are identical.

potential is not completely fixed by the modular symmetry [2, 60], there are many other terms compatible with the modular symmetry in the Kähler potential, thus the predictive power of modular invariance would be reduced. Top-down constructions in string theory typically lead to modular symmetry together with certain conventional flavor symmetry. By combining the traditional flavor symmetry with modular symmetry in the scheme of eclectic flavor group [61–65], it was shown that both Kähler potential and superpotential as well as the representations and weights of the fields would be strongly constrained. The superpotential $\mathcal{W}(\Phi_I, \tau)$ can be expanded in power series of the supermultiplets Φ_I ,

$$\mathcal{W}(\Phi_I, \tau) = \sum_n Y_{I_1 \dots I_n}(\tau) \Phi_{I_1} \dots \Phi_{I_n}, \quad (4.1)$$

where $Y_{I_1 \dots I_n}(\tau)$ is a modular form multiplet. Under the modular transformation, the supermultiplets Φ_I and $Y_{I_1 \dots I_n}(\tau)$ transform as

$$\begin{aligned} \tau &\rightarrow \gamma\tau = \frac{a\tau + b}{c\tau + d}, & \gamma &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}), \\ \Phi_I &\rightarrow (c\tau + d)^{-k_I} \rho_I(\gamma) \Phi_I, \\ Y(\tau) &\rightarrow Y(\gamma\tau) = (c\tau + d)^{k_Y} \rho_Y(\gamma) Y(\tau), \end{aligned} \quad (4.2)$$

where $-k_I$ and k_Y are the modular weights of Φ_I and $Y_{I_1 \dots I_n}$ respectively. Besides, $\rho_I(\gamma)$ and $\rho_Y(\gamma)$ are unitary representation of the representative element γ in Γ_N . The superpotential should be invariant under the modular transformation which implies

$$k_Y = k_{I_1} + \dots + k_{I_n}, \quad \rho_Y \otimes \rho_{I_1} \otimes \dots \otimes \rho_{I_n} \ni \mathbf{1}. \quad (4.3)$$

In the following, the Higgs doublets H_u and H_d are assumed to transform as trivial singlet $\mathbf{1}$ of A_4 , and their modular weights k_{H_u} and k_{H_d} are zero.

4.1 Charged lepton sector

If both left-handed and right-handed charged lepton fields transform as triplets of A_4 modular group, i.e.,

$$L \equiv (L_1, L_2, L_3)^T \sim \mathbf{3}, \quad E^c \equiv (E_1^c, E_2^c, E_3^c)^T \sim \mathbf{3}, \quad (4.4)$$

then the most general form of the superpotential in charged lepton sector is given by

$$\mathcal{W}_E = \sum_{\mathbf{r}_a} g_a^E \left(H_d (E^c L)_{\mathbf{r}'_a} Y_{\mathbf{r}_a}^{(k_L + k_{E^c})} \right)_{\mathbf{1}}, \quad (4.5)$$

where the coupling constants g_a^E are real because of the invariance under gCP, k_L and k_{E^c} denote the modular weights of L and E^c respectively. The modular invariance requires $\mathbf{r}'_a \otimes \mathbf{r}_a = \mathbf{1}$, the representation \mathbf{r}_b can be $\mathbf{1}, \mathbf{1}', \mathbf{1}'', \mathbf{3}$ and they are determined by the modular weights k_L and k_{E^c} as shown in table 1. From eq. (4.5), we can read out the

Modular weight k	Modular form $Y_{\mathbf{r}}^{(k)}$
$k = 2$	$Y_{\mathbf{3}}^{(2)}$
$k = 4$	$Y_{\mathbf{1}}^{(4)}, Y_{\mathbf{1}'}^{(4)}, Y_{\mathbf{3}}^{(4)}$
$k = 6$	$Y_{\mathbf{1}}^{(6)}, Y_{\mathbf{3}I}^{(6)}, Y_{\mathbf{3}II}^{(6)}$
$k = 8$	$Y_{\mathbf{1}}^{(8)}, Y_{\mathbf{1}'}^{(8)}, Y_{\mathbf{1}''}^{(8)}, Y_{\mathbf{3}I}^{(8)}, Y_{\mathbf{3}II}^{(8)}$

Table 1. Summary of modular forms of level 3 up to weight 8, the subscript \mathbf{r} denote the transformation property under A_4 modular symmetry. Here $Y_{\mathbf{3}I}^{(6)}$ and $Y_{\mathbf{3}II}^{(6)}$ stand for two weight 6 modular forms transforming in the representation $\mathbf{3}$ of A_4 . Similar conventions are adopted for $Y_{\mathbf{3}I}^{(8)}$ and $Y_{\mathbf{3}II}^{(8)}$.

expressions of the elements of the charged lepton mass matrix,

$$(M_E)_{ij} = v_d \sum_{a,b,c,d} \left[\left(g_{a1}^E (3\delta_{ij} - 1) + g_{a2}^E (1 - \delta_{ij}) (-1)^{\text{mod}(i-j,3)} \right) Y_{\mathbf{3}a,3-\text{mod}(i+j,3)}^{(k_L+k_{E^c})} \right. \\ \left. + g_b^E \delta_{2,\text{mod}(i+j,3)} Y_{\mathbf{1}_b}^{(k_L+k_{E^c})} + g_c^E \delta_{1,\text{mod}(i+j,3)} Y_{\mathbf{1}'_c}^{(k_L+k_{E^c})} + g_d^E \delta_{0,\text{mod}(i+j,3)} Y_{\mathbf{1}''_d}^{(k_L+k_{E^c})} \right]. \tag{4.6}$$

In this type of models, the charged lepton mass hierarchies rely on cancellations of comparable terms, and fine-tuning of the coupling constants is required. The numerical analysis indeed shows that it is impossible to obtain the charged lepton mass hierarchies from simple charged lepton mass matrices involving few coupling parameters. Thus we turn to consider another kind of assignments of the charged lepton fields.

We assume that the three generations of left-handed lepton fields transform as a triplet of A_4 modular group, while the right-handed charged leptons transform as one-dimensional representations of A_4 in order to accommodate the charged lepton masses, i.e.

$$L \equiv (L_1, L_2, L_3)^T \sim \mathbf{3}, \quad E_i^c \sim \mathbf{1}^{l_i} \quad \text{with } i = 1, 2, 3, \tag{4.7}$$

where $l_{1,2,3} = 0, 1, 2$ with $\mathbf{1}^0 \equiv \mathbf{1}$, $\mathbf{1}^1 \equiv \mathbf{1}'$, $\mathbf{1}^2 \equiv \mathbf{1}''$. The modular weights of L and $E_{1,2,3}^c$ are denoted as k_L and $k_{E_{1,2,3}^c}$ respectively. The modular invariance enforces the modular forms coupling to $E_i^c L H_d$ should be a triplet of A_4 , and its modular weight should be $k_L + k_{E_i^c}$. Thus the most general superpotential for the charged lepton masses is given by

$$\mathcal{W}_E = \sum_{i=1}^3 \sum_a \alpha_{i,a} E_i^c L H_d Y_{\mathbf{3}a}^{(k_L+k_{E_i^c})}(\tau), \tag{4.8}$$

where $\alpha_{i,a}$ are coupling constants, and they are constrained to be real by gCP symmetry. The subscript a indicates that there may exist multiple weight $k_L + k_{E_i^c}$ modular forms which transform as a triplet $\mathbf{3}$ of A_4 as shown in table 1. In the following, we will denote $\alpha_1 \equiv \alpha_e$, $\alpha_2 \equiv \beta_e$ and $\alpha_3 \equiv \gamma_e$. From eq. (4.8) we can see that the charged lepton mass matrix can be divided into three rows:

$$M_E = \begin{pmatrix} R_1 \\ R_2 \\ R_3 \end{pmatrix}, \tag{4.9}$$

$k_{E_i^c} + k_L$	Expressions of R_i	Constraints
2, 4	$\alpha_i \left(Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 0$
	$\alpha_i \left(Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 1$
	$\alpha_i \left(Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)}, Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 2$
6, 8	$\left(\alpha_{i,1} Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,1}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,3}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,2}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 0$
	$\left(\alpha_{i,1} Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,3}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,2}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,1}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 1$
	$\left(\alpha_{i,1} Y_{\mathbf{3},2}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,2}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},1}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,1}^{(k_{E_i^c}+k_L)}, \alpha_{i,1} Y_{\mathbf{3},3}^{(k_{E_i^c}+k_L)} + \alpha_{i,2} Y_{\mathbf{3}I,3}^{(k_{E_i^c}+k_L)} \right) v_d$	$l_i = 2$

Table 2. Possible structures of each row R_i of the charged lepton mass matrix for modular forms up to weight 8, where α_i , $\alpha_{i,1}$ and $\alpha_{i,2}$ are Yukawa coupling parameters.

where R_i is a 1×3 sub-matrix with $i = 1, 2, 3$. The general form of the i -th row of charged lepton mass matrix R_i is determined by modular weight $k_L + k_{E_i^c}$ and representation indices l_i . Using the contraction rules of A_4 group given in eq. (2.12), we find the general form of the (ij) element of M_E is

$$(M_E)_{ij} = \sum_a \alpha_{i,a} v_d Y_{\mathbf{3}a, 3-\text{mod}(l_i+j+1,3)}^{(k_L+k_{E_i^c})} \quad (4.10)$$

In the present work, we will be concerned with the modular forms up to weight 8, and higher weight modular forms can be discussed in a similar way. The explicit forms of R_i for $k_L + k_{E_i^c} = 2, 4, 6, 8$ are summarized in table 2. From this table, we can see that if $k_{E_i^c} + k_L = 2, 4$, there will be a single Yukawa coupling parameter α_i for each right-handed charged lepton E_i^c , while there will be two Yukawa couplings $\alpha_{i,1}$ and $\alpha_{i,2}$ for $k_{E_i^c} + k_L = 6, 8$.

We now consider the possible structures of the charged lepton mass matrix. Combining the possible forms of R_i with $i = 1, 2, 3$ given in table 2, we can directly obtain the charged lepton mass matrix. We require that the rank of the charged lepton mass matrix not less than 3 such that any two rows of the charged lepton mass matrix can not be proportional, otherwise at least one of the charged leptons would be massless. In other words, the three generations right-handed charged leptons must be distinguishable from each other by their modular weight and representations. Notice that the effect of exchanging the assignments of right-handed charged leptons is to multiply certain permutation matrices to charged lepton mass matrix from left-hand side, thus the results for the charged lepton masses and mixing matrix are not changed. Without loss of generality, we can assume $k_{E_1^c} + k_L \leq k_{E_2^c} + k_L \leq k_{E_3^c} + k_L$. Thus there are the following four possible cases.

- $k_{E_1^c} + k_L = k_{E_2^c} + k_L = k_{E_3^c} + k_L$.
 In this case, all three generations of right-handed charged lepton have the same modular weights, and the values of $k_{E_i^c} + k_L$ can be 2, 4, 6, 8. To distinguish three right-handed charged leptons, their assignments of one-dimensional representations under A_4 should be different. As a result, the values of l_1 , l_2 and l_3 can only be

chosen as

$$(l_1, l_2, l_3) = (0, 1, 2), \quad (4.11)$$

where the exchange of the values of l_i with $i = 1, 2, 3$ will not give any new results. Notice that once the values of $(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L)$ and (l_1, l_2, l_3) are fixed, the explicit structure of charged lepton mass matrix can be read off directly from table 2.

- $k_{E_1^c} + k_L = k_{E_2^c} + k_L < k_{E_3^c} + k_L$.

In this case, there are 6 allowed combinations of $(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L)$,

$$(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L) = (2, 2, 4), (2, 2, 6), (2, 2, 8), (4, 4, 6), (4, 4, 8), (6, 6, 8). \quad (4.12)$$

Since $k_{E_1^c} = k_{E_2^c}$, the values of l_1 should be distinct from l_2 , while l_3 is not constrained,

$$l_1 \neq l_2, \quad l_{1,2,3} \in \{0, 1, 2\}. \quad (4.13)$$

Consequently (l_1, l_2, l_3) can take 9 set of values,

$$(0, 1, 0), (0, 1, 1), (0, 1, 2), (0, 2, 0), (0, 2, 1), (0, 2, 2), (1, 2, 0), (1, 2, 1), (1, 2, 2). \quad (4.14)$$

- $k_{E_1^c} + k_L < k_{E_2^c} + k_L = k_{E_3^c} + k_L$.

When the last two generations of right-handed charged leptons are assigned with the same modular weight, the values of $(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L)$ can be

$$(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L) = (2, 4, 4), (2, 6, 6), (2, 8, 8), (4, 6, 6), (4, 8, 8), (6, 8, 8). \quad (4.15)$$

The second and third generations of right-handed charged leptons must transform differently under A_4 , therefore we have $l_2 \neq l_3$ while the value of l_1 is free. Similarly we can obtain 9 allowed assignments of (l_1, l_2, l_3) ,

$$(0, 0, 1), (0, 0, 2), (0, 1, 2), (1, 0, 1), (1, 0, 2), (1, 1, 2), (2, 0, 1), (2, 0, 2), (2, 1, 2). \quad (4.16)$$

- $k_{E_1^c} + k_L < k_{E_2^c} + k_L < k_{E_3^c} + k_L$.

If all the right-handed charged leptons have different modular weights, the values of $(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L)$ have 4 choices:

$$(k_{E_1^c} + k_L, k_{E_2^c} + k_L, k_{E_3^c} + k_L) = (2, 4, 6), (2, 4, 8), (2, 6, 8), (4, 6, 8). \quad (4.17)$$

In this case, the representation assignments of three right-handed charged leptons are irrelevant. As a consequence, there are 27 possible combinations of (l_1, l_2, l_3) ,

$$(0, 0, 0), (0, 0, 1), (0, 0, 2), (0, 1, 0), (0, 1, 1), (0, 1, 2), (0, 2, 0), (0, 2, 1), (0, 2, 2), \\ (1, 0, 0), (1, 0, 1), (1, 0, 2), (1, 1, 0), (1, 1, 1), (1, 1, 2), (1, 2, 0), (1, 2, 1), (1, 2, 2), \\ (2, 0, 0), (2, 0, 1), (2, 0, 2), (2, 1, 0), (2, 1, 1), (2, 1, 2), (2, 2, 0), (2, 2, 1), (2, 2, 2). \quad (4.18)$$

In short, we find there are totally $4 + 6 \times 9 + 6 \times 9 + 4 \times 27 = 220$ possible structures of charged lepton mass matrix that will lead to non-vanishing charged lepton masses.

From table 2, we can find that for the cases $k_{E_i^c} + k_L = 2, 4$, there is only one coupling α_i in the i -th row, while there are two couplings $\alpha_{i,1}$ and $\alpha_{i,2}$ for $k_{E_i^c} + k_L = 6, 8$. As a consequence, the number of real Yukawa coupling constants in charged lepton mass matrix is determined by the values of $k_{E_i^c} + k_L$ with $i = 1, 2, 3$. If all the three modular weights $k_{E_i^c} + k_L$ take the values 2 or 4, there will be 3 real coupling parameters in charged lepton mass matrix. If we change $k_{E_i^c} + k_L = 2, 4$ to $k_{E_i^c} + k_L = 6, 8$, one more real coupling in the i -th row of the mass matrix will be introduced. Thus the minimal number of real couplings in the charged lepton mass matrix M_E is 3, and the maximum number is 6. According to the number of free coupling constants in the charged lepton mass matrix, we can divide the 220 possible forms of M_E into four distinct classes: M_E^I , M_E^{II} , M_E^{III} and M_E^{IV} in which 3, 4, 5 and 6 couplings are involved in the charged lepton mass matrix respectively. Moreover, we find that M_E^I , M_E^{II} , M_E^{III} and M_E^{IV} include 20, 90, 90 and 20 possible form of the charged lepton mass matrix. The results of charged lepton mass matrix discussed above are also applicable to up and down quark mass matrices as will be shown in section 5.

4.2 Neutrino sector

We assume neutrinos to be Majorana particles and the neutrino masses are generated through the effective Weinberg operator or the type-I seesaw mechanism. If neutrino masses are described by the Weinberg operator and the three lepton doublets are assigned to an A_4 triplet $\mathbf{3}$, the general superpotential for neutrino masses is

$$\mathcal{W}_\nu = \sum_{\mathbf{r}_a} \frac{g_a^\nu}{\Lambda} (H_u H_u (LL)_{\mathbf{r}'_a} Y_{\mathbf{r}_a}^{(2k_L)}(\tau))_{\mathbf{1}}, \quad (4.19)$$

where g_a^ν are coupling constants, k_L is the modular weight of L , the modular weight of Higgs field H_u is assumed to be vanishing. The modular invariance requires $\mathbf{r}'_a \otimes \mathbf{r}_a = \mathbf{1}$, and the possible assignments of \mathbf{r}_a can be $\mathbf{1}, \mathbf{1}', \mathbf{1}'', \mathbf{3}$, and its value depends on the modular weight $2k_L$ as shown in table 1. The explicit forms of the elements of neutrino mass matrix can be denoted as

$$(M_\nu)_{ij} = \frac{v_u^2}{\Lambda} \sum_{a,b,c,d} \left[g_a^\nu (3\delta_{ij} - 1) Y_{\mathbf{3}_a, 3-\text{mod}(i+j,3)}^{(2k_L)} + g_b^\nu \delta_{2,\text{mod}(i+j,3)} Y_{\mathbf{1}_b}^{(2k_L)} + g_c^\nu \delta_{1,\text{mod}(i+j,3)} Y_{\mathbf{1}'_c}^{(2k_L)} + g_d^\nu \delta_{0,\text{mod}(i+j,3)} Y_{\mathbf{1}''_d}^{(2k_L)} \right]. \quad (4.20)$$

where $i, j = 1, 2, 3$. If neutrino masses are generated through the type-I seesaw mechanism, for the triplet assignments of both right-handed neutrinos N^c and left-handed lepton doublets L , the most general form of the superpotential in the neutrino sector is

$$\mathcal{W}_\nu = \sum_{\mathbf{r}_b} g_b^D \left(H_u (N^c L)_{\mathbf{r}'_b} Y_{\mathbf{r}_b}^{(k_L + k_{N^c})} \right)_{\mathbf{1}} + \sum_{\mathbf{r}_c} g_c^M \left(\Lambda (N^c N^c)_{\mathbf{r}'_c} Y_{\mathbf{r}_c}^{(2k_{N^c})} \right)_{\mathbf{1}}, \quad (4.21)$$

where g_b^D and g_c^M are coupling constants, k_{N^c} is the modular weight of N^c . The modular invariance requires $\mathbf{r}'_b \otimes \mathbf{r}_b = \mathbf{1}$ and $\mathbf{r}'_c \otimes \mathbf{r}_c = \mathbf{1}$, the representation \mathbf{r}_b and \mathbf{r}_c can be

$\mathbf{1}, \mathbf{1}', \mathbf{1}'', \mathbf{3}$ and they are determined by the modular weights k_L and k_{N^c} as shown in table 1. From eq. (4.21), we can read out the expressions of the elements of Dirac and Majorana neutrino mass matrices,

$$\begin{aligned}
 (M_D)_{ij} &= v_u \sum_{a,b,c,d} \left[\left(g_{a1}^D (3\delta_{ij} - 1) + g_{a2}^D (1 - \delta_{ij}) (-1)^{\text{mod}(i-j,3)} \right) Y_{\mathbf{3}_a, \mathbf{3} - \text{mod}(i+j,3)}^{(k_L+k_{N^c})} \right. \\
 &\quad \left. + g_b^D \delta_{2, \text{mod}(i+j,3)} Y_{\mathbf{1}_b}^{(k_L+k_{N^c})} + g_c^D \delta_{1, \text{mod}(i+j,3)} Y_{\mathbf{1}'_c}^{(k_L+k_{N^c})} + g_d^D \delta_{0, \text{mod}(i+j,3)} Y_{\mathbf{1}''_d}^{(k_L+k_{N^c})} \right], \\
 (M_N)_{ij} &= \Lambda \sum_{a,b,c,d} \left[g_a^M (3\delta_{ij} - 1) Y_{\mathbf{3}_a, \mathbf{3} - \text{mod}(i+j,3)}^{(2k_{N^c})} + g_b^M \delta_{2, \text{mod}(i+j,3)} Y_{\mathbf{1}_b}^{(2k_{N^c})} \right. \\
 &\quad \left. + g_c^M \delta_{1, \text{mod}(i+j,3)} Y_{\mathbf{1}'_c}^{(2k_{N^c})} + g_d^M \delta_{0, \text{mod}(i+j,3)} Y_{\mathbf{1}''_d}^{(2k_{N^c})} \right]. \tag{4.22}
 \end{aligned}$$

The effective light neutrino mass matrix in the type-I seesaw models is given by the seesaw formula,

$$M_\nu = -M_D^T M_N^{-1} M_D. \tag{4.23}$$

We are only interested in the models with less free parameters, and we list the possible neutrino models in table 3, for which the resulting light neutrino mass matrices contain less than 4 free real parameters besides the complex modulus τ .

4.3 Phenomenologically viable lepton models

In section 4.1 and section 4.2, we have systematically constructed the charged lepton and neutrino mass matrices respectively. By diagonalizing the mass matrices we can obtain the charged lepton and neutrino masses and the lepton mixing matrix. The variation of the model parameters will dynamically affect the values of these experimental observables. In order to quantitatively estimate how well a model can describe the data, we perform a χ^2 analysis to find out the best fit values of the input parameters and the corresponding predictions for lepton masses and mixing parameters. Because the inverted ordering (IO) neutrino mass spectrum is disfavored by the global data analyses [66, 67] at about 3σ confidence level, we focus on normal ordering (NO) neutrino masses in this paper. The χ^2 function is defined as

$$\chi^2 = \sum_{i=1}^n \left(\frac{P_i(x_1, x_2, \dots, x_m) - \mu_i}{\sigma_i} \right)^2, \tag{4.24}$$

where P_i are predictions for the physical observables including neutrino mixing parameters $\theta_{12}, \theta_{13}, \theta_{23}, \delta_{CP}^l$ and the mass ratios $m_e/m_\mu, m_\mu/m_\tau, \Delta m_{21}^2/\Delta m_{31}^2$ which are complex nonlinear functions of the free parameters of the model, μ_i and σ_i denote the central values and 1σ deviations respectively of the corresponding quantities given in following. From the NuFIT v5.0 with Super-Kamiokanda atmospheric data [68], the best fit values and 1σ ranges of the neutrino parameters are

$$\begin{aligned}
 \sin^2 \theta_{12}^l &= 0.304_{-0.012}^{+0.012}, & \sin^2 \theta_{13}^l &= 0.02219_{-0.00063}^{+0.00062}, & \sin^2 \theta_{23}^l &= 0.573_{-0.020}^{+0.016}, \\
 \delta_{CP}^l/\pi &= 1.0944_{-0.1333}^{+0.1500}, & \frac{\Delta m_{21}^2}{10^{-5}\text{eV}^2} &= 7.42_{-0.20}^{+0.21}, & \frac{\Delta m_{31}^2}{10^{-3}\text{eV}^2} &= 2.517_{-0.028}^{+0.026},
 \end{aligned} \tag{4.25}$$

	k_L, k_{N^c}	Neutrino mass matrix
W_1	1, —	$M_\nu = g^\nu \begin{pmatrix} 2Y_{3,1}^{(2)} & -Y_{3,3}^{(2)} & -Y_{3,2}^{(2)} \\ -Y_{3,3}^{(2)} & 2Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} \\ -Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} & 2Y_{3,3}^{(2)} \end{pmatrix} \frac{v_u^2}{\Lambda}$
W_2	2, —	$M_\nu = \left[\begin{pmatrix} 2Y_{3,1}^{(4)} & -Y_{3,3}^{(4)} & -Y_{3,2}^{(4)} \\ -Y_{3,3}^{(4)} & 2Y_{3,2}^{(4)} & -Y_{3,1}^{(4)} \\ -Y_{3,2}^{(4)} & -Y_{3,1}^{(4)} & 2Y_{3,3}^{(4)} \end{pmatrix} + g_2^\nu Y_1^{(4)} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + g_3^\nu Y_{1'}^{(4)} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right] \frac{v_u^2}{\Lambda}$
W_3	3, —	$M_\nu = \left[\begin{pmatrix} 2Y_{3l,1}^{(6)} & -Y_{3l,3}^{(6)} & -Y_{3l,2}^{(6)} \\ -Y_{3l,3}^{(6)} & 2Y_{3l,2}^{(6)} & -Y_{3l,1}^{(6)} \\ -Y_{3l,2}^{(6)} & -Y_{3l,1}^{(6)} & 2Y_{3l,3}^{(6)} \end{pmatrix} + g_2^\nu \begin{pmatrix} 2Y_{3ll,1}^{(6)} & -Y_{3ll,3}^{(6)} & -Y_{3ll,2}^{(6)} \\ -Y_{3ll,3}^{(6)} & 2Y_{3ll,2}^{(6)} & -Y_{3ll,1}^{(6)} \\ -Y_{3ll,2}^{(6)} & -Y_{3ll,1}^{(6)} & 2Y_{3ll,3}^{(6)} \end{pmatrix} + g_3^\nu Y_1^{(6)} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right] \frac{v_u^2}{\Lambda}$
S_1	2, 0	$M_D = \begin{pmatrix} 2g_1^D Y_{3,1}^{(2)} & (-g_1^D + g_2^D) Y_{3,3}^{(2)} & (-g_1^D - g_2^D) Y_{3,2}^{(2)} \\ (-g_1^D - g_2^D) Y_{3,3}^{(2)} & 2g_1^D Y_{3,2}^{(2)} & (-g_1^D + g_2^D) Y_{3,1}^{(2)} \\ (-g_1^D + g_2^D) Y_{3,2}^{(2)} & (-g_1^D - g_2^D) Y_{3,1}^{(2)} & 2g_1^D Y_{3,3}^{(2)} \end{pmatrix} v_u, \quad M_N = g^M \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \Lambda$
S_2	-1, 1	$M_D = g^D \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} v_u, \quad M_N = g^M \begin{pmatrix} 2Y_{3,1}^{(2)} & -Y_{3,3}^{(2)} & -Y_{3,2}^{(2)} \\ -Y_{3,3}^{(2)} & 2Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} \\ -Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} & 2Y_{3,3}^{(2)} \end{pmatrix} \Lambda$
S_3	1, 1	$M_D = \begin{pmatrix} 2g_1^D Y_{3,1}^{(2)} & (-g_1^D + g_2^D) Y_{3,3}^{(2)} & (-g_1^D - g_2^D) Y_{3,2}^{(2)} \\ (-g_1^D - g_2^D) Y_{3,3}^{(2)} & 2g_1^D Y_{3,2}^{(2)} & (-g_1^D + g_2^D) Y_{3,1}^{(2)} \\ (-g_1^D + g_2^D) Y_{3,2}^{(2)} & (-g_1^D - g_2^D) Y_{3,1}^{(2)} & 2g_1^D Y_{3,3}^{(2)} \end{pmatrix} v_u, \quad M_N = g^M \begin{pmatrix} 2Y_{3,1}^{(2)} & -Y_{3,3}^{(2)} & -Y_{3,2}^{(2)} \\ -Y_{3,3}^{(2)} & 2Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} \\ -Y_{3,2}^{(2)} & -Y_{3,1}^{(2)} & 2Y_{3,3}^{(2)} \end{pmatrix} \Lambda$
S_4	-2, 2	$M_D = g^D \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} v_u, \quad M_N = \left[g_1^M \begin{pmatrix} 2Y_{3,1}^{(4)} & -Y_{3,3}^{(4)} & -Y_{3,2}^{(4)} \\ -Y_{3,3}^{(4)} & 2Y_{3,2}^{(4)} & -Y_{3,1}^{(4)} \\ -Y_{3,2}^{(4)} & -Y_{3,1}^{(4)} & 2Y_{3,3}^{(4)} \end{pmatrix} + g_2^M Y_1^{(4)} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} + g_3^M Y_{1'}^{(4)} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right] \Lambda$
S_5	-3, 3	$M_D = g^D \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} v_u, \quad M_N = \left[g_1^M \begin{pmatrix} 2Y_{3l,1}^{(6)} & -Y_{3l,3}^{(6)} & -Y_{3l,2}^{(6)} \\ -Y_{3l,3}^{(6)} & 2Y_{3l,2}^{(6)} & -Y_{3l,1}^{(6)} \\ -Y_{3l,2}^{(6)} & -Y_{3l,1}^{(6)} & 2Y_{3l,3}^{(6)} \end{pmatrix} + g_2^M \begin{pmatrix} 2Y_{3ll,1}^{(6)} & -Y_{3ll,3}^{(6)} & -Y_{3ll,2}^{(6)} \\ -Y_{3ll,3}^{(6)} & 2Y_{3ll,2}^{(6)} & -Y_{3ll,1}^{(6)} \\ -Y_{3ll,2}^{(6)} & -Y_{3ll,1}^{(6)} & 2Y_{3ll,3}^{(6)} \end{pmatrix} + g_3^M Y_1^{(6)} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right] \Lambda$

Table 3. The predictions for the neutrino mass matrices, the neutrino masses are generated through the Weinberg operator for the models $W_{1,2,3}$ and the type-I seesaw mechanism for the models $S_{1,2,3,4,5}$. Here we only present the cases which involve at most three real coupling constants in the effective light neutrino mass matrix.

where $\theta_{12}^l, \theta_{13}^l$ and θ_{23}^l are the three lepton mixing angles, δ_{CP}^l is the Dirac CP phase, Δm_{21}^2 and Δm_{31}^2 are the neutrino mass squared differences. The charge lepton masses will enter in the χ^2 function in the form of their ratios, the best fit values and 1σ errors are taken from ref. [69],

$$m_e/m_\mu = 0.004737 \pm 0.000040, \quad m_\mu/m_\tau = 0.05857 \pm 0.00047, \quad (4.26)$$

at the grand unified theory (GUT) scale 2×10^{16} GeV. Notice that the charged lepton mass matrices are given at the scale where the modulus τ obtains the vacuum expectation value $\langle \tau \rangle$, which is expected to be around the GUT scale. The overall scale of the charged

lepton mass matrix doesn't affect the mass ratios and mixing parameters, and it is fixed by the central value of $m_\tau = 1.30234$ GeV at GUT scale. Similarly the overall scale of the light neutrino mass matrix is fixed by the solar neutrino mass squared difference $\Delta m_{21}^2 = 7.42 \times 10^{-5}$ eV². It has been shown that the effect of renormalization group evolution (RGE) on the neutrino masses and mixing parameters can be negligible for small value of $\tan \beta$ and NO neutrino masses [7], hence the corrections from RGE are not considered.²

The minimization algorithm in `TMinuit` [70], a package developed by CERN, is used to numerically minimize the value of χ^2 function to determine the best fit values of the input parameters. The free parameters vary in some given ranges in `TMinuit`. We regard the complex modulus τ as a random complex number in the fundamental domain $\mathcal{F} : |\text{Re}\tau| \leq \frac{1}{2}, \text{Im}\tau > 0$ and $|\tau| \geq 1$. The absolute values of all coupling constants freely run in the range of $[0, 10^6]$.

Combining the possible structures in the charged lepton sector and neutrino sector, we can obtain $220 \times 8 = 1760$ lepton models. In this paper, we are only interested in the minimal models that are compatible with the experimental data, thus only the lepton models with 7 real parameters (**A**) and 8 real parameters (**B**) are considered.³ We perform a comprehensive numerical analysis for these models one by one.

(A) Models with 7 real parameters.

There are two possibilities: 3 couplings in the charged lepton sector and 2 couplings in the neutrino sector, or 4 couplings in the charged lepton sector and 1 coupling in the neutrino sector. The former case has $20 \times 2 = 40$ (number of charged lepton models times neutrino models) lepton models, and the latter case has $90 \times 2 = 180$ possible models. We scan over all these models and optimize the χ^2 function with the `TMinuit` package. We find that 20 lepton models out of them can give very good fit to the experimental data for certain values of input parameters. All the 20 models share the same neutrino model S_2 in table 3, while the charged lepton models are different and they are listed in table 4, where the modular weights $k'_{E_i^c} \equiv k_{E_i^c} + k_L$ and the representation indices l_i with $i = 1, 2, 3$ are given explicitly. The corresponding charged lepton mass matrices can be straightforwardly read out from table 2 with the given values of $k'_{E_i^c}$ and l_i . Furthermore we show the best fit values of the input parameters and the predictions of the mixing parameters and neutrino masses in table 5 for NO neutrino mass spectrum. It is notable that the first ten models $C_{1,\dots,10}-S_2$ give quite similar predictions for the lepton observables at the best fit points, and the predictions of the other ten models $C_{11,\dots,20}-S_2$ are also very similar.

²The neutrino mass ratios and mixing angles are almost RGE invariant for hierarchical NO neutrino spectrum, while the light neutrino masses as well mass-squared differences could change under RGE. In our numerical analysis, the χ^2 involves the neutrino mixing angles, Dirac CP phase and the ratios of neutrino mass-squared differences. The overall scale of the neutrino matrix is determined by the solar neutrino mass squared difference so that the absolute neutrino masses can be fixed. Hence the effect of RGE can be absorbed into the overall factor of the neutrino matrix, and the results of fitting are not significantly changed by RGE.

³The minimal A_4 modular models for leptons contain 6 real parameters, nevertheless they are incompatible with the experimental data.

Models	$(k'_{E_1^c}, k'_{E_2^c}, k'_{E_3^c})$	(l_1, l_2, l_3)	Models	$(k'_{E_1^c}, k'_{E_2^c}, k'_{E_3^c})$	(l_1, l_2, l_3)
C_1	(2, 4, 8)	(0, 0, 1)	C_{11}	(2, 2, 6)	(1, 0, 2)
C_2	(4, 4, 8)	(1, 0, 1)	C_{12}	(2, 4, 6)	(1, 0, 2)
C_3	(4, 4, 8)	(2, 0, 1)	C_{13}	(2, 4, 6)	(1, 1, 2)
C_4	(2, 4, 8)	(1, 0, 1)	C_{14}	(2, 2, 6)	(1, 2, 2)
C_5	(2, 4, 8)	(2, 0, 1)	C_{15}	(2, 4, 6)	(1, 2, 2)
C_6	(2, 4, 8)	(0, 1, 2)	C_{16}	(2, 2, 6)	(2, 0, 0)
C_7	(4, 4, 8)	(0, 1, 2)	C_{17}	(2, 2, 6)	(2, 1, 0)
C_8	(2, 4, 8)	(1, 1, 2)	C_{18}	(2, 4, 6)	(2, 0, 0)
C_9	(4, 4, 8)	(2, 1, 2)	C_{19}	(2, 4, 6)	(2, 1, 0)
C_{10}	(2, 4, 8)	(2, 1, 2)	C_{20}	(2, 4, 6)	(2, 2, 0)

Table 4. List of the charged lepton models C_1, C_2, \dots, C_{20} , where $k'_{E_i^c} \equiv k_{E_i^c} + k_L$ is the modular weight of the modular forms in the charged lepton Yukawa coupling. We would like to remind that the right-handed charged lepton E_i^c transform as $\mathbf{1}^{l_i}$ under A_4 with $\mathbf{1}^0 \equiv \mathbf{1}$, $\mathbf{1}^1 \equiv \mathbf{1}'$ and $\mathbf{1}^2 \equiv \mathbf{1}''$.

Models	Best fit values of the input parameters for NO							χ^2_{\min}
	$\text{Re}\langle\tau\rangle$	$\text{Im}\langle\tau\rangle$	β_e/α_e	$\gamma_{e,1}/\alpha_e$	$\gamma_{e,2}/\alpha_e$	$\alpha_e v_d/\text{MeV}$	$\frac{(g^D)^2 v_u^2}{g^M \Lambda^4}/\text{meV}$	
C_1-S_2	0.19214	1.09373	230.31529	1.20754×10^3	2.04179×10^3	0.32342	24.74650	4.87172
C_2-S_2	0.19214	1.09373	183.94677	964.37095	1.63075×10^3	0.40495	24.74647	4.87191
C_3-S_2	0.19214	1.09373	62.75683	329.01303	556.35886	1.18694	24.74648	4.87191
C_4-S_2	0.19214	1.09373	132.87150	696.60007	1.17795×10^3	0.56061	24.74647	4.87191
C_5-S_2	0.19214	1.09374	72.13887	378.32788	639.58609	1.03246	24.74642	4.86955
C_6-S_2	-0.13677	1.18044	241.68127	3.75452×10^3	-2.35923×10^3	0.33938	23.58270	4.87172
C_7-S_2	-0.13677	1.18044	183.94677	2.85758×10^3	-1.79567×10^3	0.44590	23.58267	4.87191
C_8-S_2	-0.13678	1.18043	75.69807	1.17614×10^3	-739.00172	1.08340	23.58289	4.86955
C_9-S_2	-0.13677	1.18044	62.75684	974.91502	-612.62690	1.30698	23.58267	4.87191
$C_{10}-S_2$	-0.13677	1.18044	139.42871	2.166×10^3	-1.36109×10^3	0.58827	23.58267	4.87191
$C_{11}-S_2$	0.17243	1.14007	0.01051	11.53676	6.32548	67.00293	23.96081	9.61938
$C_{12}-S_2$	0.17244	1.14004	0.00526	11.53706	6.32415	67.00313	23.96145	9.58973
$C_{13}-S_2$	0.17235	1.14019	0.00504	11.53723	6.32610	67.00833	23.95954	9.59010
$C_{14}-S_2$	0.17243	1.14007	0.01917	11.53674	6.32549	67.00294	23.96081	9.61938
$C_{15}-S_2$	0.17248	1.13999	0.00503	11.53695	6.32352	67.00093	23.96217	9.59336
$C_{16}-S_2$	-0.17243	1.14007	0.01051	6.32548	11.53675	67.00297	23.96080	9.61938
$C_{17}-S_2$	-0.17243	1.14007	0.01917	6.32549	11.53674	67.00297	23.96080	9.61938
$C_{18}-S_2$	-0.17249	1.13996	0.00504	6.32449	11.53444	67.00001	23.96252	9.59010
$C_{19}-S_2$	-0.17241	1.14010	0.00526	6.32455	11.53774	67.00516	23.96072	9.58973
$C_{20}-S_2$	-0.17237	1.14015	0.00504	6.32468	11.53901	67.00708	23.95998	9.59336

Models	Predictions for mixing parameters and neutrino masses at best fit point												
	$\sin^2 \theta_{12}$	$\sin^2 \theta_{13}$	$\sin^2 \theta_{23}$	δ_{CP}/π	m_e/m_μ	m_μ/m_τ	$\Delta m_{21}^2/\Delta m_{31}^2$	α_{21}/π	α_{31}/π	m_1/meV	m_2/meV	m_3/meV	$m_{\beta\beta}/\text{meV}$
$C_{1-10}-S_2$	0.30287	0.02218	0.58073	1.41681	0.00474	0.05857	0.02957	1.43461	0.92299	10.96085	13.94060	51.27982	8.57441
$C_{11-20}-S_2$	0.33670	0.02162	0.57918	1.01474	0.00474	0.05857	0.03030	0.94333	1.85751	10.96083	13.89816	50.67471	3.52050

Table 5. The best fit values of the input parameters at the minimum of the χ^2 for NO neutrino masses. We give the predictions for neutrino mixing angles $\theta_{12}, \theta_{13}, \theta_{23}$ and Dirac CP violating phase δ_{CP} as well as Majorana CP violating phases α_{21}, α_{31} , and the light neutrino masses $m_{1,2,3}$ and the effective mass $m_{\beta\beta}$ in neutrinoless double beta decay.

The models $C_{1,\dots,5}-S_2$ only differ in the first row, the value of the coupling constant α_e associated with the first row should be smaller than β_e , $\gamma_{e,1}$ and $|\gamma_{e,2}|$ in order to be accommodate the hierarchical charged lepton masses, as can be seen from table 5. Numerically the contribution of the first row is found to be of order 10^{-3} with respect to the other two rows. Therefore the five models $C_{1,\dots,5}-S_2$ give similar numerical results and the best fit values of τ are very close to each other. Analogously the models $C_{6,\dots,10}-S_2$ are also only different in the first row whose contributions are negligible. Notice that the best fit values of τ of the models $C_{1,\dots,5}-S_2$ and $C_{6,\dots,10}-S_2$ are different, it is a numerical coincidence that they give similar predictions for the lepton masses and mixing parameters. Analogously the above reasoning also holds true for the latter ten models $C_{11,\dots,20}-S_2$ except that the contribution of the second row rather than the first row is insignificant. For illustration, we take the model C_1-S_2 as an example. We use the widely-used sampler `MultiNest` [71, 72] to scan the parameter space fully and efficiently, and the predictions for the lepton masses and mixing parameters are required to be compatible with data at 3σ level. The correlations among the input parameters and observables are shown in figure 1. We see that the CP violation phase δ_{CP}^l is predicted to be around 1.5π and the atmospheric mixing angle θ_{23}^l is in the second octant. These predictions could be tested in the forthcoming neutrino oscillation experiments.

(B) Models with 8 real parameters.

This type of models involve 6 real coupling constants, and they can be decomposed into: 3 couplings in both neutrino and charged lepton sectors, or 4 couplings in charged lepton sector and 2 couplings in neutrino sector, or 5 couplings in charged lepton sector and 1 coupling in neutrino sector. There are totally 440 such models up to weight 8. In the same fashion as previous case, we numerically minimize the χ^2 function with the `TMinuit` package. Eventually we find 360 phenomenologically viable lepton models for certain values of input parameters, and 206 models are consistent with the experimental data at 1σ level. Due to the limit of space, we present eight benchmark models, the structures of the charged lepton sector are given in table 6. We show the best fit values of the input parameters and the predictions of the mixing parameters and neutrino masses in table 7 under the assumption of NO neutrino masses. The A_4 modular model with gCP symmetry was recently studied in [30], and a lepton model with 8 parameters was proposed. It corresponds to the D_2-W_2 model in our work, see table 7 for the numerical results of this model.

5 Quark models based on A_4 modular symmetry with gCP

In this section, we proceed to consider the quark models in the framework of A_4 modular symmetry with gCP. Similar to what we have done for the charged lepton sector in section 4.1, we can systematically analyze possible quark models with A_4 modular symmetry. If the left-handed and right-handed quark fields are assigned to A_4 triplet $\mathbf{3}$, the three generations of quark fields would be treated on an equal footing. Generally more




Figure 1. The predictions for the correlations among the input free parameters, neutrino mixing angles, CP violation phases and neutrino masses in the lepton model C_1 – S_2 . The vertical and horizontal dashed lines are the 3σ bounds taken from [68].

free parameters are needed and they should be fine tuned to generate the mass hierarchies of quarks. On the other hand, if all the quark fields transform as singlets under A_4 , the quark mass matrices would be less constrained by modular symmetry. Hence we shall consider the representation assignment: the three generations of left-handed quark fields are assigned to a triplet $\mathbf{3}$ of the A_4 modular group while the right-handed quarks are singlets of A_4 , i.e.,

$$Q_L \equiv (Q_1, Q_2, Q_3)^T \sim \mathbf{3}, \quad q_i^c \sim \mathbf{1}^{l_i} \quad \text{with } q = u, d, \quad i = 1, 2, 3, \quad (5.1)$$

Models	$(k'_{E_1^c}, k'_{E_2^c}, k'_{E_3^c})$	(l_1, l_2, l_3)
D_1	(4, 6, 8)	(1, 2, 2)
D_2	(2, 2, 2)	(0, 2, 1)
D_3	(2, 4, 4)	(0, 1, 2)
D_4	(4, 4, 6)	(1, 2, 0)
D_5	(4, 6, 8)	(0, 1, 1)
D_6	(2, 2, 8)	(0, 2, 1)
D_7	(2, 2, 4)	(0, 1, 0)
D_8	(4, 4, 4)	(0, 1, 2)

Table 6. List of the charged lepton models D_1, D_2, \dots, D_8 , where $k'_{E_i^c} \equiv k_{E_i^c} + k_L$ is the weight of the modular forms in the charged lepton Yukawa coupling. The left-handed lepton fields are assumed to be an A_4 triplet and the right-handed charged leptons E_i^c transform as $\mathbf{1}^{l_i}$ under A_4 .

Best fit values of the input parameters for NO						
Model	D_1-W_1	Model	D_4-S_1	Model	D_2-W_2	D_3-W_3
$\text{Re}(\tau)$	0.05389	$\text{Re}(\tau)$	0.35597	$\text{Re}(\tau)$	0.06026	0.01908
$\text{Im}(\tau)$	2.66993	$\text{Im}(\tau)$	0.94804	$\text{Im}(\tau)$	1.01109	1.02906
$\beta_{e,1}/\alpha_e$	2.29253×10^3	β_e/α_e	2.78535×10^3	β_e/α_e	51.55285	15.57674
$\beta_{e,2}/\alpha_e$	4.22108×10^4	$\gamma_{e,1}/\alpha_e$	558.22251	γ_e/α_e	765.43156	0.03590
$\gamma_{e,1}/\alpha_e$	1.21762×10^3	$\gamma_{e,2}/\alpha_e$	-651.31064	g_2^2/g_1^2	14.03290	-0.30519
$\gamma_{e,2}/\alpha_e$	2.91946×10^4	g_2^D/g_1^D	3.54628	g_3^2/g_1^2	1.14817	7.97382
$\alpha_e v_d/\text{MeV}$	3.35046	$\alpha v_d/\text{MeV}$	0.26571	$\alpha v_d/\text{MeV}$	1.32754	60.61083
$(v_u^2/\Lambda)/\text{meV}$	28.48955	$((g_1^D)^2 v_u^2/(g^M \Lambda))/\text{meV}$	2.69366	$(g_1^D)^2 v_u^2/(g_1^M \Lambda)/\text{meV}$	3.51922	12.59531
Model	D_5-S_2	Model	D_6-S_3	Model	D_7-S_4	D_8-S_5
$\text{Re}(\tau)$	-0.17648	$\text{Re}(\tau)$	-0.08357	$\text{Re}(\tau)$	0.23585	-0.30537
$\text{Im}(\tau)$	1.12772	$\text{Im}(\tau)$	1.15143	$\text{Im}(\tau)$	1.47319	1.77322
$\beta_{e,1}/\alpha_e$	69.56803	β_e/α_e	120.79494	β_e/α_e	16.52692	3.53453×10^3
$\beta_{e,2}/\alpha_e$	158.37007	$\gamma_{e,1}/\alpha_e$	20.66057	γ_e/α_e	0.01649	209.08250
$\gamma_{e,1}/\alpha_e$	3.93725×10^3	$\gamma_{e,2}/\alpha_e$	-22.46201	g_2^M/g_1^M	-2.78133	-2.66807
$\gamma_{e,2}/\alpha_e$	-1.61631×10^3	g_2^D/g_1^D	-0.76278	g_3^M/g_1^M	5.54678	-0.07778
$\alpha v_d/\text{MeV}$	0.33971	$\alpha v_d/\text{MeV}$	4.99601	$\alpha v_d/\text{MeV}$	75.92891	0.36424
$((g^D)^2 v_u^2/(g^M \Lambda))/\text{meV}$	24.23812	$((g_1^D)^2 v_u^2/(g^M \Lambda))/\text{meV}$	16.74812	$((g^D)^2 v_u^2/(g_1^M \Lambda))/\text{meV}$	56.40935	26.95432

Models	Predictions for mixing parameters and neutrino masses at best fitting point												χ^2_{\min}	
	$\sin^2 \theta_{12}^l$	$\sin^2 \theta_{13}^l$	$\sin^2 \theta_{23}^l$	δ_{CP}^l/π	m_e/m_μ	m_μ/m_τ	$\Delta m_{21}^2/\Delta m_{31}^2$	α_{21}/π	α_{31}/π	m_1/meV	m_2/meV	m_3/meV		$m_{\beta\beta}/\text{meV}$
D_1-W_1	0.33989	0.02244	0.53179	1.11139	0.00474	0.05853	0.02989	1.06921	0.14655	27.84586	29.14261	56.98847	10.04516	8.11653
D_2-W_2	0.30933	0.02242	0.52707	1.09701	0.00474	0.05857	0.02906	0.00931	1.00870	63.29939	63.88046	80.93137	60.55211	2.99557
D_3-W_3	0.30895	0.02234	0.55798	1.24021	0.00474	0.05857	0.02917	0.14358	1.41214	18.10917	20.04600	53.49454	16.76825	1.86338
D_4-S_1	0.30993	0.02237	0.56318	1.24722	0.00474	0.05857	0.02923	1.30471	0.49605	27.07896	28.41074	57.10709	16.19580	1.68604
D_5-S_2	0.31005	0.02237	0.56302	1.33116	0.00474	0.05857	0.02924	0.14241	1.52824	10.96356	13.93197	51.45661	10.40884	3.15331
D_6-S_3	0.31244	0.02235	0.56370	1.15976	0.00474	0.05857	0.02925	1.01310	0.35970	2.57262	8.97320	50.32765	0.11038	1.03321
D_7-S_4	0.30972	0.02237	0.56394	1.19850	0.00474	0.05857	0.02924	1.05455	1.94805	9.32428	12.68236	51.13357	3.22886	1.07827
D_8-S_5	0.31123	0.02230	0.56132	1.47812	0.00474	0.05857	0.02921	0.05237	0.82489	12.08684	14.83212	51.72936	13.65139	1.35963

Table 7. The best fit values of the input parameters at the minimum of the χ^2 under the assumption of NO neutrino masses. We give the predictions for neutrino mixing angles $\theta_{12}^l, \theta_{13}^l, \theta_{23}^l$, and Dirac CP violating phase δ_{CP}^l as well as Majorana CP violating phases α_{21}, α_{31} , and the light neutrino masses $m_{1,2,3}$ and the effective mass $m_{\beta\beta}$ in neutrinoless double beta decay.

where the convention for $\mathbf{1}^l$ is the same as that in eq. (4.7). The modular weights of left- and right-handed quarks are denoted as k_{Q_L} and $k_{q_i^c}$ respectively. The superpotential for the quark masses is determined by the assignments of right-handed quarks and the values of $k_{Q_L} + k_{q_i^c}$,

$$\mathcal{W}_q = \sum_{i=1}^3 \sum_a \alpha_{i,a} q_i^c Q_L H_{u/d} Y_{\mathbf{3}a}^{(k_{Q_L} + k_{q_i^c})}, \quad (5.2)$$

where $\alpha_{i,a}$ are coupling constants. The general form of the (ij) entry of the quark mass matrix M_q is

$$(M_q)_{ij} = \sum_a \alpha_{i,a} v_{u/d} Y_{\mathbf{3}a, 3-\text{mod}(l_i+j+1,3)}^{(k_{Q_L} + k_{q_i^c})}. \quad (5.3)$$

Analogous to the charged lepton sector in section 4.1, if we are only concerned with modular forms up to weight 8, then the explicit form of R_i^q (i -th row of M_q) with $k_{Q_L} + k_{q_i^c} = 2, 4, 6, 8$ can be obtained from table 2 with the replacements $k_L + k_{E_i^c} \rightarrow k_{Q_L} + k_{q_i^c}$. As a result, the quark mass matrix can also take 220 possible structures. According to the number of free coupling parameters in quark mass matrix, we can divide the 220 possible quark mass matrices into four distinct classes: M_Q^I , M_Q^{II} , M_Q^{III} and M_Q^{IV} in which 3, 4, 5 and 6 couplings are introduced in the quark mass matrix respectively. The number of models are 20, 90, 90 and 20 respectively for these four classes.

As shown in above discussion, both up and down quark mass matrices can take 220 possible structures with proper weight and representation assignments of quark fields. In a concrete model where both up and down quark mass matrices are involved, the possible number of real couplings will range from 6 to 12 besides the complex modulus τ . There are totally 10 observables in quark sector, including six quark masses $m_{u,c,t}$, $m_{d,s,b}$, three quark mixing angles θ_{12}^q , θ_{13}^q , θ_{23}^q , and one quark CP violation phase δ_{CP}^q . The quark model will not be attractive if too many free parameters are required to fit the ten quark observables. Hence we only focus on the models with number of free parameters no more than 11. More specifically, we only consider the following four kinds of quark models.

(I) Quark model with 8 free real parameters including $\text{Re}(\tau)$ and $\text{Im}(\tau)$.

In this case, there are total 6 free couplings in the up and down quark mass matrices, which can only originated from the assignment $M_u \in M_Q^I$ and $M_d \in M_Q^I$. The triplet modular forms that enter the quark superpotentials in eq. (5.2) can only be $Y_{\mathbf{3}}^{(2)}$ and $Y_{\mathbf{3}}^{(4)}$. Combining the up and down quark mass matrices, we can obtain $20 \times 20 = 400$ models which have 8 free real parameters.

(II) Quark model with 9 free real parameters including $\text{Re}(\tau)$ and $\text{Im}(\tau)$.

For the quark models with 9 free parameters, the classifications of quark mass matrices should be

$$M_u \in M_Q^I, \quad M_d \in M_Q^{II}, \quad \text{or} \quad M_u \in M_Q^{II}, \quad M_d \in M_Q^I. \quad (5.4)$$

There are $20 \times 90 \times 2 = 3600$ distinct models with 9 free parameters.

(III) Quark model with 10 free real parameters including $\text{Re}(\tau)$ and $\text{Im}(\tau)$.

There are three types of the combinations of the up and down quark mass matrices in this case,

$$\begin{aligned}
 & M_u \in M_Q^{\text{I}}, \quad M_d \in M_Q^{\text{III}}, \\
 \text{or } & M_u \in M_Q^{\text{III}}, \quad M_d \in M_Q^{\text{I}}, \\
 \text{or } & M_u \in M_Q^{\text{II}}, \quad M_d \in M_Q^{\text{II}}.
 \end{aligned} \tag{5.5}$$

So there are total $20 \times 90 \times 2 + 90 \times 90 = 11700$ models with 10 free real parameters.

(IV) Quark model with 11 free real parameters including $\text{Re}(\tau)$ and $\text{Im}(\tau)$.

If there are 11 parameters in quark mass matrices, we find that they should satisfy the following conditions,

$$\begin{aligned}
 & M_u \in M_Q^{\text{I}}, \quad M_d \in M_Q^{\text{IV}}, \\
 \text{or } & M_u \in M_Q^{\text{IV}}, \quad M_d \in M_Q^{\text{I}}, \\
 \text{or } & M_u \in M_Q^{\text{II}}, \quad M_d \in M_Q^{\text{III}}, \\
 \text{or } & M_u \in M_Q^{\text{III}}, \quad M_d \in M_Q^{\text{II}}.
 \end{aligned} \tag{5.6}$$

Counting all the possibilities, we can obtain $2 \times 20 \times 20 + 2 \times 90 \times 90 = 17000$ models with 11 free real parameters.

Similar to the lepton sector, we need to find the phenomenologically viable quark models through the χ^2 analysis, the definition of the χ^2 function involves experimental data of the ratios of quark masses and quark mixing parameters,

$$\begin{aligned}
 m_u/m_c &= (1.9286 \pm 0.6017) \times 10^{-3}, & m_c/m_t &= (2.8213 \pm 0.1195) \times 10^{-3}, \\
 m_d/m_s &= (5.0523 \pm 0.6191) \times 10^{-2}, & m_s/m_b &= (1.8241 \pm 0.1005) \times 10^{-2}, \\
 \theta_{12}^q &= 0.22736 \pm 0.00073, & \theta_{13}^q &= 0.00349 \pm 0.00013, \\
 \theta_{23}^q &= 0.04015 \pm 0.00064, & \delta_{CP}^q/\pi &= 0.3845 \pm 0.0173.
 \end{aligned} \tag{5.7}$$

The central values and 1σ ranges at GUT scale are taken from ref. [69] with $\tan\beta = 10$ and the SUSY breaking scale $M_{\text{SUSY}} = 10 \text{ TeV}$. Analogous to the lepton sector, we use the central value of $m_t = 87.4555 \text{ GeV}$ and $m_b = 0.9682 \text{ GeV}$ at GUT scale to determine the overall coefficients of the quark mass matrices M_u and M_d respectively.

We scan the parameter space of the above quark models one by one with the `TMinuit` package. Unfortunately, we have not found a model with 8 parameters that is consistent with the experimental data, and the minimum of χ^2 is very large. There is also no model with 9 parameters which can accommodate the experimental data at 3σ confidence level. But some models with 9 parameters can achieve a relatively small χ^2 , and only the mixing angle θ_{23}^q marginally lies outside the 3σ allowed region. It's reasonable to regard these models as a good leading order approximation, we will give such an example below. For models with 10 or 11 parameters, we find thousands of models can accommodate the experimental data. It is too lengthy to show all these phenomenologically viable models,

consequently only some examples are provided in this work and they are given in section 6 within the complete models for leptons and quarks.

Here we present an example of the quark models with 9 parameters. In this model, the quark fields transform under modular symmetry A_4 as follows,

$$\begin{aligned}
 Q_L &\sim \mathbf{3}, \quad u^c \sim \mathbf{1}, \quad c^c \sim \mathbf{1}', \quad t^c \sim \mathbf{1}'', \quad d^c \sim \mathbf{1}'', \quad s^c \sim \mathbf{1}', \quad b^c \sim \mathbf{1}'', \\
 k_{Q_L} &= 2 - k_{u^c} = 2 - k_{c^c} = 2 - k_{t^c} = 2 - k_{d^c} = 4 - k_{s^c} = 6 - k_{b^c},
 \end{aligned} \tag{5.8}$$

where the modular weight k_{Q_L} is a general integer. The modular invariant superpotentials are given as

$$\begin{aligned}
 \mathcal{W}_u &= \alpha_u u_1^c (Q_L Y_{\mathbf{3}}^{(2)})_1 H_u + \beta_u c_{1'}^c (Q_L Y_{\mathbf{3}}^{(2)})_{1''} H_u + \gamma_u t_{1''}^c (Q_L Y_{\mathbf{3}'}^{(2)})_{1'} H_u, \\
 \mathcal{W}_d &= \alpha_d d_{1''}^c (Q_L Y_{\mathbf{3}}^{(2)})_{1'} H_d + \beta_d s_{1'}^c (Q_L Y_{\mathbf{3}}^{(4)})_{1''} H_d + \gamma_{d,1} b_{1''}^c (Q_L Y_{\mathbf{3}I}^{(6)})_{1'} H_d \\
 &\quad + \gamma_{d,2} b_{1''}^c (Q_L Y_{\mathbf{3}II}^{(6)})_{1'} H_d.
 \end{aligned} \tag{5.9}$$

The corresponding up and down quark mass matrices can be written as

$$\begin{aligned}
 M_u &= \begin{pmatrix} \alpha_u Y_{\mathbf{3},1}^{(2)} & \alpha_u Y_{\mathbf{3},3}^{(2)} & \alpha_u Y_{\mathbf{3},2}^{(2)} \\ \beta_u Y_{\mathbf{3},3}^{(2)} & \beta_u Y_{\mathbf{3},2}^{(2)} & \beta_u Y_{\mathbf{3},1}^{(2)} \\ \gamma_u Y_{\mathbf{3},2}^{(2)} & \gamma_u Y_{\mathbf{3},1}^{(2)} & \gamma_u Y_{\mathbf{3},3}^{(2)} \end{pmatrix} v_u, \\
 M_d &= \begin{pmatrix} \alpha_d Y_{\mathbf{3},2}^{(2)} & \alpha_d Y_{\mathbf{3},1}^{(2)} & \alpha_d Y_{\mathbf{3},3}^{(2)} \\ \beta_d Y_{\mathbf{3},1}^{(4)} & \beta_d Y_{\mathbf{3},3}^{(4)} & \beta_d Y_{\mathbf{3},2}^{(4)} \\ \gamma_{d,1} Y_{\mathbf{3}I,2}^{(6)} + \gamma_{d,2} Y_{\mathbf{3}II,2}^{(6)} & \gamma_{d,1} Y_{\mathbf{3}I,1}^{(6)} + \gamma_{d,2} Y_{\mathbf{3}II,1}^{(6)} & \gamma_{d,1} Y_{\mathbf{3}I,3}^{(6)} + \gamma_{d,2} Y_{\mathbf{3}II,3}^{(6)} \end{pmatrix} v_d.
 \end{aligned} \tag{5.10}$$

The best fit values of the input parameters are given as follows

$$\begin{aligned}
 \langle \tau \rangle &= 0.49175 + 0.88563i, & \beta_u/\alpha_u &= 518.22933, & \gamma_u/\alpha_u &= 1.83596 \times 10^5, \\
 \beta_d/\alpha_d &= 9.39751, & \gamma_{d,1}/\alpha_d &= 32.46046, & \gamma_{d,2}/\alpha_d &= -0.02697, \\
 \alpha_u v_u &= 0.00034 \text{ GeV}, & \alpha_d v_d &= 0.05081 \text{ GeV}.
 \end{aligned} \tag{5.11}$$

The quark mixing parameters and mass ratios are determined to be

$$\begin{aligned}
 \theta_{12}^q &= 0.22734, & \theta_{13}^q &= 0.00332, & \theta_{23}^q &= 0.05708, & \delta_{CP}^q &= 0.39532 \pi, \\
 m_u/m_c &= 0.00193, & m_c/m_t &= 0.00282, & m_d/m_s &= 0.05055, & m_s/m_b &= 0.01815.
 \end{aligned} \tag{5.12}$$

We find that only θ_{23}^q is somewhat large, all other observables are within the 3σ ranges of the experimental data given in eq. (5.7). Notice that if we adopt the experimental data of quark masses and mixings used in ref. [27], this model would be compatible with data very well.⁴

⁴In ref. [27], the central values as well the 1σ errors of the quark masses and mixing parameters are taken from [73]. Comparing the experimental data used in ref. [27] with those in the present work, we can

6 Unified models of leptons and quarks

In the above two sections, we have discussed the phenomenologically viable models in lepton sector and quark sector separately. Now we shall investigate whether the A_4 modular can describe the flavor structures of quark and lepton simultaneously, where the complex modulus τ works as a portal to combine these two sectors. All coupling constants are real because of the gCP symmetry, thus the vacuum expectation value of τ is the only source of CP violation phases of both quarks and leptons.

In section 4, we have found 20 lepton models with 7 parameters and 360 lepton models with 8 parameters that can accommodate the experimental data. In section 5, we have found thousands of quark models with 10 or 11 parameters that can explain the experimental data of quark masses and mixing. In each sector, the experimentally 3σ -allowed τ s are recorded in the `TMinuit` optimization processes. By showing the τ sample of lepton sector and quark sector in the same plot, we can see whether there are overlapping regions. In this way, we can easily determine whether the experimental data of quarks and leptons can be described in a given combination of lepton and quark models. It turns out that the lepton models with 7 parameters (L_7) together with the quark models with 10 parameters (Q_{10}) failed to accommodate the experimental data, while the combinations of L_7 - Q_{11} and L_8 - Q_{10} can explain the measured masses and mixings of both quarks and leptons. We can find thousands of these two kinds of unified models. Due to the limitation of the length of the article, we only give one example of L_7 - Q_{11} unification and one example of L_8 - Q_{10} unification in the following.

6.1 Unified model of L_7 - Q_{11}

In this section, we give an example of the unified model with 7 parameters in the lepton sector and 11 parameters in the quark sector. The complex modulus τ in the quark and lepton sectors is the same one, so the total number of real free parameters is $7+11-2=16$. Thus this model is relatively predictive, and it use 16 real free parameters to describe the 22 mass and mixing parameters of quarks and leptons. In the lepton sector, the model is denoted as C_{13} - S_2 , which has been described in section 4.3. For the sake of clarity, we list the details of the model in the following. The representation assignments and the modular weights of the lepton fields are,

$$\begin{aligned} L &\sim \mathbf{3}, & e^c &\sim \mathbf{1}', & \mu^c &\sim \mathbf{1}', & \tau^c &\sim \mathbf{1}'', & N^c &\sim \mathbf{3}, \\ k_L &= -1, & k_{e^c} &= 3, & k_{\mu^c} &= 5, & k_{\tau^c} &= 7, & k_{N^c} &= 1. \end{aligned} \quad (6.1)$$

The corresponding modular invariant lepton superpotentials are given by

$$\begin{aligned} \mathcal{W}_E &= \alpha_e e_{\mathbf{1}'}^c (LY_{\mathbf{3}}^{(2)})_{\mathbf{1}''} H_d + \beta_e \mu_{\mathbf{1}'}^c (LY_{\mathbf{3}}^{(4)})_{\mathbf{1}''} H_d + \gamma_{e,1} \tau_{\mathbf{1}''}^c (LY_{\mathbf{3}I}^{(6)})_{\mathbf{1}'} H_d + \gamma_{e,2} \tau_{\mathbf{1}''}^c (LY_{\mathbf{3}II}^{(6)})_{\mathbf{1}'} H_d, \\ \mathcal{W}_\nu &= g^D H_u (N^c L)_{\mathbf{1}} + g^M \Lambda \left((N^c N^c)_{\mathbf{3}_S} Y_{\mathbf{3}}^{(2)} \right)_{\mathbf{1}}. \end{aligned} \quad (6.2)$$

find that the best-fit values of quark Yukawa couplings and the CKM mixing parameters are almost the same while the 1σ ranges given in [27] are larger than the corresponding ones used in our paper. To be more specific, the 1σ errors of θ_{12}^q and θ_{23}^q in [27] are about 5 and 10 times as large as the corresponding values in this paper respectively, while the 1σ regions of other observables are about twice as large as those in eq. (5.7).

The corresponding charged lepton and neutrino mass matrices read as

$$M_E = \begin{pmatrix} \alpha_e Y_{\mathbf{3},\mathbf{3}}^{(2)} & \alpha_e Y_{\mathbf{3},\mathbf{2}}^{(2)} & \alpha_e Y_{\mathbf{3},\mathbf{1}}^{(2)} \\ \beta_e Y_{\mathbf{3},\mathbf{3}}^{(4)} & \beta_e Y_{\mathbf{3},\mathbf{2}}^{(4)} & \beta_e Y_{\mathbf{3},\mathbf{1}}^{(4)} \\ \gamma_{e,1} Y_{\mathbf{3}I,\mathbf{2}}^{(6)} + \gamma_{e,2} Y_{\mathbf{3}II,\mathbf{2}}^{(6)} & \gamma_{e,1} Y_{\mathbf{3}I,\mathbf{1}}^{(6)} + \gamma_{e,2} Y_{\mathbf{3}II,\mathbf{1}}^{(6)} & \gamma_{e,1} Y_{\mathbf{3}I,\mathbf{3}}^{(6)} + \gamma_{e,2} Y_{\mathbf{3}II,\mathbf{3}}^{(6)} \end{pmatrix} v_d,$$

$$M_D = g^D \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} v_u, \quad M_N = g^M \begin{pmatrix} 2Y_{\mathbf{3},\mathbf{1}}^{(2)} & -Y_{\mathbf{3},\mathbf{3}}^{(2)} & -Y_{\mathbf{3},\mathbf{2}}^{(2)} \\ -Y_{\mathbf{3},\mathbf{3}}^{(2)} & 2Y_{\mathbf{3},\mathbf{2}}^{(2)} & -Y_{\mathbf{3},\mathbf{1}}^{(2)} \\ -Y_{\mathbf{3},\mathbf{2}}^{(2)} & -Y_{\mathbf{3},\mathbf{1}}^{(2)} & 2Y_{\mathbf{3},\mathbf{3}}^{(2)} \end{pmatrix} \Lambda. \quad (6.3)$$

The quark fields transform under the A_4 modular symmetry as follow,

$$Q_L \sim \mathbf{3}, u^c \sim \mathbf{1}, \quad c^c \sim \mathbf{1}, \quad t^c \sim \mathbf{1}, \quad d^c \sim \mathbf{1}, \quad s^c \sim \mathbf{1}', \quad b^c \sim \mathbf{1}',$$

$$k_{Q_L} = 2 - k_{u^c} = 4 - k_{c^c} = 6 - k_{t^c} = 4 - k_{d^c} = 6 - k_{s^c} = 8 - k_{b^c}. \quad (6.4)$$

Then the Yukawa superpotentials for the quark masses are

$$\begin{aligned}
 \mathcal{W}_u &= \alpha_u u_1^c (Q_L Y_{\mathbf{3}}^{(2)})_1 H_u + \beta_u c_1^c (Q_L Y_{\mathbf{3}}^{(4)})_1 H_u + \gamma_{u,1} t_1^c (Q_L Y_{\mathbf{3}I}^{(6)})_1 H_u \\
 &\quad + \gamma_{u,2} t_1^c (Q_L Y_{\mathbf{3}II}^{(6)})_1 H_u, \\
 \mathcal{W}_d &= \alpha_d d_1^c (Q_L Y_{\mathbf{3}}^{(4)})_1 H_d + \beta_{d,1} s_{1'}^c (Q_L Y_{\mathbf{3}I}^{(6)})_{1''} H_d + \beta_{d,2} s_{1'}^c (Q_L Y_{\mathbf{3}II}^{(6)})_{1''} H_d \\
 &\quad + \gamma_{d,1} b_{1'}^c (Q_L Y_{\mathbf{3}I}^{(8)})_{1''} H_d + \gamma_{d,2} b_{1'}^c (Q_L Y_{\mathbf{3}II}^{(8)})_{1''} H_d.
 \end{aligned} \quad (6.5)$$

We find the up and down quark mass matrices take the following form

$$M_u = \begin{pmatrix} \alpha_u Y_{\mathbf{3},\mathbf{1}}^{(2)} & \alpha_u Y_{\mathbf{3},\mathbf{3}}^{(2)} & \alpha_u Y_{\mathbf{3},\mathbf{2}}^{(2)} \\ \beta_u Y_{\mathbf{3},\mathbf{1}}^{(4)} & \beta_u Y_{\mathbf{3},\mathbf{3}}^{(4)} & \beta_u Y_{\mathbf{3},\mathbf{2}}^{(4)} \\ \gamma_{u,1} Y_{\mathbf{3}I,\mathbf{1}}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,\mathbf{1}}^{(6)} & \gamma_{u,1} Y_{\mathbf{3}I,\mathbf{3}}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,\mathbf{3}}^{(6)} & \gamma_{u,1} Y_{\mathbf{3}I,\mathbf{2}}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,\mathbf{2}}^{(6)} \end{pmatrix} v_u,$$

$$M_d = \begin{pmatrix} \alpha_d Y_{\mathbf{3},\mathbf{1}}^{(4)} & \alpha_d Y_{\mathbf{3},\mathbf{3}}^{(4)} & \alpha_d Y_{\mathbf{3},\mathbf{2}}^{(4)} \\ \beta_{d,1} Y_{\mathbf{3}I,\mathbf{3}}^{(6)} + \beta_{d,2} Y_{\mathbf{3}II,\mathbf{3}}^{(6)} & \beta_{d,1} Y_{\mathbf{3}I,\mathbf{2}}^{(6)} + \beta_{d,2} Y_{\mathbf{3}II,\mathbf{2}}^{(6)} & \beta_{d,1} Y_{\mathbf{3}I,\mathbf{1}}^{(6)} + \beta_{d,2} Y_{\mathbf{3}II,\mathbf{1}}^{(6)} \\ \gamma_{d,1} Y_{\mathbf{3}I,\mathbf{3}}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,\mathbf{3}}^{(8)} & \gamma_{d,1} Y_{\mathbf{3}I,\mathbf{2}}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,\mathbf{2}}^{(8)} & \gamma_{d,1} Y_{\mathbf{3}I,\mathbf{1}}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,\mathbf{1}}^{(8)} \end{pmatrix} v_d. \quad (6.6)$$

We perform a global fit to the experimental data of quark and lepton simultaneously, the best fit values of the input parameters are determined to be

$$\begin{aligned}
 \langle \tau \rangle &= -0.17492 + 1.13648i, & \beta_e / \alpha_e &= 0.00500, & \gamma_{e,1} / \alpha_e &= 11.54025, \\
 \gamma_{e,2} / \alpha_e &= 6.27248, & \alpha_e v_d &= 0.06688 \text{ MeV}, & \frac{(g^D)^2 v_u^2}{g^M \Lambda} &= 23.94439 \text{ meV}, \\
 \beta_u / \alpha_u &= 4.62725 \times 10^2, & \gamma_{u,1} / \alpha_u &= 7.66928 \times 10^3, & \gamma_{u,2} / \alpha_u &= -1.51202 \times 10^5, \\
 \beta_{d,1} / \alpha_d &= 2.35800 \times 10^2, & \beta_{d,2} / \alpha_d &= 14.19700, & \gamma_{d,1} / \alpha_d &= 94.91290, \\
 \gamma_{d,2} / \alpha_d &= 3.84761, & \alpha_u v_u &= 0.00046 \text{ GeV}, & \alpha_d v_d &= 0.00298 \text{ GeV}.
 \end{aligned} \quad (6.7)$$

We see that the quark and lepton mass hierarchies require hierarchical values of the coupling constants, this can be naturally achieved by the weighton mechanism in which modular symmetry singlet field with non-vanishing modular weight plays the role of flavon [25]. The mass hierarchies of the charged leptons could possibly be addressed by modular symmetry around certain fixed points of the complex modulus τ , but generally more free parameters are necessary [30, 74, 75]. The corresponding predictions for masses and mixing parameters of leptons and quarks are given as

$$\begin{aligned}
 \sin^2 \theta_{12}^l &= 0.33632, & \sin^2 \theta_{13}^l &= 0.02164, & \sin^2 \theta_{23}^l &= 0.57884, & \delta_{CP}^l &= 0.98516 \pi, \\
 \alpha_{21} &= 1.05680 \pi, & \alpha_{31} &= 0.14266 \pi, & m_e/m_\mu &= 0.00474, & m_\mu/m_\tau &= 0.05857, \\
 \frac{\Delta m_{21}^2}{\Delta m_{31}^2} &= 0.03040, & m_1 &= 10.87722 \text{ meV}, & m_2 &= 13.86412 \text{ meV}, \\
 m_3 &= 50.48811 \text{ meV}, & \sum_i m_i &= 75.22945 \text{ meV}, & m_{\beta\beta} &= 3.51842 \text{ meV}, \\
 \theta_{12}^q &= 0.22719, & \theta_{13}^q &= 0.00350, & \theta_{23}^q &= 0.03988, & \delta_{CP}^q &= 0.38850 \pi, \\
 m_u/m_c &= 0.00166, & m_c/m_t &= 0.00289, & m_d/m_s &= 0.04605, & m_s/m_b &= 0.01837.
 \end{aligned} \tag{6.8}$$

We find that all of these observables are predicted to lie in the experimentally preferred 3σ ranges. We use the package `Multinest` to comprehensively scan the parameter space of lepton sector and quark sector independently, we require all the observables are in the experimentally preferred 3σ regions, and the results are plotted in figure 2. We can see there is indeed an overlapping region in the τ plane, which is colored in black. It is obvious that the allowed value of τ shrink to a small region if we include the experimental constraints of both quarks and leptons. The allowed values of all observables scatter in narrow ranges around the best fit values.

6.2 Unified model of L_8 - Q_{10}

In this section, we give another example of the unified model with 8 parameters in the lepton sector and 10 parameters in the quark sector. The total number of real free parameters in this unified model is also $8+10-2 = 16$. The lepton model is D_8 - S_5 , and the transformation properties of the lepton fields are,

$$\begin{aligned}
 L &\sim \mathbf{3}, & e^c &\sim \mathbf{1}, & \mu^c &\sim \mathbf{1}', & \tau^c &\sim \mathbf{1}'', & N^c &\sim \mathbf{3}, \\
 k_L &= -3, & k_{e^c} &= 7, & k_{\mu^c} &= 7, & k_{\tau^c} &= 7, & k_{N^c} &= 3.
 \end{aligned} \tag{6.9}$$

Then we can read out the superpotentials of charged leptons and neutrinos as

$$\begin{aligned}
 \mathcal{W}_E &= \alpha_e e_1^c (LY_{\mathbf{3}}^{(4)})_{\mathbf{1}} H_d + \beta_e \mu_{1'}^c (LY_{\mathbf{3}}^{(4)})_{\mathbf{1}'} H_d + \gamma_e \tau_{1''}^c (LY_{\mathbf{3}}^{(4)})_{\mathbf{1}''} H_d, \\
 \mathcal{W}_\nu &= g^D H_u (N^c L)_{\mathbf{1}} + g_1^M \Lambda \left((N^c N^c)_{\mathbf{3}_S} Y_{\mathbf{3}_I}^{(6)} \right)_{\mathbf{1}} + g_2^M \Lambda \left((N^c N^c)_{\mathbf{3}_S} Y_{\mathbf{3}_{II}}^{(6)} \right)_{\mathbf{1}} \\
 &\quad + g_3^M \Lambda (N^c N^c)_{\mathbf{1}} Y_{\mathbf{1}}^{(6)},
 \end{aligned} \tag{6.10}$$




Figure 2. The predictions for the correlations among the input free parameters, neutrino mixing angles, CP violation phases and neutrino masses in the unified model L_7-Q_{11} . The red and blue areas denote points compatible with experimental data of quarks and leptons respectively, and the black is the overlapping region. The vertical and horizontal dashed lines are the 3σ bounds taken from [68].

which gives rise to

$$\begin{aligned}
 M_E &= \begin{pmatrix} \alpha_e Y_{\mathbf{3},1}^{(4)} & \alpha_e Y_{\mathbf{3},3}^{(4)} & \alpha_e Y_{\mathbf{3},2}^{(4)} \\ \beta_e Y_{\mathbf{3},3}^{(4)} & \beta_e Y_{\mathbf{3},2}^{(4)} & \beta_e Y_{\mathbf{3},1}^{(4)} \\ \gamma_e Y_{\mathbf{3},2}^{(4)} & \gamma_e Y_{\mathbf{3},1}^{(4)} & \gamma_e Y_{\mathbf{3},3}^{(4)} \end{pmatrix} v_d, \quad M_D = g^D \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} v_u, \\
 M_N &= \left[g_1^M \begin{pmatrix} 2Y_{\mathbf{3}I,1}^{(6)} & -Y_{\mathbf{3}I,3}^{(6)} & -Y_{\mathbf{3}I,2}^{(6)} \\ -Y_{\mathbf{3}I,3}^{(6)} & 2Y_{\mathbf{3}I,2}^{(6)} & -Y_{\mathbf{3}I,1}^{(6)} \\ -Y_{\mathbf{3}I,2}^{(6)} & -Y_{\mathbf{3}I,1}^{(6)} & 2Y_{\mathbf{3}I,3}^{(6)} \end{pmatrix} + g_2^M \begin{pmatrix} 2Y_{\mathbf{3}II,1}^{(6)} & -Y_{\mathbf{3}II,3}^{(6)} & -Y_{\mathbf{3}II,2}^{(6)} \\ -Y_{\mathbf{3}II,3}^{(6)} & 2Y_{\mathbf{3}II,2}^{(6)} & -Y_{\mathbf{3}II,1}^{(6)} \\ -Y_{\mathbf{3}II,2}^{(6)} & -Y_{\mathbf{3}II,1}^{(6)} & 2Y_{\mathbf{3}II,3}^{(6)} \end{pmatrix} \right. \\
 &\quad \left. + g_3^M Y_{\mathbf{1}}^{(6)} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right] \Lambda. \tag{6.11}
 \end{aligned}$$

The assignments for the quark fields are

$$\begin{aligned}
 Q_L &\sim \mathbf{3}, \quad u^c \sim \mathbf{1}, \quad c^c \sim \mathbf{1}, \quad t^c \sim \mathbf{1}, \quad d^c \sim \mathbf{1}, \quad s^c \sim \mathbf{1}', \quad b^c \sim \mathbf{1}'', \\
 k_{Q_L} &= 2 - k_{u^c} = 4 - k_{c^c} = 6 - k_{t^c} = 2 - k_{d^c} = 2 - k_{s^c} = 8 - k_{b^c}. \tag{6.12}
 \end{aligned}$$

The modular invariant superpotentials for quark masses read as follows,

$$\begin{aligned}
 \mathcal{W}_u &= \alpha_u u_{\mathbf{1}}^c (Q_L Y_{\mathbf{3}}^{(2)})_{\mathbf{1}} H_u + \beta_u c_{\mathbf{1}}^c (Q_L Y_{\mathbf{3}}^{(4)})_{\mathbf{1}} H_u + \gamma_{u,1} t_{\mathbf{1}}^c (Q_L Y_{\mathbf{3}I}^{(6)})_{\mathbf{1}} H_u \\
 &\quad + \gamma_{u,2} (t_{\mathbf{1}}^c (Q_L Y_{\mathbf{3}II}^{(6)})_{\mathbf{1}})_{\mathbf{1}} H_u, \\
 \mathcal{W}_d &= \alpha_d d_{\mathbf{1}}^c (Q_L Y_{\mathbf{3}}^{(2)})_{\mathbf{1}} H_d + \beta_d s_{\mathbf{1}'}^c (Q_L Y_{\mathbf{3}}^{(2)})_{\mathbf{1}'} H_d + \gamma_{d,1} b_{\mathbf{1}''}^c (Q_L Y_{\mathbf{3}I}^{(8)})_{\mathbf{1}'} H_d \\
 &\quad + \gamma_{d,2} b_{\mathbf{1}''}^c (Q_L Y_{\mathbf{3}II}^{(8)})_{\mathbf{1}'} H_d. \tag{6.13}
 \end{aligned}$$

We find the up and down quark mass matrices are given by

$$\begin{aligned}
 M_u &= \begin{pmatrix} \alpha_u Y_{\mathbf{3},1}^{(2)} & \alpha_u Y_{\mathbf{3},3}^{(2)} & \alpha_u Y_{\mathbf{3},2}^{(2)} \\ \beta_u Y_{\mathbf{3},3}^{(4)} & \beta_u Y_{\mathbf{3},2}^{(4)} & \beta_u Y_{\mathbf{3},1}^{(4)} \\ \gamma_{u,1} Y_{\mathbf{3}I,1}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,1}^{(6)} & \gamma_{u,1} Y_{\mathbf{3}I,3}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,3}^{(6)} & \gamma_{u,1} Y_{\mathbf{3}I,2}^{(6)} + \gamma_{u,2} Y_{\mathbf{3}II,2}^{(6)} \end{pmatrix} v_u, \\
 M_d &= \begin{pmatrix} \alpha_d Y_{\mathbf{3},1}^{(2)} & \alpha_d Y_{\mathbf{3},3}^{(2)} & \alpha_d Y_{\mathbf{3},2}^{(2)} \\ \beta_d Y_{\mathbf{3},3}^{(2)} & \beta_d Y_{\mathbf{3},2}^{(2)} & \beta_d Y_{\mathbf{3},1}^{(2)} \\ \gamma_{d,1} Y_{\mathbf{3}I,2}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,2}^{(8)} & \gamma_{d,1} Y_{\mathbf{3}I,1}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,1}^{(8)} & \gamma_{d,1} Y_{\mathbf{3}I,3}^{(8)} + \gamma_{d,2} Y_{\mathbf{3}II,3}^{(8)} \end{pmatrix} v_d. \tag{6.14}
 \end{aligned}$$

The agreement between the model predictions and the experimental data is optimized for the following values of the input parameters are

$$\begin{aligned}
 \langle \tau \rangle &= -0.30537 + 1.77322i, & \beta_e / \alpha_e &= 3.53453 \times 10^3, & \gamma_e / \alpha_e &= 209.08250, \\
 g_2^M / g_1^M &= -2.66807, & g_3^M / g_1^M &= -0.07778, & \alpha_e v_d &= 0.36424 \text{ MeV}, \\
 \frac{(g^D)^2 v_u^2}{g_1^M \Lambda} &= 26.95432 \text{ meV}, & \beta_u / \alpha_u &= 43.07455, & \gamma_{u,1} / \alpha_u &= 634.51584, \\
 \gamma_{u,2} / \alpha_u &= 5.16218 \times 10^4, & \beta_d / \alpha_d &= 232.29418, & \gamma_{d,1} / \alpha_d &= 0.99699, \\
 \gamma_{d,2} / \alpha_d &= -14.39314, & \alpha_u v_u &= 0.00569 \text{ GeV}, & \alpha_d v_d &= 0.00412 \text{ GeV}. \tag{6.15}
 \end{aligned}$$

The values of the masses and mixing parameters of leptons and quarks at the above best fit point are determined to be

$$\begin{aligned}
 \sin^2 \theta_{12}^l &= 0.31123, & \sin^2 \theta_{13}^l &= 0.02230, & \sin^2 \theta_{23}^l &= 0.56132, & \delta_{CP}^l &= 1.47812 \pi, \\
 \alpha_{21} &= 0.05237 \pi, & \alpha_{31} &= 0.82489 \pi, & m_e/m_\mu &= 0.00474, & m_\mu/m_\tau &= 0.05857, \\
 \frac{\Delta m_{21}^2}{\Delta m_{31}^2} &= 0.02921, & m_1 &= 12.08684 \text{ meV}, & m_2 &= 14.83212 \text{ meV}, \\
 m_3 &= 51.72936 \text{ meV}, & \sum_i m_i &= 89.64832 \text{ meV}, & m_{\beta\beta} &= 13.65139 \text{ meV}, \\
 \theta_{12}^q &= 0.22736, & \theta_{13}^q &= 0.00348, & \theta_{23}^q &= 0.04119, & \delta_{CP}^q &= 0.38606 \pi, \\
 m_u/m_c &= 0.00193, & m_c/m_t &= 0.00282, & m_d/m_s &= 0.05057, & m_s/m_b &= 0.01824,
 \end{aligned}
 \tag{6.16}$$

which are in the experimentally favored 3σ regions. In figure 3, we show the correlations among the input free parameters, neutrino masses and mixing parameters predicted in the unified model L_8-Q_{10} . It is notable that the lepton Dirac CP phase δ_{CP}^l is close to maximal violation value 1.5π in this model.

7 Conclusion

In this paper, we have performed a systematical analysis of $\Gamma_3 \cong A_4$ modular flavor models for leptons and quarks with gCP. We intend to understand how the minimal modular symmetry A_4 can help to understand the flavor structure of quarks and leptons, and we aim to find out the minimal models. It has been established that the CP transformation consistent with modular symmetry acts on the modulus as $\tau \rightarrow -\tau^*$. The representation matrices of the generators S and T are unitary and symmetric in our working basis, therefore the CP symmetry is exactly the canonical one and all the coupling constants are constrained to be real. In this setup, both modular symmetry and gCP are spontaneously broken by the vacuum expectation value of the complex modulus τ . In particular, all weak CP violation phases arise from the real part of τ .

The left-handed charged fermions are assumed to transform as triplet of A_4 , and the right-handed charged fermions are assigned to singlets under A_4 . Here the charged fermions refer to charged leptons, up-type quarks and down-type quarks. We find out the most general form of charged lepton and quark mass matrices given by eq. (4.10) and eq. (5.3) respectively. The three generations of right-handed charged fermions are distinguished from each other by the weight and representation assignments, then we can obtain 220 possible structures of charged fermion mass matrices if the weights of involved modular forms are less than 9. Moreover, we consider the Weinberg operator and the type-I seesaw mechanism to generate neutrino masses. In type-I seesaw mechanism, the right-handed neutrinos are assigned to be a A_4 triplet as well. The most general form of neutrino mass matrix is given in eqs. (4.20), (4.22). Requiring the number of coupling constants in the effective light neutrino mass matrix is less than 4, we can obtain 3 Weinberg operator models $W_{1,2,3}$ and 5 seesaw models $S_{1,2,3,4,5}$, they are summarized in table 3. We perform




Figure 3. The predictions for the correlations among the input free parameters, neutrino mixing angles, CP violation phases and neutrino masses of the unified model L_8-Q_{10} . We adopt the same convention as figure 2.

a χ^2 analysis to estimate the goodness-of-fit of a A_4 modular model to the data. We find the minimal phenomenologically viable models for leptons have 7 real free parameters including the real and imaginary parts of τ . The best fit values of the input parameters and the corresponding predictions for neutrino masses and mixing parameters are given in table 5. For the next-to-minimal models with 8 real input parameters, we find 360 models which can explain the observed masses and mixings of leptons, and the numerical results of some typical models are presented in table 7.

In quark sector, we have systematically classified the A_4 modular quark models based on the number of free parameters. All the quark models with 8, 9, 10 and 11 input

parameters are discussed. There are no viable quark models with 8 or 9 parameters. However, we find thousands of quark models which can accommodate the experimental data of quark masses and CKM mixing matrix with 10 or 11 parameters. They can combine with the lepton models to give a unified description of both quarks and leptons. We give two predictive examples of quark-lepton unification models in which the 22 mass and mixing parameters of quarks and leptons are described in terms of 14 free real coupling constants and a common complex modulus τ .

Because the weights and representations of the matter fields are not subject to any constraint in the modular invariance approach, many possible models could be constructed, as explicit shown for the A_4 modular symmetry. We expect this drawback could possibly be overcome in the appealing framework of eclectic flavor groups which is a maximal extension of the traditional flavor group by finite discrete modular symmetries [61, 62]. The scheme of eclectic flavor symmetry can severely restrict the possible group representations and modular weights of matter fields. Thus the structure of both Kähler potential and superpotential is under control. It would be much desirable that only few models in the bottom-up approach would survive in view of the eclectic flavor symmetry. In summary, the finite modular group A_4 provides a simple and economical framework to understand the flavor structures of the quarks and leptons.

Acknowledgments

JNL and GJD are supported by the National Natural Science Foundation of China under Grant Nos. 11975224, 11835013 and 11947301. CYX is supported in part by the Grants No. NSFC-11975130, No. NSFC-12035008, No. NSFC-12047533, by the National Key Research and Development Program of China under Grant No. 2017YFA0402200 and the China Post-doctoral Science Foundation under Grant No. 2018M641621. We are grateful to Prof. Jörn Kersten for useful communication on the effect of renormalization group evolution.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] F. Feruglio and A. Romanino, *Lepton Flavour Symmetries*, [arXiv:1912.06028](https://arxiv.org/abs/1912.06028) [[INSPIRE](https://inspirehep.net/literature/1912060)].
- [2] F. Feruglio, *Are neutrino masses modular forms?*, in *From My Vast Repertoire...: Guido Altarelli's Legacy*, A. Levy, S. Forte and G. Ridolfi eds., World Scientific (2018), chapter 12, pp. 227–266 [[arXiv:1706.08749](https://arxiv.org/abs/1706.08749)] [[INSPIRE](https://inspirehep.net/literature/1706087)].
- [3] T. Kobayashi, K. Tanaka and T.H. Tatsuishi, *Neutrino mixing from finite modular groups*, *Phys. Rev. D* **98** (2018) 016004 [[arXiv:1803.10391](https://arxiv.org/abs/1803.10391)] [[INSPIRE](https://inspirehep.net/literature/1803103)].
- [4] T. Kobayashi, Y. Shimizu, K. Takagi, M. Tanimoto, T.H. Tatsuishi and H. Uchida, *Finite modular subgroups for fermion mass matrices and baryon/lepton number violation*, *Phys. Lett. B* **794** (2019) 114 [[arXiv:1812.11072](https://arxiv.org/abs/1812.11072)] [[INSPIRE](https://inspirehep.net/literature/1812110)].

- [5] T. Kobayashi, Y. Shimizu, K. Takagi, M. Tanimoto and T.H. Tatsuishi, *Modular S_3 -invariant flavor model in SU(5) grand unified theory*, *Prog. Theor. Exp. Phys.* **2020** (2020) 053B05 [[arXiv:1906.10341](#)] [[INSPIRE](#)].
- [6] H. Okada and Y. Orikasa, *Modular S_3 symmetric radiative seesaw model*, *Phys. Rev. D* **100** (2019) 115037 [[arXiv:1907.04716](#)] [[INSPIRE](#)].
- [7] J.C. Criado and F. Feruglio, *Modular Invariance Faces Precision Neutrino Data*, *SciPost Phys.* **5** (2018) 042 [[arXiv:1807.01125](#)] [[INSPIRE](#)].
- [8] T. Kobayashi, N. Omoto, Y. Shimizu, K. Takagi, M. Tanimoto and T.H. Tatsuishi, *Modular A_4 invariance and neutrino mixing*, *JHEP* **11** (2018) 196 [[arXiv:1808.03012](#)] [[INSPIRE](#)].
- [9] F.J. de Anda, S.F. King and E. Perdomo, *SU(5) grand unified theory with A_4 modular symmetry*, *Phys. Rev. D* **101** (2020) 015028 [[arXiv:1812.05620](#)] [[INSPIRE](#)].
- [10] H. Okada and M. Tanimoto, *CP violation of quarks in A_4 modular invariance*, *Phys. Lett. B* **791** (2019) 54 [[arXiv:1812.09677](#)] [[INSPIRE](#)].
- [11] P.P. Novichkov, S.T. Petcov and M. Tanimoto, *Trimaximal Neutrino Mixing from Modular A_4 Invariance with Residual Symmetries*, *Phys. Lett. B* **793** (2019) 247 [[arXiv:1812.11289](#)] [[INSPIRE](#)].
- [12] T. Nomura and H. Okada, *A modular A_4 symmetric model of dark matter and neutrino*, *Phys. Lett. B* **797** (2019) 134799 [[arXiv:1904.03937](#)] [[INSPIRE](#)].
- [13] H. Okada and M. Tanimoto, *Towards unification of quark and lepton flavors in A_4 modular invariance*, *Eur. Phys. J. C* **81** (2021) 52 [[arXiv:1905.13421](#)] [[INSPIRE](#)].
- [14] T. Nomura and H. Okada, *A two loop induced neutrino mass model with modular A_4 symmetry*, *Nucl. Phys. B* **966** (2021) 115372 [[arXiv:1906.03927](#)] [[INSPIRE](#)].
- [15] G.-J. Ding, S.F. King and X.-G. Liu, *Modular A_4 symmetry models of neutrinos and charged leptons*, *JHEP* **09** (2019) 074 [[arXiv:1907.11714](#)] [[INSPIRE](#)].
- [16] H. Okada and Y. Orikasa, *A radiative seesaw model in modular A_4 symmetry*, [arXiv:1907.13520](#) [[INSPIRE](#)].
- [17] T. Nomura, H. Okada and O. Popov, *A modular A_4 symmetric scotogenic model*, *Phys. Lett. B* **803** (2020) 135294 [[arXiv:1908.07457](#)] [[INSPIRE](#)].
- [18] T. Kobayashi, Y. Shimizu, K. Takagi, M. Tanimoto and T.H. Tatsuishi, *A_4 lepton flavor model and modulus stabilization from S_4 modular symmetry*, *Phys. Rev. D* **100** (2019) 115045 [*Erratum ibid.* **101** (2020) 039904] [[arXiv:1909.05139](#)] [[INSPIRE](#)].
- [19] T. Asaka, Y. Heo, T.H. Tatsuishi and T. Yoshida, *Modular A_4 invariance and leptogenesis*, *JHEP* **01** (2020) 144 [[arXiv:1909.06520](#)] [[INSPIRE](#)].
- [20] G.-J. Ding, S.F. King, X.-G. Liu and J.-N. Lu, *Modular S_4 and A_4 symmetries and their fixed points: new predictive examples of lepton mixing*, *JHEP* **12** (2019) 030 [[arXiv:1910.03460](#)] [[INSPIRE](#)].
- [21] D. Zhang, *A modular A_4 symmetry realization of two-zero textures of the Majorana neutrino mass matrix*, *Nucl. Phys. B* **952** (2020) 114935 [[arXiv:1910.07869](#)] [[INSPIRE](#)].
- [22] T. Nomura, H. Okada and S. Patra, *An inverse seesaw model with A_4 -modular symmetry*, *Nucl. Phys. B* **967** (2021) 115395 [[arXiv:1912.00379](#)] [[INSPIRE](#)].

- [23] X. Wang, *Lepton flavor mixing and CP-violation in the minimal type-(I+II) seesaw model with a modular A_4 symmetry*, *Nucl. Phys. B* **957** (2020) 115105 [[arXiv:1912.13284](#)] [[INSPIRE](#)].
- [24] T. Kobayashi, T. Nomura and T. Shimomura, *Type II seesaw models with modular A_4 symmetry*, *Phys. Rev. D* **102** (2020) 035019 [[arXiv:1912.00637](#)] [[INSPIRE](#)].
- [25] S.J.D. King and S.F. King, *Fermion mass hierarchies from modular symmetry*, *JHEP* **09** (2020) 043 [[arXiv:2002.00969](#)] [[INSPIRE](#)].
- [26] G.-J. Ding and F. Feruglio, *Testing Moduli and Flavon Dynamics with Neutrino Oscillations*, *JHEP* **06** (2020) 134 [[arXiv:2003.13448](#)] [[INSPIRE](#)].
- [27] H. Okada and M. Tanimoto, *Quark and lepton flavors with common modulus τ in A_4 modular symmetry*, [arXiv:2005.00775](#) [[INSPIRE](#)].
- [28] T. Nomura and H. Okada, *A linear seesaw model with A_4 -modular flavor and local $U(1)_{B-L}$ symmetries*, [arXiv:2007.04801](#) [[INSPIRE](#)].
- [29] T. Asaka, Y. Heo and T. Yoshida, *Lepton flavor model with modular A_4 symmetry in large volume limit*, *Phys. Lett. B* **811** (2020) 135956 [[arXiv:2009.12120](#)] [[INSPIRE](#)].
- [30] H. Okada and M. Tanimoto, *Spontaneous CP-violation by modulus τ in A_4 model of lepton flavors*, *JHEP* **03** (2021) 010 [[arXiv:2012.01688](#)] [[INSPIRE](#)].
- [31] P.T.P. Hutaauruk, D.W. Kang, J. Kim and H. Okada, *Muon $g - 2$ explanation in a modular A_4 symmetry*, [arXiv:2012.11156](#) [[INSPIRE](#)].
- [32] J.T. Penedo and S.T. Petcov, *Lepton Masses and Mixing from Modular S_4 Symmetry*, *Nucl. Phys. B* **939** (2019) 292 [[arXiv:1806.11040](#)] [[INSPIRE](#)].
- [33] P.P. Novichkov, J.T. Penedo, S.T. Petcov and A.V. Titov, *Modular S_4 models of lepton masses and mixing*, *JHEP* **04** (2019) 005 [[arXiv:1811.04933](#)] [[INSPIRE](#)].
- [34] I. de Medeiros Varzielas, S.F. King and Y.-L. Zhou, *Multiple modular symmetries as the origin of flavor*, *Phys. Rev. D* **101** (2020) 055033 [[arXiv:1906.02208](#)] [[INSPIRE](#)].
- [35] T. Kobayashi, Y. Shimizu, K. Takagi, M. Tanimoto and T.H. Tatsuishi, *New A_4 lepton flavor model from S_4 modular symmetry*, *JHEP* **02** (2020) 097 [[arXiv:1907.09141](#)] [[INSPIRE](#)].
- [36] S.F. King and Y.-L. Zhou, *Trimaximal TM_1 mixing with two modular S_4 groups*, *Phys. Rev. D* **101** (2020) 015001 [[arXiv:1908.02770](#)] [[INSPIRE](#)].
- [37] J.C. Criado, F. Feruglio and S.J.D. King, *Modular Invariant Models of Lepton Masses at Levels 4 and 5*, *JHEP* **02** (2020) 001 [[arXiv:1908.11867](#)] [[INSPIRE](#)].
- [38] X. Wang and S. Zhou, *The minimal seesaw model with a modular S_4 symmetry*, *JHEP* **05** (2020) 017 [[arXiv:1910.09473](#)] [[INSPIRE](#)].
- [39] X. Wang, *Dirac neutrino mass models with a modular S_4 symmetry*, *Nucl. Phys. B* **962** (2021) 115247 [[arXiv:2007.05913](#)] [[INSPIRE](#)].
- [40] P.P. Novichkov, J.T. Penedo, S.T. Petcov and A.V. Titov, *Modular A_5 symmetry for flavour model building*, *JHEP* **04** (2019) 174 [[arXiv:1812.02158](#)] [[INSPIRE](#)].
- [41] G.-J. Ding, S.F. King and X.-G. Liu, *Neutrino mass and mixing with A_5 modular symmetry*, *Phys. Rev. D* **100** (2019) 115005 [[arXiv:1903.12588](#)] [[INSPIRE](#)].
- [42] G.-J. Ding, S.F. King, C.-C. Li and Y.-L. Zhou, *Modular Invariant Models of Leptons at Level 7*, *JHEP* **08** (2020) 164 [[arXiv:2004.12662](#)] [[INSPIRE](#)].

- [43] X.-G. Liu and G.-J. Ding, *Neutrino Masses and Mixing from Double Covering of Finite Modular Groups*, *JHEP* **08** (2019) 134 [[arXiv:1907.01488](#)] [[INSPIRE](#)].
- [44] J.-N. Lu, X.-G. Liu and G.-J. Ding, *Modular symmetry origin of texture zeros and quark lepton unification*, *Phys. Rev. D* **101** (2020) 115020 [[arXiv:1912.07573](#)] [[INSPIRE](#)].
- [45] P.P. Novichkov, J.T. Penedo and S.T. Petcov, *Double cover of modular S_4 for flavour model building*, *Nucl. Phys. B* **963** (2021) 115301 [[arXiv:2006.03058](#)] [[INSPIRE](#)].
- [46] X.-G. Liu, C.-Y. Yao and G.-J. Ding, *Modular invariant quark and lepton models in double covering of S_4 modular group*, *Phys. Rev. D* **103** (2021) 056013 [[arXiv:2006.10722](#)] [[INSPIRE](#)].
- [47] X. Wang, B. Yu and S. Zhou, *Double covering of the modular A_5 group and lepton flavor mixing in the minimal seesaw model*, *Phys. Rev. D* **103** (2021) 076005 [[arXiv:2010.10159](#)] [[INSPIRE](#)].
- [48] C.-Y. Yao, X.-G. Liu and G.-J. Ding, *Fermion Masses and Mixing from Double Cover and Metaplectic Cover of A_5 Modular Group*, [arXiv:2011.03501](#) [[INSPIRE](#)].
- [49] X.-G. Liu, C.-Y. Yao, B.-Y. Qu and G.-J. Ding, *Half-integral weight modular forms and application to neutrino mass models*, *Phys. Rev. D* **102** (2020) 115035 [[arXiv:2007.13706](#)] [[INSPIRE](#)].
- [50] G.-J. Ding, F. Feruglio and X.-G. Liu, *Automorphic Forms and Fermion Masses*, *JHEP* **01** (2021) 037 [[arXiv:2010.07952](#)] [[INSPIRE](#)].
- [51] A. Baur, M. Kade, H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Siegel modular flavor group and CP from string theory*, *Phys. Lett. B* **816** (2021) 136176 [[arXiv:2012.09586](#)] [[INSPIRE](#)].
- [52] P.P. Novichkov, J.T. Penedo, S.T. Petcov and A.V. Titov, *Generalised CP Symmetry in Modular-Invariant Models of Flavour*, *JHEP* **07** (2019) 165 [[arXiv:1905.11970](#)] [[INSPIRE](#)].
- [53] A. Baur, H.P. Nilles, A. Trautner and P.K.S. Vaudrevange, *Unification of Flavor, CP, and Modular Symmetries*, *Phys. Lett. B* **795** (2019) 7 [[arXiv:1901.03251](#)] [[INSPIRE](#)].
- [54] B.S. Acharya, D. Bailin, A. Love, W.A. Sabra and S. Thomas, *Spontaneous breaking of CP symmetry by orbifold moduli*, *Phys. Lett. B* **357** (1995) 387 [*Erratum ibid.* **407** (1997) 451] [[hep-th/9506143](#)] [[INSPIRE](#)].
- [55] T. Dent, *CP violation and modular symmetries*, *Phys. Rev. D* **64** (2001) 056005 [[hep-ph/0105285](#)] [[INSPIRE](#)].
- [56] J. Giedt, *CP violation and moduli stabilization in heterotic models*, *Mod. Phys. Lett. A* **17** (2002) 1465 [[hep-ph/0204017](#)] [[INSPIRE](#)].
- [57] J.H. Bruinier, G.V.D. Geer, G. Harder and D. Zagier, *The 1–2–3 of Modular Forms*, in *Universitext*, Springer (2008).
- [58] F. Diamond and J.M. Shurman, *A first course in modular forms*, in *Graduate Texts in Mathematics* **228**, Springer (2005).
- [59] R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, *Finite Modular Groups and Lepton Mixing*, *Nucl. Phys. B* **858** (2012) 437 [[arXiv:1112.1340](#)] [[INSPIRE](#)].
- [60] M.-C. Chen, S. Ramos-Sánchez and M. Ratz, *A note on the predictions of models with modular flavor symmetries*, *Phys. Lett. B* **801** (2020) 135153 [[arXiv:1909.06910](#)] [[INSPIRE](#)].

- [61] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic Flavor Groups*, *JHEP* **02** (2020) 045 [[arXiv:2001.01736](#)] [[INSPIRE](#)].
- [62] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Lessons from eclectic flavor symmetries*, *Nucl. Phys. B* **957** (2020) 115098 [[arXiv:2004.05200](#)] [[INSPIRE](#)].
- [63] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic flavor scheme from ten-dimensional string theory — I. Basic results*, *Phys. Lett. B* **808** (2020) 135615 [[arXiv:2006.03059](#)] [[INSPIRE](#)].
- [64] A. Baur, M. Kade, H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *The eclectic flavor symmetry of the \mathbb{Z}_2 orbifold*, *JHEP* **02** (2021) 018 [[arXiv:2008.07534](#)] [[INSPIRE](#)].
- [65] H.P. Nilles, S. Ramos-Sánchez and P.K.S. Vaudrevange, *Eclectic flavor scheme from ten-dimensional string theory — II detailed technical analysis*, *Nucl. Phys. B* **966** (2021) 115367 [[arXiv:2010.13798](#)] [[INSPIRE](#)].
- [66] I. Esteban, M.C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni and T. Schwetz, *Global analysis of three-flavour neutrino oscillations: synergies and tensions in the determination of θ_{23} , δ_{CP} , and the mass ordering*, *JHEP* **01** (2019) 106 [[arXiv:1811.05487](#)] [[INSPIRE](#)].
- [67] F. Capozzi, E. Di Valentino, E. Lisi, A. Marrone, A. Melchiorri and A. Palazzo, *Global constraints on absolute neutrino masses and their ordering*, *Phys. Rev. D* **95** (2017) 096014 [*Addendum ibid.* **101** (2020) 116013] [[arXiv:2003.08511](#)] [[INSPIRE](#)].
- [68] I. Esteban, M.C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, *The fate of hints: updated global analysis of three-flavor neutrino oscillations*, *JHEP* **09** (2020) 178 [[arXiv:2007.14792](#)] [[INSPIRE](#)].
- [69] S. Antusch and V. Maurer, *Running quark and lepton parameters at various scales*, *JHEP* **11** (2013) 115 [[arXiv:1306.6879](#)] [[INSPIRE](#)].
- [70] <https://root.cern.ch/doc/master/classTMimuit.html>.
- [71] F. Feroz and M.P. Hobson, *Multimodal nested sampling: an efficient and robust alternative to Markov Chain Monte Carlo methods for astronomical data analysis*, *Mon. Not. Roy. Astron. Soc.* **384** (2008) 449 [[arXiv:0704.3704](#)] [[INSPIRE](#)].
- [72] F. Feroz, M.P. Hobson and M. Bridges, *MultiNest: an efficient and robust Bayesian inference tool for cosmology and particle physics*, *Mon. Not. Roy. Astron. Soc.* **398** (2009) 1601 [[arXiv:0809.3437](#)] [[INSPIRE](#)].
- [73] F. Björkeröth, F.J. de Anda, I. de Medeiros Varzielas and S.F. King, *Towards a complete $A_4 \times SU(5)$ SUSY GUT*, *JHEP* **06** (2015) 141 [[arXiv:1503.03306](#)] [[INSPIRE](#)].
- [74] F. Feruglio, V. Gherardi, A. Romanino and A.V. Titov, *Modular Invariant Dynamics and Fermion Mass Hierarchies around $\tau = i$* , [arXiv:2101.08718](#) [[INSPIRE](#)].
- [75] P.P. Novichkov, J.T. Penedo and S.T. Petcov, *Fermion Mass Hierarchies, Large Lepton Mixing and Residual Modular Symmetries*, [arXiv:2102.07488](#) [[INSPIRE](#)].