

Modules of the Highest Homological Dimension over a Gorenstein Ring

YASUO IWANAGA AND JUN-ICHI MIYACHI

Dedicated to Professor Kent R. Fuller on his 60th birthday

Abstract. We will study modules of the highest injective, projective and flat dimension over a Gorenstein ring. Let R be a Gorenstein ring of self-injective dimension n and $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \cdots \rightarrow E_n \rightarrow 0$ a minimal injective resolution. Then it is shown in [F-I] that the flat dimension and projective dimension of E_n is n , the highest dimension. In this note, we shall prove that if M is a left R -module of injective dimension n , then the last injective term $E^n(M)$ in a minimal injective resolution of M has projective and flat dimension n , and any indecomposable summand of $E^n(M)$ embeds in E_n . As a consequence, we obtain that if R is Auslander-Gorenstein, then $E^n(M)$ has essential socle.

1. Introduction

A Noether ring R is called **Gorenstein** if R has left and right finite self-injective dimensions. Further, a Noether ring R is called **Auslander-Gorenstein** if R is Gorenstein and in a minimal injective resolution $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow E_1 \rightarrow \cdots$, each E_i has flat dimension at most i . This concept was introduced by Auslander as

1991 *Mathematics Subject Classification.* 16E10, 18G10, 16G60.

The first author was supported by Grant-in-Aid for Scientific Research from the Ministry of Education, No.08640021.

The second author was supported by Grant-in-Aid for Scientific Research from the Ministry of Education, No.09640014.

a non-commutative version of the Gorenstein condition for commutative rings, studied by Bass [Ba]. In the non-commutative case, we can also see the ubiquity of Auslander-Gorenstein rings in several articles, for example, [B-G], [G-L], [Le], [L-S], [S-Z].

We showed in [Iw1] that for a Gorenstein ring of self-injective dimension n , finiteness of the injective, projective and flat dimensions of a module are all equivalent, and all of these dimensions are at most n . This fact motivated our interest in modules with the highest injective, projective or flat dimension. Let R be a Gorenstein ring of self-injective dimension n and $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \cdots \rightarrow E_n \rightarrow 0$ a minimal injective resolution. Then it is shown in [F-I] that any direct summand of E_n has the highest projective and flat dimension n . In this note, we will study the relationship between more general modules of projective (or flat) dimension n and the module E_n .

Throughout this note, $\text{id}(M)$, $\text{pd}(M)$ and $\text{fd}(M)$ stand for the injective, projective and flat dimension of a module M , respectively. Further, $0 \rightarrow M \rightarrow E^0(M) \rightarrow \cdots \rightarrow E^n(M) \rightarrow \cdots$ is a minimal injective resolution of M .

The results obtained in this note are the following.

Theorem 1. *Let R be a Gorenstein ring of self-injective dimension n and $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \cdots \rightarrow E_n \rightarrow 0$ a minimal injective resolution. If a left R -module M has injective dimension n , then any indecomposable direct summand E of $E^n(M)$ is isomorphic to a summand in E_n . As a consequence E has projective and flat dimension n .*

As a byproduct, [Mi2, Corollary 1.3] and [I-S2, Theorem 6] yield a generalization of [I-S2, Theorem 6] for Auslander-Gorenstein rings. Hoshino showed that an injective indecomposable module of flat dimension i over an Auslander-Gorenstein ring appears in i -th injective term of a minimal injective resolution of the ring ([Ho, Theorem 6.3]). Miyachi showed that any injective indecomposable module over a Gorenstein ring appears in some injective term of a minimal injective resolution of the ring ([Mi1, Corollary 4.7]).

Theorem 2. *If R is an Auslander-Gorenstein ring of self-injective dimension n , then any injective indecomposable left R -module of flat dimension n is isomorphic to a direct summand of E_n and is*

of the form $E(S)$ for a simple left module S . Thus if a left R -module M has injective dimension n , $E^n(M)$ has essential socle.

The final result generalizes [I-S1, Theorem; I-S2, Theorem 2]. It appears interesting to study the distribution of injective indecomposables along the terms of a minimal injective resolution of a Gorenstein ring.

Proposition 3. *Let R be a Noether ring and $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow \dots \rightarrow E_n \rightarrow \dots$ a minimal injective resolution of ${}_R R$.*

(1) *If M is a left R -module with $0 < i = \text{id}(M) < \infty$, then E_0 and $E^i(M)$ have no isomorphic direct summands in common.*

(2) *If R has left self-injective dimension $n \geq 1$, then E_0 and E_n have no isomorphic direct summands in common.*

2. The Proofs

Proof of Theorem 1.

By [Iw1, Theorem 2], M has projective dimension at most n . Thus let $0 \rightarrow P_n \rightarrow \dots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$ be a projective resolution of M and consider an injective resolution of each P_i ($0 \leq i \leq n$)

$$0 \rightarrow P_i \rightarrow E^0(P_i) \rightarrow E^1(P_i) \rightarrow \dots \rightarrow E^n(P_i) \rightarrow 0.$$

Then $E^j(P_i)$ for each j ($0 \leq j \leq n$) is a direct summand of a direct sum of copies of E_j . Hence by [Mi2, Corollary 1.3], M has an injective resolution of the following form

$$\begin{aligned} 0 \rightarrow M \rightarrow Q \rightarrow \bigoplus_{i=0}^{n-1} E^{i+1}(P_i) \rightarrow \bigoplus_{i=0}^{n-2} E^{i+2}(P_i) \rightarrow \dots \\ \rightarrow E^{n-1}(P_0) \oplus E^n(P_1) \rightarrow E^n(P_0) \rightarrow 0. \end{aligned}$$

Here Q is a direct summand of $\bigoplus_{i=0}^n E^i(P_i)$. Then $E^n(M)$ is a direct summand of $E^n(P_0)$, and so a direct summand of direct sum of copies of E_n . Since any indecomposable summand E of $E^n(M)$ is uniform, E embeds in E_n . \square

Proof of Theorem 2.

Let E be an injective indecomposable left module of flat dimension n . By [Mi1, Corollary 4.7], E is isomorphic to a direct summand in E_n . Since $\text{Soc}(E_n)$ is essential in E_n ([I-S 2, Theorem 6]), E is of the form $E(S)$ for some simple module S .

By Theorem 1 and [F-I, Proposition 1.1], any direct summand of $E^n(M)$ has flat dimension n and so has essential socle. That is, the socle of $E^n(M)$ is essential. \square

Proof of Proposition 3.

(1) Let U be any nonzero submodule of E_0 and $V = U \cap R \neq 0$. Then from the exact sequence

$$0 \rightarrow V \rightarrow R \rightarrow R/V \rightarrow 0,$$

we have an exact sequence

$$\text{Ext}_R^i(R, M) \longrightarrow \text{Ext}_R^i(V, M) \longrightarrow \text{Ext}_R^{i+1}(R/V, M).$$

Here $\text{Ext}_R^i(R, M) = 0$ from $i > 0$ and $\text{Ext}_R^{i+1}(R/V, M) = 0$ from $\text{id}(M) = i$. Hence we obtain $\text{Ext}_R^i(V, M) = 0$ and thus we see that V is not monomorphic to $E^i(M)$.

(2) is obvious from (1). \square

3. Examples

Let us conclude this note with a few examples. In the following examples, if R is a path algebra given by a quiver \mathcal{Q} with set \mathcal{Q}_0 of vertices and $i \in \mathcal{Q}_0$, then $S(i)$ denotes the simple R -module corresponding to the vertex i and $E(i)$ its injective hull.

(1) Theorem 1 and Proposition 3 prompt us to raise the following question: *Let R be a Gorenstein ring of self-injective dimension n and E an injective indecomposable R -module of projective dimension n . Then does there exist an R -module M of injective dimension n such that E embeds in $E^n(M)$?* It's easy to see that the question is affirmative if R is Auslander-Gorenstein. However, the answer is negative for Gorenstein rings. For example, let R be

a finite dimensional algebra over any field given by the following quiver

$$\begin{array}{ccccc}
 & & 1 & & \\
 & & \alpha \searrow & & \\
 & & & & \\
 & & & 3 \xrightarrow{\gamma} & 4 \xrightleftharpoons[\varepsilon]{\delta} & 5 \\
 & & & & & \\
 & & \beta \nearrow & & & \\
 & & 2 & & &
 \end{array}$$

with the relations $\gamma\alpha = \gamma\beta = \varepsilon\delta = \delta\varepsilon = 0$. Then R is a Gorenstein ring of self-injective dimension 2 and has infinite global dimension. $E(3)$ has projective dimension 2 but never appears in $E^2(M)$ for any R -module M of injective dimension 2. Also we can see from this observation that an injective indecomposable module with the highest projective dimension does not necessarily embed in the last term of a minimal injective resolution of a Gorenstein ring.

Moreover, $E(1)$ and $E(2)$ are both direct summands of the last injective term E_2 in a minimal injective resolution $0 \rightarrow {}_R R \rightarrow E_0 \rightarrow E_1 \rightarrow E_2 \rightarrow 0$ but $\text{Ext}_R^1(E(i), R) \neq 0$ ($i = 1, 2$). Hence $E(1)$ and $E(2)$ are not holonomic. Here a finitely generated module X over a Gorenstein ring R of self-injective dimension n is called **holonomic** if $\text{Ext}_R^i(X, R) = 0$ for all $i \neq n$.

Finally we can see in this example that all injective terms E_0 , E_1 and E_2 have the highest projective and flat dimensions.

(2) In [Iw2], it is proved that any holonomic module over an Auslander-Gorenstein ring has finite composition length and embeds in a direct sum of finitely many copies of the last injective term in a minimal injective resolution of the ring. However a submodule of finite composition length in the last injective term is not necessarily holonomic.

For example, let R be a finite dimensional algebra over any field given by the following quiver

$$\begin{array}{ccccc}
 & & & & 2 & & \\
 & & & & & & \\
 & & \alpha \nearrow & & & & \searrow \gamma \\
 & & 1 & \xleftarrow{\mu} & 5 & \xleftarrow{\lambda} & 4 \\
 & & \beta \searrow & & & & \nearrow \delta \\
 & & & & & & \\
 & & & & 3 & &
 \end{array}$$

with the relations $\mu\lambda = \alpha\mu = \beta\mu = 0$ and $\gamma\alpha = \delta\beta$. Then R is Auslander-Gorenstein of self-injective dimension 4.

Consider the left R -module M of dimension vector $(0, 1, 1, 1, 0)$, then M is a submodule of the last injective term of a minimal injective resolution of ${}_R R$ but not holonomic. For, we can see $\text{Ext}_R^1(M, R) \neq 0$, that is, the grade of M is one.

(3) We can see that if R is a Gorenstein ring of self-injective dimension n and S is a simple submodule of the last injective term E_n in a minimal injective resolution of ${}_R R$, then $\text{pd}(S) = \text{fd}(S) = n$ or ∞ . Conversely, if S is a simple R -module of the highest projective dimension n , S appears in the socle of E_n . There is an example of a Gorenstein ring R with a simple module of infinite projective and flat dimension not appearing in E_n .

Let R be a finite dimensional algebra over any field given by the following quiver

$$\begin{array}{ccc} 1 & \xleftarrow{\delta} & 3 \\ \alpha \searrow & & \nearrow \gamma \\ & 2 & \\ & \circlearrowleft \beta & \end{array}$$

with the relations $\alpha\delta = \gamma\alpha = \beta^2 = 0$. Then R is Auslander-Gorenstein of self-injective dimension 3. We can see

$$\text{pd}(S(1)) = 2, \quad \text{pd}(S(2)) = \infty, \quad \text{pd}(S(3)) = 3$$

and

$$E_0 = E(1)^{(4)}, \quad E_1 = E(2)^{(2)}, \quad E_2 = E(1), \quad E_3 = E(3).$$

Here, $M^{(t)}$ stands for a direct sum of t copies of a module M .

REFERENCES

- [A-R] M. Auslander and I. Reiten, *k-Gorenstein algebras and syzygy modules*, J. of Pure and Applied Algebra **92** (1994), 1-27.
- [Ba] H. Bass, *On the ubiquity of Gorenstein rings*, Math. Z. **82** (1963), 8-28.

- [B-G] K. A. Brown and K. R. Goodearl, *Homological aspects of Noetherian P.I. Hopf algebras, and irreducible modules of maximal dimension*, Preprint.
- [F-I] K. R. Fuller and Y. Iwanaga, *On n -Gorenstein rings and Auslander rings of low injective dimension*, Proceedings of ICRA VI, CMS Proceedings Series **14** (1993), 175–183.
- [G-L] K. R. Goodearl and T. H. Lenagan, *Catenarity in quantum algebras*, J. Pure and Applied Algebra **111** (1996), 123–142.
- [Ho] M. Hoshino, *Noetherian rings of self-injective dimension two*, Comm. Algebra **21** (1993), 1071–1094.
- [Iw1] Y. Iwanaga, *Rings with finite self-injective dimension II*, Tsukuba J. of Math. **4** (1980), 107–113.
- [Iw2] Y. Iwanaga, *Duality over Auslander-Gorenstein rings*, Math. Scand. **81** (1997), 184–190.
- [I-S1] Y. Iwanaga and H. Sato, *Minimal injective resolutions of Gorenstein rings*, Comm. in Algebra **18** (1990), 3835–3856.
- [I-S2] Y. Iwanaga and H. Sato, *On Auslander's n -Gorenstein rings*, J. of Pure and Applied Algebra **106** (1996), 61–76.
- [Le] T. Levasseur, *Some properties of non-commutative regular graded rings*, Glasgow Math. J. **34** (1992), 277–300.
- [L-S] T. Levasseur and J. T. Stafford, *The quantum coordinate ring of the special linear group*, J. Pure and Applied Algebra **86** (1993), 181–186.
- [Mi1] J. Miyachi, *Duality for derived categories and cotilting bi-modules*, J. Algebra **185** (1996), 583–603.
- [Mi2] J. Miyachi, *Injective resolutions of Noetherian rings and cogenerators*, Preprint, 1997.
- [S-Z] J. T. Stafford and J. J. Zhang, *Homological properties of (graded) Noetherian PI rings*, J. of Algebra **168** (1994), 988–1026.

Faculty of Education, Shinshu University, Nishi-Nagano 6,
Nagano 380, Japan

E-mail address: iwanaga@gipnc.shinshu-u.ac.jp

Department of Mathematics, Tokyo Gakugei University,
Koganei-shi, Tokyo 184, Japan

E-mail address: miyachi@u-gakugei.ac.jp