

MODULI SPACES OF COHERENT SYSTEMS OF SMALL SLOPE ON ALGEBRAIC CURVES

S. B. BRADLOW, O. GARCÍA-PRADA, V. MERCAT, V. MUÑOZ, AND P. E. NEWSTEAD

ABSTRACT. Let C be an algebraic curve of genus $g \geq 2$. A coherent system on C consists of a pair (E, V) , where E is an algebraic vector bundle over C of rank n and degree d and V is a subspace of dimension k of the space of sections of E . The stability of the coherent system depends on a parameter α . We study the geometry of the moduli space of coherent systems for $0 < d \leq 2n$. We show that these spaces are irreducible whenever they are non-empty and obtain necessary and sufficient conditions for non-emptiness.

1. INTRODUCTION

Let C be a smooth projective algebraic curve of genus $g \geq 2$. A *coherent system* on C of *type* (n, d, k) is a pair (E, V) , where E is a vector bundle on C of rank n and degree d and V is a linear subspace of the space of sections $H^0(E)$ of dimension k . Introduced in [7], [17] and [11], there is a notion of stability for coherent systems which permits the construction of moduli spaces. This notion depends on a real parameter, and thus leads to a family of moduli spaces. As described in [2], there is a useful relation between these moduli spaces and the Brill-Noether loci in the moduli spaces of semistable bundles of rank n and degree d .

In [5] we began a systematic study of the coherent systems moduli spaces, partly with a view to applications in higher rank Brill-Noether theory. This study has been continued in [4], where we have obtained substantial new information about the geometry and topology of the moduli spaces for $k \leq n$.

In the present paper, we go in a slightly different direction and consider coherent systems with $d \leq 2n$. Our results are essentially a generalisation and extension of those of [3, 12, 14]. More precisely, we show that the moduli spaces of coherent systems are irreducible whenever they are non-empty and obtain necessary and sufficient conditions

Date: 6 July 2007.

2000 Mathematics Subject Classification. 14H60, 14D20, 14H51.

Key words and phrases. Algebraic curves, moduli of vector bundles, coherent systems, Brill-Noether loci.

All authors are members of the research group VBAC (Vector Bundles on Algebraic Curves). Support was received from a grant from the European Scientific Exchange Programme of the Royal Society of London and the Consejo Superior de Investigaciones Científicas (15646) and a further grant from the Royal Society of London for an International Joint Project (2005/R3). The first author was partially supported by the National Science Foundation under grant DMS-0072073. The first, second and fourth authors were supported through MEC grant MTM2004-07090-C03-01 (Spain) and the fourth also through NSF grant DMS-0111298 (US). The fifth author was supported by the Academia Mexicana de Ciencias during a visit to CIMAT, Guanajuato, Mexico.

for non-emptiness. Even for Brill-Noether loci, the irreducibility result is stronger than those previously known. The condition for non-emptiness is identical with that for Brill-Noether loci except when C is hyperelliptic, $d = 2n$ and $k > n$. The methods are generally similar to those of [3, 12, 14], except that we make essential use of extensions of coherent systems (rather than simply extensions of bundles). In particular, at crucial points in the proof of irreducibility and, in the hyperelliptic case, that of non-emptiness, we use the methods of [5] to estimate dimensions of spaces of extensions. Some of the estimates are quite delicate and require careful use of these methods. We also make use of some of the results of [4] to handle the case $k \leq n$, although the present paper can be read independently of [4].

In order to give full statements of our main results, we need to outline some definitions and notations (for more details, see section 2). We denote by $G(\alpha; n, d, k)$ the moduli space of α -stable coherent systems of type (n, d, k) , where $\alpha \in \mathbb{R}$ is a parameter (with the necessary conditions $d > 0$, $\alpha > 0$ and $\alpha(n - k) < d$ for non-emptiness of $G(\alpha; n, d, k)$), and by $B(n, d, k)$ the *Brill-Noether locus* of stable bundles of rank n and degree d with $h^0(E) \geq k$. We write $\beta(n, d, k)$ for the “expected dimension” of $G(\alpha; n, d, k)$, namely

$$\beta(n, d, k) = n^2(g - 1) + 1 - k(k - d + n(g - 1)).$$

Note that the data for defining a coherent system (E, V) can also be expressed as an exact sequence

$$0 \longrightarrow D^* \longrightarrow V \otimes \mathcal{O} \longrightarrow E \longrightarrow F \oplus T \longrightarrow 0,$$

where D and F are vector bundles, T is a torsion sheaf and $h^0(D^*) = 0$. A coherent system (E, V) is said to be *generated* if F and T are both zero. Finally, we define $U(n, d, k)$ and $U^s(n, d, k)$ by

$$U(n, d, k) := \{(E, V) : E \text{ is stable and } (E, V) \text{ is } \alpha\text{-stable for } \alpha > 0, \alpha(n - k) < d\}$$

and

$$U^s(n, d, k) := \{(E, V) : (E, V) \text{ is } \alpha\text{-stable for } \alpha > 0, \alpha(n - k) < d\}$$

(see section 5 for further details). Clearly $U(n, d, k) \subset U^s(n, d, k)$.

We can now state our main results.

Theorem 4.4. *Suppose that $0 < d \leq 2n$ and $\alpha > 0$. If $G(\alpha; n, d, k) \neq \emptyset$, then it is irreducible. Moreover*

(a) *if $k < n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$0 \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow F \rightarrow 0,$$

where F is a vector bundle with $h^0(F^) = 0$;*

(b) *if $k = n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$0 \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow T \rightarrow 0,$$

where T is a torsion sheaf;

(c) *if $k > n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow 0,$$

i.e. (E, V) is generated;

- (d) $\dim G(\alpha; n, d, k) = \beta(n, d, k)$ except when C is hyperelliptic and $(n, d, k) = (n, 2n, n+1)$ with $n < g-1$.

Theorem 5.4. *Suppose that C is non-hyperelliptic of genus $g \geq 3$, $n \geq 2$ and $0 < d \leq 2n$. Then $U(n, d, k) \neq \emptyset$ if and only if either*

$$k \leq n + \frac{1}{g}(d - n), \quad (n, d, k) \neq (n, n, n)$$

or

$$(n, d, k) = (g - 1, 2g - 2, g).$$

In all other cases,

- $G(\alpha; n, d, k) = \emptyset$ for all $\alpha > 0$;
- $B(n, d, k) = \emptyset$.

Theorem 5.5. *Suppose that C is hyperelliptic, $n \geq 2$ and $0 < d \leq 2n$. Then*

- (a) $U(n, d, k) \neq \emptyset$ if and only if either

$$0 < d < 2n, \quad k \leq n + \frac{1}{g}(d - n), \quad (n, d, k) \neq (n, n, n)$$

or $d = 2n, k \leq n$;

- (b) if $k > n$, then

- $U(n, 2n, k) = \emptyset$;
- $U^s(n, 2n, k) \neq \emptyset$ if and only if either $k \leq n + \frac{n}{g}$ or $k = n + 1$ and $2 \leq n \leq g - 1$.

In all other cases,

- $G(\alpha; n, d, k) = \emptyset$ for all $\alpha > 0$;
- $B(n, d, k) = \emptyset$.

The case $n = 1$ is omitted from the last two statements since the results then need modifying; of course this case is very simple.

The contents of the paper are as follows. In section 2, we give definitions and notations together with some basic facts which we shall need. In section 3, we generalise the results of [3, 12, 14] to obtain a necessary condition for the existence of α -stable coherent systems. Section 4 is devoted to a proof of irreducibility (Theorem 4.4). In section 5, we state our results on non-emptiness separately for C non-hyperelliptic (Theorem 5.4) and for C hyperelliptic (Theorem 5.5); the proofs for C non-hyperelliptic are included. In the lengthy section 6 we prove Theorem 5.5; this requires some delicate constructions using the methods of [5]. Finally section 7 contains an example with $d > 2n$ to show that the situation can then be more complicated.

We suppose throughout that C is a smooth projective algebraic curve of genus $g \geq 2$ defined over the complex numbers. The cases $g = 0$ and $g = 1$ have been investigated in [8, 9, 10], where irreducibility has been proved with no restriction on the degree, but the non-emptiness results for the case $g = 0$ are still not complete. We also assume that $k \geq 1$.

2. DEFINITIONS, NOTATIONS AND BASIC FACTS

We refer the reader to [5] for the basic properties of coherent systems on algebraic curves. For convenience, we provide here a synopsis of the main definitions and facts which we shall need. Recall that the *slope* $\mu(E)$ of a vector bundle of rank n and degree d is defined by $\mu(E) := \frac{d}{n}$.

Definition 2.1. Let (E, V) be a coherent system of type (n, d, k) . For any $\alpha \in \mathbb{R}$, the α -slope $\mu_\alpha(E, V)$ is defined by

$$\mu_\alpha(E, V) := \frac{d}{n} + \alpha \frac{k}{n}.$$

A *coherent subsystem* of (E, V) is a coherent system (E', V') such that E' is a subbundle of E and $V' \subset V \cap H^0(E')$. A *quotient coherent system* of (E, V) is a coherent system (E'', V'') together with a homomorphism $(E, V) \rightarrow (E'', V'')$ such that both $E \rightarrow E''$ and $V \rightarrow V''$ are surjective.

Note that, with our definition of coherent system, a subsystem possesses a corresponding quotient system only if $V' = V \cap H^0(E')$.

Definition 2.2. A coherent system (E, V) is α -stable (α -semistable) if, for every proper coherent subsystem (E', V') ,

$$\mu_\alpha(E', V') < (\leq) \mu_\alpha(E, V).$$

There exists a moduli space $G(\alpha; n, d, k)$ of α -stable coherent systems of type (n, d, k) ; necessary conditions for non-emptiness are

$$d > 0, \quad \alpha > 0, \quad (n - k)d < \alpha.$$

Definition 2.3. A *critical value* for coherent systems of type (n, d, k) is a value of $\alpha > 0$ for which there exists a coherent system (E, V) of type (n, d, k) and a coherent subsystem (E', V') of (E, V) of type (n', d', k') such that $\frac{k'}{n'} \neq \frac{k}{n}$ but $\mu_\alpha(E', V') = \mu_\alpha(E, V)$. We also regard $\alpha = 0$ as a critical value.

It is known [5, Propositions 4.2, 4.6] that, for any (n, d, k) , there are finitely many critical values

$$0 = \alpha_0 < \alpha_1 < \dots < \alpha_L < \begin{cases} \frac{d}{n-k} & \text{if } k < n \\ \infty & \text{if } k \geq n. \end{cases}$$

Moreover, if $k < n$ and $\alpha \geq \frac{d}{n-k}$, then $G(\alpha; n, d, k) = \emptyset$. For $\alpha, \alpha' \in (\alpha_i, \alpha_{i+1})$, we have $G(\alpha; n, d, k) = G(\alpha'; n, d, k)$ and we denote this moduli space by $G_i := G_i(n, d, k)$. We shall be particularly concerned with the moduli spaces G_0 (“small” α) and G_L (“large” α). If $(E, V) \in G_0$, we say also that (E, V) is 0^+ -stable (with similar definitions for α^\pm -stable).

We denote by $M(n, d)$ the moduli space of stable bundles of rank n and degree d , and by $B(n, d, k)$ the *Brill-Noether locus*

$$B(n, d, k) := \{E \in M(n, d) : h^0(E) \geq k\}.$$

We have, for any coherent system (E, V) , [5, Proposition 2.5]

- $(E, V) \in G_0 \implies E$ semistable;

- E stable $\implies (E, V) \in G_0$.

The moduli space $G(\alpha; n, d, k)$ has the property that every irreducible component has dimension greater than or equal to the *Brill-Noether number*

$$(1) \quad \beta(n, d, k) := n^2(g-1) + 1 - k(k-d+n(g-1)).$$

This number is the “expected dimension” of $G(\alpha; n, d, k)$ in a stronger sense. For this, we define, for any coherent system (E, V) , the *Petri map* of (E, V) as the map

$$V \otimes H^0(E^* \otimes K) \longrightarrow H^0(E \otimes E^* \otimes K)$$

given by multiplication of sections. This map governs the infinitesimal behaviour of the moduli space in the following sense.

- Let (E, V) be an α -stable coherent system of type (n, d, k) . Then $G(\alpha; n, d, k)$ is smooth of dimension $\beta(n, d, k)$ at the point corresponding to (E, V) if and only if the Petri map of (E, V) is injective.

Definition 2.4. The coherent system (E, V) is *generated* if the evaluation map $V \otimes \mathcal{O} \rightarrow E$ is surjective. The bundle E is *generated* if $(E, H^0(E))$ is generated.

We shall make no explicit use of the flip loci G_i^\pm of [5, section 6], so shall not describe them here. However we make extensive use of extensions

$$(2) \quad 0 \longrightarrow (E_1, V_1) \longrightarrow (E, V) \longrightarrow (E_2, V_2) \longrightarrow 0,$$

where (E_1, V_1) , (E_2, V_2) are coherent systems of types (n_1, d_1, k_1) , (n_2, d_2, k_2) respectively. Here we use the notations and results of [5, section 3]. The extensions (2) are classified in the usual way by a group

$$\text{Ext}^1((E_2, V_2), (E_1, V_1)).$$

For dimensional reasons $\text{Ext}^q((E_2, V_2), (E_1, V_1)) = 0$ for $q \geq 3$, so we have [5, equation (8)]

$$(3) \quad \dim \text{Ext}^1((E_2, V_2), (E_1, V_1)) = C_{21} + \dim \mathbb{H}_{21}^0 + \dim \mathbb{H}_{21}^2,$$

where

$$(4) \quad C_{21} := n_1(n_2 - k_2)(g-1) + (k_2 - n_2)d_1 + d_2n_1 - k_1k_2$$

and

$$\mathbb{H}_{21}^0 = \text{Hom}((E_2, V_2), (E_1, V_1)), \quad \mathbb{H}_{21}^2 = \text{Ext}^2((E_2, V_2), (E_1, V_1)).$$

The main purpose of introducing the number C_{21} is that frequently, although not always, \mathbb{H}_{21}^0 and \mathbb{H}_{21}^2 are both zero and $\dim \text{Ext}^1((E_2, V_2), (E_1, V_1))$ is then given by the purely numerical formula (4). Of course, we can define \mathbb{H}_{12}^0 , \mathbb{H}_{12}^2 and C_{12} by interchanging the indices, and in particular

$$(5) \quad C_{12} = n_2(n_1 - k_1)(g-1) + (k_1 - n_1)d_2 + d_1n_2 - k_1k_2.$$

Note [5, Corollary 3.7] that, with the notation of (2),

$$(6) \quad \beta(n, d, k) = \beta(n_1, d_1, k_1) + \beta(n_2, d_2, k_2) + C_{12} + C_{21} - 1.$$

Note further [5, equation (11)] that, if N_2 is the kernel of the evaluation map $V_2 \otimes \mathcal{O} \rightarrow E_2$, then

$$(7) \quad \mathbb{H}_{21}^2 = H^0(E_1^* \otimes N_2 \otimes K)^*.$$

Putting $(E_1, V_1) = (E_2, V_2)$ in (3) and using (1) and (4), we get

$$\dim \text{Ext}^1((E, V), (E, V)) = \beta(n, d, k) + \dim \text{End}(E, V) + \dim \text{Ext}^2((E, V), (E, V)) - 1.$$

Now, when $\text{Ext}^2((E, V), (E, V)) = 0$, there is no obstruction to the construction of a local deformation space for (E, V) and this local deformation space has dimension

$$(8) \quad \dim \text{Ext}^1((E, V), (E, V)) = \beta(n, d, k) + \dim \text{End}(E, V) - 1$$

(see [6, Théorème 3.12] and compare [5, Proposition 3.4]).

We need one further important fact about the extensions (2).

Proposition 2.5. *If (2) is non-trivial and α_i is a critical value such that (E_1, V_1) and (E_2, V_2) are both α_i -stable with $\mu_{\alpha_i}(E_1, V_1) = \mu_{\alpha_i}(E_2, V_2)$ and $\mu_{\alpha_i^-}(E_1, V_1) < \mu_{\alpha_i^-}(E_2, V_2)$, then (E, V) is α_i^- -stable.*

Since this is not explicitly stated in either [5] or [4] (although it is used in [4]), we give a proof.

Proof. Suppose that (E', V') is a coherent subsystem of (E, V) contradicting α_i^- -stability. Then (E', V') also contradicts α_i -stability of (E, V) . It follows that either $(E', V') = (E_1, V_1)$ or (E', V') maps isomorphically to (E_2, V_2) . In the first case, α_i -stability of (E, V) is not contradicted, while in the second (2) is trivial. \square

3. COHERENT SYSTEMS FOR $d \leq 2n$

Our first object in this section is to obtain a necessary condition for the existence of α -stable coherent systems for $d \leq 2n$; it turns out that the condition is almost identical with that for stable bundles (see [12, 14]).

We start with the case $d < 2n$, when the results of [12] carry over quite easily to give a necessary condition for α -semistability.

Lemma 3.1. *Suppose that (E, V) is an α -semistable coherent system for some $\alpha > 0$ and that $0 < d < 2n$. Then*

$$(9) \quad k \leq n + \frac{1}{g}(d - n).$$

Proof. If E is semistable, the result holds by [12, Chapitre 2, Théorème A.1].

If E is not semistable, then E has a stable quotient G with $\mu(G) < \mu(E) < 2$. Again by [12, Chapitre 2, Théorème A.1], we have

$$h^0(G) \leq n_G + \frac{1}{g}(d_G - n_G),$$

where n_G and d_G denote the rank and degree of G . Let W denote the image of V in $H^0(G)$. Then, if $k > n + \frac{1}{g}(d - n)$, we have

$$\frac{\dim W}{n_G} \leq 1 + \frac{1}{g} \left(\frac{d_G}{n_G} - 1 \right) < 1 + \frac{1}{g} \left(\frac{d}{n} - 1 \right) < \frac{k}{n}.$$

It follows that the quotient coherent system (G, W) contradicts the α -semistability of (E, V) for any $\alpha > 0$. \square

This lemma has the following interesting consequence, which has relevance for coherent systems with $k > n$ in general.

Corollary 3.2. *Suppose that $k > n$ and that there exists an α -semistable coherent system (E, V) for some $\alpha > 0$. Then*

$$d \geq \min \{2n, n + g(k - n)\}.$$

Proof. If $k > n$ and $0 < d < 2n$, then the lemma implies that $d - n \geq g(k - n)$. For $d = 0$, (E, V) cannot be α -stable. The associated graded object must be a sum of coherent systems of types $(1, 0, 1)$ or $(1, 0, 0)$ and hence $k \leq n$. \square

In order to cover the case $d = 2n$, we shall make use of the dual span construction, which we briefly recall. Let (F, W) be a coherent system. Slightly modifying the notations of [5, section 5.4], we define a coherent system

$$D(F, W) = (D_W(F), W'),$$

where $D_W(F)$ is defined by the exact sequence

$$0 \longrightarrow D_W(F)^* \longrightarrow W \otimes \mathcal{O} \longrightarrow F$$

and W' is the image of W^* in $H^0(D_W(F))$. In particular we write $D(F)$ for $D_{H^0(F)}(F)$. If $W \otimes \mathcal{O} \rightarrow F$ is surjective, then the linear map $W^* \rightarrow W'$ is induced from the dual exact sequence

$$(10) \quad 0 \longrightarrow F^* \longrightarrow W^* \otimes \mathcal{O} \longrightarrow D_W(F) \longrightarrow 0.$$

Moreover, if $h^0(F^*) = 0$, then W^* maps isomorphically to W' .

For the canonical line bundle K , we obtain a bundle $D(K)$ of rank $g - 1$ and degree $2g - 2$. Taking $W = H^0(K)$, (10) becomes

$$(11) \quad 0 \longrightarrow K^* \longrightarrow H^0(K)^* \otimes \mathcal{O} \longrightarrow D(K) \longrightarrow 0.$$

It is known [16] that, if C is not hyperelliptic, then $D(K)$ is stable and $h^0(D(K)) = g$, while, if C is hyperelliptic, then $D(K) \cong L^{\oplus(g-1)}$, where L is the hyperelliptic line bundle. In both cases, we obtain new α -stable coherent systems with $d = 2n$; to describe them, we use the following general lemma.

Lemma 3.3. *Let (E, V) be a generated coherent system of type $(n, d, n + 1)$ such that E is a semistable bundle. Then (E, V) is α -stable for all $\alpha > 0$.*

Proof. Let (F, W) be a coherent subsystem of (E, V) with $0 < \text{rk } F < n$. Since E is semistable, $\mu(F) \leq \mu(E)$. To show that $\mu_\alpha(F, W) < \mu_\alpha(E, V)$, it is therefore sufficient to show $\dim W \leq \text{rk } F$. Suppose $\dim W > \text{rk } F$. Then the image of V in $H^0(E/F)$ has dimension $\leq \text{rk}(E/F)$. Now $d > 0$, hence $\deg(E/F) > 0$. It follows that the image of V does not generate E/F , which contradicts the hypothesis that V generates E . \square

Corollary 3.4. *If C is not hyperelliptic, then $D(K, H^0(K))$ is α -stable of type $(g - 1, 2g - 2, g)$ for all $\alpha > 0$.*

Proof. This follows immediately from the lemma and (11). \square

Corollary 3.5. *Suppose C is hyperelliptic and a is an integer, $1 \leq a \leq g - 1$. Let L be the hyperelliptic line bundle and W a subspace of $H^0(L^{\oplus a})$ of dimension $a + 1$ which generates $L^{\oplus a}$. Then*

- *the coherent system $(L^{\oplus a}, W)$ is α -stable of type $(a, 2a, a + 1)$ for all $\alpha > 0$;*
- *$(L^{\oplus a}, W) \cong D(L^a, H^0(L^a))$.*

Proof. The first statement follows at once from the lemma. For the second statement, note that we have an exact sequence

$$0 \longrightarrow M \longrightarrow W \otimes \mathcal{O} \longrightarrow L^{\oplus a} \longrightarrow 0,$$

where M is a line bundle. But then $M^* \cong \det L^{\oplus a} \cong L^a$. We therefore have a dual exact sequence

$$0 \longrightarrow (L^{\oplus a})^* \longrightarrow W^* \otimes \mathcal{O} \longrightarrow L^a \longrightarrow 0.$$

Since $h^0(L^a) = a + 1$, this must be the defining exact sequence for $D(L^a, H^0(L^a))$, which completes the proof. \square

Remark 3.6. In Corollary 3.5, for any subspace W of dimension $a + 1$ which generates $L^{\oplus a}$, the isomorphism class of $(L^{\oplus a}, W)$ is the same.

Lemma 3.7. *Let (E, V) be a coherent system and F a vector bundle. Suppose that F is generated and that $h^0(F^*) = 0$. Then*

$$\mathrm{Hom}(D(F, H^0(F)), (E, V))$$

is isomorphic to the kernel of the homomorphism

$$\Psi : H^0(F) \otimes V \rightarrow H^0(F \otimes E)$$

given by multiplication of sections.

Proof. We have an exact sequence of coherent systems

$$(12) \quad 0 \longrightarrow (F^*, 0) \longrightarrow (H^0(F)^* \otimes \mathcal{O}, H^0(F)^*) \longrightarrow D(F, H^0(F)) \longrightarrow 0.$$

Taking $\mathrm{Hom}((12), (E, V))$, we get an exact sequence

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}(D(F, H^0(F)), (E, V)) &\longrightarrow \mathrm{Hom}((H^0(F)^* \otimes \mathcal{O}, H^0(F)^*), (E, V)) \\ &\xrightarrow{\psi} \mathrm{Hom}((F^*, 0), (E, V)) \longrightarrow \dots \end{aligned}$$

Now ψ can be identified with the natural linear map

$$\mathrm{Hom}((H^0(F)^*, V) \longrightarrow \mathrm{Hom}(F^*, E)$$

and this in turn can be identified with Ψ . \square

Corollary 3.8. *Let (E, V) be a coherent system of type (n, d, k) with $h^0(E^*) = 0$ and let*

$$m = \dim \mathrm{Hom}(D(K, H^0(K)), (E, V)).$$

Then

$$k \leq n + \frac{1}{g}(d - n + m).$$

Proof. Apply Lemma 3.7 with $F = K$. The condition $h^0(E^*) = 0$ implies by Serre duality that $h^0(K \otimes E) = d + n(g - 1)$. \square

Remark 3.9. If E is a semistable bundle with $0 < d < 2n$, then $\text{Hom}(D(K), E) = 0$ since $D(K)$ is semistable of slope 2. So $m = 0$ and the corollary reduces to [12, Chapitre 2, Théorème A.1].

We come now to the main result of this section. Although we have already proved it in the case $d < 2n$, for completeness we state it for the whole range $d \leq 2n$.

Proposition 3.10. *Let (E, V) be an α -stable coherent system of type (n, d, k) with $0 < d \leq 2n$. Then*

$$k \leq n + \frac{1}{g}(d - n)$$

except when $d = 2n$ and

- C is not hyperelliptic and $(E, V) \cong D(K, H^0(K))$;
- C is hyperelliptic and $(E, V) \cong (L^{\oplus a}, W)$, where L is the hyperelliptic line bundle, $a \leq g - 1$ and W is a subspace of $H^0(L^{\oplus a})$ of dimension $a + 1$ which generates $L^{\oplus a}$.

Proof. For $d < 2n$, this follows at once from Lemma 3.1. So we can suppose $d = 2n$. If E is stable, the proposition follows from the results of [14]. If E is not semistable, the proof of Lemma 3.1 still works.

It remains to consider the case where E is strictly semistable with $d = 2n$. We can certainly suppose that

$$(13) \quad k > n + \frac{1}{g}(d - n) = n \left(1 + \frac{1}{g}\right).$$

By Corollary 3.8 this implies that there exists a non-zero homomorphism

$$(14) \quad D(K, H^0(K)) \longrightarrow (E, V).$$

Suppose first that C is not hyperelliptic. Then $D(K)$ is stable; since E is strictly semistable, the homomorphism (14) must be injective and indeed

$$\dim \text{Hom}(D(K, H^0(K)), (E, V)) \leq \dim \text{Hom}(D(K), E) \leq \frac{n}{g - 1}.$$

Corollary 3.8 implies that

$$k \leq n \left(1 + \frac{1}{g}\right) + \frac{n}{g(g - 1)} = \frac{ng}{g - 1}.$$

But now

$$\mu_\alpha(D(K, H^0(K))) = 2 + \alpha \frac{g}{g - 1} \geq 2 + \alpha \frac{k}{n},$$

which contradicts the α -stability of (E, V) unless $(E, V) \cong D(K, H^0(K))$.

If C is hyperelliptic, we have $D(K) \cong L^{\oplus(g-1)}$ and $h^0(L^* \otimes E) \leq n$, so Corollary 3.8 gives

$$k \leq n \left(1 + \frac{1}{g}\right) + \frac{1}{g}n(g-1) = 2n.$$

By (13), we deduce that there exists an integer a , $1 \leq a \leq g-1$ such that

$$n \left(1 + \frac{1}{a}\right) \geq k > n \left(1 + \frac{1}{a+1}\right).$$

By Clifford's Theorem (see [3, Theorem 2.1]), we have $h^0(E \otimes L^a) \leq (a+2)n$; so

$$k \cdot h^0(L^a) = k(a+1) > n(a+2) \geq h^0(E \otimes L^a).$$

Hence, by Lemma 3.7, there exists a non-zero homomorphism of coherent systems

$$D(L^a, H^0(L^a)) \longrightarrow (E, V).$$

Now

$$\mu_\alpha(D(L^a, H^0(L^a))) = 2 + \alpha \frac{a+1}{a} \geq 2 + \alpha \frac{k}{n}.$$

By Corollary 3.5, this contradicts the α -stability of (E, V) unless

$$(E, V) \cong D(L^a, H^0(L^a)) \cong (L^{\oplus a}, W),$$

where W is any subspace of $H^0(L^{\oplus a})$ of dimension $a+1$ which generates $L^{\oplus a}$. \square

4. IRREDUCIBILITY OF THE MODULI SPACE FOR $d \leq 2n$

In this section we prove that the moduli space $G(\alpha; n, d, k)$ is irreducible for $0 < d \leq 2n$. We start with two lemmas.

Lemma 4.1. *Suppose that (E, V) is a coherent system of type (n, d, k) and consider the exact sequence*

$$(15) \quad 0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow F \oplus T \rightarrow 0,$$

where $D = D_V(E)$, T is a torsion sheaf and F is a vector bundle. Suppose further that $\text{Hom}(D(K, H^0(K)), (E, V)) = 0$. Then

- (a) $h^1(D) = 0$;
- (b) if $F = 0$, the Petri map at (E, V) is injective.

Proof. (a) Suppose that $h^1(D) \neq 0$. Then there is a non-zero morphism $D \rightarrow K$. Since V^* generates D , the map $V^* \rightarrow H^0(K)$ is non-zero. Dualising we obtain a diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & K^* & \rightarrow & H^0(K)^* \otimes \mathcal{O} & \rightarrow & D(K) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & D^* & \rightarrow & V \otimes \mathcal{O} & \rightarrow & E \end{array}$$

and hence a non-zero morphism $D(K, H^0(K)) \rightarrow (E, V)$, contradicting the hypothesis.

(b) Tensor the sequence $V \otimes \mathcal{O} \rightarrow E \rightarrow T \rightarrow 0$ with D and apply cohomology to get an exact sequence

$$H^1(D \otimes V) \rightarrow H^1(D \otimes E) \rightarrow 0.$$

Proof. By Lemmas 4.1(a) and 4.2, $h^1(D) = 0$. Hence, from the cohomology sequence associated to the top row of (17), $h^1(D \otimes E_1) = 0$; thus, by (7) and Serre duality,

$$\mathrm{Ext}^2((E_1, V_1), (E_1, V_1)) = H^0(E_1^* \otimes D^* \otimes K)^* = 0.$$

It follows by (8) that the local deformation space of (E_1, V_1) has dimension

$$x_1 := \beta(n_1, d_1, k_1) + \dim \mathrm{End}(E_1, V_1) - 1.$$

On the other hand, the Petri map of (E_2, V_2) is clearly injective, so (again by (8)), the local deformation space of (E_2, V_2) has dimension

$$x_2 := \beta(n_2, d_2, k_2) + \dim \mathrm{End}(E_2, V_2) - 1.$$

We need to consider only those extensions (18) for which (E, V) is α -stable for some α . For fixed (E_1, V_1) , (E_2, V_2) , the group

$$\mathrm{Aut}(E_1, V_1) \times \mathrm{Aut}(E_2, V_2) / \{(\lambda, \lambda^{-1}) : \lambda \in \mathbb{C}^*\}$$

acts freely on these extensions. Hence, in (18), (E, V) depends on at most

$$\begin{aligned} & x_1 + x_2 + \dim \mathrm{Ext}^1((E_2, V_2), (E_1, V_1)) - (\dim \mathrm{Aut}(E_1, V_1) + \dim \mathrm{Aut}(E_2, V_2) - 1) \\ (19) \quad & = \beta(n_1, d_1, k_1) + \beta(n_2, d_2, k_2) + \dim \mathrm{Ext}^1((E_2, V_2), (E_1, V_1)) - 1 \end{aligned}$$

parameters. Now, by (3), we have

$$\dim \mathrm{Ext}^1((E_2, V_2), (E_1, V_1)) = C_{21} + \dim \mathbb{H}_{21}^0 + \dim \mathbb{H}_{21}^2.$$

Here $\mathbb{H}_{21}^0 = \mathrm{Hom}((E_2, V_2), (E_1, V_1)) = 0$, since the existence of a non-zero homomorphism would imply that (E, V) is not simple, in contradiction to [5, Proposition 2.2(ii)]. Moreover, by (7),

$$\mathbb{H}_{21}^2 = H^0(E_1^* \otimes N_2 \otimes K)^*,$$

where N_2 is the kernel of the evaluation map $V_2 \otimes \mathcal{O} \rightarrow E_2$, which is clearly 0. So (19) becomes

$$(20) \quad \beta(n_1, d_1, k_1) + \beta(n_2, d_2, k_2) + C_{21} - 1.$$

So, to prove that the number given by (20) is less than $\beta(n, d, k)$, it is enough by (6) to prove that $C_{12} \geq 1$. Now, by (5),

$$\begin{aligned} (21) \quad C_{12} &= n_2(n_1 - k_1)(g - 1) + (k_1 - n_1)d_2 + d_1n_2 - k_1k_2 \\ &= (d_1 - n_1 + (n_1 - k_1)g)n_2 + k_1(n_2 - k_2) + d_2(k_1 - n_1). \end{aligned}$$

We can now check that the third term in (21) is positive and the other two are non-negative.

- Since $h^0(G^*) = 0$ and $G \neq 0$, we have $k_1 > n_1$ and $d_1 > 0$; also $k_2 \leq n_2$, hence $\frac{k_1}{n_1} > \frac{k_2}{n_2}$. Now α -stability of (E, V) implies that $\frac{d_1}{n_1} < \frac{d_2}{n_2}$, hence $d_2 > \frac{d_1n_2}{n_1} > 0$. So $d_2(k_1 - n_1) > 0$.
- Since $k_2 \leq n_2$, $k_1(n_2 - k_2) \geq 0$.
- Since Lemma 4.2 applies to (E, V) , it follows from (18) that

$$\mathrm{Hom}(D(K, H^0(K)), (E_1, V_1)) = 0.$$

Since also $h^0(E_1^*) = 0$, it follows from Corollary 3.8 that $k_1 \leq n_1 + \frac{1}{g}(d_1 - n_1)$, so $d_1 - n_1 + (n_1 - k_1)g \geq 0$.

This completes the proof of the lemma. \square

Theorem 4.4. *Suppose that $0 < d \leq 2n$ and $\alpha > 0$. If $G(\alpha; n, d, k) \neq \emptyset$, then it is irreducible. Moreover*

(a) *if $k < n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$(22) \quad 0 \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow F \rightarrow 0,$$

where F is a vector bundle with $h^0(F^) = 0$;*

(b) *if $k = n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$(23) \quad 0 \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow T \rightarrow 0,$$

where T is a torsion sheaf;

(c) *if $k > n$, the generic element of $G(\alpha; n, d, k)$ has the form*

$$(24) \quad 0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow 0,$$

i.e. (E, V) is generated;

(d) $\dim G(\alpha; n, d, k) = \beta(n, d, k)$ *except when C is hyperelliptic and $(n, d, k) = (n, 2n, n + 1)$ with $n < g - 1$.*

Proof. Let Z be a component of $G(\alpha; n, d, k)$. Note that $\dim Z \geq \beta(n, d, k)$.

(a) Suppose $k < n$. By Lemma 4.3, the generic element of Z must have $(E_1, V_1) = 0$ in (18), so we have

$$0 \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow F \oplus T \rightarrow 0,$$

i.e. (E, V) is injective in the sense of [4, Definition 2.1]. The result now follows from [4, Theorem 3.3(iii)].

(b) Suppose $k = n$. By Lemma 4.3, the generic element of Z has the form (23). Now the proof of [5, Theorem 5.6] applies to show that $G(\alpha; n, d, k)$ is irreducible.

(c) If $k > n + \frac{1}{g}(d - n)$, then, by Proposition 3.10, $G(\alpha; n, d, k)$ consists of a single point and (24) holds. So suppose $n < k \leq n + \frac{1}{g}(d - n)$. Then, by Lemma 4.3, the generic element of Z has the form

$$(25) \quad 0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow T \rightarrow 0.$$

Moreover (25) splits into two sequences

$$(26) \quad 0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E' \rightarrow 0$$

and

$$(27) \quad 0 \rightarrow E' \rightarrow E \rightarrow T \rightarrow 0,$$

where E' is a vector bundle and T is a torsion sheaf.

Let

$$Z' := \{(E, V) \in G(\alpha; n, d, k) : (E, V) \text{ is generated}\}.$$

We shall prove that Z' is irreducible and that $G(\alpha; n, d, k) \setminus Z'$ is of dimension $< \beta(n, d, k)$. Since every component of $G(\alpha; n, d, k)$ has dimension $\geq \beta(n, d, k)$, this will complete the proof.

If $(E, V) \in Z'$, then $E' = E$ in (26). Dualising this sequence, we get

$$(28) \quad 0 \rightarrow E^* \rightarrow V^* \otimes \mathcal{O} \rightarrow D \rightarrow 0,$$

where $h^0(D^*) = 0$ from (26) and $h^1(D) = 0$ by Lemma 4.1(a). Moreover $h^0(E^*) = 0$, since otherwise there would exist a non-zero homomorphism $E \rightarrow \mathcal{O}$; since (E, V) is generated, this implies that (E, V) has a direct summand $(\mathcal{O}, H^0(\mathcal{O}))$, contradicting α -stability. The bundles D of rank $k - n$ and degree d for which $h^1(D) = h^0(D^*) = 0$ form a bounded set of bundles and are therefore parametrised (not necessarily injectively) by a variety X which is irreducible (or empty) (this follows from [1, Theorem 2], which is essentially due to Serre, see also [15, Proposition 2.6]). Let \mathcal{D} be the corresponding flat family over $X \times C$ and $\pi_X : X \times C \rightarrow X$ the projection. Since $H^1(D) = 0$ for all D in this family, $(\pi_X)_*\mathcal{D}$ is a vector bundle over X whose fibre over any point of X corresponding to D is isomorphic to $H^0(D)$. Now consider the Grassmannian bundle of subspaces V^* of dimension k of the fibres of $(\pi_X)_*\mathcal{D}$ and the open subset Y of the total space G of this bundle consisting of those V^* for which (D, V^*) is generated. We have then an exact sequence on $Y \times C$

$$0 \rightarrow \mathcal{E}^* \rightarrow \mathcal{U} \rightarrow (\pi \times \text{id}_C)^*\mathcal{D} \rightarrow 0,$$

where \mathcal{U} is the pullback to $Y \times C$ of the universal subbundle on G and $\pi : Y \rightarrow X$ is the projection. The pair $(\mathcal{E}, \mathcal{U}^*)$ is now a family of coherent systems on C parametrised by the irreducible (or empty) variety Y . Since α -stability is an open condition, it follows that Z' is the image of an open subset of Y by some morphism and is therefore irreducible or empty.

Now let

$$Z'' := \{(E, V) \in G(\alpha; n, d, k) : (E, V) \text{ is generically generated}\}.$$

Then Z'' is an open subset of $G(\alpha; n, d, k)$ consisting of those (E, V) which have the form (25). By Lemma 4.3, $G(\alpha; n, d, k) \setminus Z''$ has dimension $< \beta$. It is therefore sufficient to prove that $\dim(Z'' \setminus Z') < \beta$. In fact, if $(E, V) \in Z'' \setminus Z'$ then, in the sequences (26) and (27), T has length $t > 0$ and $\deg E' = d - t$. For fixed t , the extensions (26) are classified by an open subset of a Quot-scheme Q . Tensoring (26) by D , we see from Lemmas 4.1(a) and 4.2 that $h^1(D \otimes E') = 0$. It follows that the dimension of Q at the point corresponding to (26) is

$$h^0(D \otimes E') = k(d - t) - n(k - n)(g - 1).$$

Taking account of the action of $\text{GL}(V)$ and (27), we see that the dimension of $Z'' \setminus Z'$ at (E, V) is at most

$$k(d - t) - n(k - n)(g - 1) + nt - (k^2 - 1) = \beta(n, d, k) - (k - n)t.$$

This completes the proof.

(d) For $k \leq n$, it is clear from (22) and (23) that the Petri map is injective at the generic point of $G(\alpha; n, d, k)$. For $n < k < n + \frac{1}{g}(d - n)$, the same follows from (24) and Lemmas 4.1 (b) and 4.2. Finally, for $k > n + \frac{1}{g}(d - n)$, $G(\alpha; n, d, k)$ consists of a single point and has rank $n \leq g - 1$, $d = 2n$ and $k = n + 1$ by Proposition 3.10; moreover $\beta(n, 2n, n + 1) = 0$ if and only if $n = g - 1$. \square

Corollary 4.5. *Suppose that $0 < d \leq 2n$. If $G_L(n, d, k) \neq \emptyset$, then it is smooth, except possibly when C is hyperelliptic and $(n, d, k) = (n, 2n, n + 1)$ with $n < g - 1$.*

Proof. For $k \leq n$, this is proved in [5, Theorems 5.4 and 5.6]. For $n < k \leq n + \frac{1}{g}(d - n)$, every element of $G_L(n, d, k)$ has the form (25) by [5, Proposition 4.4]. The result follows from Lemmas 4.1(b) and 4.2.

Finally, suppose that $k > n + \frac{1}{g}(d - n)$. Then $G_L(n, d, k)$ consists of a single point (E, V) by Proposition 3.10 and we have an exact sequence (24) with D a line bundle. In the non-hyperelliptic case, $D = K$ and E is a stable bundle of positive degree, so $h^1(D \otimes E) = 0$. In the hyperelliptic case, $D = L^a$ and $E \cong L^{\oplus a}$ for some $a \leq g - 1$, where L is the hyperelliptic line bundle. Under our hypotheses, this means that $a = g - 1$, so $D \otimes E \cong (L^g)^{\oplus (g-1)}$ and again $h^1(D \otimes E) = 0$. It follows from (24) that the Petri map is injective at (E, V) ; hence $G_L(g - 1, 2g - 2, g)$ is smooth. \square

Remark 4.6. If C is hyperelliptic and $(n, d, k) = (n, 2n, n + 1)$, $n < g - 1$, the Petri map cannot be injective for dimensional reasons. In this case $\beta(n, 2n, n + 1) < 0$ and $G_L(n, 2n, n + 1)$ consists of the single point $D(L^n, H^0(L^n))$, but we do not know whether or not it is reduced.

Corollary 4.7. *Suppose $0 < d \leq 2n$. If $B(n, d, k) \neq \emptyset$, then it is irreducible.*

Proof. Suppose $0 < d \leq 2n$ and $B(n, d, k) \neq \emptyset$. Then certainly $G_0(n, d, k) \neq \emptyset$. So, by Theorem 4.4, $G_0(n, d, k)$ is irreducible. Moreover, if $g \geq 3$, then $\beta(n, d, k) \leq n^2(g - 1)$, so [5, Conditions 11.3] are satisfied and the result follows from [5, Theorem 11.4]. If $g = 2$ and $k > d - n$, the same argument works. If $g = 2$ and $k \leq d - n$, Riemann-Roch implies that $B(n, d, k) = M(n, d)$ and is therefore irreducible. \square

Remark 4.8. Corollary 4.7 is an improvement on results obtained in [12] and [4].

5. NON-EMPTYNESS

We turn now to the question of non-emptiness of the moduli spaces. We begin by defining

$$U(n, d, k) := \{(E, V) \in G_L(n, d, k) : E \text{ is stable}\}$$

and

$$U^s(n, d, k) := \{(E, V) : (E, V) \text{ is } \alpha\text{-stable for } \alpha > 0, \alpha(n - k) < d\}.$$

Note that $U(n, d, k)$ can be defined alternatively as

$$U(n, d, k) := \{(E, V) : E \text{ is stable and } (E, V) \text{ is } \alpha\text{-stable for } \alpha > 0, \alpha(n - k) < d\}$$

and in particular $U(n, d, k) \subset U^s(n, d, k)$. In the converse direction, note that, if $(E, V) \in U^s(n, d, k)$, then E is semistable. However it is not generally true that $U(n, d, k) = U^s(n, d, k)$ and we can have $U^s(n, d, k) \neq \emptyset$, $U(n, d, k) = \emptyset$. Our object in this section is to determine when these sets are non-empty.

We begin with a lemma.

Lemma 5.1. *Suppose that $0 < d \leq 2n$, $k > n$ and $B(n, d, k) \neq \emptyset$. Then $U(n, d, k) \neq \emptyset$.*

Proof. If $k > n + \frac{1}{g}(d - n)$, then, by [3, 12, 14], the only possibilities for $E \in B(n, d, k)$ are as follows:

- if C is not hyperelliptic, $E \cong D(K)$;

- if C is hyperelliptic, $E \cong L$ (the hyperelliptic line bundle).

The result follows from Corollary 3.4.

Suppose now that $n < k \leq n + \frac{1}{g}(d-n)$ and that $B(n, d, k)$ is non-empty. If (E, V) is a coherent system with $E \in B(n, d, k)$, then $(E, V) \in G_0(n, d, k)$. Hence, by Theorem 4.4(c), $G_0(n, d, k)$ is irreducible and its generic element has the form

$$(29) \quad 0 \rightarrow D^* \rightarrow V \otimes \mathcal{O} \rightarrow E \rightarrow 0,$$

where $h^0(D^*) = 0$; also $h^1(D) = 0$ by Lemmas 4.1(a) and 4.2. As shown in the proof of Theorem 4.4, these extensions are parametrised by an irreducible variety. By openness of stability, the generic extension (29) has E stable as well as $(E, V) \in G_0(n, d, k)$. Furthermore D has rank $k - n$ and its degree d satisfies

$$d \geq g(k - n) + n \geq g(k - n) + \frac{d}{2},$$

so $d \geq 2g(k - n)$. Now any stable bundle D of this rank and degree is generated by its sections and

$$h^0(D) = d - (g - 1)(k - n) = d - g(k - n) - n + k \geq k;$$

hence the generic extension (29) also has D stable.

Finally, let us see that $(E, V) \in G_L(n, d, k)$ and hence $(E, V) \in U$. Suppose that $(E, V) \notin G_L(n, d, k)$; then there is a proper coherent subsystem (E', V') such that $\frac{k'}{n'} \geq \frac{k}{n}$. We can clearly suppose that (E', V') is generically generated. Since $(E, V) \in G_0(n, d, k)$, we have $\frac{d'}{n'} < \frac{d}{n}$. Now let $D' := D_{V'}(E')$. We have $(D')^* \subset D^*$, but

$$\frac{d'}{k' - n'} < \frac{dn'}{n(k' - n')} \leq \frac{dn'}{n'k - nn'} = \frac{d}{k - n}.$$

Since $\deg D^* = -d$, $\deg(D')^* \geq -d'$, this contradicts the stability of D^* . \square

We have a corresponding result for $U^s(n, d, k)$.

Complement 5.2. *Suppose that $0 < d \leq 2n$, $k > n$ and $G_0(n, d, k) \neq \emptyset$. Then $U^s(n, d, k) \neq \emptyset$.*

Proof. If $k > n + \frac{1}{g}(d - n)$ and $(E, V) \in G_0(n, d, k)$, then E is semistable and Proposition 3.10 implies that (E, V) is generated and $k = n + 1$. The result follows from Lemma 3.3. If $n < k \leq n + \frac{1}{g}(d - n)$, the proof of Lemma 5.1 still works. \square

We are now ready to prove our main results on non-emptiness. We will state the result separately for non-hyperelliptic and hyperelliptic curves and begin with a proposition which applies in both cases.

Proposition 5.3. *Suppose $n \geq 2$ and $0 < d \leq 2n$. Then $U(n, d, k) \neq \emptyset$ if and only if one of the following three conditions applies:*

- $0 < d < 2n$, $k \leq n + \frac{1}{g}(d - n)$, $(n, d, k) \neq (n, n, n)$;
- C is non-hyperelliptic, $d = 2n$ and either $k \leq n + \frac{n}{g}$ or $(n, d, k) = (g - 1, 2g - 2, g)$;
- C is hyperelliptic, $d = 2n$ and $k \leq n$.

Proof. For $k \leq n$, this is proved in [4, Theorem 3.3(v)].

For $k > n$, the stated conditions are precisely those for which $B(n, d, k) \neq \emptyset$ [3, 12, 14] and the result follows from Lemma 5.1. \square

Theorem 5.4. *Suppose that C is non-hyperelliptic of genus $g \geq 3$, $n \geq 2$ and $0 < d \leq 2n$. Then $U(n, d, k) \neq \emptyset$ if and only if either*

$$k \leq n + \frac{1}{g}(d - n), \quad (n, d, k) \neq (n, n, n)$$

or

$$(n, d, k) = (g - 1, 2g - 2, g).$$

In all other cases,

- $G(\alpha; n, d, k) = \emptyset$ for all $\alpha > 0$;
- $B(n, d, k) = \emptyset$.

Proof. The first part is just Proposition 5.3. The last part follows from Proposition 3.10. \square

Theorem 5.5. *Suppose that C is hyperelliptic, $n \geq 2$ and $0 < d \leq 2n$. Then*

(a) $U(n, d, k) \neq \emptyset$ if and only if either

$$0 < d < 2n, \quad k \leq n + \frac{1}{g}(d - n), \quad (n, d, k) \neq (n, n, n)$$

or $d = 2n, k \leq n$;

(b) if $k > n$, then

- $U(n, 2n, k) = \emptyset$;
- $U^s(n, 2n, k) \neq \emptyset$ if and only if either $k \leq n + \frac{n}{g}$ or $k = n + 1$ and $2 \leq n \leq g - 1$.

In all other cases,

- $G(\alpha; n, d, k) = \emptyset$ for all $\alpha > 0$;
- $B(n, d, k) = \emptyset$.

We already have enough information to prove this except for showing that $U^s(n, 2n, k) \neq \emptyset$ when $n < k \leq n + \frac{n}{g}$. This will be done in the next section.

Remark 5.6. The case $n = 1$ has been explicitly excluded from these statements as the results need modification. In this case the α -stability condition is redundant and the triples for which $0 < d \leq 2$ for which $U(1, d, k) \neq \emptyset$ are $(1, 1, 1)$, $(1, 2, 1)$ and, for C hyperelliptic, $(1, 2, 2)$.

6. PROOF OF THEOREM 5.5

In this section we suppose that C is hyperelliptic and L is the hyperelliptic line bundle. We assume that $n \geq 2$ and investigate by a sequence of propositions the case

$$(30) \quad d = 2n, \quad n < k \leq n + \frac{n}{g}$$

Proposition 6.1. *Suppose C is hyperelliptic. Then $U^s(n, 2n, n+1) \neq \emptyset$.*

Proof. Let $E = L^{\oplus n}$. Then E is generated and we can choose a subspace V of $H^0(E)$ of dimension $n+1$ such that (E, V) is generated. The result follows from Lemma 3.3. \square

Remark 6.2. Proposition 6.1 applies even when $n+1 > n + \frac{n}{g}$, in which case it has already been proved in Corollary 3.5.

Now suppose that $k \geq n+2$ and write $k = n+r$, so that (30) becomes

$$d = 2n, \quad 2 \leq r \leq \frac{n}{g}.$$

Proposition 6.3. *Suppose C is hyperelliptic and $2 \leq r \leq \frac{n-2}{g}$. Then $U^s(n, 2n, n+r) \neq \emptyset$.*

Proof. We consider extensions

$$(31) \quad 0 \longrightarrow (E_1, V_1) \longrightarrow (E, V) \longrightarrow (E_2, V_2) \longrightarrow 0,$$

where (E_1, V_1) has type

$$(n_1, d_1, k_1) = (n-1, 2n-3, n+r-1)$$

and (E_2, V_2) has type $(1, 3, 1)$. Certainly $(E_2, V_2) \in U(1, 3, 1)$. On the other hand $d_1 < 2n_1$ and

$$(32) \quad k_1 = n_1 + r \leq n_1 + \frac{n-2}{g} = n_1 + \frac{1}{g}(d_1 - n_1).$$

So, by Proposition 5.3, we can choose $(E_1, V_1) \in U(n_1, d_1, k_1)$. To show that there exist non-trivial extensions (31), it is sufficient to prove that $C_{21} > 0$. In fact, by (4),

$$(33) \quad \begin{aligned} C_{21} &= n_1(n_2 - k_2)(g-1) + (k_2 - n_2)d_1 + d_2n_1 - k_1k_2 \\ &= 3(n-1) - (n+r-1) = 2n-2-r \geq 2n-2 - \frac{n-2}{g} > 0. \end{aligned}$$

Suppose now that (31) is non-trivial. If $\alpha_c = \frac{n}{r}$, then

$$\mu_{\alpha_c}(E_1, V_1) = \frac{2n-3}{n-1} + \frac{n}{r} \cdot \frac{n+r-1}{n-1} = 3 + \frac{n}{r} = \mu_{\alpha_c}(E_2, V_2).$$

Since $\mu_{\alpha_c^-}(E_1, V_1) < \mu_{\alpha_c^-}(E_2, V_2)$, it follows from Proposition 2.5 that (E, V) is α_c^- -stable.

Now consider the extension of bundles

$$(34) \quad 0 \longrightarrow E_1 \longrightarrow E \longrightarrow E_2 \longrightarrow 0$$

underlying (31) and suppose first that this extension is non-trivial. If F is a subbundle of E which contradicts semistability, then certainly $F \not\subset E_1$. Moreover, if $F \rightarrow E_2$ is not surjective, then we have an extension

$$0 \longrightarrow F_1 \longrightarrow F \longrightarrow F_2 \longrightarrow 0$$

with $\deg F_2 \leq 2$ and $F_1 \subset E_1$, so $\mu(F) < 2$. It follows that, to contradict semistability of E , we must have

$$0 \longrightarrow F_1 \longrightarrow F \longrightarrow E_2 \longrightarrow 0.$$

Moreover $F_1 \neq 0$ since (34) does not split. Since E_1 is stable, $\mu(F_1) < \mu(E_1) < 2$, so

$$\deg F_1 \leq 2 \operatorname{rk} F_1 - 1, \quad \deg F = \deg F_1 + 3 \leq 2 \operatorname{rk} F.$$

So E is semistable.

To complete the proof in this case, it is sufficient by Complement 5.2 to show that $(E, V) \in U_0(n, 2n, n+r)$. If this is not the case, there exists a proper coherent subsystem (F, W) of type (n_F, d_F, k_F) of (E, V) with $d_F = 2n_F$ and $\frac{k_F}{n_F} \geq \frac{n+r}{n}$. But in this case (E, V) cannot be α -stable for any $\alpha > 0$, contradicting the fact that (E, V) is α_c^- -stable.

It remains to prove that there exist extensions (31) such that (34) does not split. Now, by [6, Corollaire 1.6] (see also [5, equation (7)]), we have an exact sequence

$$\operatorname{Hom}(V_2, H^0(E_1)/V_1) \longrightarrow \operatorname{Ext}^1((E_2, V_2), (E_1, V_1)) \longrightarrow \operatorname{Ext}^1(E_2, E_1).$$

It is therefore sufficient to prove that

$$\dim \operatorname{Ext}^1((E_2, V_2), (E_1, V_1)) > \dim \operatorname{Hom}(V_2, H^0(E_1)/V_1).$$

Now, by (33),

$$\dim \operatorname{Ext}^1((E_2, V_2), (E_1, V_1)) \geq C_{21} = 2n - 2 - r,$$

while

$$\dim \operatorname{Hom}(V_2, H^0(E_1)/V_1) = h^0(E_1) - (n + r - 1).$$

By [12, Chapitre 2, Théorème A.1], we have

$$h^0(E_1) \leq n_1 + \frac{1}{g}(d_1 - n_1) = n - 1 + \frac{n-2}{g},$$

so

$$\dim \operatorname{Hom}(V_2, H^0(E_1)/V_1) < \frac{n-2}{g} - r < C_{21} \leq \dim \operatorname{Ext}^1((E_2, V_2), (E_1, V_1)).$$

□

Remark 6.4. It is perhaps of interest to note that the coherent systems (E, V) constructed in this proof are not themselves in $U^s(n, 2n, n+r)$. We need to use Complement 5.2 to prove the proposition. Moreover the hypothesis $r \leq \frac{n-2}{g}$ is used in an essential way (see (32)) and the method of proof does not work without it; in fact, without the hypothesis, there are no flips.

It remains to consider the cases $r = \frac{n-1}{g}$ and $r = \frac{n}{g}$. In other words, we have two cases

$$n = gr + 1, \quad r \geq 2$$

and

$$n = gr, \quad r \geq 2.$$

Proposition 6.5. *Suppose C is hyperelliptic and $r \geq 2$. Then $U^s(gr+1, 2gr+2, gr+r+1) \neq \emptyset$.*

Proof. We consider extensions

$$(35) \quad 0 \longrightarrow (E_1, V_1) \longrightarrow (E, V) \longrightarrow (E_2, V_2) \longrightarrow 0,$$

where $(E_2, V_2) \cong D(L^{g-1}, H^0(L^{g-1}))$ and

$$(E_1, V_1) \in U^s(g(r-1) + 2, 2g(r-1) + 4, g(r-1) + r + 1),$$

which is non-empty by Propositions 6.1 and 6.3. By Theorem 4.4(c), we can suppose further that (E_1, V_1) is generated. Note also that (E_2, V_2) is generated and has the form $(L^{\oplus(g-1)}, V_2)$ with $\dim V_2 = g$, and belongs to $U^s(g-1, 2g-2, g)$ by Corollary 3.5.

We show first that there exists a non-trivial extension (35). In fact, by (4),

$$\begin{aligned} C_{21} &= n_1(n_2 - k_2)(g-1) + (k_2 - n_2)d_1 + d_2n_1 - k_1k_2 \\ &= -n_1(g-1) + 2n_1g - k_1g \\ &= n_1 + g(n_1 - k_1) = g(r-1) + 2 - g(r-1) = 2. \end{aligned}$$

From now on we suppose that (35) is non-trivial. Let (E', V') be a coherent subsystem of (E, V) of type $(n', d', n' + r')$ which contradicts 0^+ -stability. Then certainly E' is semistable of slope 2, so $d' = 2n'$, and

$$(36) \quad \frac{r'}{n'} \geq \frac{r}{n} = \frac{r}{gr+1}.$$

From (35), we have an extension

$$(37) \quad 0 \longrightarrow (E'_1, V'_1) \longrightarrow (E', V') \longrightarrow (E'_2, V'_2) \longrightarrow 0.$$

Since E_1, E_2 are semistable of slope 2, so are E'_1 and E'_2 . For $i = 1, 2$, denote the type of (E'_i, V'_i) by $(n'_i, 2n'_i, n'_i + r'_i)$. Note that $n'_1 \neq 0$, for otherwise (37) would contradict (36) except when $(E'_2, V'_2) = (E_2, V_2)$, in which case it would split the sequence (35). Since (E_1, V_1) is 0^+ -stable, we have

$$(38) \quad \frac{r'_1}{n'_1} < \frac{r_1}{n_1} = \frac{r-1}{g(r-1)+2}.$$

If $(E'_2, V'_2) \neq (E_2, V_2)$, then $r'_2 \leq 0$; this, together with (38), contradicts (36). Hence (37) becomes

$$0 \longrightarrow (E'_1, V'_1) \longrightarrow (E', V') \longrightarrow (E_2, V_2) \longrightarrow 0,$$

from which it follows that

$$n'_1 = n' - g + 1, \quad r'_1 = r' - 1.$$

A simple calculation shows that equations (36) and (38) can be written as

$$(39) \quad n'_1 \leq r'_1g + 1 + \frac{r'}{r}$$

and

$$(40) \quad n'_1 > r'_1g + \frac{2r'_1}{r-1}.$$

Since $r' < r$, (39) implies that $n'_1 < r'_1 g + 2$. By equation (40), this is only possible when

$$(41) \quad n'_1 = r'_1 g + 1, \quad r'_1 < \frac{r-1}{2}.$$

Note that (E'_1, V'_1) is a coherent subsystem of (E_1, V_1) . We must have $V'_1 = V_1 \cap H^0(E'_1)$, otherwise we could replace V'_1 by a subspace of $H^0(E'_1)$ of greater dimension, which would contradict (40). Thus we have an extension

$$(42) \quad 0 \longrightarrow (E'_1, V'_1) \longrightarrow (E_1, V_1) \longrightarrow (F, W) \longrightarrow 0.$$

We now count parameters to show that the (E_1, V_1) occurring in an extension (42) are not generic.

We begin with two lemmas.

Lemma 6.6. *(E'_1, V'_1) is generically generated and 0^+ -stable.*

Proof. If $r'_1 = 0$, then, by (41), (E'_1, V'_1) has type $(1, 2, 1)$ and the result is immediate. So suppose $r'_1 \geq 1$.

If (E'_1, V'_1) is not generically generated, it possesses a coherent subsystem (E''_1, V''_1) with $\text{rk } E''_1 \leq n'_1 - 1 = r'_1 g$. By α -stability of (E_1, V_1) for large α , this implies

$$\frac{r'_1 g + 1 + r'_1}{r'_1 g} \leq \frac{g(r-1) + r + 1}{g(r-1) + 2},$$

which is evidently false. On the other hand, if (E'_1, V'_1) is not 0^+ -stable, there exists a proper coherent subsystem (E''_1, V''_1) of type $(n''_1, 2n''_1, n''_1 + r''_1)$ such that

$$\frac{r''_1}{n''_1} \geq \frac{r'_1}{n'_1} = \frac{r'_1}{r'_1 g + 1},$$

i. e.

$$(43) \quad n''_1 \leq r''_1 \left(g + \frac{1}{r'_1} \right).$$

By α -stability of (E_1, V_1) , we have also

$$\frac{r''_1}{n''_1} < \frac{r-1}{g(r-1) + 2},$$

i. e.

$$(44) \quad n''_1 > r''_1 \left(g + \frac{2}{r-1} \right).$$

Now $n''_1 < n'_1$, so

$$r''_1 \left(g + \frac{2}{r-1} \right) < n'_1 - 1 = r'_1 g.$$

Hence $r''_1 < r'_1$ and (43) and (44) give a contradiction. \square

Lemma 6.7. $\text{Hom}(D(K, H^0(K)), (F, W)) = 0$.

Proof. Suppose that $\phi : D(K, H^0(K)) \rightarrow (F, W)$ is a non-zero homomorphism. Since $D(K, H^0(K))$ is α -stable for $\alpha > 0$, and F is semistable of slope 2, the image of ϕ is a coherent subsystem (F', W') of type $(n_{F'}, 2n_{F'}, n_{F'} + r_{F'})$ of (F, W) with

$$(45) \quad n_{F'} \leq g - 1, \quad r_{F'} \geq 1.$$

The pullback of (F', W') to (E_1, V_1) in (42) has type

$$(n_{F'} + n'_1, 2(n_{F'} + n'_1), n_{F'} + r_{F'} + n'_1 + r'_1).$$

Now α -stability of (E_1, V_1) gives

$$\frac{r_{F'} + r'_1}{n_{F'} + n'_1} \leq \frac{r - 1}{g(r - 1) + 2},$$

i. e.

$$n_{F'} + n'_1 \geq (r_{F'} + r'_1) \left(g + \frac{2}{r - 1} \right).$$

Since $n'_1 = r'_1 g + 1$ by (41), this is equivalent to

$$n_{F'} + 1 \geq r_{F'} \left(g + \frac{2}{r - 1} \right) + \frac{2r'_1}{r - 1}.$$

This contradicts (45). □

For our parameter count, we now establish three claims.

Claim 6.8. *For fixed (E'_1, V'_1) , (F, W) , the non-trivial extensions (42) for which (E_1, V_1) is generated and α -stable for some α depend on at most*

$$C_{21}^{(42)} - \dim \text{Aut}(F, W)$$

parameters, where $C_{21}^{(42)}$ denotes the value of C_{21} for the extensions (42).

Proof. By (3), we have

$$\dim \text{Ext}^1((F, W), (E'_1, V'_1)) = C_{21}^{(42)} + \dim \mathbb{H}_{21}^0 + \dim \mathbb{H}_{21}^2,$$

where

$$\mathbb{H}_{21}^0 = \text{Hom}((F, W), (E'_1, V'_1)), \quad \dim \mathbb{H}_{21}^2 = \text{Ext}^2((F, W), (E'_1, V'_1)).$$

Now $\mathbb{H}_{21}^0 = 0$ since otherwise (42) would give a contradiction to the α -stability of (E_1, V_1) . On the other hand, by (7) and Serre duality,

$$\mathbb{H}_{21}^2 \cong H^1(E'_1 \otimes N_2^*),$$

where N_2 is defined by an exact sequence

$$0 \longrightarrow N_2 \longrightarrow W \otimes \mathcal{O} \longrightarrow F \longrightarrow 0.$$

(Note that (F, W) is generated since (E_1, V_1) is.) By Lemmas 6.7 and 4.1(a), we have $h^1(N_2^*) = 0$. By Lemma 6.6, we have an exact sequence

$$0 \longrightarrow N_1 \longrightarrow V'_1 \otimes \mathcal{O} \longrightarrow E'_1 \longrightarrow T_1 \longrightarrow 0,$$

where T_1 is a torsion sheaf. Hence $h^1(E'_1 \otimes N_2^*) = 0$. Finally, since we are assuming (E_1, V_1) is α -stable for some α , the action of $\text{Aut}(F, W)$ on the extensions (42) is free. The result follows. □

Claim 6.9. (E'_1, V'_1) depends on at most

$$\beta(n'_1, 2n'_1, k'_1)$$

parameters.

Proof. Since (E'_1, V'_1) is 0^+ -stable by Lemma 6.6, it is sufficient to show that the Petri map is injective. This follows from Lemmas 4.1(b) and 4.2 and (41). \square

Claim 6.10. (F, W) depends on at most

$$\beta(n_1 - n'_1, 2n_1 - 2n'_1, k_1 - k'_1) + \dim \text{Aut}(F, W) - 1$$

parameters.

Proof. By (8), it is sufficient to show that

$$\dim \text{Ext}^1((F, W), (F, W)) \leq \beta(n_1 - n'_1, 2n_1 - 2n'_1, k_1 - k'_1) + \dim \text{Aut}(F, W) - 1.$$

This is equivalent to proving that the Petri map of (F, W) is injective, which follows from Lemmas 6.7 and 4.1(b). \square

Completion of proof of Proposition 6.5. By the above claims and (6), the (E_1, V_1) for which (E'_1, V'_1) exists satisfying (39) and (40) depend on at most

$$\beta(n_1, 2n_1, k_1) - C_{12}^{(42)}$$

parameters. If $C_{12}^{(42)} > 0$, it follows that the general (E_1, V_1) contains no (E'_1, V'_1) with these properties. Hence (E, V) contains no subsystem (E', V') contradicting 0^+ -stability. The result then follows from Complement 5.2.

It remains to calculate $C_{12}^{(42)}$. In fact, by (5),

$$\begin{aligned} C_{12}^{(42)} &= (n_1 - n'_1)(n'_1 - k'_1)(g - 1) + 2(n_1 - n'_1)(k'_1 - n'_1) + 2n'_1(n_1 - n'_1) - k'_1(k_1 - k'_1) \\ &= (n_1 - n'_1)[(n'_1 - k'_1)(g - 1) + 2k'_1] - k'_1(k_1 - k'_1) \\ &= (g(r - 1 - r'_1) + 1)(-r'_1)(g - 1) + k'_1(2n_1 - 2n'_1 - k_1 + k'_1) \\ &= -(r - 1 - r'_1)(g - 1)r'_1g - r'_1(g - 1) + (r'_1(g + 1) + 1)((g - 1)(r - 1 - r'_1) + 1) \\ &= (r - 1 - r'_1)(g - 1)(r'_1 + 1) + 2r'_1 + 1 > 0. \end{aligned}$$

\square

Proposition 6.11. *Suppose C is hyperelliptic and $r \geq 2$. Then $U^s(gr, 2gr, gr + r) \neq \emptyset$.*

Proof. The proof is similar to that of Proposition 6.5; we outline below the necessary changes.

We consider sequences (35), where $(E_2, V_2) \cong D(K, H^0(K))$ as before, and now

$$(E_1, V_1) \in U^s(g(r - 1) + 1, 2g(r - 1) + 2, g(r - 1) + r).$$

This space is non-empty by Propositions 6.1 and 6.5. We have

$$C_{21} = n_1 + g(n_1 - k_1) = 1,$$

so non-trivial extensions (35) exist. After modifying (36) and (38), we proceed to (39) and (40), which become

$$n'_1 \leq r'_1 g + 1, \quad n'_1 > r'_1 g + \frac{r'_1}{r-1}.$$

So again $n'_1 = r'_1 g + 1$, where now $r'_1 < r - 1$.

The proofs of Lemmas 6.6 and 6.7 are the same as before, replacing $g(r-1) + 2$ by $g(r+1) + 1$ with consequential changes which don't affect the argument. The only remaining thing to be checked is that $C_{12}^{(42)} > 0$. In fact

$$\begin{aligned} C_{12}^{(42)} &= (n_1 - n'_1)[(n'_1 - k'_1)(g-1) + 2k'_1] - k'_1(k_1 - k'_1) \\ &= g(r-1-r'_1)(-r'_1)(g-1) + k'_1(2n_1 - 2n'_1 - k_1 + k'_1) \\ &= -(r-1-r'_1)(g-1)r'_1 g + (r'_1(g+1) + 1)(g-1)(r-1-r'_1) \\ &= (r-1-r'_1)(g-1)(r'_1 + 1) > 0. \end{aligned}$$

□

Proof of Theorem 5.5. (a) is just Proposition 5.3. (b) follows from Propositions 6.1, 6.3, 6.5 and 6.11 in the case $k \leq n + \frac{n}{g}$ and from Corollary 3.5 if $k > n + \frac{n}{g}$. The last part follows from Proposition 3.10. □

7. AN EXAMPLE WITH $d > 2n$

We have seen that, when $d < 2n$ and $k > n$ and α -stable coherent systems exist for some α (i. e. when (9) holds), then there exist coherent systems (E, V) such that E is stable and (E, V) is α -stable for all $\alpha > 0$. The same applies when $d = 2n$ if C is not hyperelliptic. If C is hyperelliptic of genus $g \geq 3$ and $a \geq 2$, the coherent systems $(L^{\oplus a}, W)$ of type $(a, 2a, a+1)$ are α -stable for all α by Corollary 3.5, but $L^{\oplus a}$ is only semistable. Moreover, when $d \leq 2n$, there is no case in which there exist semistable bundles but α -stable coherent systems do not exist for large α . The object of this section is to construct such examples, necessarily with $d > 2n$.

Lemma 7.1. *Suppose (E, V) is a coherent system of type (n, d, k) with*

$$(46) \quad n + \frac{1}{g}(d-n) < k < \frac{ng}{g-1}.$$

Then (E, V) is not α -semistable for large α .

Proof. Suppose (46) is satisfied. By Corollary 3.8, there exists a non-zero homomorphism

$$D(K, H^0(K)) \longrightarrow (E, V).$$

By Corollaries 3.4 and 3.5, $D(K, H^0(K))$ is α -stable for all $\alpha > 0$. Moreover

$$\mu_\alpha(D(K, H^0(K))) = 2 + \alpha \frac{g}{g-1} > \frac{d}{n} + \alpha \frac{k}{n}$$

for sufficiently large α by (46). This contradicts the α -semistability of (E, V) for large α . □

Proposition 7.2. *Suppose that C is not hyperelliptic and $3 \leq r \leq g - 1$. Then there exists a coherent system (E, V) of type*

$$(47) \quad (rg - r + 1, 2rg - 2r + 3, rg + 1)$$

with E stable. Moreover (46) is satisfied.

Proof. It is clear that (46) follows from (47) and the assumption $r \geq 3$.

Consider the extensions

$$(48) \quad 0 \longrightarrow D(K)^{\oplus r} \oplus \mathcal{O}(p_1, p_2) \longrightarrow E \longrightarrow \mathcal{O}_q \longrightarrow 0,$$

where $p_1, p_2, q \in C$. We take V to be the image of $H^0(D(K)^{\oplus r} \oplus \mathcal{O}(p_1, p_2))$ in $H^0(E)$. The fact that (E, V) is of type (47) is clear. Since $D(K)$ and $\mathcal{O}(p_1, p_2)$ are both stable of slope 2, E fails to be stable only if it admits either $D(K)$ or $\mathcal{O}(p_1, p_2)$ as a quotient. Equivalently E fails to be stable only if some factor of $D(K)$ or $\mathcal{O}(p_1, p_2)$ splits off (48). Now the extensions (48) are classified by elements

$$\begin{aligned} e = (e_1, \dots, e_r, e_{r+1}) &\in \text{Ext}^1(\mathcal{O}_q, D(K)^{\oplus r} \oplus \mathcal{O}(p_1, p_2)) \\ &= \text{Ext}^1(\mathcal{O}_q, D(K))^{\oplus r} \oplus \text{Ext}^1(\mathcal{O}_q, \mathcal{O}(p_1, p_2)). \end{aligned}$$

Since $D(K)$ and $\mathcal{O}(p_1, p_2)$ are stable and non-isomorphic, it follows that E is stable if e_1, \dots, e_r are linearly independent and $e_{r+1} \neq 0$. Since

$$\dim \text{Ext}^1(\mathcal{O}_x, D(K)) = g - 1,$$

it is possible to choose such e_1, \dots, e_r, e_{r+1} whenever $r \leq g - 1$. This completes the proof. \square

ACKNOWLEDGMENTS

The fourth author wishes to acknowledge Universidad Complutense de Madrid and the Institute for Advanced Study, Princeton for their hospitality and for providing excellent working conditions. The fifth author would like to thank CIMAT, Guanajuato, Mexico and California State University Channel Islands, where parts of this research were carried out.

REFERENCES

- [1] M. F. Atiyah, Vector bundles over an elliptic curve, *Proc. London Math. Soc.* **7**(3) (1957), 414–452.
- [2] S. B. Bradlow and O. García-Prada, An application of coherent systems to a Brill-Noether problem, *J. Reine Angew. Math.* **551** (2002), 123–143.
- [3] L. Brambila-Paz, I. Grzegorczyk and P. E. Newstead, Geography of Brill-Noether loci for small slopes, *J. Algebraic Geometry* **6** (1997), 645–669.
- [4] S. B. Bradlow, O. García-Prada, V. Mercat, V. Muñoz and P. E. Newstead, On the geometry of moduli spaces of coherent systems on algebraic curves, *Internat. J. Math.* **18** (2007), 411–453.
- [5] S. B. Bradlow, O. García-Prada, V. Muñoz and P. E. Newstead, Coherent Systems and Brill-Noether theory, *Internat. J. Math.* **14** (2003), 683–733.
- [6] M. He, Espaces de modules de systèmes cohérents, *Internat. J. Math.* **9** (1998), 545–598.
- [7] A. D. King and P. E. Newstead, Moduli of Brill-Noether pairs on algebraic curves, *Internat. J. Math.* **6** (1995) 733–748.
- [8] H. Lange and P. E. Newstead, Coherent systems of genus 0, *Internat. J. Math.* **15** (2004), 409–424.

- [9] H. Lange and P. E. Newstead, Coherent systems on elliptic curves, *Internat. J. Math.* **16** (2005), 787–805.
- [10] H. Lange and P. E. Newstead, Coherent systems of genus 0 II: existence results for $k \geq 3$, *Internat. J. Math.* **18** (2007), 363–393.
- [11] J. Le Potier, Faisceaux semi-stables et systèmes cohérents, *Vector Bundles in Algebraic Geometry, Durham 1993*, ed. N.J. Hitchin, P.E. Newstead and W.M. Oxbury, LMS Lecture Notes Series **208**, 179–239, Cambridge University Press, 1995.
- [12] V. Mercat, Le problème de Brill-Noether pour des fibrés stables de petite pente, *J. Reine Angew. Math.* **506** (1999), 1–41.
- [13] V. Mercat, Le problème de Brill-Noether et le théorème de Teixidor, *Manuscripta Math.* **98** (1999), 75–85.
- [14] V. Mercat, Fibrés stables de pente 2, *Bull. London Math. Soc.* **33** (2001), 535–542.
- [15] M. S. Narasimhan and S. Ramanan, Deformation of the moduli space of vector bundles over an algebraic curve, *Ann. of Math. (2)* **82** (1965), 540–567.
- [16] K. Paranjape and S. Ramanan, On the canonical ring of a curve, *Algebraic Geometry and Commutative Algebra in Honor of Masayoshi Nagata (1987)*, 503–516.
- [17] N. Raghavendra and P.A. Vishwanath, Moduli of pairs and generalized theta divisors, *Tôhoku Math. J.* **46** (1994) 321–340.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, IL 61801, USA

E-mail address: bradlow@math.uiuc.edu

INSTITUTO DE MATEMÁTICAS, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, SERRANO, 121, 28006 MADRID, SPAIN

E-mail address: oscar.garcia-prada@uam.es

5 RUE DELOUVAIN, 75019 PARIS, FRANCE

E-mail address: mercat@math.jussieu.fr

DEPARTAMENTO DE MATEMÁTICAS, INSTITUTO DE MATEMÁTICAS Y FÍSICA FUNDAMENTAL, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, SERRANO, 113 BIS, 28006 MADRID, SPAIN

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE CIENCIAS 3, 28040 MADRID, SPAIN

E-mail address: vicente.munoz@imaff.cfmac.csic.es

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF LIVERPOOL, PEACH STREET, LIVERPOOL L69 7ZL, UK

E-mail address: newstead@liverpool.ac.uk