

Research article

Open Access

Molecular characterisation of *Mycobacterium bovis* isolated from cattle slaughtered at the Bamako abattoir in Mali

Borna Müller*¹, Benjamin Steiner¹, Bassirou Bonfoh^{2,6}, Adama Fané³, Noel H Smith^{4,5} and Jakob Zinsstag¹

Address: ¹Swiss Tropical Institute, Basel, Switzerland, ²Institut du Sahel, Bamako, Mali, ³Laboratoire Centrale Vétérinaire, Bamako, Mali, ⁴Veterinary Laboratories Agency, Weybridge, UK, ⁵University of Sussex, Falmer, UK and ⁶Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire

Email: Borna Müller* - borna.mueller@unibas.ch; Benjamin Steiner - steiner@baden.ch; Bassirou Bonfoh - bassirou.bonfoh@csrs.ci; Adama Fané - afaneother@yahoo.fr; Noel H Smith - noel@sussex.ac.uk; Jakob Zinsstag - jakob.zinsstag@unibas.ch

* Corresponding author

Published: 17 July 2008

Received: 8 April 2008

BMC Veterinary Research 2008, 4:26 doi:10.1186/1746-6148-4-26

Accepted: 17 July 2008

This article is available from: <http://www.biomedcentral.com/1746-6148/4/26>

© 2008 Müller et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Mali is one of the most important livestock producers of the Sahel region of Africa. A high frequency of bovine tuberculosis (BTB) has been reported but surveillance and control schemes are restricted to abattoir inspections only. The objective of this study was to conduct, for the first time, molecular characterisation of *Mycobacterium bovis* strains isolated from cattle slaughtered at the Bamako abattoir. Of 3330 animals screened only 60 exhibited gross visible lesions. From these animals, twenty strains of *M. bovis* were isolated and characterised by spoligotyping.

Results: Organ lesions typical of BTB were most often detected in the liver, followed by the lung and the peritoneum. *M. bovis* was isolated from 20 animals and 7 different spoligotypes were observed among these 20 strains; three of the patterns had not been previously reported. Spoligotype patterns from thirteen of the strains lacked spacer 30, a characteristic common in strains of *M. bovis* found in Chad, Cameroon and Nigeria. However, unlike the other three Central African countries, the majority of spoligotype patterns observed in Mali also lacked spacer 6. Of the remaining seven strains, six had spoligotype patterns identical to strains commonly isolated in France and Spain.

Conclusion: Two groups of *M. bovis* were detected in cattle slaughtered at the Bamako abattoir. The spoligotype pattern of the first group has similarities to strains previously observed in Chad, Cameroon and Nigeria. The additional absence of spacer 6 in the majority of these strains suggests a Mali specific clone. The spoligotype patterns of the remaining strains suggest that they may have been of European origin.

Background

Bovine Tuberculosis (BTB) is considered a neglected and poverty related zoonosis [1]. It has a major economic impact on livestock productivity [2], can persist in wildlife

reservoirs and thus affect entire ecosystems [3] and it is of public health concern due to its zoonotic potential [4-6]. Although still prevalent in the developed world [7-9], BTB today mostly affects developing countries, which lack the

financial and human resources to control the disease [4,5]. The Sahel region of Africa is extremely important in terms of animal production with Mali being amongst the principal cattle producing countries [10,11]. Mali has previously reported a high frequency of BTB but does not apply specific control measures, except carcass inspection at abattoirs [4]. In a recent prevalence study in dairy cattle herds from the peri-urban region of Bamako, 19% of the animals reacted positively to the comparative tuberculin skin test [12].

Spacer oligonucleotide typing (spoligotyping) [13] has been shown to be a valuable tool for investigations of the population structure of *Mycobacterium bovis* in a number of settings [7,8,14-16]. Furthermore, the international designation of spoligotype patterns <http://www.Mbovis.org>, [17] has facilitated the comparison of results from different countries and helps elucidate the distribution and spread of strains. Assuming that spoligotype spacers can only be lost and not regained, phylogenetic relationships between strains can be suggested [18,19]. A number of *Mycobacterium tuberculosis* Complex (MTBC) strain families are readily identifiable through spoligotyping [20-22].

Variable number of tandem repeat (VNTR) typing is another simple method for *M. bovis* genotyping with a higher discriminatory power than spoligotyping [19,23]. However, extensive worldwide databases are presently not available and VNTR typing can today mainly be considered a valuable tool for sub-differentiation of strain groups initially identified by spoligotyping [19].

For *M. bovis*, previous studies in Chad, Cameroon and Nigeria have shown that virtually all spoligotype patterns lack spacer 30 with strains bearing spoligotype pattern SB0944 being the most frequent [14-16]. In Cameroon, because of the similarity to patterns of strains isolated in France [7] it was suggested that *M. bovis* could have been imported to this region during the French colonial period [15]. However, strains lacking spacer 30 were so far rarely found outside Chad, Cameroon and Nigeria.

The objectives of this study were to conduct an initial molecular characterisation of *M. bovis* in Mali using spoligotyping and to identify potential exchange of strains with other regions.

Results

At the abattoir of Bamako, Mali, a case series of 3330 slaughter animals were sequentially screened during standard meat inspection in March and April 2007. A total of 182 specimens from 60 animals with gross visible lesions (apparent lesion prevalence: 1.8%; 95% CI: 1.4 – 2.3%) were collected. Organ lesions were most often

detected in the liver (N = 22) followed by the lung (N = 14) and the peritoneum (N = 11). The specimens were put in culture and Acid-Fast Bacilli were further characterised by spoligotyping and typing of the *M. bovis* specific RD4 region. Infection with *M. bovis* was confirmed for 20 animals. From two animals, strains of the *Mycobacterium fortuitum* Complex were isolated as identified by partial sequencing of the 16S rRNA gene. In one case it appeared to be a single infection and in the other case a mixed infection of an *M. fortuitum* Complex strain and *M. bovis*. In this animal, the *M. fortuitum* Complex strain was isolated from liver lesions and *M. bovis* was isolated from lesions of the lungs and bronchial lymph nodes.

Infection with *M. bovis* was highly associated with the presence of lung lesions (N = 44, chi-squared = 23.7, $p < 0.001$); in 79% of the animals exhibiting lung lesions, *M. bovis* infection could be confirmed. The association was less strong for liver lesions (N = 48, chi-squared = 3.9, $p < 0.05$); only 41% of the animals with liver lesions were shown to be infected with *M. bovis*. However, in all 9 cases where *M. bovis* infection was detected in animals with liver lesions, lung lesions were present as well. No association was found between infection with *M. bovis* and lesions in organs other than the liver and lungs. Strains of *M. bovis* isolated in different organs of the same animal showed the same spoligotype pattern. Altogether among the 20 strains of *M. bovis* isolated, seven different spoligotypes were observed; four had been previously reported (SB0944, SB0300, SB0134 and SB0944) and the remaining three were designated SB1410, SB1411 and SB1412 by <http://www.Mbovis.org> (figure 1).

The distinctive lack of spacer 30 was observed in the spoligotype pattern of 13 strains; the majority of those in addition lacked spacer 6 (12/13; figure 1). The remaining seven strains were linked by the absence of spacers 4 and 5 (figure 1).

VNTR typing using the exact tandem repeats (ETR) A-F described by Frothingham et al. [24] allowed to further differentiate strains with the most frequent spoligotype patterns SB0300 and SB0134 (figure 1). Within the strains analysed, only VNTR loci ETR A, B and C showed variation; ETR D, E and F profiles were identical in all the strains. One isolate exhibited two different VNTR alleles (3 and 5 tandem repeats) for locus ETR C (figure 1), indicating either a mixed infection with two distinct strains or a microevolution in this population of strains.

Discussion

The apparent prevalence of 1.8% gross visible lesions in Malian slaughter cattle was surprisingly low compared to published results from other Sahelian countries [14,25,26] and also lower than the previous tuberculin

by Fulbe pastoralists [30]. However, considering the fact that, except for pattern SB0944, none of the spoligotype patterns found in Mali are present in any of the three Central African countries and considering that strains with spoligotype pattern SB0944 were rarely detected in Mali we suppose that the spread of *M. bovis* strains over this large distance is relatively slow.

The second group of related spoligotype patterns is characterized by the absence of spacers 4 and 5 (figure 1). The most often detected spoligotype belonging to this group is commonly found in strains from France and Spain (SB0134 [7,31]), suggesting a link between *M. bovis* strains from Mali and mainland Europe. VNTR profile 6 5 5 4 for ETR loci A-D identified in 2/6 Malian strains of *M. bovis* with spoligotype pattern SB0134 has also been detected in a strain isolated from French cattle in the Normandy in 1996 with the same spoligotype pattern [32]. Moreover, three other Malian *M. bovis* strains with ETR A-D profile 7 5 5 4 and spoligotype pattern SB0134 could be closely related to SB0134 *M. bovis* strains with ETR A-D profile 7 4 5 4, which is frequently found in *M. bovis* strains from the Normandy [32]. However, identical spoligotype patterns have also been found in strains from northern Algeria (unpublished results) and livestock migrations from Algeria to Mali through the Sahara desert have been reported. Comprehensive genotyping of strains from West Africa, North Africa and Europe using highly polymorphic markers would be necessary to further elucidate the interrelationship of *M. bovis* strains from these different regions.

Njanpop-Lafourcade et al., have previously suggested an influence of the French colonial history based on the similarity of the predominant spoligotype pattern (SB0944) in Cameroon to the BCG-like spoligotype pattern that is commonly seen in strains from France [15]. In a similar manner it is possible to suggest that strains of *M. bovis* with spoligotype pattern SB0134 were originally imported from Europe. If both assumptions are true this would suggest that either *M. bovis* was not present in the Central or West African region before introduction from Europe to Africa or previously existing "native" *M. bovis* strains have been largely replaced.

Due to the small sample size, the limited survey period and the sampling at only one study location, the *M. bovis* strains collected cannot reflect the country-wide bacterial population structure. Therefore, frequencies of strains with a specific spoligotype may not necessarily mirror the actual frequency of these strains in the population although more frequent strains are also more likely to be detected in a random sample. Moreover, other groups of *M. bovis* strains than the two that were observed may be present in Mali. However, due to the predominant long

distance transhumant livestock production system, we believe that the slaughter cattle encountered at the abattoir of Bamako and consequently their associated *M. bovis* strains, represent a sample from a large area of the country.

Conclusion

This study presents the first molecular characterisation of *M. bovis* strains from Mali. The results suggest that the most often detected strains are related to strains that are prevalent in Chad, Cameroon and Nigeria. A second group of strains shows spoligotype patterns similar to those abundant in mainland Europe and could have been imported directly from Europe or via Northern Africa. Our results can serve as a baseline study for future comparisons with strains from other areas in and around Mali.

Methods

Sample collection

Samples were collected from a sequential series of slaughter cattle at the Bamako abattoir, Mali in March and April 2007. The cattle population consisted of crossbreeds between N'Dama, zebu and exotic breeds. The origin of the cattle could not be traced due to poor documentation and multiple selling-on of the animals before slaughter. However, we originally assumed that the cattle originate from the peri-urban region of Bamako as well as the principle areas of cattle production throughout the country. After slaughter, animals underwent a standard meat inspection and organs showing gross visible lesions were confiscated. No ethical clearance was required for this study because it was done on slaughtered animals and organ confiscation was part of a routine monitoring. Tissue samples of 60 animals with gross visible lesions were collected. The samples were transported on ice to the Central Veterinary Laboratory in Bamako. Upon arrival, samples were immediately liberated from connective tissue and fat under a bio-safety cabinet and by use of sterile dissection instruments. The samples were seared on the outside in order to reduce superficial contamination, sealed into sterile stomacher bags and stored at -20°C for maximum one and a half months until they were shipped to Switzerland for culture. Because of frequent power cuts we cannot exclude that some of the specimen might have undergone multiple freeze-thaw cycles while they were stored. Shipment to Switzerland occurred in a refrigerated box; the temperature of the samples was monitored at all times by use of a data-logger (HoboTemp by OnsetCorp) and never exceeded -10°C.

Tissue preparation, culture and DNA extraction

At the Swiss Reference Centre for Mycobacteria, samples were stored at -80°C until processed. Specimen were dissected and approximately 2 g were homogenised for 2 minutes in 10 ml phosphate buffer saline (PBS) using the

ULTRA-TURRAX® Tube Drive homogeniser with DT-20 tubes (IKA, Staufen, Germany). A 5 ml aliquot of the suspension was decontaminated for 15 minutes with 5 ml of decontamination solution (0.5% N-acetyl-L-cystein/2% NaOH/1.45% Na-Citrate solution). The decontamination was stopped by addition of 15 ml PBS, the suspension was centrifuged at 3500 rpm for 15 minutes and the pellet was re-suspended in 2 ml PBS. Then, 0.5 ml of the suspension was added to a BBL™ MGIT™ Mycobacterium Growth Indicator Tube containing OADC enrichment and PANTA™ antibiotic mixture (BD) and incubated in a BACTEC™ MGIT™ 960 Mycobacterial Detection System. In addition, 0.25 ml of the suspension was inoculated onto Löwenstein-Jensen and 7H10 culture media and incubated at 37°C. Cultures were incubated until growth was detected or for at least 8 weeks. Presence of Acid-Fast Bacilli was tested by Ziehl-Neelson staining and Microscopy and DNA of positive cultures was extracted using the InstGene™ Matrix (Bio-Rad®).

Molecular characterisation

Spoligotyping was conducted as previously described [13]. VNTR typing was performed according to the method of Frothingham et al. [24] with adaptations described elsewhere [33]. According to the findings of Brosch et al., strains were confirmed as *M. bovis* by the absence of region RD4. [34]. The 16S rRNA gene amplification and sequencing was carried out as described by Zucol et al., (2006) [35]. Species identification was carried out by comparison with the sequences of the SmartGene Integrated Database Network System (IDNS™) 3.4.0.

Statistical analysis

Statistical analysis was carried out using Intercooled Stata 9.2 for Windows (StataCorp LP, USA). The association between the presence of lesions in each organ and confirmed *M. bovis* infection was tested by a chi-squared test.

Authors' contributions

BM: Participated in the conception and design of the study, culture of Mycobacteria, molecular analysis, statistical analysis, writing of the manuscript. BS: Participated in the conception and design of the study, sample collection, culture of Mycobacteria. BB: Participated in the conception and design of the study, acquisition of funds, principle supervision in Mali and intellectual contributions. AF: Principle organisation of sample collection NHS: Contribution in analysis of the data, important intellectual contribution for population genetical analysis. JZ: Principal supervision of the project, principle acquisition of funds, important intellectual contribution for epidemiological questions. All the authors have read and approved the final version of the manuscript.

Acknowledgements

We would like to thank Prof. Erik C. Böttger and the technicians of the Swiss National Centre for Mycobacteria in Zurich for assistance and laboratory facilities for culturing of Mycobacteria. We would also like to thank Prof. R. Glyn Hewinson and Dr. Stefan Berg from the Veterinary Laboratories Agency (VLA) in Weybridge, UK for providing technical support and laboratory facilities for spoligotyping and K. Gover for technical assistance. Moreover, we are indebted to the director of the Laboratoire Centrale Vétérinaire in Bamako and the meat inspectors at the abattoirs for providing laboratory facilities and assistance during the sample collection. Our work has received financial support from the Swiss National Science Foundation (project no. 107559).

References

1. WHO: **The Control of Neglected Zoonotic Diseases. A route to poverty alleviation.** Geneva, World Health Organization; 2006.
2. Zinsstag J, Schelling E, Roth F, Kazwala RR: **Economics of bovine tuberculosis.** In *Mycobacterium bovis Infection in Animals and Humans* 2nd edition. Edited by: Thoen CO, Steele JH and Gilsdorf MJ. Blackwell Publishing; 2006.
3. Renwick AR, White PC, Bengis RG: **Bovine tuberculosis in southern African wildlife: a multi-species host-pathogen system.** *Epidemiol Infect* 2006; **1-12**.
4. Cosivi O, Grange JM, Daborn CJ, Raviglione MC, Fujikura T, Cousins D, Robinson RA, Huchzermeyer HF, de K I, Meslin FX: **Zoonotic tuberculosis due to Mycobacterium bovis in developing countries.** *Emerg Infect Dis* 1998, **4:59-70**.
5. Ayele WY, Neill SD, Zinsstag J, Weiss MG, Pavlik I: **Bovine tuberculosis: an old disease but a new threat to Africa.** *Int J Tuberc Lung Dis* 2004, **8:924-937**.
6. Thoen C, Lobue P, de K I: **The importance of Mycobacterium bovis as a zoonosis.** *Vet Microbiol* 2006, **112:339-345**.
7. Haddad N, Ostyn A, Karoui C, Masselot M, Thorel MF, Hughes SL, Inwald J, Hewinson RG, Durand B: **Spoligotype diversity of Mycobacterium bovis strains isolated in France from 1979 to 2000.** *Journal of Clinical Microbiology* 2001, **39:3623-3632**.
8. Aranaz A, Liebana E, Mateos A, Dominguez L, Vidal D, Domingo M, Gonzalez O, Rodriguez-Ferri EF, Bunschoten AE, Vanembden JDA, Cousins D: **Spacer oligonucleotide typing of Mycobacterium bovis strains from cattle and other animals: A tool for studying epidemiology of tuberculosis.** *Journal of Clinical Microbiology* 1996, **34:2734-2740**.
9. Smith NH, Dale J, Inwald J, Palmer S, Gordon SV, Hewinson RG, Smith JM: **The population structure of Mycobacterium bovis in Great Britain: clonal expansion.** *Proc Natl Acad Sci U S A* 2003, **100:15271-15275**.
10. Dicko MS, Djitéye MA, Sangaré M: **Les systèmes de production animale au Sahel.** *Secheresse* 2006, **17:83-97**.
11. Wilson RT: **Livestock production in central Mali: Long-term studies on cattle and small ruminants in the agropastoral system.** Volume 14. Addis Ababa, Ethiopia, ILCA; 1986.
12. Sidibé SS, Dicko NA, Fané A, Doumbia RM, Sidibé CK, Kanté S, Mangané O, Konaté B, Koné AZ, Maïga MS, Fofana M: **Tuberculose bovine au Mali : résultats d'une enquête épidémiologique dans les élevages laitiers de la zone périurbaine du district de Bamako.** *Revue Elev Méd vét Pays trop* 2003, **56:115-120**.
13. Kamerbeek J, Schouls L, Kolk A, van Agterveld M, van Soolingen D, Kuijper S, Bunschoten A, Molhuizen H, Shaw R, Goyal M, Van Embden J: **Simultaneous detection and strain differentiation of Mycobacterium tuberculosis for diagnosis and epidemiology.** *J Clin Microbiol* 1997, **35:907-914**.
14. Diguimbaye-Djaïbe C, Hilty M, Ngandolo R, Mahamat HH, Pfyffer GE, Baggi F, Hewinson G, Tanner M, Zinsstag J, Schelling E: **Mycobacterium bovis isolates from tuberculous lesions in Chadian zebu carcasses.** *Emerg Infect Dis* 2006, **12:769-771**.
15. Njanpop-Lafourcade BM, Inwald J, Ostyn A, Durand B, Hughes S, Thorel MF, Hewinson G, Haddad N: **Molecular typing of Mycobacterium bovis isolates from Cameroon.** *J Clin Microbiol* 2001, **39:222-227**.
16. Cadmus S, Palmer S, Okker M, Dale J, Gover K, Smith N, Jahans K, Hewinson RG, Gordon SV: **Molecular analysis of human and bovine tubercle bacilli from a local setting in Nigeria.** *J Clin Microbiol* 2006, **44:29-34**.

17. Brudey K, Driscoll JR, Rigouts L, Prodinger WM, Gori A, Al-Hajj SA, Allix C, Aristimuno L, Arora J, Baumanis V, Binder L, Cafrune P, Cataldi A, Cheong S, Diel R, Ellermeier C, Evans JT, Fauville-Dufaux M, Ferdinand S, Garcia de Viedma D, Garzelli C, Gazzola L, Gomes HM, Guttierrez MC, Hawkey PM, van Helden PD, Kadival GV, Kreiswirth BN, Kremer K, Kubin M, Kulkarni SP, Liens B, Lillebaek T, Ly HM, Martin C, Martin C, Mokrousov I, Narvskaja O, Ngeow YF, Naumann L, Niemann S, Parwati I, Rahim Z, Rasolof-Razanamparany V, Rasolonavalona T, Rossetti ML, Rusch-Gerdes S, Sajuda A, Samper S, Shemyakin IG, Singh UB, Somoskovi A, Skuce RA, van Soolingen D, Streicher EM, Suffys PN, Tortoli E, Tracevska T, Vincent V, Victor TC, Warren RM, Yap SF, Zaman K, Portaels F, Rastogi N, Sola C: **Mycobacterium tuberculosis complex genetic diversity: mining the fourth international spoligotyping database (SpolDB4) for classification, population genetics and epidemiology.** *Bmc Microbiology* 2006, **6**.
18. van Embden JDA, van Gorkom T, Kremer K, Jansen R, van der Zeijst BAM, Schouls LM: **Genetic variation and evolutionary origin of the direct repeat locus of Mycobacterium tuberculosis complex bacteria.** *Journal of Bacteriology* 2000, **182**:2393-2401.
19. Smith NH, Gordon SV, de la Rúa-Domenech R, Clifton-Hadley RS, Hewinson RG: **Bottlenecks and broomsticks: the molecular evolution of Mycobacterium bovis.** *Nature Reviews Microbiology* 2006, **4**:670-681.
20. Streicher EM, Victor TC, van der SG, Sola C, Rastogi N, van Helden PD, Warren RM: **Spoligotype signatures in the Mycobacterium tuberculosis complex.** *J Clin Microbiol* 2007, **45**:237-240.
21. Vitol I, Driscoll J, Kreiswirth B, Kurepina N, Bennett KP: **Identifying Mycobacterium tuberculosis complex strain families using spoligotypes.** *Infect Genet Evol* 2006, **6**:491-504.
22. Rahim Z, Zaman K, van der Zanden AG, Mollers MJ, van SD, Raqib R, Zaman K, Begum V, Rigouts L, Portaels F, Rastogi N, Sola C: **Assessment of population structure and major circulating phylogeographical clades of Mycobacterium tuberculosis complex in Bangladesh suggests a high prevalence of a specific subclade of ancient M. tuberculosis genotypes.** *J Clin Microbiol* 2007, **45**:3791-3794.
23. Hilty M, Diguimbaye C, Schelling E, Baggi F, Tanner M, Zinsstag J: **Evaluation of the discriminatory power of variable number tandem repeat (VNTR) typing of Mycobacterium bovis strains.** *Vet Microbiol* 2005, **109**:217-222.
24. Frothingham R, Meeker-O'Connell WA: **Genetic diversity in the Mycobacterium tuberculosis complex based on variable numbers of tandem DNA repeats.** *Microbiology* 1998, **144**(Pt 5):1189-1196.
25. Asseged B, Woldesenbet Z, Yimer E, Lemma E: **Evaluation of abattoir inspection for the diagnosis of Mycobacterium bovis infection in cattle at Addis Ababa abattoir.** *Trop Anim Health Prod* 2004, **36**:537-546.
26. Teklul A, Asseged B, Yimer E, Gebeyehu M, Woldesenbet Z: **Tuberculous lesions not detected by routine abattoir inspection: the experience of the Hossana municipal abattoir, southern Ethiopia.** *Rev Sci Tech* 2004, **23**:957-964.
27. Ameni G, Aseffa A, Engers H, Young D, Hewinson G, Vordermeier M: **Cattle husbandry in Ethiopia is a predominant factor affecting the pathology of bovine tuberculosis and gamma interferon responses to mycobacterial antigens.** *Clin Vaccine Immunol* 2006, **13**:1030-1036.
28. Diguimbaye-Djaibe C, Vincent V, Schelling E, Hilty M, Ngandolo R, Mahamat HH, Pfyffer G, Baggi F, Tanner M, Zinsstag J: **Species identification of non-tuberculous mycobacteria from humans and cattle of Chad.** *Schweiz Arch Tierheilkd* 2006, **148**:251-256.
29. Oloya J, Kazwala R, Lund A, Opuda-Asibo J, Demelash B, Skjerve E, Johansen TB, Djonje B: **Characterisation of mycobacteria isolated from slaughter cattle in pastoral regions of Uganda.** *BMC Microbiol* 2007, **7**:95.
30. Wiese M: **Health-vulnerability in a complex crisis situation.** Edited by: H.-G.Bohle and T.Krings. Saarbrücken, Verlag für Entwicklungspolitik; 2004:1-436.
31. Aranaz A, Liebana E, Mateos A, Dominguez L, Vidal D, Domingo M, Gonzolez O, Rodriguez-Ferri EF, Bunschoten AE, van Embden JD, Cousins D: **Spacer oligonucleotide typing of Mycobacterium bovis strains from cattle and other animals: a tool for studying epidemiology of tuberculosis.** *J Clin Microbiol* 1996, **34**:2734-2740.
32. Zanella G, Durand B, Hars J, Moutou F, Garin-Bastuji B, Duvauchelle A, Ferme M, Karoui C, Boschiroli ML: **Mycobacterium bovis in wildlife in France.** *J Wildl Dis* 2008, **44**:99-108.
33. Michel AL, Hlokwé TM, Coetzee ML, Mare L, Connaway L, Rutten VP, Kremer K: **High Mycobacterium bovis genetic diversity in a low prevalence setting.** *Vet Microbiol* 2008, **126**:151-159.
34. Brosch R, Gordon SV, Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K, Garnier T, Gutierrez C, Hewinson G, Kremer K, Parsons LM, Pym AS, Samper S, van Soolingen D, Cole ST: **A new evolutionary scenario for the Mycobacterium tuberculosis complex.** *Proc Natl Acad Sci U S A* 2002, **99**:3684-3689.
35. Zucol F, Ammann RA, Berger C, Aebi C, Altwegg M, Niggli FK, Nadal D: **Real-time quantitative broad-range PCR assay for detection of the 16S rRNA gene followed by sequencing for species identification.** *J Clin Microbiol* 2006, **44**:2750-2759.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

