

MOLECULAR CONSTANTS OF THE $2^1\Pi$ STATE OF NaLi MOLECULE

NGUYEN TIEN DUNG, LE CANH TRUNG, DINH XUAN KHOA,
AND NGUYEN HUY BANG

Vinh University, 182 Le Duan Street, Vinh City, Vietnam

E-mail: bangvinhuni@gmail.com

Received 06 March 2013; revised manuscript received 25 April 2013

Accepted for publication 08 May 2013

Abstract. *We report our determination of the molecular constants for the $2^1\Pi$ state of NaLi from a set of 732 spectral lines. Using the linear least-squares fitting method, an optimum set of molecular constants, which corresponds to dimensionless root mean of squares of deviation $\sigma = 0.62$ within 0.1 cm^{-1} experimental uncertainty, have been derived.*

I. INTRODUCTION

The alkali-metal diatomic molecules with their main absorption bands in the UV-VIS region are very appropriated for modern laser spectroscopy techniques. Experimental investigations for these molecules have further promoted since the emergence of molecular Bose-Einstein condensates (see [1] and references therein). As several experimental techniques related to cold molecules involve optical excitation either in the formation and/or detection stages, a precise spectroscopic characterization of excited molecular states is necessary.

Among alkali-metal diatomic molecules NaLi is particularly interested because it is the lightest hetero-nuclear type and has permanent dipole moment, thus can be manipulated with external electric fields. Several experimental investigations for the $1^1\Sigma^+$ and $1^1\Pi$ symmetric types were performed to study up to the $9^1\Sigma^+$ and $7^1\Pi$ excited states [2-7]. Among those, the $2^1\Pi$ state was still known only at vibrational resolution [4].

In this paper we determine the molecular constants for the $2^1\Pi$ state of NaLi based our recent experimental data [8].

II. MOLECULAR CONSTANTS OF THE $2^1\Pi$ STATE

The experimental data used in this work consist of 732 spectral lines in the $2^1\Pi \leftarrow 1^1\Sigma^+$ band. The data were obtained by the polarization labeling spectroscopy technique [8] within to 0.1 cm^{-1} experimental uncertainty. Distribution of the data field corresponding




Fig. 1. Distribution of data field with the corresponding vibrational and rotational quantum numbers in the $2^1\Pi$ state.

to rotational and vibrational quantum numbers in the $2^1\Pi$ state is presented in detail in Fig. 1.

Once spectral lines had been assigned quantum numbers, their wave number $\bar{\nu}$ were fitted to the following relation:

$$\bar{\nu} = T'(v', J') - T''(v'', J'') + \delta. \quad (1)$$

In the Eq. (1), T' and T'' represent term values of the upper and lower states, respectively; δ describes the Λ -doubling in the $2^1\Pi$ state, given by

$$\delta = q_e [J(J+1) - 1], \quad (2)$$

where, q_e represents lambda-doubling coefficient for e -parity levels which gives rise to P and Q transitions. The spectroscopic term value $T(v, J)$ is represented with the conventional Dunham expansion:

$$T(v, J) = T_e + \sum_k \sum_l Y_{kl} (v + \frac{1}{2})^k [J(J+1) - 1]^l, \quad (3)$$

where T_e and Y_{kl} stand for electronic energy and Dunham coefficient, respectively. The term values for the labelled levels were calculated by using Dunham coefficients obtained from [2] and fixed during the fitting procedure. The final optimum set of Dunham coefficients for the $2^1\Pi$ state corresponding to a dimensionless root mean of squares of deviation $\sigma = 0.62$ has been obtained, and listed in Table 2.

Table 1. Molecular constants of the $2^1\Pi$ state of NaLi

Constants	Value [cm^{-1}]	Error [%]
Y_{00} (T_e)	22296.72539	0.001
Y_{10} (ω_e)	151.3553669	0.061
Y_{20} ($-\omega_e x_e$)	-1.212648477	2.097
Y_{30}	-0.1163013955	2.874
Y_{40}	0.5415371×10^2	3.810
Y_{50}	$-0.2242726086 \times 10^3$	2.138
Y_{01} (B_e)	0.2255917721	0.064
Y_{11}	$-0.2459652022 \times 10^2$	2.44
Y_{21}	0.8435599232×10^4	9.029
Y_{31}	-0.204878997×10^4	1.444
Y_{02}	-0.178619694×10^5	2.102
Y_{12}	-0.653086282×10^7	16.828
Y_{22}	$-0.1834230918 \times 10^7$	4.254
q_e	$-0.9339851354 \times 10^4$	5.715
D_e	17620	
σ	0.62	

Since the $2^1\Pi$ state associates to Na(3^2P)-Li(2^2S) atomic asymptotes, its dissociation energy can be determined by

$$D_e[2^1\Pi] = \Delta + D_e[1^1\Sigma^+] - T_e[2^1\Pi], \quad (4)$$

where, Δ is energy separation between the Na($3P$) and Na($3S$) atomic states, $D_e[1^1\Sigma^+]$ is dissociation energy of the ground electronic state $1^1\Sigma^+$. Taking electronic energy $T_e[2^1\Pi]$ from this work; $\Delta = 16965 \text{ cm}^{-1}$ from [9]; and $D_e[1^1\Sigma^+] = 7105 \text{ cm}^{-1}$ from [2]; the dissociation energy of the $2^1\Pi$ state is estimated to be $D_e = 1760 \text{ cm}^{-1}$.

From the experimental data and the dissociation energy determined above, the highest vibrational level observed here is only 17 cm^{-1} below the dissociation limit, thus the experimental data cover about 99% depth of the potential well.

Having the molecular constants, we calculated bond length R_e between the Na and Li atoms based on the following conventional relation:

$$R_e = \sqrt{\frac{\hbar}{B_e 4\pi\mu c}}, \quad (5)$$

where μ is reduced mass of NaLi, c is the light speed in vacuum. Taking the value of B_e in Table 2, the bond length is determined to be $R_e = 3.728438 \text{ \AA}$.

III. CONCLUSIONS

Spectroscopic structure of the $2^1\Pi$ state of NaLi has been characterized to rotational resolution for the first time. A set of molecular constants was determined based on the experimental data up to near the dissociation limit. The results give comprehensive spectroscopic characterization of the state and provide a benchmark to check reliability of modern theoretical calculations.

ACKNOWLEDGMENT

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 103.06.110.09. Also the supports from Ministry of Education and Training under the project code 08/2012/HĐ-HTQTSP and B2012-27-03 are acknowledged.

We would like to thank Prof. W. Jastrzebski for his supports to perform experiments in his lab.

REFERENCES

- [1] S. Jochim *et al.*, *Science* **302** (2003) 2101.
- [2] C.E. Fellows, *J. Chem. Phys.*, **94** (1991) 5855; *J. Mol. Spectrosc.*, **136** (1989) 369.
- [3] C.E. Fellows, J.Vergès, and C. Amiot, *J. Chem. Phys.* **93** (1990) 6281-6290.
- [4] M. M. Kappe, K. O. Marti, P. Radi, M. Schöp, and E. Scgumacher, *Chem. Phys. Lett.* **107** (1984) 6.
- [5] W. Jastrzebski, P. Kowalczyk, R. Nadyak, and A. Pashov, *Spectrochim. Acta, Part A*, **58A** (2002) 2193.
- [6] I. D. Petsalakis *et al.*, *Chem. Phys.*, **362** (1999) 130.
- [7] Nguyen Huy Bang *et al.* *J. Mol. Spectr.* **233** (2005) 290; *Optica. Applicata.* **36** (4) (2006) 499; *Chem. Phys. Lett.* **440** (2007) 199; *Opt. Mat.* **31**(2009) 527; *J. Chem. Phys.* **130** (2009) 124307.
- [8] Nguyen Tien Dung *et al.*, *Comm. Phys.* **21** (2011) 359.
- [9] A. A. Radzig and P. M. Smirnov, *Reference Data on Atoms, Molecules and Ions*, Springer, Berlin, 1985.