

Thermally driven escape over a barrier of arbitrary shape

Cite as: J. Chem. Phys. **110**, 2159 (1999); <https://doi.org/10.1063/1.477826>

Submitted: 13 March 1998 . Accepted: 21 October 1998 . Published Online: 12 January 1999

A. N. Drozdov, and J. J. Brey



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

[Theory of activated rate processes: Exact solution of the Kramers problem](#)

The Journal of Chemical Physics **85**, 1018 (1986); <https://doi.org/10.1063/1.451844>

[Two novel approaches to the Kramers rate problem in the spatial diffusion regime](#)

The Journal of Chemical Physics **111**, 6481 (1999); <https://doi.org/10.1063/1.479945>

[Activated rate processes: Generalization of the Kramers-Grote-Hynes and Langer theories](#)

The Journal of Chemical Physics **97**, 2422 (1992); <https://doi.org/10.1063/1.463081>

Lock-in Amplifiers
up to 600 MHz



Thermally driven escape over a barrier of arbitrary shape

A. N. Drozdov^{a)} and J. J. Brey

Física Teórica, Universidad de Sevilla, Apartado de Correos 1065, Sevilla 41080, Spain

(Received 13 March 1998; accepted 21 October 1998)

The Kramers theory for the thermally activated rate of escape of a Brownian particle from a potential well is extended to a barrier of arbitrary shape. The extension is based on an approximate solution of the underlying Fokker–Planck equation in the spatial diffusion regime. With the use of the Mel'nikov–Meshkov result for the underdamped Brownian motion an overall rate expression is constructed, which interpolates the correct limiting behavior for both weak and strong friction. It generalizes in a natural way various different rate expressions that are already available in the literature for parabolic, cusped, and quartic barriers. Applications to symmetric parabolic and cusped double-well potentials show good agreement between the theory and estimates of the rates from numerical calculations. © 1999 American Institute of Physics. [S0021-9606(99)01404-X]

I. INTRODUCTION

Ever since the pioneering contribution of Svante Arrhenius, the problem of thermally driven escape from a metastable state has become one of the most fundamental problems in physics and chemistry.¹ The modern theory of activated rate processes is essentially due to Kramers,² who provided a dynamical framework for the original concepts of Arrhenius. The underlying idea of the Kramers theory is to model the escape process by the motion of a Brownian particle with mass weighted coordinate x in a potential of mean force $V(x)$. The dynamics is governed by the following Fokker–Planck equation for the probability density $P(x, v, t)$ of finding the particle at time t at position x with velocity v :²

$$\partial_t P(x, v, t) = [-v \partial_x + V'(x) \partial_v + \gamma \partial_v (v + \beta^{-1} \partial_v)] P(x, v, t). \quad (1.1)$$

Here the prime denotes the derivative with respect to x , γ is the friction coefficient, and β the inverse energy available from the thermal bath, $\beta^{-1} = k_B T$. The potential is assumed to have a well with minimum at $x_w < 0$, separated from the continuum by a barrier at $x = 0$ of height $E = -V(x_w)$. Hereby we set for convenience $V(0) = 0$. The quantity of interest is the escape rate Γ of the particle from the well. The latter can always be written in the form

$$\Gamma = \mu \Gamma_{\text{TST}}, \quad (1.2)$$

where Γ_{TST} is the transition state theory (TST) result

$$\Gamma_{\text{TST}} = \left\{ \sqrt{2\pi\beta} \int_{-\infty}^0 dx e^{-\beta V(x)} \right\}^{-1}, \quad (1.3)$$

and μ is a transmission coefficient describing the deviation of the rate from Γ_{TST} .

Kramers studied the dependence of the escape rate on the frictional damping in two regimes, namely, for small and intermediate to large friction γ . In the former regime, the coupling between the system and the bath is assumed to be

vanishingly weak so that the rate limiting step is the transfer of energy from the bath to the particle. The transmission coefficient takes in this case the form

$$\mu(\gamma \rightarrow 0) = \Delta(\gamma \rightarrow 0) = 2\gamma\beta \int_{x_p}^0 dx \sqrt{-2V(x)}, \quad (1.4)$$

where Δ is the dimensionless loss of energy per oscillation of a particle with energy close to the barrier height, and x_p the left-hand side turning point of the asymptotic underdamped trajectory, $V(x_p) = 0$. In the intermediate to large friction regime, when the transfer of energy becomes fast enough to maintain thermal equilibrium of escaping particles, the rate limiting step is spatial diffusion across the barrier region. One of the basic assumptions of the Kramers theory² in this regime is a parabolic barrier approximation. It consists in dividing the full potential into a *parabolic* barrier part

$$U(x) = -\frac{1}{2}\omega^2 x^2, \quad (1.5)$$

with $\omega^2 = -V''(0)$, and an anharmonic correction reading

$$V(x) = U(x) + O(x^3). \quad (1.6)$$

In the immediate vicinity of the barrier top which dominates the dynamics, the nonlinearity of $V(x)$ vanishes faster than the parabolic part $-\frac{1}{2}\omega^2 x^2$ and therefore can be neglected. This yields the following expression for the transmission coefficient:

$$\mu_{\text{pb}} = \sqrt{1 + \frac{\gamma^2}{4\omega^2} - \frac{\gamma}{2\omega}}. \quad (1.7)$$

It should be noted that Eq. (1.7) is valid for $\gamma \geq \omega/(2\pi\beta E)$. Consequently, in the extreme high barrier (low temperature) limit, $\beta E \rightarrow \infty$, one will ultimately almost always be in the spatial diffusion regime.

Kramers' model, although simple, is of wide-ranging significance to a detailed understanding and evaluating the influence of the medium on reaction rates. It has found various generalizations to the full friction range,³ non-Markovian activated rate processes,^{4,5} multidimensional systems,⁶ and cases without detailed balance⁷ (for a review see Ref. 1). In

^{a)}Permanent address: Institute for High Temperatures, 13/19 Izhorskaya Street, 127412 Moscow, Russia.

all these investigations the barrier is assumed to be parabolic, though this assumption is not always met in real physical and chemical barrier crossing processes. For example, the barrier of charge transfer reactions is often of a cusp-shaped form.⁸ Kramers also derived the transmission coefficient for a cusp-shaped barrier,² $U(x) = -a|x|$,

$$\mu_{\text{cusp}}(\gamma \rightarrow \infty) = \frac{a}{\gamma} \sqrt{\frac{1}{2} \pi \beta}, \quad (1.8)$$

but his expression is valid only in the asymptotic limit of large friction where Eq. (1.1) can be approximated by a Smoluchowski equation. There are various attempts in the literature to bridge the strong friction limit result for a cusp-shaped barrier with the TST value, $\mu_{\text{TST}}=1$, at zero damping.^{9,10} An analogous interpolating formula is known for a quartic barrier.¹ Only very recently, Berezhkovskii *et al.*¹¹ have extended this formula to an arbitrary nonparabolic barrier of the form

$$U(x) = -(a/\alpha)|x|^\alpha. \quad (1.9)$$

Their generalization reads¹¹

$$\mu_\alpha = \left\{ \int_{-\infty}^{\infty} dx \exp[\beta U(x)] \right\}^{-1} \times \int_{-\infty}^{\infty} dx \exp\left\{ \beta \left[U(x) - \frac{1}{2} \gamma^2 x^2 \right] \right\}. \quad (1.10)$$

There is, however, a certain irony here; the above formula agrees with the known escape rates for nonparabolic barriers, but fails to reproduce the exact result for a parabolic barrier. In the latter case, it yields instead of Eq. (1.7) an approximate expression

$$\mu_2 = (1 + \gamma^2/\omega^2)^{-1/2}. \quad (1.11)$$

The aim of this paper is twofold. First, we want to present an approximate rate formula, which indeed is valid for arbitrarily shaped barriers and interpolates between the limits of small and large friction. And second, we wish to compare this formula with exact numerical rates in different types of potentials.

II. INTERPOLATING FORMULA

To begin with we consider the spatial diffusion regime. Our purpose is to derive an approximate solution of the Fokker–Planck equation which would allow one to recover Eqs. (1.7) and (1.10). This goal can be achieved in many different ways.¹² Here we employ the flux over population method developed by Kramers.² Within its scope, the escape rate is defined as the ratio of a stationary diffusion current at the top of the barrier to the population of the well. Accordingly, we have to look for a current carrying stationary probability density $P(x, v)$, that smoothly matches the equilibrium distribution

$$P_{\text{eq}}(x, v) = \exp[-\beta V(x) - \frac{1}{2}\beta v^2] \quad (2.1)$$

in the well and vanishes beyond the barrier. The two stationary densities are related by a form function $\xi(x, v)$,

$$P(x, v) = \xi(x, v) P_{\text{eq}}(x, v), \quad (2.2)$$

which is determined from

$$\{-v \partial_x + [V'(x) - \gamma v] \partial_v + \gamma \beta^{-1} \partial_{vv}^2\} \xi(x, v) = 0. \quad (2.3)$$

Once the form function is known, the reactive flux formula yields for the transmission coefficient

$$\mu = \beta \int_{-\infty}^{\infty} dv v \xi(0, v) \exp\left(-\frac{1}{2} \beta v^2\right). \quad (2.4)$$

Following Kramers, we approximate the potential $V(x)$ entering Eq. (2.3) by its barrier part $U(x)$. The latter is not necessarily parabolic, it may be a sum of arbitrary (parabolic and nonparabolic) terms

$$U(x) = -\frac{1}{2} \omega^2 x^2 - \frac{a}{\alpha} |x|^\alpha - \dots. \quad (2.5)$$

Moreover, we assume that $\xi(x, v)$ is a function of some linear combination of x and v ,

$$\xi(x, v) = \xi(\varrho), \quad \varrho = cx + bv. \quad (2.6)$$

Then, it is not difficult to check by direct substitution that in leading order in ϱ and $(\beta E)^{-1}$ an approximate solution to Eq. (2.3) reads

$$\xi(x, v) = Z^{-1} \int_{\varrho}^{\infty} dy e^{\beta U(y)}, \quad (2.7)$$

with

$$\varrho = \sqrt{\omega/(\gamma \mu_{\text{pb}})} [x - (\mu_{\text{pb}}/\omega)v]. \quad (2.8)$$

In the above μ_{pb} is given by Eq. (1.7), while the normalization constant Z is defined by the requirement that the form function $\xi(x, v)$ approaches unity in the initial well and zero in the product side. This immediately yields

$$Z = \int_{-\infty}^{\infty} dy e^{\beta U(y)}. \quad (2.9)$$

It will be recalled here that the barrier (temperature) is assumed to be high (low) enough so that the potential can be well approximated by its local behavior in the vicinity of the barrier top. Otherwise one can use in Eqs. (2.7) and (2.9) instead of the barrier part $U(x)$ the full potential $V(x)$ itself. In such a case, the integration has to be restricted to the barrier region with a lower limit at, say, x_w and the upper limit at a value beyond the barrier from where the recrossing probability of a particle with zero initial velocity can safely be neglected.

Inserting Eq. (2.7) into Eq. (2.4), we obtain the following expression for the transmission coefficient:

$$\mu_{\text{ab}} = Z^{-1} \int_{-\infty}^{\infty} dx \exp\left\{ \beta \left[U(x) - \frac{1}{2} (\gamma \omega / \mu_{\text{pb}}) x^2 \right] \right\}. \quad (2.10)$$

It is a simple matter to check that for a parabolic barrier the above formula coincides with the exact Kramers result, Eq. (1.7), while for a purely nonparabolic barrier ($\omega=0$) it reproduces Eq. (1.10). One may also note that it agrees in the limiting case of high friction with the transmission factor for an arbitrarily shaped barrier following from the corresponding Smoluchowski equation¹³

$$\mu(\gamma \rightarrow \infty) = \left\{ \gamma \sqrt{\frac{\beta}{2\pi}} \int_{-\infty}^{\infty} dx e^{\beta U(x)} \right\}^{-1}, \quad (2.11)$$

and reduces to unity at zero damping.

A rate expression valid in the full damping range can be obtained by making use of an elegant approach developed by Mel'nikov and Meshkov.³ This gives in a straightforward way

$$\mu = \mu_{ab} A(\Delta), \quad (2.12)$$

with

$$A(\Delta) = \exp\left(\frac{1}{\pi} \int_0^{\infty} dx \frac{\ln\{1 - \exp[-\Delta(x^2 + \frac{1}{4})]\}}{x^2 + \frac{1}{4}}\right), \quad (2.13)$$

where Δ is given by Eq. (1.4). It should be noted that the ansatz of writing a uniform formula for nonparabolic barriers as a product of a spatial diffusion expression and the depopulation factor A is *ad hoc*. It follows neither from Mel'nikov and Meshkov nor from Pollak, Grabert, and Hänggi turnover theories. It is our aim here to prove the utility of Eq. (2.12) by comparing with exact numerical rates. The latter is not so obvious as one might think. Specifically, Mel'nikov and Meshkov derived the depopulation factor (2.13) under the assumption that the escape dynamics can be described by a probabilistic integral equation in energy-action variables, whose Green function corresponds to the barrier trajectory. For a smooth potential the trajectory that leaves the barrier with the entire energy close to zero returns to it after time $T \rightarrow \infty$. This infinite time, however, is no longer true for a cusped barrier where the time is of the order of the period of particle oscillation in the well. Thus the interesting issue we shall address in our numerical applications is as follows: Does the finite period of the barrier trajectory spoil the applicability of Eq. (2.12)?

III. NUMERICAL RESULTS

The aim of this section is to present exact numerical rates for different types of potential barriers that would allow one to test analytical predictions. One might, at first, believe that this issue should have been settled long ago, mainly because of its continuous importance in many problems of chemical physics. To the best of our knowledge, however, there are no numerical solutions of such a type, other than those obtained in Refs. 10 and 11 under the assumption that the potential consists only of a barrier part. This assumption results in a *monotonic* dependence of the transmission coefficient on γ ; the coefficient increases with decreasing γ and reaches its maximal value at zero damping, when there is no coupling between the system and the bath. It is clear that the data so obtained are not suited for testing analytical predictions in the most problematic intermediate and weak damping regimes.

Here we deal with activated rate processes in a symmetric double-well potential of the form




FIG. 1. Different shapes of the potential $V(x)$, Eq. (3.1), for $a=0$ (the dashed line) and $a=1$ (the solid line).

$$V(x) = \frac{E}{1+2a} [x^4 - 4a|x| - 2(1-a)x^2], \quad a > -\frac{1}{2}. \quad (3.1)$$

Its barrier part varies with the parameter a from a purely parabolic ($a=0$) to a purely cusped ($a \geq 1$) barrier, see Fig. 1. Accordingly, the frequency ω entering our rate expression reads

$$\omega^2 = \begin{cases} 4(1-a)E/(1+2a) & -\frac{1}{2} < a \leq 1, \\ 0 & a > 1. \end{cases} \quad (3.2)$$

The method used to numerically solve Eq. (1.1) will be described elsewhere.^{14,15} Table I shows a list of the first non-zero eigenvalue in the considered potential for $\beta E = 10$ and $a = 0, 0.5$, and 1 . The calculation is performed over a large range of γ which covers all regimes of chemical interest, from the underdamped Brownian motion to the spatial diffusion regime.

Before testing the validity of the present rate expression, we note that Eq. (2.12) gives the transmission coefficient for the escape from a metastable state. Using the approach suggested by Mel'nikov and Meshkov,³ the coefficient for a symmetric double well can be written as

$$\mu = \mu_{ab} A^2(\Delta)/A(2\Delta). \quad (3.3)$$

TABLE I. First nonzero eigenvalue for symmetric double-well potentials, Eq. (3.1) with $\beta E = 10$ and $a = 0, 0.5$, and 1 . Exponential notation $-k$ means that the number preceding is to be multiplied by 10^{-k} .

γ	$a=0$	$a=0.5$	$a=1$
0.05	0.171-4	0.146-4	0.144-4
0.1	0.304-4	0.259-4	0.247-4
0.25	0.593-4	0.494-4	0.456-4
0.5	0.868-4	0.713-4	0.640-4
1	0.106-3	0.889-4	0.791-4
2	0.106-3	0.952-4	0.856-4
5	0.858-4	0.914-4	0.850-4
10	0.607-4	0.786-4	0.770-4
20	0.361-4	0.557-4	0.564-4
50	0.154-4	0.268-4	0.286-4
100	0.780-5	0.133-4	0.145-4
1000	0.783-6 ^a	0.134-5 ^a	0.147-5 ^a

^aExact estimate of the eigenvalue calculated from the respective Smoluchowski equation.




FIG. 2. Transmission coefficient and percentage error, $100 \times (\text{approximate} - \text{exact}) / \text{exact}$, made in μ by using Eq. (3.3). Exact numerical results are shown by circles. (a) $a=0$; (b) $a=0.5$; (c) $a=1$.

The least nonvanishing eigenvalue of the corresponding Fokker–Planck operator is then given by twice the rate defined by Eq. (1.2). The numerical values of the transmission coefficient extracted in this way are exhibited in Fig. 2, together with the analytical predictions obtained in terms of Eq. (3.3). As evidenced by Fig. 2, the approximate rate expression gives an upper bound to the exact result for the rate in the parabolic double-well potential. For the cusped potentials the theory overestimates the rate in both limits of weak and strong friction and underestimates it in the intermediate friction region. It is also seen that for all values of a the best agreement is achieved in the strong damping limit ($\gamma \gtrsim 100$). With decreasing γ the error made by the ansatz (3.3) increases and reaches maximal values in the weak damping region ($\gamma \lesssim 0.1$). The theoretical expression overestimates the rate in this region by 14% for a parabolic barrier ($a=0$) and by 18% for a purely cusped barrier. It should be pointed out that the same is true for the turnover theory of Pollak, Grabert, and Hänggi.⁵ As we have shown in recent papers,^{15,18} their theory also considerably overestimates the rate in the weak friction regime.

Finally, to conclude this section we note that the barrier frequency ω appearing in Eq. (2.10) may still be left even if the barrier is purely nonparabolic. In such a case, it should be treated as a variational parameter.¹⁶ Yet another way to improve the rate formula is to take into account finite-barrier corrections. These are obtainable systematically in both re-

gimes of weak¹⁷ and intermediate to strong friction.¹² A further improvement of the overall rate expression can be achieved by using in Eq. (2.10) a properly determined energy loss of the deterministic particle dynamics. In contrast to the weak friction expression for Δ proposed by Mel'nikov and Meshkov, Eq. (1.4),³ as well that suggested by Pollak, Grabert, and Hänggi⁵ in their turnover theory, the deterministic approach to this quantity yields an approximation which remains correct in the full damping range, regardless of the particular shape of the potential barrier.^{15,18}

IV. CONCLUDING REMARKS

In this paper, an approximate formula for the rate of escape over an arbitrarily shaped barrier has been constructed by means of the flux over population method and the approach by Mel'nikov and Meshkov. The resulting expression agrees in the limiting case of high friction with the rate following from the corresponding Smoluchowski equation and, in the extremely underdamped regime with the rate obtained by Kramers from a diffusion equation in energy (action) variables. It generalizes in a natural way the known rate formulas for parabolic and nonparabolic barriers.

Besides, we have presented for the first time *numerically exact* rate constants for potentials with different barrier shapes in all regimes of chemical interest, from underdamped to overdamped Brownian motion. These results pro-

vides the necessary foundation for testing various different rate expressions that already exist in the literature. Comparison with the numerical data shows that the present overall rate expression is rather accurate in the strong damping limit, underestimates the rate by $\sim 0\%$ – 18% in the intermediate friction region and overestimates the rate by $\sim 14\%$ – 23% in the weak damping regime.

ACKNOWLEDGMENTS

One of us (A.N.D.) is grateful to A. M. Berezhkovskii, P. Talkner, and V. Yu. Zitserman for many helpful discussions. We acknowledge the support of the Dirección General de Investigación Científica y Técnica of Spain for financial support (A.N.D.) and for Project No. PB95-534 (J.J.B.).

¹P. Hänggi, P. Talkner, and M. Borkovec, *Rev. Mod. Phys.* **62**, 251 (1990).

²H. Kramers, *Physica (Utrecht)* **7**, 284 (1940).

³V. I. Mel'nikov and S. V. Meshkov, *J. Chem. Phys.* **85**, 1018 (1986).

⁴R. F. Grote and J. T. Hynes, *J. Chem. Phys.* **73**, 2715 (1980); P. Hänggi and F. Mojtabai, *Phys. Rev. A* **26**, 1168 (1982).

⁵E. Pollak, H. Grabert, and P. Hänggi, *J. Chem. Phys.* **91**, 4073 (1989).

⁶H. C. Brinkman, *Physica (Utrecht)* **22**, 149 (1956); R. Landauer and J. A. Swanson, *Phys. Rev.* **121**, 1668 (1961); J. S. Langer, *Ann. Phys. (N.Y.)* **54**, 258 (1969).

⁷P. Talkner, *Z. Phys. B* **68**, 201 (1987); A. N. Drozdov, *Physica A* **187**, 329 (1992).

⁸R. A. Marcus, *Annu. Rev. Phys. Chem.* **15**, 155 (1963); L. D. Zusman, *Chem. Phys.* **49**, 295 (1980).

⁹B. J. Matkowsky and Z. Schuss, *SIAM (Soc. Ind. Appl. Math.) J. Appl. Math.* **33**, 365 (1977); D. F. Calef and P. G. Wolynes, *J. Phys. Chem.* **87**, 3387 (1983); H. Dekker, *Physica A* **136**, 124 (1986); E. Pollak, *J. Chem. Phys.* **93**, 1116 (1990).

¹⁰A. Starobinets, I. Rips, and E. Pollak, *J. Chem. Phys.* **104**, 6547 (1996).

¹¹A. M. Berezhkovskii, P. Talkner, J. Emmerich, and V. Yu. Zitserman, *J. Chem. Phys.* **105**, 10890 (1996).

¹²P. Talkner, *Chem. Phys.* **180**, 199 (1994).

¹³H. Risken, *The Fokker-Planck Equation, Methods of Solution and Applications* (Springer, New York, 1989).

¹⁴A. N. Drozdov and J. J. Brey, *Phys. Rev. E* **57**, 1284 (1998).

¹⁵A. N. Drozdov and P. Talkner, *J. Chem. Phys.* **109**, 2080 (1998).

¹⁶P. Talkner and E. Pollak, *Phys. Rev. E* **50**, 2646 (1994).

¹⁷V. I. Mel'nikov, *Phys. Rev. E* **48**, 3271 (1993).

¹⁸A. N. Drozdov and J. J. Brey, *Chem. Phys.* **235**, 147 (1998).