

## MOLECULAR GENETICS - STEP BY STEP IMPLEMENTATION IN MAIZE BREEDING

Kosana KONSTANTINOV and Snežana MLADENOVIĆ DRINIĆ

Maize Research Institute „Zemun Polje“, Belgrade

Konstantinov K. and S. Mladenović Drinić (2007): *Molecular genetics - step by step implementation in maize breeding*. – Genetika, Vol. 39, No. 2, 139-154.

Efficiency in plant breeding is determined primarily by the ability to screen for genetic polymorphism, productivity and yield stability early in program. Dependent on the knowledge about the biochemical bases of the trait and nature of its genetic control, trait could be modified either through mutagenesis of genes controlling it or through the transfer of already existing mutant genes, controlling desired trait to different plant genotypes by classic crossing. Objective of this report is to present partly results on the investigation of the possibilities to apply ionizing radiations (fast neutrons,  $\gamma$  - rays) and chemical mutagens (EI, iPMS, EMS, ENU) to get maize and wheat mutants with increased amount and improved protein quality. Besides this approach in mutation breeding, results on the very early investigation of biochemical background of *opaque -2* mutation including use of coupled cell – free RNA and protein synthesis containing components from both wild and *opaque - 2* maize genotypes (chromatin, RNA polymerase, microsomal fraction, protein bodies) will be presented. Partial results on *opaque - 2* gene incorporation in different genetic background are reviewed.

Part of report is dealing with different classes of molecular markers (proteins, RFLP, AFLP, RAPD, and SSR) application in maize genome polymorphism investigation. Besides application of different molecular markers classes in the investigation of heterosis phenomena they are useful in biochemical pathway of important traits control determination as well.

*Key words:* maize, mutagens, chromatin, proteins, molecular markers, heterosis

## INTRODUCTION

After biological sciences development, particularly molecular biology, during last decades molecular biotechnology, based on genetic engineering are widely used in genome structure and expression investigation. As consequence of the some results application, new varieties of commercially grown crops with higher yield are developed, as well as decreasing time to breed new genotypes having satisfactory resistance to different pathogens, tolerance to biotic and abiotic stress conditions etc. In spite of the fact that both information technology and molecular biotechnology are „strategic“ technologies, it is out of the question that remarkable progress in maize genetics and breeding was gained during last few decades by the use of conventional genetics and breeding methods. Although modern maize breeding is based on the concepts of SCHULL (1910) methods of breeding and evaluation have changed due to utilisation of new or improved germplasm sources, more knowledge and information on the inheritance pathway of complex traits as well as information from theoretical and computer simulation studies.

Existing or induced variation that will enable the selection of superior genotypes is at the core of a successful breeding program. KUCKUK et al. (1991) classified breeding methods into three groups: (i) selection breeding, where breeder relies on existing variation in natural populations and genotype mixtures; (ii) combination breeding where combinatorial crosses are made and  $F_1$  is not used directly, but only to generate subsequent segregation material that will form the basis for selection and finally (iii) hybrid breeding where combinatorial crosses are made to create new genotypes that will be used as  $F_1$  seed. Crucial items to have effective breeding is the prediction of the best hybrid within a large set of possible genotype combinations based on genetic distance of the parental lines. According to BOPPENMAIER et al. (1992) this was not possible for line combinations only between heterotic pools of European material. It is important to survey and choose germplasm as pure lines, cultivars, populations, clones, genes, DNA sequences etc (LEE,1995).

Development of a large number of molecular markers raises question whether only their application can further enhance the efficiency of maize breeding. It is important to collect more information on the reliable level of polymorphism in order to understand biological processes, including both genetic

control of storage protein and oil biosynthesis as well as other important traits in different genome combinations.

This review is covering partly our many years period results on the biochemical background investigation of genetic control of storage protein biosynthesis in maize endosperm, induced and existing genetic polymorphism determination by biochemical and other molecular markers application with general aim to find out what kind of genome polymorphism - at the level of genome structure or expression is reliable criteria for successful breeding program creation.

## 1. MUTATION BREEDING

1.1 Important advantage of mutations is that it may be introduced into the best commercial varieties with acceptable fulfil the demands on yield, straw-stiffness, disease resistance, winter hardness or other critical properties, to create a new variation from which selection can take place in any property whose improvement is desired. In principle, mutation breeding with these purposes may go one of two possible ways: 1- *induced mutation* and 2 - *biochemical plant breeding with special reference to existing mutants*

### 1.2 *Induced mutations*

It is known that different genes may vary strongly with respect to their relative sensitivity to different mutagenic agents (HAGBERG *et al.*, 1958) and theoretically it is possible to develop more or less specific mutagenic agents, that preferably provoke a change or variation in the gene or genes controlling the property to be improved, undesirable changes being suppressed. The use of unspecific mutagenic agents, like radiations and alkylating agents, will lead to a random variation in all kinds of properties. The success of such breeding is above all dependent on the availability of efficient screening methods and of appropriate design of experiments. Poliploidy character of hexaploid wheat makes it as promising source of good biological background for induced mutation either by ionizing radiation and chemical mutagens (EHRENBERG *et al.* 1959., MAC KEY, 1959., BHASHARAN and SWAMINATHAN, 1960., MOUTSCHEN - DAHMEN and EHRENBERG, 1968., ĐOKIĆ *et al.* 1967., BOROJEVIĆ and BOROJEVIĆ, 1969., KONSTANTINOV, 1968., KONSTANTINOV *et al.* 1970a., KONSTANTINOV and MOUTSCHEN, 1970b)

In our program on induced mutations several mutants of hexaploid wheat, with changed protein content and quality were obtained by ionizing irradiation, gama rays or fast neutrons and treated with alkylating agents, ethylenimine (EI) and ethylmethane sulphonate (EMS) (DUMANOVIĆ *et al.* 1969). All applied mutagens induced chromosomal aberrations in the first mitotic cycle of meiosis (fig 1.). There was no detectable effect of EI, EMS and gama rays on the chromosomes in M5 progeny of chemically treated or irradiated seeds. Fast neutrons induced chromosomal aberrations in M5 progeny which is in agreement with the mutagenic

behaviour of this kind of irradiation. Also there is no evidence for a direct correlation between the changed content of protein in the seeds and the induced chromosomal aberrations (JOVANOVIĆ and KONSTANTINOV, 1973).

Therefore second procedure, i.e. use of existing known mutant genes controlling desired trait must be mostly relied in mutation breeding for improved biochemical characters.




Fig 1. a) EMS induced chromosomal aberration in hexaploid wheat variety

b) Chromosomal aberration in the first meiotic mitoses of maize induced by irradiation

### 1.3 Biochemical plant breeding with special reference to existing protein mutants

The first consideration in estimating the quality of any livestock fodder or human food is the content of protein and its composition. The great achievements beneficial to food production can be expected by an increase in the relative proportion of protein, with a corresponding decrease of carbohydrate – fat production. Since the cereal grains are the main source of protein for man and animals, the best way to increase the protein supplies is to increase the protein content in cereals and to modify the proteins, i.e. to increase the biological value of crop protein, which quality is primarily determined by the content of certain essential amino acids, first-hand lysine, tryptophane and methionine (DUMANOVIĆ and EHRENBERG, 1968).

The most striking advancement in these efforts came in 1964 when the high protein quality of the *opaque-2* mutant was described (MERTZ et al. 1964). At that time primary point of *opaque-2* mutation, having as genetic effect the changes in the proportion of protein fractions in maize endosperm, has been unidentified. Zeins represent 50% of maize endosperm protein and are strictly developmentally controlled, first detected 12 days after pollination and accumulates till maturity at 45-50 days after pollination with very large increase over next 20 days. Also higher ribonuclease (RNase) activity has been reported as characteristic of *opaque-2* endosperm (MERTZ et al. 1964). In our experiments we did verified that RNase activity could be used as biochemical marker for *opaque-2* gene presence in recessive homozygous triploid endosperm, because difference in the enzyme activity between wild and mutant endosperm follows the pattern of zein

accumulation during endosperm development (DENIĆ *et al.*, 1971., DENIĆ, *et al.*, 1973).

After *opaque-2* gene incorporation in different valuable commercial inbred lines by the 4 -5 backcrossing and 2 – 3 generation of selfpollination (DUMANOVIĆ *et al.* 1974) variation of endosperm texture has been obtained, varying from standard soft *o-2* to nearly normal dent types. Interaction of *o2* gene with host genome was quite different in different genetic background. Changes in protein content and composition of endosperm in different categories of „normalized“ endosperm and changes in RNase activity at different degrees of endosperm vitreosity have been investigated. As the rule, all converted *o2* versions kept in each genetic background protein quality (aminoacid composition of endosperm proteins) similar to original *o2* type. It was concluded that, in general, the amount of protein is slightly increased in „normalized“ endosperm compared to standard *o-2* soft floury endosperm. A tendency was indicated for RNase activity to be decreased with increasing proportion of modified („norm.“) endosperm.

It has been suggested that variation of endosperm texture is consequence of polygenic character of this trait (VASAL, 1972., POPOVIĆ *et al.* 1974). Nowadays, more probable explanation is presence of different transposon elements families in inbred lines used in the experiments for the conversion to *opaque-2* phenotype. Genetic instability at the *opaque -2* locus in maize was first reported for *o2-m( r)* (SALAMINI, 1980), an allele responding to the autonomous *Bg* element (HARTINGS *et al.* 1991). MOTTO *et al.* (1986, 1988) reported *Ds*-induced alleles phenotypically indistinguishable from wild – type in the absence of *Ac*, but mutable with *opaque* sectors when *Ac* is present.

Ribonuclease activity and pattern of modification of endosperm texture have been used as genetic markers of *opaque-2* versions of converted inbred lines. Several of them were later used as parental lines in the process of several *opaque-2* hybrid creation. Results on nutritive quality and energy yield of *opaque-2* maize hybrids created at Maize Research Institute, as well as high oil and waxy maize hybrids are published by EGGUM *et al.*(1985).

Among many approaches to analyse difference between wild and *opaque-2* gene expression (DENIĆ, 1970., DENIĆ, 1971) interesting example are results on difference of protein bodies, which contain mostly or exclusively zein in homogenous matrix (BURR and BURR, 1976, DENIĆ and MILIVOJEVIĆ, 1978), fig2.

In order to study the mode of gene action on storage protein synthesis, during early of seventeens attempts have been made to isolate chromatin from the endosperm of developing seeds (KONSTANTINOV, 1975). After modification of method adopted for isolation of crude chromatin from plant tissue (BONNER *et al.* 1968) we found that it was possible to isolate chromatin from maize endosperm tissue (KONSTANTINOV and DENIĆ, 1975). Obtained results conferred that chromatin, isolated by modified method is in native, transcriptionally active form (fig 3). To approve whether the genetic activity is preserved in isolated chromatin, a chromatin dependent protein synthesizing system was set up in the coupled cell-

free protein synthesizing system from wheat embryo (ALLENDE and BRAVO, 1966). From the obtained results it has been obvious that *de novo* proteins are synthesized in response to messenger - RNA synthesized on the chromatin as a template (KONSTANTINOV, 1975, 1978, POPOVIĆ et al. 1981., POPOVIĆ, 1986). It was established in the system with chromatin, isolated from an *opaque-2* mutant endosperm 25 days after pollination as template, relatively higher incorporation of lysine in comparison with the chromatin of wild genotype.




Fig.2 Protein granules from maize endosperm 26 days after pollination




Fig 3. Transcription product of endogenous RNA polymerase activity of chromatin: 30 $\mu$ g chromatin-DNA (60°) plus 1.5U of E.coli polymerase (curve 1); 30 $\mu$ g thymus DNA plus 1.5U of E.coli polymerase (curve 2); 30 $\mu$ g chromatin-DNA (0°) plus 1.5U of E.coli polymerase (curve 1)

Based on the results obtained after analysis at the molecular level, reviewed in Mutants of Maize (NEUFER *et al.* 1997) *opaque2* (*o2*) plays an important regulatory role in the expression of zein seed storage protein genes (AUKERMAN and SCHMIDT, 1994). This gene encodes a 48-kD protein belonging to the basic leucine zipper (bZIP) class of eukaryotic transcriptional activators (HARTINGS *et al.*, 1991., SCHMIDT *et al.*, 1990). The gene was cloned by SCHMIDT *et al.* (1987) and MOTTO *et al.* (1988) and sequenced by MADDALONI *et al.* (1989).

There are very short upstream open reading frames that reduce the efficiency of *o2* translation (LOHMER *et al.* 1993). Recessive, *o2* expression appears to be endosperm – specific (SCHMIDT *et al.* 1987., MOTTO *et al.*, 1988). The protein is localized in the nuclei of developing endosperm cells (VARAGONA *et al.* 1991) and found to contain two nuclear localization signals (VARAGONA *et al.* 1992). The O2 protein is a transcriptional activator that binds to the promoter of 22 – kD zein genes (SCHMIDT *et al.* 1990, 1992) and to the promoter of the *rip1* (LOHMER *et al.* 1991). In recent years (PIRONA *et al.*, 2007), cDNA microarray is in use to investigate the transcription profiles and different gene expression of maize endosperm from two different opaque mutants (*o2* and *o7*) and in double mutant combination.

After recombinant DNA technology and different technologies for plant transformation have been developed we did maize transformation by selective marker – *NPT II* gene integration in maize genome (MLADENović, 1990, KONSTANTINOV *et al.* 1997). It has been proved that genome of transformed plants contained the selective marker gene sequence in an active form. Besides foreign gene integration and expression in maize plant, foreign integrated DNA induced several heritable changes - mutations, changing host genome function (MLADENović *et al.* 1991., KONSTANTINOV *et al.* 1997).

It is important to stress that mutation breeding must be applied along with, and partly combined with hybridization – selection methods.

## 2. MOLECULAR BREEDING

Determination and measuring of genetic variation is the most important step in any breeding program and data obtained by the methods of molecular genetics could be best interpreted by correlation with the procedure already well developed in conventional genetics. Maize is extremely diverse genus, having morphological and biological differences. Despite the nearly unlimited diversity of germplasm, the main problem is the creation of suitable new crosses, arises from divergent parental lines. To find the best method, which provides discrimination according to the purpose of selection, will remain a challenge both for classical and molecular geneticists.

Information on the genetic diversity and relationships of lines or populations is useful for choice of parents, crossing, and classification of germplasm into heterotic groups, prediction of heterosis and plant variety protection. Maize breeders are mainly concerned with the genetic diversity among

and within breeding population and elite germplasm, because it largely determines the future prospects of success in breeding programs. Comprehensive studies of genetic diversity based on molecular markers has been reported in maize (MESSMER et al., 1992; MELCHINEGER et al., 1991; LIVINI et al., 1992; AJMONE MARSAN et al., 1998; DUBREUIL et al., 1996, SRDIĆ et al., 2006).

Standard methods in maize breeding imply numerous crosses of inbred lines to different testers so as to gain information on genetic similarity, i.e. diversity of these inbreeds. According to this, inbreeds are allocated into specific heterotic groups. Just these extensive field studies are the most expensive and the most time consuming part of contemporary maize breeding and selection, and at the same time this procedure is very restrictive due to the fact that only a few inbreeds can be crossed and estimated.

An alternative could be an allocation of inbred lines in heterotic groups based on molecular markers. As this approach would provide the use of a greater number of inbreeds it would significantly accelerate the process of development of superior hybrids and decrease costs that burden the maize breeding programme.

Commercial maize germplasm is divided into 12 groups based on patterns of heterotic behavior. Members of partially heterotic groups should be detectable by their genetic relatedness. We have examined 125 maize inbred lines from different heterotic groups and miscellaneous origin using different molecular markers (protein, RAPD). Genetic distances between the genotypes were determined from the molecular marker data, and cluster analysis was used to find groups of related genotypes. The clusters found with molecular markers closely resemble the heterotic groups to which the 125 genotypes belonged. Three main groups were distinguishable: a group of BSSS lines, a group of Lancaster lines, and a set of lines with European background. Similar groups were detected despite differences in marker type and genetic distance method used (DRINIĆ et al., 2000, RADOJČIĆ 2006, DRINIĆ 2005, SRDIĆ, 2006).

Information on genetic diversity of commercially grown maize hybrids is very important for germplasm enhancement, hybrid breeding and in preventing environmental damage. Genetic uniformity implies risks of genetic vulnerability to stress factors, which may be reduced by use of unrelated single cross hybrids. It is important to choose among hybrids the one that will give highest yield and answer to environmental stress due to their existing genetic diversity (TROYER et al., 1983). Three methods of determining genetic diversity among maize hybrids are being used: a method that is based on molecular markers (MELCHINEGER et al., 1991; NAGY and MARTON, 2006), a method that is based on the genetics of inbreeding and heterosis for grain yield (TROYER et al., 1988, WILLIAMS and HALAUER, 2000) and a method based on pedigree data (SMITH et al, 2004).

Hybrid maize breeding programs in Serbia as well as at Maize Research Institute were started in the 1950s. Five periods, each characterized by introduction of the new potentially higher yielding ZP hybrids with other agronomic characteristics improved, have determined maize breeding program at the Maize Research Institute "Zemun Polje" (DRINIĆ et al., 2006).



ZP hybrids from different periods have been study by RAPD markers (ERIC, 2003, BAUER *et al.* 2005, BAUER *et al.*, 2007). Cluster analysis showed distinctive grouping of hybrids from each period. Changes in genetic background of parental genotypes during the last 50 years have a major impact on genetic diversity among ZP maize hybrids.

#### *Heterozis background investigation by molecular markers*

Comparing to other crop species maize has probably the highest level of genetic polymorphism. The most significant practical consequence of the huge genetic diversity between maize genotypes is the phenomenon of hybrid vigor or positive heterosis. Maize breeders have always been interested in choosing the parental lines which would result in positive heterotic combination without necessarily making all possible crosses among the potential parental combination. The various methods are in use to predict heterosis and can be grouped into (i) per se performance, (ii) combining ability and (iii) genetic diversity as determined through geographic origin, morphological and agronomic traits as well as molecular markers. The experimental data indicate that heterosis is a function of heterozygosity in a higher number of loci and that the increase of the heterozygous loci number by crosses to genetically distant lines or populations increases the level of heterosis in the crosses. Based on this hypothesis HALLAUER *et al.* (1988) assumed that the magnitude of heterosis could be predicted on the basis of inbred lines differences obtained after use of molecular markers.

Different classes of molecular markers have been used to analyze the genetic relationships among maize inbred lines and to examine the relationship between marker-based GD and heterosis in maize (LEE *et al.* 1989; SMITH *et al.*, 1990; BOPPENMAR *et al.*, 1992; MELCHINGER *et al.*, 1993). Correlation level varies depending on the analysed material and various types of gene effects, pointing to the complexity of the genetic background of heterosis. The general conclusion of studies based on the results of molecular markers application is that heterosis is significantly related to heterozygosity of marker loci.

In our studies on the possibility to correlate heterosis and to make prediction of heterosis by the use of molecular marker data we did obtain high and significant correlation between GD from analyses of inbred lines with protein markers (KONSTANTINOV *et al.*, 1996, DRINIC *et al.*, 2006), RAPD (SRDIĆ *et al.*, 2006; DRINIĆ *et al.*, 2006, RADOJČIĆ *et al.*, 2006, DRINIĆ *et al.*, 2007), SSR and ALFP (DRINIĆ *et al.*, 2002). The results indicating that GD based on molecular markers at the level of gene expression – polypeptides pattern in dry seed, is correlated with heterosis and that markers could be used for prediction of heterotic effect.

#### *Germplasm characterization*

The development of genetics and molecular biology has opened a new chapter in the field of describing agronomical important genotypes and providing their much more detailed characterization, not only in terms of how distant their

germplasm are from those of the existing one but also in the sense of monitoring the uniformity and stability of their characteristics relative to each other. Combining morphological, biochemical and molecular aspects in identification and description of agronomical important genotypes, it is possible to reveal their unique genetic profiles e.g. fingerprints. The most useful markers for maize germplasm characterization are proteins markers, RFLP, SSR, RAPD, and AFLP.

The SDS-PAGE method of protein profile sample mixture is included in ISTA international rules as a technique for distinguishing among and identifying commercial genotypes of different plant species. This method was used for genetic characterization of maize inbred lines as well as for distinguish sister lines at the Maize Research Institute, and for first screening of genetic purity of hybrid seed (DRINIĆ et al., 2000., ERIĆ, et al., 2003., DRINIĆ et al., 2006). All analyzed genotypes have unique protein pattern and unique code - combination of numbers and letters - have been assigned to them.

Most polypeptides identified from the electrophoretic or chromatographic profiles are mostly products of the genes expression that are unevenly distributed in the genome and expressed at the developmental stage of the tissue, the sample has been taken from. This prevents providing a sample that would cover the entire genome (GALOVIĆ et al, 2006). The complete coverage of a genome can be achieved by the use of molecular markers for the variability identification at the level of DNA - DNA polymorphism. Genetic characterization of maize inbred lines with standard kernel type as well as popcorn and sweet corn inbreds from MRI collection was done by RFLP (fig 4) (KONSTANTINOV and DENIĆ., 1985., KONSTANTINOV et al., 1985., KIDRIĆ et al., 1987., KONSTANTINOV et al., 1988) RAPD (fig 5), SSR and AFLP markers (DRINIĆ et al 2000, DRINIĆ et al. 2004., ERIĆ, 2003).




Fig. 4. Hydrolyze DNA of different maize mutants hybridize with same cDNA probe (unpublished data)




Fig. 5. RAPD of six popcorn inbred lines (unpublished data)

General conclusion, based on the results reported by different research teams, including our, is that plant breeding program demands the multidisciplinary approaches independent on plant species.

The authors feel honored to dedicate this paper to the memory of Prof. Janko Dumanovic, who gave an exceptional contribution to the development and promotion of genetics in Serbia.

Received June 22<sup>nd</sup>, 2007

Accepted August 17<sup>th</sup>, 2007

#### REFERENCES

- ALLENDE, J.E., M. BRAVO, (1966): Amino acid incorporation and amini-acyl transfer in wheat embryo. *J. Biol. Chem.* 241, 5813.
- AUKERMAN, M.J., and J.R. SCHMIDT, (1994): Regulation of a zein gene expression during maize endosperm development. In: "Results and Problems in Cell Differentiation, Vol. 20 (ed. L. Nover and E. Dennis). Pp 209 – 233. Springer – Verlag, Berlin.
- BAUER I., S.MLADENOVIC DRINIC, and G. DRINIC, D. IGNJATOVIC MICIC (2007): Assessing temporal changes in genetic diversity of maize hybrids using RAPD markers. *Cereal Research Communications*, vol. 35, no.3, 2007. (In press).
- BAUER, I., S. MLADENOVIC DRINIC, M. FILIPOVIC, and K. KONSTANTINOV (2005): Genetic characterization of early maturing maize hybrids (*Zea mays* L.) obtained by protein and RAPD markers. *Genetika*, 37, 235.
- BHASHARAN, S., and M.S. SWAMINATHAN (1960): Polyploidy and radiosensitivity in wheat and barley. 1. Cytological and cytochemical studies. Part I. *Genetics* 37, 449.
- BONNER, J., G.R. CHALKLEY, M. DAHMUS, D. FAMBROUGH, F. FUJIMURA, R.C.C. HUANG, J.HUBERMAN, K. MARUSHIGE, H. OHLENBUSCH, B. OLIVERA, J. WIDHOLM (1968): Isolation and characterization of chromosomal nucleoproteins. In: "Methods in Enzymology", XII A, Academic Press. New York, 3.
- BOPPENMAIER, J., A.E.MELCHINEGER, E.BRUNKLANS-JUNG, H.H.GEIGEN, R.G.HERRMAM (1992): *Crop Sci* 32, 895.

- BOROJEVIĆ, K., S. BOROJEVIĆ (1969): Stabilization of induced genetic variability in irradiated populations of vulgare wheat. In: "Induced Mutations in Plants. IAEA/FAO Pulman , 399.
- BURR, B., and F.A. BURR (1976): Zein synthesis in maize endosperm by polyribosomes attached to protein bodies. Proc. Natl. Acad. Sci. USA 75, 696.
- DENIĆ, M. (1970): Role on the microsomal fraction in the regulation of protein synthesis in maize endosperm. In: "Improving Plant protein by Nuclear Techniques". IAEA/Fao, Vienna.
- DENIĆ, M. (1971): Contribution to the study of biochemical background protein synthesis in wheat and maize: Ph. Thesis, Agriculture Faculty, University of Belgrade.
- DENIĆ, M., K. KONSTANTINOV, J. DUMANOVIĆ (1971): Uticaj *opaque-2* gena na neke osobine zrna kukuruza. Genetika 3 (1),35.
- DENIĆ, M., D.MILIVOJEVIĆ (1978 ): Ispitivanje aktivnosti proteinskih granula pri ugrađivanju leucina u proteine endosperma normalnog i *opaque-2* kukuruza. Arhiva za poljoprivredne nauke, godXXXI, Sv.116, 3-11.
- DENIĆ,M., K. KONSTANTINOV, J.DUMANOVIĆ (1973): Relationship between ribonuclease activity and content of basic amino acids in maize endosperm. Genetika 5, (2), 167.
- DJOKIĆ, A., J. DUMANOVIĆ, K. KONSTANTINOV (1967): Uticaj etilenimina (EI), etilmetansulfonata (EMS) i etilnitrozouree (ENU) na frekvencu hromozomskih aberacija u mitozu pšenice. Zbornik radova Poljoprivrednog fakulteta, God.XV, br. 448.
- DRINIĆ G., N. BARIŠIĆ, S.MLADENOVIĆ-DRINIĆ (2000): Genetic Relationship Among Maize Inbred Lines Revealed By Protein Molecular Markers. Maize and Sorghum EUCARPIA. XVIII International conference on Maize and Sorghum Genetics and Breeding at the end of the 20th Century. Belgrade, Yugoslavia, June 4-9, 2000.
- DRINIĆ G., G.STANKOVIĆ, Z.PAJIĆ, J VANČETOVIĆ., D.IGNJATOVIĆ MICIĆ, 2007. Sixty years of ZP® Maize Hybrids Breeding. Maydica vol.52, no3.
- DRINIC G., S. DRINIC MLADENOVIC, A.RADOJCIC, G.TODOROVIC G. (2006): Correlation of genetic distance of maize inbred lines with combining ability and heterosis. XXth international Conference of the Eucarpia Maize and sorghum section, Budapest, Hungary, 20-24 june.
- DRINIĆ MLADENOVIĆ S, A.RADOJČIĆ, M.FILIPOVIĆ, G.DRINIĆ, 2007. Genetic distance and heterosis in maize. International scientific conference „Plant genetic stocks - The basis of agriculture of today“, 13-14 june 2007, IPGR, Sadovo, Bulgaria, Book of paper in press.
- DRINIĆ MLADENOVIĆ, S., D. IGNJATOVIĆ MICIĆ, I. ERIĆ, V. ANDELKOVIĆ, D. JELOVAC and K. KONSTANTINOV ( 2004): Biotechnology in maize breeding. Genetika 36, No. 2, 93-111.
- DRINIĆ MLADENOVIĆ, S., J. SRDIĆ, G.DRINIĆ and K. KONSTANTINOV (2004): Genetic divergence of maize inbred lines based on molecular markers. In „Genetic variation for plant breeding“. Proc. XVIIth EUCARPIA General Congress. Pp. 53-57. 8-11 September 2004, Tulin, Austria.
- DRINIĆ MLADENOVIĆ,S., S. TRIFUNOVIĆ., G. DRINIĆ., and K. KONSTANTINOV (2002): Genetic divergence and its corelation to heterosis in maize as revealed by SSR based markers. Maydica 47, 8
- DUMANOVIĆ, J., M. DENIĆ, L. EHRENBERG, K.G. BERGSTRAND (1969): Radiation – induced heritable characters in wheat. Hereditas 62, 221.
- DUMANOVIĆ, J., M. DENIĆ, K. KONSTANTINOV (1974): Variation of kernel phenotype and some biochemical properties in *opaque-2* from different genetic background. Genetika Vol.6, No. 2, 211.

- DUMANOVIĆ, J., L. EHREBERG (1968): Modern breeding methods for improving protein quality and quantity in plants. In: "Isotope studies on the nitrogen chain". IAEA/FAO Vienna.
- EGGUM, B.O., J. DUMANOVIĆ, D. MIŠEVIĆ, M. DENIĆ (1985): Nutritive quality and energy yield of high oil, opaque-2 and waxy maize hybrids compared to normal maize hybrids. In: "Qual. Plant Foods Hum. Nutr. 35, 165." Martinusnilhoff/Dr. W. Junk Publishers, Dordrecht, Printed in The Netherlands.
- EHREBERG, L., A. GUSTAFSSON, U. LUNDQUIST (1959): The mutagenic effects of ionizing radiations and reactive ethylene derivatives in barley. *Hereditas* 45, 351.
- ERIĆ, I., S. MLADENOVIĆ DRINIĆ, K. KONSTANTINOV and G. STANKOVIĆ (2003): Genetic diversity among ZP maize hybrids from different selection cycles obtained by protein markers. *Maize Genetic Cooperation Newsletter MGNL* 77, 7-8.
- GALOVIĆ, V., S. DRINIĆ - MLADENOVIĆ, J. NAVALUŠIĆ, M. ZLOKOLICA (2006): Characterization methods and fingerprinting of agronomically important crop species. *Genetika*, vol.38, no.2, 83 - 96.
- HAGBERG, A., A. GUSTAFSSON, L. EHREBERG (1958): *Hereditas* 44, 523.
- HALLAUER, A.R., W.A. RUSSELL and K.R. LAMKEY (1988): Corn breeding. In *Corn and Corn Improvement*, Dudley, J.W. and Sprague, G.W. (eds), 3<sup>rd</sup> ed., ASA-CSSA-SSSA, Madison, 463-564.
- HARTINGS, H., M. MADDALONI, N. LAZZARONI, N. DI FONZO, M. MOTTO, F. SALAMINI, R.D. THOMPSON (1989): The O2 gene which regulates zein deposition in maize endosperm encodes a protein with structural homologies to transcriptional activators. *EMBO J.* 8, 2795.
- HURKMAN, W.J., D.L. SMITH, J. RICHTER, B.A. LARKINS (1981): Subcellular compartmentalization of maize storage proteins in *Xenopus* oocytes injected with zein messenger RNAs. *J. Cell. Biol.* 89, 292.
- JOVANOVIĆ, Č., K. KONSTANTINOV (1973): Effects of different mutagenic agents on the frequency of chromosomal aberrations in M5 – progeny of wheat. *Genetika* Vol. 5, No. 1, 21.
- KIDRIČ, M., K. KONSTANTINOV, M. DENIĆ and B. RISTOVIĆ (1987): Existing and induced methylation site polymorphism of maize genome. 18th FEBS Meeting, p. 145, June 28 – 3 July, Ljubljana, Yugoslavia.
- KONSTANTINOV, K, T. ČORIĆ, G. DRINIĆ and S. MLADENOVIĆ DRINIĆ (1996): F1 embryo proteins as valuable tools in better understanding of heterosis. *Maize Genetics Cooperation Newsletter* 70, pp. 72-73.
- KONSTANTINOV, K. (1968): Investigations of factors influencing seed sensitivity to some mutagenic agents. *nekim mutagenim sredstvima*. M.Sc, Faculty of Agriculture, University of Belgrade.
- KONSTANTINOV, K. (1975): Role of some chromatin components in genetic control of storage protein synthesis in maize endosperm. Ph. D. Thesis, Agriculture Faculty, University of Belgrade.
- KONSTANTINOV, K. (1978): Contribution to the study on the role of chromatin in the synthesis of protein in maize endosperm. *Genetika*, 10, 1, 115.
- KONSTANTINOV, K. and J. MOUTSCHEN (1970b): Centromeric localization of isoprophyl methanesulfonate (iPMS) induced aberrations in plant species. *Genetika* 2, 3.
- KONSTANTINOV, K. and M. DENIĆ (1985): Recombinant DNA technology in maize breeding. I. Evaluation of the relationship between maize genotypes using clones of genomic DNA. *Genetika* 17, 207.

- KONSTANTINOV, K. M. DENIĆ and J. DUMANOVIĆ (1985): Recombinant DNA technology in maize breeding. II. Genomic clones in study of maize genome organization. *Genetika* 17, 229.
- KONSTANTINOV, K., M. DENIĆ (1975): Isolation and some characteristics of chromatin from developing maize endosperm. In: „Breeding for seed protein improvement using nuclear techniques“ International Atomic Energy Agency, Vienna. 99.
- KONSTANTINOV, K., M. DENIĆ and J. DUMANOVIĆ (1988): Recombinant DNA technology in maize breeding. IV. Restriction endonuclease analysis of nuclear DNA in maize inbreds and hybrids of different maturity ratings. *Genetika* 20,63.
- KONSTANTINOV, K., S. MLADENOVIĆ DRINIĆ, T. ĆORIĆ, D. IGNJATOVIĆ, G.SARATLIĆ and J. DUMANOVIĆ (1997): Maize as Host for *Agrobacterium*. In „Genetics biotechnology and Breeding of Maize and Sorghum“ Ed. A.S.Tsaftaris, Published by The Royal Society of Chemistry, Cambridge, UK.
- KONSTANTINOV, K., M. HARMS-RINGDAHL, L. EHRENBERG, J. DUMANOVIĆ and I. BOŽOVIĆ (1970a): Influence of pH and temperature on the effects of ethyleneimine (EI) in wheat and barley seeds. *Radiation botany* 10, 499
- KUCKUCK, H., G. KOBABE, and G. WENZEL (1991): Fundamental of plant breeding. Springer – Verlag, Berlin.
- LEE, M. (1995): DNA markers and plant breeding programs. *Adv. Agron.* 55, 265.
- Lohmer, S. M., M. Maddolini, M. Motto, F. Salamini, R.D. Thompson (1993): Translation of the mRNA of the maize transcription activator Opaque -2 is inhibited by upstream open reading frames present in the leader sequence. *Plant Cell* 5, 65.
- LOHMER, S., M. MADDALONI, M. MOTTO, F. SALAMINI, R.D. THOMPSON (1991): The maize regulatory locus *Opaque -2* encodes a DNA - binding protein which activates the transcription of the *b - 32* gene. *EMBO J.* 10, 617.
- MAC KEY, J. (1959): Mutagenic response in triticum at different levels of ploidy. *Proc. 1<sup>st</sup>. int. Wheat Genet. Symp., Winnipeg 1958, Public. Press Winnipeg*, p. 88.
- MADDOLINI, M., N. DI FONZO, H. HARTINGS, H. N. LAZZARONI, F. SALAMINI, R.D. THOMPSON, M..MOTTO (1989): The sequence of the zein regulatory gene *opaque-2 (O2)* of *Zea mays*. *Nucleic Acids Res.* 17, 7532.
- MELCHINGER, A., M.M. MESSMER, M.LEE, W.L.WOODMAN, K.R. LAMKEY (1991): Diversity and relationships among U.S. maize inbreds revealed by restriction fragment length polymorphisms. *Crop Sci.* 31, 669.
- MERTZ, E., L.S. BATEST, O.E. NELSON (1964): Mutant gene that changes protein composition and increases lysine content. *Science* 145, 279.
- MESSMER, M..M., A.E. MELCHINGER, R.G. HERRMANN, and J. BOPPENMAIER (1993): Relationships among early European maize inbreds:II. Comparison of pedigree and RFLP data. *Crop Sci.* 33:944-950.
- MLADENOVIĆ S. (1990): Dobijanje transgenih biljaka kukuruza unošenjem bakterijskih gena. Magistrarska teza, Prirodno-matematički fakultet, Univerzitet u Beogradu, 1990.
- MLADENOVIĆ, S., K. KONSTANTINOV, M. DENIĆ, and B. TADIĆ (1991): Recombinant DNA technology in maize breeding. VI. Foreign gene influence on maize genome expression after plant transformation by different technologies. *Genetika* 23, 191.

- MOTTO, M., M. MADDALLONI, G. PONZIANI, M. BREMBILLA, R. MAROTTA, N. DI FONZO, C. SOAVE, R.D.THOMPSON, F. SALAMINI (1988): Molecular cloning of the *o2-m5* allele of *Zeam mays* using transposon marking. *Mol Gen. Genet.* 212,488.
- MOTTO,M., R. MAROTTA, N. DI FONZO, C. SOAVE, F. SALAMINI (1986): *Ds*-induced alleles at the *opaque-2* locus of maize. *Genetics* 112,121.
- MOUSCHEN – DAHMEN, J., L. EHRENBURG (1968): Note on the chromosome breaking activity of ethyleneoxide and ethyleneimine. *Hereditas* 60, 267.
- NAGY, E., L.C. MARTON (2006): The use of morphological traits and genetic markers to estimate genetic relationships in maize. *Cereal Research Communications* 34 (2-3) 887.
- NEUFER, M.G., E.H. COE, R.S. WESSLER (1997): Mutants of Maize. Cold Spring Laboratory Press.
- PIRONA R., H.HARTINGS, V.ROSSI and M..MOTTO (2007): Transcriptome analysis of *opaque-2* and *opaque-7* mutants in maize endosperm. *MGNL* 81, 4.
- POPOVIĆ, V. (1986): Proučavanje biohemijske osnove genetičke kontrole sinteze rezervnih proteina kukuruza. Doktorska disertacija, Prirodno matematički fakultet Univerziteta u Beogradu.
- POPOVIĆ, V., K. KONSTANTINOV, M. DENIĆ (1981): *In vitro* RNA and protein synthesis by RNA polymerase from *opaque -2* and normal *Zea mays* endosperm. *Abhdgl. Akad. Wiss. DDR, Abt. Math., Naturwiss., Techn. No. 5, p. 225.*
- POPOVIĆ, V., M. DENIĆ, J. DUMANOVIĆ (1974): Influence of the *opaque - 2* gene on proteins in maize of different genetic backgrounds. *Genetika* 6 (1), 7.
- SALAMINI, F. (1980) : Genetic instability at the *opaque -2* locus of maize. *Mol. Gen. Genet.* 179, 497.
- SCHMIDT, R.J., A.F. BURR, A.J. AUKERMANN, B. BURR (1990): Maize regulatory gene *opaque-2* encodes a protein with a “ leucine –zipper” motif that binds to zein DNA. *Proc. Natl. Acad.Sci. USA* 87,46.
- SCHMIDT, R.J., F.A. BURR, B. BURR (1987): Transposon tagging and molecular analysis of the maize regulatory locus *opaque -2*. *Science* 238, 960.
- SCHMIDT, R.J., M. KETUDAR, J.M. AUKERMANN, G. HOSCHEKI (1992): *Opaque- 2* is a transcriptional activator that recognizes a specific target site in 22-kD zein genes. *Plant Cell* 4, 689.
- SHULL, G.H. 1910. *Am. Breeders Mag.* 1, 98-107.
- SMITH, J.S., D.DUVICK, O.S. SMITH, M. COOPER, L. FENG (2004): *Changes in Pedigree Backgrounds of Pioneer Brand Maize Hybrids Widely Grown from 1930 to 1999.* *Crop Sci.* 44:1935-1946.
- SRDIĆ J., S. MLADENOVIĆ DRINIĆ, Z. PAJIĆ ( 2006): Combining abilities and genetic resemblance of maize inbred line. *Acta Agronomica Hungarica* vol.54, No.3, 337.
- TROYER, A.F., S.J. OPENSHOW, K.H. KNITLE (1988): Measurement of genetic diversity among popular commercial corn hybrids. *Crop Sci.* 28: 481-485.
- TROYER, A.F.,S.J. OPENSHOW, K.H. KNITLE (1983): Measurement of genetic diversity among commercial corn hybrids. *Method.* p83. In: *Agronomy abstracts, ASA, Madison, WI.*
- VARAGONA, M.J., R.J..SCHMIDT, V.N. RAIKHEL, (1991): Monocot regulatory protein *opaque-2* is localized in the nucleus of maize endosperm and transformed tobacco plants. *Plant Cell* 3, 105.
- VARAGONA, M.J., J.R. SCHMIDT, V.N. RAIKHEL (1992): Nuclear localization signal(s) required for nuclear targeting of the maize regulatory protein *opaque-2*. *Plant Cell* 4, 1213.
- VASIL, S.K. (1972): Use of genetic modifiers to obtain normal type kernels with *opaque-2* gene. *Proc. Symp. On Protein Quality in Maize. Mexico, El Battan.*
- WILLIAMS, T.R., A.R. HALLAUER (2000): diversity among maize hybrids. *Maydica* 45, (3) 163.

## MOLEKULARNA GENETIKA – POSTEPENO UVOĐENJE U OPLEMENJIVANJE KUKURUZA

Kosana KONSTANTINOV i Snežana MLADENOVIĆ DRINIĆ

Institut za kukuruz „Zemun Polje“, Beograd, Srbija

### Izvod

Efikasnost procesa oplemenjivanja je primarno determinisana mogućnostima utvrđivanja genetičkog polimorfizma, produktivnosti i stabilnosti prinosa početnog materijala. Poznavanjem biohemijske osnove genetičke kontrole željene osobine moguće je izvršiti poželjnu promenu ili modifikaciju ili indukovanjem mutacija ili unošenjem već postojećih poznatih gena nosioca mutacija klasičnim procesom ukrštanja sa različitim genotipovima. U radu je dat parcijalni pregled indukovanja mutacija kod pšenice i kukuruza, dobijenih korišćenjem različitih vrsta jonizujućeg zračenja (brzi neutroni,  $\gamma$  - zraci) ili hemijskih mutagena (EI, iPMS, EMS, ENU).

U drugom delu su prikazani neki od rezultata unošenja *opaque-2* gena u različite genetičke osnove kukuruza primenom klasičnih metoda oplemenjivanja. Istovremeno su navedeni rezultati istraživanja biohemijske osnove *opaque2* mutacije, uključujući korišćenje ukopčanog bezćelijskog sistema za sintezu RNK i proteina u prisustvu različitih ćelijskih komponenata (hromatin, RNK polimeraza, mikrozomalna frakcija, proteinska telašca) izolovanih iz endosperma normalnog genotipa i *opaque 2* mutanta kukuruza.

Deo rada predstavlja pregled rezultata istraživanja u oblasti molekularnog oplemenjivanja – korišćenja različitih klasa molekularnih markera (RFLP, AFLP, RAPD, SSR) u izučavanju genetičkog polimorfizma kukuruza. Dat je i prikaz rezultata ispitivanja molekularne osnove heterozisa kod kukuruza korišćenjem molekularnih markera.

Primljeno 22. VI. 2007.

Odobreno 17. VIII. 2007.