



Published in final edited form as:

Expert Rev Mol Med. ; 11: e19. doi:10.1017/S1462399409001112.

Molecular mechanisms of endothelial hyperpermeability: implications in inflammation

Puneet Kumar[†], Qiang Shen[†], Christopher D. Pivetti, Eugene S. Lee, Mack H. Wu, and Sarah Y. Yuan^{*}

Division of Research, Department of Surgery, University of California at Davis School of Medicine, Sacramento, CA 95758, USA.

Abstract

Endothelial hyperpermeability is a significant problem in vascular inflammation associated with trauma, ischaemia–reperfusion injury, sepsis, adult respiratory distress syndrome, diabetes, thrombosis and cancer. An important mechanism underlying this process is increased paracellular leakage of plasma fluid and protein. Inflammatory stimuli such as histamine, thrombin, vascular endothelial growth factor and activated neutrophils can cause dissociation of cell–cell junctions between endothelial cells as well as cytoskeleton contraction, leading to a widened intercellular space that facilitates transendothelial flux. Such structural changes initiate with agonist–receptor binding, followed by activation of intracellular signalling molecules including calcium, protein kinase C, tyrosine kinases, myosin light chain kinase, and small Rho-GTPases; these kinases and GTPases then phosphorylate or alter the conformation of different subcellular components that control cell–cell adhesion, resulting in paracellular hyperpermeability. Targeting key signalling molecules that mediate endothelial-junction-cytoskeleton dissociation demonstrates a therapeutic potential to improve vascular barrier function during inflammatory injury.

Endothelial cells lining the inner surface of microvessels form a semipermeable barrier that actively participates in blood–tissue exchange of plasma fluid, proteins and cells. The precise regulation of endothelial permeability is essential for maintaining circulatory homeostasis and the physiological function of different organs. As a result, microvascular barrier dysfunction and endothelial hyperpermeability represent crucial events in the development of a variety of disease processes, such as adult respiratory distress syndrome (ARDS), ischemia–reperfusion (I–R) injury, diabetic vascular complications, and tumour metastasis. Better insight into the molecular mechanisms underlying pathogenic conditions related to microvascular hyperpermeability is required for developing effective therapeutic strategies. Following intensive studies over the past few decades, it is now understood that endothelial permeability is mediated through a transcellular pathway (across cells) and a paracellular pathway (between cells), both of which are highly regulated by mechanical forces and biochemical signals.

Transcellular versus paracellular permeability

An important molecular mechanism underlying transcellular permeability is macromolecule transcytosis via caveoli – specialised plasmalemmal vesicles containing caveolin-1. The involvement of caveolin-1 in regulating cardiovascular functions associated with endothelial barrier properties has been demonstrated through studies using transgenic and knockout

© 2009 Cambridge University Press

*Corresponding author: Sarah Yuan, Department of Surgery, University of California Davis Medical Center, 4625 2nd Avenue, Room 3006, Sacramento, CA 95817, USA. Tel: +1 916 703 0422; Fax: +1 916 703 0421; sarahyuan@ucdavis.edu.

[†]These authors contributed equally to this work.

animals (Refs 1,2,3,4). Upon binding to plasma proteins (of size >3 nm, e.g. albumin), the cell-surface docking protein gp60 interacts with caveolin-1 and signalling intermediaries, including a G protein and Src-family tyrosine kinase. This cascade results in the formation and release of albumin- and solute-containing caveolae from the apical membrane (Ref. 5). The vesicles are subsequently transported to the basal membrane and release their contents through exocytosis (Ref. 5). Recent ultrastructural evidence suggests that caveolae-like vesiculovacuolar organelles can interconnect to each other, forming secondary, grape-like structures that function as transmembrane channels for molecular trafficking across the cell (Ref. 6). Receptors of endogenous permeability-enhancing agents have been identified on the surface of these channels, indicating the possibility of their participation in transcytosis-related endothelial permeability (Refs 7,8).

Despite the potential contribution of transcytosis to the basal permeability of the endothelium, paracellular flux of plasma fluid and proteins through endothelial cell–cell junctions has been emphasised for its pathophysiological importance in vascular inflammation during disease and injury. Among the different types of junction structures in the vascular endothelium, the tight junction and adherens junction are the best characterised with respect to their function in mediating cell–cell adhesion and thus barrier properties. Briefly, the tight junction is a zipper-like structure formed at the cell–cell contact area by a group of transmembrane proteins primarily expressed in the blood–brain barrier and retinal microvasculature, including claudins, occludins and zonular occludins (ZO-1 and ZO-2). The adherens junction has been identified in nearly all types of vascular beds, especially in the peripheral microvasculature (Fig. 1). Its molecular structure is based on VE-cadherin (vascular endothelial cadherin), a transmembrane receptor whose extracellular domain homophilically binds to the extracellular domain of another VE-cadherin molecule from an adjacent cell and whose intracellular domain is anchored to the cell cytoskeleton via a family of actin-binding proteins called catenins (α , β , γ and p120 catenins). The catenins not only serve as a structural linkage between VE-cadherin and the cytoskeleton, but also transduce biochemical signals for cell–cell communications (Refs 9, 10). Moreover, endothelial cells are tethered to the extracellular matrix through focal adhesions, which consist mainly of integrin transmembrane proteins and a family of actin-linking proteins including focal adhesion kinase (FAK), talin and paxillin (Ref. 5) (Fig. 1). The stability of this junction–cytoskeleton complex withstands fluid shear stress and is essential in maintaining the endothelial barrier function.

Many inflammatory mediators are capable of disrupting the interendothelial junction assembly, thereby causing endothelial hyperpermeability. More in-depth molecular analyses suggest that the mechanism underlying inflammation-induced endothelial paracellular hyperpermeability involves phosphorylation, internalisation or degradation of the junctional molecules (Ref. 11). In addition, the junction–cytoskeleton complex participates in other cellular processes including molecular scaffolding, intracellular trafficking, transcription and apoptosis that may directly or indirectly alter vascular barrier function (Ref. 12). Regardless of the molecular details, however, essentially all permeability responses in the vascular endothelium are initiated with receptor occupancy followed by a series of intracellular signalling cascades, some of which are described below (Fig. 2).

Intracellular signal transduction

Cytosolic calcium

Many inflammatory agonists mediate endothelial hyperpermeability via a calcium (Ca^{2+})-dependent mechanism. The signalling cascade is triggered when an agonist binds to its respective receptor expressed on the endothelial surface, which activates phospholipase C (PLC) either directly (e.g. VEGF receptor) or via a G protein (e.g. histamine receptor) (Fig. 2). This culminates in inositol 1,4,5-trisphosphate [$\text{Ins}(1,4,5)\text{P}_3$] production and Ca^{2+} release

from intracellular stores as well as influx via various channels. Elevated intracellular Ca^{2+} further targets cytoskeleton proteins or junction structures that determine the paracellular permeability property.

At the cytoskeleton, Ca^{2+} activates Ca^{2+} /calmodulin-dependent myosin light chain kinase (MLCK), which phosphorylates myosin light chain (MLC) and promotes a cross-bridge movement between actin and myosin (Fig. 1). This interaction generates a contractile force that pulls neighbouring cells apart from each other (retraction), leading to intercellular gap formation (Ref. 13). In parallel, Ca^{2+} stimulates nitric oxide synthase (NOS), which produces nitric oxide (NO) and subsequently cyclic GMP (cGMP); the latter activates cGMP-dependent protein kinase (PKG). The NO–PKG cascade is known to mediate endothelial barrier structural and functional responses to a wide spectrum of physiological and pathological mediators, including angiogenic growth factors and proinflammatory agents (Refs 14,15). Additionally, intracellular Ca^{2+} has been shown to participate in the transcellular transport of macromolecules via caveoli, as demonstrated by Ca^{2+} waves in caveolin-1-rich cell edges under shear stress or ATP stimulation (Ref. 16).

Protein kinase C

Protein kinase Cs (PKCs) are a family of serine/threonine kinases that differ in structure, expression, subcellular distribution, and function. Their activities are differentially regulated by Ca^{2+} , diacylglycerol (DAG) and phospholipids. Some isoforms can be activated by synthetic compounds such as phorbol 12-myristate 13-acetate (PMA). The downstream signals of PKCs include the Rho family of GTPases, mitogen-activated protein kinases (MAPKs) and MLCK. More specifically, PKCs induce rapid phosphorylation of the Rho-GDP dissociation inhibitor (GDI), promoting RhoGDP–GTP exchange and Rho-associated protein kinase (ROCK) activation coupled with increased actomyosin ATPase activity (Ref. 17). With respect to the MAPK cascades, PKCs trigger Ras-dependent signalling through a sequential activation of a Ras protein, Raf protein, MAPK kinase (MEK1/2) and extracellular-signal-regulated kinase (ERK1/2) (Refs 18,19). In particular, PKCs cause a time-dependent increase in the activity of Ras, which binds CRAF (Raf-1), a serine/threonine kinase capable of activating MEK1/2 and subsequently ERK1/2. Inhibition of Ras completely abolishes PMA-induced CRAF activity, and MEK1/2 inhibitors attenuate PMA-induced hyperpermeability, consistent with the role of PKC-MAPK signalling in endothelial barrier regulation (Ref. 19).

Structurally, PKCs can also directly stimulate the cytoskeletal contractile machinery by phosphorylating MLCK, a key molecule triggering actin–myosin motor function. PKCs also interact with other cytoskeletal proteins known to participate in cell contraction, such as actinin, caldesmon, and the intermediate filament protein vimentin (Ref. 20). Furthermore, PKC activation correlates with disassembly of adherens junctions (Ref. 21) and tight junctions (Ref. 22).

Tyrosine kinases

Both receptor and nonreceptor tyrosine kinases participate in the signal transduction that mediates changes in endothelial barrier structure and function. As a representative of receptor tyrosine kinases, vascular endothelial growth factor (VEGF) receptor 2 (VEGFR-2; also named KDR) undergoes dimerisation and phosphorylation at multiple tyrosine residues upon ligand binding. This provides a molecular configuration for recruiting more intracellular signals and further stimulating the receptor-linked PLC γ . The downstream reactions include PKC activation, Ca^{2+} release, NO production, and MAPK signalling (Refs ^{23,24,25,26,27,28}). In addition to these well-characterised pathways, VEGFR-2 has been shown to mediate endothelial hyperpermeability by activating p21-activated kinase (PAK) (Ref. 29) and by phosphorylating FAK, thereby triggering integrin signalling (Ref. 30). In endothelial adherens

junctions, both VE-cadherin and catenins (β -, γ - and p120-catenin) can be tyrosine phosphorylated by this receptor kinase in the presence of VEGF (Ref. 11). Recently, it became evident that the internalisation of VE-cadherin is closely regulated by VEGF signalling in a β -arrestin-dependent manner (Ref. 31).

Among multiple nonreceptor tyrosine kinases, the Src family (c-Src, Fyn, Yes, Yrk, Lyn, Hck, Fgr, Blk, Lck) has been frequently implicated in the barrier-opening action of various inflammatory mediators and growth factors (Ref. 32). The mechanisms underlying Src-induced paracellular hyperpermeability may involve cytoskeleton contraction driven by MLCK phosphorylation (Ref. 33), and/or junction opening triggered by VE-cadherin and β -catenin phosphorylation (Ref. 34). Furthermore, it has been demonstrated that Src-induced phosphorylation of caveolin-1 and dynamin is crucial for the signalling machinery in directing the polarised transportation of albumin and other solute through the transcellular pathway (Refs 5,35).

Rho GTPases

The activity of Rho GTPases is regulated through GDP–GTP cycling dependent on guanine-nucleotide-exchange factors. Three members of the Rho family (RhoA, Rac and Cdc42) have been characterised for their effects on endothelial permeability through distinct subcellular targeting and functions. In particular, RhoA promotes cell contractility and focal adhesion, whereas Cdc42 and Rac1 are major players in cell membrane protrusion or filopodia formation and thus cell migration (Ref. 36). Under physiological conditions, changes in mechanical forces (e.g. shear stress) act to relay intracellular signals to regulate RhoA activity, thus maintaining endothelial integrity, especially at intercellular junctions (Ref. 37). The importance of increased RhoA activity in endothelial barrier dysfunction is supported by studies showing that RhoA and its effector kinase ROCK mediate endothelial hyperpermeability in response to histamine, thrombin, VEGF, neutrophils, and mechanical stimuli (Refs 8,37,38,39,40,41). Upon activation, RhoA inhibits myosin-associated protein phosphatase by phosphorylating its inhibitory myosin phosphatase targeting (MYPT) subunit via ROCK, indirectly promoting MLC phosphorylation and actin–myosin contraction (Ref. 42). In addition, RhoA increases endothelial permeability by stimulating Ca^{2+} entry, which activates MLCK.

By contrast to the permeability-increasing effect of RhoA, Rac and Cdc42 have been generally accepted as barrier protectors capable of decreasing permeability. Although a recent study suggests that Rac contributes to VEGF-induced hyperpermeability (Ref. 31), activation of Rac is associated with a tightened barrier after treatment with sphingosine 1-phosphate (Ref. 43) and hepatocyte growth factor (Ref. 44). Likewise, Cdc42 has been shown to play a critical role in re-annealing adherens junctions and restoring barrier function in endothelial cells during the recovery phase of inflammatory injury (Ref. 45). While the detailed mechanisms by which these Rho members protect endothelial barrier properties remain elusive, their effects may involve PAK-dependent cytoskeleton redistribution or junction reorganisation (Ref. 43).

cAMP

The adenine nucleotide is a conserved, ubiquitous intracellular second messenger that has been well recognised for its barrier protection function (Refs 14,46). Several mechanisms have been proposed to explain the action of cyclic adenosine monophosphate (cAMP) on endothelial barrier structure. First, increased intracellular cAMP activates its cognate protein kinase, PKA, which by stabilising the actin cytoskeleton counteracts the cell retractile force, thus preventing intercellular gap formation (Ref. 46). Second, activated PKA can inhibit MLCK activity, which in turn leads to decreased MLC phosphorylation and minimises actin–myosin contraction. Finally, cAMP inhibits membrane phospholipid hydrolysis, thereby reducing DAG production and PKC activity. In accordance with these protective effects of cAMP at cell–cell junctions,

previous studies also show that cAMP analogues induce a diffuse distribution of focal adhesions of endothelium and improved barrier function (Ref. 47).

Hyperpermeability mediators

Increased circulating levels of immune cells (e.g. neutrophils) and pro-inflammatory cytokines [e.g. interleukins (ILs) and tumour necrosis factors (TNFs)] or other soluble mediators (e.g. histamine, thrombin and VEGF) are a hallmark of host defence against injury. Most of these mediators produce vasoactive and cytotoxic effects by directly targeting the vascular endothelium and activating the aforementioned signalling pathways. With recent intensive studies, the list of mediators inducing endothelial hyperpermeability continues to grow. In this review, we focus on some typical acute inflammatory mediators and discuss their effects and mechanisms in endothelial hyperpermeability (Table 1).

Histamine

Histamine has long been known as an oedematogenic factor contributing to microvascular leakage in the acute inflammatory response associated with trauma, burns, allergy and certain types of infectious disease. Endothelial paracellular hyperpermeability resulting from intercellular gaps has been accepted as the major cellular mechanism underlying histamine-induced barrier dysfunction. Four subtypes of G-protein-coupled receptors (H_1 , H_2 , H_3 and H_4) have been attributed to histamine action (Refs 48,49), among which H_1 is considered the most important with respect to vascular permeability (Ref. 50). It is well established that histamine binding to G_q -coupled H_1 receptor activates PLC β and elevates intracellular Ca^{2+} , which increases MLCK activity and triggers actin–myosin contraction (Refs 15,51). Other downstream reactions include PKG-dependent MAPK activation (Ref. 52). At the junctional level, histamine signalling causes phosphorylation and disruption of components of the adherens junction and tight junction (Refs 5,53,54).

Thrombin

This procoagulant serine protease is well known for its endothelial hyperpermeability effect, especially in vitro. Its receptors belong to the family of protease-activated receptors (PARs). Of four known isoforms, PAR-1, -3 and -4 are activated by thrombin and PAR-2 is activated by trypsin, whereas PAR-1 is best characterised with respect to endothelial permeability. Similar to the histamine H_1 receptor, PAR-1 signals through G_q -coupled Ca^{2+} mobilisation, PKC activation, and MAPK signalling (Refs 53,55,56). In addition to G_q , other G proteins may participate in PAR-1 signalling as well. For example, PAR-1 coupling of $G_{\alpha_{12/13}}$ has been shown to activate p115RhoGEF (Ref. 57), a RhoA activator that also functions downstream of PKC (Refs 18,58). Furthermore, through the PAR-1 cascades, thrombin is able to activate multiple protein kinases including Src, FAK, PKC and MAPK, as well as to stimulate their crosstalk (Ref. 59).

Vascular endothelial growth factor

VEGF is a glycoprotein originally identified as a vascular permeability factor based on the fact that it caused interstitial accumulation of intravenously injected dyes and ascites in vivo (Ref. 60). Subsequently, its role in pathological angiogenesis has been extensively studied with respect to diabetic retinopathy, I–R injury, and tumour development and metastasis, all of which involve altered endothelial permeability (Ref. 61). Three types of VEGF receptors have been characterised. Although a recent study using venom VEGF (*Tfsv*VEGF) suggests that VEGFR-1 is involved in endothelial permeability responses (Ref. 62), VEGFR-2 is still considered as the primary receptor mediating the hyperpermeability action of VEGF (Refs 40,63). As indicated above, the intracellular signal transduction triggered by this receptor tyrosine kinase involves complex interactions among multiple signalling molecules and

structural proteins (Refs 23,24,25,26,27,28,29,30). While there has been ultrastructural evidence supporting VEGF-induced transcellular permeability, recent studies suggest that VEGF also affects paracellular permeability by promoting cytoskeleton contraction, focal adhesion dynamics, and cell–cell junction opening (Refs 30,40,64).

Neutrophils and neutrophil-released cytokines

In response to injurious or inflammatory stimulation, neutrophils undergo a series of kinetic changes characterised by adherence to the venular endothelium, followed by transendothelial migration and chemotaxis to the site of stimulation (Refs 65,66,67,68,69,70,71), a process often accompanied by plasma fluid and protein leakage (Ref. 72). While the leak response is conventionally attributed to mechanical disruption of the endothelial barrier as a result of neutrophil transmigration, an emerging paradigm emphasises a dynamic and reversible interaction between the endothelium and vasoactive mediators released from neutrophils during adhesion and migration (Ref. 73). In this regard, activated neutrophils produce a variety of hyperpermeability factors, including oxidants, cytokines, lipid metabolites, leukotrienes and proteases. These factors directly or indirectly target the endothelium, inducing a series of biochemical and conformational changes in the barrier. It has been shown that microvascular endothelial cells play an active role in response to neutrophil activation through a Src- and RhoA-dependent endothelial cell–cell interaction, characterised by VE-cadherin and β -catenin phosphorylation and adherens junction disorganisation; the response is coupled with MLC phosphorylation-dependent cytoskeleton contraction (Refs 38,39,73,74). Consistently, many studies have demonstrated that neutrophil-released factors, such as oxidants and cytokines, cause endothelial hyperpermeability by activating the contractile machinery and opening cell–cell junctions through signalling pathways that involve Src, RhoA, PLC, Ca^{2+} , MAPK and MLCK (Refs 75,76,77,78).

Two of the best-studied cytokines released by activated leukocytes and endothelial cells, with respect to their impact on the endothelial barrier, are TNF and IL-1 (Ref. 5). TNF and IL-1 augment endothelial permeability and facilitate leukocyte infiltration in tissues when injected locally. This is mainly mediated through upregulation of endothelial adhesion molecules, such as intercellular adhesion molecule 1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1) and E-selectin that afterwards interact with their ligands on the activated leukocytes to mediate cell arrest and subsequent transendothelial migration (Refs 5,79,80). Recent studies have also implicated PKC, RhoA/ROCK, transient receptor potential canonical 1 (TRPC1), and matrix metalloproteinases (MMPs) in TNF-induced endothelial hyperpermeability (Ref. 5). These subcellular mediators can influence paracellular permeability at the level of adherens junction disorganisation as well as actin cytoskeleton reorganisation.

It has recently been shown that ICAM-1-mediated Src-dependent caveolin-1 phosphorylation is crucial for neutrophil-induced pulmonary vascular hyperpermeability. This finding suggests a potential role of transcellular permeability in neutrophil-mediated inflammatory responses (Ref. 81).

Clinical relevance

Vascular inflammation contributes to the development of a variety of diseases or pathological processes, including trauma, burns, I–R injury, sepsis, diabetes, atherosclerosis, thrombosis, and tumour development and metastasis. Below is a brief discussion of some clinical situations directly relevant to endothelial barrier injury.

Adult respiratory distress syndrome

ARDS is a severe pulmonary condition resulting from a major burn, trauma, sepsis or acute pancreatitis. It is characterised by diffuse pulmonary inflammation, leukocyte infiltration, alveolar-capillary barrier dysfunction, and plasma extravasation. The pathogenesis of ARDS involves multiple pro-inflammatory factors, such as histamine, TNF- α , IL-1 and IL-6 (Refs 82,83). The plasma level of histamine is significantly increased in patients and experimental animals with septic shock (Refs 84,85). Besides a strong bronchoconstrictive effect, histamine increases pulmonary microvascular permeability, leading to lung oedema (Ref. 86). This injury can be augmented by TNF- α and IL-1 β produced from activated leukocytes during the development of ARDS (Ref. 82). The levels of TNF- α in serum and bronchoalveolar lavage fluid are substantially increased in patients (Refs 87,88), whereas blocking TNF- α ameliorates pulmonary oedema in animal models of ARDS (Ref. 89). The mechanisms by which TNF- α causes barrier injury include actomyosin-contraction-driven intercellular gap formation, as well as overexpression of other proinflammatory cytokines (e.g. IL-1 β) and adhesion molecules (e.g. ICAM-1) mediated by the transcription factor NF- κ B (Refs 79,80). IL-1 β is also found in bronchoalveolar lavage fluid from ARDS patients, although its detrimental role in the disease is challenged by an in vitro study showing that IL-1 β contributes to the repairing process of injured alveolar epithelium (Ref. 90).

Ischaemia–reperfusion injury

I–R injury refers to tissue damage after blood supply returns to previously ischaemic areas, typically observed under the clinical conditions of myocardial infarction and organ transplantation. The injury is often accompanied by intracellular Ca²⁺ stress and accumulation of reactive oxygen species (ROS), which can disrupt endothelial cell–cell junctions and cause microvascular hyperpermeability (Refs 91,92). Furthermore, ROS has been implicated in the initiation and progress of the inflammatory response to I–R injury by upregulating leukotriene B₄, thromboxane A₂, and endothelial adhesion molecules that induce leukocyte activation and chemotaxis (Ref. 93). Also, oxidative stress increases VEGF production and VEGF receptor expression in endothelial cells (Ref. 94). Although it is commonly accepted that the upregulated VEGF signalling benefits collateral vessel formation after myocardial infarction, recent studies using animal models of I–R injury suggest that VEGF may aggravate pulmonary oedema and cerebral haemorrhagic transformation (Refs 95,96). In the gastrointestinal system following ischaemia, mucosal mast cells are activated by free radicals and subsequently release histamine, thus inducing microvascular permeability and leukocyte infiltration (Ref. 97). Another pathway leading to I–R injury is complement activation, which not only upregulates proinflammatory cytokines but also activates leukocytes and endothelial cells (Ref. 98). As a consequence, these cells release more oxidants, vasoactive factors and proteases (elastases and cathepsin G) that directly target the microvascular barrier resulting in plasma leakage and tissue oedema (Refs 99,100). Accumulated evidence further suggests that MMPs are also significantly upregulated in response to oxidative stress and contribute to I–R injury in multiple organs through increasing endothelial permeability (Ref. 101). Since MMPs have long been associated with degradation of extracellular matrix components, the reduced cell–matrix interaction (i.e. cell detachment) and subsequent apoptosis may partially confer microvascular barrier dysfunction.

Diabetes

Diabetes mellitus represents a group of metabolic disorders associated with defects in either insulin production or utilisation. Without proper treatments, diabetes results in a series of complications that affect multiple end organs, manifested as retinopathy, cardiomyopathy, nephropathy, and cerebral and peripheral vascular disease. Most of these problems are initiated with abnormal microcirculatory function and endothelial barrier injury; as an example, both

adherens junctions and tight junctions are diminished in the retinal and cerebral microvasculature of diabetic patients (Refs 102,103). Although the precise mechanism remains elusive, it is commonly believed that the injury is caused by polyol flux, oxidative stress, advanced glycation endproducts (AGEs) and upregulated DAG–PKC signalling, all stemming from hyperglycaemia. In diabetes, the adapted metabolism of excessive intracellular glucose results in overconsumption of nicotinamide adenine dinucleotide phosphate oxidase (NADPH) and glutathione, leading to oxidative stress that compromises vascular barrier function (Ref. 104). In parallel, AGEs at high concentration induce microvascular hyperpermeability by binding to their receptors on the endothelium as well as leukocytes (Refs 105,106), where they upregulate pro-inflammatory cytokines such as TNF- α and IL-1, thereby initiating multiple inflammatory cascades (Refs 15,107,108). AGEs may alternatively induce endothelial barrier damage by modifying extracellular matrix proteins (Ref. 109). Also, in hyperglycaemia increased de novo synthesis of DAG can activate the PKC pathway. Many PKC isoforms, including PKC α and β , promote endothelial permeability (Refs 110,111,112). Accordingly, selective inhibition of PKC β attenuates microvascular leakage in the retina, kidney and heart of diabetic patients and animals (Refs 15,112,113). Of particular interest, VEGF and its receptor VEGFR-2 are also upregulated during diabetes, which may play a critical role in the development of microvascular complications, especially diabetic retinopathy (Ref. 114).

Cancer

Tumour growth requires excessive supply of oxygen and nutrients and active removal of metabolic wastes. This extremely high level of blood–tissue exchange is supported by a dense microvasculature developed through pathological angiogenesis. Unlike other microvessels, tumour microvessels have thin walls, with defective and leaky endothelium, partially due to under developed endothelial cell–cell junctions and discontinuous smooth muscles and pericytes (Refs 115,116). The hyperpermeability property of the endothelium plays a crucial role in the initial development and continued growth of tumours as well as in tumour metastasis, as it facilitates cell transmigration and plasma accumulation in the matrix to support new vessel formation. This process is characterised by a highly orchestrated cellular response requiring interactions of multiple growth factors, adhesion molecules, and matrix proteins (Ref. 117). Typically, VEGF is considered a key signal for endothelial barrier breakdown that allows endothelial cell migration and matrix-supported capillary growth. In cancer tissues, especially breast cancer and other types of cancer, an excessive amount of VEGF is detected (Ref. 118) and its expression level correlates with the degree of tumour malignancy (Ref. 119), whereas inhibition of VEGF signalling suppresses tumour angiogenesis and malignant progression (Refs 120,121). Previously, it has been demonstrated that VEGF increases endothelial permeability and promotes angiogenesis via complex endothelial cell–cell and cell–matrix interactions (Ref. 122). Recent experiments with breast cancer cell and endothelial monolayer cocultures further indicate that matrix proteins, such as MMP-2, can be activated upon tumour cell attachment to the endothelium or after VEGF stimulation. This effect is accompanied by increased endothelial permeability and tumour cell transendothelial migration (Ref. 123).

Therapeutic implications

As described above, under the pathophysiological conditions of inflammation, microvascular hyperpermeability is elicited by circulating mediators and growth factors that can bind to their endothelial receptors and trigger further signalling reactions in the barrier structure. Selective blockades of proinflammatory agents have been used in patients with trauma and inflammatory injury. Further inhibition of their downstream signalling or terminal effectors represents a promising area of drug development for prevention and treatment of vascular disease. Below are examples of potential therapies relevant to vascular endothelial hyperpermeability.

Antihistamine therapy

Histamine receptor antagonists have been widely used in the treatment of allergic reactions. Most of them competitively bind H₁ receptors, thereby blocking histamine-induced hypersensitive responses and endothelial hyperpermeability. In animal models, the H₁ receptor antagonist diphenhydramine reduces lung microvascular permeability and ameliorates pulmonary oedema (Ref. 124). Another histamine antagonist, loratadine, exerts protective effects on microvascular barrier function during I-R injury (Ref. 97). When administered simultaneously, the H₁ receptor antagonist diphenhydramine and the H₂ receptor antagonist cimetidine prevent I-R-induced leukocyte infiltration (Ref. 125). Clinically, however, single blockades of histamine receptors often demonstrate a limited utility for they do not block the complex interactions and redundant effects of multiple inflammatory agents and pathways elaborated under injurious conditions. In the cases of severe anaphylaxis and angio-oedema, glucocorticoids are usually used in combination with antihistamine drugs. Glucocorticoids not only suppress T helper 1 (Th1)-mediated cellular immunity and systemic inflammatory response but also inhibit histamine release from mast cells (Refs 126,127).

Anti-VEGF therapy

Given the importance of VEGF in angiogenesis-associated diseases, such as tumourigenesis and diabetic retinopathy, a considerable research effort has been devoted to developing novel strategies that inhibit the VEGF pathway. Current products mainly include anti-VEGF antibodies, VEGF traps (genetically engineered soluble VEGF receptors), and VEGF receptor blockades. Animal experiments demonstrate that VEGF neutralising antibodies can suppress microvascular permeability and angiogenesis in tumours, thereby reducing tumour size (Ref. 121). Bevacizumab, the first anti-VEGF agent for cancer therapy approved by the US Food and Drug Administration, not only inhibits tumour growth in a dose-dependent fashion (Ref. 128) but also produces an antimetastatic effect (Refs 120,129). This drug has been used along with chemotherapy in patients with lung, renal or colorectal cancer. In addition to cancer therapy, anti-VEGF strategies have proven beneficial in other diseases or disorders associated with microvascular hyperpermeability. For example, in experimental diabetic retinopathy, administration of anti-VEGF antibodies or VEGF traps significantly reduces the abnormal growth of microvessels and hyperpermeability in the retina (Refs 130,131). In patients with diabetic macular oedema, the therapeutic effect of anti-VEGF agents has been tested using fluorescein angiograph and optical coherence tomography (Ref. 132). Subsequently, several VEGF receptor antagonists have been developed and are currently under preclinical or clinical trials. They act either as pan-inhibitors of VEGF receptors (e.g. pazopanib), or as specific inhibitors of VEGFR-2 (e.g. ZM323881). In animal studies, ZM323881 effectively blocks the acute vascular leak response caused by VEGF (Refs 132,133). In a Phase I trial, pazopanib demonstrates an inhibitory effect on tumour growth in patients with hypernephroma.

In addition to the strategies specifically targeting VEGF/VEGFR-2, a barrier-protective effect against VEGF-induced damage can derive from other approaches, including administration of angiopoietin-1 and inhibition of PLC, PKC, PKG, intracellular Ca²⁺, MEK and Src (Refs 14, 28,52,134). Although both angiopoietin-1 and VEGF are proangiogenic factors, angiopoietin-1 promotes vessel maturation and inhibits VEGF-induced endothelial hyperpermeability. A recent study suggests that the protective effect mediated by angiopoietin-1 is through inhibition of Src activation (Ref. 135). In support of this, it has been demonstrated that VEGF-induced vascular barrier dysfunction is specifically blocked in Src-deficient mice, with sparing of normal angiogenesis (Ref. 32). Further evidence shows that topical application of a novel VEGFR-2/Src-kinase inhibitor suppresses VEGF-mediated retinal vascular hyperpermeability in animal models (Ref. 136).

Activated protein C

Activated protein C (APC) is an endogenous anticoagulation factor that has recently been shown to possess anti-inflammatory and anti-apoptotic functions (Refs 137,138). Recombinant APC [DrotAA; drotrecogin-alpha (activated)] has been used in critical care for patients with severe sepsis and disseminated intravascular coagulation. The mechanism underlying the dual functions of APC is a subject of current investigation. It is generally accepted that the beneficial effect of APC is mediated by endothelial protein C receptor (EPCR)-dependent cleavage of PAR-1 on the vascular endothelial surface (Refs 138,139,140). Binding of APC to EPCR switches the PAR-1 signalling downstream of thrombin to a protective pathway through coupling of PAR-1 to a G_i protein (Refs 139,140). It has been shown that soluble APC-EPCR complexes with proteinase-3 and integrins expressed on leukocytes (Ref. 141), and that APC downregulates adhesion molecules such as ICAM-1, VCAM-1 and E-selectin expressed on endothelial cells, thereby inhibiting vascular inflammation (Refs 142,143). In animals, DrotAA significantly attenuates smoke-induced lung microvascular hyperpermeability (Ref. 144). It also ameliorates pulmonary oedema caused by I-R injury (Ref. 145).

PKC inhibitors

In light of the fact that PKC plays a central role in the endothelial response to various proinflammatory factors, targeting this common signalling molecule with site specificity may become an effective adjunct therapy for treating vascular inflammatory injury. In fact, it has been well documented that PKC inhibitors significantly reduce microvascular hyperpermeability caused by cytokines, vasoactive agents and growth factors (Ref. 28). In diabetic patients and animals, oral administration of PKC β inhibitors (e.g. LY333531 or Ruboxistaurin) prevents microvascular dysfunction and delays the progress of retinopathy and nephropathy (Refs 146,147). In a series of studies with porcine models of diabetes, it was demonstrated that PKC expression and enzymatic activity (especially that of the β II isoform) was upregulated in the heart and coronary system during early stages of diabetes. The abnormalities correlate with increased coronary microvascular permeability (Refs 112,113), which is significantly attenuated by PKC β inhibitors (Ref. 112). These results support the therapeutic potential of targeting PKC β for treating diabetic microvascular complications.

Rho inhibitors

Similar to PKCs, the RhoA-ROCK pathway has been frequently implicated in the pathological progression of endothelial barrier dysfunction associated with ARDS, I-R injury and diabetes mellitus. In experimental models of ARDS, systemic administration of a ROCK inhibitor, Y-27632, substantially reduces pulmonary microvascular permeability and lung injury (Refs 75,148). Fasudil, a ROCK inhibitor, has been evaluated in clinical trials for treating pulmonary hypertension, cerebral vasospasm, and angina (Ref. 149). A recent study demonstrates that fasudil inhibits VEGF-induced angiogenesis (Ref. 150). In diabetic animals, fasudil ameliorates microvascular injury associated with retinopathy and nephropathy (Refs 151, 152). In addition, some drugs that are clinically used for treating hypertension and hypercholesterolaemia have been shown to protect vascular endothelial barrier function through RhoA inhibition. For example, simvastatin can decrease thrombin-induced endothelial dysfunction by attenuating RhoA activation and stress-fibre formation (Ref. 153). Moreover, simvastatin reduces VEGF-induced glomerular endothelial hyperpermeability (Ref. 154). However, compared with other selective inhibitors of RhoA signalling such as Y-27632, statins do not seem to deliver any immediate or rapid barrier protection and their downstream effectors are relatively nonspecific as a result of a broad targeting of multiple Rho GTPases.

Future directions

Microvascular barrier dysfunction is a significant problem confronted in the clinical treatments for various diseases. In vitro and animal studies in this field have led to a better understanding of the molecular mechanisms underlying endothelial hyperpermeability, and corresponding strategies have been developed for clinical applications as described in this review. However, many puzzles still remain to be elucidated. Of particular significance is the fact that although several mediators (such as VEGF, histamine and thrombin) share similar signalling cascades in triggering endothelial hyperpermeability, their temporal effects and the permeability-recovery rates are rather different. This heterogeneity may be attributed to differential regulation of binding, trafficking, internalisation, and/or desensitisation of their receptors on the plasma membrane. It is also possible that different levels of counter-regulatory signalling (e.g. cAMP) are activated concurrently with the hyperpermeability signalling. Furthermore, despite the efficacy of antagonising individual mediator-induced hyperpermeability in certain pathological states, only moderate effectiveness is demonstrated under more complicated inflammatory conditions where multiple mediators are involved and interact with each other. Therefore, targeting their common terminal effectors (e.g. RhoA/ROCK) may represent a promising strategy against vascular injury. Within this context, site-specific drug delivery needs to be developed. A good strategy should improve the barrier function in the affected vascular bed while sparing other tissues from unwanted effects. A similar issue applies to treating cancers, where microvascular hyperpermeability may favour the delivery of chemotherapeutic drugs. Research efforts to further understand endothelial barrier structure and function should ultimately lead to improved treatment of vascular inflammation.

Acknowledgments

This work is supported by the NIH grants HL61507, HL70752, HL73324 and HL84542. We thank Ms Olesya Litovka for excellent artwork and assistance in the manuscript preparation. We also acknowledge the peer reviewers for their valuable comments.

References

1. Bauer PM, et al. Endothelial-specific expression of caveolin-1 impairs microvascular permeability and angiogenesis. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:204–209. [PubMed: 15615855]
2. Gratton JP, Bernatchez P, Sessa WC. Caveolae and caveolins in the cardiovascular system. *Circulation Research* 2004;94:1408–1417. [PubMed: 15192036]
3. Lin MI, et al. Caveolin-1-deficient mice have increased tumor microvascular permeability, angiogenesis, and growth. *Cancer Research* 2007;67:2849–2856. [PubMed: 17363608]
4. Razani B, et al. Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *Journal of Biological Chemistry* 2001;276:38121–38138. [PubMed: 11457855]
5. Mehta D, Malik AB. Signaling mechanisms regulating endothelial permeability. *Physiological Reviews* 2006;86:279–367. [PubMed: 16371600]
6. Dvorak AM, et al. The vesiculo-vacuolar organelle (VVO): a distinct endothelial cell structure that provides a transcellular pathway for macromolecular extravasation. *Journal of Leukocyte Biology* 1996;59:100–115. [PubMed: 8558058]
7. Curry FE, Adamson RH. Transendothelial pathways in venular microvessels exposed to agents which increase permeability: the gaps in our knowledge. *Microcirculation* 1999;6:3–5. [PubMed: 10100185]
8. Dvorak AM. Mast cell-derived mediators of enhanced microvascular permeability, vascular permeability factor/vascular endothelial growth factor, histamine, and serotonin, cause leakage of macromolecules through a new endothelial cell permeability organelle, the vesiculo-vacuolar organelle. *Chemical Immunology and Allergy* 2005;85:185–204. [PubMed: 15970658]

9. Iyer S, et al. VE-cadherin-p120 interaction is required for maintenance of endothelial barrier function. *American Journal of Physiology – Lung Cellular and Molecular Physiology* 2004;286:L1143–1153. [PubMed: 14672921]
10. Xiao K, et al. p120-Catenin regulates clathrin-dependent endocytosis of VE-cadherin. *Molecular Biology of the Cell* 2005;16:5141–5151. [PubMed: 16120645]
11. Esser S, et al. Vascular endothelial growth factor induces VE-cadherin tyrosine phosphorylation in endothelial cells. *Journal of Cell Science* 1998;111:1853–1865. [PubMed: 9625748]
12. Dejana E. Endothelial cell-cell junctions: happy together. *Nature Reviews Molecular Cell Biology* 2004;5:261–270.
13. Tiruppathi C, et al. Ca²⁺ signaling, TRP channels, and endothelial permeability. *Microcirculation* 2006;13:693–708. [PubMed: 17085428]
14. Yuan SY. Protein kinase signaling in the modulation of microvascular permeability. *Vascular Pharmacology* 2002;39:213–223. [PubMed: 12747961]
15. Yuan SY, et al. Microvascular permeability in diabetes and insulin resistance. *Microcirculation* 2007;14:363–373. [PubMed: 17613808]
16. Isshiki M, et al. Endothelial Ca²⁺ waves preferentially originate at specific loci in caveolin-rich cell edges. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:5009–5014. [PubMed: 9560219]
17. Maekawa M, et al. Signaling from Rho to the actin cytoskeleton through protein kinases ROCK and LIM-kinase. *Science* 1999;285:895–898. [PubMed: 10436159]
18. Mehta D, Rahman A, Malik AB. Protein kinase C- α signals rho-guanine nucleotide dissociation inhibitor phosphorylation and rho activation and regulates the endothelial cell barrier function. *Journal of Biological Chemistry* 2001;276:22614–22620. [PubMed: 11309397]
19. Verin AD, et al. Role of ras-dependent ERK activation in phorbol ester-induced endothelial cell barrier dysfunction. *American Journal of Physiology – Lung Cellular and Molecular Physiology* 2000;279:L360–370. [PubMed: 10926560]
20. Stasek JE Jr, Patterson CE, Garcia JG. Protein kinase C phosphorylates caldesmon77 and vimentin and enhances albumin permeability across cultured bovine pulmonary artery endothelial cell monolayers. *Journal of Cellular Physiology* 1992;153:62–75. [PubMed: 1522136]
21. Sandoval R, et al. Ca(2+) signalling and PKC α activate increased endothelial permeability by disassembly of VE-cadherin junctions. *Journal of Physiology* 2001;533:433–445. [PubMed: 11389203]
22. Clarke H, Soler AP, Mullin JM. Protein kinase C activation leads to dephosphorylation of occludin and tight junction permeability increase in LLC-PK1 epithelial cell sheets. *Journal of Cell Science* 2000;113(Pt 18):3187–3196. [PubMed: 10954417]
23. Aramoto H, et al. Vascular endothelial growth factor stimulates differential signaling pathways in vivo microcirculation. *American Journal of Physiology – Heart and Circulatory Physiology* 2004;287:H1590–1598. [PubMed: 15155260]
24. Bates DO, Curry FE. Vascular endothelial growth factor increases microvascular permeability via a Ca(2+)-dependent pathway. *American Journal of Physiology* 1997;273:H687–694. [PubMed: 9277485]
25. Breslin JW, et al. VEGF increases endothelial permeability by separate signaling pathways involving ERK-1/2 and nitric oxide. *American Journal of Physiology – Heart and Circulatory Physiology* 2003;284:H92–H100. [PubMed: 12388327]
26. Lal BK, et al. VEGF increases permeability of the endothelial cell monolayer by activation of PKB/akt, endothelial nitric-oxide synthase, and MAP kinase pathways. *Microvascular Research* 2001;62:252–262. [PubMed: 11678628]
27. Varma S, et al. p42/44MAPK regulates baseline permeability and cGMP-induced hyperpermeability in endothelial cells. *Microvascular Research* 2002;63:172–178. [PubMed: 11866540]
28. Wu HM, et al. Role of phospholipase C, protein kinase C, and calcium in VEGF-induced venular hyperpermeability. *American Journal of Physiology* 1999;276:H535–542. [PubMed: 9950855]
29. Stockton RA, Schaefer E, Schwartz MA. p21-activated kinase regulates endothelial permeability through modulation of contractility. *Journal of Biological Chemistry* 2004;279:46621–46630. [PubMed: 15333633]

30. Wu MH, et al. Focal adhesion kinase mediates porcine venular hyperpermeability elicited by vascular endothelial growth factor. *Journal of Physiology* 2003;552:691–699. [PubMed: 12949227]
31. Gavard J, Gutkind JS. VEGF controls endothelial-cell permeability by promoting the beta-arrestin-dependent endocytosis of VE-cadherin. *Nature Cell Biology* 2006;8:1223–1234.
32. Eliceiri BP, et al. Selective requirement for Src kinases during VEGF-induced angiogenesis and vascular permeability. *Molecular Cell* 1999;4:915–924. [PubMed: 10635317]
33. Birukov KG, et al. Differential regulation of alternatively spliced endothelial cell myosin light chain kinase isoforms by p60(Src). *Journal of Biological Chemistry* 2001;276:8567–8573. [PubMed: 11113114]
34. Wu MH, Ustinova E, Granger HJ. Integrin binding to fibronectin and vitronectin maintains the barrier function of isolated porcine coronary venules. *Journal of Physiology* 2001;532:785–791. [PubMed: 11313446]
35. Shajahan AN, et al. Role of Src-induced dynamin-2 phosphorylation in caveolae-mediated endocytosis in endothelial cells. *Journal of Biological Chemistry* 2004;279:20392–20400. [PubMed: 15007081]
36. Miao H, et al. Differential regulation of Rho GTPases by beta1 and beta3 integrins: the role of an extracellular domain of integrin in intracellular signaling. *Journal of Cell Science* 2002;115:2199–2206. [PubMed: 11973360]
37. Mammoto A, Mammoto T, Ingber DE. Rho signaling and mechanical control of vascular development. *Current Opinion in Hematology* 2008;15:228–234. [PubMed: 18391790]
38. Breslin JW, et al. Involvement of ROCK-mediated endothelial tension development in neutrophil-stimulated microvascular leakage. *American Journal of Physiology – Heart and Circulatory Physiology* 2006;290:H741–750. [PubMed: 16172166]
39. Breslin JW, Yuan SY. Involvement of RhoA and Rho kinase in neutrophil-stimulated endothelial hyperpermeability. *American Journal of Physiology – Heart and Circulatory Physiology* 2004;286:H1057–1062. [PubMed: 14630629]
40. Sun H, et al. Rho and ROCK signaling in VEGF-induced microvascular endothelial hyperpermeability. *Microcirculation* 2006;13:237–247. [PubMed: 16627366]
41. van Nieuw Amerongen GP, et al. Activation of RhoA by thrombin in endothelial hyperpermeability: role of Rho kinase and protein tyrosine kinases. *Circulation Research* 2000;87:335–340. [PubMed: 10948069]
42. Patil SB, Bitar KN. RhoA- and PKC-alpha-mediated phosphorylation of MYPT and its association with HSP27 in colonic smooth muscle cells. *American Journal of Physiology – Gastrointestinal and Liver Physiology* 2006;290:G83–95. [PubMed: 16179599]
43. Garcia JG, et al. Sphingosine 1-phosphate promotes endothelial cell barrier integrity by Edg-dependent cytoskeletal rearrangement. *Journal of Clinical Investigation* 2001;108:689–701. [PubMed: 11544274]
44. Liu F, et al. Hepatocyte growth factor enhances endothelial cell barrier function and cortical cytoskeletal rearrangement: potential role of glycogen synthase kinase-3beta. *FASEB Journal* 2002;16:950–962. [PubMed: 12087056]
45. Kouklis P, et al. Cdc42 regulates the restoration of endothelial barrier function. *Circulation Research* 2004;94:159–166. [PubMed: 14656933]
46. Moore TM, et al. Signal transduction and regulation of lung endothelial cell permeability. Interaction between calcium and cAMP. *American Journal of Physiology* 1998;275:L203–222. [PubMed: 9700080]
47. Lampugnani MG, et al. The role of integrins in the maintenance of endothelial monolayer integrity. *Journal of Cell Biology* 1991;112:479–490. [PubMed: 1899416]
48. Hill SJ, et al. International Union of Pharmacology. XIII. Classification of histamine receptors. *Pharmacological Reviews* 1997;49:253–278. [PubMed: 9311023]
49. Parsons ME, Ganellin CR. Histamine and its receptors. *British Journal of Pharmacology* 2006;147 (Suppl 1):S127–135. [PubMed: 16402096]
50. Clough GF, Bennett AR, Church MK. Effects of H1 antagonists on the cutaneous vascular response to histamine and bradykinin: a study using scanning laser Doppler imaging. *British Journal of Dermatology* 1998;138:806–814. [PubMed: 9666826]

51. Yuan SY. Signal transduction pathways in enhanced microvascular permeability. *Microcirculation* 2000;7:395–403. [PubMed: 11142336]
52. Wu MH, Yuan SY, Granger HJ. The protein kinase MEK1/2 mediate vascular endothelial growth factor- and histamine-induced hyperpermeability in porcine coronary venules. *Journal of Physiology* 2005;563:95–104. [PubMed: 15539400]
53. Guo M, et al. VE-cadherin and beta-catenin binding dynamics during histamine-induced endothelial hyperpermeability. *American Journal of Physiology - Cell Physiology* 2008;294:C977–984. [PubMed: 18287330]
54. Shasby DM, et al. Histamine stimulates phosphorylation of adherens junction proteins and alters their link to vimentin. *American Journal of Physiology – Lung Cellular and Molecular Physiology* 2002;282:L1330–1338. [PubMed: 12003790]
55. Bogatcheva NV, Garcia JG, Verin AD. Molecular mechanisms of thrombin-induced endothelial cell permeability. *Biochemistry* 2002;67:75–84. [PubMed: 11841342]
56. Borbiev T, et al. Role of CaM kinase II and ERK activation in thrombin-induced endothelial cell barrier dysfunction. *American Journal of Physiology – Lung Cellular and Molecular Physiology* 2003;285:L43–54. [PubMed: 12788788]
57. Birukova AA, et al. Novel role of microtubules in thrombin-induced endothelial barrier dysfunction. *FASEB Journal* 2004;18:1879–1890. [PubMed: 15576491]
58. Holinstat M, et al. Protein kinase C α -induced p115RhoGEF phosphorylation signals endothelial cytoskeletal rearrangement. *Journal of Biological Chemistry* 2003;278:28793–28798. [PubMed: 12754211]
59. Grand RJ, Turnell AS, Grabham PW. Cellular consequences of thrombin-receptor activation. *The Biochemical Journal* 1996;313:353–368. [PubMed: 8573065]
60. Senger DR, et al. Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 1983;219:983–985. [PubMed: 6823562]
61. Weis SM, Cheresh DA. Pathophysiological consequences of VEGF-induced vascular permeability. *Nature* 2005;437:497–504. [PubMed: 16177780]
62. Takahashi H, et al. A novel snake venom vascular endothelial growth factor (VEGF) predominantly induces vascular permeability through preferential signaling via VEGF receptor-1. *Journal of Biological Chemistry* 2004;279:46304–46314. [PubMed: 15328352]
63. Zachary I. VEGF signalling: integration and multi-tasking in endothelial cell biology. *Biochemical Society Transactions* 2003;31:1171–1177. [PubMed: 14641020]
64. Behzadian MA, et al. VEGF-induced paracellular permeability in cultured endothelial cells involves urokinase and its receptor. *FASEB Journal* 2003;17:752–754. [PubMed: 12594181]
65. Kelly M, Hwang JM, Kubes P. Modulating leukocyte recruitment in inflammation. *Journal of Allergy and Clinical Immunology* 2007;120:3–10. [PubMed: 17559914]
66. Kubes P. The complexities of leukocyte recruitment. *Seminars in Immunology* 2002;14:65–72. [PubMed: 11978078]
67. Liu Y, et al. Regulation of leukocyte transmigration: cell surface interactions and signaling events. *Journal of Immunology* 2004;172:7–13.
68. Rao RM, et al. Emerging topics in the regulation of leukocyte transendothelial migration. *Microcirculation* 2005;12:83–89. [PubMed: 15804976]
69. Rao RM, et al. Endothelial-dependent mechanisms of leukocyte recruitment to the vascular wall. *Circulation Research* 2007;101:234–247. [PubMed: 17673684]
70. Rosenkranz AR, Mayadas TN. Leukocyte-endothelial cell interactions - lessons from knockout mice. *Experimental Nephrology* 1999;7:125–136. [PubMed: 10213866]
71. Zarbock A, Ley K. Mechanisms and consequences of neutrophil interaction with the endothelium. *American Journal of Pathology* 2008;172:1–7.
72. Sarelius IH, et al. Macromolecule permeability of in situ and excised rodent skeletal muscle arterioles and venules. *American Journal of Physiology – Heart and Circulatory Physiology* 2006;290:H474–480. [PubMed: 16126813]
73. Yuan SY, et al. Myosin light chain phosphorylation in neutrophil-stimulated coronary microvascular leakage. *Circulation Research* 2002;90:1214–1221. [PubMed: 12065325]

74. Tinsley JH, et al. Src-dependent, neutrophil-mediated vascular hyperpermeability and beta-catenin modification. *American Journal of Physiology – Cell Physiology* 2002;283:C1745–1751. [PubMed: 12388068]
75. Chiba Y, et al. Activation of rho is involved in the mechanism of hydrogen-peroxide-induced lung edema in isolated perfused rabbit lung. *Microvascular Research* 2001;62:164–171. [PubMed: 11516245]
76. Sumagin R, Lomakina E, Sarelius IH. Leukocyte-endothelial cell interactions are linked to vascular permeability via ICAM-1-mediated signaling. *American Journal of Physiology – Heart and Circulatory Physiology* 2008;295:H969–H977. [PubMed: 18641276]
77. Usatyuk PV, et al. Role of Ca²⁺ in diperoxovanadate-induced cytoskeletal remodeling and endothelial cell barrier function. *American Journal of Physiology – Lung Cellular and Molecular Physiology* 2003;285:L1006–1017. [PubMed: 12882766]
78. Usatyuk PV, et al. Redox regulation of reactive oxygen species-induced p38 MAP kinase activation and barrier dysfunction in lung microvascular endothelial cells. *Antioxidants and Redox Signalling* 2003;5:723–730.
79. Blackwell TS, Christman JW. The role of nuclear factor-kappa B in cytokine gene regulation. *American Journal of Respiratory Cell and Molecular Biology* 1997;17:3–9. [PubMed: 9224203]
80. Reutershan J, Ley K. Bench-to-bedside review: acute respiratory distress syndrome - how neutrophils migrate into the lung. *Critical Care* 2004;8:453–461. [PubMed: 15566616]
81. Hu G, et al. Intercellular adhesion molecule-1-dependent neutrophil adhesion to endothelial cells induces caveolae-mediated pulmonary vascular hyperpermeability. *Circulation Research* 2008;102:e120–131. [PubMed: 18511851]
82. Bhatia M, Moochhala S. Role of inflammatory mediators in the pathophysiology of acute respiratory distress syndrome. *Journal of Pathology* 2004;202:145–156. [PubMed: 14743496]
83. Groeneveld AB. Vascular pharmacology of acute lung injury and acute respiratory distress syndrome. *Vascular Pharmacology* 2002;39:247–256. [PubMed: 12747964]
84. Hinshaw LB, Jordan MM, Vick JA. Histamine release and endotoxin shock in the primate. *Journal of Clinical Investigation* 1961;40:1631–1637. [PubMed: 13714470]
85. Nagy S. The role of histamine release in shock. *Acta Physiologica Hungarica* 1990;76:3–12. [PubMed: 1708189]
86. Breil I, et al. Effects of bradykinin, histamine and serotonin on pulmonary vascular resistance and permeability. *Acta Physiologica Scandinavica* 1997;159:189–198. [PubMed: 9079149]
87. Bauer TT, et al. Comparison of systemic cytokine levels in patients with acute respiratory distress syndrome, severe pneumonia, and controls. *Thorax* 2000;55:46–52. [PubMed: 10607801]
88. Hamacher J, et al. Tumor necrosis factor-alpha and angiostatin are mediators of endothelial cytotoxicity in bronchoalveolar lavages of patients with acute respiratory distress syndrome. *American Journal of Respiratory and Critical Care Medicine* 2002;166:651–656. [PubMed: 12204860]
89. Windsor AC, et al. Tumor necrosis factor-alpha blockade prevents neutrophil CD18 receptor upregulation and attenuates acute lung injury in porcine sepsis without inhibition of neutrophil oxygen radical generation. *Journal of Clinical Investigation* 1993;91:1459–1468. [PubMed: 8097206]
90. Geiser T, et al. Pulmonary edema fluid from patients with acute lung injury augments in vitro alveolar epithelial repair by an IL-1beta-dependent mechanism. *American Journal of Respiratory and Critical Care Medicine* 2001;163:1384–1388. [PubMed: 11371405]
91. Gilmont RR, et al. Effects of glutathione depletion on oxidant-induced endothelial cell injury. *Journal of Surgical Research* 1998;80:62–68. [PubMed: 9790816]
92. Wang Q, et al. Ethanol preconditioning protects against ischemia/reperfusion-induced brain damage: role of NADPH oxidase-derived ROS. *Free Radical Biology and Medicine* 2007;43:1048–1060. [PubMed: 17761301]
93. Aktan A, Yalcin A. Ischemia-reperfusion injury, reactive oxygen metabolites, and the Surgeon. *Turkish Journal of Medical Sciences* 1998;28:1–5.
94. Chua CC, Hamdy RC, Chua BH. Upregulation of vascular endothelial growth factor by H₂O₂ in rat heart endothelial cells. *Free Radical Biology and Medicine* 1998;25:891–897. [PubMed: 9840733]

95. Abumiya T, et al. Aggravation of hemorrhagic transformation by early intraarterial infusion of low-dose vascular endothelial growth factor after transient focal cerebral ischemia in rats. *Brain Research* 2005;1049:95–103. [PubMed: 15935998]
96. Godzich M, et al. Activation of the stress protein response prevents the development of pulmonary edema by inhibiting VEGF cell signaling in a model of lung ischemia-reperfusion injury in rats. *FASEB Journal* 2006;20:1519–1521. [PubMed: 16793871]
97. Nosal'ova V, et al. Effect of H1 antihistamines in a model of mesenteric ischaemia/reperfusion. *Inflammation Research* 2008;57(Suppl 1):S55–56. [PubMed: 18345491]
98. Collard CD, et al. Complement activation following oxidative stress. *Molecular Immunology* 1999;36:941–948. [PubMed: 10698348]
99. Carden DL, Granger DN. Pathophysiology of ischaemia-reperfusion injury. *Journal of Pathology* 2000;190:255–266. [PubMed: 10685060]
100. Hirahashi J, et al. Mac-1 signaling via Src-family and Syk kinases results in elastase-dependent thrombohemorrhagic vasculopathy. *Immunity* 2006;25:271–283. [PubMed: 16872848]
101. Alexander JS, Elrod JW. Extracellular matrix, junctional integrity and matrix metalloproteinase interactions in endothelial permeability regulation. *Journal of Anatomy* 2002;200:561–574. [PubMed: 12162724]
102. Chehade JM, Haas MJ, Mooradian AD. Diabetes-related changes in rat cerebral occludin and zonula occludens-1 (ZO-1) expression. *Neurochemical Research* 2002;27:249–252. [PubMed: 11958524]
103. Davidson MK, et al. Reduced expression of the adherens junction protein cadherin-5 in a diabetic retina. *American Journal of Ophthalmology* 2000;129:267–269. [PubMed: 10682990]
104. Lee AY, Chung SS. Contributions of polyol pathway to oxidative stress in diabetic cataract. *FASEB Journal* 1999;13:23–30. [PubMed: 9872926]
105. Bonnardel-Phu E, et al. Acute modulation of albumin microvascular leakage by advanced glycation end products in microcirculation of diabetic rats in vivo. *Diabetes* 1999;48:2052–2058. [PubMed: 10512373]
106. Otero K, et al. Albumin-derived advanced glycation end-products trigger the disruption of the vascular endothelial cadherin complex in cultured human and murine endothelial cells. *The Biochemical Journal* 2001;359:567–574. [PubMed: 11672430]
107. Schmidt AM, et al. The biology of the receptor for advanced glycation end products and its ligands. *Biochimica et Biophysica Acta* 2000;1498:99–111. [PubMed: 11108954]
108. Yamagishi S, et al. Advanced glycation end product-induced apoptosis and overexpression of vascular endothelial growth factor and monocyte chemoattractant protein-1 in human-cultured mesangial cells. *Journal of Biological Chemistry* 2002;277:20309–20315. [PubMed: 11912219]
109. Brownlee M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 2005;54:1615–1625. [PubMed: 15919781]
110. Harrington EO, et al. Role of protein kinase C isoforms in rat epididymal microvascular endothelial barrier function. *American Journal of Respiratory Cell and Molecular Biology* 2003;28:626–636. [PubMed: 12707019]
111. Tinsley JH, Teasdale NR, Yuan SY. Involvement of PKCdelta and PKD in pulmonary microvascular endothelial cell hyperpermeability. *American Journal of Physiology – Cell Physiology* 2004;286:C105–111. [PubMed: 13679307]
112. Yuan SY, et al. Protein kinase C activation contributes to microvascular barrier dysfunction in the heart at early stages of diabetes. *Circulation Research* 2000;87:412–417. [PubMed: 10969040]
113. Guo M, et al. Upregulation of PKC genes and isozymes in cardiovascular tissues during early stages of experimental diabetes. *Physiological Genomics* 2003;12:139–146. [PubMed: 12441406]
114. Caldwell RB, et al. Vascular endothelial growth factor and diabetic retinopathy: role of oxidative stress. *Current Drug Targets* 2005;6:511–524. [PubMed: 16026270]
115. Hashizume H, et al. Openings between defective endothelial cells explain tumor vessel leakiness. *American Journal of Pathology* 2000;156:1363–1380. [PubMed: 10751361]
116. Morikawa S, et al. Abnormalities in pericytes on blood vessels and endothelial sprouts in tumors. *American Journal of Pathology* 2002;160:985–1000. [PubMed: 11891196]

117. Piulats, J.; Mitjans, F. Angiogenesis switch pathways. In: Bronchud, MH., et al., editors. Principles of Molecular Oncology. 3rd edn. Humana Press; Totowa, NJ, USA: 2008. p. 239-256.
118. Anan K, et al. Vascular endothelial growth factor and platelet-derived growth factor are potential angiogenic and metastatic factors in human breast cancer. *Surgery* 1996;119:333-339. [PubMed: 8619189]
119. Dvorak HF, et al. Vascular permeability factor/vascular endothelial growth factor, microvascular hyperpermeability, and angiogenesis. *American Journal of Pathology* 1995;146:1029-1039. [PubMed: 7538264]
120. Lee FY, et al. Synergistic antitumor activity of ixabepilone (BMS-247550) plus bevacizumab in multiple in vivo tumor models. *Clinical Cancer Research* 2008;14:8123-8131. [PubMed: 19088027]
121. Yuan F, et al. Time-dependent vascular regression and permeability changes in established human tumor xenografts induced by an anti-vascular endothelial growth factor/vascular permeability factor antibody. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:14765-14770. [PubMed: 8962129]
122. Wu MH. Endothelial focal adhesions and barrier function. *Journal of Physiology* 2005;569:359-366. [PubMed: 16195317]
123. Kargozaran H, et al. A role for endothelial-derived matrix metalloproteinase-2 in breast cancer cell transmigration across the endothelial-basement membrane barrier. *Clinical and Experimental Metastasis* 2007;24:495-502. [PubMed: 17653824]
124. Brigham KL, Bowers RE, Owen PJ. Effects of antihistamines on the lung vascular response to histamine in unanesthetized sheep. Diphenhydramine prevention of pulmonary edema and increased permeability. *Journal of Clinical Investigation* 1976;58:391-398. [PubMed: 956373]
125. Cordeiro PG, et al. Prevention of ischemia-reperfusion injury in a rat skin flap model: the role of mast cells, cromolyn sodium, and histamine receptor blockade. *Plastic and Reconstructive Surgery* 2000;105:654-659. [PubMed: 10697173]
126. Elenkov IJ. Glucocorticoids and the Th1/Th2 balance. *Annals of the New York Academy of Sciences* 2004;1024:138-146. [PubMed: 15265778]
127. Svedmyr N. Effects of glucocorticoids on the airways. *European Journal of Respiratory Diseases (Suppl)* 1982;122:48-53.
128. Gerber HP, et al. Complete inhibition of rhabdomyosarcoma xenograft growth and neovascularization requires blockade of both tumor and host vascular endothelial growth factor. *Cancer Research* 2000;60:6253-6258. [PubMed: 11103779]
129. Zhang J, et al. Elevated expression of vascular endothelial growth factor correlates with increased angiogenesis and decreased progression-free survival among patients with low-grade neuroendocrine tumors. *Cancer* 2007;109:1478-1486. [PubMed: 17340592]
130. Aiello LP, et al. Suppression of retinal neovascularization in vivo by inhibition of vascular endothelial growth factor (VEGF) using soluble VEGF-receptor chimeric proteins. *Proceedings of the National Academy of Sciences of the United States of America* 1995;92:10457-10461. [PubMed: 7479819]
131. Robinson GS, et al. Nonvascular role for VEGF: VEGFR-1, 2 activity is critical for neural retinal development. *FASEB Journal* 2001;15:1215-1217. [PubMed: 11344092]
132. Haritoglou C, et al. Intravitreal bevacizumab (Avastin) therapy for persistent diffuse diabetic macular edema. *Retina* 2006;26:999-1005. [PubMed: 17151486]
133. Whittles CE, et al. ZM323881, a novel inhibitor of vascular endothelial growth factor-receptor-2 tyrosine kinase activity. *Microcirculation* 2002;9:513-522. [PubMed: 12483548]
134. van Nieuw Amerongen GP, van Hinsbergh VW. Targets for pharmacological intervention of endothelial hyperpermeability and barrier function. *Vascular Pharmacology* 2002;39:257-272. [PubMed: 12747965]
135. Gavard J, Patel V, Gutkind JS. Angiopoietin-1 prevents VEGF-induced endothelial permeability by sequestering Src through mDia. *Developmental Cell* 2008;14:25-36. [PubMed: 18194650]
136. Schepcke L, et al. Retinal vascular permeability suppression by topical application of a novel VEGFR2/Src kinase inhibitor in mice and rabbits. *Journal of Clinical Investigation* 2008;118:2337-2346. [PubMed: 18483622]

137. Dhainaut JF, Yan SB, Claessens YE. Protein C/activated protein C pathway: overview of clinical trial results in severe sepsis. *Critical Care Medicine* 2004;32:S194–201. [PubMed: 15118517]
138. Riewald M, et al. Activation of endothelial cell protease activated receptor 1 by the protein C pathway. *Science* 2002;296:1880–1882. [PubMed: 12052963]
139. Bae JS, et al. The ligand occupancy of endothelial protein C receptor switches the protease-activated receptor 1-dependent signaling specificity of thrombin from a permeability-enhancing to a barrier-protective response in endothelial cells. *Blood* 2007;110:3909–3916. [PubMed: 17823308]
140. Feistritzer C, Riewald M. Endothelial barrier protection by activated protein C through PAR1-dependent sphingosine 1-phosphate receptor-1 crossactivation. *Blood* 2005;105:3178–3184. [PubMed: 15626732]
141. Kurosawa S, Esmon CT, Stearns-Kurosawa DJ. The soluble endothelial protein C receptor binds to activated neutrophils: involvement of proteinase-3 and CD11b/CD18. *Journal of Immunology* 2000;165:4697–4703.
142. Franscini N, et al. Gene expression profiling of inflamed human endothelial cells and influence of activated protein C. *Circulation* 2004;110:2903–2909. [PubMed: 15505101]
143. Joyce DE, et al. Gene expression profile of antithrombotic protein c defines new mechanisms modulating inflammation and apoptosis. *Journal of Biological Chemistry* 2001;276:11199–11203. [PubMed: 11278252]
144. Wong SS, et al. Drotrecogin alfa (activated) prevents smoke-induced increases in pulmonary microvascular permeability and proinflammatory cytokine IL-1beta in rats. *Lung* 2004;182:319–330. [PubMed: 15765924]
145. Teke Z, et al. Activated protein C attenuates intestinal reperfusion-induced acute lung injury: an experimental study in a rat model. *American Journal of Surgery* 2008;195:861–873. [PubMed: 18367148]
146. Aiello LP, et al. Vascular endothelial growth factor-induced retinal permeability is mediated by protein kinase C in vivo and suppressed by an orally effective beta-isoform-selective inhibitor. *Diabetes* 1997;46:1473–1480. [PubMed: 9287049]
147. Ishii H, et al. Amelioration of vascular dysfunctions in diabetic rats by an oral PKC beta inhibitor. *Science* 1996;272:728–731. [PubMed: 8614835]
148. Koxsel O, et al. Rho-kinase (ROCK-1 and ROCK-2) upregulation in oleic acid-induced lung injury and its restoration by Y-27632. *European Journal of Pharmacology* 2005;510:135–142. [PubMed: 15740734]
149. Olson MF. Applications for ROCK kinase inhibition. *Current Opinion in Cell Biology* 2008;20:242–248. [PubMed: 18282695]
150. Yin L, et al. Fasudil inhibits vascular endothelial growth factor-induced angiogenesis in vitro and in vivo. *Molecular Cancer Therapeutics* 2007;6:1517–1525. [PubMed: 17513600]
151. Arita R, et al. Rho kinase inhibition by fasudil ameliorates diabetes-induced microvascular damage. *Diabetes* 2009;58:215–226. [PubMed: 18840783]
152. Peng F, et al. RhoA/Rho-kinase contribute to the pathogenesis of diabetic renal disease. *Diabetes* 2008;57:1683–1692. [PubMed: 18356410]
153. van Nieuw Amerongen GP, et al. Simvastatin improves disturbed endothelial barrier function. *Circulation* 2000;102:2803–2809. [PubMed: 11104736]
154. Zeng L, et al. HMG CoA reductase inhibition modulates VEGF-induced endothelial cell hyperpermeability by preventing RhoA activation and myosin regulatory light chain phosphorylation. *FASEB Journal* 2005;19:1845–1847. [PubMed: 16160062]

Further reading, resources and contacts

155. Mammoto A, et al. Rho signaling and mechanical control of vascular development. *Vascular Biology* 2008;15:228–234. This is an in-depth review on the impact of mechanical forces on endothelial barrier function.
156. van Nieuw Amerongen GP, et al. Targets for pharmacological intervention of endothelial hyperpermeability and barrier function. *Vascular Pharmacology* 2003;39:257–272. This is an

extensive and detailed review of different strategies used to target endothelial barrier dysfunction. [PubMed: 12747965]

157. Groeneveld AB, et al. Vascular pharmacology of acute lung injury and acute respiratory distress syndrome. *Vascular Pharmacology* 2003;39:247–256. This article reviews the pathogenesis and current therapeutic strategies for adult respiratory distress syndrome. [PubMed: 12747964]




Figure 1. Schematic diagram of microvascular endothelial barrier structure

The barrier is formed by endothelial cells that connect to each other through the junctional adhesive molecule vascular endothelial (VE)-cadherin, which binds to another VE-cadherin molecule from an adjacent cell and connects to the actin cytoskeleton via a family of catenins (α , β , γ and p120). This endothelial lining is tethered to the extracellular matrix through focal adhesions mediated by transmembrane integrins composed of α and β subunits, focal adhesion kinase (FAK), and cytoskeleton-linking proteins including paxillin and vinculin. The integrity of this barrier is maintained by VE-cadherin-mediated cell-cell adhesions and focal-adhesion-supported cell-matrix attachment. A dynamic interaction among these structural elements controls the opening and closing of the paracellular pathways for fluid, proteins and cells to move across the endothelium. In particular, the Ca^{2+} /calmodulin (CaM)-dependent myosin light chain kinase (MLCK) catalyses phosphorylation of myosin light chains (small red circles), triggering binding of the myosin heavy chain motor domains to actin and their cross-bridge movement. This reaction promotes cytoskeleton contraction and cell retraction. In parallel, phosphorylation of VE-cadherin and/or catenins may cause the junction complex to dissociate from its cytoskeletal anchor, leading to weakened cell-cell adhesion. The cytoskeletal and junctional responses act in concert causing paracellular hyperpermeability. These structural changes are caused by signalling reactions depicted in Figure 2.




Figure 2. Signal transduction in endothelial hyperpermeability

(Legend; see previous page for figure) Multiple cascades of intracellular signalling reactions are initiated when an inflammatory agonist binds to its respective receptor expressed on the endothelial surface [e.g. thrombin binds the protease-activated receptor 1 (PAR-1), histamine binds its receptor H₁, and vascular endothelial growth factor (VEGF) binds its receptor VEGFR-2 (KDR)]. Occupancy of G-protein-coupled receptors activates RhoA and its effector kinase ROCK (left), or it triggers phospholipase (PLC)-catalysed protein kinase C (PKC) activation and elevated intracellular calcium, which stimulates nitric oxide production and cGMP-dependent protein kinase (PKG) activation (right). Agonist binding of receptor tyrosine kinase also activates the mitogen-activated protein (MAP) kinase cascades characterised by

phosphorylation of extracellular-signal-regulated kinases (ERK1/2) (middle). The three pathways (Rho GTPases, MAP kinases and protein kinases) interact with each other, causing changes in the endothelial barrier structure. Abbreviations: cGMP, cyclic guanosine monophosphate; CRAF, Raf-1; DAG, diacylglycerol; ER, endoplasmic reticulum; GC, guanylate cyclase; GDP, guanosine diphosphate; GEF-H1, guanine-nucleotide-exchange factor H1; GRB2, growth factor receptor-bound protein 2; GTP, guanosine triphosphate; JNK, c-Jun N-terminal kinase; JNKK, c-Jun N-terminal kinase kinase; L-Arg, L-arginine; NO, nitric oxide; NOS, nitric oxide synthase; p115RhoGEF, 115 kDa guanine-nucleotide-exchange factor; p190RhoGAP, p190 Rho GTPase-activating protein; Ras, *ras* gene product; Rho-GDI, GDP dissociation inhibitor (GDI); Rho-GDI, Rho GDP-dissociation inhibitor 1; ROCK, Rho kinase; RTK, receptor tyrosine kinase; SAPK, stress-activated protein kinase; SEK, stress-activated protein kinase/ERK kinase; Sos, Son of sevenless; Src, *src* gene product.

Table 1

Acute inflammatory stimuli capable of inducing endothelial hyperpermeability

Stimulus	Receptor	Intracellular signalling	Subcellular targeting	Endothelial cell response	Refs
Histamine	H ₁ , H ₂	Gq, Gs, Gi, PLC/ Ca ²⁺ , PKC, MEK1/2, PKG, RhoA, eNOS, cAMP, PKA	MLC phosphorylation, actin polymerisation, VE-cadherin and β-catenin phosphorylation	Actomyosin contraction, stress-fibre formation, VE-cadherin and β-catenin dissociation, adherens junction disassembly	5, 26, 40, 41, 42, 43, 44, 45, 46
Thrombin	PAR-1, -3 and -4; EPCR	Gq, G _{12/13} , Gi, PLC, Ca ²⁺ , PKC, CaMKII, ERK1/2, p115RhoGEF, ROCK, Src, FAK, cAMP, PKA	MLC phosphorylation, VE-cadherin and β-catenin phosphorylation	Actomyosin contraction, stress-fibre formation, adherens junction disassembly, focal adhesion reorganisation	8, 47, 48, 49, 50, 51
VEGF	VEGFR-2 (KDR)	PLCγ, Ca ²⁺ , PKC, eNOS, PKG, p38 and ERK MAPKs, PI3K, Akt, FAK, Src	MLC phosphorylation, VE-cadherin and β-catenin dissociation, integrin activation	Actomyosin contraction, adherens junction disassembly, focal adhesion formation and redistribution	13, 14, 15, 16, 17, 18, 19, 20, 32, 54, 55, 61, 62, 63
Neutrophils, neutrophil- released cytokines (e.g. TNF, IL-1)	Selectins, ICAMs VCAMs	PLC, Ca ²⁺ , Src, RhoA PKC, MAPK, FAK	ROS generation, VE-cadherin and β-catenin phosphorylation, integrin activation, focal adhesion activation	Adherence to neutrophils, cytoskeleton contraction, adherens junction dissociation	30, 31, 56, 57, 58, 59, 64, 65, 66, 67, 68, 69, 70

Abbreviations: CaMKII, Ca²⁺/calmodulin-dependent protein kinase type II; cAMP, cyclic adenosine monophosphate; eNOS, nitric oxide synthase, endothelial; EPCR, endothelial protein C receptor; ERK1/2, extracellular-signal-regulated kinases; FAK, focal adhesion kinase; G_{12/13}, guanine-nucleotide-binding protein G_{12/13}; Gi, guanine-nucleotide-binding protein G(i) subunit alpha-i; Gq, guanine-nucleotide-binding protein G(q) subunit alpha-q; Gs, guanine-nucleotide-binding protein G(s) subunit alpha short; ICAM, intracellular adhesion molecule; IL-1, interleukin 1; MAPK, mitogen-activated protein kinase; MEK1/2, MAP kinase kinase or ERK kinase; MLC, myosin light chain; MMP, matrix metalloproteinase; p115RhoGEF, 115 kDa guanine-nucleotide-exchange factor; PAR-1, -3, -4, protease-activated receptor 1, 3, 4; PI3K, phosphoinositide 3-kinase; PKA, protein kinase A; PKC, protein kinase C; PKG, cGMP-dependent protein kinase; PLC, phospholipase C; ROCK, Rho kinase; ROS, reactive oxygen species; TNF, tumour necrosis factor; VCAM, vascular cell adhesion molecule; VE-cadherin, vascular endothelial cadherin; VEGF, vascular endothelial growth factor; VEGFR-2, vascular endothelial growth factor receptor 2 (also known as KDR).