

MOLECULAR SIMILARITIES BETWEEN THE Qa-2
ALLOANTIGEN AND OTHER GENE
PRODUCTS OF THE 17TH CHROMOSOME OF THE MOUSE*

By JAMES MICHAELSON, LORRAINE FLAHERTY, ELLEN VITETTA, AND
MIROSLOV D. POULIK

(From the Memorial Sloan-Kettering Institute for Cancer Research, New York 10021; the Division of Laboratories and Research, New York State Department of Health, Albany, New York 12201; the Department of Microbiology, University of Texas Southwestern Medical School, Dallas, Texas 75235; the Department of Immunology and Microbiology, Wayne State University Medical School, Detroit, Michigan 48202; and the Department of Immunochemistry, William Beaumont Hospital Research Institute, Royal Oak, Michigan 48072.)

Recent immunogenetic analysis of the region between *H-2D* and *Tla* on the 17th chromosome of the mouse has revealed the presence of previously unrecognized genes which specify cell surface antigens (1). One part of this area, the *Qa-2* region, contains two such genes, one of which codes for an antigen on thymus, spleen, and lymph node, and the other which is expressed on spleen and lymph node only (2). It has been of interest to us to examine biochemically these gene products. We report here that the molecule which carries *Qa-2* on lymph node cells (LNC) and spleen is composed of a large subunit of approximately the same molecular weight as *H-2* and *TL* molecules, and a small subunit which can be serologically identified as $\beta 2$ -microglobulin ($\beta 2M$).

Materials and Methods

Preparation of Radioiodinated Cells. Cells from the thymus, spleen, or lymph nodes were prepared and iodinated as described previously (3). Cells were lysed in phosphate-buffered saline (PBS) containing 0.5% Nonidet P40 (NP40) (Shell Chemical Corp., New York) and the lysates were centrifuged at 1,200 *g*. Samples were then dialyzed overnight at 4°C against PBS. After dialysis, protein-associated radioactivity was determined (4).

Sera

RABBIT ANTI-MOUSE Ig. Rabbit anti-mouse Ig (RAMIg) (5) contained specificities against μ , γ , κ , α , and λ chains and was a pool of several sera prepared against purified myeloma proteins.

GOAT ANTI-MOUSE Ig. Goat anti-mouse Ig (GAMIg) contained specificities against γ and L chains.

GOAT ANTI-RABBIT Ig. Goat anti-rabbit Ig (GARIG) (6) contained antibodies against rabbit γ and L chains.

αK^b (7). This serum was a gift from Dr. Jan Klein (University of Texas Southwestern Medical School) and was produced in (D2.GD \times B10.D2)F₁ mice against C57BL/6 lymphoid cells.

αD^b . This serum was produced in (HTI \times B6-H-2^K)F₁ mice against the C57BL/6 leukemia EL4, and detects H-2D^b when tested against cells from HTH mice.

* This work was supported by grants from the National Institutes of Health (nos. AI-12603, AI-11650, AI-11335, AI-13448, and CA-08748-12); the Children's Leukemia Foundation of Michigan, the William Beaumont Hospital Research Institute, and The Rockefeller Foundation (no. RF#7404-03).




FIG. 1. SDS-polyacrylamide-gel electrophoresis of radioiodinated Qa-2 antigen. Immunoprecipitates were reduced and electrophoresed on a 7.5% gel. (A) Precipitate of α Qa-2 and C57BL/6 LNC. (B). Precipitate of α Qa-2 and B⁶-H-2^k LNC (control).

α Qa-2. This serum was produced by immunizing B6.K1 mice with C57BL/6 lymphocytes and thymocytes (2).

$\alpha\beta$ 2M. Rabbit anti-rat β 2M (5) was produced as described previously and was cross-reactive with mouse β 2M. Previous studies (5) have shown that this serum can precipitate the β 2M-containing molecules H-2 and TL.

Immunoprecipitation. Dialyzed lysates were depleted of B-cell Ig by treatment with RAMIg and GARIg as described previously (5, 6). The alloantigens were then precipitated with alloantisera and GAMIg. Precipitates were washed and solubilized at 56°C in 1% sodium dodecyl sulfate (SDS) containing 2% 2-mercaptoethanol (2ME) and 6 M urea at pH 6.8. Samples were electrophoresed along with markers of ³H- μ and L chains on 10 cm SDS polyacrylamide gels at 25 V for 18 h. Gels were sliced into 1-mm fragments with a Mickle gel slicer (Brinkmann Instruments, Inc., Westbury, N. Y.). Samples were counted as described previously (3).

Results and Discussion

Using α Qa-2 to precipitate antigens from Qa-2⁺ LNC (C57BL/6, HTH, or B6.K2 mice) two proteins were resolved on 7.5 and 10% acrylamide gels, with apparent mol wt of approximately 43,000 and 12,000 daltons (Fig. 1). The precipitate contained approximately 0.15% of the total labeled protein as compared to 1–4% for H-2. Immunoprecipitates from lysates of spleen cells also show these peaks, but in about 50% of the quantity of Qa-2 found on LNC. No Qa-2 was found on thymocytes by this method. (The reaction of α Qa-2 serum with thymocytes in the cytotoxicity assay is probably due to a second antigen specified by the Qa-2 region.) The amounts of Qa-2 recovered from thymus, spleen, and lymph node cells parallel the number of mature T cells found in these lymphoid organs. LNC from Qa-2⁻ (B6-H-2^k or B6.K1) mice did not yield peaks with the α Qa-2. Since B6.K1 and B6.K2 are genetically identical except for a small portion of the 17th chromosome adjacent to Qa-2, the peaks detected can




FIG. 2. The effect of preclearing radioiodinated lysate of C57BL/6 LNC with αK^b on Qa-2 antigen precipitated with $\alpha Qa-2$. (A) Precleared with αK^b . (B) Precleared with normal mouse serum (control).

be ascribed to the Qa-2 antigen which is detected on LNC with the cytotoxicity assay.

Qa-2 was shown to reside on a molecule separate from H-2D by sequential precipitation of HTH LNC lysate with αD^b antiserum followed by $\alpha Qa-2$. The first precipitation removed 85% of the H-2D peak but did not reduce the quantity of the Qa-2 precipitated subsequently (results not shown). By an analogous procedure the Qa-2 molecule was shown to be distinct from H-2K because prior precipitation of K^b did not reduce the amount of Qa-2 subsequently precipitated (Fig. 2).

Vitetta et al. (5) have recently demonstrated that treatment of splenic or thymic lysates with rabbit anti-rat $\beta 2M$ removes both H-2 and TL molecules. Likewise, prior treatment of lymph node lysates with $\alpha \beta 2M$ completely eliminated the capacity of $\alpha Qa-2$ to precipitate a Qa-2 peak from the NP40 lysate, while prior treatment with normal rabbit serum had no effect (Fig. 3). In the previous studies performed by Vitetta et al. (5), prior treatment of splenic lysates with anti-H-2 depleted all radioactivity which could be subsequently recognized by $\alpha \beta 2M$. However, since the Qa-2 antigen contains less than 5% of the radioactivity found in H-2, these molecules went undetected.

The molecular similarity between H-2D, H-2K, TL, and Qa-2 is remarkable in that they are all molecules of approximately 45,000 daltons which are associated with $\beta 2M$ (5, 8-11). Moreover, to the left of *H-2K* is the *T/t* locus in which map the genes specifying the F9 antigen. This antigen also has a mol wt of 44,000 and a 12,000 subunit (12) which does not carry immunologically recognizable $\beta 2M$ determinants (13). This chromosome therefore contains a family of molecules, related by size, subunit structure, genetic linkage, membrane location, and antigenicity, having most likely arisen from a common ancestor gene by tandem duplication.

It is of interest that one of the two antigens determined by the *Qa-2* region appears to be expressed on peripheral T cells and is absent from thymocytes (2), thus being the first known alloantigen with this pattern of expression. The transition from thymocyte to peripheral T cell in a TL^+ , $Qa-2^+$ mouse therefore involves the loss of TL and the acquisition of Qa-2. This is reminiscent of the loss




FIG. 3. The effect of preclearing radioiodinated lysate of C57BL/6 LNC with $\alpha\beta 2M$ on Qa-2 antigen precipitation with $\alpha Qa-2$. (A) Precleared with normal rabbit sera (control). (B) Precleared with $\alpha\beta 2M$.

of F9 and gain of H-2 (14) which occurs during embryogenesis, and of the reduction of H-2D which accompanies the expression of TL during thymocyte development (15). Presumably, all of these molecules are not only biochemically but functionally analogous. In light of evidence suggesting that F9 functions in cellular recognition in the early embryo (16) and that the *H-2* region is important in cellular interaction in the immune response (17, 18) it is tempting to speculate that a variety of cell to cell interactions may be mediated by the family of molecules to which Qa-2 belongs.

Summary

The alloantigen Qa-2, whose gene is located on the 17th chromosome between *H-2D* and *Tla*, is identified as a molecule of 43,000 daltons which is associated with $\beta 2$ -microglobulin. Qa-2 comprises approximately 0.15% of the iodinateable cell surface protein of lymph node cells. Sequential precipitations demonstrated that Qa-2 is distinct from H-2D and H-2K molecules.

We wish to express our thanks to Doctors E. A. Boyse, D. Bennett, and K. Artzt whose advice and comments have been invaluable.

Received for publication 10 January 1977.

References

1. Stanton, T. H., and E. A. Boyse. 1976. A new serologically defined locus, *Qa-1*, in the *Tla*-region of the mouse. *Immunogenetics*. 3:525.
2. Flaherty, L. 1976. The *Tla* region of the mouse: identification of a new serologically defined locus, *Qa-2*. *Immunogenetics*. 3:533.

3. Baur, S., E. S. Vitetta, C. J. Sherr, I. Schenkein, and J. W. Uhr. 1971. Isolation of heavy and light chains of immunoglobulin from the surfaces of lymphoid cells. *J. Immunol.* 106:1133.
4. Vitetta, E. S., J. Forman, and J. R. Kettman. 1976. Cell surface immunoglobulin. XVIII. Functional differences of B lymphocytes bearing different surface immunoglobulin isotypes. *J. Exp. Med.* 143:1055.
5. Vitetta, E. S., M. D. Poulik, J. Klein, and J. W. Uhr. 1976. Beta 2-microglobulin is selectively associated with H-2 and TL alloantigens on murine lymphoid cells. *J. Exp. Med.* 144:179.
6. Vitetta, E. S., J. W. Uhr, and E. A. Boyse. 1975. Association of a β 2-microglobulin-like subunit with H-2 and TL alloantigens on murine thymocytes. *J. Immunol.* 114:252.
7. Capra, J. D., E. S. Vitetta, D. G. Klapper, J. W. Uhr, and J. Klein. 1976. Structural studies on protein products of murine chromosome 17: partial amino acid sequence of an H-2K^b molecule. *Proc. Natl. Acad. Sci. U. S. A.* 73:3661.
8. Silver, J., and L. Hood. 1974. Detergent-solubilised H-2 alloantigen is associated with a small molecular weight polypeptide. *Nature (Lond.)*. 249:764.
9. Natori, T., M. Katagiri, N. Tanigaki, and D. Pressman. 1974. The 11,000-dalton component of mouse H-2: isolation and identification. *Transplantation (Baltimore)*. 18:550.
10. Rask, L., J. B. Lindblom, and P. A. Peterson. 1974. Subunit structure of H-2 alloantigens. *Nature (Lond.)*. 249:833.
11. Ostberg, L., L. Rask, H. Wigzell, and P. A. Peterson. 1975. Thymus leukemia antigen contains β 2-microglobulin. *Nature (Lond.)*. 253:735.
12. Vitetta, E. S., K. Artzt, D. Bennett, E. A. Boyse, and F. Jacob. 1975. Structural similarities between a product of the T/t-locus isolated from sperm and teratoma cells, and H-2 antigens isolated from splenocytes. *Proc. Natl. Acad. Sci. U.S.A.* 73:3215.
13. Dubois, P., M. Fellous, G. Gachelin, F. Jacob, R. Kemler, D. Pressman, and N. Tanigaki. 1976. Absence of a serologically detectable association of murine β 2-microglobulin with the embryonic F9 antigen. *Transplantation (Baltimore)*. 22:467.
14. Artzt, K., and D. Bennett. 1975. Analogies between embryonic (T/t) antigens and adult major histocompatibility (H-2) antigens. *Nature (Lond.)*. 256:545.
15. Boyse, E. A., E. Stockert, and L. J. Old. 1968. Isoantigens of the H-2 and Tla loci of the mouse: interactions affecting their representation on thymocytes. *J. Exp. Med.* 128:85.
16. Artzt, K., D. Bennett, and F. Jacob. 1974. Primitive teratocarcinoma cells express a differentiation antigen specified by a gene at the T-locus in the mouse. *Proc. Natl. Acad. Sci. U. S. A.* 71:811.
17. Shearer, G. M. 1974. Cell-mediated cytotoxicity to trinitrophenyl-modified syngeneic lymphocytes. *Eur. J. Immunol.* 4:527.
18. Zinkernagel, R. M., and P. C. Doherty. Restriction of *in vitro* T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature (Lond.)*. 248:701.