

# Monge–Ampère equations in big cohomology classes

by

SÉBASTIEN BOUCKSOM

*CNRS–Université Pierre et Marie Curie  
Paris, France*

PHILIPPE EYSSIDIEUX

*Université Grenoble 1  
Saint-Martin d’Hères, France*

VINCENT GUEDJ

*Université Aix-Marseille I  
Marseille, France*

AHMED ZERIAHI

*Université Paul Sabatier  
Toulouse, France*

## Contents

Introduction . . . . .	199
1. Non-pluripolar products of closed positive currents . . . . .	202
2. Weighted energy classes . . . . .	223
3. Non-pluripolar measures are Monge–Ampère . . . . .	236
4. $L^\infty$ a-priori estimate . . . . .	244
5. Regularity of solutions: the case of nef classes . . . . .	249
6. Singular Kähler–Einstein metrics . . . . .	253
References . . . . .	260

## Introduction

The primary goal of the present paper is to extend the results obtained by the last two authors on complex Monge–Ampère equations on compact Kähler manifolds in [27] to the case of *arbitrary* cohomology classes.

More specifically, let  $X$  be a compact  $n$ -dimensional Kähler manifold and let  $\omega$  be a Kähler form on  $X$ . An  $\omega$ -*plurisubharmonic* (*psh* for short) function is an upper semi-continuous function  $\varphi$  such that  $\omega + dd^c\varphi$  is positive in the sense of currents, and [27] focused on the Monge–Ampère type equation

$$(\omega + dd^c\varphi)^n = \mu, \tag{0.1}$$

where  $\mu$  is a positive measure on  $X$  of total mass  $\mu(X) = \int_X \omega^n$ . As is well known, it is not always possible to make sense of the left-hand side of (0.1), but it was observed

in [27] that a construction going back to Bedford–Taylor [4] enables one, in this global setting, to define the non-pluripolar part of the would-be positive measure  $(\omega + dd^c\varphi)^n$  for an arbitrary  $\omega$ -psh function  $\varphi$ .

In the present paper we first give a more systematic treatment of these issues. We explain how to define, in a simple and canonical way, the *non-pluripolar product*

$$\langle T_1 \wedge \dots \wedge T_p \rangle$$

of arbitrary closed positive  $(1, 1)$ -currents  $T_1, \dots, T_p$  with  $1 \leq p \leq n$ . The resulting positive  $(p, p)$ -current  $\langle T_1 \wedge \dots \wedge T_p \rangle$  puts no mass on pluripolar subsets. It is also shown to be *closed* (Theorem 1.8), generalizing a classical result of Skoda–El Mir. We relate its cohomology class to the *positive intersection class*

$$\langle \alpha_1 \dots \alpha_p \rangle \in H^{p,p}(X, \mathbb{R})$$

of the cohomology classes  $\alpha_j := \{T_j\} \in H^{1,1}(X, \mathbb{R})$  in the sense of [12] and [13]. In particular we show that  $\int_X \langle T^n \rangle \leq \text{vol}(\alpha)$ , where the right-hand side denotes the *volume* of the class  $\alpha := \{T\}$  [10], which implies in particular that  $\langle T^n \rangle$  is non-trivial only if the cohomology class  $\alpha$  is *big*.

An important aspect of the present approach is that the non-pluripolar Monge–Ampère measure  $\langle T^n \rangle$  is well defined for *any* closed positive  $(1, 1)$ -current  $T$ . In the second section we study the continuity properties of the mapping  $T \mapsto \langle T^n \rangle$ : it is continuous along decreasing sequences (of potentials) if and only if  $T$  has *full Monge–Ampère mass* (Theorem 2.17), i.e. when

$$\int_X \langle T^n \rangle = \text{vol}(\alpha).$$

We prove this fact by defining and studying weighted energy functionals in this general context, extending the case of a Kähler class [27]. The two main new features are a generalized comparison principle (Corollary 2.3) and an asymptotic criterion to check whether a current  $T$  has full Monge–Ampère mass (Proposition 2.19).

In the third part of the paper we obtain our first main result (Theorem 3.1):

**THEOREM A.** *Let  $\alpha \in H^{1,1}(X, \mathbb{R})$  be a big class on a compact Kähler manifold  $X$ . If  $\mu$  is a positive measure on  $X$  that puts no mass on pluripolar subsets and satisfies the compatibility condition  $\mu(X) = \text{vol}(\alpha)$ , then there exists a unique closed positive  $(1, 1)$ -current  $T \in \alpha$  such that*

$$\langle T^n \rangle = \mu.$$

The existence part extends the main result of [27], which corresponds exactly to the case where  $\alpha$  is a Kähler class. In fact the proof of Theorem A consists of reducing to the Kähler case via approximate Zariski decompositions. Uniqueness is obtained by adapting the proof of S. Dinew [22] (which also deals with the Kähler-class case).

When the measure  $\mu$  satisfies some additional regularity condition, we show how to adapt Kołodziej’s pluripotential theoretic approach to the sup-norm a-priori estimates, see [31], to get *global* information on the singularities of  $T$ .

**THEOREM B.** *Assume that the measure  $\mu$  in Theorem A furthermore has  $L^{1+\varepsilon}$  density with respect to Lebesgue measure for some  $\varepsilon > 0$ . Then the solution  $T \in \alpha$  to  $\langle T^n \rangle = \mu$  has minimal singularities.*

Currents with minimal singularities were introduced by Demailly. When  $\alpha$  is a Kähler class, the positive currents  $T \in \alpha$  with minimal singularities are exactly those with *locally bounded* potentials. When  $\alpha$  is merely big, all positive currents  $T \in \alpha$  will have poles in general, and the minimal singularity condition on  $T$  essentially says that  $T$  has the least possible poles among all positive currents in  $\alpha$ . Currents with minimal singularities have in particular locally bounded potentials on the *ample locus*  $\text{Amp}(\alpha)$  of  $\alpha$ , which is roughly speaking the largest Zariski open subset where  $\alpha$  locally behaves like a Kähler class.

Regarding local regularity properties, we obtain the following result.

**THEOREM C.** *In the setting of Theorem A, assume that  $\mu$  is a smooth strictly positive volume form. Assume also that  $\alpha$  is nef. Then the solution  $T \in \alpha$  to the equation  $\langle T^n \rangle = \mu$  is  $C^\infty$  on  $\text{Amp}(\alpha)$ .*

The expectation is of course that Theorem C holds whether or not  $\alpha$  is nef, but we are unfortunately unable to prove this for the moment. It is perhaps worth emphasizing that currents with minimal singularities can have a non-empty polar set even when  $\alpha$  is nef and big (see Example 5.4).

In the last part of the paper we consider Monge–Ampère equations of the form

$$\langle (\theta + dd^c \varphi)^n \rangle = e^\varphi dV, \tag{0.2}$$

where  $\theta$  is a smooth representative of a big cohomology class  $\alpha$ ,  $\varphi$  is a  $\theta$ -psh function and  $dV$  is a smooth positive volume form. We show that (0.2) admits a unique solution  $\varphi$  such that  $\int_X e^\varphi dV = \text{vol}(\alpha)$ . Theorem B then shows that  $\varphi$  has minimal singularities, and we obtain as a special case the following result.

THEOREM D. *Let  $X$  be a smooth projective variety of general type. Then  $X$  admits a unique singular Kähler–Einstein volume form of total mass equal to  $\text{vol}(K_X)$ . In other words, the canonical bundle  $K_X$  can be endowed with a unique non-negatively curved metric  $e^{-\phi_{\text{KE}}}$  whose curvature current  $dd^c\phi_{\text{KE}}$  satisfies*

$$\langle (dd^c\phi_{\text{KE}})^n \rangle = e^{\phi_{\text{KE}}} \quad (0.3)$$

and is such that

$$\int_X e^{\phi_{\text{KE}}} = \text{vol}(K_X). \quad (0.4)$$

The weight  $\phi_{\text{KE}}$  furthermore has minimal singularities.

Since the canonical ring  $R(K_X) = \bigoplus_{k \geq 0} H^0(kK_X)$  is now known to be finitely generated [7], this result can be obtained as a consequence of [25] by passing to the canonical model of  $X$ . But one of the points of the proof presented here is to avoid the use of the difficult result of [7].

The existence of  $\phi_{\text{KE}}$  satisfying (0.3) and (0.4) was also recently obtained by J. Song and G. Tian in [38], building on a previous approach of H. Tsuji [43]. It is also shown in [38] that  $\phi_{\text{KE}}$  is an *analytic Zariski decomposition* in the sense of Tsuji, which means that every pluricanonical section  $\sigma \in H^0(mK_X)$  is  $L^\infty$  with respect to the metric induced by  $\phi_{\text{KE}}$ . The main new information we add is that  $\phi_{\text{KE}}$  actually has minimal singularities, which is strictly stronger than being an analytic Zariski decomposition for general big line bundles (cf. Proposition 6.5).

*Acknowledgement.* We would like to thank J.-P. Demailly and R. Berman for several useful discussions.

## 1. Non-pluripolar products of closed positive currents

In this section  $X$  denotes an arbitrary  $n$ -dimensional complex manifold unless otherwise specified.

### 1.1. Plurifine topology

The *plurifine topology* on  $X$  is defined as the coarsest topology with respect to which all psh functions  $u$  on all open subsets of  $X$  become continuous (cf. [4]). Note that the plurifine topology on  $X$  is always strictly finer than the ordinary one.

Since psh functions are upper semi-continuous, subsets of the form

$$V \cap \{u > 0\}$$

(as usual, we write  $\{u > 0\}$  for the set  $\{x \in V : u(x) > 0\}$ ) with  $V \subset X$  open and  $u$  psh on  $V$  obviously form a basis for the plurifine topology, and  $u$  can moreover be assumed to be locally bounded by maxing it with 0.

When  $X$  is furthermore compact and Kähler, a bounded local psh function defined on an open subset  $V$  of  $X$  can be extended to a quasi-psh function on  $X$  (possibly after shrinking  $V$  a little bit, see [26]), and it follows that the plurifine topology on  $X$  can alternatively be described as the coarsest topology with respect to which all bounded quasi-psh functions  $\varphi$  on  $X$  become continuous. It therefore admits sets of the form  $V \cap \{\varphi > 0\}$  with  $V \subset X$  open and  $\varphi$  quasi-psh and bounded on  $X$  as a basis.

## 1.2. Local non-pluripolar products of currents

Let  $u_1, \dots, u_p$  be psh functions on  $X$ . If the  $u_j$ 's are locally bounded, then the fundamental work of Bedford–Taylor [3] enables one to define

$$dd^c u_1 \wedge \dots \wedge dd^c u_p$$

on  $X$  as a closed positive  $(p, p)$ -current. The wedge product only depends on the closed positive  $(1, 1)$ -currents  $dd^c u_j$ , and not on the specific choice of the potentials  $u_j$ .

A very important property of this construction is that it is *local in the plurifine topology*, in the following sense. If  $u_j$  and  $v_j$  are locally bounded psh functions on  $X$  and  $u_j = v_j$  (pointwise) on a plurifine open subset  $O$  of  $U$ , then

$$\mathbf{1}_O dd^c u_1 \wedge \dots \wedge dd^c u_p = \mathbf{1}_O dd^c v_1 \wedge \dots \wedge dd^c v_p.$$

This is indeed an obvious generalization of Corollary 4.3 of [4].

In the case of a possibly unbounded psh function  $u$ , Bedford–Taylor have observed (cf. [4, p. 236]) that it is always possible to define the *non-pluripolar* part of the would-be positive measure  $(dd^c u)^n$  as a Borel measure. The main issue however is that this measure is not going to be locally finite in general (see Example 1.3 below). More generally, suppose that we are trying to associate with any  $p$ -tuple  $u_1, \dots, u_p$  of psh functions on  $X$  a positive (a priori not necessarily closed)  $(p, p)$ -current

$$\langle dd^c u_1 \wedge \dots \wedge dd^c u_p \rangle$$

putting no mass on pluripolar subsets, in such a way that the construction is local in the plurifine topology in the above sense. Then we have no choice: since  $u_j$  coincides with the locally bounded psh function  $\max\{u_j, -k\}$  on the plurifine open subset

$$O_k := \bigcap_{j=1}^p \{u_j > -k\}$$

we must have

$$\mathbf{1}_{O_k} \left\langle \bigwedge_{j=1}^p dd^c u_j \right\rangle = \mathbf{1}_{O_k} \bigwedge_{j=1}^p dd^c \max\{u_j, -k\}. \quad (1.1)$$

Note that the right-hand side is non-decreasing in  $k$ . This equation completely determines  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$ , since the latter is required not to put mass on the pluripolar set

$$X \setminus \bigcup_{k=1}^{\infty} O_k = \{u = -\infty\}.$$

*Definition 1.1.* If  $u_1, \dots, u_p$  are psh functions on the complex manifold  $X$ , we shall say that the non-pluripolar product  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  is well defined on  $X$  if for each compact subset  $K$  of  $X$  we have

$$\sup_k \int_{K \cap O_k} \omega^{n-p} \wedge \bigwedge_{j=1}^p dd^c \max\{u_j, -k\} < \infty. \quad (1.2)$$

Here  $\omega$  is an auxiliary (strictly) positive  $(1,1)$ -form on  $X$  with respect to which masses are being measured, the condition being of course independent of  $\omega$ . When (1.2) is satisfied, equation (1.1) indeed defines a positive  $(p,p)$ -current  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  on  $X$ . We will show below that it is automatically *closed* (Theorem 1.8).

Condition (1.2) is always satisfied when  $p=1$ , and in fact it is not difficult to show that

$$\langle dd^c u \rangle = \mathbf{1}_{\{u > -\infty\}} dd^c u.$$

There are however examples where non-pluripolar products are not well defined as soon as  $p \geq 2$ . This is most easily understood in the following situation.

*Definition 1.2.* A psh function  $u$  on  $X$  will be said to have *small unbounded locus* if there exists a (locally) complete pluripolar closed subset  $A$  of  $X$  outside which  $u$  is locally bounded.

Assume that  $u_1, \dots, u_p$  have small unbounded locus, and let  $A$  be a closed complete pluripolar subset of  $X$  such that each  $u_j$  is locally bounded outside  $A$  (recall that complete pluripolar subsets are stable under finite unions). Then  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  is well defined if and only if the Bedford–Taylor product  $\bigwedge_{j=1}^p dd^c u_j$ , which is defined on the open subset  $X \setminus A$ , has locally finite mass near each point of  $A$ . In that case  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  is nothing but the trivial extension of  $\bigwedge_{j=1}^p dd^c u_j$  to  $X$ .

*Example 1.3.* Consider Kiselman’s example (see [28])

$$u(x, y) := (1 - |x|^2)(-\log |y|)^{1/2}$$

for  $(x, y) \in \mathbb{C}^2$  near 0. The function  $u$  is psh near 0, it is smooth outside the  $y=0$  axis, but the smooth measure  $(dd^c u)^2$ , defined outside  $y=0$ , is not locally finite near any point of  $y=0$ . This means that the positive Borel measure  $\langle (dd^c u)^2 \rangle$  is not locally finite.

We now collect some basic properties of non-pluripolar products.

PROPOSITION 1.4. *Let  $u_1, \dots, u_p$  be psh functions on  $X$ .*

(a) *The operator*

$$(u_1, \dots, u_p) \mapsto \left\langle \bigwedge_{j=1}^p dd^c u_j \right\rangle$$

*is local in the plurifine topology whenever it is well defined.*

(b) *The current  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  and the fact that it is well defined, both only depend on the currents  $dd^c u_j$ , not on the specific potentials  $u_j$ .*

(c) *Non-pluripolar products, which are obviously symmetric, are also multilinear in the following sense: if  $v_1$  is another psh function, then*

$$\left\langle (dd^c u_1 + dd^c v_1) \wedge \bigwedge_{j=2}^p dd^c u_j \right\rangle = \left\langle dd^c u_1 \wedge \bigwedge_{j=2}^p dd^c u_j \right\rangle + \left\langle dd^c v_1 \wedge \bigwedge_{j=2}^p dd^c u_j \right\rangle,$$

*in the sense that the left-hand side is well defined if and only if both terms in the right-hand side are, and equality holds in this case.*

*Proof.* (a) If  $u_j$  and  $v_j$  are psh functions on  $X$  such that  $u_j = v_j$  on a given plurifine open subset  $O$ , then the locally bounded psh functions  $\max\{u_j, -k\}$  and  $\max\{v_j, -k\}$  also coincide on  $O$ . If we set

$$E_k := \bigcap_{j=1}^p \{u_j > -k\} \cap \bigcap_{j=1}^p \{v_j > -k\},$$

then we infer that

$$\mathbf{1}_{O \cap E_k} \bigwedge_{j=1}^p dd^c \max\{u_j, -k\} = \mathbf{1}_{O \cap E_k} \bigwedge_{j=1}^p dd^c \max\{v_j, -k\}.$$

Hence in the limit

$$\mathbf{1}_O \left\langle \bigwedge_{j=1}^p dd^c u_j \right\rangle = \mathbf{1}_O \left\langle \bigwedge_{j=1}^p dd^c v_j \right\rangle,$$

as desired by Lemma 1.5 below.

(b) Let  $w_j$  be pluriharmonic on  $X$ , and let  $K$  be a compact subset. We can find  $C > 0$  such that  $w_j \leq C$  on an open neighborhood  $V$  of  $K$ . On the plurifine open subset

$$O_k := \bigcap_{j=1}^p \{u_j + w_j > -k\} \cap V \subset \bigcap_{j=1}^p \{u_j > -k - C\} \cap V$$

the following locally bounded psh functions coincide:

$$\max\{u_j + w_j, -k\} = \max\{u_j, -w_j - k\} + w_j = \max\{u_j, -k - C\} + w_j.$$

Since  $dd^c w_j = 0$ , it follows that

$$\begin{aligned} \int_{O_k} \omega^{n-p} \wedge \bigwedge_{j=1}^p dd^c \max\{u_j + w_j, -k\} &= \int_{O_k} \omega^{n-p} \wedge \bigwedge_{j=1}^p dd^c \max\{u_j, -k - C\} \\ &\leq \int_{\bigcap_{j=1}^p \{u_j > -k - C\} \cap V} \omega^{n-p} \wedge \bigwedge_{j=1}^p dd^c \max\{u_j, -k - C\}, \end{aligned}$$

which is uniformly bounded in  $k$  by assumption, and the second point is proved.

The proof of (c) is similarly easy but tedious and will be left to the reader. □

LEMMA 1.5. *Assume that the non-pluripolar product  $\langle \bigwedge_{j=1}^p dd^c u_j \rangle$  is well defined. Then for every sequence of Borel subsets  $E_k$  such that*

$$E_k \subset \bigcap_{j=1}^p \{u_j > -k\}$$

and  $X \setminus \bigcup_{k=1}^\infty E_k$  is pluripolar, we have

$$\lim_{k \rightarrow \infty} \mathbf{1}_{E_k} \bigwedge_{j=1}^p dd^c \max\{u_j, -k\} = \left\langle \bigwedge_{j=1}^p dd^c u_j \right\rangle$$

against all bounded measurable functions.

*Proof.* This follows by dominated convergence from

$$(\mathbf{1}_{\bigcap_{j=1}^p \{u_j > -k\}} - \mathbf{1}_{E_k}) \bigwedge_{j=1}^p dd^c \max\{u_j, -k\} \leq (1 - \mathbf{1}_{E_k}) \left\langle \bigwedge_{j=1}^p dd^c u_j \right\rangle. \quad \square$$

A crucial point for what follows is that non-pluripolar products of globally defined currents are *always* well defined on compact Kähler manifolds.

PROPOSITION 1.6. *Let  $T_1, \dots, T_p$  be closed positive  $(1, 1)$ -currents on a compact Kähler manifold  $X$ . Then their non-pluripolar product  $\langle T_1 \wedge \dots \wedge T_p \rangle$  is well defined.*



*Proof.* Let  $\omega$  be a Kähler form on  $X$ . In view of Proposition 1.4 (c), upon adding a large multiple of  $\omega$  to the  $T_j$ 's, we may assume that their cohomology classes are Kähler classes. We can thus find Kähler forms  $\omega_j$  and  $\omega_j$ -psh functions  $\varphi_j$  such that  $T_j = \omega_j + dd^c \varphi_j$ . Let  $U$  be a small open subset of  $X$  on which  $\omega_j = dd^c \psi_j$ , where  $\psi_j \leq 0$  is a smooth psh function on  $U$ , so that  $T_j = dd^c u_j$  on  $U$  with  $u_j := \psi_j + \varphi_j$ . The bounded psh functions on  $U$ ,

$$\psi_j + \max\{\varphi_j, -k\} \quad \text{and} \quad \max\{u_j, -k\}$$

coincide on the plurifine open subset  $\{u_j > -k\} \subset \{\varphi_j > -k\}$ , thus we have

$$\begin{aligned} \int_{\bigcap_{j=1}^p \{u_j > -k\}} \omega^{n-p} \wedge \bigwedge_{j=1}^p dd^c \max\{u_j, -k\} &= \int_{\bigcap_{j=1}^p \{u_j > -k\}} \omega^{n-p} \wedge \bigwedge_{j=1}^p (\omega_j + dd^c \max\{\varphi_j, -k\}) \\ &\leq \int_X \omega^{n-p} \wedge \bigwedge_{j=1}^p (\omega_j + dd^c \max\{\varphi_j, -k\}). \end{aligned}$$

But the latter integral is computed in cohomology, and hence is independent of  $k$ . This shows that (1.2) is satisfied on  $U$ .  $\square$

*Remark 1.7.* The same property extends to the case where  $X$  is a compact complex manifold in the Fujiki class, i.e. bimeromorphic to a compact Kähler manifold. Indeed there exists in that case a modification  $\mu: X' \rightarrow X$  with  $X'$  compact Kähler. Since  $\mu$  is an isomorphism outside closed analytic (hence pluripolar) subsets, it is easy to deduce that  $\langle T_1 \wedge \dots \wedge T_p \rangle$  is well defined on  $X$  from the fact that  $\langle \mu^* T_1 \wedge \dots \wedge \mu^* T_p \rangle$  is well defined on  $X'$ , and in fact

$$\langle T_1 \wedge \dots \wedge T_p \rangle = \mu_* \langle \mu^* T_1 \wedge \dots \wedge \mu^* T_p \rangle.$$

On the other hand, it seems to be unknown whether finiteness of non-pluripolar products holds on arbitrary compact complex manifolds.

Building on the proof of the Skoda–El Mir extension theorem, we will now prove the less trivial closedness property of non-pluripolar products.

**THEOREM 1.8.** *Let  $T_1, \dots, T_p$  be closed positive  $(1, 1)$ -currents on a complex manifold  $X$  whose non-pluripolar product is well defined. Then, the positive  $(p, p)$ -current*

$$\langle T_1 \wedge \dots \wedge T_p \rangle$$

*is closed.*

*Proof.* The result is of course local, and we can assume that  $X$  is a small neighborhood of  $0 \in \mathbb{C}^n$ . The proof will follow rather closely Sibony's proof of the Skoda–El Mir theorem [35], cf. also [18, pp. 159–161]. Let  $u_j \leq 0$  be a local potential of  $T_j$  near  $0 \in \mathbb{C}^n$ , and for each  $k$  consider the closed positive current of bidimension  $(1, 1)$ ,

$$\Theta_k := \varrho \wedge \bigwedge_{j=1}^p dd^c \max\{u_j, -k\}$$

and the plurifine open subset

$$O_k := \bigcap_{j=1}^p \{u_j > -k\}$$

so that  $\mathbf{1}_{O_k} \Theta_k$  converges towards

$$\varrho \wedge \langle T_1 \wedge \dots \wedge T_p \rangle$$

by (1.1). Here  $\varrho$  is a positive  $(n-p-1, n-p-1)$ -form with constant coefficients, so that  $\Theta_k$  has bidimension  $(1, 1)$ . It is enough to show that

$$\lim_{k \rightarrow \infty} d(\mathbf{1}_{O_k} \Theta_k) = 0$$

for any choice of such a form  $\varrho$ .

Let also  $u := \sum_{j=1}^p u_j$ , so that  $u \leq -k$  outside  $O_k$ , and set

$$w_k := \chi(e^{u/k}),$$

where  $\chi(t)$  is a smooth convex and non-decreasing function of  $t \in \mathbb{R}$  such that  $\chi(t) = 0$  for  $t \leq \frac{1}{2}$  and  $\chi(1) = 1$ . We thus see that  $0 \leq w_k \leq 1$  is a non-decreasing sequence of bounded psh functions defined near  $0 \in \mathbb{C}^n$  with  $w_k = 0$  outside  $O_k$  and  $w_k \rightarrow 1$  pointwise outside the pluripolar set  $\{u = -\infty\}$ . Finally let  $0 \leq \theta(t) \leq 1$  be a smooth non-decreasing function of  $t \in \mathbb{R}$  such that  $\theta(t) = 0$  for  $t \leq \frac{1}{2}$  and  $\theta \equiv 1$  near  $t = 1$ . The functions  $\theta(w_k)$  are bounded, non-decreasing in  $k$  and we have

$$\theta(w_k) \leq \mathbf{1}_{O_k}$$

since  $\theta(w_k) \leq 1$  vanishes outside  $O_k$ . Note also that  $\theta'(w_k)$  vanishes outside  $O_k$ , and converges to 0 pointwise outside  $\{u = -\infty\}$ .

Our goal is to show that

$$\lim_{k \rightarrow \infty} d(\mathbf{1}_{O_k} \Theta_k) = 0.$$

But we have

$$0 \leq (\mathbf{1}_{O_k} - \theta(w_k)) \Theta_k \leq (1 - \theta(w_k)) \langle T_1 \wedge \dots \wedge T_p \rangle,$$

and the latter converges to 0 by dominated convergence since  $\theta(w_k) \rightarrow 1$  pointwise outside the polar set of  $u$ , which is negligible for  $\langle T_1 \wedge \dots \wedge T_p \rangle$ . It is thus equivalent to show that

$$\lim_{k \rightarrow \infty} d(\theta(w_k)\Theta_k) = 0.$$

Since  $w_k$  is a bounded psh function, Lemma 1.9 below shows that the chain rule applies, that is

$$d(\theta(w_k)\Theta_k) = \theta'(w_k) dw_k \wedge \Theta_k.$$

Recall that  $dw_k \wedge \Theta_k$  has order 0 by the Bedford–Taylor theorem, so that the right-hand side makes sense. Now let  $\psi$  be a given smooth 1-form compactly supported near 0 and let  $\tau \geq 0$  be a smooth cut-off function with  $\tau \equiv 1$  on the support of  $\psi$ . The Cauchy–Schwarz inequality implies that

$$\left| \int_X \theta'(w_k) \psi \wedge dw_k \wedge \Theta_k \right|^2 \leq \left( \int_X \tau dw_k \wedge d^c w_k \wedge \Theta_k \right) \left( \int_X \theta'(w_k)^2 \psi \wedge \bar{\psi} \wedge \Theta_k \right).$$

But we have

$$2 \int_X \tau dw_k \wedge d^c w_k \wedge \Theta_k \leq \int_X \tau dd^c w_k^2 \wedge \Theta_k = \int_X w_k^2 dd^c \tau \wedge \Theta_k = \int_{O_k} w_k^2 dd^c \tau \wedge \Theta_k,$$

since  $w_k$  vanishes outside  $O_k$ , and the last integral is uniformly bounded, as  $0 \leq w_k \leq 1$  and  $\mathbf{1}_{O_k} \Theta_k$  has uniformly bounded mass by (1.2). On the other hand, we have

$$\theta'(w_k)^2 \Theta_k = \theta'(w_k)^2 \mathbf{1}_{O_k} \Theta_k \leq \theta'(w_k)^2 \langle T_1 \wedge \dots \wedge T_p \rangle,$$

since  $\theta'(w_k)$  also vanishes outside  $O_k$ , and we conclude that

$$\lim_{k \rightarrow \infty} \int_X \theta'(w_k)^2 \psi \wedge \bar{\psi} \wedge \Theta_k = 0$$

by dominated convergence as  $\theta'(w_k) \rightarrow 0$  pointwise outside the polar set of  $u$ , which is negligible for  $\langle T_1 \wedge \dots \wedge T_p \rangle$ . The proof is thus complete.  $\square$

LEMMA 1.9. *Let  $\Theta$  be a closed positive  $(p, p)$ -current on a complex manifold  $X$ ,  $f$  be a smooth function on  $\mathbb{R}$  and  $v$  be a bounded psh function. Then we have*

$$d(f(v)\Theta) = f'(v) dv \wedge \Theta.$$

*Proof.* This is a local result, and we can thus assume that  $\Theta$  has bidimension  $(1, 1)$  by multiplying it by constant forms as above. The result is of course true when  $v$  is smooth. As we shall see the result holds true in our case basically because  $v$  belongs

to the Sobolev space  $L_1^2(\Theta)$ , in the sense that  $dv \wedge d^c v \wedge \Theta$  is well defined. Indeed the result is standard when  $\Theta = [X]$ , and proceeds by approximation. Here we let  $v_k$  be a decreasing sequence of smooth psh functions converging pointwise to  $v$ . We then have  $f(v_k)\theta \rightarrow f(v)\Theta$  by dominated convergence. Thus it suffices to show that

$$\lim_{k \rightarrow \infty} f'(v_k) dv_k \wedge \Theta = f'(v) dv \wedge \Theta.$$

We write

$$f'(v_k) dv_k \wedge \Theta - f'(v) dv \wedge \Theta = (f'(v_k) - f'(v)) dv_k \wedge \Theta + f'(v)(dv_k - dv) \wedge \Theta.$$

Let  $\psi$  be a test 1-form and  $\tau \geq 0$  be a smooth cut-off function with  $\tau \equiv 1$  on the support of  $\psi$ . The Cauchy–Schwarz inequality implies that

$$\left| \int_X (f'(v_k) - f'(v)) \psi \wedge dv_k \wedge \Theta \right|^2 \leq \left( \int_X (f'(v_k) - f'(v))^2 \psi \wedge \bar{\psi} \wedge \Theta \right) \left( \int_X \tau dv_k \wedge d^c v_k \wedge \Theta \right).$$

The second factor is bounded since  $dv_k \wedge d^c v_k \wedge \Theta$  converges to  $dv \wedge d^c v \wedge \Theta$  by Bedford–Taylor’s monotone convergence theorem, and the first one converges to 0 by dominated convergence. We similarly have

$$\left| \int_X f'(v)(dv_k - dv) \wedge \psi \wedge \Theta \right|^2 \leq \left( \int_X f'(v)^2 \psi \wedge \bar{\psi} \wedge \Theta \right) \left( \int_X \tau d(v_k - v) \wedge d^c(v_k - v) \wedge \Theta \right),$$

where now the first factor is bounded while the second one tends to 0 by Bedford–Taylor’s theorem once again, and the proof is complete.  $\square$

*Remark 1.10.* Injecting as above Lemma 1.9 in the proof of Skoda–El Mir’s extension theorem presented in [18, pp. 159–161] shows that Skoda–El Mir’s result remains true for complete pluripolar subsets that are not necessarily *closed*, in the following sense: let  $\Theta$  be a closed positive  $(p, p)$ -current and let  $A$  be a complete pluripolar subset of  $X$ . Then  $\mathbf{1}_{X \setminus A} \Theta$  and thus also  $\mathbf{1}_A \Theta$  are closed.

We conclude this section with a log-concavity property of non-pluripolar products. Let  $T_1, \dots, T_n$  be closed positive  $(1, 1)$ -currents with locally bounded potentials near  $0 \in \mathbb{C}^n$  and let  $\mu$  be a positive measure. Suppose also given non-negative measurable functions  $f_j$  such that

$$T_j^n \geq f_j \mu, \quad j = 1, \dots, n. \tag{1.3}$$

Theorem 1.3 of [21] then implies that

$$T_1 \wedge \dots \wedge T_n \geq (f_1 \dots f_n)^{1/n} \mu. \tag{1.4}$$

It is in fact a standard variation of the Brunn–Minkowski inequality that (1.3) implies (1.4) when the whole data is smooth, and [21] reduces to this case by an adequate regularization process. As an easy consequence of Dinew’s result, we get the following version for non-pluripolar products.

PROPOSITION 1.11. *Let  $T_1, \dots, T_n$  be closed positive  $(1, 1)$ -currents, let  $\mu$  be a positive measure and assume given for each  $j=1, \dots, n$  a non-negative measurable function  $f_j$  such that*

$$\langle T_j^n \rangle \geq f_j \mu.$$

*Then we have*

$$\langle T_1 \wedge \dots \wedge T_n \rangle \geq (f_1 \dots f_n)^{1/n} \mu.$$

Here non-pluripolar products are considered as (possibly locally infinite) positive Borel measures.

*Proof.* Write  $T_j = dd^c u_j$ , with  $u_j$  psh, and consider the plurifine open subset

$$O_k := \bigcap_{j=1}^n \{u_j > -k\}.$$

Since non-pluripolar products are local in plurifine topology, we get for each  $j$  and  $k$ ,

$$(dd^c \max\{u_j, -k\})^n \geq \mathbf{1}_{O_k} f_j \mu.$$

Hence

$$dd^c \max\{u_1, -k\} \wedge \dots \wedge dd^c \max\{u_n, -k\} \geq \mathbf{1}_{O_k} (f_1 \dots f_n)^{1/n} \mu$$

by Dinew's result, or equivalently by plurifine locality

$$\langle dd^c u_1 \wedge \dots \wedge dd^c u_n \rangle \geq \mathbf{1}_{O_k} (f_1 \dots f_n)^{1/n} \mu.$$

Since this is valid for all  $k$  and since the complement of  $\bigcup_{k=1}^{\infty} O_k$  is pluripolar, the result follows.  $\square$

From now on we will work in the global setting where  $X$  is a compact Kähler manifold.

### 1.3. Positive cohomology classes

Let  $X$  be a compact Kähler manifold and let  $\alpha \in H^{1,1}(X, \mathbb{R})$  be a real  $(1, 1)$ -cohomology class.

Recall that  $\alpha$  is said to be *pseudo-effective* (*psef* for short) if it can be represented by a closed positive  $(1, 1)$ -current  $T$ . If  $\theta$  is a given smooth representative of  $\alpha$ , then any such current can be written as  $T = \theta + dd^c \varphi$ , where  $\varphi$  is thus a  $\theta$ -psh function by definition, and will sometimes be referred to as a *global potential* of  $T$ . Global potentials only depend on the choice of  $\theta$  up to a smooth function on  $X$ .

On the other hand,  $\alpha$  is *nef* if it lies in the closure of the Kähler cone. Such a class can thus be represented by *smooth* forms with arbitrarily small negative parts, and a compactness argument then shows that  $\alpha$  contains a positive current, i.e. nef implies psef. Note however that the negative part of smooth representatives cannot be taken to be 0 in general. In fact, Demailly, Peternell and Schneider have shown in [20] that a classical construction of Serre yields an example of a smooth curve  $C$  on a projective surface whose cohomology class is nef but contains only one positive current, to wit (the integration current on)  $C$  itself.

The set of all nef classes is by definition a closed convex cone in  $H^{1,1}(X, \mathbb{R})$ , whose interior is none but the Kähler cone. The set of all psef classes also forms a closed convex cone, and its interior is by definition the set of all *big* cohomology classes. In other words, a class  $\alpha$  is big if and only if it can be represented by a *strictly positive* current, i.e. a closed current  $T$  that dominates some (small enough) smooth strictly positive form on  $X$ .

#### 1.4. Comparison of singularities

If  $T$  and  $T'$  are two closed positive currents on  $X$ , then  $T$  is said to be *more singular* than  $T'$  if their global potentials satisfy  $\varphi \leq \varphi' + O(1)$ . By the *singularity type* of  $T$ , we will mean its equivalence class with respect to the above notion of comparison of singularities.

A positive current  $T$  in a given psef class  $\alpha$  is now said to have *minimal singularities* (inside its cohomology class) if it is less singular than any other positive current in  $\alpha$ , and its  $\theta$ -psh potentials  $\varphi$  will correspondingly be said to have minimal singularities. Such  $\theta$ -psh functions with minimal singularities always exist, as was observed by Demailly. Indeed, the upper envelope

$$V_\theta := \sup\{\varphi \text{ } \theta\text{-psh} : \varphi \leq 0 \text{ on } X\}$$

of all non-positive  $\theta$ -psh functions obviously yields a positive current

$$\theta + dd^c V_\theta$$

with minimal singularities (note that  $V_\theta$  is upper semi-continuous since  $V_\theta^*$  is a candidate in the envelope).

We stress that currents with minimal singularities in a given class  $\alpha$  are in general far from unique. Indeed currents with minimal singularities in a class  $\alpha$  admitting a smooth non-negative representative  $\theta \geq 0$  (for instance a Kähler class) are exactly those with bounded potentials.

Currents with minimal singularities are stable under pull-back.

PROPOSITION 1.12. *Let  $\pi: Y \rightarrow X$  be a surjective morphism between compact Kähler manifolds and  $\theta$  be a smooth closed  $(1, 1)$ -form on  $X$ . If  $\varphi$  is a  $\theta$ -psh function with minimal singularities on  $X$ , then the  $\pi^*\theta$ -psh function  $\varphi \circ \pi$  also has minimal singularities.*

*Proof.* Let  $\psi$  be a  $\pi^*\theta$ -psh function. For a regular value  $x \in X$  of  $\pi$  we set

$$\tau(x) := \sup_{\pi(y)=x} \psi(y).$$

It is standard to show that  $\tau$  uniquely extends to a  $\theta$ -psh function on  $X$ , so that  $\tau \leq \varphi + O(1)$ . But this clearly implies that  $\psi \leq \varphi \circ \pi + O(1)$  as was to be shown.  $\square$

If  $\pi$  furthermore has connected fibers then any  $\pi^*\theta$ -psh function on  $Y$  is of the form  $\varphi \circ \pi$ , where  $\varphi$  is a  $\theta$ -psh function on  $X$ .

A positive current  $T = \theta + dd^c \varphi$  and its global potential  $\varphi$  are said to have *analytic singularities* if there exists  $c > 0$  such that (locally on  $X$ )

$$\varphi = \frac{c}{2} \log \sum_{j=1}^N |f_j|^2 + u,$$

where  $u$  is smooth and  $f_1, \dots, f_N$  are local holomorphic functions. The coherent ideal sheaf  $\mathcal{I}$  locally generated by these functions (in fact, its integral closure) is then globally defined, and the singularity type of  $T$  is thus encoded by the data formally denoted by  $\mathcal{I}^c$ .

Demailly’s fundamental regularization theorem [16] states that a given  $\theta$ -psh function  $\varphi$  can be approximated from above by a sequence  $\varphi_k$  of  $(\theta + \varepsilon_k \omega)$ -psh functions with analytic singularities,  $\omega$  denoting some auxiliary Kähler form. In fact the singularity type of  $\varphi_k$  is described by the  $k$ th root of the multiplier ideal sheaf of  $k\varphi$ .

This result implies in particular that a big class  $\alpha$  always contains strictly positive currents with analytic singularities. It follows that there exists a Zariski open subset  $\Omega$  of  $X$  on which global potentials of currents with minimal singularities in  $\alpha$  are all locally bounded. The following definition is extracted from [11].

*Definition 1.13.* If  $\alpha$  is a big class, we define its *ample locus*  $\text{Amp}(\alpha)$  as the set of points  $x \in X$  such that there exists a strictly positive current  $T \in \alpha$  with analytic singularities which is smooth around  $x$ .

The ample locus  $\text{Amp}(\alpha)$  is a Zariski open subset by definition, and it is non-empty thanks to Demailly’s regularization result. In fact it is shown in [11] that there exists a strictly positive current  $T \in \alpha$  with analytic singularities whose smooth locus is precisely  $\text{Amp}(\alpha)$ . Note that  $\text{Amp}(\alpha)$  coincides with the complement of the so-called *augmented base locus*  $\mathbb{B}_+(\alpha)$  (see [23]) when  $\alpha = c_1(L)$  is the first Chern class of a big line bundle  $L$ .

### 1.5. Global currents with small unbounded locus

Let  $X$  be a compact Kähler manifold, and let  $T_1, \dots, T_p$  be closed positive  $(1, 1)$ -currents on  $X$  with small unbounded locus (see Definition 1.2). We can then find a closed complete pluripolar subset  $A$  such that each  $T_j$  has locally bounded potentials on  $X \setminus A$ . The content of Proposition 1.6 in this case is that the Bedford–Taylor product  $T_1 \wedge \dots \wedge T_p$ , which is well defined on  $X \setminus A$ , has finite total mass on  $X \setminus A$ :

$$\int_{X \setminus A} T_1 \wedge \dots \wedge T_p \wedge \omega^{n-p} < \infty,$$

for any Kähler form  $\omega$  on  $X$ .

We now establish a somewhat technical-looking integration-by-parts theorem in this context that will be crucial in order to establish the basic properties of weighted energy functionals in Proposition 2.8.

**THEOREM 1.14.** *Let  $A \subset X$  be a closed complete pluripolar subset, and let  $\Theta$  be a closed positive current on  $X$  of bidimension  $(1, 1)$ . Let  $\varphi_i$  and  $\psi_i$ ,  $i=1, 2$ , be quasi-psh functions on  $X$  that are locally bounded on  $X \setminus A$ . If  $u := \varphi_1 - \varphi_2$  and  $v := \psi_1 - \psi_2$  are globally bounded on  $X$ , then*

$$\int_{X \setminus A} u dd^c v \wedge \Theta = \int_{X \setminus A} v dd^c u \wedge \Theta = - \int_{X \setminus A} dv \wedge d^c u \wedge \Theta.$$

Note that the (signed) measure  $dd^c \varphi_i \wedge \Theta$  defined on  $X \setminus A$  has finite total mass, hence so does  $dd^c u \wedge \Theta$ , and  $\int_{X \setminus A} v dd^c u \wedge \Theta$  is therefore well defined since  $v$  is bounded (and defined everywhere). The last integral is also well defined by Lemma 1.15 below and the Cauchy–Schwarz inequality.

*Proof.* As is well known, if  $\varphi \geq 0$  is a bounded psh function, the identity

$$dd^c \varphi^2 = 2 d\varphi \wedge d^c \varphi + 2\varphi dd^c \varphi$$

enables one to define  $d\varphi \wedge d^c \varphi \wedge \Theta$  as a positive measure, which shows that

$$d^c \varphi \wedge \Theta := d^c(\varphi \Theta)$$

is a current of order 0 by the Cauchy–Schwarz inequality. By linearity, we can therefore make sense of  $v d^c u \wedge \Theta$  and  $du \wedge d^c u \wedge \Theta$  as currents of order 0 on  $X \setminus A$ . By Lemma 1.15 below, the positive measure  $du \wedge d^c u \wedge \Theta$  on  $X \setminus A$  has finite total mass, and hence so does  $v d^c u \wedge \Theta$ , by the Cauchy–Schwarz inequality.

We claim that

$$d[v d^c u \wedge \Theta] = dv \wedge d^c u \wedge \Theta + v dd^c u \wedge \Theta \quad \text{on } X \setminus A. \quad (1.5)$$



To show this, we argue locally. We can then assume that  $u$  and  $v$  are locally bounded psh functions by bilinearity. Let  $u_k$  and  $v_k$  be smooth psh functions decreasing towards  $u$  and  $v$ , respectively. Since  $d^c u \wedge \Theta$  has order 0, we see that  $v_k d^c u \wedge \Theta$  converges to  $v d^c u \wedge \Theta$  by monotone convergence. As the right-hand side of the desired formula is continuous along decreasing sequences of bounded psh functions by Bedford–Taylor’s theorem, we can thus assume that  $v$  is smooth and psh. Now  $u_k \Theta$  converges to  $u \Theta$ , and thus  $v d^c u_k \wedge \Theta \rightarrow v d^c u_k \wedge \Theta$ , and we are done by another application of Bedford–Taylor’s theorem to the right-hand side.

In particular (1.5) shows that  $d[v d^c u \wedge \Theta]$  is a current of order 0 on  $X \setminus A$ . Let us somewhat abusively denote by  $\mathbf{1}_{X \setminus A} v d^c u \wedge \Theta$  and  $\mathbf{1}_{X \setminus A} d[v d^c u \wedge \Theta]$  the trivial extensions to  $X$ . We are going to show that

$$d[\mathbf{1}_{X \setminus A} v d^c u \wedge \Theta] = \mathbf{1}_{X \setminus A} d[v d^c u \wedge \Theta] \tag{1.6}$$

as currents on  $X$ . By Stokes’ theorem we will thus have

$$\int_{X \setminus A} d[v d^c u \wedge \Theta] = 0,$$

and the desired formulae will follow by (1.5).

The proof of (1.6) will again follow rather closely Sibony’s proof of the Skoda–El Mir extension theorem as presented in [18, pp. 159–161]. Note that (1.6) is a local equality, and that it clearly holds on  $X \setminus A$ . We will thus focus on a fixed small enough neighborhood  $U$  of a given point  $0 \in A$ . Since  $A$  is complete pluripolar around 0, we may by definition find a psh function  $\tau$  on  $U$  such that  $\{\tau = -\infty\} = A$  near 0. Pick a smooth non-decreasing convex function  $\chi$  on  $\mathbb{R}$  such that  $\chi(t) = 0$  for  $t \leq \frac{1}{2}$  and  $\chi(1) = 1$ , and set

$$w_k := \chi(e^{\tau/k}).$$

Then  $0 \leq w_k \leq 1$  is a bounded psh function on  $U$  vanishing on  $A$ , and the sequence  $w_k$  increases to 1 pointwise on  $U \setminus A$  as  $k \rightarrow \infty$ . Now choose a smooth non-decreasing function  $0 \leq \theta(t) \leq 1$ ,  $t \in \mathbb{R}$ , such that  $\theta \equiv 0$  near 0 and  $\theta \equiv 1$  near 1 (we of course cannot require  $\theta$  to be convex here). It then follows that  $\theta(w_k)$  vanishes near  $A$  and  $\theta(w_k) \rightarrow 1$  pointwise on  $U \setminus A$ , whereas  $\theta'(w_k) \rightarrow 0$  pointwise on  $U$ . As a consequence we get

$$\theta(w_k) v d^c u \wedge \Theta \rightarrow \mathbf{1}_{U \setminus A} v d^c u \wedge \Theta$$

and

$$\theta(w_k) d[v d^c u \wedge \Theta] \rightarrow \mathbf{1}_{U \setminus A} d[v d^c u \wedge \Theta]$$

on  $U$  as  $k \rightarrow \infty$ . Since

$$d[\theta(w_k)v d^c u \wedge \Theta] = \theta'(w_k)v dw_k \wedge d^c u \wedge \Theta + \theta(w_k) d[v d^c u \wedge \Theta],$$

by Lemma 1.9, we are thus reduced to showing that

$$\theta'(w_k)v dw_k \wedge d^c u \wedge \Theta \rightarrow 0$$

on  $U$  as  $k \rightarrow \infty$ . But if  $\psi \geq 0$  is a given smooth function on  $U$  with compact support, the Cauchy–Schwarz inequality implies that

$$\left| \int_X \psi \theta'(w_k)v dw_k \wedge d^c u \wedge \Theta \right|^2 \leq \left( \int_X \psi dw_k \wedge d^c w_k \wedge \Theta \right) \left( \int_X \psi \theta'(w_k)^2 v^2 du \wedge d^c u \wedge \Theta \right).$$

Since

$$2 \int_X \psi dw_k \wedge d^c w_k \wedge \Theta \leq \int_X \psi dd^c w_k^2 \wedge \Theta = \int_X w_k^2 dd^c \psi \wedge \Theta$$

is bounded, whereas

$$\int_X \psi \theta'(w_k)^2 v^2 du \wedge d^c u \wedge \Theta \rightarrow 0$$

by dominated convergence, as  $\theta'(w_k)^2 \rightarrow 0$  pointwise, one concludes the proof.  $\square$

LEMMA 1.15. *Let  $\varphi_1$  and  $\varphi_2$  be two quasi-psh functions on  $X$  that are locally bounded on an open subset  $U$ . If the difference  $u := \varphi_1 - \varphi_2$  is globally bounded on  $X$ , then*

$$\int_U du \wedge d^c u \wedge \Theta < \infty.$$

*Proof.* By Demailly’s approximation theorem [16] we can find decreasing sequences of smooth functions  $\varphi_i^{(k)}$  decreasing towards  $\varphi_i$ , as  $k \rightarrow \infty$ , and such that  $dd^c \varphi_i^{(k)} \geq -\omega$  for some fixed large enough Kähler form  $\omega$  (see also [9]). Since  $u = \varphi_1 - \varphi_2$  is bounded by assumption, it follows that  $u_k := \varphi_1^{(k)} - \varphi_2^{(k)}$  can be taken to be uniformly bounded with respect to  $k$ . We can thus assume that  $0 \leq u_k \leq C$  for all  $k$ .

Now the formula  $dd^c v^2 = 2v dd^c v + 2dv \wedge d^c v$  and Bedford–Taylor’s continuity theorem imply by polarization identities that

$$\lim_{k \rightarrow \infty} d\varphi_i^{(k)} \wedge d^c \varphi_j^{(k)} \wedge \Theta = d\varphi_i \wedge d^c \varphi_j \wedge \Theta$$

weakly on  $U$  for any two indices  $i$  and  $j$ . Hence also

$$du_k \wedge d^c u_k \wedge \Theta \rightarrow du \wedge d^c u \wedge \Theta$$

weakly on  $U$ . We thus get

$$\int_U du \wedge d^c u \wedge \Theta \leq \liminf_{k \rightarrow \infty} \int_X du_k \wedge d^c u_k \wedge \Theta,$$

and our goal is to show that the right-hand integrals are uniformly bounded. Now Stokes' theorem yields

$$\int_X du_k \wedge d^c u_k \wedge \Theta = - \int_X u_k dd^c u_k \wedge \Theta = \int_X u_k dd^c \varphi_2^{(k)} \wedge \Theta - \int_X u_k dd^c \varphi_1^{(k)} \wedge \Theta.$$

For each  $i=1, 2$  we have

$$0 \leq u_k(\omega + dd^c \varphi_i^{(k)}) \leq C(\omega + dd^c \varphi_i^{(k)}).$$

Hence

$$0 \leq \int_X u_k(\omega + dd^c \varphi_i^{(k)}) \wedge \Theta \leq C \int_X \omega \wedge \Theta$$

so that

$$-C \int_X \omega \wedge \Theta \leq \int_X u_k dd^c \varphi_i^{(k)} \wedge \Theta \leq C \int_X \omega \wedge \Theta$$

for all  $k$ , and the uniform boundedness of  $\int_X du_k \wedge d^c u_k \wedge \Theta$  follows.  $\square$

The following crucial result shows that non-pluripolar masses are basically non-increasing with singularities, at least for currents with small unbounded locus.

**THEOREM 1.16.** *For  $j=1, \dots, p$ , let  $T_j$  and  $T'_j$  be two cohomologous closed positive  $(1, 1)$ -currents with small unbounded locus, and assume also that  $T_j$  is less singular than  $T'_j$ . Then the cohomology classes of their non-pluripolar products satisfy*

$$\{\langle T_1 \wedge \dots \wedge T_p \rangle\} \geq \{\langle T'_1 \wedge \dots \wedge T'_p \rangle\}$$

in  $H^{p,p}(X, \mathbb{R})$ , with  $\geq$  meaning that the difference is pseudo-effective, i.e. representable by a closed positive  $(p, p)$ -current.

The statement of course makes sense for arbitrary closed positive  $(1, 1)$ -currents, and we certainly expect it to hold true in general, but we are only able for the moment to prove it in the special case of currents with small unbounded locus.

*Proof.* By duality, this is equivalent to showing that

$$\int_{X \setminus A} T_1 \wedge \dots \wedge T_p \wedge \tau \geq \int_{X \setminus A} T'_1 \wedge \dots \wedge T'_p \wedge \tau$$

for every positive smooth  $dd^c$ -closed  $(n-p, n-p)$ -form  $\tau$ , where  $A$  is a closed complete pluripolar set outside of which all currents  $T_i$  and  $T'_i$  have locally bounded potentials.

Replacing successively  $T_i$  by  $T'_i$ , we can assume that  $T_i=T'_i$  for  $i>1$ . The bidimension- $(1, 1)$  current

$$\Theta := \langle T_2 \wedge \dots \wedge T_p \rangle \wedge \tau$$

is  $dd^c$ -closed and positive. Writing  $T_1=\theta+dd^c\varphi$  and  $T'_1=\theta+dd^c\varphi'$ , where  $\theta$  is a smooth form in their common cohomology class, we are reduced to showing that

$$\int_{X \setminus A} dd^c\varphi \wedge \Theta \geq \int_{X \setminus A} dd^c\varphi' \wedge \Theta$$

if  $\varphi \geq \varphi'$  are quasi-psh functions on  $X$ , locally bounded on  $X \setminus A$ , and  $\Theta$  is a  $dd^c$ -closed positive bidimension- $(1, 1)$  current on  $X$ . Let  $\psi$  be a quasi-psh function such that  $A = \{\psi = -\infty\}$ . Replacing  $\varphi'$  by  $\varphi' + \varepsilon\psi$  and letting  $\varepsilon \rightarrow 0$  in the end, we may further assume that  $\varphi' - \varphi \rightarrow -\infty$  near  $A$ . For each  $k$  the function  $\psi_k := \max\{\varphi', \varphi - k\}$  thus coincides with  $\varphi - k$  in some neighborhood of  $A$ , so that

$$\int_{X \setminus A} dd^c\psi_k \wedge \Theta = \int_{X \setminus A} dd^c\varphi \wedge \Theta$$

by Stokes' theorem, since (the trivial extension of)  $dd^c(\varphi - \psi_k) \wedge \Theta$  is the exterior derivative of a current with compact support on  $X \setminus A$ . On the other hand

$$\lim_{k \rightarrow \infty} dd^c\psi_k \wedge \Theta = dd^c\varphi' \wedge \Theta$$

weakly on  $X \setminus A$ , and the result follows. □

As a consequence, we can give the following definition.

*Definition 1.17.* Let  $\alpha_1, \dots, \alpha_p \in H^{1,1}(X, \mathbb{R})$  be big cohomology classes, and for all  $i=1, \dots, p$ , let  $T_{i,\min} \in \alpha_i$  be a positive current with minimal singularities. Then the cohomology class of the non-pluripolar product  $\langle T_{1,\min} \wedge \dots \wedge T_{p,\min} \rangle$  is independent of the choice of  $T_{i,\min} \in \alpha_i$  with minimal singularities. It will be denoted by

$$\langle \alpha_1 \dots \alpha_p \rangle \in H^{p,p}(X, \mathbb{R})$$

and called the *positive product* of the  $\alpha_i$ . If  $\alpha_1, \dots, \alpha_n$  are merely psef, we set

$$\langle \alpha_1 \dots \alpha_p \rangle := \lim_{\varepsilon \rightarrow 0} \langle (\alpha_1 + \varepsilon\beta) \dots (\alpha_p + \varepsilon\beta) \rangle,$$

where  $\beta \in H^{1,1}(X, \mathbb{R})$  is an arbitrary Kähler class.

As a special case, given a big class  $\alpha$ , the positive number

$$\text{vol}(\alpha) := \langle \alpha^n \rangle$$

is called the *volume* of  $\alpha$ .

Some explanations are in order. The positive product  $\langle \alpha_1 \dots \alpha_p \rangle$  is *not* multilinear, since the fact that  $T \in \alpha$  and  $T' \in \alpha'$  both have minimal singularities (in their respective cohomology classes) does not imply that  $T + T'$  has minimal singularities in  $\alpha + \alpha'$ . Positive products are however homogeneous and non-decreasing in each variable  $\alpha_i$  due to Theorem 1.16 when the  $\alpha_i$ 's are big (with respect to the partial order on cohomology induced by positive currents). These two properties imply in a completely formal way (cf. [13, Proposition 2.9]) that  $\langle \alpha_1 \dots \alpha_p \rangle$  depends *continuously* on the  $p$ -tuple  $(\alpha_1, \dots, \alpha_p)$  of *big* classes. We can thus extend positive products to psef classes  $\alpha_i$  as indicated, the existence of the limit and its independence on  $\beta$  being easy to check by monotonicity of positive products.

As an important special case, note that the positive product of *nef* classes  $\alpha_i$  coincides with their ordinary cup product, that is

$$\langle \alpha_1 \dots \alpha_p \rangle = \alpha_1 \dots \alpha_p.$$

The volume  $\text{vol}(\alpha)$  considered here coincides (as it should!) with the one introduced in [10]. In fact, the following result holds.

PROPOSITION 1.18. *Let  $\alpha_1, \dots, \alpha_p$  be big cohomology classes. Then there exist sequences  $T_j^{(k)} \in \alpha_j$  of strictly positive currents with analytic singularities such that*

$$\lim_{k \rightarrow \infty} \{ \langle T_1^{(k)} \wedge \dots \wedge T_p^{(k)} \rangle \} = \langle \alpha_1 \wedge \dots \wedge \alpha_p \rangle$$

in  $H^{p,p}(X, \mathbb{R})$ .

*Proof.* To keep notation simple we assume that  $\alpha_1 = \dots = \alpha_p$ . Let  $T_{\min} \in \alpha$  be a positive current with minimal singularities, let  $T_+ \in \alpha$  be a given strictly positive current with analytic singularities, and let  $\omega$  be a Kähler form such that  $T_+ \geq \omega$ . By Demailly's regularization theorem, there exists a sequence  $S_k \in \alpha$  of currents with analytic singularities whose global potentials decrease to that of  $T_{\min}$  and such that  $S_k \geq -\varepsilon_k \omega$ . Now let

$$T_k := (1 - \varepsilon_k)S_k + \varepsilon_k T_+,$$

so that  $T_k \geq \varepsilon_k^2 \omega$  is indeed strictly positive with analytic singularities. It is clear that  $T_k$  converges to  $T_{\min}$  as  $k \rightarrow \infty$ . By continuity of mixed Monge–Ampère operators along decreasing sequences of locally bounded psh functions, we get

$$T_k^p \rightarrow T_{\min}^p$$

weakly on  $X \setminus A$  with  $A$  as above. We infer that

$$\int_{X \setminus A} T_{\min}^p \wedge \omega^{n-p} \leq \liminf_{k \rightarrow \infty} \int_{X \setminus A} T_k^p \wedge \omega^{n-p}.$$

On the other hand, Theorem 1.16 yields

$$\int_{X \setminus A} T_{\min}^p \wedge \omega^{n-p} \geq \int_{X \setminus A} T_k^p \wedge \omega^{n-p},$$

since  $T_{\min}$  has minimal singularities, and we thus get

$$\lim_{k \rightarrow \infty} \langle T_k^p \rangle = \langle T_{\min}^p \rangle$$

as  $(p, p)$ -currents on  $X$ . The result follows.  $\square$

The next result corresponds to [10, Theorem 4.15], which generalizes Fujita's approximation theorem to arbitrary  $(1, 1)$ -classes. We reprove it here for convenience of the reader.

**PROPOSITION 1.19.** *Let  $\alpha$  be a big class on  $X$ . Then, for each  $\varepsilon > 0$ , there exist a modification  $\pi: X' \rightarrow X$ , where  $X'$  is a compact Kähler manifold, a Kähler class  $\beta$  on  $X'$  and an effective  $\mathbb{R}$ -divisor  $E$  such that*

- (a)  $\pi^* \alpha = \beta + \{E\}$ ;
- (b)  $\text{vol}(\alpha) - \varepsilon \leq \text{vol}(\beta) \leq \text{vol}(\alpha)$ .

*Proof.* By Proposition 1.18, there exists a positive current  $T \in \alpha$  with analytic singularities described by  $\mathcal{I}^c$  for some  $c > 0$  and some coherent ideal sheaf  $\mathcal{I}$  and a Kähler form  $\omega$  such that  $\int_X \langle T^n \rangle > \text{vol}(\alpha) - \varepsilon$  and  $T \geq \omega$ . By Hironaka's theorem, there exists a finite sequence of blow-ups with smooth centers  $\pi: X' \rightarrow X$  such that  $\pi^{-1} \mathcal{I} \cdot \mathcal{O}_{X'}$  is a principal ideal sheaf. The Siu decomposition of  $\pi^* T$  thus is

$$\pi^* T = \theta + [D],$$

where  $D$  is an effective  $\mathbb{R}$ -divisor and  $\theta \geq 0$  is a smooth form. It is also straightforward to see that  $\int_X \langle T^n \rangle = \int_{X'} \theta^n$ . The condition  $T \geq \omega$  yields  $\theta \geq \pi^* \omega$ , but this is not quite enough to conclude that the class of  $\theta$  is Kähler, in which case we would be done. However, since  $\pi$  is a finite composition of blow-ups with smooth centers, it is well known that there exists a  $\pi$ -exceptional effective  $\mathbb{R}$ -divisor  $F$  such that  $\pi^* \{\omega\} - \{F\}$  is a Kähler class (which shows by the way that  $X'$  is indeed Kähler). It follows that  $\gamma := \{\theta\} - \{F\}$  is also a Kähler class. It remains to set  $\beta := (1 - \varepsilon)\{\theta\} + \varepsilon\gamma$  and  $E := D + \varepsilon F$  with  $\varepsilon > 0$  small enough so that  $(1 - \varepsilon) \text{vol}(\alpha) \leq \text{vol}(\beta) \leq \text{vol}(\alpha)$  also holds (recall that  $\text{vol}$  is continuous on big classes).  $\square$

As a final remark, if  $\alpha = c_1(L)$  is the first Chern class of a big line bundle  $L$ , then it follows from a theorem of Fujita that  $\text{vol}(\alpha)$  as defined above coincides with the volume of  $L$  (see [10]), defined by

$$\text{vol}(L) = \lim_{k \rightarrow \infty} \frac{n!}{k^n} \dim H^0(X, L^{\otimes k}).$$

**1.6. Currents with full Monge–Ampère mass**

As we saw in Proposition 1.6, if  $T_1, \dots, T_p$  are arbitrary closed positive  $(1, 1)$ -currents on  $X$ , their non-pluripolar product  $\langle T_1 \wedge \dots \wedge T_p \rangle$  is always well defined. We want to give here a more convenient description of such global non-pluripolar products.

As a first observation, note that for any Kähler form  $\omega$  on  $X$  we have

$$\langle T_1 \wedge \dots \wedge T_p \rangle = \lim_{\varepsilon \rightarrow 0} \left\langle \bigwedge_{j=1}^p (T_j + \varepsilon \omega) \right\rangle,$$

simply by multilinearity of non-pluripolar products.

We can thus assume that the cohomology classes  $\alpha_j := \{T_j\}$  are big. Let  $T_{j,\min} \in \alpha_j$  be a positive current with minimal singularities, and recall that  $T_{j,\min}$  has small unbounded locus by Demailly’s regularization theorem, since  $\alpha_j$  is big. If we pick a smooth representative  $\theta_j \in \alpha_j$  and write

$$T_j = \theta_j + dd^c \varphi_j$$

and

$$T_{j,\min} = \theta_j + dd^c \varphi_{j,\min},$$

then we have the following useful description of the non-pluripolar product:

$$\langle T_1 \wedge \dots \wedge T_p \rangle = \lim_{k \rightarrow \infty} \mathbf{1}_{\bigcap_{j=1}^p \{\varphi_j > \varphi_{\min} - k\}} \left\langle \bigwedge_{j=1}^p (\theta_j + dd^c \max\{\varphi_j, \varphi_{j,\min} - k\}) \right\rangle, \tag{1.7}$$

where the right-hand side is in fact non-decreasing, so that convergence holds against any bounded measurable function. This fact is a straightforward consequence of the local character of non-pluripolar products in the plurifine topology, which in fact yields

$$\mathbf{1}_{\bigcap_{j=1}^p \{\varphi_j > \varphi_{\min} - k\}} \langle T_1 \wedge \dots \wedge T_p \rangle = \mathbf{1}_{\bigcap_{j=1}^p \{\varphi_j > \varphi_{\min} - k\}} \langle (\theta_j + dd^c \max\{\varphi_j, \varphi_{j,\min} - k\}) \rangle.$$

As a consequence, Theorem 1.16 yields the following result.

PROPOSITION 1.20. *If  $\alpha_1, \dots, \alpha_p \in H^{1,1}(X, \mathbb{R})$  are psef cohomology classes, then*

$$\{\langle T_1 \wedge \dots \wedge T_p \rangle\} \leq \langle \alpha_1 \dots \alpha_p \rangle$$

in  $H^{p,p}(X, \mathbb{R})$  for all positive currents  $T_i \in \alpha_i$ .

*Proof.* We can assume that each  $\alpha_i$  is big. By duality, we have to show that

$$\int_X \langle T_1 \wedge \dots \wedge T_p \rangle \wedge \tau \leq \langle \alpha_1 \dots \alpha_p \rangle \{\tau\}$$

for every positive  $dd^c$ -closed positive form  $\tau$ . But (1.7) shows that

$$\begin{aligned} \int_X \langle T_1 \wedge \dots \wedge T_p \rangle \wedge \tau &= \lim_{k \rightarrow \infty} \int_{\bigcap_{j=1}^p \{\varphi_j > \varphi_{\min} - k\}} \left\langle \bigwedge_{j=1}^p (\theta_j + dd^c \max\{\varphi_j, \varphi_{j,\min} - k\}) \right\rangle \wedge \tau \\ &\leq \lim_{k \rightarrow \infty} \int_X \left\langle \bigwedge_{j=1}^p (\theta_j + dd^c \max\{\varphi_j, \varphi_{j,\min} - k\}) \right\rangle \wedge \tau. \end{aligned}$$

The integrals in question are equal to

$$\langle \alpha_1 \dots \alpha_p \rangle \{\tau\}$$

for all  $k$ . Hence the result follows. □

When the  $\alpha_i$ 's are big, equality holds in Proposition 1.20 when each  $T_i \in \alpha_i$  has minimal singularities. In the general psef case, there might however not exist positive currents achieving equality. Indeed [20] provides an exemple of a nef class  $\alpha$  on a surface such that the only closed positive current in the class is the integration current  $T$  on a curve. It follows that  $\langle T \rangle = 0$ , whereas  $\langle \alpha \rangle = \alpha$  is non-zero.

*Definition 1.21.* A closed positive  $(1, 1)$ -current  $T$  on  $X$  with cohomology class  $\alpha$  will be said to have *full Monge–Ampère mass* if

$$\int_X \langle T^n \rangle = \text{vol}(\alpha).$$

We will have a much closer look at such currents in the coming sections.

**PROPOSITION 1.22.** *If  $T$  is a closed positive  $(1, 1)$ -current on  $X$  such that  $\langle T^n \rangle \neq 0$ , then its cohomology class  $\alpha = \{T\}$  is big.*

This follows immediately from [10, Theorem 4.8], which was itself an elaboration of the key technical result of [19].

We conjecture the following log-concavity property of total masses.

*Conjecture 1.23.* If  $T_1, \dots, T_n$  are closed positive  $(1, 1)$ -currents on  $X$ , then

$$\int_X \langle T_1 \wedge \dots \wedge T_n \rangle \geq \left( \int_X \langle T_1^n \rangle \right)^{1/n} \dots \left( \int_X \langle T_n^n \rangle \right)^{1/n}.$$

In particular, the function

$$T \mapsto \left( \int_X \langle T^n \rangle \right)^{1/n}$$

is concave.



Note that the conjecture holds true for currents with *analytic singularities*, since passing to a log-resolution of the singularities reduces it to the Khovanski–Teissier inequalities for nef classes proved in [17]. Reasoning as in [15, Theorem 5.5], the general case would follow from the Hodge index-type inequality

$$\int_X \langle T_1 \wedge T_2 \wedge \Theta \rangle \geq \left( \int_X \langle T_1^2 \wedge \Theta \rangle \right)^{1/2} \left( \int_X \langle T_2^2 \wedge \Theta \rangle \right)^{1/2}$$

with  $\Theta := T_3 \wedge \dots \wedge T_n$ . An important special case of the conjecture will be proved in Corollary 2.15 below.

## 2. Weighted energy classes

Let  $X$  be a compact Kähler manifold, and let  $T$  be a closed positive  $(1, 1)$ -current on  $X$ . We write  $T = \theta + dd^c \varphi$ , where  $\theta$  is a smooth representative of the cohomology class  $\alpha := \{T\}$  and  $\varphi$  is a  $\theta$ -psh function. Recall (Definition 1.21) that  $T$  is said to have full Monge–Ampère mass if

$$\int_X \langle T^n \rangle = \text{vol}(\alpha).$$

Note that this is always the case when  $\alpha$  is not big, since  $\text{vol}(\alpha) = 0$  in that case. Assuming from now on that  $\alpha$  is big, our goal will be to characterize the full Monge–Ampère mass property as a finite (weighted) energy condition for  $\varphi$  (Proposition 2.11). As a consequence we will prove continuity results for non-pluripolar products.

Since we focus on  $\varphi$  rather than  $T$ , we give the following definition.

*Definition 2.1.* Let  $\theta$  be a smooth closed  $(1, 1)$ -form on  $X$  and let  $\varphi$  be a  $\theta$ -psh function. The *non-pluripolar Monge–Ampère measure* of  $\varphi$  (with respect to  $\theta$ ) is

$$\text{MA}(\varphi) := \langle (\theta + dd^c \varphi)^n \rangle.$$

We say that  $\varphi$  has *full Monge–Ampère mass* if  $\theta + dd^c \varphi$  has full Monge–Ampère mass, that is if and only if the measure  $\text{MA}(\varphi)$  satisfies

$$\int_X \text{MA}(\varphi) = \text{vol}(\alpha).$$

Let us stress that  $\text{MA}(\varphi)$  is well defined for *any*  $\theta$ -psh function  $\varphi$ . Note that  $\theta$ -psh functions with minimal singularities have full Monge–Ampère mass.

From now on we fix a smooth closed  $(1, 1)$ -form  $\theta$  whose cohomology class  $\alpha \in H^{1,1}(X, \mathbb{R})$  is big. We also fix the choice of a  $\theta$ -psh function with minimal singularities  $\varphi_{\min}$ . If  $\varphi$  is a given  $\theta$ -psh function, we set

$$\varphi^{(k)} := \max\{\varphi, \varphi_{\min} - k\}.$$

The  $\theta$ -psh functions  $\varphi^{(k)}$  have minimal singularities, and they decrease pointwise to  $\varphi$ . We will call them the *canonical* approximants of  $\varphi$ . As was explained above, for each Borel subset  $E$  of  $X$  we have

$$\int_E \text{MA}(\varphi) := \lim_{k \rightarrow \infty} \int_{E \cap \{\varphi > \varphi_{\min} - k\}} \text{MA}(\varphi^{(k)}),$$

where the limit is increasing.

### 2.1. Comparison principles

We begin with the following generalized comparison principle, which will be a basic tool in what follows.

**PROPOSITION 2.2.** (Generalized comparison principle) *Let  $T_j = \theta_j + dd^c \varphi_j$  be a closed positive  $(1, 1)$ -current with cohomology class  $\{T_j\} =: \alpha_j$ , for every  $j = 0, \dots, p$ . Let also  $S_0 = \theta_0 + dd^c \psi_0$  be another positive current in  $\alpha_0$ . Then we have*

$$\begin{aligned} \int_{\{\varphi_0 < \psi_0\}} \langle S_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle &\leq \int_{\{\varphi_0 < \psi_0\}} \langle T_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle + \langle \alpha_0^{n-p} \cdot \alpha_1 \dots \alpha_p \rangle \\ &\quad - \int_X \langle T_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle. \end{aligned}$$

*Proof.* Since non-pluripolar products are local in the plurifine topology, we have

$$\begin{aligned} \langle \alpha_0^{n-p} \alpha_1 \dots \alpha_p \rangle &\geq \int_X \langle (\theta_0 + dd^c \max\{\varphi_0, \psi_0 - \varepsilon\})^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle \\ &\geq \int_{\{\varphi_0 < \psi_0 - \varepsilon\}} \langle S_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle + \int_{\{\varphi_0 > \psi_0 - \varepsilon\}} \langle T_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle \\ &\geq \int_{\{\varphi_0 < \psi_0 - \varepsilon\}} \langle S_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle + \int_X \langle T_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle \\ &\quad - \int_{\{\varphi_0 < \psi_0\}} \langle T_0^{n-p} \wedge T_1 \wedge \dots \wedge T_p \rangle, \end{aligned}$$

and the result follows by letting  $\varepsilon \rightarrow 0$  by monotone convergence. □

**COROLLARY 2.3.** *For any two  $\theta$ -psh functions  $\varphi$  and  $\psi$  we have*

$$\int_{\{\varphi < \psi\}} \text{MA}(\psi) \leq \int_{\{\varphi < \psi\}} \text{MA}(\varphi) + \text{vol}(\alpha) - \int_X \text{MA}(\varphi).$$

The main new feature of this generalized comparison principle is the additional “error term”  $\text{vol}(\alpha) - \int_X \text{MA}(\varphi)$ , which is non-negative by Proposition 1.20.

*Remark 2.4.* If  $\varphi$  and  $\psi$  have small unbounded locus (and conjecturally for arbitrary  $\theta$ -psh functions), the same proof plus Theorem 1.16 yield

$$\int_{\{\varphi < \psi\}} \text{MA}(\psi) \leq \int_{\{\varphi < \psi\}} \text{MA}(\varphi)$$

as soon as  $\varphi$  is less singular than  $\psi$ , that is  $\varphi \geq \psi + O(1)$ . Indeed Theorem 1.16 implies

$$\int_X \text{MA}(\max\{\varphi, \psi - \varepsilon\}) = \int_X \text{MA}(\varphi)$$

in that case, and the result follows from the above proof.

We deduce from the comparison principle its usual companion.

**COROLLARY 2.5.** (Domination principle) *Let  $\psi$  and  $\varphi$  be  $\theta$ -psh functions. If  $\varphi$  has minimal singularities and  $\psi \leq \varphi$  holds a.e. with respect to  $\text{MA}(\varphi)$ , then  $\psi \leq \varphi$  holds everywhere.*

*Proof.* Since  $\{\theta\}$  is big by assumption, we can choose a  $\theta$ -psh function  $\varrho$  such that  $\theta + dd^c \varrho \geq \omega$  for some (small enough) Kähler form  $\omega$ . It follows that  $\text{MA}(\varrho)$  dominates Lebesgue measure. We may also assume that  $\varrho \leq \varphi$ , as  $\varphi$  has minimal singularities.

Now let  $\varepsilon > 0$ . Since  $\varphi$  has minimal singularities, we have  $\text{vol}(\alpha) = \int_X \text{MA}(\varphi)$  and the comparison principle thus yields

$$\varepsilon^n \int_{\{\varphi < (1-\varepsilon)\psi + \varepsilon\varrho\}} \text{MA}(\varrho) \leq \int_{\{\varphi < (1-\varepsilon)\psi + \varepsilon\varrho\}} \text{MA}((1-\varepsilon)\psi + \varepsilon\varrho) \leq \int_{\{\varphi < (1-\varepsilon)\psi + \varepsilon\varrho\}} \text{MA}(\varphi).$$

But the latter integral is zero by assumption, as  $\varrho \leq \varphi$  implies that

$$\{\varphi < (1-\varepsilon)\psi + \varepsilon\varrho\} \subset \{\varphi < \psi\}.$$

Since  $\text{MA}(\varrho)$  dominates Lebesgue measure, we conclude that for each  $\varepsilon > 0$  we have

$$\varphi \geq (1-\varepsilon)\psi + \varepsilon\varrho$$

a.e. with respect to Lebesgue measure, and the result follows. □

*Remark 2.6.* We do not know whether the result still holds if we replace the assumption that  $\varphi$  has minimal singularities by  $\varphi \geq \psi + O(1)$ .

## 2.2. Weighted energy functionals

By a *weight function*, we mean a smooth increasing function  $\chi: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\chi(-\infty) = -\infty \quad \text{and} \quad \chi(t) = t \quad \text{for } t \geq 0.$$

In order to generalize to the present setting the energy classes defined in [14] and [27], we introduce the following weighted energy functionals.

*Definition 2.7.* Let  $\chi$  be a weight function. We define the  $\chi$ -energy of a  $\theta$ -psh function with minimal singularities  $\varphi$  as

$$E_\chi(\varphi) := \frac{1}{n+1} \sum_{j=0}^n \int_X (-\chi)(\varphi - \varphi_{\min}) \langle T^j \wedge T_{\min}^{n-j} \rangle, \quad (2.1)$$

with  $T = \theta + dd^c \varphi$  and  $T_{\min} = \theta + dd^c \varphi_{\min}$ .

Recall that we have fixed an arbitrary  $\theta$ -psh function  $\varphi_{\min}$  with minimal singularities on  $X$ . Note that the  $\chi$ -energy functional for  $\chi(t) = t$  is nothing but the Aubin–Mabuchi energy functional (up to a minus sign, cf. [2], [33], and [6] for the extension to the singular setting).

The next proposition contains important properties of the  $\chi$ -energy.

**PROPOSITION 2.8.** *Let  $\chi$  be a convex weight function. Then the following properties hold.*

(i) *For any  $\theta$ -psh function  $\varphi$  with minimal singularities the sequence*

$$j \mapsto \int_X (-\chi)(\varphi - \varphi_{\min}) \langle T^j \wedge T_{\min}^{n-j} \rangle$$

*as in (2.1) is non-decreasing. In particular we have*

$$E_\chi(\varphi) \leq \int_X (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) \leq (n+1)E_\chi(\varphi)$$

*if  $\varphi \leq \varphi_{\min}$ .*

(ii) *The  $\chi$ -energy  $\varphi \mapsto E_\chi(\varphi)$  is non-increasing.*

(iii) *If  $\varphi_k$  decreases to  $\varphi$  with minimal singularities, then  $E_\chi(\varphi_k) \rightarrow E_\chi(\varphi)$ .*

*Proof.* Let  $u := \varphi - \varphi_{\min} \leq 0$ . Note that  $\chi(u) + \varphi_{\min}$  is also a  $\theta$ -psh function with minimal singularities, since

$$dd^c \chi(u) = \chi'(u) dd^c u + \chi''(u) du \wedge d^c u \geq -(\theta + dd^c \varphi_{\min}). \quad (2.2)$$

It follows that the bounded functions  $u$  and  $\chi(u)$  are both differences of quasi-psh functions which are locally bounded on the complement  $X \setminus A$  of a closed complete pluripolar subset  $A$ . We can thus apply the integration-by-parts formula of Theorem 1.14 to get

$$\begin{aligned} \int_X (-\chi)(\varphi - \varphi_{\min}) \langle T^{j+1} \wedge T_{\min}^{n-j-1} \rangle &= \int_{X \setminus A} (-\chi)(u) (T_{\min} + dd^c u) \wedge T^j \wedge T_{\min}^{n-j-1} \\ &= \int_{X \setminus A} (-\chi)(\varphi - \varphi_{\min}) T^j \wedge T_{\min}^{n-j} \\ &\quad + \int_{X \setminus A} \chi'(u) du \wedge d^c u \wedge T^j \wedge T_{\min}^{n-j-1} \\ &\geq \int_X (-\chi)(\varphi - \varphi_{\min}) \langle T^j \wedge T_{\min}^{n-j} \rangle \end{aligned}$$

for  $j=0, \dots, n-1$ , and (i) follows.

In order to prove (ii), let  $v \geq 0$  be a bounded function such that  $\varphi + v$  is still  $\theta$ -psh. We will show that the derivative at  $t=0$  of the function  $t \mapsto e(t) := E_\chi(\varphi + tv)$  is  $\leq 0$ . Indeed integration by parts yields

$$\begin{aligned} -e'(0) &= \sum_{j=0}^n \int_{X \setminus A} (v\chi'(u) T^j \wedge T_{\min}^{n-j} + j\chi(u) dd^c v \wedge T^{j-1} \wedge T_{\min}^{n-j}) \\ &= \sum_{j=0}^n \int_{X \setminus A} (v\chi'(u) T^j \wedge T_{\min}^{n-j} + jv dd^c \chi(u) \wedge T^{j-1} \wedge T_{\min}^{n-j}) \\ &\geq \sum_{j=0}^n \int_{X \setminus A} (v\chi'(u) T^j \wedge T_{\min}^{n-j} + jv\chi'(u) (T - T_{\min}) \wedge T^{j-1} \wedge T_{\min}^{n-j}) \\ &= (n+1) \int_{X \setminus A} v\chi'(u) T^n \\ &\geq 0, \end{aligned}$$

by (2.2).

Finally, let us prove (iii). If  $\varphi_k$  decreases to  $\varphi$ , then  $\chi(\varphi_k - \varphi_{\min}) + \varphi_{\min}$  is a sequence of  $\theta$ -psh functions decreasing towards  $\chi(\varphi - \varphi_{\min}) + \varphi_{\min}$ . Thus Bedford–Taylor’s theorem implies that

$$\lim_{k \rightarrow \infty} T_k^j \wedge T_{\min}^{n-j} = T^j \wedge T_{\min}^{n-j}$$

and

$$\lim_{k \rightarrow \infty} \chi(\varphi_k - \varphi_{\min}) T_k^j \wedge T_{\min}^{n-j} = \chi(\varphi - \varphi_{\min}) T^j \wedge T_{\min}^{n-j}$$

weakly on  $X \setminus A$ . But Theorem 1.16 implies that the total masses also converge, since in fact

$$\int_{X \setminus A} T_k^j \wedge T_{\min}^{n-j} = \int_{X \setminus A} T^j \wedge T_{\min}^{n-j}$$

for all  $k$ , the thus trivial extensions converge on the whole of  $X$ , i.e.

$$\lim_{k \rightarrow \infty} \langle T_k^j \wedge T_{\min}^{n-j} \rangle = \langle T^j \wedge T_{\min}^{n-j} \rangle.$$

As  $\chi(\varphi_k - \varphi_{\min})$  is uniformly bounded, it follows from Bedford–Taylor’s monotone convergence theorem [4] that

$$\lim_{k \rightarrow \infty} \int_X \chi(\varphi_k - \varphi_{\min}) \langle T_k^j \wedge T_{\min}^{n-j} \rangle = \int_X \chi(\varphi - \varphi_{\min}) \langle T^j \wedge T_{\min}^{n-j} \rangle$$

for all  $j$ , which concludes the proof.  $\square$

*Definition 2.9.* Let  $\chi$  be a convex weight function. If  $\varphi$  is an arbitrary  $\theta$ -psh function, its  $\chi$ -energy is defined as

$$E_\chi(\varphi) := \sup_{\psi \geq \varphi} E_\chi(\psi) \in ]-\infty, \infty]$$

over all  $\psi \geq \varphi$  with minimal singularities. We say that  $\varphi$  has finite  $\chi$ -energy if

$$E_\chi(\varphi) < \infty.$$

We will see in Corollary 2.18 how to extend (2.1) to this more general picture.

**PROPOSITION 2.10.** *Let  $\chi$  be a convex weight.*

(i) *The  $\chi$ -energy functional  $\varphi \mapsto E_\chi(\varphi)$  is non-increasing and lower semi-continuous on  $\theta$ -psh functions for the  $L^1(X)$  topology.*

(ii)  *$\varphi \mapsto E_\chi(\varphi)$  is continuous along decreasing sequences.*

(iii) *If  $\varphi_j$  are  $\theta$ -psh functions converging to  $\varphi$  in the  $L^1(X)$  topology, then*

$$\sup_j E_\chi(\varphi_j) < \infty$$

*implies that  $\varphi$  has finite  $\chi$ -energy.*

*Proof.* Statements (ii) and (iii) are trivial consequences of (i). Let  $\varphi_j \rightarrow \varphi$  be a convergent sequence of  $\theta$ -psh functions, and set  $\tilde{\varphi}_j := (\sup_{k \geq j} \varphi_k)^*$ , so that  $\tilde{\varphi}_j \geq \varphi_j$  decreases pointwise to  $\varphi$ . If  $\psi \geq \varphi$  is a given  $\theta$ -psh function with minimal singularities, then  $\max\{\tilde{\varphi}_j, \psi\} \geq \max\{\varphi_j, \psi\}$  decreases to  $\max\{\varphi, \psi\} = \psi$ . Hence

$$E_\chi(\psi) = \lim_{j \rightarrow \infty} E_\chi(\max\{\tilde{\varphi}_j, \psi\}) \leq \liminf_{j \rightarrow \infty} E_\chi(\varphi_j),$$

by Proposition 2.8, and the result follows by the definition of  $E_\chi(\varphi)$ .  $\square$

As an important consequence, we get

$$E_\chi(\varphi) = \sup_k E_\chi(\varphi^{(k)})$$

for the “canonical” approximants  $\varphi^{(k)} := \max\{\varphi, \varphi_{\min} - k\}$  of  $\varphi$ . The connection between weighted energies and full Monge–Ampère mass is given by the following result.

PROPOSITION 2.11. *Let  $\varphi$  be a  $\theta$ -psh function.*

(i) *The function  $\varphi$  has full Monge–Ampère mass if and only if  $E_\chi(\varphi) < \infty$  for some convex weight  $\chi$ .*

(ii) *If  $\varphi$  has full Monge–Ampère mass and  $\chi$  is a given convex weight, then*

$$E_\chi(\varphi) < \infty \quad \text{if and only if} \quad \int_X (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) < \infty.$$

*Proof.* Upon subtracting a constant, we may assume that  $\varphi \leq \varphi_{\min}$ . Let

$$\varphi^{(k)} := \max\{\varphi, \varphi_{\min} - k\}.$$

In view of Proposition 2.8, we have to show that

$$m_k := \int_{\{\varphi \leq \varphi_{\min} - k\}} \text{MA}(\varphi^{(k)})$$

tends to 0 if and only if

$$\sup_k \int_X (-\chi)(\varphi^{(k)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) < \infty \tag{2.3}$$

for some convex weight  $\chi$ . But we have

$$\begin{aligned} \int_X (-\chi)(\varphi^{(k)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) &= |\chi(-k)| m_k + \int_{\{\varphi > \varphi_{\min} - k\}} (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) \\ &= |\chi(-k)| m_k + O(1) \end{aligned}$$

for some convex weight  $\chi$ . Indeed we may always choose  $\chi$  such that

$$\int_X (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) < \infty,$$

simply because  $\text{MA}(\varphi)$  puts no mass on the pluripolar set  $\{\varphi = -\infty\}$ . It follows that (2.3) holds for some convex weight if and only if there exists a convex weight  $\chi$  such that  $\chi(-k) = O(1/m_k)$ , which is indeed the case if and only if  $m_k \rightarrow 0$ .

Let us now prove (ii). Assume that  $E_\chi(\varphi) < \infty$ . Since  $\text{MA}(\varphi)$  is the increasing limit in the strong Borel topology of

$$\mathbf{1}_{\{\varphi > \varphi_{\min} - k\}} \text{MA}(\varphi^{(k)}),$$

and since  $\chi(\varphi^{(l)} - \varphi_{\min})$  is a bounded measurable function, we infer that

$$\begin{aligned} \int_X (-\chi)(\varphi^{(l)} - \varphi_{\min}) \text{MA}(\varphi) &= \lim_{k \rightarrow \infty} \int_{\{\varphi > \varphi_{\min} - k\}} (-\chi)(\varphi^{(l)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) \\ &\leq \lim_{k \rightarrow \infty} \int_{\{\varphi > \varphi_{\min} - k\}} (-\chi)(\varphi^{(k)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) \\ &\leq E_\chi(\varphi), \end{aligned}$$

as  $\varphi^{(k)} \leq \varphi^{(l)}$  for  $k \geq l$ , and we infer that

$$\int_X (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) < \infty$$

by monotone convergence as desired.

Conversely, if  $(-\chi)(\varphi - \varphi_{\min}) \in L^1(\text{MA}(\varphi))$  and  $\varphi$  has full Monge–Ampère mass, then Lemma 2.12 below implies that  $E_\chi(\varphi) = \sup_k E_\chi(\varphi^{(k)}) < \infty$ .  $\square$

**LEMMA 2.12.** *Let  $\varphi$  be a  $\theta$ -psh function with canonical approximants  $\varphi^{(k)}$ . Then for every weight function  $\chi$  we have*

$$\begin{aligned} \int_X (-\chi)(\varphi^{(k)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) &\leq \int_X (-\chi)(\varphi - \varphi_{\min}) \text{MA}(\varphi) \\ &\quad + (-\chi)(-k) \left( \text{vol}(\alpha) - \int_X \text{MA}(\varphi) \right). \end{aligned}$$

*Proof.* We have

$$\begin{aligned} \int_X (-\chi)(\varphi^{(k)} - \varphi_{\min}) \text{MA}(\varphi^{(k)}) &= (-\chi)(-k) \int_{\{\varphi \leq \varphi^{(k)}\}} \text{MA}(\varphi^{(k)}) \\ &\quad + \int_{\{\varphi > \varphi_{\min} - k\}} \chi(\varphi - \varphi_{\min}) \text{MA}(\varphi) \end{aligned}$$

(since  $\mathbf{1}_{\{\varphi > \varphi_{\min} - k\}} \text{MA}(\varphi^{(k)}) = \mathbf{1}_{\{\varphi > \varphi_{\min} - k\}} \text{MA}(\varphi)$  by plurifine locality)

$$\begin{aligned} &\leq (-\chi)(-k) \int_{\{\varphi \leq \varphi^{(k)}\}} \text{MA}(\varphi) \\ &\quad + (-\chi)(-k) \left( \text{vol}(\alpha) - \int_X \text{MA}(\varphi) \right) \\ &\quad + \int_{\{\varphi > \varphi_{\min} - k\}} \chi(\varphi - \varphi_{\min}) \text{MA}(\varphi), \end{aligned}$$

by the generalized comparison principle, and the result follows.  $\square$



**2.3. Energy classes of positive currents**

Since  $\chi$  is convex with  $0 \leq \chi' \leq 1$ , we have

$$\chi(t) \leq \chi(t+C) \leq \chi(t) + C$$

for all  $C > 0$ , and it easily follows that the condition that  $\varphi$  has finite  $\chi$ -energy does not depend on the choice of the function with minimal singularities  $\varphi_{\min}$  with respect to which  $E_\chi$  is defined.

*Definition 2.13.* Given a convex weight  $\chi$ , we denote by  $\mathcal{E}_\chi(\alpha)$  the set of positive currents  $T \in \alpha$  with finite  $\chi$ -energy, and by  $\mathcal{E}(\alpha)$  the set of positive currents with full Monge–Ampère mass.

Proposition 2.11 can thus be reformulated by saying that

$$\mathcal{E}(\alpha) = \bigcup_{\chi} \mathcal{E}_\chi(\alpha),$$

where  $\chi$  ranges over all convex weights.

**PROPOSITION 2.14.** *The energy classes satisfy the following properties:*

- (i) *If  $T \in \mathcal{E}_\chi(\alpha)$ , then also  $S \in \mathcal{E}_\chi(\alpha)$  for all positive currents  $S \in \alpha$  which are less singular than  $T$ .*
- (ii) *The set  $\mathcal{E}(\alpha)$  is an extremal face (in particular a convex subset) of the closed convex set of all positive currents in  $\alpha$ .*

Recall that an *extremal face*  $F$  of a convex set  $C$  is a convex subset of  $C$  such that for all  $x, y \in C$ ,  $tx + (1-t)y \in F$  for some  $0 < t < 1$  implies that  $x, y \in F$ .

*Proof.* Statement (i) follows by Proposition 2.11 above, since  $E_\chi$  is non-increasing. To prove (ii), we first take care of the extremality property. Let  $T, T' \in \alpha$  be two closed positive currents and  $0 < t < 1$  be such that  $tT + (1-t)T'$  has finite energy. We have to show that  $T$  also has finite energy (then exchange the roles of  $T$  and  $T'$ ). If  $T_{\min} \in \alpha$  is a positive current with minimal singularities, then  $tT + (1-t)T_{\min}$  is less singular than  $tT + (1-t)T'$ , thus it a fortiori has finite energy, i.e. we can assume that  $T' = T_{\min}$  has minimal singularities. Now write  $T = \theta + dd^c \varphi$  and  $T_{\min} = \theta + dd^c \varphi_{\min}$ , and set  $\varphi_t := t\varphi + (1-t)\varphi_{\min}$ . The assumption is that

$$\lim_{k \rightarrow \infty} \int_{\{\varphi_t \leq \varphi_{\min} - k\}} \text{MA}(\max\{\varphi_t, \varphi_{\min} - k\}) = 0,$$

and we are to show that the same holds with  $\varphi$  in place of  $\varphi_t$ . But note that

$$\max\{\varphi_t, \varphi_{\min} - k\} = \max\{t(\varphi - \varphi_{\min}), -k\} + \varphi_{\min} = t \max\{\varphi, \varphi_{\min} - k/t\} + (1-t)\varphi_{\min}.$$

Hence

$$\int_{\{\varphi_t \leq \varphi_{\min} - k\}} \text{MA}(\max\{\varphi_t, \varphi_{\min} - k\}) \geq t^n \int_{\{\varphi \leq \varphi_{\min} - k/t\}} \text{MA}(\max\{\varphi, \varphi_{\min} - k/t\}),$$

and the extremality follows.

The convexity of  $\mathcal{E}(\alpha)$  now follows exactly as in the proof of [27, Proposition 1.6].  $\square$

The convexity of  $\mathcal{E}(\alpha)$  would also follow from Conjecture 1.23 above. Conversely the following important special case of the conjecture can be settled using convexity.

**COROLLARY 2.15.** *Let  $\alpha$  be a big class, and let  $T_1, \dots, T_n \in \alpha$  be positive currents with full Monge–Ampère mass. Then*

$$\int_X \langle T_1 \wedge \dots \wedge T_n \rangle = \langle \alpha^n \rangle.$$

*Proof.* Upon scaling  $\alpha$ , we may assume that  $\langle \alpha^n \rangle = 1$ . For any  $t = (t_1, \dots, t_n) \in \mathbb{R}_+^n$  set

$$P(t_1, \dots, t_n) := \int_X \langle (t_1 T_1 + \dots + t_n T_n)^n \rangle,$$

which defines a homogeneous polynomial of degree  $n$  by multilinearity of non-pluripolar products. Since  $T_i \in \mathcal{E}(\alpha)$  for each  $i$ , convexity implies that  $t_1 T_1 + \dots + t_n T_n \in \mathcal{E}(\alpha)$  when the  $t_i$ 's lie on the simplex

$$\Delta := \{t \in \mathbb{R}_+^n : t_1 + \dots + t_n = 1\}.$$

We thus see that the polynomial

$$P(t_1, \dots, t_n) - (t_1 + \dots + t_n)^n$$

vanishes identically on  $\Delta$ . Since this polynomial is homogeneous of degree  $n$ , it therefore vanishes on  $\mathbb{R}_+^n$ , and this implies that

$$P(t_1, \dots, t_n) = (t_1 + \dots + t_n)^n$$

for all  $t$ . Identifying the coefficients of the monomial  $t_1 \dots t_n$  on both sides now yields the result.  $\square$

*Remark 2.16.* It is quite plausible that each  $\mathcal{E}_\chi(\alpha)$  is itself convex, and indeed for  $\chi(t) = t$  it simply follows from the convexity of the Aubin–Mabuchi energy functional  $E_1$  (cf. [6] for a proof in this singular situation). The functional  $E_\chi$  is however *not* convex when  $\chi$  is not affine, since  $E_\chi(\varphi + c)$  is rather a concave function of  $c \in \mathbb{R}$ .

**2.4. Continuity properties of Monge–Ampère operators**

The Monge–Ampère operator on functions of full Monge–Ampère mass satisfies the usual continuity properties along monotonic sequences, as shown in the following result.

**THEOREM 2.17.** *If  $\varphi$  is a  $\theta$ -psh function with full Monge–Ampère mass and  $\varphi_j$  is a sequence of  $\theta$ -psh functions with full Monge–Ampère mass decreasing (resp. increasing a.e.) to  $\varphi$ , then*

$$\text{MA}(\varphi_j) \rightarrow \text{MA}(\varphi)$$

as  $j \rightarrow \infty$ . If  $\varphi$  furthermore has finite  $\chi$ -energy for a given convex weight  $\chi$ , then

$$\chi(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) \rightarrow \chi(\varphi - \varphi_{\min}) \text{MA}(\varphi)$$

weakly as  $j \rightarrow \infty$ .

*Proof.* Theorem 2.17 has already been proved for functions with minimal singularities during the proof of Proposition 2.8. Let us first prove the first point in the general case. Choose  $\chi$  such that  $\varphi$  and  $\varphi_1$  both have finite  $\chi$ -energy, so that  $E_\chi(\varphi_j) \leq C$  uniformly by the monotonicity of  $E_\chi$ . Let  $h$  be a continuous function on  $X$ . For each  $k$  we have

$$\lim_{j \rightarrow \infty} \int_X h \text{MA}(\varphi_j^{(k)}) = \int_X h \text{MA}(\varphi^{(k)})$$

by the minimal singularities case. Thus it is enough to establish that

$$\lim_{k \rightarrow \infty} \int_X h \text{MA}(\psi^{(k)}) = \int_X h \text{MA}(\psi)$$

uniformly for all  $\psi \geq \varphi$ . We have

$$\left| \int_X h(\text{MA}(\psi^{(k)}) - \text{MA}(\psi)) \right| \leq \sup_X |h| \left( \int_{\{\psi \leq \varphi_{\min} - k\}} \text{MA}(\psi^{(k)}) + \int_{\{\psi \leq \varphi_{\min} - k\}} \text{MA}(\psi) \right),$$

since  $\text{MA}(\psi^{(k)}) = \text{MA}(\psi)$  on  $\{\psi > \varphi_{\min} - k\}$ . The first term is controlled by

$$|\chi|(-k)^{-1} \int_X (-\chi)(\psi^{(k)}) \text{MA}(\psi^{(k)}) \leq |\chi|(-k)^{-1} E_\chi(\varphi),$$

and hence tends to 0 uniformly with respect to  $\psi \geq \varphi$ . On the other hand, since  $\text{MA}(\psi)$  is the monotone and thus *strong* limit of  $\mathbf{1}_{\{\psi > \varphi_{\min} - l\}} \text{MA}(\psi^{(l)})$ , the second term can be written as

$$\int_{\{\psi \leq \varphi_{\min} - k\}} \text{MA}(\psi) = \lim_{l \rightarrow \infty} \int_{\{\varphi_{\min} - l < \psi \leq \varphi_{\min} - k\}} \text{MA}(\psi^{(l)}),$$

which is similarly controlled by

$$\limsup_{l \rightarrow \infty} |\chi|(-k)^{-1} E_{\chi}(\psi^{(l)}) \leq |\chi|(-k)^{-1} E_{\chi}(\varphi),$$

and the first part is thus proved. The same reasoning shows that

$$\chi(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) \rightarrow \chi(\varphi - \varphi_{\min}) \text{MA}(\varphi)$$

as soon as  $\varphi$  has finite  $\tilde{\chi}$ -energy for some convex weight  $\tilde{\chi} \gg \chi$ . But this is in fact automatic, since  $E_{\chi}(\varphi) < \infty$  is equivalent to  $(-\chi)(\varphi - \varphi_{\min}) \in L^1(\text{MA}(\varphi))$  by Proposition 2.11, as  $\varphi$  has full Monge–Ampère mass, and this integrability condition can always be improved to  $\tilde{\chi}(\varphi - \varphi_{\min}) \in L^1(\text{MA}(\varphi))$  for  $\tilde{\chi} \gg \chi$ , by a standard measure-theoretic result (see [27] for more details on such properties).  $\square$

As a consequence, we get a more explicit formula for the  $\chi$ -energy.

**COROLLARY 2.18.** *If  $\varphi$  has full Monge–Ampère mass and  $\chi$  is a given convex weight, then setting  $T = \theta + dd^c \varphi$  and  $T_{\min} = \theta + dd^c \varphi_{\min}$  we have*

$$E_{\chi}(\varphi) = \int_X (-\chi)(\varphi - \varphi_{\min}) \sum_{j=0}^n \langle T^j \wedge T_{\min}^{n-j} \rangle,$$

*in the sense that the left-hand side is finite if and only if the right-hand side is, and equality holds in this case.*

## 2.5. A useful asymptotic criterion

In the continuity theorem above, it is crucial that the functions dealt with have finite energy in order to ensure that no mass is lost towards the polar set in the limit process. For an arbitrary  $\theta$ -psh function, the condition  $\chi(\varphi - \varphi_{\min}) \in L^1(\text{MA}(\varphi))$  alone does not imply that  $\varphi$  has finite energy. For instance, if  $\omega$  denotes the Fubini–Study form on  $\mathbb{P}^1$ , a global potential of the current  $T := \delta_0$  yields a counterexample since  $\langle T \rangle = 0!$  This example is in fact a simple instance of the type of currents we will have to handle in the proof of Theorem 3.1.

It is a consequence of Proposition 2.10 that any  $L^1(X)$ -limit  $\varphi = \lim_{j \rightarrow \infty} \varphi_j$  of functions  $\varphi_j$  with *full Monge–Ampère mass* and such that

$$\sup_j \int_X (-\chi)(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) < \infty$$

has finite  $\chi$ -energy. This still holds for sequences with asymptotically full Monge–Ampère mass as shown by our next result which yields a practical criterion to check the finite-energy condition.

PROPOSITION 2.19. *Let  $\chi$  be a convex weight and  $\varphi_j \rightarrow \varphi$  be a sequence of  $\theta$ -psh functions converging in  $L^1(X)$  to a  $\theta$ -psh function  $\varphi$ . Assume that*

- (1)  $\int_X \text{MA}(\varphi_j) \rightarrow \text{vol}(\alpha)$ ;
- (2)  $\sup_j \int_X (-\chi)(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) < \infty$ .

*Then  $\varphi$  has finite  $\chi$ -energy.*

*Proof.* Lemma 2.12 above yields two constants  $A, B > 0$  such that

$$E_\chi(\varphi_j^{(k)}) \leq A + B(-\chi)(-k) \left( \text{vol}(\alpha) - \int_X \text{MA}(\varphi_j) \right).$$

Hence

$$E_\chi(\varphi^{(k)}) \leq \liminf_{j \rightarrow \infty} E_\chi(\varphi_j^{(k)}) \leq A$$

for all  $k$ , by Proposition 2.10, and the result follows. □

PROPOSITION 2.20. *Let  $\varphi_j$  be a sequence of arbitrary  $\theta$ -psh functions uniformly bounded above, and let  $\varphi := (\sup_j \varphi_j)^*$ . Suppose that  $\mu$  is a positive measure such that  $\text{MA}(\varphi_j) \geq \mu$  for all  $j$ . If  $\varphi$  has full Monge–Ampère mass, then  $\text{MA}(\varphi) \geq \mu$ .*

*Proof.* Set  $\psi_j := \max_{1 \leq i \leq j} \varphi_i$ , so that  $\psi_j$  increases a.e. to  $\varphi$ . It is a standard matter to show that  $\text{MA}(\psi_j) \geq \mu$  using plurifine locality (cf. [27, Corollary 1.10]). The desired result does however not follow directly from Theorem 2.17, since  $\psi_j$  does not a priori have full Monge–Ampère mass. But since  $\text{MA}(\psi_j)$  and  $\text{MA}(\psi_j^{(k)})$  coincide on  $\{\psi_j > \varphi_{\min} - k\}$  as Borel measures, we get

$$\text{MA}(\psi_j^{(k)}) \geq \mathbf{1}_{\{\psi_j > \varphi_{\min} - k\}} \text{MA}(\psi_j) \geq \mathbf{1}_{\{\psi_j > \varphi_{\min} - k\}} \mu \geq \mathbf{1}_{\{\psi_1 > \varphi_{\min} - k\}} \mu.$$

For each fixed  $k$ , Theorem 2.17 (in fact simply Bedford–Taylor’s theorem on a Zariski open subset plus preservation of total masses) yields

$$\lim_{j \rightarrow \infty} \text{MA}(\psi_j^{(k)}) = \text{MA}(\varphi^{(k)}).$$

Hence

$$\text{MA}(\varphi^{(k)}) \geq \mathbf{1}_{\{\psi_1 > \varphi_{\min} - k\}} \mu.$$

Now the left-hand side converges to  $\text{MA}(\varphi)$  weakly as  $k \rightarrow \infty$ , since  $\varphi$  is assumed to have finite Monge–Ampère energy, whereas the right-hand side tends to  $\mu$ , as the latter puts no mass on the pluripolar set  $\{\psi_1 = -\infty\}$ , being dominated by non-pluripolar measures  $\text{MA}(\varphi_j)$ . □

COROLLARY 2.21. *Let  $T_j \rightarrow T$  be a convergent sequence of positive currents in a given cohomology class  $\alpha$ , and assume that the limit current  $T$  has full Monge–Ampère mass. Assume that a positive measure  $\mu$  and non-negative functions  $f_j$  are given such that*

$$\langle T_j^n \rangle \geq f_j \mu$$

*holds for all  $j$ . Then we have*

$$\langle T^n \rangle \geq \left( \liminf_{j \rightarrow \infty} f_j \right) \mu.$$

*Proof.* Write  $T_j = \theta + dd^c \varphi_j$ , with  $\varphi_j$  normalized in some fixed way, and set

$$\tilde{\varphi}_j := \left( \sup_{k \geq j} \varphi_k \right)^*,$$

so that  $\tilde{\varphi}_j$  decreases pointwise to  $\varphi$  with  $T = \theta + dd^c \varphi$ . Since  $\varphi$  has full Monge–Ampère mass, so does each  $\tilde{\varphi}_j \geq \varphi$ , and hence  $\text{MA}(\tilde{\varphi}_j) \geq (\inf_{k \geq j} f_k) \mu$ , by Proposition 2.20. The result now follows by Theorem 2.17.  $\square$

### 3. Non-pluripolar measures are Monge–Ampère

We will say for convenience that a positive measure  $\mu$  is *non-pluripolar* if it puts no mass on pluripolar subsets. The range of the non-pluripolar Monge–Ampère operator is described by the following result.

THEOREM 3.1. *Let  $X$  be a compact Kähler manifold and let  $\alpha \in H^{1,1}(X, \mathbb{R})$  be a big cohomology class. Then for any non-pluripolar measure  $\mu$  such that  $\mu(X) = \text{vol}(\alpha)$ , there exists a unique closed positive current  $T \in \alpha$  such that*

$$\langle T^n \rangle = \mu.$$

Note that  $T$  automatically has full Monge–Ampère mass since

$$\int_X \langle T^n \rangle = \mu(X) = \text{vol}(\alpha)$$

by assumption.

In the sequel  $\theta$  denotes a smooth representative of  $\alpha$ . As before, we denote the non-pluripolar Monge–Ampère measure of a  $\theta$ -psh function  $\varphi$  by  $\text{MA}(\varphi) := \langle (\theta + dd^c \varphi)^n \rangle$ . We also choose once and for all a  $\theta$ -psh function  $\varphi_{\min}$  with minimal singularities.

**3.1. Outline of the existence proof**

We first present an outline of the existence proof, and provide technical details below. By Proposition 1.19, given  $\varepsilon > 0$ , there exists a decomposition

$$\pi^* \alpha = \beta + \{E\}$$

on a modification  $\pi: X' \rightarrow X$ , where  $\beta$  is a Kähler class such that

$$\text{vol}(\alpha) - \varepsilon \leq \text{vol}(\beta) \leq \text{vol}(\alpha)$$

and  $E$  is an effective  $\mathbb{R}$ -divisor on  $X'$  (the whole data depends on  $\varepsilon$ ). By the main result of [27] (i.e. the existence part of Theorem 3.1 for a Kähler class), there exists a positive current with finite energy  $S \in \beta$  such that

$$\langle S^n \rangle = c\mu'$$

holds on  $X'$  with

$$c := \frac{\text{vol}(\beta)}{\text{vol}(\alpha)} \leq 1$$

and  $\mu'$  denoting the non-pluripolar measure on  $X'$  induced by lifting  $\mu$  to the Zariski open subset where  $\pi$  is an isomorphism. We will first show that the pushed-forward current

$$\pi_*(S + [E]) =: \theta + dd^c \varphi$$

satisfies  $\text{MA}(\varphi) = \langle (\theta + dd^c \varphi)^n \rangle = c\mu$ , which is at least plausible since  $[E]$  vanishes outside a pluripolar subset. We would already be done at this stage if  $c$  were 1, i.e.  $\text{vol}(\beta) = \text{vol}(\alpha)$ , but this actually *never* happens unless  $\alpha$  is a Kähler class in the first place. On the other hand, we have obtained, at this point, a sequence of  $\theta$ -psh functions  $\varphi_j$  such that

$$\text{MA}(\varphi_j) = (1 - \varepsilon_j)\mu$$

with  $\varepsilon_j > 0$  decreasing towards 0. We will next prove that some  $L^1(X)$ -limit  $\varphi_\infty$  of the  $\varphi_j$ 's (appropriately normalized) solves  $\text{MA}(\varphi_\infty) = \mu$ , and setting  $T := \theta + dd^c \varphi_\infty$  then proves the theorem.

It is in fact enough to show that  $\varphi_\infty$  has full Monge–Ampère mass. Indeed Corollary 2.21 then implies that  $\text{MA}(\varphi_\infty) \geq \mu$ , which must be an equality since both measures have the same mass, and we will be done. As

$$\lim_{j \rightarrow \infty} \int_X \text{MA}(\varphi_j) = \text{vol}(\alpha),$$

Proposition 2.19 shows that it is enough to find a convex weight  $\chi$  such that

$$\sup_j \int_X (-\chi)(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) < \infty.$$

But we will prove in Proposition 3.2 below that non-pluripolarity of  $\mu$  yields a convex weight  $\chi$  such that  $\int_X (-\chi)(\psi) d\mu$  is uniformly bounded for all normalized  $\theta$ -psh functions  $\psi$ . Hence the result follows, since  $\text{MA}(\varphi_j) \leq \mu$  for all  $j$ .

**3.2. Existence proof: technical details**

Using the above notation, let  $\omega \in \beta$  be a Kähler form and write  $S = \omega + dd^c \psi$ . We first have to show that

$$\langle (\pi_*(S + [E]))^n \rangle = \pi_* \langle S^n \rangle,$$

that is

$$\langle (\theta + dd^c \varphi)^n \rangle = \pi_* \langle (\omega + dd^c \psi)^n \rangle. \tag{3.1}$$

At this stage everything depends on  $\varepsilon$ , which is however fixed for the moment. Let  $\Omega$  be a Zariski open subset of  $X$  such that  $\varphi_{\min}$  is locally bounded on  $\Omega$ ,  $\pi$  induces an isomorphism  $\Omega' := \pi^{-1}(\Omega) \rightarrow \Omega$  and every component of  $E$  is contained in the complement of  $\Omega'$ . Since the measures involved in equation (3.1) put no mass on  $X \setminus \Omega$ , it is enough to show the result on  $\Omega$ . Let thus  $f$  be a given smooth function with compact support in  $\Omega$ . We then have, by definition,

$$\begin{aligned} \int_X f \langle (\theta + dd^c \varphi)^n \rangle &= \lim_{k \rightarrow \infty} \int_{\{\varphi > \varphi_{\min} - k\} \cap \Omega} f(\theta + dd^c \max\{\varphi, \varphi_{\min} - k\})^n \\ &= \lim_{k \rightarrow \infty} \int_{\{\varphi' > \varphi'_{\min} - k\} \cap \Omega'} f'(\theta' + dd^c \max\{\varphi', \varphi'_{\min} - k\})^n, \end{aligned}$$

where primed objects have been pulled back by  $\pi$ . Note that  $\varphi'_{\min}$  is a  $\theta'$ -psh function with minimal singularities, by Proposition 1.12.

Now let  $\theta_E := \theta' - \omega$ , so that

$$[E] = \theta_E + dd^c \varphi_E$$

for some  $\theta_E$ -psh function  $\varphi_E$ . For cohomological reasons we have

$$\theta' + dd^c \varphi' = \pi^* \pi_*(\omega + dd^c \psi + [E]) = \omega + dd^c \psi + [E] = \theta' + dd^c(\psi + \varphi_E),$$

as the cohomology class of the right-hand current is  $\pi^* \theta$ . We thus see that  $\varphi' = \psi + \varphi_E$  up to an additive constant, which can be chosen to be 0, and we are reduced to showing that

$$I_k := \lim_{k \rightarrow \infty} \int_{\{\psi + \varphi_E > \varphi'_{\min} - k\} \cap \Omega'} f'(\omega + \theta_E + dd^c \max\{\psi + \varphi_E, \varphi'_{\min} - k\})^n$$

coincides with

$$\lim_{k \rightarrow \infty} \int_{\{\psi > -k\} \cap \Omega'} f'(\omega + dd^c \max\{\psi, -k\})^n.$$

Now  $\varphi_E$  is smooth on  $\Omega'$ , since  $E$  does not meet the latter set by assumption, and  $\varphi'_{\min} - \varphi_E \geq -C$  on  $X'$ , since the  $\theta'$ -psh function  $\varphi'_{\min}$  has minimal singularities, and hence is less singular than the  $\theta_E$ -psh function  $\varphi_E$ , with  $\theta_E \leq \theta'$ . We thus see that  $\varphi'_{\min} - \varphi_E$  is



bounded near the support of  $f'$ . On the other hand,  $\theta_E + dd^c \varphi_E = [E]$  vanishes on  $\Omega'$ , so that

$$(\omega + \theta_E + dd^c \max\{\psi + \varphi_E, \varphi'_{\min} - k\})^n = (\omega + dd^c \max\{\psi, \varphi'_{\min} - \varphi_E - k\})^n$$

there. Hence

$$I_k = \int_{\{\psi > v - k\} \cap \Omega'} f'(\omega + dd^c \max\{\psi, v - k\})^n,$$

with  $v := \varphi'_{\min} - \varphi_E$  bounded near the support of  $f'$ . We thus conclude that the latter integral equals

$$\lim_{k \rightarrow \infty} \int_{\{\psi > -k\} \cap \Omega'} f'(\omega + dd^c \max\{\psi, -k\})^n$$

as desired, which finishes the proof of (3.1).

At this point, we have thus obtained a sequence of  $\theta$ -psh functions  $\varphi_j$  such that  $\text{MA}(\varphi_j) = (1 - \varepsilon_j)\mu$  for some sequence  $\varepsilon_j > 0$  decreasing towards 0. If we normalize them in some way, say by  $\sup_X \varphi_j = 0$ , we can also assume by compactness that  $\varphi_j$  converges in  $L^1(X)$  to some  $\theta$ -psh function  $\varphi_\infty$ , as  $j \rightarrow \infty$ .

As explained above, proving that  $\varphi_\infty$  has full Monge–Ampère mass is enough to conclude that  $\text{MA}(\varphi_\infty) \geq \mu$  by Corollary 2.21, and hence  $\text{MA}(\varphi_\infty) = \mu$  since both measures then have the same mass.

As  $\int_X \text{MA}(\varphi_j) \rightarrow \text{vol}(\alpha)$ , we are reduced by Proposition 2.19 to finding a convex weight  $\chi$  such that

$$\sup_j \int_X (-\chi)(\varphi_j - \varphi_{\min}) \text{MA}(\varphi_j) \leq \sup_j \int_X (-\chi)(\varphi_j - \varphi_{\min}) d\mu < \infty,$$

which is taken care of by the next proposition. This concludes the proof of Theorem 3.1.

**PROPOSITION 3.2.** *Let  $\mu$  be a non-pluripolar measure and let  $\mathcal{K}$  be a given compact set of quasi-psh functions on  $X$ . Then there exists a convex weight function  $\chi$  and  $C > 0$  such that*

$$\int_X (-\chi)(\varphi) d\mu \leq C$$

for all  $\varphi \in \mathcal{K}$ .

*Proof.* Functions in  $\mathcal{K}$  are uniformly bounded from above by compactness, and we can thus assume that they are all non-positive. Now let  $\omega$  be a given Kähler form, and let  $\mathcal{M}$  be the (weakly) closed convex set of positive measures generated by all Monge–Ampère measures  $(\omega + dd^c \psi)^n$ , where  $0 \leq \psi \leq 1$  is  $\omega$ -psh. By the Chern–Levine–Nirenberg inequality, there exists a constant  $C > 0$  such that

$$\int_X (-\varphi)(\omega + dd^c \psi)^n \leq C$$

for all  $\varphi \in \mathcal{K}$  and all  $0 \leq \psi \leq 1$  as above, and this remains true on  $\mathcal{M}$ , by the convexity and the lower semi-continuity of  $\nu \mapsto \int_X (-\varphi) d\nu$  (since  $-\varphi$  is lower semi-continuous—just write it as the supremum of all continuous functions below it).

Now recall that pluripolar subsets  $A$  are characterized by the condition that  $\nu(A) = 0$  for all Monge–Ampère measures  $\nu$  as above. By an extension of Radon–Nikodym’s theorem proved in [34], there exist measures  $\nu \in \mathcal{M}$  and  $\nu' \perp \mathcal{M}$  such that our given measure  $\mu$  can be written as

$$\mu = g\nu + \nu',$$

where  $g$  is a non-negative element of  $L^1(\nu)$ . Since  $\mu(A) = 0$  for all pluripolar subsets by assumption, we infer that  $\nu' = 0$ , and we have thus proved that  $\mu \ll \nu$  for some  $\nu \in \mathcal{M}$ . We now conclude the proof by the following result, certainly a standard one in the theory of Orlicz spaces.  $\square$

LEMMA 3.3. *Let  $\mu \ll \nu$  be two positive measures with finite mass on a measured space  $X$ . Then there exists a concave function  $h: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and  $C > 0$  such that*

$$\int_X h(f) d\mu \leq C + \int_X f d\nu$$

for all measurable functions  $f \geq 0$ .

*Proof.* Let  $g \geq 0$  be the function in  $L^1(\nu)$  such that  $d\mu = g d\nu$ . By an easy measure-theoretic result that we have already used, there exists a convex increasing function  $\tau$  on  $\mathbb{R}^+$  such that  $\tau(x) \gg x$ , as  $x \rightarrow \infty$ , but still  $\int_X \tau(g) d\nu < \infty$ . Now let

$$\tau^*(y) = \sup_{x \geq 0} (xy - \tau(x))$$

be the Legendre transform of  $\tau$  and let  $h$  be the reciprocal function of  $\tau^*$ . By definition, we thus have the following generalized form of Young’s inequality

$$h(y)x \leq \tau(x) + y.$$

It follows that

$$\int_X h(f) d\mu = \int_X h(f)g d\nu \leq \int_X \tau(g) d\nu + \int_X f d\nu,$$

which yields the desired result with  $C := \int_X \tau(g) d\nu$ .  $\square$

Remark 3.4. When  $\mathcal{K}$  consists of a single function, Proposition 3.2 follows from standard measure-theoretic considerations since  $\mu(\{\varphi = -\infty\}) = 0$ . The general case would thus follow by compactness provided that  $\varphi \mapsto \int_X \varphi d\mu$  were known to be lower semi-continuous on  $\theta$ -psh functions in the  $L^1(X)$  topology (it is always *upper* semi-continuous by Fatou’s lemma). But already when  $n = 1$  the lower semi-continuity condition is not automatic: it holds if and only if  $\mu$  has *continuous* local potentials.

**3.3. Uniqueness**

In this section we show how to adapt Dinew’s tricky proof of uniqueness in [22] to our more general setting. As we shall see, his arguments carry over *mutatis mutandis*. On the other hand, we will make one of his arguments more precise, namely

$$\int_X \langle T_1 \wedge \dots \wedge T_n \rangle = \langle \alpha^n \rangle$$

for every set of positive currents  $T_1, \dots, T_n \in \alpha$  with full Monge–Ampère mass, which is the content of Corollary 2.15 above. This fact is non-obvious even when  $\alpha$  is Kähler, which is the case considered in [22], and seems to have been implicitly taken for granted there.

Let thus  $T_1 = \theta + dd^c \varphi_1$  and  $T_2 = \theta + dd^c \varphi_2$  be two positive currents in  $\alpha$  such that

$$\langle T_1^n \rangle = \langle T_2^n \rangle = \mu.$$

As in [21], we first remark that the log-concavity property of non-pluripolar products (Proposition 1.11) implies that

$$\langle T_1^k \wedge T_2^{n-k} \rangle \geq \mu,$$

and hence

$$\langle T_1^k \wedge T_2^{n-k} \rangle = \mu, \tag{3.2}$$

for  $k=0, \dots, n$ , since both measures have the same total mass  $v := \text{vol}(\alpha)$ .

*Step 1.* As in the first part of the proof of Theorem 1 in [22], we are first going to show by contradiction that there exists  $t \in \mathbb{R}$  such that  $\varphi_1 = \varphi_2 + t$   $\mu$ -almost everywhere.

Since  $\mu$  puts no mass on pluripolar subsets, the set of  $t \in \mathbb{R}$  such that

$$\mu(\{\varphi_1 = \varphi_2 + t\}) > 0$$

coincides with the discontinuity locus of the non-decreasing function

$$t \mapsto \mu(\{\varphi_1 < \varphi_2 + t\})$$

and is thus at most countable. We now assume by contradiction that  $\mu(\{\varphi_1 = \varphi_2 + t\}) < v$  for all  $t$ , so that we can find  $t \in \mathbb{R}$  such that

$$0 < \mu(\{\varphi_1 < \varphi_2 + t\}) < v.$$

By monotone convergence, upon slightly perturbing  $t$  we can furthermore arrange so that

$$\mu(\{\varphi_1 = \varphi_2 + t\}) = 0.$$

Replacing  $\varphi_2$  by  $\varphi_2+t$ , we can finally assume that  $t=0$ , so that both  $\mu(\{\varphi_1<\varphi_2\})$  and  $\mu(\{\varphi_2<\varphi_1\})$  are less than  $v$ , while  $\mu(\{\varphi_1=\varphi_2\})=0$ . We can thus pick  $\varepsilon>0$  small enough such that the mass of  $(1+\varepsilon)^n\mu$  on both  $\{\varphi_1<\varphi_2\}$  and  $\{\varphi_2<\varphi_1\}$  is still less than  $v$ .

By the existence part of Theorem 3.1 (which is already proved!), we get a positive current  $T=\theta+dd^c\varphi$  such that

$$\langle T^n \rangle = \mathbf{1}_{\{\varphi_1<\varphi_2\}}(1+\varepsilon)^n\mu + \mathbf{1}_{\{\varphi_1>\varphi_2\}}t\mu,$$

where  $t>0$  is taken so that the left-hand side has total mass  $v$ . As  $\varphi_{\min}$  has minimal singularities, we can assume that  $\varphi\leq\varphi_{\min}$ . Proposition 1.11 thus implies that

$$\langle T \wedge T_j^{n-1} \rangle \geq (1+\varepsilon)\mu$$

on  $\{\varphi_1<\varphi_2\}$ .

We now consider the subsets

$$O_\delta := \{(1-\delta)\varphi_1 + \delta\varphi_{\min} < (1-\delta)\varphi_2 + \delta\varphi\} \subset \{\varphi_1 < \varphi_2\}.$$

Since both  $T$  and  $T_j$  have full Monge–Ampère mass, Corollary 2.15 yields

$$\int_X \langle T_{\min} \wedge T_j^{n-1} \rangle = \langle \alpha^n \rangle \tag{3.3}$$

and the generalized comparison principle (Proposition 2.2) thus implies that

$$\int_{O_\delta} \langle ((1-\delta)T_2 + \delta T) \wedge T_j^{n-1} \rangle \leq \int_{O_\delta} \langle ((1-\delta)T_1 + \delta T_{\min}) \wedge T_j^{n-1} \rangle$$

for  $j=1, 2$ . By (3.2) it follows that

$$\delta(1+\varepsilon) \int_{O_\delta} \mu \leq \delta \int_{O_\delta} \langle T \wedge T_j^{n-1} \rangle \leq \delta \int_{\{\varphi_1<\varphi_2\}} \langle T_{\min} \wedge T_j^{n-1} \rangle.$$

Dividing by  $\delta>0$  and letting  $\delta$  tend to 0, we infer that

$$(1+\varepsilon) \int_{\{\varphi_1<\varphi_2\}} \mu \leq \int_{\{\varphi_1<\varphi_2\}} \langle T_{\min} \wedge T_j^{n-1} \rangle$$

for  $j=1, 2$ , by monotone convergence.

We can now study  $\{\varphi_2<\varphi_1\}$  (with a possibly different  $T$ ) similarly, to conclude that

$$(1+\varepsilon) \int_{\{\varphi_2<\varphi_1\}} \mu \leq \int_{\{\varphi_2<\varphi_1\}} \langle T_{\min} \wedge T_j^{n-1} \rangle,$$

and we finally reach a contradiction by summing these two inequalities, which yields

$$(1+\varepsilon)v \leq \int_X \langle T_{\min} \wedge T_j^{n-1} \rangle = \langle \alpha^n \rangle = v$$

by (3.3), using that  $\mu(\{\varphi_1 = \varphi_2\}) = 0$ .

*Step 2.* At this point we have proved that  $\varphi_1 = \varphi_2$  a.e. with respect to  $\text{MA}(\varphi_1) = \text{MA}(\varphi_2)$ . One might be tempted to use the domination principle to conclude that  $\varphi_2 \leq \varphi_1$ , and hence  $\varphi_1 = \varphi_2$  by symmetry. But Corollary 2.5 cannot be applied here since  $\varphi_2$  does not a priori have minimal singularities. The rest of the proof will circumvent this difficulty by using some additional cancellation.

By the existence part of the proof, we can choose a positive current  $T = \theta + dd^c \varphi$  such that  $\langle T^m \rangle$  is a smooth positive volume form. We normalize  $\varphi$  so that  $\varphi \leq \varphi_{\min}$ .

Let  $O := \{\varphi_1 < \varphi_2\}$ . We are going to show, by descending induction on  $m$ , that

$$\int_O \langle T_1^k \wedge T_2^l \wedge T^m \rangle = 0 \tag{3.4}$$

for all  $k, l$  and  $m$  such that  $k+l+m=n$ . The result holds for  $m=0$  by the first part of the proof, and the result for  $m=n$  will imply that  $\varphi_2 \leq \varphi_1$  a.e. with respect to Lebesgue measure, and will conclude the proof by symmetry.

Let thus  $m$  be given, and assume that (3.4) holds true for all  $k$  and  $l$  such that  $k+l+m=n$ . Pick  $k$  and  $l$  such that  $k+l+m+1=n$  and consider as usual the current with minimal singularities

$$T_1^{(j)} := \theta_1 + dd^c \varphi_1^{(j)},$$

where

$$\varphi_1^{(j)} = \max\{\varphi_1, \varphi_{\min} - j\}.$$

We introduce the plurifine open subsets

$$O_{j,\varepsilon} := \left\{ \left(1 - \frac{\varepsilon}{j}\right) \varphi_1 + \frac{\varepsilon}{j} \varphi_1^{(j)} + \varepsilon < \left(1 - \frac{\varepsilon}{j}\right) \varphi_2 + \frac{\varepsilon}{j} \varphi \right\}.$$

Note that  $O_{j,\varepsilon} \subset O$ , since

$$\frac{\varepsilon}{j} \varphi_1^{(j)} + \varepsilon \geq \frac{\varepsilon}{j} \varphi_{\min} \geq \frac{\varepsilon}{j} \varphi,$$

so that neither  $\langle T_k \wedge T_2^{l+1} \wedge T^m \rangle$  nor  $\langle T_1^{k+1} \wedge T_2^l \wedge T^m \rangle$  put any mass on  $O_{j,\varepsilon}$  by induction.

As  $T_1, T_2, T$  and  $T_1^{(j)}$  all have full Monge–Ampère mass, Corollary 2.15 yields

$$\int_X \langle T_1^{(j)} \wedge T_1^k \wedge T_2^l \wedge T^m \rangle = \langle \alpha^n \rangle.$$

The generalized comparison principle therefore implies that

$$\int_{O_{j,\varepsilon}} \left\langle \left( \left( 1 - \frac{\varepsilon}{j} \right) T_2 + \frac{\varepsilon}{j} T \right) \wedge T_1^k \wedge T_2^l \wedge T^m \right\rangle \leq \int_{O_{j,\varepsilon}} \left\langle \left( \left( 1 - \frac{\varepsilon}{j} \right) T_1 + \frac{\varepsilon}{j} T_1^{(j)} \right) \wedge T_1^k \wedge T_2^l \wedge T^m \right\rangle.$$

The leading terms on both sides vanish and we end up with

$$\int_{O_{j,\varepsilon}} \langle T_1^k \wedge T_2^l \wedge T^{m+1} \rangle \leq \int_O \langle T_1^{(j)} \wedge T_1^k \wedge T_2^l \wedge T^m \rangle.$$

By Corollary 2.15 again

$$\int_X \langle T_1^{k+1} \wedge T_2^l \wedge T^m \rangle = \langle \alpha^n \rangle,$$

thus

$$\lim_{j \rightarrow \infty} \int_O \langle T_1^{(j)} \wedge T_1^k \wedge T_2^l \wedge T^m \rangle = \int_O \langle T_1^{k+1} \wedge T_2^l \wedge T^m \rangle = 0.$$

On the other hand, we clearly have

$$\mathbf{1}_{\{\varphi_1 + \varepsilon < \varphi_2\}} \leq \liminf_{j \rightarrow \infty} \mathbf{1}_{O_{j,\varepsilon}}$$

pointwise outside a pluripolar set, and Fatou's lemma thus yields

$$\int_{\{\varphi_1 + \varepsilon < \varphi_2\}} \langle T_1^k \wedge T_2^l \wedge T^{m+1} \rangle = 0,$$

and hence also

$$\int_O \langle T_1^k \wedge T_2^l \wedge T^{m+1} \rangle = 0$$

by monotone convergence. The induction is thus complete, and the proof of uniqueness is finally over.

#### 4. $L^\infty$ a-priori estimate

We now consider the case of measures with density in  $L^{1+\varepsilon}$ ,  $\varepsilon > 0$ , with respect to Lebesgue measure. The goal of this section is to show how to adapt Kołodziej's pluripotential theoretic approach to the  $L^\infty$  a-priori estimates [30] to the present context of big cohomology classes. Fix an arbitrary (smooth positive) volume form  $dV$  on  $X$ .

Given a smooth representative  $\theta$  of the big cohomology class  $\alpha \in H^{1,1}(X, \mathbb{R})$ , we introduce the extremal function

$$V_\theta := \sup\{\psi \text{ } \theta\text{-psh} : \psi \leq 0\}. \quad (4.1)$$

Note that  $V_\theta$  has minimal singularities. Of course we have  $V_\theta = 0$  if (and only if)  $\theta$  is semi-positive.

**THEOREM 4.1.** *Let  $\mu=f dV$  be a positive measure with density  $f \in L^{1+\varepsilon}$ ,  $\varepsilon>0$ , such that  $\mu(X)=\text{vol}(\alpha)$ . Then the unique closed positive current  $T \in \alpha$ , such that  $\langle T^n \rangle = \mu$ , has minimal singularities.*

*More precisely, there exists a constant  $M$  only depending on  $\theta$ ,  $dV$  and  $\varepsilon$  such that the unique  $\theta$ -psh function  $\varphi$  with*

$$\text{MA}(\varphi) = \mu$$

*and normalized by*

$$\sup_X \varphi = 0$$

*satisfies*

$$\varphi \geq V_\theta - M \|f\|_{L^{1+\varepsilon}}^{1/n}. \tag{4.2}$$

The whole section is devoted to the proof of this result. As mentioned above, we follow Kołodziej’s approach [30]. After introducing the appropriate Monge–Ampère (pre)capacity  $\text{Cap}$ , we will estimate the capacity decay of sublevel sets  $\{\varphi < V_\theta - t\}$  as  $t \rightarrow \infty$  in order to show that actually  $\text{Cap}(\{\varphi < V_\theta - M\}) = 0$  for some  $M > 0$ , under control as above, which will prove the theorem.

**4.1. The Monge–Ampère capacity**

We define the *Monge–Ampère capacity*  $\text{Cap}$  as the upper envelope of the family of all Monge–Ampère measures  $\text{MA}(\psi)$  with  $\psi$   $\theta$ -psh such that  $V_\theta \leq \psi \leq V_\theta + 1$ , that is for each Borel subset  $B$  of  $X$  we set

$$\text{Cap}(B) := \sup \left\{ \int_B \text{MA}(\psi) : V_\theta \leq \psi \leq V_\theta + 1 \right\}. \tag{4.3}$$

By definition we have  $\text{Cap}(B) \leq \text{Cap}(X) = \text{vol}(\alpha)$ .

If  $K$  is a compact subset of  $X$ , we define its *extremal function* by

$$V_{K,\theta} := \sup \{ \psi \text{ } \theta\text{-psh} : \psi \leq 0 \text{ on } K \}.$$

Note that

$$V_\theta = V_{X,\theta} \leq V_{K,\theta} \tag{4.4}$$

by definition. Standard arguments show that the upper semi-continuous regularization  $V_{K,\theta}^*$  of  $V_{K,\theta}$  is either  $\theta$ -psh with minimal singularities when  $K$  is non-pluripolar, or  $V_{K,\theta}^* \equiv \infty$  when  $K$  is pluripolar. It is also a standard matter to show that  $\text{MA}(V_{K,\theta}^*)$  is concentrated on  $K$  using local solutions to the homogeneous Monge–Ampère equation on

small balls in the complement of  $K$  (cf. [26, Theorem 4.2]). It follows from the definition of  $V_{K,\theta}$  that the following maximum principle holds:

$$\sup_K \varphi = \sup_X (\varphi - V_{K,\theta}) \tag{4.5}$$

for any  $\theta$ -psh function  $\varphi$ .

Finally we introduce the *Alexander–Taylor capacity* of  $K$  in the following guise:

$$M_\theta(K) := \sup_X V_{K,\theta}^* \in [0, \infty], \tag{4.6}$$

so that  $M_\theta(K) = \infty$  if and only if  $K$  is pluripolar. We then have the following comparison theorem.

LEMMA 4.2. *For every non-pluripolar compact subset  $K$  of  $X$ , we have*

$$1 \leq \left( \frac{\text{vol}(\alpha)}{\text{Cap}(K)} \right)^{1/n} \leq \max\{1, M_\theta(K)\}.$$

*Proof.* The left-hand inequality is trivial. In order to prove the right-hand inequality we consider two cases. If  $M_\theta(K) \leq 1$ , then  $V_{K,\theta}^*$  is a candidate in the definition (4.3) of  $\text{Cap}(K)$  by (4.5). Since  $\text{MA}(V_{K,\theta}^*)$  is supported on  $K$ , we thus have

$$\text{Cap}(K) \geq \int_K \text{MA}(V_{K,\theta}^*) = \int_X \text{MA}(V_{K,\theta}^*) = \text{vol}(\alpha)$$

and the desired inequality holds in that case.

On the other hand, if  $M := M_\theta(K) \geq 1$  we have

$$V_\theta \leq \frac{1}{M} V_{K,\theta}^* + \left(1 - \frac{1}{M}\right) V_\theta \leq V_\theta + 1$$

by (4.4), and it follows by definition of the capacity again that

$$\text{Cap}(K) \geq \int_K \text{MA} \left( \frac{1}{M} V_{K,\theta}^* + \left(1 - \frac{1}{M}\right) V_\theta \right).$$

But since

$$\text{MA} \left( \frac{1}{M} V_{K,\theta}^* + \left(1 - \frac{1}{M}\right) V_\theta \right) \geq \frac{1}{M^n} \text{MA}(V_{K,\theta}^*),$$

we deduce that

$$\int_K \text{MA} \left( \frac{1}{M} V_{K,\theta}^* + \left(1 - \frac{1}{M}\right) V_\theta \right) \geq \frac{1}{M^n} \int_X \text{MA}(V_{K,\theta}^*) = \frac{1}{M^n} \text{vol}(\alpha),$$

and the result follows. □



Using this fact, we will now prove, as in [30], that any measure with  $L^{1+\varepsilon}$ -density is nicely dominated by the capacity.

PROPOSITION 4.3. *Let  $\mu = f dV$  be a positive measure with  $L^{1+\varepsilon}$ -density with respect to Lebesgue measure, where  $\varepsilon > 0$ . Then there exists a constant  $C$  only depending on  $\theta$  and  $\mu$  such that*

$$\mu(B) \leq C \text{Cap}(B)^2$$

for all Borel subsets  $B$  of  $X$ . In fact we can take

$$C := \varepsilon^{-2n} A \|f\|_{L^{1+\varepsilon}(dV)}$$

for a constant  $A$  only depending on  $\theta$  and  $dV$ .

*Proof.* By the inner regularity of  $\mu$ , it is enough to consider the case where  $B = K$  is compact. We can also assume that  $K$  is non-pluripolar, since  $\mu(K) = 0$  otherwise and the inequality is then trivial.

Now set

$$\nu_\theta := 2 \sup_{T,x} \nu(T, x), \tag{4.7}$$

the supremum ranging over all positive currents  $T \in \alpha$  and all  $x \in X$ , and  $\nu(T, x)$  denoting the Lelong number of  $T$  at  $x$ . As all Lelong numbers of  $\nu_\theta^{-1} T$  are  $\leq \frac{1}{2} < 1$  for each positive current  $T \in \alpha$ , the uniform version of Skoda's integrability theorem together with the compactness of normalized  $\theta$ -psh functions yields a constant  $C_\theta > 0$  only depending on  $dV$  and  $\theta$  such that

$$\int_X \exp\left(-\frac{1}{\nu_\theta} \psi\right) dV \leq C_\theta$$

for all  $\theta$ -psh functions  $\psi$  normalized by  $\sup_X \psi = 0$  (see [45]). Applying this to

$$\psi = V_{K,\theta}^* - M_\theta(K)$$

(which has the right normalization by (4.6)), we get

$$\int_X \exp\left(-\frac{1}{\nu_\theta} V_{K,\theta}^*\right) dV \leq C_\theta \exp\left(-\frac{1}{\nu_\theta} M_\theta(K)\right).$$

On the other hand, we have  $V_{K,\theta}^* \leq 0$  on  $K$  a.e. with respect to Lebesgue measure. Hence

$$\text{vol}(K) \leq C_\theta \exp\left(-\frac{1}{\nu_\theta} M_\theta(K)\right). \tag{4.8}$$

Hölder's inequality yields

$$\mu(K) \leq \|f\|_{L^{1+\varepsilon}(dV)} \text{vol}(K)^{\varepsilon/(1+\varepsilon)}. \tag{4.9}$$

We may also assume that  $M_\theta(K) \geq 1$ . Otherwise Lemma 4.2 implies  $\text{Cap}(K) = \text{vol}(\alpha)$ , and the result is thus clear in this case. By Lemma 4.2, (4.8) and (4.9) together, we thus get

$$\mu(K) \leq C_\theta^{\varepsilon/(1+\varepsilon)} \|f\|_{L^{1+\varepsilon}(dV)} \exp\left(-\frac{\varepsilon}{(1+\varepsilon)\nu_\theta} \left(\frac{\text{Cap}(K)}{\text{vol}(\alpha)}\right)^{-1/n}\right),$$

and the result follows since  $\exp(-t^{-1/n}) = O(t^2)$  when  $t \rightarrow 0_+$ . □

**4.2. Proof of Theorem 4.1**

We first apply the comparison principle to get the following result.

LEMMA 4.4. *Let  $\varphi$  be a  $\theta$ -psh function with full Monge–Ampère mass. Then for all  $t > 0$  and  $0 < \delta < 1$  we have*

$$\text{Cap}(\{\varphi < V_\theta - t - \delta\}) \leq \frac{1}{\delta^n} \int_{\{\varphi < V_\theta - t\}} \text{MA}(\varphi).$$

*Proof.* Let  $\psi$  be a  $\theta$ -psh function such that  $V_\theta \leq \psi \leq V_\theta + 1$ . We then have

$$\{\varphi < V_\theta - t - \delta\} \subset \{\varphi < \delta\psi + (1-\delta)V_\theta - t - \delta\} \subset \{\varphi < V_\theta - t\}.$$

Since  $\delta^n \text{MA}(\psi) \leq \text{MA}(\delta\psi + (1-\delta)V_\theta)$  and  $\varphi$  has finite Monge–Ampère energy, Proposition 2.2 yields

$$\begin{aligned} \delta^n \int_{\{\varphi < V_\theta - t - \delta\}} \text{MA}(\psi) &\leq \int_{\{\varphi < \delta\psi + (1-\delta)V_\theta - t - \delta\}} \text{MA}(\delta\psi + (1-\delta)V_\theta) \\ &\leq \int_{\{\varphi < \delta\psi + (1-\delta)V_\theta - t - \delta\}} \text{MA}(\varphi) \leq \int_{\{\varphi < V_\theta - t\}} \text{MA}(\varphi), \end{aligned}$$

and the proof is complete. □

Now set

$$g(t) := \text{Cap}(\{\varphi < V_\theta - t\})^{1/n}.$$

Our goal is to show that  $g(M) = 0$  for some  $M$  under control. Indeed we will then have  $\varphi \geq V_\theta - M$  on  $X \setminus P$  for some Borel subset  $P$  such that  $\text{Cap}(P) = 0$ . But it then follows in particular from Proposition 4.3 (applied to the Lebesgue measure itself) that  $P$  has Lebesgue measure zero, and hence  $\varphi \geq V_\theta - M$  will hold everywhere.

Now, since  $\text{MA}(\varphi) = \mu$ , it follows from Proposition 4.3 and Lemma 4.4 that

$$g(t+\delta) \leq \frac{C^{1/n}}{\delta} g(t)^2 \quad \text{for all } t > 0 \text{ and } 0 < \delta < 1.$$

We can thus apply [25, Lemma 2.3] which yields  $g(M)=0$  for  $M:=t_0+4C^{1/n}$  (see also [5]). Here  $t_0>0$  has to be chosen so that

$$g(t_0) < \frac{1}{2C^{1/n}}.$$

Lemma 4.4 (with  $\delta=1$ ) implies that

$$\begin{aligned} g(t)^n &\leq \mu(\{\varphi < V_\theta - t + 1\}) \leq \frac{1}{t-1} \int_X |V_\theta - \varphi| f \, dV \\ &\leq \frac{1}{t-1} \|f\|_{L^{1+\varepsilon}(dV)} (\|\varphi\|_{L^{1+1/\varepsilon}(dV)} + \|V_\theta\|_{L^{1+1/\varepsilon}(dV)}), \end{aligned}$$

by Hölder’s inequality. Since  $\varphi$  and  $V_\theta$  both belong to the compact set of  $\theta$ -psh functions normalized by  $\sup_X \varphi=0$ , their  $L^{1+1/\varepsilon}(dV)$ -norms are bounded by a constant  $C_2$  only depending on  $\theta, dV$  and  $\varepsilon$ . It is thus enough to take

$$t_0 > 1 + 2^{n-1} C_2 C \|f\|_{L^{1+\varepsilon}(dV)}.$$

*Remark 4.5.* We have already mentioned that  $\text{vol}(\alpha)$  is continuous on the big cone. On the other hand, it is easy to see that  $\nu_\theta$  remains bounded as long as  $\theta$  is  $C^2$ -bounded. As a consequence, if  $\mu=f \, dV$  is fixed, we see that the constant  $C$  in Proposition 4.3 can be taken uniform in a small  $C^2$ -neighborhood of a given form  $\theta_0$ . We similarly conclude that  $M$  in Theorem 4.1 is uniform around a fixed form  $\theta_0, \mu$  being fixed.

*Remark 4.6.* As in [30], Theorem 4.1 holds more generally when the density  $f$  belongs to some Orlicz class. Indeed it suffices to replace (4.9) by the appropriate Orlicz-type inequality in the proof of Proposition 4.3.

### 5. Regularity of solutions: the case of nef classes

Let  $\alpha$  be a big cohomology class and let  $\mu$  be a smooth positive volume form of total mass equal to  $\text{vol}(\alpha)$ . Let  $T \in \alpha$  be the unique positive current such that  $\langle T^n \rangle = \mu$ . By Theorem 4.1,  $T$  has minimal singularities, which implies in particular that  $T$  has locally bounded potentials on the Zariski open subset  $\text{Amp}(\alpha)$ , the ample locus of  $\alpha$ . Note that the equation  $\langle T^n \rangle = \mu$  then simply means that  $T^n$ , which is well defined on  $\text{Amp}(\alpha)$  by Bedford–Taylor’s theorem, satisfies  $T^n = \mu$  on this open subset.

If  $\alpha$  is a Kähler class, then  $\text{Amp}(\alpha)=X$  and Yau’s theorem [44] implies that  $T$  is smooth. For an arbitrary big class  $\alpha$  the expectation is that  $T$  is smooth on  $\text{Amp}(\alpha)$ . By Evans–Trudinger’s general regularity theory for fully non-linear elliptic equations [24], [40] it is enough to show that  $T$  has  $L^\infty_{\text{loc}}$  coefficients on  $\text{Amp}(\alpha)$ , or equivalently that the trace measure of  $T$  has  $L^\infty_{\text{loc}}$ -density with respect to Lebesgue measure (cf. the discussion in [8]). We are unfortunately unable to prove this in general, but we can handle the following special case. The arguments are similar to those used by Sugiyama in [39].

THEOREM 5.1. *Let  $\alpha$  be a nef and big class. Let  $\mu$  be a smooth positive volume of total mass equal to  $\text{vol}(\alpha)=\alpha^n$ . Then the positive current  $T\in\alpha$  such that  $\langle T^n\rangle=\mu$  is smooth on  $\text{Amp}(\alpha)$ .*

*Proof.* Write  $T=\theta+dd^c\varphi$  with  $\sup_X\varphi=0$ . The  $\theta$ -psh function  $\varphi$  is locally bounded on  $\Omega:=\text{Amp}(\alpha)$  and satisfies

$$(\theta+dd^c\varphi)^n=\mu$$

there. If  $\pi:X'\rightarrow X$  is a modification that is isomorphic over  $\Omega$ , then  $\varphi':=\pi^*\varphi$ ,  $\theta':=\pi^*\theta$  and  $\mu':=\pi^*\mu$  obviously satisfy

$$(\theta'+dd^c\varphi')^n=\mu'$$

on  $\Omega':=\pi^{-1}(\Omega)$ . Note also that  $\varphi'$  is a  $\theta'$ -psh function with minimal singularities by Proposition 1.12. Now  $\text{Amp}(\alpha)$  is covered by Zariski open subsets of the form  $X\setminus\pi(E)$ , where  $\pi$  is as above,  $E$  is an effective  $\mathbb{R}$ -divisor on  $X'$  and  $\theta'$  is cohomologous to  $\omega+[E]$  for some Kähler form  $\omega$  on  $X'$ . We thus fix such a data, and our goal is to show that  $\Delta\varphi'$  belongs to  $L_{\text{loc}}^\infty$  on  $X'\setminus E$  (the Laplacian being computed with respect to  $\omega$ ). We can find a quasi-psh function  $\varphi_E$  on  $X'$  such that

$$[E]=\theta'-\omega+dd^c\varphi_E,$$

so that  $\varphi_E$  is smooth on  $X'\setminus E$  and satisfies  $\theta'=\omega-dd^c\varphi_E$  there. Since  $\varphi_E$  is in particular  $\theta$ -psh and  $\varphi'$  has minimal singularities, we have  $\varphi_E\leq\varphi'+O(1)$ . Upon replacing  $E$  by  $(1+\varepsilon)E$ , with  $0<\varepsilon\ll 1$ , in the above construction and shifting  $\varphi_E$  by a constant, we can in fact assume that

$$\varphi'\geq(1-\varepsilon)\varphi_E \tag{5.1}$$

for some  $\varepsilon>0$ .

On the other hand, let us write

$$\mu'=e^F\omega^n.$$

The function  $e^F$  is smooth but vanishes on the critical locus of  $\pi$ . Indeed  $F$  is quasi-psh (in particular bounded from above) and  $dd^cF$  is equal to the integration current on the relative canonical divisor of  $\pi$  modulo a smooth form. In particular the Laplacian  $\Delta F$  (again with respect to  $\omega$ ) is *globally* bounded on  $X'\setminus E$ .

For each  $t\geq 0$  we consider the smooth  $(1,1)$ -form

$$\theta_t:=\theta'+t\omega.$$

Note that a  $\theta_t$ -psh function a fortiori is  $\theta_s$ -psh for  $s\geq t$ , as  $\omega\geq 0$ . We set  $V_t:=V_{\theta_t}$ , where the latter is the extremal  $\theta_t$ -psh function defined by (4.1). The proof of the following lemma is straightforward.

LEMMA 5.2. *The quasi-psh functions  $V_t$  decrease pointwise to  $V_0$ , as  $t \rightarrow 0$ .*

Since  $V_0$  is a  $\theta_0$ -psh function with minimal singularities, inequality (5.1) together with Lemma 5.2 imply that  $V_t - \varphi_E$  tends to  $\infty$  near  $E$  for each  $t \geq 0$ .

By Theorem 4.1, for each  $t \geq 0$  there exists a unique  $\theta_t$ -psh function  $\varphi_t$  with minimal singularities such that

$$(\theta_t + dd^c \varphi_t)^n = e^F \omega^n \tag{5.2}$$

which is normalized by  $\sup_X \varphi_t = 0$ . By Theorem 4.1 and Remark 4.5, there exists  $M > 0$  such that

$$\varphi_t \geq V_t - M \tag{5.3}$$

for all  $t \geq 0$ . We thus have  $\varphi_0 = \varphi'$ . The class of  $\theta_t$  is Kähler for  $t > 0$  since the class of  $\theta$  is nef. As  $F$  has analytic singularities, Theorem 3 in [44, p. 374] implies that  $\varphi_t$  is smooth on  $X' \setminus E$  and  $\Delta \varphi_t$  is globally bounded on  $X'$  for each  $t > 0$  (but of course no uniformity is claimed with respect to  $t$ ).

LEMMA 5.3. *The normalized solutions  $\varphi_t$  satisfy*

$$\lim_{t \rightarrow 0} \varphi_t = \varphi_0.$$

*Proof.* The normalized quasi-psh functions  $\varphi_t$  live in a compact subset of  $L^1(X)$ . By the uniqueness part of Theorem 3.1, it is thus enough to show that any limit  $\psi$  of a sequence  $\varphi_k := \varphi_{t_k}$ , with  $t_k \rightarrow 0$ , satisfies  $\langle (\theta_0 + dd^c \psi)^n \rangle = e^F \omega^n$  and  $\sup_X \psi = 0$ . The latter property follows from Hartogs' lemma. To prove the former, we introduce

$$\psi_k := \left( \sup_{j \geq k} \varphi_j \right)^*$$

so that  $\psi_k$  is  $\theta_k$ -psh and decreases pointwise to  $\psi$ . As in Proposition 2.20, we get

$$(\theta_k + dd^c \psi_k)^n \geq e^F \omega^n$$

on  $X' \setminus E$  and the result follows by continuity of the Monge–Ampère operator along decreasing sequences of bounded psh functions.  $\square$

We are now going to prove that  $\Delta \varphi_t$  is uniformly bounded on compact subsets of  $X' \setminus E$ , which will imply that  $\Delta \varphi_0$  is  $L^\infty_{\text{loc}}$  on  $X' \setminus E$  by Lemma 5.3. This will be accomplished by using Yau's pointwise computations [44, p. 350]. In order to do so, we rely on Tsuji's trick [41]: we introduce

$$u_t := \varphi_t - \varphi_E,$$

which is smooth on  $X' \setminus E$  and satisfies

$$(\omega + dd^c u_t)^n = \mu$$

there, since  $\theta' = \omega - dd^c \varphi_E$  on  $X' \setminus E$ . Note that  $u_t$  is *not* quasi-psh on  $X'$ . Indeed, we have  $u_t \rightarrow \infty$  near  $E$ , as  $\varphi_t = V_t + O(1)$  on  $X'$ . Since  $\Delta \varphi_E$  is globally bounded on  $X' \setminus E$ , as was already noted, to bound  $\Delta \varphi_t$  is equivalent to bounding  $\Delta u_t$ .

We now basically follow the argument in [44, p. 350]. By inequalities (2.18) and (2.20) in [44, p. 350] (which only depend on pointwise computations), we have

$$\begin{aligned} e^{Au_t} \Delta_t (e^{-Au_t} (n + \Delta u_t)) \\ \geq -An(n + \Delta u_t) + (A + b_\omega) e^{-F/(n-1)} (n + \Delta \psi)^{n/(n-1)} + \Delta F - n^2 b_\omega \end{aligned}$$

for every  $A > 0$  such that  $A + b_\omega > 1$ . Here  $\Delta$  and  $\Delta_t$  denote the (negative) Laplacians associated with the Kähler forms  $\omega$  and  $\omega + dd^c u_t$ , respectively, and  $b_\omega$  denotes the (pointwise) minimal holomorphic bisectional curvature of  $\omega$ . Now  $b_\omega$  is globally bounded,  $F$  is bounded from above and  $\Delta F$  is globally bounded on  $X' \setminus E$ . We can thus find  $B, C, D > 0$  (independent of  $t$ ) such that

$$e^{Au_t} \Delta_t (e^{-Au_t} (n + \Delta u_t)) \geq -B(n + \Delta u_t) + C(n + \Delta u_t)^{n/(n-1)} - D \tag{5.4}$$

holds on  $X' \setminus E$ . As  $\Delta u_t = \Delta \varphi_t - \Delta \varphi_E$  is smooth and globally bounded on  $X' \setminus E$ , and since  $u_t$  tends to  $\infty$  near  $E$ , we see that

$$e^{-Cu_t} (n + \Delta u_t)$$

achieves its maximum on  $X' \setminus E$  at some point  $x_t \in X' \setminus E$ . Inequality (5.4) applied at  $x_t$  thus yields

$$C(n + \Delta u_t)^{n/(n-1)} \leq B(n + \Delta u_t) + D$$

at  $x_t$ . As  $n/(n-1) > 1$ , we thus see that there exists  $C_1$  (independent of  $t$ ) such that

$$(n + \Delta u_t)(x_t) \leq C_1.$$

We thus get

$$n + \Delta u_t \leq C_1 \exp\left(C\left(u_t - \inf_{X' \setminus E} u_t\right)\right) \tag{5.5}$$

by the definition of  $x_t$  (cf. (2.24) in [44, p. 351]). By (5.1) and (5.3), we have

$$-M \leq V_t - \varphi_E - M \leq \varphi_t - \varphi_E = u_t \leq \left(\sup_X V_1\right) - \varphi_E$$

for  $0 \leq t \leq 1$ , which shows that  $u_t - \inf_{X' \setminus E} u_t$  is uniformly bounded from above on compact subsets of  $X' \setminus E$ , and the proof is thus complete in view of (5.5). □

It is a consequence of Demailly’s regularization theorem that currents with minimal singularities  $T \in \alpha$  in a nef and big class have identically zero Lelong numbers (cf. [11]). Here is an example where such currents however have poles.

*Example 5.4.* Start with a famous example due to Serre: let  $E$  be the (flat, but not unitary flat) rank-2 vector bundle over the elliptic curve  $C := \mathbb{C}/\mathbb{Z}[i]$  associated with the representation  $\pi_1(C) = \mathbb{Z}[i] \rightarrow \mathrm{SL}(2, \mathbb{C})$  sending 1 to the identity and  $i$  to

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

(Here, but not elsewhere in this paper,  $i$  denotes the imaginary unit.)

The ruled surface  $S := \mathbb{P}(E) \rightarrow C$  of hyperplanes of  $E$  has a section  $C'$  with flat normal bundle, which lies in the linear system  $|\mathcal{O}_E(1)|$ . The original point of this construction of Serre was that  $X \setminus C'$  is Stein but not affine, and the reason for that is that  $C'$  is rigid in  $X$  despite having a non-negative normal bundle. In fact, Demailly–Peternell–Schneider [20] have proved that  $C'$  is rigid in the very strong sense that the only closed positive current cohomologous to  $C'$  is  $C'$  itself.

Now let  $X := \mathbb{P}(V)$  be the projective bundle of hyperplanes in  $V := E \oplus A$ , where  $A$  is a given ample line bundle on  $C$ , and let  $L := \mathcal{O}_V(1)$  be the tautological bundle. The line bundle  $L$  is nef since  $E$  and  $A$  are nef, and it is also big since  $A$  is big. It is easy to show that the non-ample locus of  $L$  is exactly  $S = \mathbb{P}(E) \subset \mathbb{P}(V) = X$ . But the restriction of  $L$  to  $S$  is  $\mathcal{O}_E(1)$ , and positive currents can be restricted to any subvariety not entirely contained in their polar set. It follows that any positive current in the nef and big class  $\alpha := c_1(L)$  has poles along  $C'$ .

## 6. Singular Kähler–Einstein metrics

### 6.1. More Monge–Ampère equations

In this section we show how to use the a-priori estimate of Theorem 4.1 and Schauder’s fixed point theorem to solve Monge–Ampère equations with non-constant right-hand side.

We let as before  $dV$  be a smooth volume form on  $X$  and  $\theta$  be a smooth representative of a big cohomology class.

**THEOREM 6.1.** *There exists a unique  $\theta$ -psh function  $\varphi$  with full Monge–Ampère mass such that*

$$\mathrm{MA}(\varphi) = e^\varphi dV.$$

*Furthermore  $\varphi$  has minimal singularities.*

Since  $\varphi$  is bounded from above, the measure  $e^\varphi \mu$  has  $L^\infty$ -density with respect to Lebesgue measure. Hence Theorem 4.1 implies that any  $\theta$ -psh solution  $\varphi$  with full Monge–Ampère mass has minimal singularities.

We now proceed with the existence proof, and uniqueness will be taken care of by Proposition 6.3 below.

*Proof.* We may assume that  $dV$  has total mass 1. Let  $\mathcal{C}$  be the compact convex subset of  $L^1(X)$  consisting of all  $\theta$ -psh functions  $\psi$  normalized by  $\sup_X \psi = 0$ . By compactness, there exists  $C > 0$  such that

$$\int_X \psi dV \geq -C$$

for all  $\psi \in \mathcal{C}$  (cf. [26, Proposition 1.7]). On the other hand, since  $dV$  is a probability measure, we get by convexity that

$$\log \int_X e^\psi dV \geq -C \tag{6.1}$$

for all  $\psi \in \mathcal{C}$ .

As noticed above, since  $e^\psi dV$  has  $L^\infty$ -density with respect to Lebesgue measure, it follows from Theorem 3.1 that for each  $\psi \in \mathcal{C}$  there exists a unique  $\Phi(\psi) \in \mathcal{C}$  such that

$$\text{MA}(\Phi(\psi)) = e^{\psi + c_\psi} dV,$$

where

$$c_\psi := \log \frac{\text{vol}(\alpha)}{\int_X e^\psi dV}$$

is a normalizing constant making the total masses fit. Note that

$$c_\psi \leq \log \text{vol}(\alpha) + C$$

for all  $\psi \in \mathcal{C}$  by (6.1). Theorem 4.1 therefore gives us a constant  $M > 0$  independent of  $\psi \in \mathcal{C}$  such that

$$\Phi(\psi) \geq V_\theta - M.$$

LEMMA 6.2. *The mapping  $\Phi: \mathcal{C} \rightarrow \mathcal{C}$  is continuous.*

*Proof.* Let  $\psi_j \rightarrow \psi$  be a convergent sequence in  $\mathcal{C}$ . Upon extracting, we may assume by compactness that

$$\varphi_j := \Phi(\psi_j) \rightarrow \varphi$$

for some  $\varphi \in \mathcal{C}$ , and also that  $\psi_j \rightarrow \psi$  almost everywhere. By the definition of  $c_\psi$ , this implies  $c_{\psi_j} \rightarrow c_\psi$  by dominated convergence. We claim that  $\varphi$  has full Monge–Ampère mass and satisfies  $\text{MA}(\varphi) = e^{\psi + c_\psi} dV$ . By uniqueness, we will then get  $\varphi = \Phi(\psi)$  as desired.

Indeed, as  $\Phi(\psi_j) \geq V_\theta - M$  for all  $j$ , it follows that also  $\varphi \geq V_\theta - M$ , and in particular  $\varphi$  has full Monge–Ampère mass. Corollary 2.21 thus yields  $\text{MA}(\varphi) \geq e^{\psi + c_\psi} \mu$ , and equality follows as desired since the total masses are both equal to  $\text{vol}(\alpha)$ .  $\square$



By Schauder’s fixed point theorem,  $\Phi$  has a fixed point  $\psi \in \mathcal{C}$ , and we then get a solution by setting  $\varphi := \psi + c_\psi$ . □

PROPOSITION 6.3. *If  $\varphi_1$  and  $\varphi_2$  are two  $\theta$ -psh functions with small unbounded locus such that*

$$\text{MA}(\varphi_i) = e^{\varphi_i} dV, \quad i = 1, 2,$$

*then  $\varphi_1 \leq \varphi_2 + O(1)$  already implies that  $\varphi_1 \leq \varphi_2$ .*

*Proof.* By the refined comparison principle for functions with small unbounded locus explained in Remark 2.4, we get

$$\int_{\{\varphi_2 < \varphi_1 - \varepsilon\}} \text{MA}(\varphi_1) \leq \int_{\{\varphi_2 < \varphi_1 - \varepsilon\}} \text{MA}(\varphi_2),$$

that is

$$\int_{\{\varphi_2 < \varphi_1 - \varepsilon\}} e^{\varphi_1} dV \leq \int_{\{\varphi_2 < \varphi_1 - \varepsilon\}} e^{\varphi_2} dV.$$

This implies that  $\varphi_2 \geq \varphi_1 - \varepsilon$  almost everywhere, and hence everywhere, and the result follows. □

As an immediate corollary to Theorem 6.1, we get the following result.

COROLLARY 6.4. *Let  $X$  be a projective manifold of general type, that is with a big canonical bundle  $K_X$ . Then there exists a unique singular non-negatively curved metric  $e^{-\phi_{\text{KE}}}$  on  $K_X$  satisfying the Kähler–Einstein equation*

$$\langle (dd^c \phi_{\text{KE}})^n \rangle = e^{\phi_{\text{KE}}}$$

*and such that*

$$\int_X e^{\phi_{\text{KE}}} = \text{vol}(X).$$

*Furthermore  $\phi_{\text{KE}}$  has minimal singularities.*

### 6.2. Comparison with previous results

We compare Corollary 6.4 with previous results and discuss the difficulties raised by the regularity issue on  $\text{Amp}(K_X)$ .

*Ample case.* When  $K_X$  is ample we have  $\text{Amp}(K_X) = X$ , and Aubin–Yau’s theorem [1], [44] can of course be reformulated by saying that  $\phi_{\text{KE}}$  constructed in Corollary 6.4 is smooth.

*Nef and big case.* Assume that  $K_X$  is nef and big. It is then semi-ample by the base-point-free theorem, so that  $\text{Amp}(K_X)$  coincides with the degeneracy locus of the birational morphism

$$X \longrightarrow X_{\text{can}} := \text{Proj } R(K_X)$$

and minimal singularities means locally bounded. Theorem 1 of [41] constructs a “canonical” psh weight  $\phi$  on  $K_X$  which is smooth outside  $\text{Amp}(K_X)$  and satisfies  $(dd^c\phi)^n = e^\phi$ . Theorem 3.8 of [39] yields the additional information that  $\phi$  has globally bounded Laplacian on  $X$  (and hence in  $C^{2-\varepsilon}(X)$  for each  $\varepsilon > 0$ ).

*Big case.* If we are willing to use finite generation of the canonical ring  $R(K_X)$  [7] (see also [42] and [37] for more analytic approaches) let  $\mu: Y \rightarrow X$  and  $\nu: Y \rightarrow X_{\text{can}}$  be a resolution of the graph of  $X \dashrightarrow X_{\text{can}}$ . By the negativity lemma ([29, Lemma 3.39]), we have

$$\mu^* K_X = \nu^* K_{X_{\text{can}}} + E,$$

where  $E$  is an effective  $\nu$ -exceptional  $\mathbb{Q}$ -divisor. This decomposition is thus the Zariski decomposition of  $K_X$ , and it follows that every positive current  $T$  in  $c_1(K_X)$  satisfies

$$\mu^* T = \nu^* S + [E]$$

for a unique positive current  $S$  in  $c_1(K_{X_{\text{can}}})$  (cf. for instance [11]). It is then immediate to see that the Kähler–Einstein metric  $e^{-\phi_{\text{KE}}}$  on  $K_X$  constructed in Corollary 6.4 corresponds in this way to the the Kähler–Einstein metric on  $K_{X_{\text{can}}}$  constructed in [25, Theorem 7.8].

Suppose now that we do not use the existence of the canonical model  $X_{\text{can}}$ . Theorem 5.1 of [43] claims the existence of a psh weight  $\phi$  on  $K_X$  such that  $\phi$  is smooth on a Zariski open subset  $U$  of  $X$ ,  $(dd^c\phi)^n = e^\phi$  holds on  $U$  and  $\phi$  is an *analytic Zariski decomposition*. This property means that the length with respect to  $\phi$  of every pluricanonical section is in  $L^2$  on  $X$  (and hence is implied by the stronger condition that such length functions are bounded).

Tsuji’s argument essentially proceeds as follows. As in the proof of Theorem 3.1 above we consider approximate Zariski decompositions, that is we let  $\pi_k: X_k \rightarrow X$  be an increasing sequence of modifications such that

$$\pi_k^* K = A_k + E_k$$

with  $A_k$  ample,  $E_k$  effective and  $E_{k+1} < E_k$  (on  $X_{k+1}$ ), and such that

$$\lim_{k \rightarrow \infty} \text{vol}(A_k) = \text{vol}(K_X).$$

We may assume that  $\pi_0$  is the identity.

For each  $k$ , we can use [44] as in the beginning of the proof of Theorem 3.1 to get a psh weight  $\phi_k$  on  $K_X$  such that

$$\pi_k^* dd^c \phi_k = \omega_k + [E_k]$$

for some Kähler form  $\omega_k$  in the class of  $A_k$  and  $(dd^c \phi_k)^n = e^{\phi_k}$  holds on  $X \setminus E_1$ .

Tsuji shows, using the comparison principle (cf. Proposition 6.3), that  $\phi_k$  is non-decreasing and locally uniformly bounded from above. The weight  $\phi := (\sup_k \phi_k)^*$  is thus psh and satisfies

$$\phi_0 \leq \phi_k \leq \phi$$

for all  $k$ . It follows by construction that the length with respect to  $\phi$  of every pluricanonical section is  $L^\infty$  on  $X$ , so that  $\phi$  is in particular an analytic Zariski decomposition. These arguments of Tsuji have also been expanded in [38, §4.3].

Note that by monotone convergence we have

$$\int_X e^\phi = \lim_{k \rightarrow \infty} \int_X e^{\phi_k}.$$

Now  $\langle (dd^c \phi_k)^n \rangle = e^{\phi_k}$  and it is easy to see that

$$\int_X \langle (dd^c \phi_k)^n \rangle = \text{vol}(A_k).$$

It thus follows that  $\int_X e^\phi = \text{vol}(K_X)$ , so that the psh weight  $\phi$  constructed in [43] and [38] coincides with  $\phi_{\text{KE}}$  of Corollary 6.4. As was already noted, it follows by construction that the length with respect to  $\phi$  of any section  $\sigma \in H^0(kL)$  is bounded, which implies in particular that  $\phi$  is an analytic Zariski decomposition in the sense of Tsuji. On the other hand, Corollary 6.4 says that  $\phi_{\text{KE}}$  has *minimal singularities*. As we explain below, the latter condition is strictly stronger than the former for a general big line bundle  $L$  (but a posteriori not for  $K_X$  due to finite generation of the canonical ring).

We now discuss Tsuji's approach to the smoothness of  $\phi$  on  $X \setminus E_0$ , which unfortunately seems to encounter a difficulty that we were not able to fix. As in the proof of Theorem 5.1 above, set  $u_k := \phi_k - \phi_0$ , which is smooth on  $X \setminus E_0$  and satisfies

$$(\omega_0 + dd^c u_k)^n = e^{u_k + F} \omega_0^n$$

on  $X \setminus E_0$ , with  $F$  such that  $e^F \omega_0^n = e^{\phi_0}$ . Following the arguments in [44, p. 350], as we did in the proof of Theorem 5.1, Tsuji claims that  $\Delta u_k$  is uniformly bounded on compact subsets of  $X \setminus E_0$  (which would indeed imply that  $\phi$  is smooth on  $X \setminus E_0$ ).

In order to apply the maximum principle to (5.4), one needs to choose  $A > 0$  such that

$$e^{-Au_k}(n + \Delta u_k)$$

is bounded on  $X \setminus E_0$ . But as opposed to the nef case, the positive form  $\omega_0 + dd^c u_k$  is just the push-forward of a form with bounded coefficients, so that it a priori acquires poles along  $X \setminus E_0$  (as was pointed out to us by Demailly) and it does not seem to be obvious to show that the poles in question can be controlled by  $e^{Au_k}$  for some uniform  $A > 0$ .

### 6.3. Analytic Zariski decompositions versus minimal singularities

Let  $L$  be a big line bundle on  $X$ . For each  $m$  we choose a basis of sections  $\sigma_j^{(m)}$  of  $H^0(m!L)$  and let

$$\phi_m := \frac{1}{m!} \log \sum_j |\sigma_j^{(m)}|^2$$

be the associated psh weight on  $L$ . Upon scaling the sections, we can arrange that

$$\phi_m \leq \phi_{m+1} \leq \psi \tag{6.2}$$

for some weight  $\psi$  with minimal singularities on  $L$ .

On the other hand, let  $\sum_{m \geq 1} \varepsilon_m$  be a convergent series of positive numbers. Then one can consider the psh weight

$$\varrho := \log \sum_{m \geq 1} \varepsilon_m e^{\phi_m}.$$

This kind of weight was first introduced by Siu in [36]. Note that we have

$$\varrho \geq \phi_m + \log \varepsilon_m$$

for each  $m$ , which implies that the length with respect to  $\varrho$  of any section  $\sigma \in H^0(kL)$  is bounded on  $X$ .

The following observation was explained to us by J.-P. Demailly.

**PROPOSITION 6.5.** *The psh weight  $\varrho$  of Siu-type has minimal singularities if and only if the algebra*

$$R(L) = \bigoplus_{k \geq 0} H^0(kL)$$

*is finitely generated.*

In order to appreciate this fact, it should be recalled that  $R(L)$  is generally *not* finitely generated when  $L$  is big. In fact if  $L$  is nef and big for instance, then  $R(L)$  is finitely generated if and only if some multiple of  $L$  is base-point free. Classical constructions of Zariski and Cutkosky yield examples in dimension 2 (cf. [32]). Note that Example 5.4 also yields, in particular, an example (in dimension 3).

*Proof.* One direction is clear. If we conversely assume that  $R(L)$  is not finitely generated, then Lemma 6.6 below implies that no  $\phi_m$  can have minimal singularities. Therefore, given any sequence  $C_m > 0$ , we can find, for each  $m$ , some point  $x_m \in X$  such that

$$\phi_m(x_m) \leq \psi(x_m) - C_m. \tag{6.3}$$

It thus follows from (6.2) and (6.3) that

$$e^{\varrho(x_m)} \leq (\varepsilon_1 + \dots + \varepsilon_m) e^{-C_m} e^{\psi(x_m)} + \delta_m e^{\psi(x_m)}$$

with

$$\delta_m := \sum_{l > m} \varepsilon_l.$$

If we choose  $C_m \rightarrow \infty$  fast enough to ensure that

$$(\varepsilon_1 + \dots + \varepsilon_m) e^{-C_m} \leq \delta_m,$$

then we get

$$(\varrho - \psi)(x_m) \leq \log 2\delta_m$$

for all  $m$ , which shows that we cannot have  $\varrho = \psi + O(1)$ . □

The following fact is completely standard modulo the language used.

LEMMA 6.6.  *$R(L)$  is finitely generated if and only if there exists  $m$  such that for each  $l \geq m$  we have  $\phi_l = \phi_m + O(1)$ .*

*Proof.* Let  $\pi: X_m \rightarrow X$  be an increasing sequence of log-resolutions of the base scheme of  $|m!L|$ , so that  $\pi^* m!L = M_m + F_m$ . The assumption amounts to saying that  $F_l$  coincides with the pull-back of  $F_m$  to  $X_l$ , which is in turn equivalent to saying that every section of  $H^0(X_m, l! \pi^* L)$  vanishes along  $F_m$ . But this implies that the canonical inclusion

$$H^0(kM_m) \longrightarrow H^0(k\pi^* m!L)$$

is an isomorphism for each  $k$ , and the result follows since  $R(L)$  is finitely generated if and only if  $R(aL)$  is finitely generated for some integer  $a$ . □

## References

- [1] AUBIN, T., Équations du type Monge–Ampère sur les variétés kähleriennes compactes. *C. R. Acad. Sci. Paris Sér. A-B*, 283 (1976), Aiii, A119–A121.
- [2] — Réduction du cas positif de l'équation de Monge–Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité. *J. Funct. Anal.*, 57 (1984), 143–153.
- [3] BEDFORD, E. & TAYLOR, B. A., A new capacity for plurisubharmonic functions. *Acta Math.*, 149 (1982), 1–40.
- [4] — Fine topology, Shilov boundary, and  $(dd^c)^n$ . *J. Funct. Anal.*, 72 (1987), 225–251.
- [5] BENELKOURCHI, S., GUEDJ, V. & ZERIAHI, A., A priori estimates for weak solutions of complex Monge–Ampère equations. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, 7 (2008), 81–96.
- [6] BERMAN, R. & BOUCKSOM, S., Growth of balls of holomorphic sections and energy at equilibrium. *Invent. Math.*, 181 (2010), 337–394.
- [7] BIRKAR, C., CASCINI, P., HACON, C. D. & MCKERNAN, J., Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.*, 23 (2010), 405–468.
- [8] BŁOCKI, Z., On the regularity of the complex Monge–Ampère operator, in *Complex Geometric Analysis in Pohang* (1997), Contemp. Math., 222, pp. 181–189. Amer. Math. Soc., Providence, RI, 1999.
- [9] BŁOCKI, Z. & KOŁODZIEJ, S., On regularization of plurisubharmonic functions on manifolds. *Proc. Amer. Math. Soc.*, 135 (2007), 2089–2093.
- [10] BOUCKSOM, S., On the volume of a line bundle. *Internat. J. Math.*, 13 (2002), 1043–1063.
- [11] — Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup.*, 37 (2004), 45–76.
- [12] BOUCKSOM, S., DEMAILLY, J.-P., PĂUN, M. & PETERNELL, T., The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. To appear in *J. Algebraic Geom.*
- [13] BOUCKSOM, S., FAVRE, C. & JONSSON, M., Differentiability of volumes of divisors and a problem of Teissier. *J. Algebraic Geom.*, 18 (2009), 279–308.
- [14] CEGRELL, U., Pluricomplex energy. *Acta Math.*, 180 (1998), 187–217.
- [15] — The general definition of the complex Monge–Ampère operator. *Ann. Inst. Fourier (Grenoble)*, 54 (2004), 159–179.
- [16] DEMAILLY, J.-P., Regularization of closed positive currents and intersection theory. *J. Algebraic Geom.*, 1 (1992), 361–409.
- [17] — A numerical criterion for very ample line bundles. *J. Differential Geom.*, 37 (1993), 323–374.
- [18] — *Complex Analytic and Algebraic Geometry*. Book available at <http://www-fourier.ujf-grenoble.fr/~demailly/books.html>.
- [19] DEMAILLY, J.-P. & PĂUN, M., Numerical characterization of the Kähler cone of a compact Kähler manifold. *Ann. of Math.*, 159 (2004), 1247–1274.
- [20] DEMAILLY, J.-P., PETERNELL, T. & SCHNEIDER, M., Compact complex manifolds with numerically effective tangent bundles. *J. Algebraic Geom.*, 3 (1994), 295–345.
- [21] DINEW, S., An inequality for mixed Monge–Ampère measures. *Math. Z.*, 262 (2009), 1–15.
- [22] — Uniqueness in  $\mathcal{E}(X, \omega)$ . *J. Funct. Anal.*, 256 (2009), 2113–2122.
- [23] EIN, L., LAZARSFELD, R., MUSTATA, M., NAKAMAYE, M. & POPA, M., Asymptotic invariants of base loci. *Ann. Inst. Fourier (Grenoble)*, 56 (2006), 1701–1734.
- [24] EVANS, L. C., Classical solutions of fully nonlinear, convex, second-order elliptic equations. *Comm. Pure Appl. Math.*, 35 (1982), 333–363.
- [25] EYSSIDIEUX, P., GUEDJ, V. & ZERIAHI, A., Singular Kähler–Einstein metrics. *J. Amer. Math. Soc.*, 22 (2009), 607–639.

- [26] GUEDJ, V. & ZERIAHI, A., Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.*, 15 (2005), 607–639.
- [27] — The weighted Monge–Ampère energy of quasiplurisubharmonic functions. *J. Funct. Anal.*, 250 (2007), 442–482.
- [28] KISELMAN, C. O., Sur la définition de l’opérateur de Monge–Ampère complexe, in *Complex Analysis* (Toulouse, 1983), Lecture Notes in Math., 1094, pp. 139–150. Springer, Berlin–Heidelberg, 1984.
- [29] KOLLÁR, J. & MORI, S., *Birational Geometry of Algebraic Varieties*. Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998.
- [30] KOŁODZIEJ, S., The complex Monge–Ampère equation. *Acta Math.*, 180 (1998), 69–117.
- [31] — The complex Monge–Ampère equation and pluripotential theory. *Mem. Amer. Math. Soc.*, 178:840 (2005).
- [32] LAZARSEFELD, R., *Positivity in Algebraic Geometry*. I, II. Ergebnisse der Mathematik und ihrer Grenzgebiete, 49. Springer, Berlin–Heidelberg, 2004.
- [33] MABUCHI, T.,  $K$ -energy maps integrating Futaki invariants. *Tohoku Math. J.*, 38 (1986), 575–593.
- [34] RAINWATER, J., A note on the preceding paper. *Duke Math. J.*, 36 (1969), 799–800.
- [35] SIBONY, N., Quelques problèmes de prolongement de courants en analyse complexe. *Duke Math. J.*, 52 (1985), 157–197.
- [36] SIU, Y.-T., Invariance of plurigena. *Invent. Math.*, 134 (1998), 661–673.
- [37] — A general non-vanishing theorem and an analytic proof of the finite generation of the canonical ring. Preprint, 2006. [arXiv:math/0610740v1 \[math.AG\]](#).
- [38] SONG, J. & TIAN, G., Canonical measures and Kähler–Ricci flow. Preprint, 2008. [arXiv:0802.2570v1 \[math.DG\]](#).
- [39] SUGIYAMA, K., Einstein–Kähler metrics on minimal varieties of general type and an inequality between Chern numbers, in *Recent Topics in Differential and Analytic Geometry*, Adv. Stud. Pure Math., 18, pp. 417–433. Academic Press, Boston, MA, 1990.
- [40] TRUDINGER, N. S., Regularity of solutions of fully nonlinear elliptic equations. *Boll. Un. Mat. Ital. A*, 3 (1984), 421–430.
- [41] TSUJI, H., Existence and degeneration of Kähler–Einstein metrics on minimal algebraic varieties of general type. *Math. Ann.*, 281 (1988), 123–133.
- [42] — Finite generation of canonical rings. Preprint, 1999. [arXiv:math/9908078v8 \[math.AG\]](#).
- [43] — Dynamical constructions of Kähler–Einstein metrics. Preprint, 2006. [arXiv:math/0606626v3 \[math.AG\]](#).
- [44] YAU, S. T., On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation. I. *Comm. Pure Appl. Math.*, 31 (1978), 339–411.
- [45] ZERIAHI, A., Volume and capacity of sublevel sets of a Lelong class of plurisubharmonic functions. *Indiana Univ. Math. J.*, 50 (2001), 671–703.

SÉBASTIEN BOUCKSOM  
Institut de Mathématiques de Jussieu  
CNRS–Université Pierre et Marie Curie  
FR-75251 Paris Cedex 05  
France  
boucksom@math.jussieu.fr

PHILIPPE EYSSIDIEUX  
Institut Fourier  
Université Grenoble 1  
FR-38402 Saint-Martin d’Hères  
France  
Philippe.Eyssidieux@ujf-grenoble.fr

VINCENT GUEDJ  
Institut de Mathématiques de Toulouse  
Université Paul Sabatier  
FR-31062 Toulouse Cedex 09  
France  
guedj@math.ups-tlse.fr

AHMED ZERIAHI  
Institut de Mathématiques de Toulouse  
Université Paul Sabatier  
FR-31062 Toulouse Cedex 09  
France  
zeriahi@math.ups-tlse.fr

*Received January 23, 2009*