

# Monomial subdigraphs of reachable and controllable positive discrete-time systems

Rafael Bru<sup>a</sup>, Louis Caccetta<sup>b</sup> and Ventsi G. Rumchev<sup>b</sup>

<sup>a</sup>Dept. de Matemàtica Aplicada, Univ. Politècnica de València,

Camí de Vera s/n, 46071 València, Spain, rbru@mat.upv.es

<sup>b</sup>Dept. of Mathematics and Statistics, Curtin Univ. of Technology,

P.O. Box U 1987, Perth, WA 6845, Australia, {caccetta,rumchev}@maths.curtin.edu.au

## Abstract

Generic structure of reachable and controllable positive linear systems is given in terms of some characteristic components (monomial subdigraphs) of the digraph of a non-negative pair. The properties of monomial subdigraphs are examined and used to derive reachability and controllability criteria in digraph form for the general case when the system matrix  $A$  may contain zero columns. The graph-theoretic nature of these criteria makes them computationally more efficient than their known equivalents. The criteria not only identify the reachability and controllability properties of positive linear systems but also their reachable and controllable parts (subsystems) when the system does not possess such properties.

**Keywords:** Positive linear systems; reachability; controllability; system structure; monomial subdigraphs

## 1 Introduction

Positive discrete-time linear control systems are described by the equation

$$x(t+1) = Ax(t) + Bu(t), \quad t = 0, 1, 2, \dots \quad (1)$$

where  $A = [a_{ij}] \in \mathbb{R}_+^{n \times n}$ ,  $B = [b_{ij}] \in \mathbb{R}_+^{n \times m}$ ,  $x \in \mathbb{R}_+^n$  is the state vector and  $u \in \mathbb{R}_+^m$  is the control vector. The system (1) is denoted by the pair  $(A, B)$  and, when the system is positive, by  $(A, B) \geq 0$ .

A common property of positive systems is that their state evolution is always positive (or at least nonnegative) whenever the initial state is positive (or at least nonnegative). Note that  $A$  and  $B$  being nonnegative matrices is a necessary and sufficient condition for a discrete-time linear system to have nonnegative state evolution for any nonnegative initial state, given that the controls are also nonnegative.

The system (1) is said to be *reachable (controllable from the origin)* if, for any final state  $x_f \geq 0$ , there exist  $k \in \mathbb{N}$  and a nonnegative control sequence  $u(t) \geq 0$ ,  $t = 0, 1, 2, \dots, k$ , transferring the system from  $x_0 = 0$  at  $t = 0$  to  $x_f$  at  $t = k$ . System (1) is called *null-controllable (controllable to the origin)* if, for any initial state  $x_p \geq 0$ , there exist  $k \in \mathbb{N}$  and

a nonnegative control sequence  $u(t) \geq 0$ ,  $t = 0, 1, 2, \dots, k - 1$ , transferring the system from  $x_0 = x_p$  at  $t = 0$  to  $x_f = 0$  at  $t = k$ . The system (1) is *controllable* when it is reachable and null-controllable, (see [11]). Controllability is a fundamental property of the system that shows its ability to move in space. It has direct implications in many control problems such as optimal control, feedback stabilization, nonnegative realizations and system minimality among others.

Characterizations of the reachability of the system (1) can be given in terms of the reachability matrix of the pair  $(A, B)$ . The reachability matrix at time  $k$  is given by

$$\mathcal{R}_k(A, B) = [B|AB|A^2B|\dots|A^{k-1}B]. \quad (2)$$

It is well-known that the pair  $(A, B) \geq 0$  is reachable if and only if the reachability matrix  $\mathcal{R}_k(A, B)$  has a monomial submatrix of order  $n$ , for some  $k \leq n$ . We recall that an  $n$ -dimensional vector is called *i-monomial* if it is a nonzero multiple of the  $i$ th-unit vector  $e_i$  of  $\mathbb{R}^n$ . A monomial matrix consists of  $n$  linearly independent monomial vectors. Throughout this paper we consider nonnegative vectors only.

Different authors have contributed to the characterization of positive reachability and controllability properties, these include Coxson and Shapiro [5], Coxson, Larson and Schneider [4], Rumchev and James [11], Murthy [9], Muratori and Rinaldi [8], Bru, Romero and Sanchez [2] and Caccetta and Rumchev [3]. At the same time, digraphs have been widely used in control theory. It is sufficient to mention only that the notion of structural controllability of linear systems [7] and the criteria to test this property have been formulated in terms of digraphs. However, algebraic methods have been used for the same problems, see for example [5], [8] and [11]. An overview of these results in both forms - algebraic and graph-theoretic, can be found in the very recent monograph by Kaczorek [6]. Moreover, original results on reachability and controllability of continuous-time positive linear systems are also provided in that monograph.

In this paper, in order to increase the understanding of the reachability and controllability properties of positive linear systems, the generic structure of reachable and controllable pairs  $(A, B) \geq 0$ , for the general case when  $A$  may contain zero columns, is given in terms of the digraph of  $A$ . In this way, all possible structures (subdigraphs) of the digraph of  $A$  that can have a reachable or controllable pair  $(A, B)$  are detected and studied.

The paper is organized as follows. In section 2 some basic combinatorial concepts are given as well as the reiteration of a basic but known lemma. The algebraic properties of all different monomial subdigraphs, which can be in the digraph of a reachable pair, are studied in section 3. The characterization of reachability and controllability properties of the system (1) is obtained in section 4. Finally, section 5 contains the conclusions.

## 2 Some preliminaries

Let  $A = [a_{ij}]$  be an  $n \times n$  nonnegative matrix. The *digraph* of  $A$ , denoted by  $D(A)$ , is defined as follows. The set of vertices of  $D(A)$  is denoted as  $N = \{1, 2, \dots, n\}$  and there is an arc in  $D(A)$  from vertex  $i$  to vertex  $j$  if  $a_{ji} > 0$ . The set of all arcs is denoted by  $U$ . A *walk* in  $D(A)$ , from vertex  $i_1$  to vertex  $i_k$ , is an alternating sequence of vertices and arcs, and we will denote it by  $(i_1, \dots, i_k)$ . A walk is called closed if the initial and final vertices coincide. The *length* of a walk is the number of arcs it contains. A walk is said to be a *path* if all its vertices are distinct, and a *cycle* if it is a closed path. The number of arcs directed away from a vertex  $i$  is called the *outdegree* of  $i$  and is written  $\text{od}(i)$ . The number of ingoing arcs of a vertex  $i$  is called

the *indegree* of  $i$  and is denoted by  $\text{id}(i)$ . Note that the number of nonzero entries in the  $i$ th column of  $A$  is  $\text{od}(i)$ , while  $\text{id}(i)$  coincides with the number of nonzero entries of the  $i$ th row.

The positive entries of the columns of matrix  $B$  are associated with the corresponding vertices in  $D(A)$ . Vertices associated with the monomial columns of  $B$  are referred to as *origins*.

We have the following simple but basic result (see [3]).

**Lemma 1.** *Let  $M$  be an  $n \times n$  matrix whose  $j$ th column is  $i$ -monomial. Let  $b$  be an  $n$ -dimensional  $j$ -monomial vector. Then, the product  $Mb$  is an  $i$ -monomial vector. In particular, if  $M^s b$  is  $j$ -monomial, then  $M^{s+1}b$  will be  $i$ -monomial.*

The above lemma tells us that if  $\text{od}(i) = 1$  and  $b$  is an  $i$ -monomial vector then  $Mb$  is a monomial vector as well. However, if  $\text{od}(i) > 1$ , then  $Mb$  is not monomial anymore; in fact, the number of nonzero entries of that product is exactly  $\text{od}(i)$ .

### 3 Monomial subdigraphs

In this section we construct special subdigraphs of the digraph  $D(A)$  called monomial subdigraphs. The common property of monomial subdigraphs is that from the column of  $B$  associated with the initial vertex of a monomial subdigraph one can obtain a maximal sequence of linearly independent monomial vectors  $b, Ab, A^2b, \dots, A^{p-1}b$ , where  $p$  is the number of vertices of the subdigraph. Now, given a nonnegative matrix  $A$  and a path  $(i_1, i_2, \dots, i_{p-1}, i_p)$  of length  $p - 1$  of a digraph  $D(A)$ , we will consider the following special paths.

**Definition 1.** (i) *The above path is said to be an  $i_1$ -**monomial** path if its vertices have outdegree  $\text{od}(i_j) = 1$ , for all  $j = 1, 2, \dots, p-1$  and  $\text{od}(i_p)$  is arbitrary, but  $i_p$  cannot be connected with any other vertex of the path.*

(ii) *When the last vertex of the monomial path has  $\text{od}(i_p) = 0$  then we have a **single** monomial path.*

(iii) *The path  $(i_1, i_2, \dots, i_{p-1}, i_p)$  of length  $p - 1$  with  $\text{od}(i_k) = 1$  for all  $k = 1, 2, \dots, p$  is called a (**monomial**) **cycle** if  $i_1 = i_p$ .*

A monomial path  $(i_1, \dots, i_p)$  of length  $p - 1$  is represented in Figure 1.




Figure 1: Monomial Path

When  $D(A)$  consists of a monomial path or a (monomial) cycle, we have the following result.

**Lemma 2.** Let  $(A, B) \geq 0$  and let  $D(A)$  be a (single) monomial path, with vertices  $(i_1, i_2, \dots, i_p)$  of length  $p - 1$ , where  $p \leq n$  and let  $B$  has an  $i_1$ -monomial column  $b$ . Then,  
(i) the  $p$  vectors

$$b, Ab, A^2b, \dots, A^{p-1}b \tag{3}$$

are linearly independent monomial vectors, and  
(ii) these  $p$  vectors are the maximal number of linearly independent monomial vectors generated by any column of  $B$ .

*Proof.* (i) Apply Lemma 1.

(ii) Since  $\text{od}(i_p) = 0$  (single monomial path) or  $\text{od}(i_p) > 1$  (monomial path), then the vector  $A^p b$  is a zero vector or has, respectively, more than one positive entry, that is it is not monomial anymore. By Lemma 4 and Remark 6 of [3], any nonmonomial column  $b$  cannot generate in (3) as many monomial vectors as the  $i_1$ -monomial column. It is readily seen that the  $i_1$ -monomial column yields at least as many monomial columns as any other monomial column.  $\square$

**Remark.** The results in Lemma 2 hold for monomial cycles. It is not difficult to see that monomial cycles raise a  $p$ -periodic sequence (3). That is,  $A^{k+lp}b = A^k b$  (up to a scalar),  $0 \leq k \leq p - 1$  and  $l = 0, 1, 2, \dots$

**Definition 2.** A subdigraph  $T$  of a digraph  $D(A)$  is called a **monomial tree** if it is a union of different monomial paths, originating at different vertices and connected one to the other from the last vertices only (in  $D(A)$ ) without forming cycles.

Note that the existence of at least a single monomial path in a monomial tree results from the fact that there are no cycles. As cycles are not permitted in monomial trees, the monomial paths of any monomial tree can be grouped in levels as follows. At level  $T_1$  we consider all single monomial paths of that monomial tree. Any monomial path connected from its last vertex only with that of a single monomial path will belong to level  $T_2$ . Any monomial path connected from its last vertex only with that of a monomial path from  $T_2$ , and possibly  $T_1$ , is in level  $T_3$ . By recursion, all levels in the monomial tree can be defined up to the last, which is denoted as  $T_{n-1}$ . The following digraph is a monomial tree of three levels:




Figure 2: Monomial tree

Let  $\mathcal{T}$  be the index set of all initial vertices of all monomial paths of  $T$ . A similar result to Lemma 2 can be obtained for a monomial tree.

**Lemma 3.** Let  $(A, B) \geq 0$  and let  $D(A)$  be a monomial tree  $T$ . Suppose that  $B$  has the  $i$ -monomial columns, for all  $i \in \mathcal{T}$ . Then,

- (i) the vectors generated along each monomial path, as in (3), form a set of linearly independent monomial vectors; the union of all these sets is also linearly independent, and
- (ii) this union is the maximal set of linearly independent monomial vectors generated by any column of  $B$ .

**Definition 3.** Let a digraph  $D(A)$  contains at least one monomial path, one cycle and a tree  $T$ . A subdigraph  $F \subseteq D(A)$  is said to be a **flower** if it consists of a monomial path  $(i_1, i_2, \dots, i_p)$  of length  $p - 1$ , linked to a (monomial) cycle  $(i_{p+1}, i_{p+2}, \dots, i_{p+k+1})$  with the arc  $(i_p, i_{p+1})$ , and moreover, from the vertex  $i_p$  of the monomial path, there are arcs  $(i_p, t)$  for some  $t \in T$ .

Observe that there must be a tree in the digraph  $D(A)$  for a flower to exist, however the flower itself contains only a monomial path and a connected (monomial) cycle. All vertices of a flower have  $\text{od}(i_s) = 1$ , except for the vertex  $i_p$ , in which case  $\text{od}(i_p) \geq 2$ . The figure below is a flower.




Figure 3: Flower

Again, the following result is similar to Lemma 2.

**Lemma 4.** Let  $(A, B) \geq 0$  and let  $D(A)$  be the digraph of  $A$  containing a flower  $F$  connected to a monomial tree  $T$  with  $q$  vertices. Assume that the flower has a monomial path  $(i_1, i_2, \dots, i_p)$ ,  $p \leq n$  linked to the (monomial) cycle  $(i_{p+1}, i_{p+2}, \dots, i_{p+k+1})$ ,  $k \leq n - p - q - 1$ . Suppose that  $B$  has an  $i_1$ -monomial column, namely  $b$ . Then,

- (i) the  $p$  vectors  $\{b, Ab, A^2b, \dots, A^{p-1}b\}$  are linearly independent and monomial. In addition, the  $k + 1$  vectors  $\{A^{q+p}b, A^{q+p+1}b, \dots, A^{q+p+(k+1)}b\}$  are linearly independent and monomial, and
- (ii) the union of both sets gives the maximal number of linearly independent monomial vectors generated by any column of  $B$  along the flower  $F$ .

*Proof.* (i) Since  $\text{od}(i_r) = 1$ ,  $r = 1, 2, \dots, p-1$ , it is clear that the  $p$  vectors  $\{b, Ab, A^2b, \dots, A^{p-1}b\}$  are linearly independent monomial vectors, (see Lemma 2). The vector  $A^p b$  will have at least two positive entries since  $\text{od}(i_p) \geq 2$ . These positive entries correspond to the arcs present from  $i_p$  to a vertex of  $T$ , and to a vertex of the cycle. The vectors  $\{A^p b, A^{p+1}b, \dots, A^{p+k}b\}$  will have at least a positive entry in addition to the  $i_s$ th entry,  $s = p+1, p+2, \dots, p+k+1$ , produced by the cycle. This additional positive entry, namely the  $j$ th, is yielded by the link from  $i_p$  to the monomial tree. However, the  $j$ th entry will eventually become zero, at least for the  $(q + p)$ th power of  $A$ . This is because the entry will ultimately correspond to the final vertex of a single monomial path of  $T$ . So, the  $k + 1$  monomial vectors  $\{A^{q+p}b, A^{q+p+1}b, \dots, A^{q+p+(k+1)}b\}$  will be linearly independent. Clearly, the set  $\{b, Ab, A^2b, \dots, A^{p-1}b, A^{q+p}b, A^{q+p+1}b, \dots, A^{q+p+(k+1)}b\}$  is

formed by linearly independent monomial vectors, since vertices in the flower are distinct.

(ii) Similar to the proof of part (ii) of Lemma 2.  $\square$

Cycles associated with monomial columns of  $B$  produce linearly independent monomial vectors, see Remark after Lemma 2. In addition, (monomial) cycles may yield similar periodic sequences of linearly independent monomial vectors when they are associated with some special columns of  $B$ , called proper, as stated in Lemma 5, the proof of which is similar to that of Lemma 4.

**Lemma 5.** *Let  $(A, B) \geq 0$  and let  $C$  be a (monomial) cycle with vertices  $(i_1, i_2, \dots, i_p = i_1)$ , where  $p < n$ . Also, let  $T$  be a monomial tree with  $q$  vertices in  $D(A)$ . Suppose that  $B$  has a proper column, which can be written as  $b = e_{i_k} + w$ , where  $i_k$  is one of the indices of the cycle, e.g.  $i_k = i_1$ , and when  $w_j > 0$ , then  $j$  is a vertex of the monomial tree  $T$ . Then,*

(i) *the  $p$  vectors  $\{A^q b, A^{q+1} b, \dots, A^{q+p-1} b\}$  are linearly independent monomial vectors, and*

(ii) *these  $p$  vectors are the maximal number of linearly independent monomial vectors generated by any column of  $B$  associated with the cycle  $C$ .*

We can weaken the definition of a monomial tree in order to obtain linearly independent monomial vectors.

**Definition 4.** *A subdigraph  $P$  of a digraph  $D(A)$  is called a **palm** if it is a path  $(i_1, i_2, \dots, i_p)$ , such that  $\text{od}(i_k) = 1$ ,  $k = 1, 2, \dots, p-1$ , and an arbitrary subset of arcs  $(i_p, i_k)$ ,  $k = 1, 2, \dots, p$ .*

It follows from Definition 4 that monomial paths can be considered as a special type of palm without  $\{(i_p, i_k), k = 1, 2, \dots, p\}$ , but not any palm is a monomial path. If  $\text{od}(i_p) = 0$ , the path is, indeed, a single monomial path in a monomial tree. Note that any connection from  $i_{p-1}$  to any monomial tree is excluded. It seems that a flower can be viewed as a particular type of palm, in which the last vertex  $i_p$  is connected with only one vertex  $i_k, k = 2, \dots, p$ . For the existence of flowers, the links  $(i_{k-1}, t), t \in T$  must exist. But such links are not permitted in the palm. For this reason, we have considered the digraph flower independently. In addition, note that any monomial path or cycle that is not in a monomial tree, flower or (monomial) cycle is considered as a palm.

A palm looks like:




Figure 4: Palm

The following properties can be deduced in a similar way as Lemma 2.

**Lemma 6.** *Let  $(A, B) \geq 0$  and let  $D(A)$  be a palm with vertices  $i_1, i_2, \dots, i_p$ , where  $p \leq n$ . Suppose that  $B$  has a column  $b$  which is  $i_1$ -monomial. Then,*

(i) *the  $p$  vectors  $\{b, Ab, A^2 b, \dots, A^{p-1} b\}$  are linearly independent monomial vectors, and*

(ii) *the set of those  $p$  vectors is the maximal set of linearly independent monomial vectors generated by any column of  $B$ .*

Palms can be linked among themselves by arcs  $(i_p, p)$  for some vertex  $p$  of another palm  $P$ , forming a family of palms in which case *other cycles* can appear. Palms can also be linked to any other  $i_1$ -monomial subdigraph by arcs  $(i_p, t)$  for some vertex  $t$  in a monomial tree or flower or (monomial) cycle.

The above lemma is theorem 3 of [4]. In addition, for multi input systems  $(A, B)$  a composition of palms is used in [10] (theorem 1) to study the case where  $A$  does not have any null columns.

## 4 Positive reachability and controllability

It is clear that when a pair  $(A, B)$  is such that  $D(A)$  is one of the monomial subdigraphs introduced in the previous section and  $B$  contains all the columns needed to generate the maximal number of linearly independent monomial vectors on  $D(A)$ , then the pair  $(A, B)$  is reachable. This is because one can obtain a monomial matrix of order  $n$  in the reachability matrix.

With the above results, a characterization of reachable positive systems  $(A, B)$  is given in this section.

For this purpose, consider the nonnegative pair  $(A, B)$  and the associated digraph  $D(A)$ . Recall that the positive entries of the monomial columns of  $B$  are identified with the corresponding vertices in  $D(A)$  called origins. From these origins, construct the maximal monomial subdigraphs, without repeating vertices, in the following order: (i) all possible monomial trees; the initial vertices of all monomial paths of the monomial trees form the index set of origins  $\mathcal{T}$ ; (ii) all possible flowers; the initial vertices of all monomial paths of the flowers form the index set of origins  $\mathcal{F}$ ; (iii) all possible palms; the initial vertices of all paths of the palms form the index set of origins  $\mathcal{P}$ ; (iv) all possible (monomial) cycles from the *proper columns* of  $B$ ,  $b_{l_r} = e_{l_r} + w$ , where the indices of the positive components of vector  $w$  are vertices of a monomial tree; indices  $l_r$  form the set of origins  $\mathcal{C}$ .

Let  $L = \{(i_p, t), i_p \in F, t \in T \text{ and } (i_p, t), i_p \in P, t \in T \text{ or } t \in F \text{ or } t \in P \text{ or } t \in C\}$  be the set of all arcs linking the formed monomial subdigraphs. Define  $D'(A) = D(A) \setminus L = (N', U')$ , where  $N' = N$  and  $U' = U \setminus L$ . Thus, the monomial subdigraphs in  $D'(A)$  are disjoint. The following characterization follows from this construction.

**Theorem 1.** *Let  $A \geq 0$  and let  $D(A)$  be the associated digraph. Let  $\mathcal{T}$ ,  $\mathcal{F}$ ,  $\mathcal{P}$  and  $\mathcal{C}$  be the index sets of origins of the monomial subdigraphs, respectively, monomial trees  $T$ , flowers  $F$ , palms  $P$ , and (monomial) cycles  $C$  of  $D(A)$  formed from the monomial and proper columns of  $B$ . Then, the pair  $(A, B)$  is reachable if and only if  $D'(A)$  is a union of these monomial subdigraphs, that is*

$$D'(A) = \bigcup_{t=1}^{c_t} T_t \bigcup_{f=1}^{c_f} F_f \bigcup_{p=1}^{c_p} P_p \bigcup_{c=1}^{c_c} C_c, \quad (4)$$

where  $c_t, c_f, c_k$  and  $c_p$  stand for the number of monomial trees, flowers, palms and (monomial) cycles, respectively.

*Proof.* Assume that all possible monomial subdigraphs are formed, without repeating vertices, in the order stated above, from the origins  $\mathcal{T}$ ,  $\mathcal{F}$ ,  $\mathcal{P}$  and  $\mathcal{C}$  obtained from the monomial and proper ( $b_{l_r} = e_{l_r} + w$ ) columns of  $B$ .

Suppose that  $D'(A)$  is a union of monomial subdigraphs:

$$D'(A) = \bigcup_{t=1}^{c_t} T_t \bigcup_{f=1}^{c_f} F_f \bigcup_{p=1}^{c_p} P_p \bigcup_{c=1}^{c_p} C_c.$$

Since  $B$  has the monomial columns corresponding to indices of  $\mathcal{T}$ ,  $\mathcal{F}$ , and  $\mathcal{P}$ , and columns of the type  $b_{l_r} = e_{l_r} + w_{l_r}$  corresponding to  $\mathcal{C}$ , then by Lemmas 3–6 we obtain for each monomial subdigraph a maximal set of linearly independent monomial vectors. Since  $D'(A)$  is a union of these monomial subdigraphs and it contains all the vertices of  $D(A)$ , the union of all these vectors is a set of  $n$  linearly independent monomial vectors. This occurs because each vertex of  $D'(A)$  is in one and only one monomial subdigraph, and so the pair  $(A, B)$  is reachable.

Conversely, assume now that (4) does not hold, that is

$$\bigcup_{t=1}^{c_t} T_t \bigcup_{f=1}^{c_f} F_f \bigcup_{p=1}^{c_p} P_p \bigcup_{c=1}^{c_p} C_c \subset D'(A).$$

Then, either  $D'(A)$  has at least one vertex or an arc not included in the union. Denote the digraph formed by the union of those monomial subdigraphs by  $(N'', U'')$ , where  $N''$  is the vertex set and  $U''$  is the arc set.

*Case 1.* Suppose  $N'' \subset N'$ . Hence, the number of vertices in the union is strictly less than the  $n$  vertices of  $D'(A)$  and  $D(A)$ . Then, the maximal number of linearly independent monomial vectors produced by the union of all monomial subdigraphs is strictly less than  $n$ . Since the monomial subdigraphs give the maximal number of such vectors, according to Lemmas 3–6 the reachability matrix  $\mathcal{R}_n$  will not contain a monomial submatrix of order  $n$ . Columns of  $B$ , which are not monomial or proper, are not used in the formation of the monomial subdigraphs of the union. They might produce with columns of  $A$  (corresponding to vertices of a certain subdigraphs, which are in  $D'(A)$  and not in the union of monomial subdigraphs) some linearly independent monomial vectors in the sequence  $b, Ab, A^2b, \dots, A^k b$ . However, the number of such vectors is less than the number of linearly independent monomial vectors generated by the monomial column of  $B$ , corresponding to the origins, applied to the same subdigraph, see the proof of Lemma 2. Therefore,  $\mathcal{R}_n$  will not contain an  $n \times n$  monomial submatrix and the pair  $(A, B)$  is not reachable.

*Case 2.* Now  $U'' \subset U'$ . Since all arcs connecting monomial subdigraphs are in  $L$ , the strict inclusion is due to the existence of an arc not included in  $L$ . Such an arc has a vertex in  $D'(A)$ , but not in the union, that is  $N'' \subset N'$ , and then we can proceed as in Case 1.  $\square$

The following examples illustrate the monomial subdigraphs of the digraph of a pair  $(A, B) \geq 0$ .

**Example 1.** Let

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad B_1 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$



The digraph of  $A$  is




Figure 5:  $D(A)$

First note that the matrix  $B_1$  has the unit vectors  $e_4$ ,  $e_8$  and  $e_3$ . It is clear that starting from:

- vector  $e_4$  the single monomial path (4), of length zero, is found and it will be in the monomial tree  $T$ ;
- vector  $e_3$  a flower,  $F$ , is found and it is formed by the monomial path (3), of length zero, and the (monomial) cycle (7, 2); note that vertex 3 is connected with this cycle and there is an arc from vertex 3 to  $T$ , the arc (3, 4);
- vector  $e_8$  the monomial path (8, 5, 1), of length two, is obtained and it will be in the set of palms  $P$ ; it cannot be in  $T$  because this would produce a cycle in  $T$ ;
- vector  $b = e_6 + w$ , where the positive components of  $w$  are just the 4th component and this index is a vertex of  $T$ ; so, from that vector, one can consider the cycle (6, 9), which is a (monomial) cycle  $C$ .

It is clear that  $D'(A) = T \cup P \cup F \cup C$  and thus the pair  $(A, B_1)$  is reachable. The arcs of  $D(A)$  not included in  $D'(A)$  are  $L = \{(1, 4), (1, 6), (3, 4)\}$ . It is worth noticing that the same decomposition can be obtained if the matrix  $B_1$  has the monomial column  $e_6$  instead of the column  $b_4$ .

**Example 2.** Let  $A$  be the matrix of Example 1 and let

$$B_2 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

In this case, the monomial tree  $T$ , the flower  $F$  and the palm  $P$  previously described are obtained starting from the vertices which correspond to the first three columns of  $B_2$ . However, from the fourth column  $b = e_6 + e_8 + e_1$  one can not obtain monomial vectors because vectors  $A^k b$ ,  $k = 0, 1, 2, \dots$  have more than one positive component. The cycle (6, 9) cannot be obtained and  $T \cup P \cup F \subset D'(A)$ . Hence, the pair  $(A, B_2)$  is not reachable.

Following the approach proposed in this paper we can identify reachable parts (monomial trees, flowers, palms, cycles) of the system matrix  $A$  even when the pair  $(A, B)$  is not reachable. The positive system can be made reachable by applying suitable controls (see Examples 1 and 2).

As it is well known, reachability from zero plus nilpotence is equivalent to controllability (see [5] and [11]). It is thereby sufficient to eliminate all possible monomial subdigraphs of  $D(A)$  with cycles for obtaining the controllability property. In fact, we can establish the following result.

**Theorem 2.** *Let  $(A, B) \geq 0$  and let  $D(A)$  be the associated digraph of  $A$ . Then, the pair  $(A, B)$  is controllable if and only if,  $D(A) = \bigcup_{t=1}^{c_T} T_t$  and  $B$  contains all monomial columns corresponding to the origins of the monomial trees.*

## 5 Conclusion

In this paper, reachability and controllability properties of discrete-time positive linear systems, in the more general case when the system matrix contains zero columns are established in terms of the digraph of the pair  $(A, B)$ . Monomial subgraphs of reachable and controllable nonnegative pairs  $(A, B)$  are identified and their properties studied. Criteria in digraph form recognising the reachability and controllability properties of such pairs are obtained in the paper. These criteria give a better understanding of the structure of reachable and controllable discrete-time positive linear systems than the corresponding criteria in algebraic form. The results obtained in this paper can be used to develop computationally efficient combinatorial algorithms for revealing such fundamental properties of discrete-time positive linear systems as reachability and controllability.

*Acknowledgement:* The authors wish to thank to The Australian Research Council for providing support for this work under grant A00001126. The work of the first author was also supported by Spanish grant DGI number BFM2001-2783 and Universitat Politècnica de València under its research incentive program.

## References

- [1] A. Berman and R.J. Plemmons, *Nonnegative matrices in the mathematical sciences*, SIAM, Philadelphia, 1994.
- [2] R. Bru, S. Romero and E. Sánchez, “Canonical forms for positive discrete-time linear control systems”, *Linear Algebra and its Applications*, 310, 49–71, 2000.
- [3] L. Caccetta and V.G. Rumchev, “Reachable discrete-time positive systems with minimal dimension control sets”, *Dynamics of Continuous, Discrete and Impulsive Systems*, 4, 539–552, 1998.
- [4] P.G. Coxson, I.C. Larson and H. Schneider, “Monomial patterns in the sequence  $A^k b$ ”, *Linear Algebra and its Applications*, 94, 89–101, 1987.
- [5] P.G. Coxson and H. Shapiro, “Positive input reachability and controllability of positive systems”, *Linear Algebra and its Applications*, 94, 35–53, 1987.

- [6] T. Kaczorek, *Positive 1D and 2D systems*, Springer–Verlag, Heidelberg, 2002.
- [7] C.T. Lin, “Structural controllability”, *IEEE Transactions on Automatic Control*, vol. AC-19, no. 3, pp.201–208, 1974.
- [8] S. Muratori and S. Rinaldi, “Excitability, stability, and sign of equilibria in positive linear systems”, *Systems and Control Letters*, 16, 59–63, 1991.
- [9] D.N.P. Murthy, “Controllability of a linear positive dynamic system”, *Int. J. Systems Sci.*, vol. 17, no. 1, 49–54, 1986.
- [10] V.G. Rumchev, “On controllability of discrete–time positive systems”, in *Proc. The Sixth Intern. Conference on Control, Automation, Robotics and Vision*, Singapore, December, 2000.
- [11] V.G. Rumchev and D.J.G. James, “Controllability of positive discrete–time systems”, *International Journal of Control*, vol. 50, no. 3, 845–857, 1989.