

1 **More green and less blue water in the Alps during warmer summers**

2 Theodoros Mastrotheodoros¹, Christoforos Pappas², Peter Molnar¹, Paolo Burlando¹, Gabriele
3 Manoli^{1,3}, Juraj Parajka⁴, Riccardo Rigon⁵, Borbala Szeles⁴, Michele Bottazzi⁵, Panagiotis
4 Hadjidoukas⁶, and Simone Fatichi^{1*}

5 ¹ETH Zurich, Institute of Environmental Engineering, Zürich, Switzerland.

6 ²Département de géographie and Centre d'études nordiques, Université de Montréal, Montréal,
7 QC, Canada.

8 ³Department of Civil, Environmental and Geomatic Engineering, University College London,
9 London, UK.

10 ⁴Centre for Water Resource Systems, Vienna University of Technology, Vienna, Austria.

11 ⁵Department of Civil, Environmental and Mechanical Engineering, DICAM, University of
12 Trento, Trento, Italy.

13 ⁶IBM, Zurich Research Laboratory, Zürich, Switzerland.

14 **Climate change can reduce surface-water supply by enhancing evapotranspiration in**
15 **forested mountains, especially during heatwaves. We investigate this “drought paradox”**
16 **for the European Alps using a 1212-station database and hyper-resolution**
17 **ecohydrological simulations to quantify blue (runoff) and green (evapotranspiration)**
18 **water fluxes. During the 2003 heatwave, evapotranspiration in large areas over the Alps**
19 **was above average despite low precipitation, amplifying the runoff deficit by 32% in the**
20 **most runoff-productive areas (1300-3000 m above sea level). A 3 °C air temperature**
21 **increase could enhance annual evapotranspiration by up to 100 mm (45 mm on average),**
22 **which would reduce annual runoff at a rate similar to a 3% precipitation decrease. This**

* Corresponding author

23 **suggests that green water feedbacks--often poorly represented in large-scale model**
24 **simulations--pose an additional threat to water resources, especially in dry summers.**
25 **Despite uncertainty in the validation of the hyper-resolution ecohydrological modelling**
26 **with observations, this approach permits more realistic predictions of mountain region**
27 **water availability.**

28 Although relatively small in size, the European Alps (hereafter “Alps”) contribute a
29 disproportionally large amount of water, especially during summer, to four major European
30 rivers¹, in the basins of which reside more than 170 million people². For this reason, they are
31 referred to as “the water towers of Europe”³. At the same time, water scarcity and droughts in
32 central Europe are becoming more frequent⁴. The summer droughts of 2003, 2010, 2015 and
33 2018 have raised concerns about the vulnerability of the European water budget to climate
34 change^{2,5} as these events have affected more than 17% of the European population with an
35 annual economic impact exceeding € 6.2 billion between 2001 and 2006⁶. Temperature in the
36 Alps is increasing at a fast pace⁷, relative humidity is generally decreasing⁸, evapotranspiration
37 (ET) is increasing⁹, Alpine glaciers are shrinking and the distribution of snow is shifting to
38 higher elevation¹⁰, while climatic extremes are becoming more frequent¹¹. The complex
39 topography, the interactions between water and vegetation and the multiple processes shaping
40 the water cycle in mountainous areas hinder the quantification of the different water budget
41 components in traditional large-scale climate change impact assessment studies¹². For example,
42 climate change can shift the partitioning of water fluxes in the hydrosphere and biosphere
43 moving blue water (runoff and streamflow) into green water (ET)^{13,14}. Quantifying how these
44 fluxes change with elevation, seasonally, and interannually is an important and challenging
45 scientific question.

46 Large uncertainties are associated with the vegetation response to water stress^{15,16}. Studies in
47 different parts of the Alps have found contrasting impacts of droughts on vegetation¹⁷, spanning

48 from increased mortality in dry inner-Alpine valleys¹⁸ to enhanced productivity in wet pre-
49 Alpine hills¹⁹. These discrepancies emphasize that extrapolating results of specific case
50 studies²⁰ to the entire Alpine domain or downscaling results of coarse resolution modelling
51 studies¹⁴ is problematic²¹. The largest component of land ET is plant transpiration²², which is
52 poorly quantified due to the large variability in plant water use strategies and stomata sensitivity
53 to water stress^{23,24}. Land-surface models often represent the links between soil moisture and
54 transpiration in a simplistic way, without mechanistic representation of the soil-plant-
55 atmosphere continuum^{5,25,26}. Most importantly, they do not resolve land-surface energy and
56 water fluxes at sufficient spatial resolution to capture local topographic and microclimatic
57 effects, and often ignore lateral flows of water^{12,27-29}.

58 Here, to overcome these limitations, we combined a new pan-Alpine database with hyper-
59 resolution ecohydrological simulations to test the “drought paradox” hypothesis, i.e. that during
60 droughts ET may increase at high elevations in a large part of the Alps, thus amplifying the
61 runoff deficit. Plot-scale observations provided evidence of the drought paradox
62 occurrence^{17,30}, but the extent and implications of ET changes during heatwaves at the pan-
63 Alpine scale remain unexplored. We compiled a new dataset comprising meteorological,
64 discharge, and snow depth measurements at 1212 stations and combined it with distributed
65 hydrometeorological products to generate high-resolution forcing (precipitation (P), air
66 temperature, solar radiation, wind speed, and relative humidity; see Methods) to drive and
67 validate the ecohydrological model Tethys-Chloris (T&C)²¹. The model resolves the water,
68 carbon and energy budgets at the hourly time scale in a physically based and spatially explicit
69 manner, accounting for lateral water transfer and topographic effects on radiation. The model
70 has been extensively validated in many regions worldwide^{21,22,31}, including Alpine
71 ecosystems³². To account for the high spatial heterogeneity of the region, we performed
72 massively parallel simulations (6.1×10^5 CPU hours) at an unprecedented high resolution (250
73 m grid) for the entire Alpine arch (257,045 km², 4.12 million pixels, Supplementary Fig. 1).

74 The simulation period consists of three hydrological years (2001-2003, Fig. 1 and
75 Supplementary Tables 1-4), including a very wet and a very dry year (2001 and 2003,
76 respectively, Supplementary Fig. 2). We used these simulations for partitioning the pan-Alpine
77 water budget into blue and green water fluxes and we quantified the sensitivity of each
78 component to changes in precipitation and temperature. We validated the model output with
79 daily discharge and snow depth measurements, as well as with CO₂, water, and energy fluxes
80 observed by eddy covariance flux towers located in the Alps or nearby (see Methods and
81 Supplementary Table 5). This validation yielded very satisfactory results, considering that
82 model parameterization was only based on literature values and calibration with previous plot-
83 scale model applications in the Alpine region (see Methods, Extended Data Figure 1 and 2, and
84 Supplementary Figs 3, 4, and 5). We further compared the simulated distributions of ET and
85 gross primary production (GPP) over the entire domain with other products (TRENDY^{33,34},
86 FLUXCOM³⁵ and ERA Interim³⁶). Averaged in space, these products provide estimates
87 comparable to the T&C simulations (Extended Data Figure 3, Supplementary Figs 6, see
88 Methods). However, due to their coarse resolution compared to our analysis, they largely
89 underestimate the spatial variability of ET and GPP and they provide spatial patterns that are
90 poorly resolved. This reinforces the need for hyper-resolution simulations in complex terrain.

91 **Figure 1 | Simulation results highlight the spatial heterogeneity in latent heat**
92 **(evapotranspiration in energy units). a,** The spatial extent of the European Alps. **b,**
93 November 2000-October 2003 average latent heat flux (W m⁻²) for the entire 257,045 km²
94 domain simulated with Tethys-Chloris. **c-d** zoom on the Bernese highlands, Switzerland,
95 and illustration of the small-scale spatial heterogeneity captured with the hyper-resolution
96 simulation (250 m x 250 m pixels).

97 **Dissecting the water towers of Europe**

98 Simulation results indicate that latent heat, i.e. total ET expressed in units of equivalent energy,
99 reaches its maximum values in wetter areas (e.g., in the north in Fig. 1c, where annual
100 precipitation exceeds 2000 mm), especially on south-facing slopes (Fig. 1d) confirming that
101 energy is the dominant driver of ET in this area. In drier regions, such as in the SE valley in
102 Fig. 1c (upper Rhone valley), latent heat is overall lower because P ($\sim 500 \text{ mm yr}^{-1}$) becomes
103 the critical constraint for annual ET. High-elevation areas are clearly distinguishable because
104 rocks, snow, and ice emit low latent heat (Fig. 1b and c).

105 Analyzing the three-year average water fluxes across the entire elevation range, we found that
106 the areas between 200 and 300 m a.s.l. contribute proportionally the largest fraction to total ET.
107 At higher elevations the contribution to ET declines faster than the fractional area, despite the
108 slightly increasing contribution of precipitation (Fig. 2a). The elevational distribution of ET
109 varies considerably between different catchments, due to climatic and vegetation heterogeneity
110 and the interplay between water and energy limitations (Supplementary Fig. 7)³⁷. We used
111 three-year average $P-ET$ as a proxy for runoff (Extended Data Figure 2)³⁸, since changes in soil
112 and snow water storage over three hydrological years could be considered small, and ice melt
113 only marginally contributed to the total water budget (less than 3%) at the annual scale at a rate
114 of roughly $4 \text{ km}^3 \text{ yr}^{-1}$ (Extended Data Figure 4). Runoff production, $P-ET$, peaks at around 800
115 m a.s.l. (Fig. 2a). However, more than 50% of the blue water originates from the areas between
116 1300 and 3000 m a.s.l., which correspond to only 35% of the total Alpine domain. This can be
117 explained by the sharp decrease of ET with elevation due to temperature constraints (Fig. 2a).
118 The runoff production shows a great spatial variability, and even neighboring catchments, such
119 as the upper Rhone and the Aare catchment in Switzerland, may exhibit distinct $P-ET$ patterns
120 (Fig. 2b).

121 **Blue vs. green water during an exceptionally dry and warm summer**

122 The Alpine water budget also displays high temporal variability (Fig. 2c); P-ET in 2001 was
123 53% higher than in 2003 (Extended Data Figure 4), which can be explained by both higher
124 precipitation and lower ET. More specifically, the Alps received 225 mm more P in 2001
125 compared to 2003 (1363 and 1138 mm, respectively, averaged over the entire domain) while
126 ET was 30 mm lower on average (Fig. 2c).

127 We used discharge measurements from 381 catchments across the entire domain to validate the
128 simulations and assess the severity of the 2003 drought. We focused on analyzing runoff
129 deficits, thus we excluded 47 locations in which observed 2003 runoff was higher than the long-
130 term average and/or catchment P in 2003 was higher than the 2001-2003 mean. For the
131 remaining 334 catchments, we computed how much ET contributed to amplifying the effect of
132 precipitation deficit on runoff during the 2003 growing season (here defined as the May-
133 September period to isolate the period with active vegetation, when green water feedback can
134 be pronounced, Supplementary Fig. 8). We found that in 75% of the catchments, ET amplified
135 the drought impact on runoff. The remaining 25% of the catchments - mostly located in the SW
136 and NE of the pan-Alpine domain - experienced dry conditions with water-stressed vegetation
137 and reduced ET. Considering the entire domain, ET increased during the drought in an area
138 covering more than 144,000 km² (Supplementary Fig. 9). Overall, the increase in green water
139 flux amplified the precipitation-driven deficit by roughly 22% (Fig. 2c and Supplementary Fig.
140 8). In the areas between 1300 and 3000 m a.s.l., enhanced ET created an additional water loss
141 of almost 4 km³ during the 2003 growing season compared with the 2001-2003 growing season
142 average, amplifying the runoff decrease due to precipitation by 32% (mean weighted by the
143 area of each catchment).

144 **Figure 2 | Relationship between elevation and blue and green water fluxes. a,** Percent
145 contribution of each elevation class (grouped in 100 m elevation bins) to precipitation (P),

146 evapotranspiration (ET) and P-ET (the fluxes are averaged over the entire simulation period,
147 i.e. 2001-2003), including the fractional area of each class for the entire domain. For the
148 latter, the areas between 1300 and 3000 m a.s.l. (from which more than 50% of the total pan-
149 Alpine runoff originates) are highlighted. **b**, P-ET in mm yr^{-1} . The dashed blue line and the
150 shaded area represent the median P-ET (averaged over the entire simulation period) over the
151 entire domain and the interquartile range, respectively. Colored lines show the median P-ET
152 elevational distribution for selected catchments to illustrate the spatial variability. A locator
153 map is included. **c**, ET anomalies during the 2003 growing season and for the +3 °C scenario
154 (solid lines; left y-axis) and ET contribution to the runoff deficits compared with the
155 precipitation deficit during the 2003 growing season (solid points; right y-axis). The
156 anomalies were computed based on the 2001-2003 mean and the fluxes were averaged in
157 space based on the 100 m elevation bins. Each point represents one of the 334 catchments
158 (out of the 381 in total) for which catchment-averaged precipitation in 2003 was lower than
159 in the period 2001-2003 and runoff in 2003 was below the long-term average (the latter was
160 computed using the entire available record for each station, as summarized in the
161 Supplementary Table 3). For ET anomalies, the interquartile ranges are also shown (shaded
162 cyan and magenta areas).

163 To quantify the sensitivity of ET to warming only and remove the effect of reduced precipitation
164 (2003 was not only the warmest summer but also exceptionally dry, see Supplementary Fig. 2),
165 we performed a scenario analysis exploiting the correlation between the spatial distribution of
166 simulated mean annual ET and mean annual temperature (space-for-time substitution, see
167 Methods). This procedure was necessary because multiple simulations with T&C at a so high
168 spatial resolution are computationally too expensive. We found that with a 3 °C increase in air
169 temperature, annual ET would increase on average by 6% (evaporation will increase by 9% and
170 transpiration will increase by 5%) while P-ET would decrease by roughly 5% under the
171 assumption of constant precipitation (Fig. 2c). A similar effect on runoff is expected if annual

172 precipitation is reduced by only 3% (change in P-ET for unchanged ET), which is likely to
173 happen by the end of the century³⁹. The sensitivity of annual ET to temperature shifts from
174 positive to negative below 700 m a.s.l., which implies that in a warmer climate ET will likely
175 decrease at low elevations (Fig. 2c and Extended Data Figure 4), because of higher water stress
176 and earlier senescence of grass and leaf-shedding in deciduous forests. A recent study found
177 similar patterns with a general positive ET sensitivity to temperature increases in the Sierra
178 Nevada mountains in California⁴⁰.

179 During the growing season, precipitation is still the main source of blue water (81% on
180 average), snowmelt comes second (16%), and ice melt accounts for the remaining 3%
181 (Extended Data Figure 4). Below 500 m a.s.l., the contribution of soil water stored during winter
182 and early spring and depleted during the growing season is substantial (99 mm, compared with
183 464 mm of ET, Extended Data Figure 4). The simulated growing season ET for the +3 °C
184 scenario is generally higher than the simulated ET during 2003 (when the growing season
185 temperature anomaly was roughly +3 °C) because precipitation during 2003 was very low
186 (Supplementary Fig. 2), but overall the two patterns agree well (Extended Data Figure 4). This
187 reinforces our confidence on the space-for-time approach employed.

188 Blue and green water fluxes averaged over the entire domain show that ET sporadically
189 exceeded precipitation during the growing season of 2001 and 2002. For 2003, however, ET
190 was higher than precipitation (which was 32% lower than the long-term average) already before
191 the beginning of the growing season (Supplementary Fig. 10). The earlier snowmelt, which
192 peaked before the start of the growing season and plummeted afterwards, amplified this
193 precipitation deficit. Early snowmelt is becoming more frequent with rising temperature¹⁰ and
194 in 2003 it was only partly compensated by increased ice melt. The simulated ice melt during
195 August 2003 was 38% of the total Alpine runoff in August, corresponding to about 2 km³ of
196 water.

197 **Figure 3 | Analysis of anomalies in blue and green water fluxes during the 2003**
198 **drought. a**, Histogram of observed May-September 2003 total runoff anomalies in mm for
199 381 locations. **b**, Histogram of observed May-September 2003 runoff anomalies in
200 percentage for the same locations colored according to the magnitude of the anomaly (<-
201 75%: yellow, -50 to -75%: cyan, -25 to -50%: red and >-25%: black), growing season 2003
202 is compared with the mean of each station for the period 2001-2003. **c**, Spatial distribution
203 of the simulated ET anomaly in mm during the 2003 growing season (May-September, the
204 reference period for ET is also 2001-2003). The dots represent the 381 locations with
205 hydrological measurements and are colored as described in (b). The three insets in the lower
206 right part show the boxplots of simulated ET anomaly in May-September 2003 for three
207 different vegetation types (evergreen, deciduous, and grassland) in three elevation classes
208 (<1000 m a.s.l., 1000-2000 m a.s.l. and >2000 m a.s.l.). The box length provides the
209 interquartile range (I_{QR}), the bottom of the box the 25th percentile (first quartile, q₁), the top
210 of the box the 75th percentile (third quartile, q₃), and the horizontal line within the box the
211 median value. The lower whisker corresponds to q₁ - 1.5I_{QR}, and the upper whisker
212 corresponds to q₃ + 1.5I_{QR}.

213 Considering all 381 discharge stations, the observed runoff in May-September 2003 was on
214 average 50% lower than the long-term mean at each station in the period May-September (Fig.
215 3 and Supplementary Fig. 10). Higher-than-average runoff occurred at a few locations, mostly
216 in highly-glacierized catchments (Fig. 3)⁴¹. During this period, precipitation over the Alps was
217 the lowest recorded between 1992-2008 and mean temperature was record-breaking high
218 (Supplementary Fig. 2). The detailed vegetation scheme in T&C allows an analysis of different
219 vegetation responses to the 2003 drought (Fig. 3). Most evergreen forests strongly benefited
220 from the increased radiation and temperature, and did more so as the drought intensified during
221 the summer months, mostly at high elevations, where ET increased. Grasslands and deciduous
222 forests were water-stressed below 1000 m a.s.l., but benefited from the drought above 2000 m

223 a.s.l., especially at the beginning of the growing season, when monthly ET increased by up to
224 60 mm (Fig. 3). In most areas of the Northern Alps the simulated ET anomaly was positive
225 throughout the summer, in agreement with local measurements¹⁹. In all dry inner Alpine valleys
226 (such as Valle d’Aosta and Val d’Adige in Italy, Valais in Switzerland and Murtal in Austria),
227 lateral subsurface water redistribution leads to higher soil moisture and ET only in proximity
228 of the streams (Extended Data Figure 5), but it cannot outweigh the low precipitation over the
229 entire valley. These conditions render irrigation necessary to sustain vegetation productivity
230 during summer in these areas⁴². This becomes even clearer during the 2003 drought, when very
231 low soil moisture (Extended Data Figure 5) led to increased water stress and reduced ET for all
232 vegetation types in the valleys⁴³. However, the fractional area occupied by valleys is
233 disproportionally smaller than the area with increased ET at high elevation (Fig 3), which
234 implies that the role of these valleys for the overall Alpine water budget is minimal (Fig. 4).
235 We remark that due to poor knowledge of subsurface conditions, we assumed 1 m uniform soil
236 depth over the entire domain (except rocky areas and lakes), which leads to a simplified
237 simulation of subsurface unsaturated and saturated dynamics and likely de-emphasizes the role
238 of deep-storage and groundwater flow^{44,45}. This could explain the excessive drying during
239 summer in certain locations. This simplification together with the use of invariable rooting
240 depths for the various vegetation types across space are potentially important limitations, but
241 the relatively good performance of T&C for discharge, even during the most extreme low-flow
242 conditions of August 2003 (Extended Data Figure 6), suggests that our inferences about ET are
243 also robust, despite some discrepancy in the ET anomaly with other distributed products
244 (Extended Data Figure 7).

245 **Discussion**

246 Our results –derived from a single mechanistic model– indicate that ET considerably
247 contributed to reduce water yield during the 2003 growing season because vegetation benefited

248 from the unusually warm and sunny conditions in a large part of the Alpine region at higher
249 elevations. At the annual timescale, however, the temperature-driven ET feedbacks on runoff
250 are less important than the direct effect of changes in precipitation; a 3% reduction in annual
251 precipitation would affect runoff production over the entire pan-Alpine domain similarly to a
252 hypothetical increase in mean annual air temperature by 3 °C. Note, however, that the scenario
253 of +3 °C change in air temperature is simply based on a space-for-time analysis. Thus, it could
254 overestimate the response because other meteorological variables covary with elevation. It
255 further does not account for non-climatic factors (e.g., plant acclimation) that might lead to
256 different vegetation responses. Similarly to our results, previous research in the California's
257 Sierra Nevada showed that the 2012-2015 drought in that region caused severe runoff deficits,
258 while ET was less influenced⁴⁶. However, for unchanged precipitation, increases in temperature
259 are projected to mostly enhance ET⁴⁰, which implies a negative feedback on runoff. Annual
260 precipitation in the Alps has shown no long-term trends so far⁴⁷, but summer precipitation in
261 central Europe has decreased and could further do so in the future due to changes in atmospheric
262 circulation patterns leading to more intense summer droughts⁴⁸. Combined with the expected
263 decrease in ice melt and earlier snowmelt¹⁰, our results demonstrate that blue water could be
264 considerably reduced in the European Alps during warmer summers, but green water will
265 continue to increase⁹, leading to the oxymoron “lush vegetation-drier rivers” (Fig. 4). In the
266 near future events like the 2003 drought may be no longer classified as extreme¹¹, but the
267 combined effects of spatially variable changes in the timing and magnitude of precipitation will
268 likely be more complicated, which could lead to different ET changes and potentially affect
269 plant water stress patterns. On the other hand, the expected increase in plant water use efficiency
270 with higher levels of CO₂ concentration than present⁴⁹ as well as large-scale disturbances (e.g.
271 forest mortality⁵⁰), species changes, and plant acclimation, which are not considered here, may
272 partially offset this ET feedback during warmer summers in the long-term but they will unlikely
273 have a major role in the near future. Furthermore, in certain regions of the Alps, vegetation

274 management is intense and past disturbances such as wildfires or forest logging may have
275 influenced vegetation composition and function in ways that are not accounted for in the model
276 initialization.

277 While the presented concepts are general, the extension of the results to other mountain regions
278 strongly depends on the relative magnitude of P and ET at the annual scale and during summer
279 (Fig. 4). Important factors are also the elevation at which P-ET shifts from positive to negative
280 during warm and dry summers and the areal extent covered by different elevation bands and
281 vegetation types (Fig. 4). Nevertheless, results from Sierra Nevada^{40,46} largely agree with our
282 findings.

283 **Figure 4 | Conceptual representation of the drought paradox in the Alps.** **a**, Elevation
284 profile and P-ET fluxes for average (in light blue) and dry and warm (in yellow) growing
285 seasons across contrasting subregions of the Alps (1, plains and hills; 2, dry valleys; 3,
286 forested mountains; 4, rocks and ice). **b**, Blue (P and P-ET) and green (ET) water fluxes in
287 average (filled bars) and warm and dry (empty bars) growing seasons in contrasting Alpine
288 subregions (as explained in (a) and shown in the map in (c), in mm). **c**, Map of the four
289 distinct areas and the contribution of each area to runoff production, the circle size is
290 proportional to the km³. Contribution to runoff production was computed as P-ET +
291 snowmelt + ice melt - ΔS , where ΔS denotes the difference in soil water storage from the
292 beginning to the end of the growing season. The filled circles show the water volumes for
293 average growing seasons and the empty circles show the water volumes for warm and dry
294 growing seasons for the four subregions.

295 Understanding the partitioning of green and blue water fluxes and their spatial distribution from
296 few square kilometers to the entire Alps is essential to manage the European water resources
297 under current and future climatic conditions^{2,14}. This partition has implications on ecosystem
298 functioning, energy production, and water supply. We showed that ecohydrological simulations

299 driven by high-resolution hydrometeorological forcing improve the quantification and
300 understanding of the water budget in mountainous areas and its vulnerability to climate,
301 providing insights into processes that coarser-scale approaches fail to reproduce^{12,27,51}. The
302 need for more realistic, high-resolution quantifications of water availability is urgent⁵². Our
303 study demonstrates that recent advances in ecohydrological modeling, combined with large
304 scale datasets and new computational capabilities, offer the possibility to address this urgent
305 need, thus helping to define strategies to counteract or adapt to climate change impacts on water
306 resources.

307 **References**

- 308 1. Weingartner, R. & Viviroli, D. Water resources in mountain regions: a methodological
309 approach to assess the water balance in a highland-lowland-system. *Hydrol. Process.*
310 **21**, 578–585 (2007).
- 311 2. European Environment Agency (EEA). *Regional climate change and adaptation. The*
312 *Alps facing the challenge of changing water resources.* (2009).
- 313 3. Viviroli, D., Dürri, H. H., Messerli, B., Meybeck, M. & Weingartner, R. Mountains of
314 the world, water towers for humanity: Typology, mapping, and global significance.
315 *Water Resour. Res.* **43**, 1–13 (2007).
- 316 4. Briffa, K. R., van der Schrier, G. & Jones, P. D. Wet and dry summers in Europe since
317 1750: evidence of increasing drought. *Int. J. Climatol.* **29**, 1894–1905 (2009).
- 318 5. Teuling, A. J. A hot future for European droughts. *Nat. Clim. Chang.* **8**, 364–365
319 (2018).
- 320 6. European Commission. *Water scarcity and droughts - In-depth assessment.* (2007).
- 321 7. Brunetti, M. *et al.* Climate variability and change in the Greater Alpine Region over the

- 322 last two centuries based on multi-variable analysis. *Int. J. Climatol.* **29**, 2197–2225
323 (2009).
- 324 8. Fatichi, S., Molnar, P., Mastrotheodoros, T. & Burlando, P. Diurnal and seasonal
325 changes in near-surface humidity in a complex orography. *J. Geophys. Res.* **120**, 2358–
326 2374 (2015).
- 327 9. Duethmann, D. & Blöschl, G. Why has catchment evaporation increased in the past 40
328 years? A data-based study in Austria. *Hydrol. Earth Syst. Sci.* **22**, 5143–5158 (2018).
- 329 10. Beniston, M. *et al.* The European mountain cryosphere: a review of its current state,
330 trends, and future challenges. *Cryosph.* **12**, 759–794 (2018).
- 331 11. Samaniego, L. *et al.* Anthropogenic warming exacerbates European soil moisture
332 droughts. *Nat. Clim. Chang.* **8**, 421–426 (2018).
- 333 12. Fan, Y. *et al.* Hillslope Hydrology in Global Change Research and Earth System
334 Modeling. *Water Resour. Res.* 1–36 (2019). doi:10.1029/2018WR023903
- 335 13. Falkenmark, M. & Rockström, J. The New Blue and Green Water Paradigm: Breaking
336 New Ground for Water Resources Planning and Management. *J. Water Resour. Plan.*
337 *Manag.* **132**, 129–132 (2006).
- 338 14. Orth, R. & Destouni, G. Drought reduces blue-water fluxes more strongly than green-
339 water fluxes in Europe. *Nat. Commun.* **9**, (2018).
- 340 15. Xu, C., McDowell, N. G., Sevanto, S. & Fisher, R. A. Our limited ability to predict
341 vegetation dynamics under water stress. *New Phytol.* **200**, 298–300 (2013).
- 342 16. Fatichi, S., Pappas, C. & Ivanov, V. Y. Modeling plant-water interactions: an
343 ecohydrological overview from the cell to the global scale. *Wiley Interdiscip. Rev.*
344 *Water* **3**, 327–368 (2016).

- 345 17. Jolly, W. M., Dobbertin, M., Zimmermann, N. E. & Reichstein, M. Divergent
346 vegetation growth responses to the 2003 heat wave in the Swiss Alps. *Geophys. Res.*
347 *Lett.* **32**, 1–4 (2005).
- 348 18. Rigling, A. *et al.* Driving factors of a vegetation shift from Scots pine to pubescent oak
349 in dry Alpine forests. *Glob. Chang. Biol.* **19**, 229–240 (2013).
- 350 19. Seneviratne, S. I. *et al.* Swiss prealpine Rietholzbach research catchment and lysimeter:
351 32 year time series and 2003 drought event. *Water Resour. Res.* **48**, 1–20 (2012).
- 352 20. Laghari, A. N., Vanham, D. & Rauch, W. To what extent does climate change result in
353 a shift in Alpine hydrology? A case study in the Austrian Alps. *Hydrol. Sci. J.* **57**, 103–
354 117 (2012).
- 355 21. Mastrotheodoros, T. *et al.* Ecohydrological dynamics in the Alps: Insights from a
356 modelling analysis of the spatial variability. *Ecohydrology* **12**, 1–18 (2019).
- 357 22. Fatichi, S. & Pappas, C. Constrained variability of modeled T:ET ratio across biomes.
358 *Geophys. Res. Lett.* **44**, 6795–6803 (2017).
- 359 23. Teuling, A. J. *et al.* Contrasting response of European forest and grassland energy
360 exchange to heatwaves. *Nat. Geosci.* **3**, 722–727 (2010).
- 361 24. Lin, Y. S. *et al.* Optimal stomatal behaviour around the world. *Nat. Clim. Chang.* **5**,
362 459–464 (2015).
- 363 25. Clark, D. B. *et al.* Model Development The Joint UK Land Environment Simulator (
364 JULES), model description – Part 2 : Carbon fluxes and vegetation dynamics. 701–722
365 (2011). doi:10.5194/gmd-4-701-2011
- 366 26. Sheffield, J., Wood, E. F. & Roderick, M. L. Little change in global drought over the
367 past 60 years. *Nature* **491**, 435–438 (2012).

- 368 27. Wood, E. F. *et al.* Hyperresolution global land surface modeling: Meeting a grand
369 challenge for monitoring Earth’s terrestrial water. *Water Resour. Res.* **47**, 1–10 (2011).
- 370 28. Bierkens, M. F. P. *et al.* Hyper-resolution global hydrological modelling: what is next?
371 *Hydrol. Process.* **29**, 310–320 (2015).
- 372 29. Ivanov, V. Y., Bras, R. L. & Vivoni, E. R. Vegetation-hydrology dynamics in complex
373 terrain of semiarid areas: 2. Energy-water controls of vegetation spatiotemporal
374 dynamics and topographic niches of favorability. *Water Resour. Res.* **44**, 1–20 (2008).
- 375 30. Teuling, A. J. *et al.* Evapotranspiration amplifies European summer drought. *Geophys.*
376 *Res. Lett.* **40**, 2071–2075 (2013).
- 377 31. Manoli, G., Ivanov, V. Y. & Fatichi, S. Dry-Season Greening and Water Stress in
378 Amazonia: The Role of Modeling Leaf Phenology. *J. Geophys. Res. Biogeosciences*
379 **123**, 1909–1926 (2018).
- 380 32. Fatichi, S., Zeeman, M. J., Fuhrer, J. & Burlando, P. Ecohydrological effects of
381 management on subalpine grasslands: From local to catchment scale. *Water Resour.*
382 *Res.* **50**, 148–164 (2014).
- 383 33. Le Quéré, C. *et al.* Global carbon budget 2014. *Earth Syst. Sci. Data* **7**, 47–85 (2015).
- 384 34. Sitch, S. *et al.* Recent trends and drivers of regional sources and sinks of carbon
385 dioxide. *Biogeosciences* **12**, 653–679 (2015).
- 386 35. Jung, M. *et al.* The FLUXCOM ensemble of global land-atmosphere energy fluxes.
387 *Sci. Data* **6**, 74 (2019).
- 388 36. Dee, D. P. *et al.* The ERA-Interim reanalysis: Configuration and performance of the
389 data assimilation system. *Q. J. R. Meteorol. Soc.* **137**, 553–597 (2011).
- 390 37. Chiesa, D. D. *et al.* Modelling changes in grassland hydrological cycling along an

- 391 elevational gradient in the Alps. *Ecohydrology* **7**, 1453–1473 (2014).
- 392 38. Goulden, M. L. & Bales, R. C. Mountain runoff vulnerability to increased
393 evapotranspiration with vegetation expansion. *Proc. Natl. Acad. Sci.* **111**, 14071–
394 14075 (2014).
- 395 39. Gobiet, A. *et al.* 21st century climate change in the European Alps-A review. *Sci. Total*
396 *Environ.* **493**, 1138–1151 (2014).
- 397 40. Gilbert, J. M. & Maxwell, R. M. Contrasting warming and drought in snowmelt-
398 dominated agricultural basins: Revealing the role of elevation gradients in regional
399 response to temperature change. *Environ. Res. Lett.* **13**, (2018).
- 400 41. Zappa, M. & Kan, C. Extreme heat and runoff extremes in the Swiss Alps. *Nat.*
401 *Hazards Earth Syst. Sci.* **7**, 375–389 (2007).
- 402 42. Leibundgut, C. & Kohn, I. European traditional irrigation in transition part II:
403 Traditional irrigation in our time - decline, rediscovery and restoration perspectives.
404 *Irrig. Drain.* **63**, 294–314 (2014).
- 405 43. Dobbertin, M. *et al.* The decline of *Pinus sylvestris* L. forests in the Swiss Rhone
406 valley - A result of drought stress? *Phyt. - Ann. Rei Bot.* **45**, 153–156 (2005).
- 407 44. Maxwell, R. M. & Kollet, S. J. Interdependence of groundwater dynamics and land-
408 energy feedbacks under climate change. *Nat. Geosci.* **1**, 665–669 (2008).
- 409 45. Fan, Y. Groundwater in the Earth’s critical zone: Relevance to large-scale patterns and
410 processes. *Water Resour. Res.* **51**, 3052–3069 (2015).
- 411 46. Bales, R. C. *et al.* Mechanisms controlling the impact of multi-year drought on
412 mountain hydrology. *Sci. Rep.* **8**, 1–8 (2018).
- 413 47. Casty, C., Wanner, H., Luterbacher, J., Esper, J. & Böhm, R. Temperature and

- 414 precipitation variability in the European Alps since 1500. *Int. J. Climatol.* **25**, 1855–
415 1880 (2005).
- 416 48. Pal, J. S., Giorgi, F. & Bi, X. Consistency of recent European summer precipitation
417 trends and extremes with future regional climate projections. *Geophys. Res. Lett.* **31**,
418 20–23 (2004).
- 419 49. Mastrotheodoros, T. *et al.* Linking plant functional trait plasticity and the large increase
420 in forest water use efficiency. *J. Geophys. Res. Biogeosciences* **122**, 2393–2408 (2017).
- 421 50. Dupire, S., Curt, T. & Bigot, S. Spatio-temporal trends in fire weather in the French
422 Alps. *Sci. Total Environ.* **595**, 801–817 (2017).
- 423 51. Maxwell, R. M. & Condon, L. E. Connections between groundwater flow and
424 transpiration partitioning. *Science*. **353**, 377–380 (2016).
- 425 52. Barnett, T. P., Adam, J. C. & Lettenmaier, D. P. Potential impacts of a warming climate
426 on water availability in snow-dominated regions. *Nature* **438**, 303–9 (2005).

427 **Acknowledgements**

428 T.M. wants to thank Daniela Anghileri for her support regarding the computational part of this
429 study. Numerical simulations were performed on the ETH Euler cluster. T.M. and S.F. thank
430 the support of the Stavros Niarchos Foundation and the ETH Zurich Foundation (Grant ETH-
431 29 14-2). C.P. acknowledges the support of the Swiss National Science Foundation (SNSF),
432 the Stavros Niarchos Foundation and the ETH Zurich Foundation (Grants P2EZP2_162293 and
433 P300P2_174477). G.M. was supported by the “The Branco Weiss Fellowship - Society in
434 Science” administered by ETH Zurich. We thank the participants of the TRENDY project,
435 namely, P. Levy (Hyland), S. Sitch and C. Huntingford (JULES/ TRIFFID), B. Poulter (LPJ),
436 A. Ahlström (LPJ-GUESS), S. Levis (NCAR-CLM4), N. Viovy, S. Zaehle (OCN), M. Lomas
437 (SDGVM) and N. Zeng (VEGAS), who made their simulations results (TRENDY v.1,

438 experiment S2) freely available. This work used eddy covariance data acquired and shared by
439 the FLUXNET community, including CarboEuropeIP and CarboItaly. The FLUXNET eddy
440 covariance data processing and harmonization was carried out by the European Fluxes Database
441 Cluster, AmeriFlux Management Project, and Fluxdata project of FLUXNET, with the support
442 of CDIAC and ICOS Ecosystem Thematic Center, and the OzFlux, ChinaFlux and AsiaFlux
443 offices.

444 **Author contributions**

445 S.F. and T.M. designed the study, J.P., R.R., B.S. and M.B. contributed with data. P.H.
446 contributed to the optimization of the parallel simulations. T.M. performed the simulations and
447 the analyses. T.M. designed the figures with contributions from G.M and S.F. The results were
448 synthesized by T.M., S.F., C.P. and P.M. T.M. and S. F. wrote the manuscript with contributions
449 from all other authors.

450 **Correspondence and requests for materials** should be addressed to Simone Fatichi.

451 **Competing financial interests**

452 The authors declare no competing financial interests

Methods

The Tethys-Chloris model (T&C). We performed simulations with the distributed
ecohydrological model Tethys-Chloris (T&C). T&C simulates the coupled dynamics of energy,
water, and vegetation and has been successfully applied to a very large spectrum of ecosystems
and environmental conditions as summarized elsewhere^{31,32,49,53–55}.

The model simulates the energy, water, and carbon exchanges between the land surface and the
atmospheric surface layer accounting for aerodynamic, undercanopy, and leaf boundary layer
resistances, as well as for stomatal and soil resistances⁵⁶. In each simulated grid cell, vegetation

can occupy two vertical layers to mimic the coexistence of trees and bushes/grasses. Horizontal composition of vegetation is also possible since each element can account for multiple species or plant functional types^{54,57}. Dynamics of water content in the soil profile are solved using a quasi 3-D approach: the one-dimensional (1-D) Richards equation is used for vertical flow and the kinematic wave equation is used for lateral subsurface flow. Saturated and unsaturated parts of the soil column are explicitly identified. Surface overland flow and channel flow are also solved by the kinematic equation. Snowpack dynamics are computed using the energy balance: snow can be intercepted by the vegetation or fall to the ground, where it accumulates and successively melts. Runoff generation occurs via saturation excess and infiltration excess mechanisms and depends on lateral moisture fluxes in the unsaturated and saturated zones as well as on overland flow⁵⁸. Soil water content changes, infiltration, and runoff production are estimated with an adaptive time step based on the stability of Richards equation solution with the method of lines⁵⁹. Overland and channel flow routing are computed with an adaptive time step that satisfies the Courant-Friedrichs-Lewy condition (seconds to 5 min). The version of T&C used in this study does not include soil freezing and thawing processes and the water present in soil pores is always considered to be in a liquid state. The soil biogeochemistry module is not activated, thus there is no modeling of soil carbon and ground evaporation does not occur from litter, but only from soil.

Photosynthesis is simulated using the Farquhar biochemical model⁶⁰ adapted with subsequent modifications and temperature dependence of biochemical parameters. The model follows the two big leaves scheme, where sunlit and shaded leaves are treated separately for estimating net assimilation and stomatal resistance⁶¹. Leaf maintenance respiration is assumed equal to the leaf dark respiration and acclimation effects are not accounted for. For upscaling photosynthesis from leaf to plant scale, photosynthetic capacity is assumed to decay exponentially with canopy depth⁶². The stomatal conductance parameterization accounts for net assimilation rate, leaf internal CO₂ concentration and vapor pressure deficit following the Leuning model⁶³. The

dynamics of seven carbon pools are explicitly simulated in the model and include leaves, living sapwood, fine roots, carbohydrate reserve (non-structural carbohydrates), reproductive tissues (fruits and flowers), standing dead leaves and heartwood/dead sapwood. The carbon assimilated through photosynthetic activity is used for maintenance and growth respiration, otherwise it is allocated to one of the first five pools. The different pools are undergoing tissue turnover in function of tissue longevity and environmental stresses, i.e., drought and low temperatures. Carbon allocation is a dynamic process that accounts for resource availability (light and water) and allometric constraints⁶⁴, e.g., a minimum ratio of fine root to foliage carbon; and an upper limit for the storage of carbohydrate reserves⁶⁵. Carbon allocated to reserves can be subsequently translocated to favor leaf and fine root expansion at the onset of the growing season or after severe disturbances⁵⁴. Phenology for extratropical species is simulated considering four states⁶⁶: dormant, maximum growth, normal growth, and senescence. Patterns of plant allocation are influenced by the phenological phase⁶⁷. Transition between phenological phases are prognostic in the model and controlled by soil temperature, soil moisture and photoperiod. The model assumes that vegetation is in a mature phase and in equilibrium with its nutritional environment. Further details of model computational set-up, structure, and description of process parameterizations are presented elsewhere⁵⁴. Vegetation dynamics are solved at the daily time scale, energy fluxes at the hourly time scale.

Domain setup. The land surface for the simulations was defined based on the 90 m resolution STRM digital elevation model (DEM) following the definition of the Alpine Region of the European Soil Data Center (ESDAC, project name: Eco-pedological Map for the Alpine Territory, https://esdac.jrc.ec.europa.eu/projects/Alpsis/Ecalp_data.html)⁶⁸. This includes the entire alpine arch from the French coast in the southwest to the Austrian lowlands in the north east and from the Swiss plateau on the northwest to the Slovenian coast in the southeast. This DEM was resampled to the final resolution of the simulations (250 m). The flow matrix used by T&C was calculated based on the D-infinity method⁶⁹. To compute the stream network, we

set the threshold of upslope area above which a pixel is considered to belong to the channel network to 40 km². Therefore, small streams are not explicitly resolved.

Soil texture data were derived from the SoilGrids product at 250 m resolution⁷⁰. We obtained sand, silt, clay, and organic content for seven different depths (at the ground surface and at 0.05, 0.10, 0.30, and 1 m below the ground surface). Average soil properties were then computed for each grid cell considering the vertical discretization of the soil and assuming linearity. A uniform soil depth equal to 1 m was used for the entire area. This is a simplification of the real system, since variability in soil depth can affect runoff generation mechanisms, but our choice is considered reasonable for large parts of the pan-Alpine domain, in the absence of consistent information about soil depth. We used land cover data from Corine⁷¹, using seven different classes to summarize the information provided in the dataset. The classification we used is shown in the Supplementary Table 1. Glaciers were initialized with a 50 m depth. Lake depth was not explicitly simulated. In other words, we assured that there was always water available for evaporation for every lake-pixel (by starting with a very large lake depth), but the rivers flowing into the lakes did not interact with this “lake” water.

For reasons of computational efficiency, the domain was divided into eight subdomains, roughly equal in size, following the divides (Supplementary Fig. 1); these subdomains were run independently of each other. No lateral exchange between the eight domains is anyhow expected because of the kinematic wave assumption (this is indeed a simplification because groundwater rarely follows the surface topography). Further, each domain was run in parallel with 24-36 cores on the Euler cluster at ETH (<https://scicomp.ethz.ch/wiki/Euler>). For this parallelization, each subdomain was automatically split into smaller subdomains according to the number of cores. The vegetation and hydrology modules of T&C were run independently for each of these small subdomains. Lateral exchanges were then performed serially by a master process (e.g. flow routing, avalanches) at each time step. Each completed job on the cluster

automatically saved a report where the CPU time was stated. The total computational demand for the three-year simulation for each of these sub-regions was roughly 76,000 CPU hours (information about the cores available can be found at the following link: <https://scicomp.ethz.ch/wiki/Euler#> - Euler III and IV). We initialized the carbon pools in the model using values for the different vegetation types corresponding to mid-summer conditions (e.g., leaf biomass for evergreen forests was set to 270 gC m⁻²) derived from decadal simulations at the plot-scale run for each vegetation type³². Furthermore, we started from a relatively dry domain and we run the distributed model twice with observed meteorological forcing for the period September-October 2000. The final values of all state variables (e.g., soil moisture, channel water storage, surface temperature, snow water equivalent, carbon pools) were considered the initial conditions for the numerical experiment presented here, which started on the 1st of November 2000.

Meteorological input. Obtaining hourly meteorological fields at the desired 250 m resolution for an area spanning 257,000 km² and seven countries (Austria, France, Germany, Italy, Lichtenstein, Slovenia and Switzerland) required the combination of distributed products and station data as described below.

Precipitation

We used the daily Alpine precipitation grid dataset⁷², which has a grid spacing of 5 km and daily time resolution. To obtain the required model input at hourly temporal resolution and 250 m spatial resolution we used ground observations from 111 stations in Austria, France, Italy, Liechtenstein, and Switzerland (Supplementary Table 2). The computation of hourly precipitation was performed as the simulation was running, because of the impracticability of saving hourly precipitation fields for the entire period. Prior to the simulation, Thiessen polygons were defined based on the 111 stations, so that each pixel of the simulated domain was assigned to a single station. Each pixel was also assigned to the corresponding grid of the

Alpine precipitation. Then, during the simulation, in hour 1:00 of each day, the daily sum of the distributed product was disaggregated to hourly precipitation proportionally to the hourly measurements of the corresponding station but preserving the amount of precipitation based on the gridded daily product. In case that the corresponding station recorded no precipitation in a day in which the gridded product did record some, the daily gridded sum was assigned uniformly to the hours 18:00-22:00. These cases were however rare and mostly occurring at low rainfall intensities.

Air Temperature

For air temperature we used the ECMWF ERA INTERIM product³⁶ using time 00:00:00 and 12:00:00 and step 3/6/9/12. This corresponds to the forecasts that are issued twice a day for the next 12 hours with 3-hourly time step. Thus, we obtained 3-hourly data. We compared this to the product that includes analyses of four times a day and only 3-hourly forecasts and the difference was minimal. We used the finest resolution available (0.125°, data downloaded from <https://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/> on 23/2/2018). We first assigned the ERA INTERIM temperature to all the T&C cells within the corresponding ERA INTERIM grid and then we readjusted the temperature for each T&C grid cell to account for the elevation variability within each grid cell of the ERA INTERIM. For this readjustment, we used a single, temporally dynamic lapse-rate over each subdomain. The lapse-rate was computed every three hours by using all the ERA INTERIM grid cells within the subdomain. The two hours between the three-hourly ERA INTERIM data were computed using linear interpolation.

Wind Speed

For wind speed, we used the corresponding variable of ERA INTERIM, as described for temperature above. A constant lapse rate was used for downscaling from ERA INTERIM to T&C grid cells throughout the entire simulation based on observations in Switzerland (0.48 m

$\text{s}^{-1}/100 \text{ m}$). A lower limit to wind speed was also imposed to avoid numerical instabilities in the computation of turbulent fluxes at the surface (0.01 m s^{-1}).

Atmospheric Pressure

For surface pressure, we used the corresponding ERA INTERIM variable, as described for temperature and wind speed above. Between the three-hourly data, we performed linear interpolation and applied a specific exponential correction to account for the variations of atmospheric pressure with elevation within each ERA INTERIM grid cell.

Relative Humidity

To compute the relative humidity, which is an additional necessary model input, we used dew temperature data from the ERA INTERIM product, and extrapolated it in space and time as described above for air temperature. Then, we combined hourly air temperature and dew point temperature to compute the hourly values of relative humidity for each cell in the T&C domain.

Solar Radiation

For computing distributed hourly radiation components for each pixel, we used measurements from 90 stations across the entire domain (stations with an asterisk (*) in the Supplementary Table 1). We used the inverse distance weight to compute radiation in each T&C grid cell from the meteorological stations, which means that more than one station was taken into account for each T&C grid cell. Only direct radiation was corrected based on constant lapse rates that varied according to the radiation wavelength (lapse rate $0.0015 \text{ W m}^{-2} \text{ m}^{-1}$ for direct shortwave radiation between 0.29 and $0.70 \mu\text{m}$, $0.0027 \text{ W m}^{-2} \text{ m}^{-1}$ for $0.70 - 4.0 \mu\text{m}$ and $0.0014 \text{ W m}^{-2} \text{ m}^{-1}$ for direct photosynthetically active radiation). Diffuse shortwave radiation was not rescaled with elevation. Subsequently, T&C modified the incoming radiation to account for local and remote topographic effects, (i.e., shading and backscatter from nearby terrain, slope and aspects

of the grid cell). Sky view factor and terrain configuration factor scaling between 0 and 1 are used for this purpose⁷³.

Longwave Radiation

To compute the incoming longwave radiation we followed an empirical formula for the clear sky emissivity⁷⁴, based on air temperature, vapor pressure (computed from relative humidity) and cloud cover. The latter was estimated for each of the stations with solar radiation measurements by comparing observed radiation and clear-sky radiation, simulated by a weather generator⁷⁵.

Space-for-time substitution. We estimated ET for a +3 °C change in mean annual temperature using a space-for-time substitution⁷⁶. For each vegetation type (i.e. grassland, evergreen, deciduous, and mixed forest) we fit linear (for E) and non-linear (for T) models to describe the current spatial relationship between mean annual evapotranspiration and mean annual temperature. We performed separate regressions for evaporation and transpiration against temperature for each subdomain (Supplementary Fig. 1). Once the new reference values of ET were computed for each cell at the higher temperature, we also added the residuals of the regressions to preserve the spatial variability and the non-temperature-driven effects.

Model validation. We validated the model against remote sensing MODIS snow cover estimates (<https://nsidc.org/data/mod10a2>)⁷⁷. Using the distributed dataset with temporal resolution of 8 days, we computed the time fraction with snow cover over the entire domain and compared it with the simulation results (Supplementary Fig. 3). The model tends to underestimate snow cover at lower elevations, and especially on the north facing slopes of the inner Alpine valleys. It also overestimates snow cover in the very wet pre-Alpine mountains, which might be partly an artefact of the model spin-up (precipitation in October 2000 was very high). Overall, results are satisfactory, considering that the model was run with parameters derived from previous plot-scale studies^{22,32} and MODIS may tend to overestimate snow cover

at low elevations, because it assigns to all eight days the largest snow cover recorded in that period. In fact, comparison of simulated snow cover with station observations (Extended Data Figure 1) suggests that the bias in snow cover duration only sporadically exceeds 30 days (Supplementary Fig. 3) and much larger values are likely to be a remote sensing artefact.

We further validated our analysis by comparing T&C output with gross primary production (GPP) and latent heat (LE) observations in 10 FLUXNET sites in the Alps or in the nearby areas (Supplementary Figs 4 and 5, and Supplementary Tables 5 and 6). Since the periods with available data vary between these sites and most often do not cover the 2001-2003 period, we performed an independent plot-scale simulation with T&C with hourly time step for each site. For these simulations, we used local meteorological forcing (downloaded from the FLUXNET database, see Supplementary Tables 5 and 6); these data were gap-filled whenever needed with linear regression when data gaps lasted less than a few hours or using the mean of the corresponding hour and day of the year over the entire period with available data. For all the sites we applied the same parameterization (according to the vegetation type) of the pan-Alpine setup.

We further compared the model with ET and GPP estimates from three distributed products: FLUXCOM³⁵ for ET (0.0833° spatial resolution) and GPP (0.5° spatial resolution), TRENDY^{33,34} for GPP (spatial resolution 1°, simulations S2: CO₂ and climate (time-invariant present-day land use mask, models used: CLM4CN, HYLAND, LPJ_GUESS, LPJ, OCN, ORCHIDEE, SDGVM, TRIFFID VEGAS-2.1), and ERA Interim³⁶ for ET (derived from monthly estimates of latent heat at 0.125° spatial resolution, Extended Data Figure 3 and Supplementary Figs 6). The comparison shows that all products provide estimates close to the T&C average, but they all underestimate spatial variability due to their coarse spatial resolution. For instance, a single grid cell corresponding to the spatial resolution of TRENDY multimodel ensemble, includes areas with annual precipitation ranging from less than 700 mm to more than

3300 mm, while the entire domain is represented by only 52 grid cells. All of these distributed products cannot represent the complex topography of the Alps, which implies that other methods are needed for a robust quantification of green and blue water fluxes in mountainous regions as shown here.

We validated the model against daily runoff measurements in 381 stations (Supplementary Table 3) and daily snow depth measurements in 720 stations in Switzerland, Austria, and Italy (Supplementary Table 4). The results of the validation against station measurements are summarized in Extended Data Figure 1. Since T&C simulates only natural flows, i.e., it does not account for human regulation, which is quite important in the Alps, we removed heavily regulated catchments. To do so, in the absence of a pan-Alpine database indicating which station is affected by regulation, dams, or water withdrawals, we visually inspected all the observed hydrographs to flag each station as regulated or natural. We found that the model shows a considerably higher performance when excluding the regulated catchments (Extended Data Figure 2). The correlation between simulated and observed runoff time series is higher for the natural catchments, compared with the regulated ones (R^2 equals 0.69 and 0.47, respectively). Mean bias is less than 0.01 mm hr^{-1} for both groups of catchments, but the sign is opposite. As expected, T&C tends to overestimate runoff in heavily regulated catchments due to water abstractions, which were not simulated. Since in our analysis we used P-ET as a proxy for runoff, the inset of Extended Data Figure 2 compares area-averaged P-ET with measured runoff and shows that this assumption is largely valid, especially for the natural and non-glacierized catchments. For glacierized catchments, icemelt should be added to P-ET to obtain a reliable estimate of runoff.

Data availability. Data sources are summarized in the Supplementary Table 6. The DEM, land cover product and MODIS snow cover product used for this study are publicly available (https://esdac.jrc.ec.europa.eu/projects/Alpsis/Ecalp_data.html,

<https://www.eea.europa.eu/data-and-maps/data/clc-2006-raster-4>, and <https://doi.org/10.5067/MODIS/MOD10A2.006>, respectively). FLUXNET data for running and validating T&C plot-scale simulations were downloaded from <https://fluxnet.fluxdata.org/>. FLUXCOM GPP and ET estimates were downloaded from <http://www.fluxcom.org/>, ERA Interim latent heat estimates were downloaded from <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim> and TRENDY estimates of GPP were downloaded from <http://www-lscedods.cea.fr/invsat/RECCAP/>. Hydrological and meteorological/snow depth data from Switzerland are publicly available (<https://www.hydrodaten.admin.ch/>, and <https://gate.meteoswiss.ch/idaweb/more.do>, respectively). Daily data from the Austrian network are provided by the Austrian hydrological service (<https://ehyd.gv.at/>), and are publicly available. Hydrological and meteorological data for the French stations are publicly available (<http://www.hydro.eaufrance.fr/>, and <https://donneespubliques.meteofrance.fr/>, respectively). Hydrological data for the Slovenian stations are publicly available (<http://www.arso.gov.si/vode/podatki/>). Hydrological, meteorological and snow depth data for the Piemonte region and the Friuli-Venezia-Giulia region (Italy) are publicly available (<http://www.arpa.piemonte.it/> and <http://www.meteo.fvg.it/archivio.php?ln=&p=dati>, respectively). Hydrological, meteorological and snow depth data for the Trentino-Alto Adige region (Italy) are provided by the University of Trento and are available upon request.

Code availability. Tethys-Chloris source code used for these simulations (11/2015 version) is available at <https://hyd.ifu.ethz.ch/forschung/models.html>, saved simulations results and the analysis scripts are available upon request to the corresponding author (S.F.). The software used to generate all the results is MATLAB 2016b®.

References

53. Fatichi, S., Leuzinger, S. & Körner, C. Moving beyond photosynthesis: From carbon

- source to sink-driven vegetation modeling. *New Phytol.* **201**, 1086–1095 (2014).
54. Fatichi, S., Ivanov, V. Y. & Caporali, E. A mechanistic ecohydrological model to investigate complex interactions in cold and warm water-controlled environments: 1. Theoretical framework and plot-scale analysis. *J. Adv. Model. Earth Syst.* **4**, M05002 (2012).
55. Wu, D. *et al.* Asymmetric responses of primary productivity to altered precipitation simulated by ecosystem models across three long-term grassland sites. *Biogeosciences* **15**, 3421–3437 (2018).
56. Sellers, P. J. *et al.* Modeling the exchanges of energy, water, and carbon between continents and the atmosphere. *Science.* **275**, 502–509 (1997).
57. Fatichi, S. & Leuzinger, S. Reconciling observations with modeling: The fate of water and carbon allocation in a mature deciduous forest exposed to elevated CO₂. *Agric. For. Meteorol.* **174–175**, 144–157 (2013).
58. Loague, K., Heppner, C. S., Ebel, B. A. & VanderKwaak, J. E. The quixotic search for a comprehensive understanding of hydrologic response at the surface: Horton, Dunne, Dunton, and the role of concept-development simulation. *Hydrol. Process.* **24**, 2499–2505 (2010).
59. Lee, H. S., Matthews, C. J., Braddock, R. D., Sander, G. C. & Gandola, F. A MATLAB method of lines template for transport equations. *Environ. Model. Softw.* **19**, 603–614 (2004).
60. Collatz, G. J., Ribas-carbo, M. & Berry, J. A. Coupled Photosynthesis-Stomatal Conductance Model for Leaves of C₄ Plants. *Aust. J. Plant Physiol.* **19**, 519 (1992).
61. Dai, Y., Dickinson, R. E. & Wang, Y. P. A two-big-leaf model for canopy temperature, photosynthesis, and stomatal conductance. *J. Clim.* **17**, 2281–2299 (2004).

62. Bonan, G. B. *et al.* Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data. *J. Geophys. Res.* **116**, 1–22 (2011).
63. Leuning, R. Modelling Stomatal Behaviour and Photosynthesis of *Eucalyptus grandis*. *Funct. Plant Biol.* **17**, 159 (2006).
64. Krinner, G. *et al.* A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochem. Cycles* **19**, 1–33 (2005).
65. Kozlowski, T. T. & Pallardy, S. G. *Physiology of wood plants*. (Academic Press, 1997).
66. Arora, V. K. & Boer, G. J. A parameterization of leaf phenology for the terrestrial ecosystem component of climate models. *Glob. Chang. Biol.* **11**, 39–59 (2005).
67. Farquhar, G. D., von Caemmerer, S. & Berry, J. A. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* **149**, 78–90 (1980).
68. Panagos, P., Van Liedekerke, M., Jones, A. & Montanarella, L. European Soil Data Centre: Response to European policy support and public data requirements. *Land use policy* **29**, 329–338 (2012).
69. Tarboton, D. G. A new method for the determination of flow directions and upslope areas in grid digital elevation models. *Water Resour. Res.* **33**, 309–319 (1997).
70. Hengl, T. *et al.* SoilGrids1km - Global soil information based on automated mapping. *PLoS One* **9**, (2014).
71. European Environment Agency. *CORINE Land Cover 2006*. (2006).
72. Isotta, F. A. *et al.* The climate of daily precipitation in the Alps: Development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data. *Int. J.*

- Climatol.* **34**, 1657–1675 (2014).
73. Fatichi, S., Ivanov, V. Y. & Caporali, E. A mechanistic ecohydrological model to investigate complex interactions in cold and warm water-controlled environments: 1. Theoretical framework and plot-scale analysis. *J. Adv. Model. Earth Syst.* **4**, 1–31 (2012).
74. Dilley, A. C. & O'Brien, D. M. Estimating downward clear sky long-wave irradiance at the surface from screen temperature and precipitable water. *Q. J. R. Meteorol. Soc.* **124**, 1391–1401 (1998).
75. Fatichi, S., Ivanov, V. Y. & Caporali, E. Simulation of future climate scenarios with a weather generator. *Adv. Water Resour.* **34**, 448–467 (2011).
76. Dunne, J. A., Saleska, S. R., Fischer, M. L. & Harte, J. Integrating experimental and gradient methods in ecological climate change research. *Ecology* **85**, 904–916 (2004).
77. Hall, D. K. & Riggs, G. A. *MODIS/Terra Snow Cover 8-Day L3 Global 500m Grid, Version 6.MOD10A2*. (2016). doi:<https://doi.org/10.5067/MODIS/MOD10A2.006>