

More on complexity of operators in quantum field theory

Run-Qiu Yang,^a Yu-Sen An,^{b,c} Chao Niu,^d Cheng-Yong Zhang^e and Keun-Young Kim^f

^aQuantum Universe Center, Korea Institute for Advanced Study,
Seoul 130-722, Korea

^bKey Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Science, Beijing 100190, China

^cSchool of physical Science, University of Chinese Academy of Science,
Beijing 100049, China

^dDepartment of Physics and Siyuan Laboratory, Jinan University,
Guangzhou 510632, China

^eDepartment of Physics and Center for Field Theory and Particle Physics, Fudan University,
Shanghai 200433, China

^fSchool of Physics and Chemistry, Gwangju Institute of Science and Technology,
Gwangju 61005, Korea

E-mail: aqiu@kias.re.kr, anyusen@itp.ac.cn, chaoniu09@gmail.com,
zhangchengyong@fudan.edu.cn, fortoe@gist.ac.kr

ABSTRACT: Recently it has been shown that the complexity of $SU(n)$ operator is determined by the geodesic length in a bi-invariant Finsler geometry, which is constrained by some symmetries of quantum field theory. It is based on three axioms and one assumption regarding the complexity in continuous systems. By relaxing one axiom and an assumption, we find that the complexity formula is naturally generalized to the Schatten p -norm type. We also clarify the relation between our complexity and other works. First, we show that our results in a bi-invariant geometry are consistent with the ones in a right-invariant geometry such as k -local geometry. Here, a careful analysis of the sectional curvature is crucial. Second, we show that our complexity can concretely realize the conjectured pattern of the time-evolution of the complexity: the linear growth up to saturation time. The saturation time can be estimated by the relation between the topology and curvature of $SU(n)$ groups.

KEYWORDS: Gauge-gravity correspondence, Holography and condensed matter physics (AdS/CMT)

ARXIV EPRINT: [1809.06678](https://arxiv.org/abs/1809.06678)

Contents

1	Introduction	1
2	Review on the complexity in quantum field theory	4
3	Generalization of axioms	8
3.1	Generalized parallel decomposition rule	8
3.2	Generalized smoothness	8
3.3	Emergence of the Finsler metric	9
4	Constraints on the Finsler metric	9
4.1	Adjoint invariance	10
4.1.1	Adjoint invariance from the independence of left/right generators	10
4.1.2	Adjoint invariance from symmetric transformations	12
4.2	Reversibility from the CPT symmetry	14
5	Finsler metric and complexity of $SU(n)$ operators	15
5.1	Finsler metric of $SU(n)$ operators	15
5.2	Complexity of $SU(n)$ operators	17
6	Properties of operator complexity	18
6.1	Geodesic deviation and chaos	18
6.1.1	Comparison with “ k -local” metrics	20
6.2	Linear growth and quantum recurrence	21
6.3	Complexity of precursors	23
7	Conclusion	25
A	Relationship between the time order and left/right orders	27
B	Two kinds of special symmetric transformations	28
B.1	Adding a divergent term	28
B.2	Canonical transformation	29
C	Explicit form of $\mathfrak{g}_s^{(\infty)}$	32
D	“Local” and “nonlocal” Hamiltonians	33
E	Jacobi field and sectional curvature	34

1 Introduction

Recently, many concepts in quantum information theory have been applied to investigate gravity and quantum field theory (QFT). In particular, a concept named “complexity”, which comes from the quantum circuit complexity in quantum information theory [1], was introduced to study some properties of black holes, in particular for the physics inside the black hole horizon. From the perspective of holographic principle, it is expected to have a physical meaning of “complexity” in QFT.

In gravity side, the first motivation to consider the complexity was to investigate the firewalls in the black hole horizon [2] and the growth rate of the Einstein-Rosen bridge in AdS black holes [3, 4]. There are two widely studied conjectures to compute the complexity for some particular quantum states which are dual to boundary time slices of the asymptotic AdS black hole. One states that the complexity is proportional to the maximum volume of some time-like hypersurface (CV conjecture) [3, 5, 6] and the other one states that the complexity is proportional to the on-shell action in a particular spacetime region so called the wheeler dewitt (WdW) patch (CA conjecture) [7, 8]. Along these lines, there have been a lot of developments and we refer to some of them, [9–22], for examples. There are also other interesting conjectures proposed in gravity, for example, subregion complexity [23–27] and thermodynamic volume [28].

Because the holographic duality connects the gravity and QFT, it is important and timely to develop the theory of complexity in QFT. Compared with much progress for the complexity in gravity side, the precise meaning or a proper definition of the complexity in QFT side, is still absent. However, recently some interesting ideas have been proposed and promising preliminary results have been obtained in [29–49].

To make a progress towards the complexity in QFT we may rely on the intuitions from the circuit complexity in computer science, where the concept of the complexity is well developed. However, in the quantum circuits, the complexity is defined in *discrete* and *finite* Hilbert spaces. The intuition based on the complexity in terms of the *number* of quantum gates may be ideal for quantum circuits but may not be enough for QFT, of which operations are *continuous*. The first clue to define the complexity in *continuous* systems appeared in Nielsen and collaborators’ works [50–52]. Here, as a continuum approximation of the circuit complexity, a kind of “complexity geometry” was introduced and the complexity is identified with the geodesic length in the geometry.

Inspired by this idea, a few proposals to define the complexity in QFT appeared in [16, 29–32, 48]. Ref. [29] first introduced Nielsen and collaborators’ idea to the study of complexity in QFT and [30, 31] argued the conditions for the complexity geometry to satisfy. Ref. [32] obtained the complexity of the ground state of a free scalar field theory by computing the length of geodesic in a complexity geometry. See also [33–38] for related developments. In these works it was shown that the UV divergence structures are consistent with the holographic results [12–14]. Refs. [16, 48] computed the time evolution of the complexity between thermofield double states in a complexity geometry and made a comparison with the holographic results. In [16] another proposal for QFT complexity based on the Fubini-Study metric [39] was also investigated.

However, in all these attempts [16, 29–32, 48] there are ambiguities in choosing complexity geometry. These works mostly focus on the behaviour of geodesics in an *assumed* geometry. For example, the geometry may be chosen to match the holographic results. However, using holographic results as a guide for the complexity geometry in field theory may have the following issues:

1. There are some ambiguities in the definition of holographic conjectures. For example, in both CV and CA conjectures we do not know what the reference states are, and in CA conjecture there is an ambiguity in choosing the parameterization on null boundaries [10].
2. Even after fixing those ambiguities there is still a possibility that the definition of the holographic conjectures is not complete. The concept of the complexity in field theory independent of holography will be useful to identify the holographic complexity and check its consistency in the holographic duality.

Therefore, it will be more satisfactory if we can first *determine* the complexity geometry by some field theory principles.

We addressed this issue in ref. [49] and proposed how to determine the complexity geometry and the complexity of the $SU(n)$ operators. The basic ideas is as follows.

1. Start with three minimal axioms that the complexity in any system should satisfy. These axioms are extracted from the circuit complexity which is a *discrete* system.
2. Add a certain smoothness assumption to deal with the complexity in *continuous* systems.
3. From these considerations (three axioms and one assumption) the Finsler geometry is naturally emerges and the complexity is identified with the length of the geodesic between the identity and the target operator in the Finsler geometry.
4. Because we want to investigate the complexity in QFT it is natural to impose the *symmetry of QFT* such as unitary and CPT invariance on the complexity. By these symmetries, the structure of the Finsler geometry is more constrained. It turns out the constraints enable us to determine the metric of the complexity geometry.
5. Based on these considerations, we finally obtain the complexity of the operator \hat{O} in $SU(n)$ group as follows.

$$\mathcal{C}(\hat{O}) = \lambda \text{Tr} \sqrt{\bar{H} \hat{H}^\dagger}, \quad \forall \bar{H} = \ln \hat{O}. \quad (1.1)$$

In this process, there are two important features that we want to emphasize again. First, the Finsler geometry emerges naturally from three axioms and a smoothness assumption. The Finsler geometry is not an input in our formalism but an output, which is different from other works. Second, we impose the symmetry of QFT to the complexity, so the some of our results may not be compatible with the circuit complexity. Indeed, this is not a shortcoming at all, because we want to study the complexity in QFT not of the quantum circuit.

In this paper we generalize eq. (1.1) by relaxing one axiom (so called ‘parallel decomposition rule’) and a smoothness assumption. We show that the complexity of an operator \hat{O} in $SU(n)$ group is still given by the geodesic between the identity and the operator \hat{O} in a bi-invariant Finsler geometry. By imposing some symmetries of QFT we finally obtain

$$\mathcal{C}(\hat{O}) = \lambda(w) \left\{ \text{Tr} \left[\left(\bar{H} \bar{H}^\dagger \right)^{p/2} \right] \right\}^{1/p}, \quad \forall \bar{H} = \ln \hat{O}. \quad (1.2)$$

There are two changes compared to eq. (1.1). It is generalized to the Schatten p -norm [35, 36, 53] and the overall constant is a function of additional parameter w , which may represent a penalty factor. Because w enters into the overall constant there is no essential effect of the complexity. We will present detailed explanation and motivation to introduce p and w (by relaxing one axiom and the smoothness condition) in the main text.

Another important goal of this paper is to investigate the property of the complexity (1.2) and clarify the relation to other works. It is classified as three sub-goals.

First, our work may look different from refs. [29–31] in the sense that our Finsler geometry is bi-invariant while it is only right-invariant in refs. [29–31]. However, we will show that two results are consistent with each other. The key point of the resolution lies on the careful analysis of the section curvature and its relation to geodesic deviations. In essence, we find that even though our Finsler geometry is bi-invariant, its sub-manifold relevant to geodesic deviation may not be bi-invariant and only right-invariant. It implies that the sub-manifold may have negative curvature in part which makes chaotic behavior of the geodesics possible.

Second, we find that our complexity (1.2) realize the pattern of the time-evolution of the complexity conjectured in [5, 31]. The linear growth in early time is explained by the geodesic generated by a *constant* generator, which is originated from the *bi-invariance* of the Finsler geometry. The complexity reaches its maximum value in the exponential time ($t \sim e^{\mathcal{O}(d)}$), where, d is the size of the classical phase space, due to the relation between the topology and curvature of $SU(n)$ groups. To our knowledge, this is the first concrete realization of the time-evolution of the complexity conjectured in [5, 31].

Third, we have investigated the complexity of the precursor operators and compare with the results in refs. [3, 29]. We find that (i) the complexity of the precursor operator grows linearly at early time, which is similar to the result in ref. [29]. (ii) the complexity of precursors for infinitesimal operators corresponds to the sectional curvature.

This paper is organized as follows. In section 2, we first make a brief review on the principles and symmetries of complexity in QFT proposed by ref. [49]. Here, the complexity in QFT is defined based on some axioms inspired by the circuit complexity and constraints imposed on QFT. The complexity of $SU(n)$ operator is given by the geodesic length in a Finsler geometry. In section 3, we relax some requirements in axioms in section 2 (or ref. [49]) and find a Finsler geometry is still relevant. In section 4, by imposing some symmetries of QFT on the complexity, we constrain the structure of a Finsler geometry and, in section 5, determine the complexity of $SU(n)$ group operator uniquely. In section 6, we make comments on the properties of the complexity such as i) negative sectional curvature and chaos, ii) the complexity growth, saturation, and quantum recurrence iii) complexity of precursors. We conclude in section 7.

2 Review on the complexity in quantum field theory

In this section, we review the complexity proposed in ref. [49]. Here, we briefly summarize the main results and refer to ref. [49] for more details.

One of the minimal requirements of the operators in complexity is the operators need to be associative to construct a composite operators (bigger quantum circuits). Therefore, at least, the operators should belong to a monoid (a semigroup with an identity $\hat{\mathbb{I}}$). Let us denote a complexity of an operator \hat{x} in an arbitrary monoid \mathcal{O} by $\mathcal{C}(\hat{x})$.

Based on the essential and minimal properties of the complexity in the quantum circuit, it was proposed that the complexity should satisfy the following three axioms.

G1 [*Nonnegativity*]

$$\forall \hat{x} \in \mathcal{O}, \mathcal{C}(\hat{x}) \geq 0 \text{ and } \mathcal{C}(\hat{\mathbb{I}}) = 0.$$

G2 [*Series decomposition rule (triangle inequality)*]

$$\forall \hat{x}, \hat{y} \in \mathcal{O}, \mathcal{C}(\hat{x}\hat{y}) \leq \mathcal{C}(\hat{x}) + \mathcal{C}(\hat{y}).$$

G3a [*Parallel decomposition rule*]

$$\forall (\hat{x}_1, \hat{x}_2) \in \mathcal{N} = \mathcal{O}_1 \times \mathcal{O}_2 \subseteq \mathcal{O}, \mathcal{C}((\hat{x}_1, \hat{x}_2)) = \mathcal{C}(\hat{x}_1, \hat{\mathbb{I}}_2) + \mathcal{C}(\hat{\mathbb{I}}_1, \hat{x}_2).$$

Here, in **G3a**, we consider the case that there is a sub-monoid $\mathcal{N} \subseteq \mathcal{O}$ which can be decomposed into the Cartesian product of two monoids, i.e., $\mathcal{N} = \mathcal{O}_1 \times \mathcal{O}_2$. $\hat{\mathbb{I}}_1$ and $\hat{\mathbb{I}}_2$ are the identities of \mathcal{O}_1 and \mathcal{O}_2 . The Cartesian product of two monoids implies that $(\hat{x}_1, \hat{x}_2)(\hat{y}_1, \hat{y}_2) = (\hat{x}_1\hat{y}_1, \hat{x}_2\hat{y}_2)$ for arbitrary $(\hat{x}_1, \hat{x}_2), (\hat{y}_1, \hat{y}_2) \in \mathcal{N}$.

The first axiom **G1** is obvious by definition. The axioms **G2** and **G3a** states the relation between the composite operator and its component operators. In the quantum circuit, the operator $\hat{x}\hat{y}$ corresponds to the series circuit and (\hat{x}_1, \hat{x}_2) corresponds to the parallel circuit. The axiom **G2** will imply the triangle inequality once the complexity is interpreted as a distance in some metric space. In **G3a**, we stress that we consider only the case that the operators \hat{x}_1 and \hat{x}_2 are completely independent. It may not be possible for some systems. See figure 1 for a graphical explanation. In terms of the computer science, **G3a** states the relationship between the total complexity and the complexities of parallel sub-tasks.

The Cartesian product of two monoids is represented by the direct sum in a matrix representation. For example, if matrixes M_1 and M_2 are two representations of monoids \mathcal{O}_1 and \mathcal{O}_1 , then the representation of their Cartesian product is $M_1 \oplus M_2$ rather than $M_1 \otimes M_2$. Thus, in a matrix representation, **G3a** says, for arbitrary operators M_1 and M_2 ,

$$\mathcal{C}(M_1 \oplus M_2) = \mathcal{C}(M_1) + \mathcal{C}(M_2).$$

This equation can be generalized to the direct sum of more operators. For operators M_1, M_2, \dots, M_k

$$\mathbf{G3a} \Leftrightarrow \mathcal{C}\left(\bigoplus_{i=1}^k M_i\right) = \sum_{i=1}^k \mathcal{C}(M_i). \tag{2.1}$$




Figure 1. Schematic diagram for **G3a** in quantum circuits. Two operators \hat{x}_1 and \hat{x}_2 are independent so it is natural that $\mathcal{C}((\hat{x}_1, \hat{x}_2)) = \mathcal{C}((\hat{x}_1, \hat{\mathbb{I}}_2)) + \mathcal{C}((\hat{\mathbb{I}}_1, \hat{x}_2))$.

The axioms **G1**, **G2** and **G3a** are valid for both discrete and continuous systems. Next, we turn to the axiom for continuous systems, in particular, $SU(n)$ groups. For a given operator $\hat{O} \in SU(n)$,¹ as $SU(n)$ is connected, there is a curve $c(s)$ connecting \hat{O} and identity $\hat{\mathbb{I}}$, where the curve may be parameterized by s with $c(0) = \hat{\mathbb{I}}$ and $c(1) = \hat{O}$. The tangent of the curve, $\dot{c}(s)$, is assumed to be given by a right generator $H_r(s)$ or a left generator $H_l(s)$:

$$\dot{c}(s) = H_r(s)c(s) \quad \text{or} \quad \dot{c}(s) = c(s)H_l(s). \tag{2.2}$$

As shown in figure 2, this curve can be approximated by discrete forms:

$$\hat{O}_n = c(s_n) = \delta\hat{O}_n^{(r)}\hat{O}_{n-1} = \hat{O}_{n-1}\delta\hat{O}_n^{(l)}, \tag{2.3}$$

where $s_n = n/N$, $n = 1, 2, 3, \dots, N$, $\hat{O}_0 = \hat{\mathbb{I}}$ and $\delta\hat{O}_n^{(\alpha)} = \exp[H_\alpha(s_n)\delta s]$ with $\alpha = r$ or l and $\delta s = 1/N$. In general, the two generators $H_r(s)$ and $H_l(s)$ at the same point of the same curve can be different, i.e., $H_l(s) \neq H_r(s)$. Indeed, $H_r(s)$ is an adjoint transformation of $H_l(s)$,

$$H_r(s) = c(s)H_l(s)c(s)^{-1}, \tag{2.4}$$

from eq. (2.2).

For an arbitrary infinitesimal operator in $SU(n)$ group, the fourth axiom was proposed:

G4a [*Smoothness*] The complexity of any infinitesimal operator in $SU(n)$, $\delta\hat{O}^{(\alpha)} = \exp(H_\alpha\delta s)$, is a smooth function of only $H \neq 0$ and $\delta s \geq 0$, i.e.,

$$\mathcal{C}(\delta\hat{O}^{(\alpha)}) = \mathcal{C}(\hat{\mathbb{I}}) + \tilde{F}(H_\alpha)\delta s + \mathcal{O}(\delta s^2), \tag{2.5}$$

This axiom implies $\mathcal{C}(\delta\hat{O}^{(l)}) = \mathcal{C}(\delta\hat{O}^{(r)})$ if $\delta\hat{O}^{(l)} = \delta\hat{O}^{(r)}$, which means that an infinitesimal operator will give the same complexity contribution to the complexity regardless that it is added to the left-side or right-side.

¹In this paper, by $SU(n)$ and $U(n)$, we mean finite dimensional groups. For infinite dimensional cases, we will use the notation $SU(\infty)$ and $U(\infty)$.




Figure 2. A continuous curve $c(s)$ connects the identity ($c(0) = \hat{\mathbb{I}}$) and a particular operator \hat{O} ($c(1) = \hat{O}$). It can be approximated by a discrete form, where every intermediate point (\hat{O}_n) is also an operator.

All information regarding the complexity has been encoded in the function \tilde{F} so the cost ($L_\alpha[c]$) of a curve c can be defined as

$$L_\alpha[c] := \sum_{i=1}^N \mathcal{C}(\delta \hat{O}_i^{(\alpha)}) \xrightarrow{N \rightarrow \infty} \int_0^1 \tilde{F}(H_\alpha(s)) ds, \quad (2.6)$$

where we assume that the curve is constructed by only $\delta \hat{O}_n^{(r)}$ or only $\delta \hat{O}_n^{(l)}$. In a geometric picture, it is the length of a given curve and $\tilde{F} ds$ is considered as a line element in a geometry. Thus, a natural question is what kind of geometry can be allowed for complexity? It has been shown that it is the Finsler geometry as follows.

First, it has been proven that three axioms **G1**, **G2**, and **G4a** implies that the function \tilde{F} introduced in **G4a** satisfies

F1 (Nonnegativity) $\tilde{F}(H_\alpha) \geq 0$ and $\tilde{F}(0) = 0$

F2 (Positive homogeneity) $\forall \lambda \in \mathbb{R}^+, \tilde{F}(\lambda H_\alpha) = \lambda \tilde{F}(H_\alpha)$

F3 (Triangle inequality) $\tilde{F}(H_{\alpha,1}) + \tilde{F}(H_{\alpha,2}) \geq \tilde{F}(H_{\alpha,1} + H_{\alpha,2})$,

which are the defining properties of so called *Minkowski norm* in mathematics literature.² The conditions, **F1-F3**, characterize properties of a norm of the vector (the generators (H_α)) in the Lie algebra, the tangent space at the identity. The *Finsler metric* is nothing but a Minkowskia norm defined at all points on the base manifold. There are two natural but different ways to extend the Minkowski norm \tilde{F} at the identity to every point on the base manifold:

$$F_r(c, \dot{c}) := \tilde{F}(H_r) = \tilde{F}(\dot{c}c^{-1}), \quad \text{or} \quad F_l(c, \dot{c}) := \tilde{F}(H_l) = \tilde{F}(c^{-1}\dot{c}). \quad (2.7)$$

Here, we introduced ' $F_\alpha(c, \dot{c})$ ', a standard notation for the *Finsler metric* in mathematics literatures. In this paper, we will simply call both $\tilde{F}(H_\alpha)$ and $F_\alpha(c, \dot{c})$ a 'Finsler metric'

²Strictly speaking, the requirements of the Minkowski norm are a little more stronger than **F1-F3**. However, these differences will not affect our results in this paper.

if there is no confusion. We refer to refs. [54–57] for more detailed explanation of the Minkowski norm and Finsler geometry.

Note that there is symmetry in the Finsler metrics. $F_r(c, \dot{c})$ ($F_l(c, \dot{c})$) is *right(left)-invariant* because $H_r(H_l)$ is invariant under the right(left)-translation $c \rightarrow c\hat{x}$ ($c \rightarrow \hat{x}c$) for $\forall \hat{x} \in \text{SU}(n)$.

At this stage, any right or left invariant Finsler metric will be allowed for ‘complexity geometry’. By requiring that the complexity obeys some symmetries of quantum field theory the Finsler metric \tilde{F} can be more constrained. Indeed, the unitary invariance and the CPT symmetry of QFT respectively impose the following constraints on \tilde{F}

$$\text{[adjoint invariance]} \quad \tilde{F}(H_\alpha) = \tilde{F}(\hat{U}H_\alpha\hat{U}^\dagger), \quad (2.8)$$

$$\text{[reversibility]} \quad \tilde{F}(H_\alpha) = \tilde{F}(-H_\alpha), \quad (2.9)$$

for $\forall U \in \text{SU}(n)$. It has been shown that eq. (2.8) implies

$$\text{[Independence of left/right generators]} \quad \tilde{F}(H_l) = \tilde{F}(H_r), \quad (2.10)$$

so the Finsler metric is *bi-invariant*, which means both left and right invariant.

The constraints eq. (2.8) and eq. (2.9) (together with the axioms **G1-G4a**) turn out to be strong enough to determine the complexity of $\text{SU}(n)$ operators uniquely (up to an overall constant λ) as

$$F(c(s), \dot{c}(s)) = \tilde{F}(H) = \lambda \text{Tr} \sqrt{H(s)H^\dagger(s)}, \quad H(s) := H_l(s) \text{ or } H_r(s), \quad (2.11)$$

where $c(s)$ is a curve in $\text{SU}(n)$ group and λ is a positive constant. Note that the subscript α in H_α is dropped because H_r or H_l give the same Finsler metric (see also eq. (2.10)).

Finally, the complexity of an operator is defined by the minimal length (or minimal cost) of the curves connecting $\hat{\mathbb{I}}$ and \hat{O} :

$$\mathcal{C}(\hat{O}) := \min \left\{ L[c] \mid \forall c(s), c(0) = \hat{\mathbb{I}}, c(1) = \hat{O} \right\}. \quad (2.12)$$

Therefore, we are left with the variational problem minimizing

$$\int_0^1 \text{Tr} \sqrt{H(s)H^\dagger(s)} ds, \quad \text{with } c(0) = \hat{\mathbb{I}}, \text{ and } c(1) = \hat{O}. \quad (2.13)$$

This minimization problem is simplified thanks to *bi-invariance* of the Finsler metric explained below eq. (2.10). It has been shown in refs. [58, 59] that, if the Finsler metric is bi-invariant, the curve $c(s)$ is a geodesic if and only if there is a *constant* generator $H(s) = \bar{H}$ such that

$$\dot{c}(s) = \bar{H}c(s) \text{ or } c(s) = \exp(s\bar{H}). \quad (2.14)$$

In our case $\bar{H} = \ln \hat{O}$, by the boundary condition $\hat{O} = c(1) = \exp(\bar{H})$. Therefore,

$$\mathcal{C}(\hat{O}) = \min \left\{ \text{Tr} \sqrt{\bar{H}\bar{H}^\dagger} \mid \forall \bar{H} = \ln \hat{O} \right\}, \quad (2.15)$$

where ‘min’ is not in the sense of a variational problem. It picks up the minimal value among the multi-values of $\ln \hat{O}$.

3 Generalization of axioms

The axioms **G1** and **G2** are general enough but the axioms **G3a** and **G4a** may be relaxed more.

3.1 Generalized parallel decomposition rule

First, we generalize the parallel decomposition rule in the axiom **G3a** as follows

G3b [*Parallel decomposition rule*]

$$\forall(\hat{x}_1, \hat{x}_2) \in \mathcal{N} = \mathcal{O}_1 \times \mathcal{O}_2 \subseteq \mathcal{O}, [\mathcal{C}((\hat{x}_1, \hat{x}_2))]^p = [\mathcal{C}((\hat{x}_1, \hat{\mathbb{I}}_2))]^p + [\mathcal{C}((\hat{\mathbb{I}}_1, \hat{x}_2))]^p,$$

where p is a positive integer.

Here, the only difference from the axiom **G3a** is the existence of a positive integer number p . If $p = 1$, **G3b** becomes **G3a**. This new parallel decomposition rule may look less natural if $p \neq 1$ because we assumed that \hat{x}_1 and \hat{x}_2 are totally independent (figure 2). However, this p -deformed rule will open an interesting possibility that realize the Schatten norm eq. (1.2). If we use the matrix representation for the monoid, then the axiom **G3b** can be presented as

$$\mathbf{G3b} \Leftrightarrow \left[\mathcal{C} \left(\bigoplus_{i=1}^k M_i \right) \right]^p = \sum_{i=1}^k \mathcal{C}(M_i)^p. \quad (3.1)$$

When we take $p = 1$, this equation just recovers into the eq. (2.1). Note again that 1) $p = 1$ is the most natural choice intuitively and 2) we may ‘explain’ what the Schatten norm really means from the axiomatic point of view.

3.2 Generalized smoothness

In the axiom **G4a**, it is assumed that the complexity of any infinitesimal operator depends only on this infinitesimal operator itself. However, in principle, the complexity may depend on other factors, which are *independent* of any specific operator. In order to take into account this possibility we introduce w_α in the smoothness axiom **G4a**.

G4b The complexity of any infinitesimal operator in $SU(n)$, $\delta\hat{O}^{(\alpha)} = \exp(H_\alpha \delta s)$, satisfies,

$${}_p\mathcal{C}(\delta\hat{O}^{(\alpha)}, w_\alpha) = {}_p\mathcal{C}_\alpha(\hat{\mathbb{I}}) + {}_p\tilde{F}(H_\alpha, w_\alpha)\delta s + \mathcal{O}(\delta s^2), \quad (3.2)$$

where ${}_p\tilde{F}(H_\alpha, w_\alpha) := \frac{\partial}{\partial \delta s} [{}_p\mathcal{C}(\delta\hat{O}^{(\alpha)}, w_\alpha)]|_{\delta s=0}$ and ${}_p\mathcal{C}_\alpha(\hat{\mathbb{I}}) = 0$ by **G1**.

Here, the quantities with the index $\alpha = r, l$ are related to the right generator H_r or left generator H_l respectively. We use the left subscript p to distinguish a difference choice of p in the axiom **G3b**. The difference from the axiom **G4a** is the existence of $w_\alpha = \{w_\alpha^{(1)}, w_\alpha^{(2)}, \dots\}$. They stand for all the other possible variables defined at the Lie algebra $\mathfrak{su}(n)$. For example, w_α may stand for penalty or weight in the previous works [29, 31, 32, 50–52]. In general $w_r \neq w_l$ so it is possible ${}_p\mathcal{C}(\delta\hat{O}, w_r) \neq {}_p\mathcal{C}(\delta\hat{O}, w_l)$. It means that an infinitesimal operator $\delta\hat{O}$ can contribute to the complexity differently depending on whether it is added to the left-side or right-side. It is one difference compared with the axiom **G4a**. (See the comment below eq. (2.5).)

3.3 Emergence of the Finsler metric

Similar to eq. (3.3) we define the *cost* (${}_pL_\alpha[c, w_\alpha]$) of a curve c as

$${}_pL_\alpha[c, w_\alpha] := \sum_{i=1}^N {}_p\mathcal{C}(\delta\hat{O}_i^{(\alpha)}, w_\alpha) \xrightarrow{N \rightarrow \infty} \int_0^1 {}_p\tilde{F}_\alpha(H_\alpha(s), w_\alpha) ds, \quad (3.3)$$

where $\alpha = r$ or l denoting the curve constructed by only $\delta\hat{O}_n^{(r)}$ or only $\delta\hat{O}_n^{(l)}$ respectively. Similar to the case $p = 1$ and $w_\alpha = 0$ in ref. [49], we can prove that ${}_p\tilde{F}$ satisfies three defining properties of the Minkowski norm by using **G1**, **G2** and **G4b**:

F1 (Nonnegativity) ${}_p\tilde{F}(H_\alpha, w_\alpha) \geq 0$ and ${}_p\tilde{F}(H_\alpha, w_\alpha) = 0$ iff $H_\alpha = 0$

F2 (Positive homogeneity) $\forall \lambda \in \mathbb{R}^+, {}_p\tilde{F}(\lambda H_\alpha, w_\alpha) = \lambda {}_p\tilde{F}(H_\alpha, w_\alpha)$

F3 (Triangle inequality) ${}_p\tilde{F}(H_{\alpha,1}, w_\alpha) + {}_p\tilde{F}(H_{\alpha,2}, w_\alpha) \geq {}_p\tilde{F}(H_{\alpha,1} + H_{\alpha,2}, w_\alpha)$

Roughly speaking, **F1** is equivalent to **G1**, **F3** is equivalent to **G2**, and **F2** can be read from **G4b**. A detailed proof is essentially the same as the case of $p = 1$ and $w_\alpha = 0$ and provided in appendix A in ref. [49], where we only need to replace $\tilde{F}(H) \rightarrow {}_p\tilde{F}(H_\alpha, w_\alpha)$. The *Finsler metric* is a Minkowski norm defined at all points on the base manifold. Similarly to the case $p = 1, w_\alpha = 0$, there are two natural ways to extend the Minkowski norm \tilde{F} at the identity to every point on the base manifold:

$${}_pF_\alpha(c, \dot{c}, w_\alpha) := {}_p\tilde{F}(H_\alpha, w_\alpha), \quad \text{with } H_r = \dot{c}c^{-1} \text{ and } H_l = c^{-1}\dot{c}, \quad (3.4)$$

where we introduce a standard notation for the Finsler metric ‘ ${}_pF_\alpha(c, \dot{c})$ ’. Thus, we conclude the complexity is still given by a Finsler geometry with generalized axioms: **G3.G4** \rightarrow **G3b, G4b**.

Note also that ${}_pF_r(c, \dot{c}, w_r)$ is *right-invariant*, because ${}_p\tilde{F}$ is invariant under the right-translation $c \rightarrow c\hat{x}$ for $\forall \hat{x} \in \text{SU}(n)$. Similarly ${}_pF_l(c, \dot{c}, w_l)$ is *left-invariant*. Finally, the left or right complexity of an operator \hat{O} is identified with the minimal cost of the curves connecting $\hat{\mathbb{I}}$ and \hat{O} :

$${}_p\mathcal{C}_\alpha(\hat{O}, w_\alpha) := \min\{{}_pL_\alpha[c, w_\alpha] \mid \forall c(s), c(0) = \hat{\mathbb{I}}, c(1) = \hat{O}\}. \quad (3.5)$$

4 Constraints on the Finsler metric

So far, any Finsler metric will be allowed for the complexity geometry. In this section, we give constraints on the Finsler metric by taking into account some physical requirement or symmetry properties of quantum field theory.

First, we propose that the Finsler metric should be invariant under a adjoint transformation $H_\alpha \rightarrow \hat{U}H_\alpha\hat{U}^\dagger$, i.e.,

$${}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(\hat{U}H_\alpha\hat{U}^\dagger, w_\alpha), \quad \forall \hat{U} \in \text{SU}(n), \forall H_\alpha \in \mathfrak{su}(n), \quad (4.1)$$

which we call ‘adjoint invariance’. We provide two arguments: i) (mathematical/geometric reason) independence of left/right generators in subsection 4.1.1 ii) (physical reason) gauge

invariance in subsection 4.1.2. As a corollary, it will be shown that the adjoint invariance leads to the bi-invariance of the Finsler geometry. Next, by requiring the CPT symmetry of QFT we propose that the Finsler metric should satisfy

$${}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(-H_\alpha, w_\alpha), \tag{4.2}$$

which is called ‘reversibility’.

When it comes to the final results (4.1) and (4.2), this subsection generalizes the result in ref. [49] to the case with $p \neq 1$ and $w_\alpha \neq 0$. However, the supporting arguments here are different from ref. [49]. The arguments in ref. [49] and this paper are complementary and strengthen each other.

4.1 Adjoint invariance

4.1.1 Adjoint invariance from the independence of left/right generators

If the complexity is an intrinsic property of an operator in a given physical system, the length (cost) of the curve $c(s)$ should depend on the curve itself for given w_α .

However, as discussed in section 2 and ref. [49], at an arbitrary point $c(s_0)$ on an arbitrary curve c , there are two different ways to compute the length between $c(s_0)$ and $c(s_0 + \delta s)$:

$${}_p\tilde{F}(H_l, w_l)\delta s, \quad \text{or} \quad {}_p\tilde{F}(H_r, w_r)\delta s. \tag{4.3}$$

In the Nielsen’s original works, it is argued that the right-invariance is a natural condition. It is based on an operator (in discrete circuits) constructed as $U_n \cdot U_{n-1} \cdots U_1$. However, this is not the only possible way. For example, if we construct the operator as $U_1 \cdots U_{n-1} U_n$, by the same reason as Nielsen’s, the left-invariance will be the natural condition. Because, there is no a priori reason to choose among $U_n \cdot U_{n-1} \cdots U_1$ and $U_1 \cdots U_{n-1} U_n$ we just open both possibilities. See appendix A for more details. Because, there is no a priori reason to choose either the right-invariance or left-invariant complexity, we require they give the same physics, which means

$${}_p\tilde{F}(H_l, w_l) = {}_p\tilde{F}(H_r, w_r). \tag{4.4}$$

At first sight, it looks that eq. (4.4) is a very weak condition, because, for a given ${}_p\tilde{F}(H_r, w_r)$, we can always choose w_l so that eq. (4.4) holds. However, we will show that, because the operators form a group, eq. (4.4) leads to eq. (4.1).

Let us start with the following relation implied by eq. (2.2)

$$H_l = \hat{U}^{-1} H_r \hat{U}. \tag{4.5}$$

Here, because \hat{U} is arbitrary, eq. (4.5) can be written as

$$\begin{aligned} H_l^{(1)} &= \hat{U}_1^{-1} H_r \hat{U}_1. \\ H_l^{(2)} &= \hat{U}_2^{-1} H_r \hat{U}_2. \end{aligned} \tag{4.6}$$

with arbitrary operators \hat{U}_1 and \hat{U}_2 . In other words, the left generator corresponding to a given right generator H_r is not unique. This fact also can be interpreted for the generators




Figure 3. The schematic figure to show that the left generator is not unique for a given right generator H_r . The two left generators can be connected as: $H_l^{(2)} = \hat{U} H_l^{(1)} \hat{U}^{-1}$ with $\hat{U} = \hat{U}_2^{-1} \hat{U}_1$.

on the curves in the figure 3. A right generator H_r can appear at \hat{U}_1 and \hat{U}_2 in $SU(n)$ group. Depending on the positions \hat{U}_1 and \hat{U}_2 there will be different left generators.³ Finally, eq. (4.6) yields

$$H_l^{(2)} = \hat{U} H_l^{(1)} \hat{U}^{-1}, \quad \hat{U} := \hat{U}_2^{-1} \hat{U}_1 \tag{4.7}$$

where \hat{U} is an arbitrary operator in $SU(n)$ because \hat{U}_1 and \hat{U}_2 are arbitrary.

By using eq. (4.7) in eq. (4.4) we obtain

$${}_p\tilde{F}(H_r, w_r) = {}_p\tilde{F}(H_l^{(1)}, w_l) = {}_p\tilde{F}(H_l^{(2)}, w_l) = {}_p\tilde{F}(\hat{U} H_l^{(1)} \hat{U}^{-1}, w_l), \tag{4.8}$$

which proves that ${}_p\tilde{F}(H_l, w_l)$ is invariant under the adjoint transformation

$${}_p\tilde{F}(H_l, w_l) = {}_p\tilde{F}(\hat{U} H_l \hat{U}^{-1}, w_l), \quad \forall H_l \in \mathfrak{su}(n), \quad \forall \hat{U} \in SU(n). \tag{4.9}$$

Similarly, ${}_p\tilde{F}_r(H_r, w_r)$ is invariant under the adjoint transformation too.

Eq. (4.1) or (4.9) also implies the bi-invariance of our Finsler metric. Under the left translation $c(s) \rightarrow \hat{U}c(s)$

$${}_p\tilde{F}(H_r(s), w_r) \rightarrow {}_p\tilde{F}(\hat{U} H_r(s) \hat{U}^{-1}, w_r) = {}_p\tilde{F}(H_r(s), w_r) \tag{4.10}$$

where we used the adjoint-invariance for equality. Eq. (4.10) means that the right-invariant Finsler metric is also left invariant so it is bi-invariant. Similarly, the left-invariant Finsler metric is also bi-invariant.

Furthermore, if the Finsler metric is bi-invariant we can show that the Finsler metric is adjoint-invariance. For example, let us consider the right-invariant form of the Finsler metric, ${}_p\tilde{F}(H_r(s), w_r)$. In the left-translation, it becomes ${}_p\tilde{F}(\hat{U} H_r(s) \hat{U}^{-1}, w_r)$ and because it must be invariant we conclude ${}_p\tilde{F}(H_r(s), w_r) = {}_p\tilde{F}(\hat{U} H_r(s) \hat{U}^{-1}, w_r)$. In summary we have the following equivalence.

$$\text{left or right invariance} + \text{adjoint invariance} \quad \Leftrightarrow \quad \text{bi invariance} \tag{4.11}$$

³The generator is not the tangent vector of the curve. The tangent vector of the curve and generators are connected by eq. (2.2). Thus, the tangent vector of a curve at a point is unique but the generator is not.

4.1.2 Adjoint invariance from symmetric transformations

Let us consider a time evolution operator $c(t)$ of which generator is given by $H_\alpha := -i\mathbb{H}$ with a Hamiltonian \mathbb{H} (not the Hamiltonian density). We set $\hbar = 1$ for convenience. Suppose that two seemingly different Hamiltonians are related by a transformation S and both Hamiltonians give the same physical properties (except complexity yet). Now the question is “do they give the same complexity or not?” It will be natural to expect they also give the same complexity. Mathematically it means

$${}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(S(H_\alpha), w_\alpha), \quad S(H_\alpha) := -iS(\mathbb{H}), \quad (4.12)$$

where $S(\mathbb{H})$ denotes the Hamiltonian related to \mathbb{H} by the transformation S . w_α does not change because it is the parameter which is introduced in the definition of the complexity, not related with a Hamiltonian \mathbb{H} or a generator H_α .

Since the Hamiltonians $S(\mathbb{H})$ and \mathbb{H} are supposed to describe equivalent physics, they have the same observables such as energy and have the same eigenvalues. This means $S(\mathbb{H}) = \hat{U}\mathbb{H}\hat{U}^\dagger$ with a unitary operator \hat{U} .⁴ The adjoint invariance (4.1) or (4.12) is a *sufficient* condition to insure that the complexities given by \mathbb{H} and $S(\mathbb{H})$ are the same. We now want to show that the adjoint invariance is also the *necessary* condition to have this symmetry (the invariance of complexity under the transformation S). The main idea in following proof has three steps:

- (1) start with the generators for two simple symmetric transformations;
 - (2) by adding their commutators and linear combinations, construct more symmetric transformations;
 - (3) show that almost all the unitary transformations can be obtained by the above way.
- (1) Without loss of generality, we will consider a one-particle quantum mechanical system as an example.⁵ In appendix B, we obtain two kinds of special symmetric transformations, which are given by the following Lie algebras

$$\mathfrak{g}_s := \left\{ i\varphi(\vec{x}) \mid \forall C^\infty \text{ scalar field } \varphi(\vec{x}) \right\}, \quad (4.13)$$

and

$$\tilde{\mathfrak{g}}_s := \left\{ ic^{jl}\hat{p}_j\hat{p}_l \mid \forall c^{jl} \in \mathbb{R} \right\}. \quad (4.14)$$

The first is associated with the fact that we have a freedom to add divergent terms to a Lagrangian. The second is associated with some canonical transformations.

⁴We distinguish this from ‘representation/bases transformations.’ In representation/bases transformations, we fix the Hamiltonian operator but change its matrix components by choosing different bases in a Hilbert space. Here we change the Hamiltonian operator itself but do not change the bases in a Hilbert space. For example, we refer to the transformations \hat{U}_φ and \hat{W}_f in eqs. (B.7) and (B.32).

⁵Strictly speaking, this Hamiltonian may not have a finite dimensional representation, which cannot be covered by this paper. However, we expect that the general symmetries of the complexity should hold for both finite and infinite dimensional cases.

(2) Let us now consider the following series based on two special symmetric transformations defined in eqs. (4.13) and (4.14)

$$\mathfrak{g}_s^{(0)} := \{ag_1 + bg_2 \mid \forall g_1 \in \mathfrak{g}_s, \forall g_2 \in \tilde{\mathfrak{g}}_s, \forall a, b \in \mathbb{R}\}, \quad (4.15)$$

and

$$\mathfrak{g}_s^{(n)} := \left\{ ag_1 + bg_2 \mid \forall g_1 \in \mathfrak{g}_s^{(n-1)}, \forall g_2 \in [\mathfrak{g}_s^{(n-1)}, \mathfrak{g}_s^{(n-1)}], \forall a, b \in \mathbb{R} \right\}, \quad (4.16)$$

for $n \geq 1$. Here, the commutator $[\mathfrak{g}_s^{(n-1)}, \mathfrak{g}_s^{(n-1)}]$ is defined as

$$[\mathfrak{g}_s^{(n-1)}, \mathfrak{g}_s^{(n-1)}] := \left\{ [g_1, g_2] \mid \forall g_1, g_2 \in \mathfrak{g}_s^{(n-1)} \right\}. \quad (4.17)$$

Because the generators in \mathfrak{g}_s and $\tilde{\mathfrak{g}}_s$ generate symmetric transformations of Finsler metric, the generators in $[\mathfrak{g}_s^{(n-1)}, \mathfrak{g}_s^{(n-1)}]$ also generate symmetric transformations of Finsler metric. Thus, the elements of $\mathfrak{g}_s^{(n)}$ are all anti-Hermit operators generating symmetric transformations of ${}_p\tilde{F}(-i\mathbb{H}, w_\alpha)$. For example, in one-dimensional case,

$$\mathfrak{g}_s^{(1)} = \left\{ i\varphi_1(\hat{x}) + i(\varphi_2(\hat{x})\hat{p} + \hat{p}\varphi_2(\hat{x})) + ic\hat{p}^2 \mid \forall C^\infty \text{ scalar fields } \varphi_1, \varphi_2 \text{ and } c \in \mathbb{R} \right\}, \quad (4.18)$$

and

$$\mathfrak{g}_s^{(2)} = \left\{ i\varphi_1(\hat{x}) + i(\varphi_2(\hat{x})\hat{p} + \hat{p}\varphi_2(\hat{x})) + i(\varphi_3(\hat{x})\hat{p}^2 + \hat{p}^2\varphi_3(\hat{x})) \mid \forall C^\infty \text{ scalar fields } \varphi_1, \varphi_2 \text{ and } \varphi_3 \right\}. \quad (4.19)$$

In general,

$$\mathfrak{g}_s^{(n)} \subsetneq \mathfrak{g}_s^{(n+1)} \quad (4.20)$$

for all $n \geq 0$ so the set $\mathfrak{g}_s^{(n)}$ will be bigger and bigger when we increase n . In the limit $n \rightarrow \infty$, we define

$$\mathfrak{g}_s^{(\infty)} := \lim_{n \rightarrow \infty} \mathfrak{g}_s^{(n)}, \quad (4.21)$$

which is closed under the commutators and forms a Lie algebra.

(3) In appendix C we have shown

$$\mathfrak{g}_s^{(\infty)} = \left\{ iH(\hat{x}, \hat{p}) \mid \forall H(\hat{x}, \hat{p}) = H(\hat{x}, \hat{p})^\dagger, \right. \\ \left. H(x, p) \text{ is smooth and has a Taylor's expansion with respect to } p \text{ at } p = 0 \right\}.$$

Interestingly, $\mathfrak{g}_s^{(\infty)}$ contains almost all the possible anti-Hermit operators of a particle in one-dimensional space, although our starting point only contains two kinds of very special generators, eqs. (4.13) and (4.14). It can be generalized to higher dimensional cases so we obtain one symmetric group for ${}_p\tilde{F}(-i\mathbb{H}, w_\alpha)$:

$$G_s^{(\infty)} = \left\{ \exp[iH(\vec{\hat{x}}, \vec{\hat{p}})] \mid \forall H(\vec{\hat{x}}, \vec{\hat{p}}) = H(\vec{\hat{x}}, \vec{\hat{p}})^\dagger, \right. \\ \left. H(\vec{x}, \vec{p}) \text{ is smooth and has a Taylor's expansion with respect to } \vec{p} \text{ at } \vec{p} = 0 \right\}.$$

This is enough to show that the Finsler metric should be invariant under all unitary transformation.

Recently, ref. [60] provided a different supporting argument for the adjoint symmetry (4.1) based on the symmetry of the partition function/generating functional of quantum systems.

4.2 Reversibility from the CPT symmetry

Let us consider the effect of the CPT symmetry⁶ of the quantum field theory on the Finsler metric. For a quantum field Φ , the time evolution is given by $\Phi(\vec{x}, t) := c(t)^\dagger \Phi(\vec{x}, 0) c(t)$, where $c(t)$ is an arbitrary curve in the $SU(n)$ group. By denoting the CPT partner of $\Phi(\vec{x}, t)$ by $\bar{\Phi}(\vec{x}, t)$ we have

$$\begin{aligned} \bar{\Phi}(\vec{x}, t) &= C \circ P \circ T [c(t)^\dagger \Phi(\vec{x}, 0) c(t)] \\ &= c(-t)^\dagger \bar{\Phi}(\vec{x}, 0) c(-t), \end{aligned} \tag{4.22}$$

where $c(t)$ does not have charge and spatial variables \vec{x} . Thus, the evolution of the CPT partner is $c(-t) =: \bar{c}(t)$. Given the CPT symmetry of the theory, it is natural to expect that the costs of $c(t)$ and $\bar{c}(t)$ is also the same, i.e.,

$${}_p L_\alpha [c, w_\alpha] = {}_p L_\alpha [\bar{c}, w_\alpha]. \tag{4.23}$$

Because the generator of $\bar{c}(s)$ is given by $\bar{H}_\alpha(t) = -H_\alpha(t)$,⁷ eq. (4.23) yields

$$\int_0^1 {}_p \tilde{F}(H_\alpha(t), w_\alpha) dt = \int_0^1 {}_p \tilde{F}(-H_\alpha(t), w_\alpha) dt. \tag{4.24}$$

Because it is valid for arbitrary generators we have

$$\tilde{F}_\alpha(H_\alpha, w_\alpha) = \tilde{F}_\alpha(-H_\alpha, w_\alpha). \tag{4.25}$$

Strictly speaking, our argument here applies for $SU(\infty)$. We use the intuition from the $SU(\infty)$ case to give a plausibility argument for the ‘reversibility’ (4.23) of the $SU(n)$ case with finite n .

Path-reversal symmetry. By using the adjoint invariance eq. (4.1) and the reversibility eq. (4.2) we can prove the ‘path-reversal symmetry’ of the cost for an arbitrary curve:

$$L_\alpha [c, w_\alpha] = L_\alpha [c^{-1}, w_\alpha], \quad \forall c(s). \tag{4.26}$$

Note that if the curve $c(s)$ is generated by $H_\alpha(s)$, $c^{-1}(s) := [c(s)]^{-1}$ is not generated by $-H_\alpha(s)$ but by $-c^{-1} H_r(c) c$ for $\alpha = r$ and $-c H_l(c) c^{-1}$ for $\alpha = l$. For example, for the right generator, $H_r(c^{-1}) = (dc^{-1}/ds)c = -c^{-1}(\dot{c}c^{-1})c = -c^{-1} H_r(c) c$. Thus

$$\tilde{F}(H_r(c^{-1}), w_r) = \tilde{F}(-c^{-1} H_r(c) c, w_r) = \tilde{F}(H_r(c), w_r), \tag{4.27}$$

⁶The ‘charge conjugation’ C, ‘parity transformation’ (‘space inversion’) P and ‘time reversal’ T.

⁷For example, for the right generator, $\bar{H}_r(t) = (d\bar{c}/dt)\bar{c}^{-1} = -(dc/dt)c^{-1} = -H_r(t)$.

which gives eq. (4.26). Here, we used the adjoint invariance eq. (4.1) and the reversibility eq. (4.2) in the second equality. The left-generator case works similarly.

In fact, the reverse also holds, i.e. eq. (4.26) implies eqs. (4.1) and (4.2). First, by considering a special case $c = e^{Hs}$ with a constant H , eqs. (4.2) can be derived from eq. (4.26) and eq. (4.27). Thus, we are left with

$$\int_0^1 {}_p\tilde{F}(H_\alpha(t), w_\alpha) dt = \int_0^1 {}_p\tilde{F}(c^{-1}H_\alpha(t)c, w_\alpha) dt. \quad (4.28)$$

It is valid for arbitrary H_α and c so we have eq. (4.1). As a result, we have the following equivalence between the path-reversal symmetry of the cost and the adjoint invariance plus reversibility of the Finsler metric:

$$\begin{aligned} \text{Path reversal symmetry: } & \forall c(s), \quad L_\alpha[c, w_\alpha] = L_\alpha[c^{-1}, w_\alpha] \\ \Leftrightarrow \left\{ \begin{array}{l} \text{adjoint invariance: } \quad {}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(\hat{U}H_\alpha\hat{U}^\dagger, w_\alpha); \\ \text{reversibility: } \quad \quad \quad {}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(-H_\alpha, w_\alpha). \end{array} \right. & (4.29) \end{aligned}$$

5 Finsler metric and complexity of $SU(n)$ operators

5.1 Finsler metric of $SU(n)$ operators

So far, we have found, for the complexity geometry, we need the Finsler metric satisfying two constraints eqs. (4.1) and (4.2). It turns out that these constraints with **G3b** are strong enough to determine the Finsler metric in the operator space of any $SU(n)$ groups uniquely (up to an overall constant λ)

$${}_p\tilde{F}(H(s), w_\alpha) = \lambda \left\{ \text{Tr} \left[\left(H(s)H(s)^\dagger \right)^{p/2} \right] \right\}^{1/p}, \quad (5.1)$$

where $H(s) = H_r(s)$ or $H_l(s)$ for the curve $c(s)$ and $\lambda := \lambda_r(w_r) = \lambda_l(w_l)$ is arbitrary constant. The proof is similar to the case with $p = 1$ and $w_\alpha = 0$ in ref. [49] and consists of the following four steps.

- ① Note that, by using a unitary matrix \hat{U} , H_α always can be diagonalized and the position of eigenvalues can be exchanged. Thus, the adjoint invariance (4.1), ${}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(\hat{U}H_\alpha\hat{U}^\dagger, w_\alpha)$, implies that ${}_p\tilde{F}(H_\alpha, w_\alpha)$ is only a function of eigenvalues of H_α and independent of the permutations of these eigenvalues. Therefore, without loss of generality, we can say

$$H_\alpha = \text{diag}(i\gamma_1, i\gamma_2, \dots, i\gamma_n) =: \bigoplus_{j=1}^n i\gamma_j, \quad (5.2)$$

where H_α is anti-Hermitian and we separate i from the eigenvalues $i\gamma_j$ with $\gamma_j \in \mathbb{R}$. There is no index for α in the eigenvalues because H_r and H_l is related by a unitary matrix \hat{U} (see eq. (4.5)) and their eigenvalues are the same.

② **G3b** or eq. (3.1) implies that if $H_\alpha = H_1 \oplus H_2$

$$\left({}_p\tilde{F}(H_\alpha, w_\alpha)\right)^p = \left({}_p\tilde{F}(H_1 \oplus \mathbf{0}_{n-k}, w_\alpha)\right)^p + \left({}_p\tilde{F}(\mathbf{0}_n \oplus H_2, w_\alpha)\right)^p. \quad (5.3)$$

With the generator (5.2), eq. (5.3) reads

$$\left({}_p\tilde{F}(H_\alpha, w_\alpha)\right)^p = \sum_{j=1}^n (f_\alpha(i\gamma_j), w_\alpha)^p = \sum_{j=1}^n (f_\alpha(i|\gamma_j|), w_\alpha)^p \quad (5.4)$$

where f_α is a function of the eigenvalues independent of their order. In the second equality, the reversibility (4.2) ${}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(-H_\alpha, w_\alpha)$ was used.

③ Using the positive homogeneity **F2** below eq. (3.3), ${}_p\tilde{F}(\beta H_\alpha, w_\alpha) = \beta {}_p\tilde{F}(H_\alpha, w_\alpha)$ for $\beta \in \mathbb{R}^+$, we obtain ${}_p\tilde{F}(H_\alpha, w_\alpha) = \lambda_\alpha(w_\alpha) \left[\sum_{j=1}^n |\gamma_j|^p\right]^{1/p}$, where $\lambda_\alpha(w_\alpha)$ is an overall constant. Thus,

$${}_p\tilde{F}(H_\alpha(s), w_\alpha) = \lambda_\alpha(w_\alpha) \left[\sum_{j=1}^n |\gamma_j|^p\right]^{1/p} = \lambda_\alpha(w_\alpha) \left\{ \text{Tr} \left[\left(H_\alpha(s) H_\alpha(s)^\dagger \right)^{p/2} \right] \right\}^{1/p}. \quad (5.5)$$

④ By eq. (4.5) the trace part of the last term in eq. (5.5) are the same for $\alpha = r$ and $\alpha = l$. By eq. (4.4), we obtain $\lambda_r(w_r) = \lambda_l(w_l) =: \lambda$. Thus, eq. (5.1) is proven.⁸ \square

Note that our final results (5.1) does not depend on $\alpha = r, l$ as expected from eq. (4.4). Thus, from here, we will omit the indexes r, l and the symbol w_α in the Finsler metric. Without loss of generality, we may set

$$\lambda = \sqrt{2} n^{\frac{1}{2} - \frac{1}{p}}, \quad (5.6)$$

so that ${}_p\tilde{F}(\hat{\mathbb{I}}) = {}_2\tilde{F}(\hat{\mathbb{I}})$.⁹ The unimportant factor $\sqrt{2}$ was introduced for future convenience. For example, by this factor, eq. (5.11) becomes simplified.

There are two values of p of special interest in eq. (5.1). For $p = 1$ we obtain the result in ref. [49],

$${}_1\tilde{F}(H(s)) = \sqrt{\frac{2}{n}} \text{Tr} \sqrt{H(s) H(s)^\dagger}. \quad (5.7)$$

For $p = 2$

$${}_2\tilde{F}(H(s)) = \sqrt{2} \sqrt{\text{Tr} [H(s) H(s)^\dagger]}. \quad (5.8)$$

Interestingly enough, it turns out that the $p = 2$ case gives just the bi-invariant Riemannian metric ('standard' metric) of $SU(n)$ group. To show it, let us consider two tangent vectors V_1 and V_2 at the point \hat{U} can be written as

$$V_k = H_k \hat{U} = i H_k^a T_a \hat{U}, \quad k = 1, 2, \quad (5.9)$$

⁸A penalization of different generator directions for other groups than $SU(n)$ may be possible in principle. For example, for non-unitary representations of some groups, as they are not related to quantum mechanical processes, there is no physical principle to restrict the complexity for them and their penalties may be chosen arbitrarily.

⁹This choice is just one convention.

where $H_k := iH_k^a T_a$ is generator for V_k , H_k^a is a real number, and $\{T_a, a = 1, 2, \dots, n^2 - 1\}$ are bases of Lie algebra $\mathfrak{su}(n)$ in the fundamental representation. The basis satisfy

$$T_a T_b = \frac{1}{2n} \delta_{ab} \hat{\mathbb{1}} + \frac{1}{2} (i f_{ab}{}^c + d_{ab}{}^c) T_c, \quad T_a^\dagger = T_a, \quad \text{Tr}(T_a) = 0, \quad (5.10)$$

where $f_{ab}{}^c$ are the structure constants antisymmetric in all indices and the d -coefficients are symmetric in all indices. The metric tensor at the identity is given by

$$\tilde{g}_{ab} := \frac{1}{2} \frac{\partial^2 \tilde{F}^2}{\partial H^a \partial H^b} = \delta_{ab}. \quad (5.11)$$

We may compare the metric (5.11) with the Killing form of $\mathfrak{su}(n)$. For semi-simple Lie algebra, the bi-invariant metric must be proportional to its Killing form. The $\mathfrak{su}(n)$ Lie algebra is semisimple and the Killing form of $\mathfrak{su}(n)$ is [61]

$$B(H_1, H_2) = -n \delta_{ab} H_1^a H_2^b. \quad (5.12)$$

Noting that it is a unique candidate of the metric near identity up to proportionality constant [61], we conclude that our metric eq. (5.11) is consistent with the Killing form and fixes the proportionality constant.

5.2 Complexity of $SU(n)$ operators

Now we have the precise Finsler metric, the next step is to compute the complexity by finding the path of minimal cost (length) as shown in eq. (3.5). This minimization becomes straightforward thanks to the *bi-invariance* proven in eq. (4.10). It has been shown that, in bi-invariant Finsler geometry, the curve $c(s)$ is a geodesic if and only if there is a constant generator $H(s) = \bar{H}$ such that [58, 59]

$$\dot{c}(s) = \bar{H} c(s) \quad \text{or} \quad c(s) = \exp(s \bar{H}). \quad (5.13)$$

With the boundary condition $\hat{O} = c(1) = \exp(\bar{H})$, $\bar{H} = \ln \hat{O}$, which is the same as ref. [49]. Because \bar{H} is constant, eqs. (3.3) yields

$${}_p L[c] = {}_p \tilde{F}(\bar{H}) = \lambda \left\{ \text{Tr} \left[\left(\bar{H} \bar{H}^\dagger \right)^{p/2} \right] \right\}^{1/p}, \quad (5.14)$$

where λ is defined in eq. (5.6). Finally, the complexity of \hat{O} in eq. (3.5) is given by

$${}_p \mathcal{C}(\hat{O}) = \min \left\{ {}_p \tilde{F}(\bar{H}), \forall \bar{H} = \ln \hat{O} \right\}. \quad (5.15)$$

Here ‘min’ means the minimal value among multi-values of $\ln \hat{O}$, which corresponds to the possibility that the geodesic is not unique.

Note that the geodesics are independent of the value of p . It is because the geodesic is determined only by the *constant* generator in the bi-invariant Finsler geometry, no matter what specific metrics are given. The value of p only affects the metric so the numerical value of the length of the geodesics. Therefore, all different choices of p will give the same

qualitative results for the complexity apart from the numerical values of the complexity. Thanks to this property, it is enough to choose a specific value of p to investigate the property of the complexity of $SU(n)$ operators. We choose $p = 2$ because in this case the Finsler metric becomes the ‘standard’ bi-invariant Riemannian metric in $SU(n)$ groups and we can use a well-developed mathematics for the Riemannian geometry. In the following, we will mostly focus on $p = 2$ but all the conclusion are still valid for arbitrary $p > 0$.

It is well known that the ‘standard’ bi-invariant Riemannian metric in $SU(n)$ groups is given *uniquely* by the Killing form of $\mathfrak{su}(n)$ up to overall constants. However, this is not true if we allow the geometry to be general Finsler geometry. For $SU(n)$ groups, there are infinitely many inequivalent Finsler geometries. For example, we refer to ref. [62] for the way to construct a series of infinite inequivalent Finsler geometries.

6 Properties of operator complexity

In this section, we discuss some properties which can be derived from our results, eq. (5.15). First, we investigate the geodesic deviation and chaos and compare our results with previous works based on “k-local” metrics in refs. [29–31]. Next, we show that the pattern of the time-evolution of the complexity conjectured in [5, 31] can be concretely realized in our formalism of the complexity. Finally, we study the complexity related properties of the precursor operators and compare with the results in refs. [3, 29].

6.1 Geodesic deviation and chaos

Refs. [29–31] considered the geodesic deviation of two geodesics generated by the generators H and $H + \Delta\delta\theta$ respectively, where $\delta\theta$ is an infinitesimal parameter and $[H, \Delta] \neq 0$.¹⁰ It was argued that, in order to reflect the quantum chaos, the geodesics generated by H and $H + \Delta\delta\theta$ must diverge exponentially in a exponential time region. Whether the geodesics diverge or converge has been quantified by the ‘sectional curvature’: if the sectional curvature is negative (positive) the geodesics diverge (converge). We justify this criteria by using the Jacobi field (a vector quantifying geodesic deviation) in appendix E. It has been argued in refs. [29–31] that because the sectional curvatures of the bi-invariant metric are all nonnegative the bi-invariant geometry cannot exhibit any chaotic behavior. However, in this subsection we provide an argument for the possibility that even a bi-invariant geometry may allow the diverging geodesics and so “chaotic” behaviours.

The key point of our argument is the careful analysis of the ‘sectional curvature’ quantifying the geodesic deviation. Let us consider a Riemannian manifold M with a metric $g(\cdot, \cdot)$ and two linearly independent tangent vectors $\{X, Y\}$ at the same point in M . The sectional curvature ($k_M(X, Y)$) is defined as

$$k_M(X, Y) := \frac{g(R_M(X, Y)X, Y)}{g(X, X)g(Y, Y) - [g(X, Y)]^2}, \tag{6.1}$$

where R_M is the Riemannian curvature tensor of M with the metric $g(\cdot, \cdot)$.

¹⁰ $[H, \Delta] \neq 0$ is necessary to insure the sectional curvature, which will be explain below, is nonzero.

However, the evolution of the geodesic deviation between two neighboring geodesics $\xi_1(s) = \exp(Hs)$ and $\xi_2(s) = \exp[(H + \Delta\delta\theta)s]$ is not determined by this sectional curvature eq. (6.1). Instead, it is determined by the sectional curvature ($k_{\mathcal{M}}(X, Y)$) of a two dimensional manifold $\mathcal{M} \subset M$ where two geodesic $\xi_1(s)$ and $\xi_2(s)$ belongs to:

$$k_{\mathcal{M}}(X, Y) := \frac{\tilde{g}(R_{\mathcal{M}}(X, Y)X, Y)}{\tilde{g}(X, X)\tilde{g}(Y, Y) - [\tilde{g}(X, Y)]^2}. \quad (6.2)$$

Here $R_{\mathcal{M}}$ is the Riemannian curvature tensor of \mathcal{M} and $\tilde{g}(\cdot, \cdot)$ is the induced metric in \mathcal{M} .

Let us denote two orthonormal vector fields by $\{e_{\parallel}, e_{\perp}\}$ tangent to \mathcal{M} embedded in an N -dimensional manifold M . Let us also denote $N - 2$ independent orthonormal vector fields perpendicular to \mathcal{M} by e_i where $i = 1, 2, \dots, N - 2$. By the *Gauss-Codazzi* equation we have the following relation between $k_M(e_{\parallel}, e_{\perp})$ and $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$,

$$k_M(e_{\parallel}, e_{\perp}) = k_{\mathcal{M}}(e_{\parallel}, e_{\perp}) + \sum_{i=1}^{N-2} [K_i(e_{\parallel}, e_{\perp})^2 - K_i(e_{\perp}, e_{\perp})K_i(e_{\parallel}, e_{\parallel})], \quad (6.3)$$

where $K_i(\cdot, \cdot)$ is the second fundamental form associated with the normal vector field e_i . For more details for eq. (6.3), we refer to chapter 11.4c of ref. [63].

As shown in appendix E it is $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$ that governs the geodesic deviation, not $k_M(e_{\parallel}, e_{\perp})$. Note that it is possible

$$\sum_{i=1}^{N-2} K_i(e_{\parallel}, e_{\perp})^2 - K_i(e_{\perp}, e_{\perp})K_i(e_{\parallel}, e_{\parallel}) > k_M(e_{\parallel}, e_{\perp}), \quad (6.4)$$

in some regions of a special \mathcal{M} , if K_i is not positive definite. As a result, it is possible that $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$ is negative even if $k_M(e_{\parallel}, e_{\perp}) > 0$. For example, it is known that, for a bi-invariant metric, $k_M(X, Y)$ is always nonnegative for any pair of tangent vectors $\{X, Y\}$ [61], but even in this case, it is possible that the neighboring geodesics converge in a region where $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$ is positive and diverge in a region where $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$ is non-positive.

We may understand this possibility in a different way. For a general H and Δ , the 2-dimensional sub-manifold \mathcal{M} may not form a subgroup so in this case $\tilde{g}(\cdot, \cdot)$ is no longer a bi-invariant metric of \mathcal{M} . Thus, the bi-invariant metric $g(\cdot, \cdot)$ can insure $k_M(e_{\parallel}, e_{\perp}) \geq 0$ but cannot insure $k_{\mathcal{M}}(e_{\parallel}, e_{\perp}) \geq 0$. For some particular choices of H and Δ , it is possible that the sectional curvature of \mathcal{M} is negative in some regions along $\xi_1(s)$.

In particular, for a $SU(n)$ group with large n , there are many linear independent generators to choose, so there are more possibilities to find two generators $\{H, \Delta\}$ such that two geodesics $\exp(Hs)$ and $\exp[(H + \Delta\delta\theta)s]$ can lay in a 2-dimensional sub-manifold \mathcal{M} of which sectional curvature is negative along $\xi_1(s)$ for some range of s . For finite n , it is not so easy to find an example to support our claims explicitly because the group manifold is high dimensional and its metric is very complicated. However, for $SU(\infty)$ representation, ref. [60] has shown an AdS_3 spacetime (rather than only a time slices in AdS_3) emerges as a complexity geometry, which implies the negative sectional curvature.

6.1.1 Comparison with “ k -local” metrics

In this subsection, we compare physics of the geodesic deviations from our Finsler metric with the ones from the “ k -local” metrics [29–31]

First of all, let us start with an important statement which has not been clarified in previous works. “It is impossible to make the curve $\xi_1(s) = \exp(Hs)$ (where H is constant) to be a geodesic completely lay in a negatively curved space no matter what metric we choose.” The reason is as follows. i) The geodesic in a negatively curved 2-dimensional space will never approach to the original point again as s increases as we explained in section 6.1, ii) $\exp(Hs)$ can approach arbitrarily to the original point again and again as s increases due to the quantum recurrence.¹¹ iii) Therefore, $\exp(Hs)$ cannot be a geodesic which completely lays in a negatively curved space. Thus, the correspondence between the complexity geometry and hyperbolic geometry argued in ref. [30] will be valid only *locally* in some regions of $\xi_1(s)$.

As a corollary, the sectional curvature can be negative only in some regions of $\xi_1(s)$ rather than in the whole $\xi_1(s)$ in *both* our bi-invariant metric and the “ k -local” metric case, where the geodesic is also given by constant generators. The “ k -local” metric with some suitable penalties can insure some sectional curvatures to be negative near the identity [31] but positive sectional curvature must appear somewhere else; the bi-invariant metric makes the sectional curvature positive near the identity¹² but the negative sectional curvature can appear somewhere else. In this sense, the bi-invariant metric and “ k -local” metric have no essential difference on the aspect of geodesic deviation.

Compared with the negative sectional curvature discussed in refs. [29–31], the negative sectional curvature appearing in our bi-invariant metric is determined intrinsically by the $SU(n)$ group itself rather than caused by some artificial penalties. The same $SU(n)$ group can appear in many different physical systems. Thus, the “chaos” appearing in the bi-invariant complexity can be seen as the universal property of complexity geometry across the various different physical systems. On the contrary, the “chaos” discussed by refs. [29–31] is not universal, as it depends on how to give the penalties to different generators. For example, in ref. [31], the sectional curvature at the identity can be negative only if we choose penalty $\mathcal{I}_2 = 1$ “2-local” generators and $\mathcal{I}_n > 4/3$ for all other generators.

Finally, the main motivation of refs. [29–31] to require the negative sectional curvature for the complexity geometry is to reflect the chaos of a time dependent quantum state. In our opinion, this quantum chaos should be achieved by the combination of the Hamilton and initial state rather than by the Hamiltonian itself. For example, for chaotic nonlinear

¹¹This point can be understood as follows. Supposed $i\gamma_1, i\gamma_2, \dots, i\gamma_n$ are the eigenvalues of H . Then $\text{Tr}[(e^{Ht} - \hat{\mathbb{1}})(e^{Ht} - \hat{\mathbb{1}})^\dagger] = 2 \sum_{i=1}^n [1 - \cos(\gamma_i t)]$ can be made arbitrarily small for a large enough time t , which follows from the existence of large t such that

$$0 < (\gamma_i t \bmod 2\pi) < \delta$$

for all γ_i with arbitrary small δ .

¹²At the identity, we always have $k_M(e_{\parallel}, e_{\perp}) = k_M(e_{\perp}, e_{\parallel})$ for any two generators H and Δ due to the fact that surface \mathcal{M} is locally geodesic at identity, where all K_i vanish. See the chapter 11.4c of ref. [63] for details.




Figure 4. The conjectured schematic diagram for the complexity evolution of the operator $\exp(Hs)$, where H is a constant generator in $\mathfrak{su}(n)$. The complexity first grows linearly when $s < s_c$ and reaches its maximum at the time $s = s_c \leq \mathcal{O}(\sqrt{n}) \propto \mathcal{O}(e^{d/2})$, where d is proportional to the classical degree of freedom of the system i.e., the size of classical phase space. At a very large time $s = s_r \propto \mathcal{O}(\exp(e^d))$, the quantum recurrence occurs and the complexity goes down to zero.

systems, the evolutions may be chaotic for some initial conditions but may not for other initial conditions. Therefore, also in quantum chaos, it seems to be better to focus on the trajectories in the Hilbert space rather than the trajectories of $SU(n)$ group operators. From this perspective, the quantum chaos can be read from the complexity between the *states* rather than operators, for example between two states: $|\psi(t)\rangle := \exp(Ht)|\psi_0\rangle$ and initial state $|\psi_0\rangle$ or a small perturbation $|\psi'(t)\rangle := \exp((H + \delta H)t)|\psi_0\rangle$. Indeed, in ref. [64], we show that the chaos can be presented by the exponential growth of the complexity between $|\psi(t)\rangle = \exp(Ht)|\psi_0\rangle$ and $|\psi_0\rangle$ rather than $\exp(Ht)$ and $\exp((H + \delta H)t)$. To obtain this result, we first defined the complexity between states based on the complexity of operators in this paper, which is an important result developed in ref. [64].

6.2 Linear growth and quantum recurrence

In this section, we show how the complexity of “time-dependent” operator $\hat{U}(s) = \exp(Hs)$ evolves as time “ s ” goes on. It was argued that the complexity of $\exp(Hs)$ should show the property in the figure 4 based on the quantum circuits. The complexity of $\exp(Hs)$ first grows linearly until it saturates the maximum value. Such a linear growth time in general is of order e^d , where d is the number of the classical degrees of freedom. After then there appears some fluctuations until $s = s_r \sim \exp(e^d)$. At $s \sim s_r$ the quantum recurrence occurs, i.e, $\exp(Hs_r) \approx \hat{\mathbb{I}}$ and $\mathcal{C} \approx 0$ [5, 31].

Let us show that our complexity based on the Finsler metric (5.1) can realize these properties. First, because of the bi-invariance, the curve $\exp(Hs)$ with constant H is a geodesic. Thus, *naively* we may conclude

$${}_p\mathcal{C}(\exp(Hs)) = {}_p\tilde{F}(H)s, \tag{6.5}$$

where we relaxed the range of our parameterization s , which is $s \geq 0$. Any value of s can be the ending point of the path, where our target operator is located at.

However, it turns out that this linear growth behavior of the complexity will be valid only for $s < s_c$ as shown in figure 4, where s_c is some critical time scale to be estimated below. For $s = s_0 > s_c$, $\exp(Hs)$ is still the geodesic connecting $\hat{\mathbb{I}}$ to $O_0 = e^{Hs_0}$ but will not be the shortest geodesic due to the multi-valuedness of $\log O$. In general, there may be a shorter geodesic $e^{H's'}$ connecting $\hat{\mathbb{I}}$ and O_0 , which gives the complexity.

In the case $p = 2$, the geometry is bi-invariant Riemannian geometry and the existence of the maximum complexity can be anticipated by the relation between the topology of a manifold and curvature of $SU(n)$ groups. According to the Bonnet-Myers theorem the largest distance between two arbitrary points is given by (see the theorem 2.19 in ref. [61].)

$$\frac{\pi}{\sqrt{\Xi}}, \tag{6.6}$$

where Ξ is defined by the following relation between the Ricci tensor $\text{Ric}(\cdot, \cdot)$ and metric $g(\cdot, \cdot)$:

$$\text{Ric}(\cdot, \cdot) = [(n^2 - 1) - 1]\Xi g(\cdot, \cdot). \tag{6.7}$$

Here $n^2 - 1$ is the dimension of $SU(n)$ group. By using the relation between the Ricci tensor and the Killing form [61]

$$\text{Ric}(\cdot, \cdot) = -\frac{1}{4}B(\cdot, \cdot), \tag{6.8}$$

and the relation between the metric and the Killing form in eqs. (5.11) and (5.12)

$$B(\cdot, \cdot) = -ng(\cdot, \cdot), \tag{6.9}$$

we obtain

$$\Xi = \frac{n}{4(n^2 - 2)}. \tag{6.10}$$

Thus we have

$${}_2\mathcal{C}_m \sim \sqrt{n}, \quad \text{for } n \gg 1. \tag{6.11}$$

The critical time s_c is proportional to the maximum complexity as follows.

$$s_c = \frac{{}_2\mathcal{C}_m}{{}_2\tilde{F}(H)} \sim \mathcal{O}(\sqrt{n}) \sim \mathcal{O}(e^{d/2}) \sim e^{\mathcal{O}(d)}, \tag{6.12}$$

where we used the fact that if the classical phase space has size of order d the corresponding quantum Hilbert space have dimension: $n \sim \mathcal{O}(e^d)$.

After $s = s_c$ the complexity will stop grow and may decrease or oscillate. After a long enough time, the quantum recurrence can appear as argued in [5, 31]. The quantum recurrence theorem says that, for large n , the recurrent time is of order $\exp(n)$. Thus, there is a time $s = s_r \propto \exp(n)$ to make $\exp(Hs) \approx \hat{\mathbb{I}}$:

$${}_2\mathcal{C}(\exp(Hs_r)) \approx 0, \quad s_r \sim \mathcal{O}(\exp(e^d)), \tag{6.13}$$

which is the recurrent region in the figure 4. For systems with large degrees of freedom, the dimension of Hilbert space will be the exponential of entropy S ($n \sim e^S$) so

$${}_2\mathcal{C}_m \sim s_c \sim e^{\mathcal{O}(S)}, \quad s_r \sim \exp\left(e^{\mathcal{O}(S)}\right), \tag{6.14}$$

which is consistent with the conclusions in ref. [3].




Figure 5. Left panel: geometric explanation of $\hat{W}(t) = \hat{U}(t)\hat{W}_0\hat{U}(-t)$ and distance $d(t)$ (relative complexity). Right panel: $c_0(s)$ and $c_t(s)$ are two geodesics connecting from $\hat{\mathbb{I}}$ to \hat{W}_0 and $\hat{W}(t)$ respectively. It is possible that two geodesics $c_0(s)$ and $c_t(s)$ have the same length even though $\hat{W}(t) \neq \hat{W}_0$.

6.3 Complexity of precursors

It is also worth while to investigate the “complexity of precursors” studied in refs. [3, 29] and make some comparisons. For a unitary operator \hat{W}_0 and a time-dependent unitary operator $U(t)$, a precursor operator $\hat{W}(t)$ is defined as

$$\hat{W}(t) := \hat{U}(t)\hat{W}_0\hat{U}(-t), \tag{6.15}$$

where note that \hat{W}_0 is a unitary operator rather than a Hermitian observable. A geometrical explanation of $\hat{W}(t)$ on a group manifold is shown in figure 5.

To quantify the time-dependent property of $\hat{W}(t)$ related to complexity, the complexity ${}_p\mathcal{C}(\hat{W}(t))$ itself is not a good quantity because ${}_p\mathcal{C}(\hat{W}(t))$ may not change because of the adjoint invariance even if $\hat{W}(t)$ changes. See the right panel of figure 5 for a schematic example. A better one to characterize the evolution of precursor operator $\hat{W}(t)$ is the distance $d(t)$ between \hat{W}_0 and $\hat{W}(t)$ defined as

$$d(t) := \min \int_0^1 {}_pF(c(s), \dot{c}(s)) ds, \quad c(0) = \hat{W}_0, \quad c(1) = \hat{W}(t), \tag{6.16}$$

which is the minimal length of the geodesic connecting \hat{W}_0 and $\hat{W}(t)$. We may call this ‘relative complexity’ from \hat{W}_0 to $\hat{W}(t)$. Because of the right-invariance of the Finsler metric, eq. (6.16) is equivalent to

$$d(t) = \min \int_0^1 {}_pF(c(s), \dot{c}(s)) ds, \quad c(0) = \hat{\mathbb{I}}, \quad c(1) = \hat{W}(t)\hat{W}_0^{-1}, \tag{6.17}$$

which is just the complexity of $\hat{W}(t)\hat{W}_0^{-1}$.

By taking $\hat{W}_0 = \exp(H_w)$ and $U = \exp(Ht)$ we have

$$d(t) = {}_p\mathcal{C}(\hat{W}(t)\hat{W}_0^{-1}) = {}_p\mathcal{C}(\exp(Ht)\exp(H_w)\exp(-Ht)\exp(-H_w)). \tag{6.18}$$

While it is not easy to write down the explicit function of $d(t)$ in term of t , for $t \ll 1$ the expression may be written as

$$\exp(Ht) \exp(H_w) \exp(-Ht) \exp(-H_w) \approx \exp(\Lambda t + \mathcal{O}(t^2)), \quad (6.19)$$

where $\Lambda = \Lambda_1 + \Lambda_2$ and Λ_1 and Λ_2 are given by the Baker-Cambell-Hausdorff formula

$$\begin{aligned} \Lambda_1 &= \frac{1}{2}[H, H_w] + \frac{1}{12}[H_w, [H_w, H]] + \dots \\ \Lambda_2 &= \frac{1}{2}[H, H_w] - \frac{1}{12}[H_w, [H_w, H]] + \dots \end{aligned} \quad (6.20)$$

According to eq. (5.15), we obtain

$$d(t) = {}_p\tilde{F}(\Lambda)|t| + \mathcal{O}(t^2), \quad (6.21)$$

which means the difference between $\hat{W}(t)$ and \hat{W}_0 will increase linearly in time at the beginning. This result is similar to the result in ref. [29] when the time satisfies $t \ll 1$.

In particular, for $H_w = \epsilon \tilde{H}$, where \tilde{H} and H are two generators and $\epsilon \sim t \ll 1$, we have

$$\exp(Ht) \exp(H_w) \exp(-Ht) \exp(-H_w) \approx \exp([H, \tilde{H}]\epsilon^2). \quad (6.22)$$

The distance (relative complexity) becomes

$$d(\epsilon) = {}_p\mathcal{C}(\exp([H, \tilde{H}]\epsilon^2)) = \epsilon^2 {}_p\tilde{F}([H, \tilde{H}]). \quad (6.23)$$

On the other hand, if \tilde{H} and H are orthonormal generators, the sectional curvature spanned by H and \tilde{H} in a bi-invariant Finsler geometry can be expressed as [59]

$$K_{[H, \tilde{H}]}(H, \tilde{H}) = \frac{1}{4}\tilde{g}([H, \tilde{H}], [H, \tilde{H}]) = \frac{1}{4}{}_p\tilde{F}([H, \tilde{H}])^2, \quad (6.24)$$

where the notation $K_Y(H, \tilde{H})$ means that the sectional curvature spanned by $\{H, \tilde{H}\}$ with a reference vector Y . In eq. (6.24), the reference vector is $Y = [H, \tilde{H}]$. To distinguish the sectional curvature in general Finsler geometry from the one in Riemannian geometry defined in eq. (6.1), we use a different symbol in eq. (6.24). For $p = 2$, $K_Y(H, \tilde{H})$ becomes the sectional curvature in Riemannian geometry defined by eq. (6.1), which depends only on the plane spanned by $\{H, \tilde{H}\}$. For $p \neq 2$, the $K_Y(H, \tilde{H})$ depends not only on the plane spanned by $\{H, \tilde{H}\}$ but also on a choice of reference vector Y . For more details about the sectional curvatures in bi-invariant Finsler geometry, we refer to refs. [54, 55, 58, 59]

Thus, we obtain a geometrical interpretation of the relative complexity of the precursor operator:

$$d(\epsilon) = \epsilon^2 2\sqrt{K_{[H, \tilde{H}]}(H, \tilde{H})}. \quad (6.25)$$

For two orthonormal generators H and \tilde{H} , the square root of sectional curvature $K_{[H, \tilde{H}]}(H, \tilde{H})$ quantifies the minimal required quantum gates to change the infinitesimal operator $\exp(\tilde{H}\epsilon)$ to its infinitesimal precursor $\exp(H\epsilon) \exp(\tilde{H}\epsilon) \exp(-H\epsilon)$.

7 Conclusion

In this paper we generalize the result in ref. [49] in two ways. First, we generalize the parallel decomposition rule: **G3a** \rightarrow **G3b**. It opens a possibility to consider the p -norm. Second, we allow the complexity of the infinitesimal operator can depend on other factors w_α than the generator itself: **G4** \rightarrow **G4a**. For example, $w_\alpha(\alpha = r, l)$ may represent the penalty factors. We showed that the, even with two generalizations, the complexity geometry is still Finsler geometry.

We further constrain the Finsler geometry by these two properties. For $\forall \hat{U} \in \text{SU}(n)$ and $\forall H_\alpha \in \mathfrak{su}(n)$,

$$\begin{aligned} \text{[adjoint invariance]} \quad & {}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(\hat{U}H_\alpha\hat{U}^\dagger, w_\alpha), \\ \text{[reversibility]} \quad & {}_p\tilde{F}(H_\alpha, w_\alpha) = {}_p\tilde{F}(-H_\alpha, w_\alpha), \end{aligned} \tag{7.1}$$

The first is implied by the geometric idea of the complexity (i.e. independence of the curve length on the left-right generator) and supported by the physical symmetric transformations of the complexity. The second is supported by the CPT invariance of the complexity.

The complexity of an operator is given by minimal geodesic length in the Finsler geometry satisfying the constraints (7.1). Note that the adjoint invariance implies the Finsler metric is bi-invariant. Thanks to this bi-invariance the geodesic is given by constant generators and can be computed easily. We have shown that the complexity of $\text{SU}(n)$ operator, \hat{O} , is given by eq. (5.15):

$$\mathcal{C}(\hat{O}) = \lambda(w) \left\{ \text{Tr} \left[\left(\bar{H} \bar{H}^\dagger \right)^{p/2} \right] \right\}^{1/p}, \quad \forall \bar{H} = \ln \hat{O}, \tag{7.2}$$

where we choose the minimal value among multi-values of $\ln \hat{O}$. Note that eq. (7.2) depends on $w = w_r = w_l$ and p . If $w = 0$ and $p = 1$ it becomes the result in ref. [49]. A new ingredient w affects the complexity but only as an overall constant, so it is not essential. Note that even though different p gives different complexity, the geodesics are the same for all ps . i.e. the qualitative geometric and topological properties of the complexity based on the geodesic will not be changed for different ps . Note that if $p = 2$ the complexity geometry is given by the bi-invariant Riemannian metric and easier to handle. Thus, by analyzing $p = 2$ case, we can figure out qualitative properties of the complexity for $p \neq 2$.

We have also discussed a few interesting properties of (7.2). First, we investigated the geodesic deviation and chaos. In a manifold M with a bi-invariant metric, the sectional curvature $k_M(X, Y)$ is always nonnegative for any pair of tangent vectors $\{X, Y\}$, which may imply that the geodesics converge and there cannot be any chaotic behavior [61]. However, we pointed out that the geodesic deviation is determined by the sectional curvature ($k_{\mathcal{M}}(X, Y)$) of a two dimensional submanifold $\mathcal{M} \subset M$ where two geodesic $\xi_1(s)$ and $\xi_2(s)$ belongs to. Therefore, even in a manifold M with a bi-invariant metric, it is possible that the neighboring geodesics diverge in a region where $k_{\mathcal{M}}(X, Y)$ is non-positive, which may lead a chaotic behavior. As another supporting argument, we note that the induced metric in \mathcal{M} may not be bi-invariant even though the metric in M is bi-invariant.

We also showed that if the geodesic is given by a constant generator, the sectional curvature can be negative only in some part of $\xi_1(s)$, not in the whole $\xi_1(s)$.¹³ This statement is applied to *both* our bi-invariant metric and the “ k -local” metric case, because the geodesics are given by constant generators for both cases. Therefore, the bi-invariant metric and “ k -local” metric have no essential difference on the aspect of geodesic deviation. Furthermore, because the geodesics both in “ k -local” metric of refs. [29–31] and in a Finsler geometry of this paper (also ref. [49]) are generated by a constant generator, most properties in refs. [29–31] will also appear in our Finsler geometry and the predictions given by two theories have no contradictions. In particular, all the results given by “ k -local” subspace in ref. [31] will also appear in our bi-invariant Finsler geometry.

Next, we have shown that the pattern of the time-evolution of the complexity conjectured in [5, 31] is concretely realized in our complexity. i) the complexity grows linearly because the generator is constant thanks to the bi-invariance of Finsler geometry ii) the complexity reaches its maximum value in the exponential time ($t \sim e^{d/2}$) because of the compactness of $SU(n)$ group and the relation between the topology and curvature of $SU(n)$ groups. Finally, we have investigated the complexity of the precursor operators and found (i) the complexity of the precursor operator grows linearly at early time (ii) the complexity of precursors for infinitesimal operators corresponds to the sectional curvature.

In this paper we considered the complexity of the operator in the $SU(n)$ group with finite n . This result is extended to the case of $SU(\infty)$ and a non-compact group $Sp(2N, R)$ in [60]. To study the complexity of operators in quantum mechanics, we need to choose Hamiltonians for the generators in unitary representations. Non-compact groups have ‘infinite’ dimensional unitary representations. Therefore, to study non-compact groups we need to generalize our method in this paper to deal with ‘infinite’ dimensional unitary groups, which is a main technical achievement in ref. [60]. Interestingly, ref. [60] has shown an AdS_3 spacetime (rather than only a time slices in AdS_3) emerges as a complexity geometry for $Sp(2,1)$ or $SU(1,1)$ operator.

Another important future research direction is about the complexity between states. Based on our work in this paper we have developed the way to compute the complexity between states [64], where we have found a good agreement with the holographic results.

Acknowledgments

The work of K.-Y. Kim and C. Niu was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (NRF-2017R1A2B4004810) and GIST Research Institute (GRI) grant funded by the GIST in 2019. C. Niu is also supported by the Natural Science Foundation of China under Grant No. 11805083. C.-Y. Zhang is supported by

¹³There is another way to see it. If we study some topological properties such as the conjugate points, which do not depend on the exact values of curve lengths, we can set $p = 2$ and use bi-invariant Riemannian metric in $SU(n)$ groups (as discussed below eq. (7.2)). In this case it can be shown explicitly that even a bi-invariant complexity geometry could have 2-dimensional sub-manifolds with negative sectional curvature in some regions.

Project funded by China Postdoctoral Science Foundation. We also would like to thank the APCTP (Asia-Pacific Center for Theoretical Physics) focus program, “Holography and geometry of quantum entanglement” in Seoul, Korea for the hospitality during our visit, where part of this work was done.

A Relationship between the time order and left/right orders

Let us consider the time sequences $(t_1 < t_2 < \dots < t_n)$ and \hat{U}_n , the operator acting on the system at $t = t_n$. First, let us distinguish the time-order from the left or right order.

The time order, denoted by $(\mathcal{T} \prod_{k=1}^n) \hat{U}_k$, determines how to add a new operator to the old operators. For example, it is possible, for $n = 4$, that

$$\left(\mathcal{T} \prod_{k=1}^n\right) \hat{U}_k = \hat{U}_4 \hat{U}_3 \cdots \hat{U}_1 \hat{U}_2, \tag{A.1}$$

where at $t = t_2$ the operator \hat{U}_2 is added to the right, at $t = t_3$ the operator \hat{U}_3 is added to the right and at $t = t_4$ the operator \hat{U}_4 is added to the right. This kind of time order looks not so natural but can be allowed for discrete quantum circuits. However, from the perspective of group theory, only the following ‘left-order’ or ‘right-order’ are *equally* natural for a ‘time-order’.

- 1 Left-order \mathcal{L} : a new operator can appear only at the left-side of old operators.

$$\left(\mathcal{T} \prod_{k=1}^n\right) U_k = \hat{U}_n \hat{U}_{n-1} \cdots \hat{U}_2 \hat{U}_1 := \left(\mathcal{L} \prod_{k=1}^n\right) \hat{U}_k \tag{A.2}$$

- 2 Right-order \mathcal{R} : a new operator can appear only at the right-side of old operators.

$$\left(\mathcal{T} \prod_{k=1}^n\right) U_k = \hat{U}_1 \hat{U}_2 \cdots \hat{U}_{n-1} \hat{U}_n =: \left(\mathcal{R} \prod_{k=1}^n\right) \hat{U}_k \tag{A.3}$$

For the left-order the complexity is right-invariant while for the right-order the complexity is left-invariant.

Let us recall that both the left-order and right-order naturally appear in quantum mechanics. For example, the Schrödinger’s equation can be written as¹⁴

$$\frac{d}{dt} |\psi(t)\rangle = H(t) |\psi(t)\rangle, \tag{A.4}$$

which yields the evolution operator $c(t)$ given by the left-order product:

$$c(t) = \exp\left(\overleftarrow{\mathcal{P}} \int_0^t H(s) ds\right). \tag{A.5}$$

¹⁴Here $H(s)$ is anti-Hermitian.

This shows that the time order corresponds to the left-order. However, it is just a convention to write the Schrödinger's equation in terms of “ket” state $|\psi\rangle$. We can equally use “bra” state $\langle\psi|$ to express the Schrödinger's equation:

$$\frac{d}{dt}\langle\psi(t)| = -\langle\psi(t)|H(s), \tag{A.6}$$

which gives that evolution operator $[c(t)]^{-1}$ given by the right-order product:

$$[c(t)]^{-1} = \exp\left(-\vec{\mathcal{P}} \int_0^t H(s)ds\right) \tag{A.7}$$

This shows that the time order corresponds to the right-order.

Although we usually use the left-order to present the time-order in quantum theories, we can also use the right-order. Now a question arises: for a given unitary operator \hat{U} , can we know if it is built according to the “ $c(t)$ ” manner (adding the new operator at the left) or according to the “ $[c(t)]^{-1}$ ” manner (adding the new operator at the right side)? In quantum circuits, the answer is yes because a real quantum circuit is created by ourselves (we can decide which manner will be used.) However, for physically evolving systems in nature, there seems no a priori reason to choose either of them. If a complexity theory gives two different values for \hat{U} by these two ways, which way is employed by nature? A natural assumption is that two ways give the same complexity:

$${}_p\tilde{F}(H_l, w_l) = {}_p\tilde{F}(H_r, w_r), \tag{A.8}$$

which is shown to be true if the complexity is bi-invariant, in this paper.

B Two kinds of special symmetric transformations

In this appendix we will introduce two kinds of special symmetric transformations for the complexity, which are used in section 4.1.2.

B.1 Adding a divergent term

Let us consider a Lagrangian for a particle in flat space

$$\mathcal{L} := \mathcal{L}(\vec{x}, \dot{\vec{x}}, t), \tag{B.1}$$

the canonical momentum \vec{p} is defined by

$$p_j := \frac{\partial \mathcal{L}}{\partial \dot{x}^j}, \quad j = 1, 2, 3, \tag{B.2}$$

from which we can obtain $\dot{\vec{x}}$ as a function of \vec{x}, \vec{p} , and t , i.e. $\dot{\vec{x}} = \dot{\vec{x}}(\vec{x}, \vec{p}, t)$. The corresponding Hamiltonian reads

$$\mathbb{H} = \dot{\vec{x}} \cdot \vec{p} - \mathcal{L} = \mathbb{H}(\vec{x}, \vec{p}, t). \tag{B.3}$$

For a transformation

$$S_\varphi : \mathcal{L} \rightarrow \mathcal{L} + \vec{\nabla}\varphi(\vec{x}) \cdot \dot{\vec{x}}, \tag{B.4}$$

with an arbitrary smooth scalar field $\varphi(\vec{x})$, the Hamiltonian \mathbb{H} eq. (B.3) is transformed as

$$S_\varphi(\mathbb{H}) = \mathbb{H}(\vec{x}, \vec{p} - \vec{\nabla}\varphi(\vec{x}), t). \tag{B.5}$$

This transformation S_φ does not change physics such as the Feynman propagator and has a unitary representation

$$S_\varphi(\mathbb{H}) = \hat{U}_\varphi \mathbb{H} \hat{U}_\varphi^\dagger, \tag{B.6}$$

with

$$\hat{U}_\varphi = \exp[i\varphi(\hat{x})]. \tag{B.7}$$

Any composition of S_φ also gives a new Hamiltonian yielding the same physics:

$$(S_{\varphi_1} \circ S_{\varphi_2})(\mathbb{H}) = \mathbb{H}(\vec{x}, \vec{p} - \vec{\nabla}(\varphi_1 + \varphi_2), t). \tag{B.8}$$

Thus, S_φ and their products (defined by the composition of maps) form a symmetric group G_s for ${}_p\tilde{F}(-i\mathbb{H}, w_\alpha)$. G_s is an *infinite* dimensional Lie group with the following representation for its Lie algebra \mathfrak{g}_s

$$\mathfrak{g}_s := \left\{ i\varphi(\hat{x}) \mid \forall C^\infty \text{ scalar field } \varphi \right\}. \tag{B.9}$$

B.2 Canonical transformation

Next, let consider another kind of symmetric transformations, special canonical transformations. For example, let us take one-dimensional case:

$$S_f : (x, p) \mapsto (X(x, p), P(x, p)). \tag{B.10}$$

We will show that if (X, P) and (x, p) are transformed by a *constant* linear transformation

$$\begin{pmatrix} X \\ P \end{pmatrix} = \begin{pmatrix} f_1 & f_2 \\ f_3 & f_4 \end{pmatrix} \begin{pmatrix} x \\ p \end{pmatrix}, \tag{B.11}$$

with

$$f_1 f_4 - f_2 f_3 = 1, \tag{B.12}$$

then two Hamiltonians $\mathcal{H}(\hat{x}, \hat{p})$ and

$$S_f(\mathcal{H}) := \mathcal{H}(X(\hat{x}, \hat{p}), P(\hat{x}, \hat{p})), \tag{B.13}$$

describes the equivalent physics so should give the same complexity.¹⁵

First, these two Hamiltonians describe the same classical systems. This can be understood by noting that a symplectic structure is invariant under the transformation

$$dx \wedge dp = dX \wedge dP, \tag{B.14}$$

which is equivalent to

$$\{X, P\}_{P.B.} := \frac{\partial X}{\partial x} \frac{\partial P}{\partial p} - \frac{\partial X}{\partial p} \frac{\partial P}{\partial x} = 1 = \{x, p\}_{P.B.}. \tag{B.15}$$

¹⁵For example, if $\mathcal{H}(\hat{x}, \hat{p}) = \hat{x}^2 + \hat{p}^2$ then $S_f(\mathcal{H}) = (f_1^2 + f_3^2)\hat{x}^2 + (f_1 f_2 + f_3 f_4)(\hat{x}\hat{p} + \hat{p}\hat{x}) + (f_2^2 + f_4^2)\hat{p}^2$.

Thus, two Hamiltonians \mathcal{H} and $S_f(\mathcal{H})$ are transformed by a canonical transformation so describe the same physics in classical mechanics.

Next, in order to show that two Hamiltonians \mathcal{H} and $S_f(\mathcal{H})$ are equivalent in quantum level, let us consider the “phase-space formulation of quantum mechanics” [65, 66]. In this formulation, the quantum state is described by a quasi-probability distribution: the Wigner quasi-probability distribution $W(x, p)$ [67], which is the Wigner transform of the density matrix $\hat{\rho}$

$$W(x, p) := \frac{1}{\pi\hbar} \int \langle x + y | \hat{\rho} | x - y \rangle e^{-2ipy/\hbar} dy. \quad (\text{B.16})$$

If $\hat{A}(\hat{x}, \hat{p})$ is an operator representing an observable, its expectation value with respect to the phase-space state distribution $W(x, p)$ is,

$$\langle \hat{A} \rangle := \text{Tr}(\rho \hat{A}) = \int A(x, p) W(x, p) dx dp. \quad (\text{B.17})$$

Let us define the star product “ \star ” and the Moyal bracket “ $\{\{\cdot, \cdot\}\}$ ” as follows

$$h \star g := h \exp \left[\frac{1}{i\hbar} \left(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x \right) \right] g, \quad (\text{B.18})$$

and

$$\{\{h, g\}\} := -\frac{1}{i\hbar} (h \star g - g \star h) = -\frac{2}{\hbar} h \sin \left[\frac{\hbar}{2} \left(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x \right) \right] g. \quad (\text{B.19})$$

The eigenvalue a and the eigenstate distribution $W_a(x, p)$ of the observable \hat{A} are given by the following \star -eigenvalue equation

$$A(x, p) \star W_a(x, p) = a W_a(x, p). \quad (\text{B.20})$$

For a given state distribution $W(x, p)$, the possibility of obtaining eigenvalue a is given by

$$R_a = 2\pi\hbar \int W_a(x, p) W(x, p) dx dp. \quad (\text{B.21})$$

For a given Hamiltonian \mathcal{H} , the time evolution equation of an arbitrary state distribution $W(x, p)$ is given by

$$\frac{\partial W}{\partial t} = -\{\{W, \mathcal{H}\}\}. \quad (\text{B.22})$$

Now let us show that the transformation S_f does not change physics in quantum level. The proof consists of the following three steps:

(1) The expectation values of observables. An observable \hat{A} and a state distribution $W(x, p)$ will be transformed as

$$S_f(\hat{A}) = \hat{A}(X(\hat{x}, \hat{p}), P(\hat{x}, \hat{p})), \quad S_f(W)|_{(x,p)} = W(X(x, p), P(x, p)). \quad (\text{B.23})$$

The expectation value $\langle \hat{A} \rangle$ in eq. (B.17) is transformed as

$$\begin{aligned}
 S_f(\langle \hat{A} \rangle) &= \int S_f(A) S_f(W) dx dp = \int A(X(x, p), P(x, p)) W(X(x, p), P(x, p)) dx dp \\
 &= \int A(X, P) W(X, P) \det \mathbf{J} dX dP = \int A(X, P) W(X, P) dX dP \quad (\text{B.24}) \\
 &= \int A(x, p) W(x, p) dx dp = \langle \hat{A} \rangle.
 \end{aligned}$$

In the second line of eq. (B.24), we have made a variables transformation $(x, p) \rightarrow (X, P)$ in the integration and $\det \mathbf{J}$ is the Jacobi determinant of this variables transformation. Eqs. (B.11) and (B.12) imply

$$\det \mathbf{J} = f_1 f_4 - f_2 f_3 = 1. \quad (\text{B.25})$$

Eq. (B.24) shows that the expectation value of observable is invariant under the transformation S_f .

(2) The possible measurable values of observables. For arbitrary functions $h = h(x, p)$ and $g = g(x, p)$, eqs. (B.11) and (B.12) imply

$$S_f(h) (\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)^n S_f(g) \Big|_{x=x_0, p=p_0} = h (\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)^n g \Big|_{x=X(x_0, p_0), p=P(x_0, y_0)}, \quad (\text{B.26})$$

with $n = 0, 1, 2, 3, \dots$ so

$$S_f(h) \star S_f(g) \Big|_{x=x_0, p=p_0} = h \star g \Big|_{x=X(x_0, p_0), p=P(x_0, y_0)}. \quad (\text{B.27})$$

For an arbitrary observable \hat{A} with its eigenvalue a and eigenstate distribution $W_a(x, p)$,

$$\begin{aligned}
 S_f(A) \star S_f(W_a) \Big|_{x=x_0, p=p_0} &= A \star W_a \Big|_{x=X(x_0, p_0), p=P(x_0, y_0)} \\
 &= a W_a \Big|_{x=X(x_0, p_0), p=P(x_0, y_0)} = a S_f(W_a) \Big|_{x=x_0, p=p_0}.
 \end{aligned} \quad (\text{B.28})$$

This means that a and $S_f(W_a)$ are the eigenvalue and eigenstate distribution of observable $S_f(\hat{A})$. Thus, S_f will not change the possible measurable values of any observable.

(3) The possibility of measurable values. For an arbitrary observable \hat{A} and a given state distribution $W(x, p)$, let us assume that the possibility of obtaining a measurable value a is R_a . Under the transformation S_f , a is still the possible value of measurement but the corresponding eigenstate distribution is $S_f(W)$. Under the transformation S_f , the possibility of obtaining a is

$$S_f(R_a) = 2\pi\hbar \int S_f(W_a) S_f(W) dx dp. \quad (\text{B.29})$$

Using similar steps in eq. (B.24), we find that

$$S_f(R_a) = R_a. \quad (\text{B.30})$$

Thus, the possibility of obtaining a measurable value a is also invariant under the transformation S_f .

These three results show that the transformation S_f will not change the results of all measurements in quantum level. Therefore, \mathcal{H} and $S_f(\mathcal{H})$ describe two equivalent physical systems in quantum mechanics. \square

In the above proof, it is important that the coefficients in eq. (B.11) are constants and satisfy (B.12). Non-constant transformations satisfying eq. (B.12) but does not satisfying (B.26) will generate equivalent classical systems but inequivalent in quantum level. For our purpose, it is enough to specify S_f by setting $f_1 = f_4 = 1, f_3 = 0$ and $f_2 = f$ and we obtain

$$X(\hat{x}, \hat{p}) = \hat{x} + f\hat{p} = \hat{W}_f \hat{x} \hat{W}_f^\dagger, \quad P(\hat{x}, \hat{p}) = \hat{p} = \hat{W}_f \hat{p} \hat{W}_f^\dagger, \quad S_f(\mathcal{H}) = \hat{W}_f \mathcal{H} \hat{W}_f^\dagger, \quad (\text{B.31})$$

with

$$\hat{W}_f = \exp(if\hat{p}^2/2). \quad (\text{B.32})$$

Here, \hat{W}_f can form a one-dimensional Lie group with Lie algebra $\tilde{\mathfrak{g}}_s := \{ic\hat{p}^2 \mid \forall c \in \mathbb{R}\}$. This result can be generalized to higher dimensional cases and we obtain another symmetric group of the Finsler metric with the following Lie algebra

$$\tilde{\mathfrak{g}}_s := \left\{ ic^{jl} \hat{p}_j \hat{p}_l \mid \forall c^{jl} \in \mathbb{R} \right\}. \quad (\text{B.33})$$

C Explicit form of $\mathfrak{g}_s^{(\infty)}$

In this appendix, we will show the following two results for one-dimensional case:

$$\forall n \geq 0, \quad \forall \text{smooth } \varphi(x), \quad i\varphi(\hat{x})\hat{p}^n + i\hat{p}^n\varphi(\hat{x}) \in \mathfrak{g}_s^{(\infty)}, \quad (\text{C.1})$$

and

$$\mathfrak{g}_s^{(\infty)} = \left\{ iH(\hat{x}, \hat{p}) \mid \forall H(\hat{x}, \hat{p}) = H(\hat{x}, \hat{p})^\dagger, \right. \\ \left. H(x, p) \text{ is smooth and has a Taylor's expansion with respect to } p \text{ at } p = 0 \right\}. \quad (\text{C.2})$$

Proof. It is obvious that eq. (C.1) is true for $n = 0$. For $n \geq 1$, eq. (C.1) can be proven by a mathematical induction. First, $\forall n \geq 0$

$$\begin{aligned} [i\varphi(\hat{x})\hat{p}^n + i\hat{p}^n\varphi(\hat{x}), i\hat{p}^2] &= -[\varphi(\hat{x}), \hat{p}^2]\hat{p}^n - \hat{p}^n[\varphi(\hat{x}), \hat{p}^2] \\ &= -i\varphi' \hat{p}^{n+1} - i\hat{p}\varphi' \hat{p}^n - i\hat{p}^n\varphi' \hat{p} - i\hat{p}^{n+1}\varphi' \\ &= -2i(\varphi' \hat{p}^{n+1} + \hat{p}^{n+1}\varphi') + i([\varphi', \hat{p}]\hat{p}^n - \hat{p}^n[\varphi', \hat{p}]) \\ &= -2i(\varphi' \hat{p}^{n+1} + \hat{p}^{n+1}\varphi') + [i\varphi'', i\hat{p}^n], \end{aligned} \quad (\text{C.3})$$

where we set $\hbar = 1$. Note that $i\varphi(\hat{x})$ and $i\hat{p}^2$ belong to $\mathfrak{g}_s^{(\infty)}$, which is closed under commutators. If eq. (C.1) is true for $n = k$ then we have $i\hat{p}^k \in \mathfrak{g}_s^{(\infty)}$ and so

$$\forall \varphi(x), \quad [i\varphi(\hat{x}), i\hat{p}^k] \in \mathfrak{g}_s^{(\infty)}, \quad [i\varphi(\hat{x})\hat{p}^k + i\hat{p}^k\varphi(\hat{x}), i\hat{p}^2] \in \mathfrak{g}_s^{(\infty)}. \quad (\text{C.4})$$

Thus, eq. (C.3) implies

$$i\varphi(\hat{x})\hat{p}^{k+1} + i\hat{p}^{k+1}\varphi(\hat{x}) = -\frac{1}{2}[i\Phi(\hat{x})\hat{p}^k + i\hat{p}^k\Phi(\hat{x}), i\hat{p}^2] + \frac{1}{2}[i\varphi'(\hat{x}), i\hat{p}^k] \in \mathfrak{g}_s^{(\infty)}, \quad (\text{C.5})$$

where $\Phi(x) := \int \varphi(x)dx$. This proves eq. (C.1). As any linear combination of elements in $\mathfrak{g}_s^{(\infty)}$ is still in $\mathfrak{g}_s^{(\infty)}$, we can find that for arbitrary smooth functions $\{\varphi_n(\hat{x}), n = 0, 1, 2, \dots\}$

$$i \sum_{n=0}^{\infty} [\varphi_n(\hat{x})\hat{p}^n + \hat{p}^n\varphi_n(\hat{x})] \in \mathfrak{g}_s^{(\infty)},$$

which proves eq. (C.2). □

D “Local” and “nonlocal” Hamiltonians

In some references such as ref. [31], the “locality” plays an important role. It was argued that the nonlocal interactions should be “more complex” than local interactions. In general, by a unitary transformation, a Hamiltonian may change its “locality”, but the symmetry (4.1) implies the complexity is invariant. This seems to be contradictory. In this appendix, we will argue that this contradictory is due to the ambiguity of “locality” itself. Let us explain it by an example.

Before starting let us first make a clarification on our terminology. The words “non-locality/locality” appear in many different areas of physics and may stand for various different meanings. To avoid misunderstanding, we emphasize that the “nonlocality/locality” we discuss here are only about the interactions in a given Hamiltonian. The “nonlocality/locality” of quantum states and quantum correlations do not have any relationship to the following discussions.

Let us consider a Hamiltonian for N -lattices in 3-dimensional space

$$\mathcal{H}_1 = \frac{1}{2} \sum_{n=1}^{3N} \frac{p_n^2}{2m_n} + \frac{1}{2} \sum_{n,l=1}^{3N} f_{nl} q_n q_l, \quad (\text{D.1})$$

where $\{q_n, p_n\}$ (with $n = 1, 2, \dots, 3N$) are the canonical coordinates of lattices, m_n are the masses of the lattices, f_{nl} are the components of a symmetric non-diagonal matrix which does not have negative eigenvalues. It seems that the Hamiltonian \mathcal{H}_1 is “non-local” in the sense that every lattice has interactions with all others. However, we know that there is a unitary transformation making Hamiltonian \mathcal{H}_1 “local” as follows

$$\mathcal{H}_2 = \frac{1}{2} \sum_{n=1}^{3N} \frac{p_n^2}{2m_n} + \frac{1}{2} \sum_{n=1}^{3N} \lambda_n q_n^2, \quad (\text{D.2})$$

where λ_n are the eigenvalues of the matrix of f_{nl} . The “locality” looks changed when we transform \mathcal{H}_1 to \mathcal{H}_2 , although they are the same systems. Thus, the “nonlocality” in Hamiltonian \mathcal{H}_1 should not be “intrinsic”.

To make this concept clear, we will call a Hamiltonian is *apparently nonlocal (local)* if it contains nonlocal (only local) interactions in a given canonical variables. However,

we will call this Hamiltonian *intrinsically local* if there is a unitary transformation which renders it local. Thus, \mathcal{H}_1 is apparently nonlocal, \mathcal{H}_2 is apparently local and both are intrinsically local.

Notice again that \mathcal{H}_1 and \mathcal{H}_2 describe the same physical system and the apparent locality will not be physical. Thus, we may expect that the complexity is also invariant under the unitary transformation.

E Jacobi field and sectional curvature

In this subsection, we review on the Jacobi field and sectional curvature in Riemannian geometry. We will be brief and restrict ourselves to the only relevant part to this paper, referring to some textbooks, for example, refs. [63, 68] for more details.

In a N -dimensional Riemannian manifold $(M, g(\cdot, \cdot))$ let us consider two neighboring geodesics $\xi_1(s)$ and $\xi_2(s)$ laying in a 2-dimensional sub-manifold \mathcal{M} . Locally, such a 2-dimensional sub-manifold is determined uniquely by the neighboring geodesics $\xi_1(s)$ and $\xi_2(s)$. They start at the same point $s = 0$, i.e. $\xi_1(0) = \xi_2(0)$, and the angle $\delta\theta$ between their tangent vectors at $s = 0$ is infinitesimal. The *Jacobi field* along the geodesic $\xi_1(s)$ (or the geodesic deviation vector) is defined as

$$J(s) := \lim_{\delta\theta \rightarrow 0} \frac{\xi_2(s) - \xi_1(s)}{\delta\theta} = \lim_{\delta\theta \rightarrow 0} \frac{\exp[(H + \Delta\delta\theta)s] - \exp(Hs)}{\delta\theta}, \quad (\text{E.1})$$

which is a tangent vector in \mathcal{M} . In the second equality a specific case is considered: $\xi_1(s) = \exp(Hs)$ and $\xi_2(s) = \exp[(H + \Delta\delta\theta)s]$. If there is an induced Riemannian metric $\tilde{g}(\cdot, \cdot)$ in \mathcal{M} , the Jacobi field $J(s)$ satisfies the following equation

$$\frac{\tilde{D}}{ds} \frac{\tilde{D}}{ds} J(s) + R_{\mathcal{M}}(T(s), J(s))T(s) = 0, \quad (\text{E.2})$$

with the initial condition $J(0) = 0$. Here $T(s)$ is the tangent vector of $\xi_1(s)$ (i.e. $T(s) := \dot{\xi}_1(s) = \exp(Hs)H$). $\frac{\tilde{D}}{ds}$ is the directional covariant derivative along $T(s)$ (i.e. in a local coordinate system, $T^\mu \tilde{\nabla}_\mu$ where $\tilde{\nabla}_\mu$ is covariant derivative corresponding to metric $\tilde{g}(\cdot, \cdot)$). $R_{\mathcal{M}}$ is the Riemannian curvature tensor corresponding to metric $\tilde{g}(\cdot, \cdot)$.

Using the metric $\tilde{g}(\cdot, \cdot)$, we can make a unique decomposition such that $J(s) = J_\perp(s) + J_\parallel(s)$, where $J_\parallel(s)$ is parallel to $T(s)$ and $J_\perp(s)$ is orthogonal to $T(s)$. With this decomposition, the general solution for eq. (E.2) with condition $J(0) = 0$ may be expressed as

$$J(s) = \alpha_0 s T(s) + J_\perp(s), \quad (\text{E.3})$$

where α_0 is constant. Plugging (E.3) into eq. (E.2), we find that transverse component $J_\perp(s)$ satisfies

$$\frac{\tilde{D}}{ds} \frac{\tilde{D}}{ds} J_\perp(s) + R_{\mathcal{M}}(T(s), J_\perp(s))T(s) = 0, \quad J_\perp(0) = 0. \quad (\text{E.4})$$

Next, let us consider the vector $e_\perp(s)$, the parallel transport of $e_\perp(0)$ along $\xi_1(s)$ in \mathcal{M} , where $e_\perp(0)$ is the unit vector orthogonal to the tangent vector $T(0)$. i.e.

$$\frac{\tilde{D}}{ds} e_\perp(s) = 0, \quad \tilde{g}(e_\perp(0), T(0)) = 0, \quad (\text{E.5})$$

which implies that $\tilde{g}(e_{\perp}(s), T(s)) = 0$ for arbitrary s . The inner product of $e_{\perp}(s)$ and eq. (E.4) yields

$$\frac{\tilde{D}}{ds} \frac{\tilde{D}}{ds} \tilde{g}(J_{\perp}, e_{\perp}) + \tilde{g}(R_{\mathcal{M}}(T, J_{\perp})T, e_{\perp}) = 0, \quad (\text{E.6})$$

where eq. (E.5) was used.

Because \mathcal{M} is 2-dimensional there is a scalar function $f(s)$ such that

$$J_{\perp}(s) = f(s)e_{\perp}(s), \quad f(0) = 0, \quad (\text{E.7})$$

so eq. (E.6) yields

$$f''(s) + k_{\mathcal{M}}(e_{\parallel}, e_{\perp})\tilde{g}(T, T)f(s) = 0, \quad (\text{E.8})$$

where

$$e_{\parallel} = \frac{T}{\sqrt{\tilde{g}(T, T)}}, \quad (\text{E.9})$$

and

$$k_{\mathcal{M}}(e_{\parallel}, e_{\perp}) := \frac{\tilde{g}(R_{\mathcal{M}}(e_{\parallel}, e_{\perp})e_{\parallel}, e_{\perp})}{\tilde{g}(e_{\parallel}, e_{\parallel})\tilde{g}(e_{\perp}, e_{\perp}) - [\tilde{g}(e_{\parallel}, e_{\perp})]^2}, \quad (\text{E.10})$$

which is the *sectional curvature* of the section spanned by $\{e_{\parallel}, e_{\perp}\}$ embedded in \mathcal{M} . Here the denominator is indeed unity because $\{e_{\parallel}, e_{\perp}\}$ are an orthonormal set, but we keep this form for an easy comparison with eq. (6.2).

Note that the behavior of J_{\perp} is characterized by $f(s)$ in eq. (E.7) and eq. (E.8). Therefore, it is governed by $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$ not by $k_{\mathcal{M}}(e_{\parallel}, e_{\parallel})$, eq. (6.1). Depending on the sign of the sectional curvature $k_{\mathcal{M}}(e_{\parallel}, e_{\perp})$, the neighboring geodesics $\xi_1(s)$ and $\xi_2(s)$ may converge or diverge as s increases: in the regions for the sectional curvature is positive (negative) the geodesics approach to each other (go far away from each other). In particular, if the sectional curvature along $\xi_1(s)$ is positive almost everywhere¹⁶ $\xi_1(s)$ and $\xi_2(s)$ must intersect with each other at a point $s > 0$; if the sectional curvature along $\xi_1(s)$ is non-positive everywhere then $\xi_1(s)$ and $\xi_2(s)$ cannot intersect for $s > 0$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Aaronson, *The Complexity of Quantum States and Transformations: From Quantum Money to Black Holes*, [arXiv:1607.05256](https://arxiv.org/abs/1607.05256) [[INSPIRE](https://inspirehep.net/literature/1607052)].
- [2] D. Harlow and P. Hayden, *Quantum Computation vs. Firewalls*, *JHEP* **06** (2013) 085 [[arXiv:1301.4504](https://arxiv.org/abs/1301.4504)] [[INSPIRE](https://inspirehep.net/literature/1301450)].
- [3] L. Susskind, *Computational Complexity and Black Hole Horizons*, *Fortsch. Phys.* **64** (2016) 44 [[arXiv:1403.5695](https://arxiv.org/abs/1403.5695)] [[INSPIRE](https://inspirehep.net/literature/1403569)].
- [4] L. Susskind, *Entanglement is not enough*, *Fortsch. Phys.* **64** (2016) 49 [[arXiv:1411.0690](https://arxiv.org/abs/1411.0690)] [[INSPIRE](https://inspirehep.net/literature/1411069)].

¹⁶This means that the region where the sectional curvature is non-positive is of zero measure.

- [5] D. Stanford and L. Susskind, *Complexity and Shock Wave Geometries*, *Phys. Rev. D* **90** (2014) 126007 [[arXiv:1406.2678](#)] [[INSPIRE](#)].
- [6] D.A. Roberts, D. Stanford and L. Susskind, *Localized shocks*, *JHEP* **03** (2015) 051 [[arXiv:1409.8180](#)] [[INSPIRE](#)].
- [7] A.R. Brown, D.A. Roberts, L. Susskind, B. Swingle and Y. Zhao, *Holographic Complexity Equals Bulk Action?*, *Phys. Rev. Lett.* **116** (2016) 191301 [[arXiv:1509.07876](#)] [[INSPIRE](#)].
- [8] A.R. Brown, D.A. Roberts, L. Susskind, B. Swingle and Y. Zhao, *Complexity, action and black holes*, *Phys. Rev. D* **93** (2016) 086006 [[arXiv:1512.04993](#)] [[INSPIRE](#)].
- [9] R.-G. Cai, S.-M. Ruan, S.-J. Wang, R.-Q. Yang and R.-H. Peng, *Action growth for AdS black holes*, *JHEP* **09** (2016) 161 [[arXiv:1606.08307](#)] [[INSPIRE](#)].
- [10] L. Lehner, R.C. Myers, E. Poisson and R.D. Sorkin, *Gravitational action with null boundaries*, *Phys. Rev. D* **94** (2016) 084046 [[arXiv:1609.00207](#)] [[INSPIRE](#)].
- [11] S. Chapman, H. Marrochio and R.C. Myers, *Complexity of Formation in Holography*, *JHEP* **01** (2017) 062 [[arXiv:1610.08063](#)] [[INSPIRE](#)].
- [12] D. Carmi, R.C. Myers and P. Rath, *Comments on Holographic Complexity*, *JHEP* **03** (2017) 118 [[arXiv:1612.00433](#)] [[INSPIRE](#)].
- [13] A. Reynolds and S.F. Ross, *Divergences in Holographic Complexity*, *Class. Quant. Grav.* **34** (2017) 105004 [[arXiv:1612.05439](#)] [[INSPIRE](#)].
- [14] R.-Q. Yang, C. Niu and K.-Y. Kim, *Surface Counterterms and Regularized Holographic Complexity*, *JHEP* **09** (2017) 042 [[arXiv:1701.03706](#)] [[INSPIRE](#)].
- [15] D. Carmi, S. Chapman, H. Marrochio, R.C. Myers and S. Sugishita, *On the Time Dependence of Holographic Complexity*, *JHEP* **11** (2017) 188 [[arXiv:1709.10184](#)] [[INSPIRE](#)].
- [16] R.-Q. Yang, C. Niu, C.-Y. Zhang and K.-Y. Kim, *Comparison of holographic and field theoretic complexities for time dependent thermofield double states*, *JHEP* **02** (2018) 082 [[arXiv:1710.00600](#)] [[INSPIRE](#)].
- [17] B. Swingle and Y. Wang, *Holographic Complexity of Einstein-Maxwell-Dilaton Gravity*, *JHEP* **09** (2018) 106 [[arXiv:1712.09826](#)] [[INSPIRE](#)].
- [18] S. Chapman, H. Marrochio and R.C. Myers, *Holographic complexity in Vaidya spacetimes. Part I*, *JHEP* **06** (2018) 046 [[arXiv:1804.07410](#)] [[INSPIRE](#)].
- [19] S. Chapman, H. Marrochio and R.C. Myers, *Holographic complexity in Vaidya spacetimes. Part II*, *JHEP* **06** (2018) 114 [[arXiv:1805.07262](#)] [[INSPIRE](#)].
- [20] Y.-S. An, R.-G. Cai and Y. Peng, *Time Dependence of Holographic Complexity in Gauss-Bonnet Gravity*, *Phys. Rev. D* **98** (2018) 106013 [[arXiv:1805.07775](#)] [[INSPIRE](#)].
- [21] K. Nagasaki, *Complexity growth of rotating black holes with a probe string*, *Phys. Rev. D* **98** (2018) 126014 [[arXiv:1807.01088](#)] [[INSPIRE](#)].
- [22] S. Mahapatra and P. Roy, *On the time dependence of holographic complexity in a dynamical Einstein-dilaton model*, *JHEP* **11** (2018) 138 [[arXiv:1808.09917](#)] [[INSPIRE](#)].
- [23] M. Alishahiha, *Holographic Complexity*, *Phys. Rev. D* **92** (2015) 126009 [[arXiv:1509.06614](#)] [[INSPIRE](#)].
- [24] O. Ben-Ami and D. Carmi, *On Volumes of Subregions in Holography and Complexity*, *JHEP* **11** (2016) 129 [[arXiv:1609.02514](#)] [[INSPIRE](#)].

- [25] B. Chen, W.-M. Li, R.-Q. Yang, C.-Y. Zhang and S.-J. Zhang, *Holographic subregion complexity under a thermal quench*, *JHEP* **07** (2018) 034 [[arXiv:1803.06680](#)] [[INSPIRE](#)].
- [26] Y. Ling, Y. Liu and C.-Y. Zhang, *Holographic Subregion Complexity in Einstein-Born-Infeld theory*, *Eur. Phys. J. C* **79** (2019) 194 [[arXiv:1808.10169](#)] [[INSPIRE](#)].
- [27] C.A. Agón, M. Headrick and B. Swingle, *Subsystem Complexity and Holography*, *JHEP* **02** (2019) 145 [[arXiv:1804.01561](#)] [[INSPIRE](#)].
- [28] J. Couch, W. Fischler and P.H. Nguyen, *Noether charge, black hole volume and complexity*, *JHEP* **03** (2017) 119 [[arXiv:1610.02038](#)] [[INSPIRE](#)].
- [29] L. Susskind and Y. Zhao, *Switchbacks and the Bridge to Nowhere*, [arXiv:1408.2823](#) [[INSPIRE](#)].
- [30] A.R. Brown, L. Susskind and Y. Zhao, *Quantum Complexity and Negative Curvature*, *Phys. Rev. D* **95** (2017) 045010 [[arXiv:1608.02612](#)] [[INSPIRE](#)].
- [31] A.R. Brown and L. Susskind, *Second law of quantum complexity*, *Phys. Rev. D* **97** (2018) 086015 [[arXiv:1701.01107](#)] [[INSPIRE](#)].
- [32] R. Jefferson and R.C. Myers, *Circuit complexity in quantum field theory*, *JHEP* **10** (2017) 107 [[arXiv:1707.08570](#)] [[INSPIRE](#)].
- [33] A.P. Reynolds and S.F. Ross, *Complexity of the AdS Soliton*, *Class. Quant. Grav.* **35** (2018) 095006 [[arXiv:1712.03732](#)] [[INSPIRE](#)].
- [34] R. Khan, C. Krishnan and S. Sharma, *Circuit Complexity in Fermionic Field Theory*, *Phys. Rev. D* **98** (2018) 126001 [[arXiv:1801.07620](#)] [[INSPIRE](#)].
- [35] L. Hackl and R.C. Myers, *Circuit complexity for free fermions*, *JHEP* **07** (2018) 139 [[arXiv:1803.10638](#)] [[INSPIRE](#)].
- [36] M. Guo, J. Hernandez, R.C. Myers and S.-M. Ruan, *Circuit Complexity for Coherent States*, *JHEP* **10** (2018) 011 [[arXiv:1807.07677](#)] [[INSPIRE](#)].
- [37] A. Bhattacharyya, A. Shekar and A. Sinha, *Circuit complexity in interacting QFTs and RG flows*, *JHEP* **10** (2018) 140 [[arXiv:1808.03105](#)] [[INSPIRE](#)].
- [38] H.A. Camargo, P. Caputa, D. Das, M.P. Heller and R. Jefferson, *Complexity as a novel probe of quantum quenches: universal scalings and purifications*, *Phys. Rev. Lett.* **122** (2019) 081601 [[arXiv:1807.07075](#)] [[INSPIRE](#)].
- [39] S. Chapman, M.P. Heller, H. Marrochio and F. Pastawski, *Toward a Definition of Complexity for Quantum Field Theory States*, *Phys. Rev. Lett.* **120** (2018) 121602 [[arXiv:1707.08582](#)] [[INSPIRE](#)].
- [40] P. Caputa, N. Kundu, M. Miyaji, T. Takayanagi and K. Watanabe, *Anti-de Sitter Space from Optimization of Path Integrals in Conformal Field Theories*, *Phys. Rev. Lett.* **119** (2017) 071602 [[arXiv:1703.00456](#)] [[INSPIRE](#)].
- [41] P. Caputa, N. Kundu, M. Miyaji, T. Takayanagi and K. Watanabe, *Liouville Action as Path-Integral Complexity: From Continuous Tensor Networks to AdS/CFT*, *JHEP* **11** (2017) 097 [[arXiv:1706.07056](#)] [[INSPIRE](#)].
- [42] T. Takayanagi, *Holographic Spacetimes as Quantum Circuits of Path-Integrations*, *JHEP* **12** (2018) 048 [[arXiv:1808.09072](#)] [[INSPIRE](#)].
- [43] K. Hashimoto, N. Iizuka and S. Sugishita, *Time evolution of complexity in Abelian gauge theories*, *Phys. Rev. D* **96** (2017) 126001 [[arXiv:1707.03840](#)] [[INSPIRE](#)].

- [44] K. Hashimoto, N. Iizuka and S. Sugishita, *Thoughts on Holographic Complexity and its Basis-dependence*, *Phys. Rev. D* **98** (2018) 046002 [[arXiv:1805.04226](#)] [[INSPIRE](#)].
- [45] J.M. Magán, *Black holes, complexity and quantum chaos*, *JHEP* **09** (2018) 043 [[arXiv:1805.05839](#)] [[INSPIRE](#)].
- [46] P. Caputa and J.M. Magan, *Quantum Computation as Gravity*, [arXiv:1807.04422](#) [[INSPIRE](#)].
- [47] M. Flory and N. Miekley, *Complexity change under conformal transformations in AdS_3/CFT_2* , [arXiv:1806.08376](#) [[INSPIRE](#)].
- [48] R.-Q. Yang, *Complexity for quantum field theory states and applications to thermofield double states*, *Phys. Rev. D* **97** (2018) 066004 [[arXiv:1709.00921](#)] [[INSPIRE](#)].
- [49] R.-Q. Yang, Y.-S. An, C. Niu, C.-Y. Zhang and K.-Y. Kim, *Principles and symmetries of complexity in quantum field theory*, *Eur. Phys. J. C* **79** (2019) 109 [[arXiv:1803.01797](#)] [[INSPIRE](#)].
- [50] M.A. Nielsen, M.R. Dowling, M. Gu and A.C. Doherty, *Quantum computation as geometry*, *Science* **311** (2006) 1133.
- [51] M.A. Nielsen, *A geometric approach to quantum circuit lower bounds*, *Quant. Inf. Comput.* **6** (2006) 213.
- [52] M.R. Dowling and M.A. Nielsen, *The geometry of quantum computation*, *Quant. Inf. Comput.* **8** (2008) 861.
- [53] J. Watrous, *Theory of quantum information*, Cambridge University Press (2018).
- [54] D. Bao, S.-S. Chern and Z. Shen, *An Introduction to Riemann-Finsler Geometry*, Springer New York, New York (2000) [[DOI:10.1007/978-1-4612-1268-3](#)].
- [55] Z. Shen, *Lectures on Finsler geometry*, Series on Multivariate Analysis, Wspc (2001).
- [56] M. Xiaohuan, *An introduction to Finsler geometry*, World Scientific, Singapore Hackensack, NJ (2006).
- [57] G.S. Asanov, *Finsler Geometry, Relativity and Gauge Theories*, Springer Netherlands, Dordrecht (1985).
- [58] D. Latifi and M. Toomanian, *On the existence of bi-invariant finsler metrics on lie groups*, *Math. Sci.* **7** (2013) 37.
- [59] D. Latifi and A. Razavi, *Bi-invariant finsler metrics on lie groups*, *J. Basic Appl. Sci.* **5** (2011) 607.
- [60] R.-Q. Yang and K.-Y. Kim, *Complexity of operators generated by quantum mechanical Hamiltonians*, *JHEP* **03** (2019) 010 [[arXiv:1810.09405](#)] [[INSPIRE](#)].
- [61] M.M. Alexandrino and R.G. Bettiol, *Lie Groups with Bi-invariant Metrics*, Springer International Publishing, Cham (2015), pp. 27–47.
- [62] S. Deng and Z. Hou, *Positive definite minkowski lie algebras and bi-invariant finsler metrics on lie groups*, *Geom. Dedicata* **136** (2008) 191.
- [63] T. Frankel, *The geometry of physics: an introduction*, Cambridge University Press, Cambridge, New York (2012).
- [64] R.-Q. Yang, Y.-S. An, C. Niu, C.-Y. Zhang and K.-Y. Kim, *Complexity between states in quantum field theories*, work in progress.

- [65] C. Zachos, D. Fairlie and T. Curtright, *Quantum Mechanics in Phase Space: An Overview with Selected Papers*, World Scientific (2005).
- [66] T.L. Curtright and C.K. Zachos, *Quantum Mechanics in Phase Space*, *Asia Pac. Phys. Newslett.* **1** (2012) 37 [[arXiv:1104.5269](#)] [[INSPIRE](#)].
- [67] E.P. Wigner, *On the quantum correction for thermodynamic equilibrium*, *Phys. Rev.* **40** (1932) 749 [[INSPIRE](#)].
- [68] S.S. Chern, *Lectures on differential geometry*, World Scientific, Singapore River Edge, NJ (1999).